The AIRE G228W mutation disturbs the interaction of AIRE with its partner molecule SIRT1

Jadson C. Santos [1#]**, Mariangela Dametto** [2#]**, Ana Paula Masson** [3]**, Vitor M. Faça** [3]**, Rodrigo Bonacin** [2]**, Eduardo A. Donadi** [4]**, Geraldo Aleixo Passos** [1,5,6,*]

Molecular Immunogenetics Group, Department of Genetics, Ribeirão Preto Medical School, University of São Paulo (USP), Ribeirão Preto, SP, Brazil.

Renato Archer Information Technology Center (CTI Brazil), Ministry of Science, Technology and Innovation (MCTI), Campinas, SP, Brazil.

Department of Biochemistry and Immunology, Ribeirão Preto Medical School, University of São Paulo (USP), Ribeirão Preto, SP, Brazil.

Division of Clinical Immunology, Department of Medicine, Ribeirão Preto Medical School, USP, Ribeirão Preto, SP, Brazil.

Laboratory of Genetics and Molecular Biology, Department of Basic and Oral Biology, School of Dentistry of Ribeirão Preto, USP, Ribeirão Preto, SP, Brazil.

Center for Cell-Based Therapy in Dentistry, School of Dentistry of Ribeirão Preto, USP, Ribeirão Preto, SP, Brazil.

* Corresponding author at: Molecular Immunogenetics Group, Department of Genetics, Ribeirão Preto Medical School, University of São Paulo (USP), Ribeirão Preto, SP, Brazil. E-mail: passos@usp.br

# Equal contribution.

Abstract

The autoimmune regulator (AIRE) protein functions as a tetramer, interacting with partner proteins to form the "AIRE complex," which relieves RNA Pol II stalling in the chromatin of medullary thymic epithelial cells (mTECs). AIRE is the primary mTEC transcriptional controller, promoting the expression of a large set of peripheral tissue antigen genes implicated in the negative selection of self-reactive thymocytes. Under normal conditions, the SIRT1 protein temporarily interacts with AIRE and deacetylates K residues of the AIRE SAND domain. Once the AIRE SAND domain is deacetylated, the binding with SIRT1 is undone, allowing the AIRE complex to proceed downstream with the RNA Pol II to the elongation phase of transcription. Considering that the *in silico* and *in vitro* binding of the AIRE SAND domain with SIRT1 provides a powerful model system for studying the dominant SAND G228W mutation mechanism, which causes the autoimmune polyglandular syndrome-1, we integrated computational molecular modeling, docking, dynamics between the whole SAND domain with SIRT1, and surface plasmon resonance using a peptide harboring the 211 to 230 residues of the SAND domain, to compare the structure and energetics of binding/release between AIRE G228 (wild-type) and W228 (mutant) SAND domain to SIRT1. We observed that the G228W mutation in the SAND domain negatively influences the AIRE-SIRT1 interaction. The disturbed interaction might cause a disruption in the binding of the AIRE SAND domain with the SIRT1 catalytic site, impairing the AIRE complex to proceed downstream with RNA Pol II.

Key words AIRE, SAND domain, SIRT1 interaction, APS-1 mutation, molecular dynamics

Introduction

The human autoimmune regulator (*AIRE*) gene spans 14 exons at chromosome 21q22.3, encoding a 1635-nucleotide mRNA and a 545-amino acid protein (Aaltonen et al., 1994; Nagamine et al., 1997; Blechschmidt et al., 1999). Nearly 145 single-nucleotide mutations or large deletions have been observed along the *AIRE* gene (Wolff and Oftedal, 2019; Besnard et al., 2021; Human Gene Mutation Database available at https://www.hgmd.cf.ac.uk/ac/). Some mutations in the *AIRE* gene are related to autoimmune polyendocrinopathy-candidiasis ectodermal dystrophy (APECED) (Finnish-German APECED Consortium 1997; Nagamine et al., 1997), also known as autoimmune polyglandular syndrome type 1 (APS-1) (OMIM entry # 240300). Most APS-1 mutations are inherited in an autosomal recessive manner, except for a group of monoallelic mutations clustered within the zinc finger domain of the first plant homeodomain (PHD1) (Oftedal et al., 2015; Wolff and Oftedal, 2019) and a G228W mutation at the SAND domain, which follows a dominant inheritance pattern (Cetani et al., 2001; Ilmarinen et al., 2005; Su et al., 2008).

The 56-kDa AIRE protein is a transcriptional regulator in medullary thymic epithelial cells (Giraud et al., 2012; Passos et al., 2018; Perniola, 2018; Giraud and Peterson, 2019), exhibiting five domains and four LXXLL motifs. The caspase activation and recruitment (CARD) domain is essential for AIRE dimerization. The nuclear localization signal (NLS) domain presents a nuclear localization signal that mediates AIRE migration from the cytoplasm to the nucleus. The Sp100, AIRE-1, NucP41/75, and DEAF-1 (SAND) domains promote protein-protein interactions and

chromatin binding, which play a role in the nuclear speckle localization of AIRE, and the PHD1 and PHD2 domains function as histone code readers, interacting and supporting the process of chromatin decondensation (Anderson et al., 2016; Kumar et al., 2001; Perniola et al., 2014; Giraud and Peterson, 2019). During its action in chromatin, AIRE interacts with several partner proteins forming the "AIRE complex," in which SIRT1 is one of the most critical deacetylating K residues in the SAND domain, permitting the proper function of AIRE (Incani et al., 2014; Abramson et al., 2010; Abramson and Husebye, 2016).

In addition to chromatin binding, the SAND domain may also have a regulatory role in the homomultimerization of AIRE (Ramsey et al., 2002a; Halonen et al., 2004). The G228W SAND domain mutation was the first dominant mutation associated with the aggressiveness of APS-1 (Cetani et al., 2001; Ilmarinen et al., 2005; Su et al., 2008; Wolff and Oftedal, 2019); however, the reason for its dominant character and the molecular mechanism of APS-1 pathogenesis are still elusive.

Considering that i) the deacetylation of SAND domain K residues mediated by the partner SIRT1 molecule is essential for AIRE function in releasing RNA Pol II and allowing it to proceed with the elongation phase of transcription (Vaquero et al., 2004; Giraud et al., 2012, Saare et al., 2012, Incani et al., 2014, Chuprin et al., 2015, Bheda et al., 2016), and ii) the G228W mutation occurs at a position located at six amino acid residues from one of the acetylated K residues (K222), which is further deacetylated by SIRT1 (Saare et al., 2012), in this study, we hypothesized that the proximity of the G228W mutation to the K222 residue might interfere with the interaction between the SAND domain and the SIRT1 protein.

We integrated computational molecular modeling, docking, and dynamics between the whole AIRE SAND domain with the SIRT1 protein to test this hypothesis. Additionally, we evaluated the binding of a peptide harboring the 211 to 230 residues of the SAND domain with the SIRT1 protein to compare the binding/release between AIRE G228 (wild-type peptide) and W228 (mutant peptide) using surface plasmon resonance. The *in silico* analyses showed differences in the physical interaction between the AIRE SAND domain and the SIRT1 catalytic site for both wild-type and mutant domains. On the other hand, the surface plasmon resonance evaluation showed a stronger affinity (as seen through binding and dissociation) of the wild-type SAND domain peptide with the SIRT1 protein compared to a weaker association observed for the mutant peptide.

Materials and Methods

Protein structure homology modeling

We used the SWISS-MODEL workspace for protein modeling (http://swissmodel.expasy.org/workspace). The amino acid sequence in the FASTA format of the SIRT1 target protein was used as input. The SWISS-MODEL algorithms performed BLASTp and provided the best homology models based on global model quality estimation (GMQE) and qualitative model energy analysis (QMEAN) statistical parameters. The GMQE provided the quality by combining the target-template alignment properties, estimating a range between 0 and 1. The scoring function of

134 QMEAN consisted of a linear combination of structural descriptors, which is related to
135 the high-resolution X-ray crystal structure, estimating a Z-score scheme.
136
137 ***Molecular docking***
138
Molecular docking analysis of the modeled AIRE SAND domain, wild-type or
G228W mutant AIRE SAND domain, and the SIRT1 protein (PDB code 4I5I) was
performed to study the stability of the protein-protein interaction. The ClusPro 2.0
protein-protein docking server (Alekseenko et al., 2020) (available at
https://cluspro.org/) was used. In the ClusPro server, the rigid body docking phase uses
the PIPER docking program (https://www.schrodinger.com/products/piper), which
relies on the fast Fourier transform (FFT) correlation approach. PIPER represents the
interaction energy between two proteins using the
formula $E = w1Erep+w2Eattr+w3Eelec+w4EDARS$, where E_{rep} and E_{attr} denote the
attractive and repulsive contributions to the van der Waals interaction energy,
and E_{elec} represents the electrostatic energy. *EDARS* is a pairwise structure-based
potential; it primarily means desolvation contributions, i.e., free energy change by
removing the water molecules from the interface. The coefficients w_1, w_2, w_3,
and w_4 define the molecular weights of the corresponding amino acid residues.
153
154 ***Systems setup and molecular dynamics simulations***
155
156   As mentioned above, the three-dimensional (3D) structures of the complexes
157 formed with AIRE wild-type or with AIRE mutant peptide and SIRT1 generated in the
158 docking section were used as initial configurations for the molecular dynamics
159 simulations. In this work, calculations were performed to verify the stability and affinity
160 of the complexes.
161
162   To mimic the interaction between the two peptides and SIRT1 protein, the K222
163 residue was acetylated using the parameters given in the AMBER Molecular Dynamics
164 Package (https://ambermd.org/) for this residue. ACK and the connectivity library file
165 for K222 residue were generated by tleap.
166
The two systems were solvated using the TIP3P water model and neutralized
with four Na+ ions in the tleap tool implemented in AMBERTOOLS21
(https://ambermd.org/AmberTools.php). The whole system was truncated in a box with
15 Å between the edges of the periodic box and the molecules analyzed. An 8 Å cutoff
particle mesh Ewald (PME) method was used to treat the long-range electrostatic
interactions. AMBER molecular dynamics software and the ff19SB force field were
used to prepare the systems and run the simulations.
174
Before running, the complexes were subjected to minimization in two stages
using the steepest descent method followed by a conjugate gradient. In the first stage,
the elements, peptides, and SIRT1 were fixed, and the position of the water molecules
was minimized. After that, the whole system was minimized. A restrained
thermalization phase with the microcanonical ensemble (NVE) was performed,
increasing the temperature by 0.1 K until it reached 300 K using the Langevin
thermostat and SHAKE algorithm (https://ambermd.org/) to constrain hydrogen bonds.
182

The production dynamics were carried out at constant pressure (NPT ensemble) with a time step of 2 fs. The total simulation time for each complex was 2700 ns, and frames were saved every 20 ps. The analyses were performed in the last 2650 ns after the systems were equilibrated. All calculations shown in the current work were performed using AMBERTOOLS21 scripts. The visualization programs used to create the figures of the molecules were VMD for LINUXAMD64, version 1.9.3, and PYMOL (https://pymol.org/2/).

Surface plasmon resonance

The surface plasmon resonance (SPR) experiments were performed on Biacore Model T200 equipment (GE Healthcare Biosciences, Uppsala, Sweden). For the immobilization of human SIRT1 (catalog number S8846, Sigma–Aldrich, Saint Louis, MO), a CM5 sensor chip S series (GE Healthcare Biosciences) has carboxylic groups available for covalent coupling was used. All solutions used were purchased from the equipment supplier (GE Healthcare Biosciences). The procedures described here were performed automatically with the Biacore instrument using pre-established programs for binding amines in peptides or proteins.

Briefly, the carboxylic groups on the CM5 sensor chip surface were activated with EDC + NHS, forming the NHS ester reactive intermediate, which, after washing with HBS-N, readily reacted with amines dissolved in the immobilization solution. The human SIRT1 ligand was dissolved in 10 mM sodium acetate pH 4.0 plus 20 µM NAD+ (catalog number N1636, beta-nicotinamide adenine dinucleotide hydrate, Sigma–Aldrich), which was immobilized onto the Biacore sensor chip according to the manufacturer´s instructions.

To detect interactions between SIRT1 and AIRE WT or AIRE G228W mutant SAND domain peptides, different concentrations of peptides were injected onto the CM5 chip with immobilized SIRT1 protein. Peptide samples were injected sequentially from lowest to highest concentrations (20, 100, and 200 μM) with a contact time of 300 s, a 20 µl/min flow, and a dissociation time of 1200 s. The surface was regenerated between samples by injecting 2.0 M glycine pH 2.0 (30 s, 30 µl/min) followed by injection of 2.0 M NaCl (30 s, 30 µl/min). The stabilization period between injections after regeneration was 200 s.

The respective sequences of the twenty-amino acid peptide residues (Synbio Technologies, Monmouth Junction, NJ, USA) comprising positions 211 to 230 of the SAND domain of the human AIRE protein are as follows: from the C to N-terminus, AIRE wild-type (IQQVFESGGSK<u>K</u>CIQVGGEF) and AIRE G228W mutant (IQQVFESGGSK<u>K</u>CIQVGWEF). The underlined K residue represents the position where the acetyl radical was added. The experiment determined whether the K residue underlined was acetylated or not. The AIRE wild-type or AIRE G228W mutation data are available through GenBank NCBI (https://www.ncbi.nlm.nih.gov/) accession numbers (NM_000383.4): c.682G>T G [GGG] > W [TGG], coding sequence variant autoimmune regulator (NP_000374.1): p. Gly228TrpG (Gly) > W missense variant, DbSNP (https://www.ncbi.nlm.nih.gov/snp/rs121434257) and ClinVar (https://www.ncbi.nlm.nih.gov/clinvar/variation/3313/).

232   Kinetic data were analyzed with Anabel Analysis of Binding Events software
233 (Krämer et al., 2019), available at (www.skscience.org/anabel). Data obtained with the
234 different concentrations of AIRE WT or mutant peptides were combined in the same
235 graph.
236
237

238 **Results**

240 ***Homology molecular modeling of the*** AIRE *SAND domain*

241   To obtain a separate 3D human AIRE SAND domain structure, we used the
SwissModel tool to model proteins through homology. We used the SAND domain
sequence of the human SP100 protein (PDB code 1h5p) for the modeling, which
features 39.62% sequence identity to the human AIRE SAND domain. The SP100
protein SAND domain conserves the SKKCIQ amino acid sequence in the AIRE SAND
domain, corresponding to the accurate interaction site between SAND and the SIRT1
protein. Thus, obtaining a 3D structure of the wild-type AIRE SAND domain and the
AIRE SAND domain carrying the G228W mutation was possible. Considering that the
G228W mutation is located in a loop region of the AIRE SAND domain (Figure 1), we
may infer that this substitution does not cause local protein misfolding.

Blind molecular docking shows that the SAND W228 mutant residue was targeted to
the SIRT1 *catalytic residue*

253   To obtain an interaction model between the AIRE SAND domain (wild-
254 type or G228W mutant) and SIRT1 and assess the affinity of the mutant SAND domain,
255 we used the ClusPro 2.0 blind molecular docking tool. The docking was performed with
256 nonacetylated K222 to understand whether the modeled AIRE SAND domain would
257 have an affinity for the catalytic region of the SIRT1 protein.

258   Through blind molecular docking, it was possible to observe an affinity between
259 the AIRE SAND domain WT G228 or W228 mutant domain and the catalytic
260 region/site (H363 residue) of the SIRT1 protein, even without any previous targeting or
261 chemical modification of the K222 residue, which is a target of the SIRT1 protein
262 (Figure 2). Important to note that when K222 is nonacetylated, the structure formed is
263 incompatible with the expected native structure.

264   Comparing the structural conformations, we observed that the SAND domain
265 mobilizes the W228 mutant residue that interacts with the SIRT1 H363 catalytic residue
266 in the docking analyses (Figure 3).

267 *Molecular dynamics*

268   The molecular dynamics results with the AIRE SAND domain (wild-type or
G228W mutant) showed that even the acetylated K222 residue at 1 µs of simulation did
not allow the wild-type domain to form a stable structure, in which the K222 residue
would interact with the H363 residue from the catalytic site of SIRT1. On the other
hand, the G228W mutant domain exhibited a stable interaction with the SIRT1 catalytic
site at the H363 residue. The analysis of the dynamics was performed to understand if
K222 acetylation would be able to direct it to the catalytic residue of SIRT1.

The RMSD results showed that the WT SAND domain and SIRT1 complex is
slightly less stable along the simulation time (1200 ns) than the complex formed
between the SAND G228W mutant. Nevertheless, the G228W mutant domain complex
remained stable in the RMSD [Figure 4, where higher angstrom distance values (Y-
axis) indicate less stability]. The analysis of structures from molecular dynamics
showed that even having inserted an acetyl radical at K222 residue in the WT or mutant
domains, there was no targeting to the catalytic residue H363 of SIRT1 along the
molecular dynamics simulation time. For molecular docking analyses, we used the
SAND domain without chemical modification. Still, to understand the behavior of this
domain as the SIRT1 protein recognizes it, we acetylated K222 for dynamic analysis
(Figure 5).

Surface plasmon resonance indicates that the SAND G228W mutation modifies the
interaction kinetics of AIRE with SIRT1

Considering that, the results from molecular docking and molecular dynamics
exhibited slight differences when comparing WT and mutant SAND domains during the
formation of complexes with SIRT1, next, we performed surface plasmon resonance
(SPR) to study the behavior of the two types of peptides when they bind to SIRT1. The
SPR results indicated a differential association and dissociation dynamics between the
wild-type SAND G228 or mutant SAND W228 peptides and the SIRT1 protein. The
interaction kinetics showed that the wild-type SAND G228 peptide has a more
significant association capacity as observed at a concentration range of 20 to 200 µM.
The interaction between the wild-type G228 SAND peptide and SIRT1 was almost
completely dissociated after 1200 seconds of measurement. Conversely, the mutant
W228 SAND peptide was less able to associate with SIRT1 (Figure 6). These results,
combined with the *in silico* analyses, show that the G228W mutation changes the
SAND-SIRT1 interaction, which is necessary to activate the AIRE protein. Of note,
SPR results were in contrast with the blind docking and molecular dynamics, i.e.,
instead, the WT complex was more stable and mutant less stable. The SPR results were
reproduced with six different concentrations of each of the acetylated peptides, WT and
mutant (range 10 to 200 µM), in addition to the control (0 µM) (Supplementary Figure
1). For clarity of the results, we chose three concentrations (20, 100 and 200 µM) in
addition to the control (0 µM), whose results we plot the graph in Figure 6.

Discussion

We reported that the dominant G228W mutation observed at the SAND domain
of the human AIRE protein found in APS-1 patients changes interaction with the
deacetylase SIRT1 compared to the AIRE wild-type protein. Because of the interference
of the W228 mutant residue, the kinetics of the association and dissociation processes
differ compared to the wild-type, which might explain the nuclear accumulation of the
AIRE complex, as observed in AIRE mutant cells. Since the mutant AIRE protein does
not properly stabilize in the AIRE complex, it may abnormally accumulate in the
nucleus, as previously observed (Ilmarinen et al., 2005). The binding/dissociation
kinetics between AIRE-SIRT1 is crucial, and any disturbance in these processes may
destabilize the catalytic action of SIRT1 for AIRE protein deacetylation. Thus, the
model system employed in this work improves the understanding of the molecular
mechanism of the G228W mutation.
320

The study of the molecular mechanisms of primary immunodeficiencies
associated with the development of autoimmune disorders has contributed to
understanding the mechanisms of tolerance loss to self-antigens. Although recessive
AIRE mutations at the PHD1 domain may occur at a ratio of 1:1000 in the general
population (Anderson et al. 2014; Su et al. 2016), AIRE SAND G228W is the first
dominant mutation associated with APS-1 (Cetani et al., 2001; Ilmarinen et al., 2005;
Wolff and Oftedal, 2019). In the context of dominant inheritance, the central question is
how these mutations influence the function of AIRE at the molecular level. Halonen et
al. (2004) and Ilmarinen et al. (2005) observed that the G228W mutation quantitatively
alters the nuclear location of the AIRE protein, impairing its transactivating property.
Furthermore, Anderson and Su (2016) demonstrated that this mutation interferes with
the localization of the AIRE protein, reducing the transcription of peripheral tissue
antigens (PTAs).
334
This study evaluated whether the G228W mutation changes the interaction with
SIRT1, an essential partner protein of the AIRE complex. We accessed the 3D structure
for the AIRE SAND domain by homology modeling to the human SP100 protein SAND
domain, which is available in a public database. We observed that the G228W
replacement, which occurs in a loop region, might not result in misfolding of the AIRE
SAND 3D structure since loop regions do not present secondary structures (Donate et
al., 1996; Bottomley et al., 2001). The wild-type and the mutant AIRE SAND domain
may not differ in their respective 3D structures, as performed by homology. This may
be due to the segment's characteristics in which the G228W mutation is present, taking
the form of a flexible loop region (SAND amino acid residues 226-228). Therefore, the
G228W mutation must involve another mechanism. Next, we asked whether the
mutation could interfere with AIRE binding to one of its partner proteins. To initiate the
downstream cascade of interactions, AIRE needs to be deacetylated, relieving RNA Pol
II stalling in the chromatin. As SIRT-1 is the only deacetylase that integrates the AIRE
complex, we next evaluated the binding of the AIRE SAND domain to SIRT1.
350
351　　　　For this purpose, the structures of the wild-type SAND domain, the G228W
352　mutant output from homology modeling, and the structure of the SIRT1 catalytic
353　domain (PDB code 4I5I) were used. With the domains modeled, we used blind docking
354　to understand how the complex between the SAND domains of AIRE WT and mutant
355　would be formed with the SIRT1 protein.
356
We observed that the SAND domain mobilizes the W228 mutant residue that
interacts with the SIRT1 H363 catalytic residue, changing with the interaction kinetics
between these proteins, followed by SPR measurements. As observed in this study, due
to the SAND G228W mutation, the AIRE protein might bind weakly to SIRT1. The
AIRE complex, including RNA Pol II, formed with an AIRE mutant protein, might be
dysfunctional and keep stalled to chromatin. The dysfunctional AIRE complex might
occur due to the non-deacetylation of K222 residue, which is essential for the proper
function of the AIRE protein. In addition, to signify a non-physiological situation, this
finding might explain the accumulation of the AIRE complex in the nucleus of mutant
cells, as previously reported (Halonen et al. 2004; Ilmarinen et al. 2005).
367
368　　　　The AIRE protein plays a role in the cell nucleus as a tetramer (Abramson et al.,
369　2010; Incani et al., 2014), in which the SAND domain of each subunit has K residues
370　SIRT1 deacetylates. SIRT1 exerts its enzymatic activity, which depends on the amino

acid residues near the acetylated K residues in the AIRE SAND domain (Garske et al., 2006). Accordingly, we hypothesized that the G228W mutation close to acetylated K residues could interfere with the SIRT1 interaction since it is an enzyme necessary to convert the acetylated to the nonacetylated and functional form of AIRE.

Although there is a prediction that the K222 residue is only moderately acetylated (Chuprin et al., 2015), previous work (Saare et al., 2012) showed that this amino acid residue is acetylated in the presence of the CBP and p300 II proteins, using mass spectrometry. Several lines of evidence stress the importance of K deacetylation in the proper interaction of AIRE with SIRT1: i) functional analyses of the G228W AIRE SAND mutation show that the subunit-containing acetylated K residue could form a hybrid tetramer unable to promote AIRE transactivation activity (Halonen et al. 2004; Ilmarinen et al. 2005), ii) AIRE mutations mimicking acetylated K243 or K253 residues in the SAND domain also reduced the transactivation activity and accumulated as nuclear bodies, whereas mutations mimicking nonacetylated K residue were functionally similar to wild-type AIRE (Saare et al. 2012), and iii) the AIRE SAND R247C mutation retains the ability to form complexes with wild-type AIRE and localize into the nucleus; however, it reduces the activation of downstream transcription through dominant-negative inhibition (Abbott et al. 2018). Notably, the R247C mutation occurs close to other K residues (K243 and K253) in the SAND domain that SIRT1 can deacetylate, also harming the transactivation function of AIRE.

This approach confirmed the hypothesis raised in this study, in which we consider that the AIRE G228W mutation changes the interaction with SIRT1, mainly when evaluated using SPR analysis. Sensor-based SPR is a label-free, sensitive, precise, and *in vitro* tool used for determining binding kinetics between biological molecules, including antigen-antibody or protein-protein interactions, among other molecular interactions (Nguyen et al., 2015; Wang et al., 2019; Soltermann et al., 2021; Ribeiro et al., 2021).

The complementary approaches used in this study (*in silico* and SPR) allowed us to evaluate the structure, energetic affinity, and association/dissociation kinetics of the complete AIRE SAND domain and peptides (wild-type and G228W mutation) with the SIRT1 protein. We observed that: i) both complete SAND domain can bind to the SIRT1 catalytic domain, ii) the binding kinetics of the G228W mutant peptide exhibited a weaker association when compared with its wild-type counterpart, and as a result, the wild-type SAND domain peptide bound faster and dissociated longer, and iii) taken together, these approaches provided evidence that the G228W mutation acts as a destabilizer for AIRE's interaction with its partner SIRT1 and offers a more accurate molecular basis for clarifying how the mutant G228W AIRE protein loses its function.

References

Abbott J. K., Huoh Y. S., Reynolds P. R., Yu L., Rewers M., Reddy M., et al. Dominant-negative loss of function arises from a second more frequent variant within the SAND domain of autoimmune regulator (AIRE). J Autoimmun. 2018; 88:114–20.

Abramson J., Giraud M., Benoist C., Mathis D. Aire's partners in the molecular control of immunological tolerance. Cell. 2010; 8;140(1):123-35.

Abramson J., Husebye E. S. Autoimmune regulator and self-tolerance – molecular and clinical aspects. Immunol. Rev. 2016; 271:127-140.

Anderson M. S., Su M. A. AIRE expands: new roles in immune tolerance and beuond. Nat. Rev. Immunol. 2016; 16: 247-258.

Alekseenko A., Ignatov M., Kozakov D. Protein-protein and protein-peptide docking with ClusPro server. Methods Mol. Biol. 2020; 2165:157–174.

Besnard M., Padonou F., Provin N., Giraud M., Guillonneau C. AIRE deficiency, from preclinical models to human APECED disease. Dis. Model. Mech. 2021; 14)2): dmm046359.

Bheda P., Jing H., Wolberger C., and Lin H. The Substrate Specificity of Sirtuins. Annual Review of Biochemistry. 2016; 85, 405–429.

Blechschmidt K., Schweiger M., Wertz K., Poulson R., Christensen H. M., Rosenthal A., et al The mouse Aire gene: comparative genomic sequencing, gene organization, and expression. Genome Res 1999; 9:158–66.

Bottomley M.J., Collard M.W., Huggenvik J.I., et al. The SAND domain structure defines a novel DNA-binding fold in transcriptional regulation. Nat Struct Biol. 2001; 8:626-33.

Case D. A. et al. AMBER. (2022), University of California, San Francisco (https://ambermd.org/AmberTools.php).

Cetani F., Barbesino G., Borsari S., Pardi E., Cianferotti E., Pinchera A., Marcocci C. A novel mutation of the autoimmune regulator gene in an Italian kindred with autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy, acting in a dominant fashion and strongly cosegregating with hypothyroid autoimmune thyroiditis J. Clin. Endocrinol. Metab. 2006; 86:4747-4752.

Chuprin A., Avin A., Goldfarb Y. et al. The deacetylase Sirt1 is an essential regulator of Aire-mediated induction of central immunological tolerance. Nat Immunol. 2015; 16:737–745.

Dai H., Case A., Riera T. et al. Crystallographic structure of a small molecule SIRT1 activator-enzyme complex. Nat Commun. 2015; 6:7645.

Donate L.E., Rufino S.D., Canard L.H., Blundell T.L. Conformational analysis and clustering of short and medium size loops connecting regular secondary structures: a database for modeling and prediction. Protein Sci. 1996 12:2600-16.

Finnish-German APECED Consortium (1997) An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PDH-type zinc-finger domains. Nat. Genet. 1997; 17(4): 399-403.

Garske A. L., Denu J. M. SIRT1 top 40 hits: use of one-bead, one-compound acetyl-peptide libraries and quantum dots to probe deacetylase specificity. Biochemistry. 2006; 10;45(1):94-101.

Glozak M. A., Sengupta N., Zhang X., Seto E. Acetylation and deacetylation of non-histone proteins. Gene. 2005; 363:15-23.

Giraud M., Peterson P. The autoimmune regulator (AIRE) gene, the master activator of self-antigen expression in the thymus. In: Thymus Transcriptome and Cell Biology. (Passos GA, Editor). 2019; Springer-Nature Switzerland AG.

Halonen M., Kangas H., Rüppell T., Ilmarinen T., Ollila J., Kolmer M., Vihinen M., Palvimo J., Saarela J., Ulmanen I., Eskelin P. APECED-causing mutations in AIRE reveal the functional domains of the protein. Hum Mutat. 2004; 23:245–257.

Humphrey, W., Dalke, A. and Schulten, K.VMD - Visual Molecular Dynamics. J. Molec. Graphics. 1996; 14.1:33-38

Ilmarinen T., Eskelin P., Halonen M., Ruppell T., Kilpikari R., Torres G. D. et al. Functional analysis of SAND mutations in AIRE supports dominant inheritance of the G228W mutation. Hum Mutat. 2005;26(4):322–331. doi: 10.1002/humu.20224.

Incani F., Serra M., Meloni A., Cossu C., Saba L., Cabras T., Messana I., Rosatelli M. C. AIRE acetylation and deacetylation: effect on protein stability and transactivation activity. J Biomed Sci. 2014; 27:21-85.

Kozakov D. The ClusPro web server for protein–protein docking. Nat Protoc. 2017; 12(2):255.

Krämer SD., Wöhrle J., Rath C., Roth G. Anabel: An Online Tool for the Real-Time Kinetic Analysis of Binding Events. Bioinform Biol Insights. 2019; 9;13:1177932218821383.

Kumar P. G., Laloraya M., Wang C. Y., Ruan Q. G., Davoodi Semiromi A., Kao K. J. et al. The autoimmune regulator (AIRE) is a DNA binding protein. J Biol Chem 2001; 276:41357–64.

Nagamine, K. et al. Positional cloning of the APECED gene. Nat. Genet., 1997; 17 (4), pp. 393-398.

Nguyen H. H., Pank J., Kang S., Kim M. Surface plasmon resonance: a versatile technique for biosensor applications. Sensors, 2015; 15(5), 10481-10510. doi: 10.3390/s150510481.

Oftedal B. E., Hellesen A., Erichsen M.M., Bratland E., Vardi A., Perheentupa J. et al. Dominant mutations in the autoimmune regulator AIRE are associated with common organ-specific autoimmune diseases. Immunity. 2015; 42:1185-1196.

Passos G. A., Speck-Hernandez C. A., Assis A. F., Mendes-da-Cruz D. A. Update on Aire and thymic negative selection. Immunology 2018; 10-20.

Perniola R., Musco G. The biophysical and biochemical properties of the autoimmune
regulator (AIRE) protein. Biochim Biophys Acta 2014; 1842:326–37.

Ramsey C., Bukrinsky A., Peltonen L. Systematic mutagenesis of the functional
domains of AIRE reveals their role in intracellular targeting. Hum Mol Genet. 2002;
11:3299–3308.

Ribeiro, J. A., Sales, M. G., Pereira, C. M. Electrochemistry-assisted surface plasmon
resonance biosensor for detection of CA 15-3. Anal Chem. 2021; 93(22): 7815-7824.
doi: 1021/acs.analchem.0c05367.

Saare M., Rebane A., Rajashekar B., Vilo J., Peterson P. Autoimmune regulator is
acetylated by transcription coactivator CBP/p300. Exp Cell Res. 2012; 15:1767-78.

Samanta U., Pal D., Chakrabarti P. Packing of aromatic rings against tryptophan
residues in proteins. Acta Crystallogr D Biol Crystallogr. 1999; 55:1421-1427.

Soltermann F., Struwe W. B., Kukura, P. Labe-free methods for optical *in vitro*
characterization of protein-protein interactions. Phys Chem Chem Phys. 2021; 23(31):
16488-16500. doi: 10.1039/d1cp01072g.

Su M. A., Giang K., Zumer K., Jiang H., Oven I. et al. Mechanisms of an autoimmunity
syndrome in mice caused by a dominant mutation in Aire. J. Clin. Invest. 2008; 118(5):
1712-1726.

Vaquero A., Scher M., Lee D., Erdjument-Bromage H., Tempst P., Reinberg D. Human
SirT1 interacts with histone H1 and promotes formation of facultative heterochromatin.
Mol. Cell. 2004; 16: 93 105.

Wang D., Loo J. F., Chen J., Yam Y., Chen S-C., He H., Kong S. K., Ho H. P. Recent
advances in surface plasmon resonance imaging sensors. Sensors 2019; 19(6): 1266.
doi: 10.3390/s19061266.

Wolff S. B., Oftedal B. E. Aire mutations and autoimmune diseases. In: Thymus
Transcriptome and Cell Biology. (Passos GA, Editor). 2019; Springer-Nature
Switzerland AG.

Acknowledgments

This work was funded by the following Brazilian research support agencies; Fundação
de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grant # 17/10780-4 to GAP
and EAD), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq,
grants # 305787/2017-9 and # 311304/2021-4 to GAP, and # 302060/2019-7 to EAD),
and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Finance
Code 001).

Contributions

JCS: raised hypothesis, discussed the analysis tools, performed molecular docking,
analyzed data, interpreted the results, and wrote the manuscript.

MD: discussed the analysis tools, performed molecular dynamics, analyzed data and
interpreted the results.
APM: performed protein binding kinetics (SPR), and interpreted the results.
VMF: performed protein binding kinetics (SPR), and interpreted the results.
RB: performed molecular dynamics, analyzed data.
EAD: raised hypothesis, interpreted the results and wrote the manuscript.
GAP: conceived the study, raised hypothesis, organized the study pipeline, interpreted
the results and wrote the manuscript.
576
577   **Competing interests**: The authors declare no competing interests.
578
579   **Figures**
580

583   **Figure 1**. Molecular modeling of the G228 (Gly228) wild-type (A and B) or mutant W228 (Trp228)
584   AIRE SAND domain (C and D) as acquired through homology modeling using the Swiss – Model with
585   the human SP100 SAND domain as a template. The G228W mutation is located in a loop region and does
586   not directly interfere with the formation of a secondary structure.

Figure 2. The blind docking output of the wild-type AIRE SAND domain shows that this domain interacts with the catalytic region of the SIRT1 protein. In A and B, the structures of the WT SAND domain (in red) and SIRT1 (in dark gray) are represented. In this representation, the AIRE SAND domain interacts with the catalytic region through its terminal loop, and the glycine 228 (Gly228) and lysine 222 (Lys222) residues are presented with their side chains. Without the Lys222 acetylated, this formation is incompatible with the native structure of the complex formed with the complete AIRE protein, in which this loop does not exist. In C and D, the same complex is represented with the surface of SIRT1 highlighted. D represents the catalytic region of SIRT1 and its catalytic residue, His363.

Figure 3. The blind docking output of the mutant AIRE SAND domain shows that this domain interacts with the catalytic region of the SIRT1 (His363 in green) protein through its mutant residue W228 (Trp228). The structures of the mutant SAND domain (in red) and SIRT1 (in dark gray) are represented in A and B. In this representation, the mutant AIRE SAND domain interacts with the catalytic region through its mutant residue W228 (Trp228), and the lysine 222 (Lys222) residues are presented with its side chain. In C and D, the same complex is represented with the surface of SIRT1 highlighted. D represents the catalytic region of SIRT1 and its catalytic residue, His363.

608 **Figure 4**. Root-mean square-deviation (RMSD) plot as a result of the molecular dynamics analysis of the
609 wild-type (red) or mutant W228 (black) AIRE SAND domain with the SIRT1 protein. The RMSD plot
610 shows that both WT and mutant SAND domains present a similar behavior along the first 600 ns. Then,
611 the WT SAND domain exhibited a slight offset from 600 to 900 ns, and both complexes tended to
612 stability at approximately 1200 ns.

Sirt1 and AIRE SAND Domain (WT)

Sirt1 and AIRE SAND Domain (Mutant)

Figure 5. Molecular dynamics structures formed between (A and B) the WT AIRE SAND domain (in red) and SIRT1 protein (in dark grey) and (C and D) between Mutant AIRE SAND domain (in red) and SIRT1 (in dark grey). (A and B) Note that the acetylated K222 residue from the AIRE SAND domain (in red) is not directed to the SIRT1 H363 catalytic residue (side chain in green) after the analyzed dynamics time (1.0 microsecond) (C and D). Considering the complex formed between the mutant AIRE SAND domain, at 1.0 microseconds of dynamics, the mutant residue W228, in the AIRE SAND domain, remained close to the catalytic H363 residue (side chain in green) of SIRT1.

Figure 6. Surface plasmon resonance sensorgrams for binding the AIRE SAND domain with SIRT1
protein immobilized onto a Biacore sensor chip. The SAND domain peptides were assayed at various
concentrations (20, 100, and 200 µM) with a contact time of 300 seconds, a 20 µl/min flow, and a
dissociation time of 1200 seconds. Note that the WT AIRE SAND peptide (Gly228) associated and
dissociated with SIRT1 throughout the analysis, while the mutant AIRE SAND peptide (Trp228)
exhibited lesser association affinity with SIRT1 when compared to the WT peptide.

Supplementary Figure 1. Surface plasmon resonance sensorgram analysis of the binding between WT or
G228W mutant AIRE SAND domain peptides with SIRT1 protein immobilized onto a Biacore sensor
chip. Six different peptide concentrations were assayed (range 10, 15, 20, 50, 100, and 200 µM) plus the
control (0 µM) for both WT and mutant peptides. Binding analysis shows a progressive increase in the
association with the WT peptide, which was not observed with the mutant peptide.

649   * END OF THE MANUSCRIPT *