Feedforward regulatory logic underlies robustness of the specification-to-differentiation transition and fidelity of terminal cell fate during *C. elegans* endoderm development

Chee Kiang Ewe[1,†], Erica M. Sommermann[1,†], Josh Kenchel[2,3], Sagen E. Flowers[1], Morris F. Maduro[4], Joel H. Rothman[1,2]*

[1] Department of MCD Biology and Neuroscience Research Institute, University of California Santa Barbara, Santa Barbara, CA 93106

[2] Program in Biomolecular Science and Engineering, University of California Santa Barbara, Santa Barbara, CA 93106

[3] Chemical and Biomolecular Engineering Department, University of California Los Angeles, Los Angeles, CA 90095

[4] Molecular, Cell and Systems Biology Department, University of California Riverside, Riverside, CA 92521

[†] These authors contributed equally to this work and are listed alphabetically.

* Correspondence: rothman@lifesci.ucsb.edu

Running title: The gene regulatory logic of endoderm development

Keywords: gene regulatory network, GATA factors, cell specification, morphological differentiation, developmental robustness, endoderm

ORCiD:

CKE - 0000-0003-1973-1308

JK - 0000-0001-6378-6723

SEF - 0000-0002-7818-2188

MFM - 0000-0001-6257-7612

JHR - 0000-0002-6844-1377

Graphic abstract

Progressive cell fate lockdown | Robust developmental outcome

Specification-to-differentiation transition

Abstract

Development is driven by gene regulatory networks (GRNs) that progressively dictate

specification and differentiation of cell fates. The architecture of GRNs directly determines the

specificity and accuracy of developmental outcomes. We report here that the core regulatory

circuitry for endoderm development in *C. elegans* is comprised of a recursive series of

interlocked feedforward modules linking a cascade of six sequentially expressed GATA-type

transcription factors. This structure results in a reiterated sequential redundancy, in which

removal of a single factor or alternate factors in the cascade results in no, or a mild, effect on

endoderm development and gut differentiation, while elimination of any two factors that are

sequentially deployed in the cascade invariably results in a strong phenotype. The strength of

the observed phenotypes is successfully predicted by a computational model based on the

timing and levels of transcriptional states. The feedforward regulatory logic in the GRN appears

to ensure timely onset of terminal differentiation genes and allows rapid and robust lockdown of

cell fate during early embryogenesis**.** We further found that specification-to-differentiation

transition is linked through a common regulator, the END-1 GATA factor that straddles the two

processes. Finally, we revealed roles for key GATA factors in establishing spatial regulatory

state domains by acting as transcriptional repressors that appear to define the boundaries of the

digestive tract. Our findings support a comprehensive model of the core gene network that

describes how robust endoderm development is achieved during *C. elegans* embryogenesis.

Introduction

Development is driven by progressive activation of transcriptional programs, or gene

regulatory networks (GRNs) that direct cell specification and subsequent differentiation (Boveri,

1899; Davidson and Levine, 2008). The sequential restriction of cell identity and developmental

potential through dynamic changes in transcriptional states was anticipated by Waddington's

epigenetic landscape, a graphic metaphor describing canalization and robustness during

development (Waddington, 1957).

The generation of diverse animal forms largely relies on a common genetic toolkit. The

GATA transcription factors play a conserved role in the development of diverse cell types,

including those of the endoderm, the first of the three germ layers to have evolved during the

late Precambrian era (Hashimshony et al., 2015; Rodaway and Patient, 2001). In the

diploblastic phyla Poriferans and Cnidarians, GATA factors have been found to be specifically

expressed in the endoderm, suggesting that these transcriptional regulators may have driven

the invention of the endoderm germ layer and gastrulation at the dawn of metazoan evolution

(Martindale et al., 2004; Nakanishi et al., 2014). This association of GATA factors with

endoderm development persists throughout metazoan phylogeny, generally through successive

deployment of multiple GATA factors. This sequential use of GATA factors is particularly striking

in *C. elegans*, in which a cascade of six GATA factor-like transcription factors regulate

specification and differentiation of the endoderm (Maduro, 2017; Maduro and Rothman, 2002;

McGhee, 2007). In *Drosophila*, the GATA factor Serpent specifies endodermal fate and

activates the expression of a second GATA factor, dGATAe, which is essential for the terminal

differentiation of the intestine. These factors can act across wide phylogenetic spans, as first

demonstrated with the *C. elegans* END-1 GATA factor, which ectopically activates endoderm

development when expressed in the prospective ectoderm of *Xenopus* (Shoichet et al., 2000).

Similarly, overexpression of *serpent* or *dGATAe* causes ectopic endoderm differentiation in non-

endodermal lineages in *Drosophila*, as well as in *Xenopus*, further supporting the functional

conservation of the GATA factors (Murakami et al., 2005; Okumura et al., 2005). In sea urchin,

Blimp1/Krox1 activates *otx1*, the product of which activates *gatae* expression (Davidson et al.,

2002). Gatae in turn provides a positive input to *otx1*, in addition to activating the transcriptional

program for endoderm development, thereby forming a stable circuit in the GRN (Davidson et

al., 2002; Yuh et al., 2004). Accordingly, knocking down *gatae* severely blocks endoderm

development and gastrulation (Davidson et al., 2002).

The endoderm in *C. elegans*, which arises from a single blastomere born at the 8-cell

stage, the E cell (Boveri, 1899; Sulston et al., 1983), provides a highly tractable system for

investigating the mechanisms of cell specification, differentiation, and organogenesis. This

progenitor cell gives rise to a clone 20 cells comprising the intestine, which are arranged in nine

rings (int1-9) spanning the length of the animal (Supplementary Figure 1A). Endoderm

development is driven by pairs of duplicated genes encoding GATA-like transcription factors:

the divergent MED-1/2 factors and the canonical END-1/3 and ELT-2/7 factors. The maternally

provided SKN-1/Nrf transcription factor activates MED-1 and -2, which activate the specification

of mesendodermal fate in the EMS blastomere (Bowerman et al., 1992). In the anterior daughter

of EMS, the MS cell, the Wnt effector POP-1/Tcf represses the expression of *end-1/3*, and

MED-1 and -2 activate *tbx-35*, whose product specifies mesodermal fate. In the E cell, the

posterior daughter of EMS, a triply redundant signaling system (Wnt, Src, and MAPK) leads to

the phosphorylation of POP-1 by nemo-like kinase LIT-1 (Bei et al., 2002; Maduro et al., 2002;

Meneghini et al., 1999; Shin et al., 1999; Thorpe et al., 1997). Together with MED-1/2, Wnt-

signaled POP-1 activates genes encoding the transiently expressed endoderm specification

factors END-1 and -3, which in turn activate the expression of *elt-7* and *-2*, orthologues of

vertebrate *GATA4/5/6*. Expression of the ELT factors is sustained throughout the remainder of

the animal's life via a positive autoregulatory loop that "locks down" the differentiated state of

the intestine (Supplemental Figure 1B) (Maduro, 2017; Maduro and Rothman, 2002; McGhee,

2007). While *elt-7* loss-of-function mutants do not show a discernible phenotype, animals

lacking ELT-2 arrest at early larvae stage (L1) owing to severely obstructed gut (Sommermann

et al., 2010). Nonetheless, *elt-2(-)* mutant animals contain a well-defined intestinal lumen and

the intestinal cells appear to be fully differentiated (Fukushige et al., 1998; Sommermann et al.,

2010). In the absence of both ELT-2 and -7, however, the intestinal lumen is completely

abolished and differentiation appears to proceed in only a subset of the endoderm-derived cells

in a sporadic manner (Sommermann et al., 2010). This suggests that ELT-2 and -7 function

synergistically in mediating morphological differentiation of the intestine, and additional input(s)

mediate a bistable switch in the endodermal differentiation program in the absence of ELT-2

and -7 (Dineen et al., 2018; Sommermann et al., 2010).

In this study, we sought to decipher the functional requirements and interactions

between the GATA transcription factors and how they allow rapid and faithful deployment of the

endoderm GRN. We found that the endoderm GRN consists of a series of interlocking

feedforward loops, creating "sequential redundancy" in the cascade and culminating in the rapid

lockdown of cell fate. We further report that END-1 acts at a transition point, participating in both

specification and differentiation. Finally, we demonstrated the important roles of GATA factors in

safeguarding intestinal cell fate and defining the boundaries of the digestive tract. Overall, our

findings reveal the nature of the extensive genetic redundancy in the regulatory circuitry and

how this GRN architecture dictates robust cell specification and differentiation during embryonic

development.

Methods and Materials

C. elegans cultivation and genetics

Worm strains were cultured using standard procedure (Brenner, 1974) and all experiments were performed at room temperature (20-23^{O}C). All genetic manipulations were performed according to standard techniques (Fay, 2013). *him-5(-)* or *him-8(-)* was introduced into some strains to generate males and facilitate crosses. See Supplementary Table 1 for a complete list of strains used in this study.

Immunofluorescence analysis

Antibody staining with methanol-acetone fixation was performed as previously described (Sommermann et al., 2010). Antibodies MH27 (AB_531819), MH33, and 455-2A4 (AB_2618114) were used to detect AJM-1, IFB-2, and ELT-2, respectively. Alexa Fluor® 488 goat anti-mouse secondary antibody was used at 1:1000 dilution.

RNAi

RNAi feeding clones were obtained from the Ahringer (Kamath et al., 2003) or the Vidal (Rual et al., 2004) library. The bacterial strain was inoculated overnight at 37^{O}C in LB containing 50 µg/ml ampicillin. The bacterial culture was then diluted 1:10 and incubated for an additional 4 h. Next, 1 mM of IPTG was added to the bacterial culture and 100 µL was seeded onto 35 mm NGM agar plates containing 1 mM IPTG and 25 µg/mL carbenicillin. For simultaneous knockdown of *elt-2* and *elt-7*, the two bacterial strains, each expressing dsRNA for one gene, were concentrated and resuspended in 1 mL of LB in 1:1 ratio before seeding the NGM plates. Seeded plates were allowed to dry for 48 h before use. Next, 10-20 L4 animals were placed on the RNAi plates. 24 h later, the animals were transferred to fresh RNAi plates to lay eggs. The progeny was then collected for analyses.

Imaging and fluorescence quantification

The animals were immobilized using 10 mM Levamisole and mounted on 4% agarose pads. Images were acquired, typically at 60X, using Nikon Eclipse Ti-E inverted microscope fitted with ORCA-Flash2.8 camera. For expression studies, maximum intensity Z-projection was generated on the Nikon NIS-Elements AR v4.13.05. Images were then analyzed using ImageJ or Imaris v9.7.2.

RT-qPCR

RNA was extracted from synchronized L1 animals using Monarch® Total RNA Miniprep Kit (#TS010S). cDNA synthesis was performed using SuperScript™ III First-Strand Synthesis SuperMix (Thermo Fisher Scientific, #18080400). Quantitative PCR was performed using BioRad CFX96 Real-Time System. Each 15 µL reaction contained cDNA, primers, and PowerUp™ SYBR™ Green Master Mix (Thermo Fisher Scientific, #A25743). The data were analyzed using the standard $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001).

The primers sequences are:

act-1 - 5'-TCCATTGTCGGAAGACCACG-3' and 5'-GGTGACGATACCGTGCTCAA-3'

act-5 - 5'-GTCACTCACACCGTTCCAATC-3' and 5'-GTGAGGATCTTCATCATGTAGTCG-3'

Modeling endoderm gene regulatory circuits

The topology of the gene circuits with temporal information was written as a system of differential equations, with expression of each factor dependent on the concentration of its activators (See Supplementary Materials and Supplementary File 1). The gene cascade is initiated by SKN-1, which was modelled as a square wave in the EMS blastomere. Similarly, the positive inputs of (phosphorylated) POP-1 into *end-3* and *end-1* were modelled as a square wave in the E blastomere (23 mins after the four-cell stage). Model runs were calculated as

time-discretized Euler approximations (Hahn, 1991) with time steps of 0.01 s. An iterative least-

squares algorithm following a modified Gauss-Newton method (Ruhe, 1979; Yip et al., 2010)

was used to fit the model parameters to published transcriptomics data (Baugh et al., 2003;

Tintori et al., 2016). The performance of the final model was then evaluated by comparing the

predicted phenotypes of the single mutants with published results (Boeck et al., 2011; Dineen et

al., 2018; Maduro et al., 2005a; Maduro et al., 2015). Finally, predictions of *elt-2* activation (a

readout for the commitment to E fate) in the mutant combinations were generated by holding the

concentration of knockout genes at zero and otherwise running the model as described. The

source code is available on https://github.com/RothmanLabCode/endoderm_GRN_model.

Statistics and figure preparation

Statistics were performed using R software v3.4.1 (https://www.r-project.org/). The specific

statistical tests were reported in the figure legends. Plots were generated using R package

ggplot2 or Microsoft Excel. Figures were assembled in Inkscape v0.92.4 (https://inkscape.org/).

Results

Sequential redundancy suggests feedforward regulatory circuitry in the endoderm GRN

Owing to the extensive genetic redundancy, single mutants of most of the genes
throughout the *C. elegans* endoderm GRN either show no overt phenotype (single *med* mutants,
end-1(-), *elt-7(-)*) or an extremely weakly penetrant phenotype (*end-3(-)*). Moreover, while the
elt-2(-) mutant undergoes larval arrest immediately after hatching with a dysfunctional gut, the
gut appears fully differentiated in these mutants. A strong gut differentiation defect requires the
removal of both *elt-2* and *elt-7* (Figure 1A) (Fukushige et al., 1998; Sommermann et al., 2010).
Similarly, strong phenotypes are observed only when both *meds* or both *ends* are removed in
pairs. In this study, we sought to examine the basis for this extensive redundancy in the
pathway and to illuminate how it might contribute to faithful specification and differentiation in
endoderm GRN.

Temporally resolved transcriptomic analyses revealed that the GATA transcription
factors are sequentially activated in the mesendoderm regulatory pathway (Baugh et al., 2003;
Tintori et al., 2016). This observation suggests that, rather than pairs of factors (MEDs, ENDs,
ELTs) acting together at specific tiers in the cascade (Supplementary Figure 1B), each factor is
redundant with its immediate upstream or downstream factor in a continuously sequential
cascade ("sequential redundancy"). For example, as is often seen in forward-driven biological
switches, each may act through a feedforward motif in which a given factor activates both its
immediate target gene and the target of that gene (Mangan and Alon, 2003).

To test the model that endoderm differentiation is directed by a series of sequentially
redundant feedforward regulatory motifs, we constructed a series of double mutants and
compared the penetrance of lethality and the extent of gut differentiation. We found that unlike
the mild or undetectable phenotypes observed with single mutants, removing pairs of genes that

are expressed at sequential steps in the cascade invariably results in severely diminished

viability (Figure 1A) and defects in endoderm specification/differentiation, as revealed by

disrupted gut lumen morphology (Figure 1A, B). This effect is associated with sporadic

expression and in some cases the complete absence of gut-specific rhabditin granules (Clokey

and Jacobson, 1986) (Figure 1C). We found that the expression of the gut-specific IFB

(Intermediate Filament, B)-2 protein is strongly diminished in the double mutants

(Supplementary Figure 2A). Moreover, reduced AJM (Apical Junction Molecule)-1 expression

suggests gut epithelialization defects in double mutants of sequential members in the endoderm

GRN (Supplementary Figure 2B).

In *med-2(-); med-1(-)* double mutants, all embryos die as embryos as MS blastomere

fails to be specified and its fate is transformed to that of the C-like mesectodermal progenitor in

the early embryo (Figure 1A), as is also the case in embryos lacking the initiating, maternally

provided SKN-1 transcription factor (Bowerman et al., 1992). The E cell similarly adopt a C cell

fate in some, but not all, of the *med-2(-); med-1(-)* arrested embryos (Maduro et al., 2007). As

previously reported (Maduro et al., 2005a), many *end-1(-) end-3(-)* mutants also die as embryos

owing to transformation of the E cell into a C-like cell (Figure 1A). Thus, END-1 and -3 act as

redundant selector genes that promote specification of endoderm over other cell fates.

Consistent with the sequential redundancy model, removal of two genes that function at

alternate steps in the endoderm cascade does not result in the strong synergistic effect that we

observed with removal of pairs of sequentially expressed genes (Figure 1A, E, F;

Supplementary Figure 3). We observed normal expression of IFB-2 and AJM-1 along the length

of the gut region in all such double mutants (Supplementary Figure 3). Further, while the *end-3(-*

); med-1(-) double mutant, in which sequential genes in the cascade are removed, shows

severe embryonic lethality, most (89.5%; n = 219) of the *end-1(-); med-1(-)* double mutants, in

which alternate genes in the cascade are removed, develop into fertile adults (Figure 1A). This

observation is also consistent with the reported divergent roles of END-1 and -3 in endoderm

specification (Boeck et al., 2011). Moreover, the *elt-7(-) end-3(-)* double mutants are largely

viable, in contrast to the *end-3(-); med-1(-)* sequential double mutants, which show a strongly

penetrant embryonic lethal phenotype (Figure 1A). Of the hatched L1 larvae, 31% of *end-3(-);*

med-1(-) sequential double mutant animals exhibit no overt signs of gut differentiation, while

only 2% of *elt-7(-) end-3(-)* alternate double mutants completely lack gut (Fisher's exact test; p <

0.001) (Figure 1D, G). However, many *elt-7(-) end-3(-)* mutants contain a partially differentiated

gut (Figure 1G). We reasoned that the developmental defects observed in the most affected

animals are the result of suboptimal expression of *end-1* in the absence of *end-3* (Maduro et al.,

2007).

While neither single mutant shows a discernible phenotype, animals lacking both END-1

and ELT-7 are 100% inviable, and the arrested L1 larvae show a striking gut differentiation

defect (Figure 1A, Supplementary Figure 2). The mutant worms contain patches of apparently

differentiated gut as evidenced by expression of gut granules and immunoreactive IFB-2, similar

to *elt-7(-); elt-2(-)* double mutants, which exhibit an all-or-none block to differentiation event

along the length of the animals (Figure 1B, C; Supplementary Figure 2) (Sommermann et al.,

2010). However, the differentiation defect observed in *elt-7(-) end-1(-)* double mutants appears

to be somewhat milder from that in *elt-7(-); elt-2(-)*: unlike the latter, we observed defined, albeit

sporadic, lumen and brush border, and the undifferentiated patches are more frequently

interspersed with differentiated patches (Figure 1B, C; Supplementary Figure 2A) (more below).

Analysis of most alternate and sequential double mutant combinations show similar

effects (Fig. 1A): sequential double mutants are invariably much more severely affected than

alternate double mutants. Together, our data support a sequentially redundant cascade,

comprising a recursive series of feedforward regulatory steps (Figure 1H). It is conceivable that

such a feedforward system creates a strongly forward-driven, rapidly deployed switch that

ensures timely and robust cell fate commitment and robust lockdown of endoderm cell fate

during embryogenesis.

Computational model predicts phenotypes of sequential and alternative double mutants

We took a complementary approach to testing the sequential feedforward model through

a computational strategy. We constructed a mathematical model based on the network topology,

in which the interactions between the GATA factors, as well as the additional POP-1-dependent

activation of *end-3* and *end-1*, were written as a series of ordinary differential equations and the

model parameters (Supplementary File 1) were determined by fitting to published

transcriptomics data (Baugh et al., 2003; Tintori et al., 2016) using a custom algorithm that

follows an iterative least-squares method. We then carried out *in silico* perturbations of the

feedforward circuits (see Methods and Materials and Supplementary Materials) to provide

predictions of these effects on the relative timing and levels of *elt-2* activation and therefore the

final output of the endoderm GRN. Our computed results revealed that *elt-2* expression is

predicted to occur, but with delayed onset in all single mutants and in mutants lacking alternate

pairs of GATA factors (Figure 1I; Supplementary File 1). However, as observed with the

experimental outcomes, *elt-2* expression is predicted to be completely abrogated in most double

mutant combinations in which sequential members of the endoderm cascade are removed

(Figure 1I), consistent with their pronounced developmental defects (Figure 1A). The predicted

elt-2 expression level strikingly correlates (Spearman's Rank Correlation ρ = 0.96; p < 0.001)

with the experimentally observed penetrance of the phenotypes of the single and multiple

mutants (Figure 1I): single and alternate double mutant combinations, which show no, or weak

phenotypes, are predicted to express high levels of *elt-2* early, while sequential double mutants,

which show strong defects in gut development and inviability, are predicted not to express *elt-2*

at substantial levels at its normal time of onset. These findings, based on modeling with gene

expression data, bolster the proposed recursive feedforward structure of the GATA factor

cascade.

Figure 1: Evidence for recursive feedforward loops in the endoderm GRN.

(A) Comparison of viability and gut lumen morphology in mutant combinations lacking one or
two GATA factors in the endoderm regulatory cascade. Mutants missing sequential members of
the endoderm cascade exhibit more penetrant developmental defects than mutants with
alternate steps disrupted. We note the two loss-of-function alleles of *end-1*, *ox134* (14822 bp
deletion; also removes the adjacent *ric-7* gene) and *ok558* (879 bp deletion), do not result in
significant loss in viability (Fisher's exact test; p = 0.31) or other discernible phenotypes. We
used *ox134* in subsequent analyses unless stated otherwise (see Supplementary Table 1). The
total number of animals scored for each genotype is indicated. No L1 animals were scored for
med-2(-); med-1(-) (indicated by *) as all double mutants arrest as embryos owing to
misspecification of MS (and E in majority but not all of the embryos). The frequencies of (B, E)
visible lumen and (C, F) gut granules are dramatically reduced in double mutants missing (B, C)
sequential GATA pairs compared to those lacking (E, F) alternate members of the endoderm
cascade. (D, G) *end-3(-); med-1(-)* shows a more severe defect in intestinal differentiation than
elt-7(-) end-3(-). 31% of *end-3(-); med-1(-)* and 2% of *elt-7(-) end-3(-)* show no overt signs of
endoderm differentiation. (G) Many *elt-7(-) end-3(-)* larvae contain a partial gut as observed by
DIC microscopy and birefringent gut granules, presumably due to sub-threshold levels of END-
1-dependent activation of *elt-2*. Scale bar = 10 μm. (H) The sequential redundancy model of the
endoderm GRN. (I) Computed *elt-2* expression levels predict the phenotype severity of the
mutant combinations. All *med-2(-); med-1(-)* mutants show severe embryonic lethality despite
elt-2 activation owing to a fully penetrant MS→C misspecification. We excluded *med-2(-); med-
1(-)* from this analysis.
304

Variation in temporal expression explains distinct functions for MED-1 and -2

It has been shown previously that *end-3* expression is activated slightly earlier than *end-1* (Supplementary Figure 4) (Maduro et al., 2007; Tintori et al., 2016), as is also consistent with our genetic analyses (Figure 1). Additionally, the END paralogues have diverged considerably with variable DNA-binding domains that perform overlapping but distinct functions (Boeck et al., 2011). Unlike the ENDs, MED-1 and -2 are 98% identical (Maduro et al., 2001). Nonetheless, our genetic evidence demonstrates distinguishable contributions of the two nearly identical paralogues, with MED-2 functions preceding MED-1 (Figure 1; Supplementary Figure 2, 3). By examining published lineage-resolved single-cell RNA-seq data (Tintori et al., 2016), we observed that *med-2* transcripts are undetectable by the 8-cell stage, while *med-1* expression persists briefly in the endoderm precursors (Supplementary Figure 4A). The differential temporal expression of the MEDs was recapitulated using protein-fusion reporters: unlike MED-1, which is expressed in 16E embryos, MED-2 protein expression is largely diminished by the 8E embryonic stage (Supplementary Figure 4B). Together, our data suggest *med-1* and *-2* genes are differentially regulated and MED-2 functions upstream of MED-1, further supporting the feedforward structure at the beginning of the endoderm regulatory cascade.

Synergistic requirements and cross-regulatory interactions of END-1, ELT-7, and ELT-2

As described above, while both *elt-7(-)* and *elt-2(-)* single mutants contain a fully differentiated gut with a contiguous lumen from the pharynx to the rectum surrounded by cells of normal differentiated morphology, *elt-7(-); elt-2(-)* double mutants invariably lack both a defined gut lumen as well as some gut cells, and show a sporadic, all-or-none, block to gut differentiation along the length of the animals (Figure 1A-C; Figure 2A; Supplementary Figure 2, 5) (Sommermann et al., 2010). Although differentiation is highly defective in the absence of ELT-2 and -7, patches of well-differentiated gut are nonetheless evident. Moreover, many terminal differentiation genes remain activated in the absence of ELT-2 and -7 (Dineen et al.,

2018). For example, eliminating the functions of ELT-2 or -7 has little effect on the expression of

act-5, a gene encoding an actin isoform required for microvilli formation (Dineen et al., 2018;

MacQueen et al., 2005). These observations suggest that at least one additional factor, in

addition to the ELTs, may activate gut differentiation. One such candidate is END-1, which acts

in specification of E cell identity immediately upstream of the *elt* genes. Indeed, although *end-1(-)*

and *elt-7(-)* mutants are both phenotypically silent, we found that the *elt-7(-) end-1(-)* double

mutant shows extensive gut differentiation defects, with sporadic expression of rhabditin

granules (Figure 1A-C; Figure 2B) and *ifb-2* (Figure 2C, D; Supplementary Figure 2), as well as

reduced number of differentiated gut cells (Figure 2E; Supplementary Figure 6), reminiscent of

elt-7(-); elt-2(-) double mutant animals.

We further tested whether END-1 could account for the residual gut-promoting activity by

removing it from animals also lacking ELT-2/7. Strikingly, simultaneously eliminating END-1,

ELT-7, and ELT-2 results in an apparent elimination of intestinal differentiation (Figure 2F-I). We

found that 19.6% of *elt-7(-) end-1(-); elt-2(-)* mutants undergo embryonic arrest, and the

remainder die as L1 larvae (n = 143). The triple mutant animals exhibit no morphological signs

of gut differentiation, showing no gut granules (Figure 2F, G, H), no detectable intestinal brush

border or lumen (Figure 2F, G, I), and no differentiated gut nuclei expressing a gut-specific

peptidase transporter, OPT-2 (Figure 2J; Supplementary Figure 7), suggesting a near total

block to gut differentiation. We found that knocking out *end-1* strongly reduces the expression

level of *act-5* (Figure 2K, L), and *act-5* expression is further downregulated in *elt-7(-) end-1(-)*

and *elt-7(-) end-1(-); elt-2(RNAi)* animals (Figure 2K; Supplementary Figure 8), suggesting that

END-1, ELT-7 and ELT-2 act collaboratively and redundantly to mediate *act-5* expression, and

that END-1 may compensate for the loss of ELT-2 and -7 inputs.

A challenge to the notion of a role for END-1 as the gut-promoting factor acting in the

absence of the ELTs is that its expression in wildtype embryos is transient, such that its product

is largely undetectable by the 16E embryonic stage (Li et al., 2019; Zhu et al., 1997); yet gut

differentiation seen in the differentiated patches in the *elt-7(-); elt-2(-)* double mutant appears

strong and robust late in development. These findings led us to hypothesize that ELT-2 and/or

ELT-7 may normally repress *end-1* transcription through negative feedback and that in the

absence of the ELTs, *end-1* may be upregulated and drive differentiation. Indeed, we found that

the intensity of an *end-1* endogenous protein fusion reporter is modestly elevated in 8E embryos

when *elt-2* is knocked down by RNAi (Figure 3A). Moreover, the expression level of *end-1* is

upregulated in both 4E and 8E embryos when ELT-7 and -2 are simultaneously depleted

(Figure 3B). While these findings provide evidence for the hypothesized feedback inhibition, the

effect is rather weak, as we did not observe obvious longer-term perdurance of *end-1*

expression (see Discussion).

366   Taken together, our results suggest END-1 is poised in the cascade at the interface

367   between specification and differentiation, placing it at the crux of this key transition. END-1,

368   acting with END-3, regulates *specification* of the E lineage, whereas END-1, acting with ELT-7

369   and ELT-2, controls *differentiation* of the intestine.

Figure 2: Synergistic actions of END-1, ELT-7 and ELT-2 mediate morphological differentiation of endoderm.

(A, B) *elt-7(-) end-1(-)* contains a grossly defective gut with sporadic patches of rhabditin granules interspersed with apparently undifferentiated regions, similar to that observed in *elt-7(-); elt-2(-)*. Scale bars = 20 µm. (C, D) Balanced *elt-7(-/-) end-1(-/+)* larva shows uniform expression of IFB-2 (*kcIs6* transgene) along the length of the animal, while *elt-7(-/-) end-1(-/-)* shows sporadic expression of the transgene. Scale bars = 10 µm. (E) Wildtype and *elt-7(-/-) end-1(-/+)* animals contain an average of 20 intestinal cells, marked by *elt-2*::GFP reporter. The number of differentiated gut cells is markedly reduced in *elt-7(-/-) end-1(-/-)* larvae (mean = 13.5

cells). (F, G) Representative micrographs of *elt-7(-) end-1(+/-); elt-2(-)* and *elt-7(-) end-1(-/-); elt-*

2(-) triple mutants. (F, F') *elt-7(-) end-1(+/-); elt-2(-)* lacks a visible lumen and contains sporadic

birefringent granules. (G, G') *elt-7(-) end-1(-/-); elt-2(-)* exhibits absolutely no signs of endoderm

differentiation. (G") Magnified view of the animal present in (G). Scale bars = 10 µm. (H, I) The

average frequencies of (H) gut granule and (I) lumen are dramatically reduced in *elt-7(-) end-1(-*

/-); elt-2(-) compared to *elt-7(-) end-1(+/-); elt-2(-)* animals. (J) Wildtype and *elt-2(-)* show similar

number of differentiated intestinal cells, although the variance in *elt-2(-)* is significantly increased

(F-test p < 0.001). The number of *opt-2p::mCherry* (*irSi24*) expressing cells is strongly reduced

in *elt-2(-); elt-7(RNAi)*. No gut cells were detected in vast majority of *elt-7(-) end-1(-); elt-2(-)*

triple mutants. (K) The expression of *act-5::GFP* translational reporter *(jyIs13)* in various mutant

combinations. END-1, ELT-7, and ELT-2 appear to act together to regulate *act-5* expression in

the intestine. (L) *act-5* is downregulated in *end-1(-)* compared to wildtype as detected by RT-

qPCR. *act-1* was used as the internal reference. Three replicates were performed for each

genotype. Error bars represent standard deviation. * p ≤ 0.05 by two-tail t-test. For panels E and

J, NS p > 0.05, ** p ≤ 0.01, *** p ≤ 0.001 by non-parametric Kruskal-Wallis test and pairwise

Wilcoxon tests with Benjamini & Hochberg correction. For panel K, NS p > 0.05, * p ≤ 0.05, ** p

≤ 0.01, *** p ≤ 0.001 by parametric one-way ANOVA followed by pairwise t-tests with Benjamini

& Hochberg correction.

Figure 3: ELT-2 antagonizes *end-1* expression.

(A) Expression of endogenously tagged *end-1* increases in 8E embryos upon *elt-2* RNAi treatment. NS p > 0.05, *** p ≤ 0.001 by Kruskal-Wallis test and pairwise Wilcoxon tests with Benjamini & Hochberg correction. (B) Knocking down both *elt-7* and *elt-2* further elevates END-1 expression in both 4E and 8E embryos. *** p ≤ 0.001 by Wilcoxon tests.

ELT-2 and ELT-7 collaborate to safeguard intestinal cell fate

When improperly specified, E has been shown to undergo wholesale lineage conversion

to either a C-like mesectodermal fate (when SKN-1, the MEDs, or END-1/3 are removed),

thereby inappropriately generating epidermis, or an MS-like mesodermal fate (for example, in

the absence of Wnt signaling), thereby inappropriately giving rise to pharyngeal tissue (details

described in Figure 4A). Given that END-1 straddles the transition from specification of the

endoderm progenitor and differentiation of the gut, we sought to investigate whether

specification and differentiation involve distinct regulatory events by examining non-endodermal

(epidermal and pharyngeal) gene activity when differentiation is impaired after this transition is

thought to occur. In *end-1(-) end-3(-)* double mutants, misspecification of the E cell and its

conversion to a C-like mesectodermal progenitor, results in severe morphological defects as a

result of the supernumerary epidermal cells and consequent deformation of the epidermis

(Figure 4B; Supplementary Figure 9) (Maduro et al., 2005a). In contrast, the *elt-7(-) end-1(-); elt-*

2(-) triple mutant animals that, like the *end-1(-) end-3(-)* double mutants, do not make a

discernible gut, are not substantially defective for overall body morphogenesis (Supplementary

Figure 9), suggesting that E is initially specified. Consistent with this observation, unlike *end-1(-)*

end-3(-) double mutant animals, *elt-7(-) end-1(-); elt-2(-)* triple mutant larvae contain a wildtype

number of epidermal cells, implying that E→C misspecification does not occur (Figure 4B).

These results imply that END-3 alone is sufficient to promote endoderm specification over C-like

mesectodermal cell fate. However, we were surprised to observe mis-expression of the

pharyngeal muscle-specific myosin gene, *myo-2*, in the gut region of many *elt-7(-); elt-2(-)* and

elt-7(-) end-1(-); elt-2(-) mutants (Figure 4C, D). Additionally, knocking down *elt-7* in *end-1(-);*

elt-2(-) results in ectopic expression of *ceh-22*, which encodes a pharynx-specific NK-2-type

homeodomain protein, in the otherwise undifferentiated gut (Supplementary Figure 10). As the

Wnt/POP-1-dependent polarization of EMS is unperturbed in the mutants, we reasoned that

inappropriate expression of pharyngeal genes is unlikely to be the result of wholesale E→MS

transformation, but may reflect later errors in the fidelity of differentiation.

PHA-4/FoxA is the organ selector gene that specifies pharyngeal identity, regulating

myo-2 and *ceh-22* among thousands of other targets in the pharynx (Mango et al., 1994; Zhong

et al., 2010). We found that the ectopic expression of *myo-2* in *elt-7(-); elt-2(-)* is at least partially

suppressed in *pha-4(RNAi)* animals (Figure 4E). In wildtype animals, PHA-4 is expressed at

high levels in the pharynx and rectum and at low levels in the intestine (Figure 4F, F') (Horner et

al., 1998; Kalb et al., 1998). It was previously found that ELT-2 positively regulates *pha-4* as

forced expression of *elt-2* causes widespread activation of *pha-4* (Kalb et al., 1998).

Paradoxically, we found that *pha-4* is upregulated in the intestine in *elt-2(-)* animals (Figure 4G,

G'), and depleting *elt-7* in *elt-2(-)* animals further enhances this effect (Figure 4H, H'). These

findings raise the possibility that ELT-2 serves dual roles as both an activator and a repressor

depending on its expression level. Thus, it appears that upregulation of *pha-4* in the midgut in

the absence of ELT-2 and -7 activates sporadic ectopic pharyngeal gene activity. Supporting

our model, PHA-4 target genes have been shown to be regulated in part by PHA-4 binding

affinity and occupancy (Fakhouri et al., 2010; Gaudet et al., 2004). Taken together, our results

show that the boundaries of regulator state domains along the digestive tract of *C. elegans* are

established, at least partly, by transcriptional repression mediated by ELT-2 and -7 in the

intestine (Figure 4I).

Figure 4: ELT-2 and ELT-7 repress pharyngeal fate in the intestine.

(A) E/C or E/MS binary fate choice during early embryogenesis. The Caudal homolog PAL-1 is required for the specification of mesectodermal C blastomere, which gives rise to epidermis and body wall muscles (Hunter and Kenyon, 1996). Maternally provided *pal-1* is specifically translated in the EMS and P_2 cells. In MS and E, PAL-1 activity is blocked by TBX-35 and END-1/3, respectively (Broitman-Maduro et al., 2006). Thus, depleting TBX-35, END-1/3, or their upstream activators, MED-1/2 and SKN-1, causes excess skin and muscle owing to the misspecification of MS and/or E as C, the somatic descendant of P_2 (Hunter and Kenyon, 1996). As EMS divides, Wnt, MAPK, and Src signaling from P_2 polarizes EMS, leading to the phosphorylation and nucleocytoplasmic redistribution of POP-1, and activation of *end-1/3* in E, but not MS. When the polarizing signal from P_2 is disrupted, POP-1 is unphosphorylated, and MED-1 and -2, instead, activate the development of mesoderm (MS), which gives rise to the posterior pharynx and body wall muscles (Maduro and Rothman, 2002; Maduro et al., 2002;

Rocheleau et al., 1999; Shin et al., 1999). (B) Wildtype and *end-1(-); elt-2(-); elt-7(RNAi)* larvae

contain ~73 epidermal cells, while *end-1(-) end-3(-)* larvae contain ~80 epidermal cells marked

by *dpy-7p::GFP* expression. NS p > 0.05, *** p ≤ 0.001 by parametric one-way ANOVA followed

by pairwise t-tests with Benjamini & Hochberg correction. (C) Representative DIC and

fluorescent micrographs showing different mutation combinations expressing *myo-2p::GFP*.

Ectopic expression of *myo-2* is evident in *elt-7(-); elt-2(-)* and *elt-7(-) end-1(-); elt-2(-)* mutants

(arrowheads). (D) The frequency of animals showing mis-expression of *myo-2* as shown in (C).

Number of animals scored for each genotype is indicated. (E) Knocking down *pha-4* partially

rescues ectopic expression of *myo-2* in *elt-7(-); elt-2(-)* animals. * p ≤ 0.05 by Fisher's exact

test. (F-H) The expression of endogenously tagged *pha-4* reporter in (F, F') wildtype, (G, G') *elt-*

2(-), and (H, H') *elt-2(-); elt-7(RNAi)* worms. The white horizontal lines in (F'-H') mark the

posterior end of the pharynx. Exposure time = 195 ms. All scale bars = 10 µm. (I) Model of

spatial repression and fate exclusion in the digestive tract. ELT-2 and -7 repress *pha-4* and

prevent the expression pharyngeal genes in the midgut.

END-1 and ELT-7 establish the boundary between the valve and intestinal tubes

The foregoing results suggest that the core regulators involved in gut differentiation (END-1, ELT-2, and ELT-7) regulate faithful differentiation of cells in the digestive tract. The pharynx and the intestine are linked by the pharyngeal-intestinal valve (vpi), which consists of six cells arranged into three rings (Rasmussen et al., 2013) (Figure 5A). In wildtype worms, *ajm-1*::*GFP* (*jcIs1* transgene) is strongly expressed through the adherens junctions lining the lumen of the pharynx and vpi, and the expression drops off sharply to low levels starting at the anteriormost ring of the intestine, and continuing throughout the entire length of the animal (Köppen et al., 2001; Sommermann et al., 2010). However, while *end-1(-)* and *elt-7(-/-) end-1(+/-)* animals show wildtype *ajm-1* expression pattern, *ajm-1* signal is markedly elevated in the anterior intestinal terminus of *elt-7(-/-) end-1(-/-)* animals (Figure 5B-E). Additionally, we observed ectopic expression of two vpi markers, *cdf (cation diffusion facilitator)-1* (Figure 5F-H) and *hum (heavy chain, unconventional myosin)-1* (Figure 5I-K), in the anterior terminus of *end-1(-/-) elt-7(-/-)* larvae.

It has been previously shown that worms lacking *elt-2*, like *elt-7(-/-) end-1(-/-)* animals, exhibit striking caudal extension of the valve cell markers (Sommermann et al., 2010). We found that ELT-2 is expressed at wildtype levels in *elt-7(-/-) end-1(-/-)* larvae, owing to its positive autoregulation (Figure 5L). This suggests that the expansion of vpi gene expression in the intestine that we observed in *elt-7(-/-) end-1(-/-)* animals may be independent of ELT-2 function. It is currently unclear whether the expression of vpi reporters in the intestine reflects *bona fide* transformation of gut cells into valve-like cells or aberrant development of vpi and mispositioning of excess valve cells. Regardless, our results demonstrate the important roles of intestinal GATA factors in the development of a properly patterned digestive tract.

Figure 5: ELT-7 and END-1 function synergistically to repress ectopic expression of valve cell markers in the anterior gut.

(A) Drawing highlighting the anatomy of pharynx, vpi, and intestine. The expression of *jcIs1[ajm-1::GFP]* transgene appears wildtype in (B, B') *end-1(-)* and (C, C') balanced *elt-7(-/-) end-1(+/-)* mutants. (D, D') *elt-7(-) end-1(-)* shows caudal expansion of *jcIs1* expression into the anterior gut (bracket). (E) The frequency of animals exhibiting intense *jcIs1* reporter expression in the anterior gut. (F, F') *cdf-1::GFP* expression is restricted to the vpi in *elt-7(-/-) end-1(+/-)*. (G, G') Ectopic expression of *cdf-1* reporter is observed in *elt-7(-/-) end-1(-/-)* animals (arrowhead). (H) The frequency of animals with ectopic *cdf-1::GFP* expression in the anterior gut. (I) HUM-1 is highly expressed in the vpi in wildtype animal as revealed by an endogenously tagged reporter. (J) Ectopic expression of *hum-1* is observed in *elt-7(-/-) end-1(-/-)* (arrowhead).

540   The intestinal cells are marked by *elt-2::GFP*. (K) The frequency of animals with ectopic *hum-*

541   *1::RFP* expression in the anterior gut. (L) The expression of ELT-2 is not altered in *elt-7(-/-) end-*

542   *1(+/-)* and *elt-7(-/-) end-1(-/-)*, compared to wildtype L1 larvae. All scale bars = 10µm. For panels

543   E, H, K, the number of animals scored is indicated in each graph. *** $p \leq 0.001$ by Fisher's exact

544   test.

Discussion

The development of *C. elegans* endoderm provides a powerful system to study the regulatory logic underlying cell specification and differentiation. In this study, we reported four major findings: (1) the hierarchical organization and feedforward regulation of GATA factors promote robust and rapid lockdown of endodermal cell fate during *C. elegans* early embryogenesis. (2) END-1 participates in both specification and differentiation and mediates a smooth regulatory state transition. (3) ELT-2 and -7 repress expression of *pha-4* in the midgut to establish regulatory state boundary between the pharynx and the intestine. (4) END-1, ELT-7, and ELT-2 repress the characteristics of pharyngeal valve cell fate at the anterior gut terminus, further defining the spatial domains of the foregut and midgut. Our study therefore provides an important insight into the regulatory circuits that direct specification-to-differentiation transition and subsequent restriction and maintenance of differentiation pattern during development.

Architecture of the *C. elegans* Endoderm Regulatory Cascade

This study, together with findings reported in other publications, provided a comprehensive view of the core endoderm regulatory cascade, with six GATA factors acting through reiterated sequential feedforward loops (Figure 6). At the top of the cascade, maternally provided SKN-1 binds to and turns on the *meds* and the *ends* (Maduro et al., 2001; Maduro et al., 2005b). Although MED-1 and MED-2 protein sequences are nearly identical, we found distinguishable contributions between the two paralogs, with MED-2 acting upstream of MED-1. Indeed, embryos lacking MED-1 show a weaker loss-of-gut phenotype than those lacking MED-2, when SKN-1 function is debilitated (Maduro et al., 2007). Moreover, *med-2* is expressed slightly earlier than *med-1* (Maduro et al., 2007; Tintori et al., 2016), suggesting they are differentially regulated as we have observed in this study.

In the E blastomere, SKN-1 and the MEDs collaboratively activate END-3. MED-1/2 and

END-3 in turn activate END-1, the next step in the cascade. While the two endoderm-specifying

factors, END-1 and -3, perform largely overlapping functions, our data suggest that END-3

alone is sufficient to direct endoderm specification and suppress inappropriate mesectodermal

development, consistent with the lack of detectable phenotype of in *end-1(-)* single mutants.

Interesting, we found that END-1, poised at the interface between specification and

differentiation, works synergistically with ELT-2 and -7 to activate the endoderm differentiation

program (Figure 6). Supporting our model, in vitro gel-shift assays demonstrate the binding of

END-1, ELT-7, and ELT-2 to TGATAA sites, which are highly enriched in the promotors of

intestinal genes (Du et al., 2016; McGhee et al., 2009; Wiesenfahrt et al., 2016). Remarkably,

END-1 is able to initiate endoderm differentiation in *Xenopus* embryos, demonstrating its role as

a potent organ selector (Shoichet et al., 2000). Hence, it appears that specification and

differentiation involve a *bona fide* handoff of regulatory events by END-1.

Regulatory Logic of Developmental GRN

Our results support the conclusion that the *C. elegans* endoderm GRN comprises a

recursive series of feedforward steps, culminating with rapid terminal differentiation (Figure 6).

Each GATA factor in the entire cascade receives redundant activating inputs acting through an

"OR" logic gate. Consequently, any single mutation in the regulatory cascade is largely

phenotypically silent, with the exception of *elt-2(-)*; however, even *elt-2(-)* animals contain what

appears to be a well-differentiated, albeit dysfunctional, intestine. Coherent feedforward motifs

of the type we observe reiteratively in the endoderm GRN are ubiquitous in developmental

GRNs. Such a network configuration appears to result in a rapid response to an activating

signal and a delayed response when the inputs are removed (termed sign-sensitive delay),

thereby prolonging the effect of a transient activator (Mangan and Alon, 2003). Additionally,

such feedforward loops are effective at buffering the system against stochastic noise, thereby

ensuring developmental robustness (Chepyala et al., 2016; Gui et al., 2016; Maduro, 2015).

This design principal appears to be crucial to ensure timely and robust activation of *elt-7* and *-2*,

as we and others have shown (Boeck et al., 2011; Dineen et al., 2018; Maduro et al., 2015).

Timely onset of *elt-2* appears to be critical, as its delay in early embryos has been shown to

cause sustained metabolic defects in larvae despite later reattainment of wildtype ELT-2 levels

(Maduro et al., 2015).

Our results suggest that END-1 mediates an all-or-none switch in the differentiation

program in the absence of ELT-2 and -7. END-1 straddles both specification and differentiation,

being buttressed by END-3 upstream and ELT-2/7 downstream. The only difference in these

END-1 functions appears to be its timing of action and partnership with another regulatory factor.

Regulatory nodes in early specification can indeed directly control morphogenetic events in

various contexts (Davidson, 2010; Zhu and Rosenfeld, 2004). Interestingly, although END-1

expression decays shortly after gastrulation, we observed a persistent reduction in *act-5*

expression in postembryonic larvae lacking *end-1*. One possible explanation for this observation,

and the finding that patches of well-differentiated gut arise in *elt-7(-); elt-2(-)* double mutants

long after END-1 becomes undetectable, might be that END-1 targets an unidentified factor (X;

Figure 6) that directs differentiation in at least a subset of endodermal progenitors. Like ELT-2

and -7, expression of the putative X factor would be expected to be maintained through a

positive autoregulatory loop, thereby continuing to promote gut differentiation after the

expression of END-1 has subsided (Figure 6) (Sommermann et al., 2010). Alternatively, END-1

may directly activate differentiation gene batteries in early embryos, and that regulatory state

might be sustained through the propagation of epigenetic memory. This priming mechanism has

recently been demonstrated in the specification of the ASE sensory neurons in *C. elegans*

(Charest et al., 2020). Two transiently expressed T-box factors, TBX-37 and -38, lock their

target, *lsy-6*, in a transcriptionally active state during early embryogenesis, priming it for

activation in restricted neuronal lineage (Charest et al., 2020). In mammals, Pax-7 (paired-box-7)

initiates myogenic specification and differentiation. Interestingly, many enhancers of Pax-7

target genes retain epigenetic signatures and remain active even in the absence of the initial

activator (Zhang et al., 2020).

We found that knocking down *elt-2* causes a slight but significant increase in *end-1*

expression in early embryos; however, we did not observe obvious perdurance of END-1 when

ELT-2 and ELT-7 are depleted, though we cannot rule out the possibility that the modest

negative feedback we observed resulted from incomplete RNAi penetrance. Nevertheless, we

propose that this modest feedback inhibition may function to facilitate the transition of regulatory

states and ensure that developmental process moves inexorably forward. For example,

repression of an early cardiac specification factor, Bmp2, by the Nkx2-5 homeodomain factor is

necessary for proper morphological development of the heart in mice (Prall et al., 2007). Such

transcriptional repression is also frequently used to install spatial subdivision of regulatory states

(Davidson, 2010). As we have shown above, ELT-2, ELT-7 and END-1 may repress alternate

cell fates in the midgut and define the boundaries of the digestive tract. It is noteworthy that

structurally similar regulatory circuits are repeatedly deployed in biological networks of different

contexts while performing similar functions. Thus, the functional output of a GRN depends not

only on the specificity of the transcription factors, but also the underlying circuit architecture

(Davidson, 2010; Peter, 2020).

Rapid Rewiring of the Endoderm GRN in *Caenorhabditis*

How might a regulatory system of the type described here evolve? Effectors acting on

terminal differentiation gene batteries, such as ELT-2 and PHA-4, are widely conserved across

the animal kingdom, while the upstream inputs into GRNs appear to be recent innovations that

arose during the radiation of the Elegans supergroup within the *Caenorhabditis* genus (Maduro,

2020). The *ends* and *meds* have been proposed to have arisen from the duplication of *elt-2*.

644 Hence, gene duplication, coupled with *cis*-regulatory changes, may have led to the emergence

645 of new circuitry and rewiring of the endoderm GRN in nematodes through intercalation of these

646 factors in the regulatory network.

647   Intercalation of the MEDs and ENDs in the cascade may serve to buffer the system

648 against environmental variation and developmental noise by freeing ELT-2 from the direct

649 control of SKN-1, which has been shown to play conserved pleiotropic roles in stress response

650 and lifespan regulation (reviewed in Ewe et al., 2020; Ewe et al., 2021). As we and others have

651 shown, robust induction of *elt-2* is critical to ensure the viability and fitness of the animals

652 (Maduro et al., 2015; Raj et al., 2010). Moreover, the deployment of MEDs and ENDs in the

653 sequential hierarchy may allow canalization of the endodermal lineage by rapidly establishing its

654 regulatory state in the E blastomere (Peter and Davidson, 2011). Consequently, this structure

655 may enable increased developmental speed and early specification of the founder cells in

656 *Caenorhabditis* species.

Figure 6: Current model for the *C. elegans* endoderm GRN.

Solid lines indicate known interactions identified biochemically or implied genetically, while dashed lines represent proposed interactions. Maternal factor SKN-1 initiates the endoderm cascade which consists of GATA transcription factors arranged in interlocking feedforward circuits. END-1 and -3 are also regulated by non-GATA transcriptional factors, including SPTF-3 (Sullivan-Brown et al., 2016), PAL-1 (Maduro et al., 2005b), PLP-1 (Witze et al., 2009), and POP-1 (Maduro et al., 2005b), which are omitted from this model for simplicity. The observation of residual gut differentiation in *elt-7(-) end-(-)* larvae suggests that END-3 may directly act on ELT-2 and initiate gut differentiation program in some, but not all, of the endodermal progenitors in the double mutants. END-1 may activate an unidentified transcription factor (X) which

functions to promote gut differentiation after END-1 decays at ~16E embryonic stage (see text

for details).

Acknowledgments

We thank members of Rothman lab for helpful advice and feedback, and Cricket Wood for

excellent technical support. We thank Dr. Irene Chen (University of California Los Angeles) for

her helpful inputs and Dr. Zhuo Du (University of Chinese Academy of Sciences) for providing

strain SYS549. We would also like to thank Dr. Rubayn Goh, Dr. Francesca Zappa, Nerea

Muniozguren, and Soham Chowdhury for technical assistance. We acknowledge the use of the

UCSB NRI-MCDB Microscopy Facility supported by the NIH Grant S10OD010610-01A1, as well

as WormBase. Some nematode strains used in this work were provided by the Caenorhabditis

Genetics Center (CGC), which is funded by NIH Office of Research Infrastructure Programs

Grant P40 OD010440.

Competing Interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: CKE, EMS, JHR; Resources: CKE, EMS, SEF, MFM; Formal analysis: CKE,

EMS, JK; Investigation: CKE, EMS, JK; Writing - original draft: CKE; Writing – review & editing:

CKE, EMS, JK, SEF, MFM, JHR; Supervision: JHR; Funding acquisition: JHR

Funding

This work is supported by National Institute of Health ((#1R01HD082347 and # 1R01HD081266)

to JHR and NSF (IOS#1258054) to MM.

Data Availability

All data generated or analyzed during this study are included in this article and accompanied

supplementary materials. The source code for the GRN computational model is available on

https://github.com/RothmanLabCode/endoderm_GRN_model.

References

Baugh, L. R., Hill, A. A., Slonim, D. K., Brown, E. L. and Hunter, C. P. (2003). Composition
and dynamics of the Caenorhabditis elegans early embryonic transcriptome. *Development*
130, 889–900.

Bei, Y., Hogan, J., Berkowitz, L. A., Soto, M., Rocheleau, C. E., Pang, K. M., Collins, J. and
Mello, C. C. (2002). SRC-1 and Wnt signaling act together to specify endoderm and to
control cleavage orientation in early C. elegans embryos. *Dev. Cell* **3**, 113–25.

Boeck, M. E., Boyle, T., Bao, Z., Murray, J., Mericle, B. and Waterston, R. (2011). Specific
roles for the GATA transcription factors end-1 and end-3 during C. elegans E-lineage
development. *Dev. Biol.* **358**, 345–355.

Boveri, T. (1899). *Die Entwickelung von Ascaris megalocephala mit besonderer Rücksicht auf*
die Kernverhältnisse. Jena, Gustav Fischer (Opitz Library).

Bowerman, B., Eaton, B. A. and Priess, J. R. (1992). skn-1, a maternally expressed gene
required to specify the fate of ventral blastomeres in the early C. elegans embryo. *Cell* **68**,
1061–75.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94.

Broitman-Maduro, G., Lin, K. T.-H., Hung, W. W. K. and Maduro, M. F. (2006). Specification
of the C. elegans MS blastomere by the T-box factor TBX-35. *Development* **133**, 3097–106.

Charest, J., Daniele, T., Wang, J., Bykov, A., Mandlbauer, A., Asparuhova, M., Röhsner, J.,
Gutiérrez-Pérez, P. and Cochella, L. (2020). Combinatorial Action of Temporally
Segregated Transcription Factors. *Dev. Cell* **55**, 483-499.e7.

Chepyala, S. R., Chen, Y. C., Yan, C. C. S., Lu, C. Y. D., Wu, Y. C. and Hsu, C. P. (2016).
Noise propagation with interlinked feed-forward pathways. *Sci. Rep.* **6**, 1–15.

Choi, H., Broitman-Maduro, G. and Maduro, M. F. (2017). Partially compromised specification
causes stochastic effects on gut development in C. elegans. *Dev. Biol.* **427**, 49–60.

Clokey, G. V and Jacobson, L. A. (1986). The autofluorescent "lipofuscin granules" in the
intestinal cells of Caenorhabditis elegans are secondary lysosomes. *Mech. Ageing Dev.* **35**,
79–94.

Davidson, E. H. (2010). Emerging properties of animal gene regulatory networks. *Nature* **468**,

911–920.

Davidson, E. H. and Levine, M. S. (2008). Properties of developmental gene regulatory
networks. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 20063–6.

Davidson, E. H., Rast, J. P., Oliveri, P., Ransick, A., Calestani, C., Yuh, C. H., Minokawa, T.,
Amore, G., Hinman, V., Arenas-Mena, C., et al. (2002). A genomic regulatory network for
development. *Science (80-.).* **295**, 1669–1678.

Dineen, A., Osborne Nishimura, E., Goszczynski, B., Rothman, J. H. and McGhee, J. D.
(2018). Quantitating transcription factor redundancy: The relative roles of the ELT-2 and
ELT-7 GATA factors in the C. elegans endoderm. *Dev. Biol.* **435**, 150–161.

Du, L., Tracy, S. and Rifkin, S. A. (2016). Mutagenesis of GATA motifs controlling the
endoderm regulator elt-2 reveals distinct dominant and secondary cis-regulatory elements.
Dev. Biol. **412**, 160–170.

Ewe, C. K., Torres Cleuren, Y. N. and Rothman, J. H. (2020). Evolution and developmental
system drift in the endoderm gene regulatory network of Caenorhabditis and other
nematodes. *Front. Cell Dev. Biol.* **8**, 170.

Ewe, C. K., Alok, G. and Rothman, J. H. (2021). Stressful development: integrating endoderm
development, stress, and longevity. *Dev. Biol.* **471**, 34–48.

Fakhouri, T. H. I., Stevenson, J., Chisholm, A. D. and Mango, S. E. (2010). Dynamic
Chromatin Organization during Foregut Development Mediated by the Organ Selector
Gene PHA-4/FoxA. *PLOS Genet.* **6**, e1001060.

Fay, D. S. (2013). Classical genetic methods. *WormBook* 1–58.

Fukushige, T., Hawkins, M. G. and McGhee, J. D. (1998). The GATA-factor elt-2 is essential
for formation of the Caenorhabditis elegans intestine. *Dev. Biol.* **198**, 286–302.

Gaudet, J., Muttumu, S., Horner, M. and Mango, S. E. (2004). Whole-Genome Analysis of
Temporal Gene Expression during Foregut Development. *PLOS Biol.* **2**, e352.

Gilleard, J. S. and McGhee, J. D. (2001). Activation of Hypodermal Differentiation in the
Caenorhabditis elegans Embryo by GATA Transcription Factors ELT-1 and ELT-3 . *Mol.*
Cell. Biol. **21**, 2533–2544.

Gui, R., Liu, Q., Yao, Y., Deng, H., Ma, C., Jia, Y. and Yi, M. (2016). Noise decomposition

principle in a coherent feed-forward transcriptional regulatory loop. *Front. Physiol.* **7**, 600.

Hahn, G. D. (1991). A modified Euler method for dynamic analyses. *Int. J. Numer. Methods Eng.*
32, 943–955.

Hashimshony, T., Feder, M., Levin, M., Hall, B. K. and Yanai, I. (2015). Spatiotemporal
transcriptomics reveals the evolutionary history of the endoderm germ layer. *Nature* **519**,
219–222.

Horner, M. A., Quintin, S., Domeier, M. E., Kimble, J., Labouesse, M. and Mango, S. E.
(1998). pha-4, an HNF-3 homolog, specifies pharyngeal organ identity in
Caenorhabditis elegans. *Genes Dev.* **12**, 1947.

Hunter, C. P. and Kenyon, C. (1996). Spatial and temporal controls target pal-1 blastomere-
specification activity to a single blastomere lineage in C. elegans embryos. *Cell* **87**, 217–
226.

Kalb, J. M., Lau, K. K., Goszczynski, B., Fukushige, T., Moons, D., Okkema, P. G. and
McGhee, J. D. (1998). pha-4 is Ce-fkh-1, a fork head/HNF-3alpha,beta,gamma homolog
that functions in organogenesis of the C. elegans pharynx. *Development* **125**, 2171–80.

Kamath, R. S., Fraser, A. G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le
Bot, N., Moreno, S., Sohrmann, M., et al. (2003). Systematic functional analysis of the
Caenorhabditis elegans genome using RNAi. *Nature* **421**, 231–237.

Köppen, M., Simske, J. S., Sims, P. A., Firestein, B. L., Hall, D. H., Radice, A. D., Rongo, C.
and Hardin, J. D. (2001). Cooperative regulation of AJM-1 controls junctional integrity in
Caenorhabditis elegans epithelia. *Nat. Cell Biol.* **3**, 983–991.

Li, X., Zhao, Z., Xu, W., Fan, R., Xiao, L., Ma, X. and Du, Z. (2019). Systems Properties and
Spatiotemporal Regulation of Cell Position Variability during Embryogenesis. *Cell Rep.* **26**,
313-321.e7.

Livak, K. J. and Schmittgen, T. D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408.

MacQueen, A. J., Baggett, J. J., Perumov, N., Bauer, R. A., Januszewski, T., Schriefer, L.
and Waddle, J. A. (2005). ACT-5 is an essential Caenorhabditis elegans actin required for
intestinal microvilli formation. *Mol. Biol. Cell* **16**, 3247–3259.

Maduro, M. F. (2015). Developmental robustness in the *Caenorhabditis elegans* embryo. *Mol.*
Reprod. Dev. **82**, 918–931.

Maduro, M. F. (2017). Gut development in C. elegans. *Semin. Cell Dev. Biol.* **66**, 3–11.

Maduro, M. F. (2020). Evolutionary dynamics of the Skn-1/Med/end-1,3 regulatory gene
cascade in Caenorhabditis endoderm specification. *G3 Genes, Genomes, Genet.* **10**, 333–
356.

Maduro, M. F. and Rothman, J. H. (2002). Making Worm Guts: The Gene Regulatory Network
of the Caenorhabditis elegans Endoderm. *Dev. Biol.* **246**, 68–85.

Maduro, M. F., Meneghini, M. D., Bowerman, B., Broitman-Maduro, G. and Rothman, J. H.
(2001). Restriction of mesendoderm to a single blastomere by the combined action of SKN-
1 and a GSK-3beta homolog is mediated by MED-1 and -2 in C. elegans. *Mol. Cell* **7**, 475–
85.

Maduro, M. F., Lin, R. and Rothman, J. H. (2002). Dynamics of a Developmental Switch:
Recursive Intracellular and Intranuclear Redistribution of Caenorhabditis elegans POP-1
Parallels Wnt-Inhibited Transcriptional Repression. *Dev. Biol.* **248**, 128–142.

Maduro, M. F., Hill, R. J., Heid, P. J., Newman-Smith, E. D., Zhu, J., Priess, J. R. and
Rothman, J. H. (2005a). Genetic redundancy in endoderm specification within the genus
Caenorhabditis. *Dev. Biol.* **284**, 509–522.

Maduro, M. F., Kasmir, J. J., Zhu, J. and Rothman, J. H. (2005b). The Wnt effector POP-1
and the PAL-1/Caudal homeoprotein collaborate with SKN-1 to activate C. elegans
endoderm development. *Dev. Biol.* **285**, 510–523.

Maduro, M. F., Broitman-Maduro, G., Mengarelli, I. and Rothman, J. H. (2007). Maternal
deployment of the embryonic SKN-1 → MED-1,2 cell specification pathway in C. elegans.
Dev. Biol. **301**, 590–601.

Maduro, M. F., Broitman-Maduro, G., Choi, H., Carranza, F., Wu, A. C.-Y. and Rifkin, S. A.
(2015). MED GATA factors promote robust development of the C. elegans endoderm. *Dev.*
Biol. **404**, 66–79.

Mangan, S. and Alon, U. (2003). Structure and function of the feed-forward loop network motif.
Proc. Natl. Acad. Sci. U. S. A. **100**, 11980–11985.

Mango, S. E., Lambie, E. J. and Kimble, J. (1994). The pha-4 gene is required to generate the
pharyngeal primordium of Caenorhabditis elegans. *Development* **120**, 3019–3031.

Martindale, M. Q., Pang, K. and Finnerty, J. R. (2004). Investigating the origins of triploblasty:
"mesodermal" gene expression in a diploblastic animal, the sea anemone Nematostella
vectensis (phylum, Cnidaria; class, Anthozoa). *Development* **131**, 2463–74.

McGhee, J. (2007). The C. elegans intestine. *WormBook* 1–36.

McGhee, J. D., Fukushige, T., Krause, M. W., Minnema, S. E., Goszczynski, B., Gaudet, J.,
Kohara, Y., Bossinger, O., Zhao, Y., Khattra, J., et al. (2009). ELT-2 is the predominant
transcription factor controlling differentiation and function of the C. elegans intestine, from
embryo to adult. *Dev. Biol.* **327**, 551–565.

Meneghini, M. D., Ishitani, T., Carter, J. C., Hisamoto, N., Ninomiya-Tsuji, J., Thorpe, C. J.,
Hamill, D. R., Matsumoto, K. and Bowerman, B. (1999). MAP kinase and Wnt pathways
converge to downregulate an HMG-domain repressor in Caenorhabditis elegans. *Nature*
399, 793–797.

Murakami, R., Okumura, T. and Uchiyama, H. (2005). GATA factors as key regulatory
molecules in the development of Drosophila endoderm. *Dev. Growth Differ.* **47**, 581–589.

Nakanishi, N., Sogabe, S. and Degnan, B. M. (2014). Evolutionary origin of gastrulation:
Insights from sponge development. *BMC Biol.* **12**,.

Okumura, T., Matsumoto, A., Tanimura, T. and Murakami, R. (2005). An endoderm-specific
GATA factor gene, dGATAe, is required for the terminal differentiation of the Drosophila
endoderm. *Dev. Biol.* **278**, 576–586.

Owraghi, M., Broitman-Maduro, G., Luu, T., Roberson, H. and Maduro, M. F. (2010). Roles
of the Wnt effector POP-1/TCF in the C. elegans endomesoderm specification gene
network. *Dev. Biol.* **340**, 209–21.

Peter, I. S. (2020). The function of architecture and logic in developmental gene regulatory
networks. In *Current Topics in Developmental Biology*, pp. 267–295. Academic Press Inc.

Peter, I. S. and Davidson, E. H. (2011). Evolution of gene regulatory networks controlling body
plan development. *Cell* **144**, 970–85.

Prall, O. W. J., Menon, M. K., Solloway, M. J., Watanabe, Y., Zaffran, S., Bajolle, F., Biben,

C., McBride, J. J., Robertson, B. R., Chaulet, H., et al. (2007). An Nkx2-5/Bmp2/Smad1
Negative Feedback Loop Controls Heart Progenitor Specification and Proliferation. *Cell*
128, 947–959.

Raj, A., Rifkin, S. A., Andersen, E. and van Oudenaarden, A. (2010). Variability in gene
expression underlies incomplete penetrance. *Nature* **463**, 913–918.

Rasmussen, J. P., Feldman, J. L., Reddy, S. S. and Priess, J. R. (2013). Cell Interactions
and Patterned Intercalations Shape and Link Epithelial Tubes in C. elegans. *PLoS Genet.* **9**,
1003772.

Rocheleau, C. E., Yasuda, J., Shin, T. H., Lin, R., Sawa, H., Okano, H., Priess, J. R., Davis,
R. J. and Mello, C. C. (1999). WRM-1 Activates the LIT-1 Protein Kinase to Transduce
Anterior/Posterior Polarity Signals in C. elegans. *Cell* **97**, 717–726.

Rodaway, A. and Patient, R. (2001). Mesendoderm: An Ancient Germ Layer? *Cell* **105**, 169–
172.

Rual, J.-F., Ceron, J., Koreth, J., Hao, T., Nicot, A.-S., Hirozane-Kishikawa, T.,
Vandenhaute, J., Orkin, S. H., Hill, D. E., van den Heuvel, S., et al. (2004). Toward
Improving Caenorhabditis elegans Phenome Mapping With an ORFeome-Based RNAi
Library. *Genome Res.* **14**, 2162–2168.

Ruhe, A. (1979). Accelerated Gauss-Newton algorithms for nonlinear least squares problems.
BIT Numer. Math. 1979 193 **19**, 356–367.

Shin, T. H., Yasuda, J., Rocheleau, C. E., Lin, R., Soto, M., Bei, Y., Davis, R. J. and Mello, C.
C. (1999). MOM-4, a MAP kinase kinase kinase-related protein, activates WRM-1/LIT-1
kinase to transduce anterior/posterior polarity signals in C. elegans. *Mol. Cell* **4**, 275–80.

Shoichet, S. A., Malik, T. H., Rothman, J. H. and Shivdasani, R. A. (2000). Action of the
Caenorhabditis elegans GATA factor END-1 in Xenopus suggests that similar mechanisms
initiate endoderm development in ecdysozoa and vertebrates. *Proc. Natl. Acad. Sci. U. S.*
A. **97**, 4076–81.

Sommermann, E. M., Strohmaier, K. R., Maduro, M. F. and Rothman, J. H. (2010).
Endoderm development in Caenorhabditis elegans: The synergistic action of ELT-2 and -7
mediates the specification→differentiation transition. *Dev. Biol.* **347**, 154–166.

Sullivan-Brown, J. L., Tandon, P., Bird, K. E., Dickinson, D. J., Tintori, S. C., Heppert, J. K.,

Meserve, J. H., Trogden, K. P., Orlowski, S. K., Conlon, F. L., et al. (2016). Identifying
regulators of morphogenesis common to vertebrate neural tube closure and
Caenorhabditis elegans gastrulation. *Genetics* **202**, 123–139.

Sulston, J. E., Schierenberg, E., White, J. G. and Thomson, J. N. (1983). The embryonic cell
lineage of the nematode Caenorhabditis elegans. *Dev. Biol.* **100**, 64–119.

Thorpe, C. J., Schlesinger, A., Carter, J. C. and Bowerman, B. (1997). Wnt signaling
polarizes an early C. elegans blastomere to distinguish endoderm from mesoderm. *Cell* **90**,
695–705.

Tintori, S. C., Osborne Nishimura, E., Golden, P., Lieb, J. D. and Goldstein, B. (2016). A
Transcriptional Lineage of the Early C. elegans Embryo. *Dev. Cell* **38**, 430–444.

Waddington, C. (1957). *The strategy of the genes.* London: Volume George Allen and Unwin.

Wiesenfahrt, T., Berg, J. Y., Osborne Nishimura, E., Robinson, A. G., Goszczynski, B.,
Lieb, J. D. and McGhee, J. D. (2016). The function and regulation of the GATA factor
ELT-2 in the *C. elegans* endoderm. *Development* **143**, 483–491.

Witze, E. S., Field, E. D., Hunt, D. F. and Rothman, J. H. (2009). C. elegans pur alpha, an
activator of end-1, synergizes with the Wnt pathway to specify endoderm. *Dev. Biol.* **327**,
12–23.

Yip, K. Y., Alexander, R. P., Yan, K.-K. and Gerstein, M. (2010). Improved Reconstruction of
In Silico Gene Regulatory Networks by Integrating Knockout and Perturbation Data. *PLoS*
One **5**, e8121.

Yuh, C. H., Dorman, E. R., Howard, M. L. and Davidson, E. H. (2004). An otx cis-regulatory
module: a key node in the sea urchin endomesoderm gene regulatory network. *Dev. Biol.*
269, 536–551.

Zhang, N., Mendieta-Esteban, J., Magli, A., Lilja, K. C., Perlingeiro, R. C. R., Marti-Renom,
M. A., Tsirigos, A. and Dynlacht, B. D. (2020). Muscle progenitor specification and
myogenic differentiation are associated with changes in chromatin topology. *Nat. Commun.*
11, 1–18.

Zhong, M., Niu, W., Lu, Z. J., Sarov, M., Murray, J. I., Janette, J., Raha, D., Sheaffer, K. L.,
Lam, H. Y. K., Preston, E., et al. (2010). Genome-Wide Identification of Binding Sites
Defines Distinct Functions for Caenorhabditis elegans PHA-4/FOXA in Development and

Environmental Response. *PLoS Genet.* **6**, e1000848.

Zhu, X. and Rosenfeld, M. G. (2004). Transcriptional control of precursor proliferation in the
early phases of pituitary development. *Curr. Opin. Genet. Dev.* **14**, 567–574.

Zhu, J., Hill, R. J., Heid, P. J., Fukuyama, M., Sugimoto, A., Priess, J. R. and Rothman, J. H.
(1997). end-1 encodes an apparent GATA factor that specifies the endoderm precursor in
Caenorhabditis elegans embryos. *Genes Dev.* **11**, 2883–96.

Supplementary Materials

<u>Description of computational model</u>

The interactions between the GATA factors (Figure 6), as well as POP-1 activation of *end-3* and

end-1, were written as a system of differential equations. Each factor is expressed according to

the concentration of its activators multiplied by respective coefficients *a*, representing the

strength of activation (see Supplementary File 1). The activation is not instant, but delayed by a

time interval τ, which accounts for the time required for transcription, translation, and re-

localization to the nucleus. SKN-1 and POP-1 activation occur as square waves in EMS and E

blastomere, respectively. ELT-7 and ELT-2 self-activate via feedback loops of strength f_1 and f_2,

respectively, after surpassing arbitrary concentration thresholds Φ_1 and Φ_2, respectively. Finally,

all factors are degraded at the same rate δ. The resulting system of equations is as follows.

(1) $\frac{d[MED2]}{dt} = a_1[SKN1]_t - \delta[MED2]_t$

(2) $\frac{d[MED1]}{dt} = a_2[SKN1]_t + a_5[MED2]_{t-\tau} - \delta[MED1]_t$

(3) $\frac{d[END3]}{dt} = a_3[SKN1]_t + a_9[POP1]_t + a_6[MED2]_{t-\tau} + a_7[MED1]_{t-\tau} - \delta[END3]_t$

(4) $\frac{d[END1]}{dt} = a_4[SKN1]_t + a_{10}[POP1]_t + a_8[MED1]_{t-\tau} + a_{11}[END3]_{t-\tau} - \delta[END1]_t$

(5) $\frac{d[ELT7]}{dt} = a_{12}[END3]_{t-\tau} + a_{13}[END1]_{t-\tau} + f_1([ELT7]_{t-\tau} > \Phi_1)[ELT7]_{t-\tau} - \delta[ELT7]_t$

(6) $\frac{d[ELT2]}{dt} = a_{14}[END1]_{t-\tau} + a_{15}[ELT7]_{t-\tau} + f_2([ELT2]_{t-\tau} > \Phi_2)[ELT2]_{t-\tau} - \delta[ELT2]_t$

Model parameters (Supplementary File 1) were determined by fitting to transcriptomics data by

a custom algorithm written in R following an iterative least-squares method. Parameter guesses

were provided, and the model was run using an Euler approximation with 0.01 s time steps. The

model fit was measured as the sum of squared differences between model predictions and

transcriptomics data. Randomly selected parameter values were randomly changed, and the

model was run again. If the resulting sum of squares was lower, the new parameter values were

kept. This process was iterated ~10^6 times. Algorithm-selected parameter values were validated

by checking model predictions against the published phenotypes of the single mutants (Boeck

et al., 2011; Dineen et al., 2018; Maduro et al., 2015). Two parameters (a_8 and a_{14}) were

adjusted manually such that the predicted phenotype of *end-3(-)* matched that reported in

published studies (Boeck et al., 2011; Maduro et al., 2005a). The source code is available on

https://github.com/RothmanLabCode/endoderm_GRN_model.

Supplementary Figure 1: *C. elegans* **mesendoderm development.**

(A) At the four-cell stage, EMS (blue) undergoes asymmetrical division to produce anterior MS (green) and posterior E (orange) blastomeres. Newly hatched L1s contain 20 intestinal cells arranged in nine rings (Ints). Int1 contains four cells, while the remaining eight rings contain two cells each. The MS cell gives rise mesodermal cell types, including the posterior pharynx. (B) Simplified mesendoderm GRN showing three sequential tiers of paired redundant GATA transcription factors (2 redundant MEDs → 2 redundant ENDs → 2 redundant ELTs). Maternally provided SKN-1 activates *med-1* and *-2*, which have both a maternal and zygotic component. In MS, POP-1 represses the expression of *end-1* and *-3* genes. MED-1 and -2 then directly activate the expression of mesoderm-specifying factor, TBX-35. In E, however, Wnt, MAPK and Src signaling from neighboring P2 cell ultimately leads to the phosphorylation of POP-1 (indicated by *), converting it from a repressor to an activator of gut fate. END-1 and -3 subsequently activate the expression of ELT-7 and -2, both of which promote gut morphological differentiation.

A

B

955 **Supplementary Figure 2: Severe gut defects in mutants lacking sequential GATA pairs.**

956 Eliminating sequential GATA pairs causes impaired gut differentiation and aberrant expression

957 of immunoreactive (A) IFB-2 and (B) AJM-1.

Supplementary Figure 3: Mutants lacking alternate GATA pairs do not show apparent gut defects.

(A) Mutants lacking alternate GATA pairs contain fully differentiated lumen (top row) and gut granules (bottom row) along the length of the animals. The same set of double mutants show wildtype expression of (B, C) immunoreactive IFB-2 and (D, E) AJM-1.

966 **Supplementary Figure 4: The endoderm GATA factors are deployed in temporal order.**

967 (A) Temporal expression of *med-2*, *med-1*, *end-3,* and *end-1* revealed by single-cell

968 transcriptomic analysis (http://tintori.bio.unc.edu/) (Tintori et al., 2016). (B) Expression of MED-2

969 and MED-1 protein-fusion reporters in staged embryos.

Supplementary Figure 5: Reduced number of differentiated intestinal cells *in elt-2(-); elt-7(RNAi)* animals.

On average, *elt-2(-)* animals contain 19.3 cells, while *elt-2(-); elt-7(RNAi)* animals contain 14.1 cells. The number of gut cells were scored by the expression of *elt-2p::GFP* transcriptional reporter *wIs84*. *** p ≤ 0.001 by Wilcoxon tests.

Supplementary Figure 6: Reduced number of differentiated intestinal cells in *elt-7(-) end-1(-)* animals.

(A-C) The expression of *elt-2::GFP* translational reporter in (A, A') wildtype, (B, B') *elt-7(-/-) end-1(+/-)*, and (C, C') *elt-7(-/-) end-1(-/-)* L1 larvae. (D) *elt-7(-/-) end-1(-/-)* double mutants contain fewer cells expressing immunoreactive ELT-2 than *elt-7(-/-) end-1(+/-)* animals. *** p ≤ 0.001 by Wilcoxon tests.

Supplementary Figure 7: Reduced number of differentiated gut cells in differentiation-defective mutants.

(A-D) DIC images of (A) wildtype, (B) *elt-2(-)*, (C) *elt-2(-); elt-7(RNAi),* and (D) *elt-7(-) end-1(-); elt-2(-)* animals. While wildtype and *elt-2(-)* worms contain a differentiated gut, *elt-2(-); elt-7(RNAi)* animals lack evident lumen and contains sporadic patches of gut granules. *elt-7(-) end-1(-); elt-2(-)* triple mutants show no apparent signs of differentiation. (A'-D') Fluorescent images of worms in (A-D) show expression of *opt-2p::mCherry*. The number of *opt-2*-expressing cells is markedly reduced in *elt-2(-); elt-7(RNAi)* (arrowhead)*. opt-2* expression is completely abolished in *elt-7(-) end-1(-); elt-2(-).* Scale bars = 10 µm.

Supplementary Figure 8: END-1, ELT-7, and ELT-2 regulate *act-5* expression.

act-5 transgene (*jyIs13*) is expressed strongly in the intestine and weakly in the excretory canal

cells. *act-5::GFP* signals appear sporadic in *elt-7(-/-) end-1(+/-); elt-2(RNAi)* and *elt-7(-/-) end-*

1(-/-) animals, and are almost completely missing in *elt-7(-/-) end-1(-/-); elt-2(RNAi)* mutant. The

residual *act-5* expression in *elt-7(-/-) end-1(-/-); elt-2(RNAi)* may be due to incomplete RNAi

penetrance*.* Scale bars = 10 µm.

1001

Supplementary Figure 9: Gross epidermal defects in GATA mutants.

(A) A representative *end-1(-) end-3(-)* arrested L1 shows gross epidermal defects due to E→C

misspecification. (B) *elt-7(-) end-1(-); elt-2(-)* mutant does not show obvious epidermal defects,

as observed by DIC microscopy. Scale bars = 10 µm. (C) A large fraction of *end-1(-) end-3(-)*

mutants showed deformations of the epidermis, which was rarely observed in *elt-7(-) end-1(-);*

elt-2(-) worms. Number of animals scored for each genotype is indicated.

1008

Supplementary Figure 10: Eliminating *end-1*, *elt-7* and *elt-2* causes ectopic expression of *ceh-22* reporter.

(A) A representative *end-1(-); elt-2(-)* worm contains a differentiated gut with defined lumen (top) and wildtype expression pattern of *ceh-22* that is restricted to the pharynx (bottom). On the other hand, *end-1(-); elt-2(-); elt-7(RNAi)* mutant shows no sign of gut differentiation as observed by DIC microscopy (top), and ectopic expression of *ceh-22p::GFP* reporter (bottom). Scale bars = 10 μm. (B) Knocking out *elt-7* in *end-1(-); elt-2(-)* worms causes ectopic expression of *ceh-22p::GFP* marker, as shown in (A). Number of animals scored for each genotype is indicated.

1018

1019 **Supplementary Table 1: Worm strains used in this study.**

Strain	Genotype	Source
MS1893	*unc-119(ed9?) III; irSi24[opt-2p::mCherry::H2B, Cb-unc-119(+)] IV*	(Choi et al., 2017)
IA105	*unc-76(e911)V; ijIs12[dpy-7p::GFP::lacZ, unc-76(+)]*	(Gilleard and McGhee, 2001)
SYS549	*ujIs113[pie-1p::mCherry::H2B; nhr-2p::mCherry::HIS-24-let858UTR;unc119(+)] II; end-1(dev162[end-1::mNeonGreen]) V*	(Li et al., 2019)
JR1188	*wIs95[med-1p::GFP::MED-1+rol-6(su1006)]*	(Maduro et al., 2001)
MS1362	*med-2(cx9744) III; med-1(ok804) X; irEx581[T24D3, sur-5::dsRed]*	(Maduro et al., 2007)
MS386	*med-2(cx9744) III; dpy-11(e224) end-1(ok558) V*	(Maduro et al., 2007)
MS387	*dpy-11(e224) end-1(ok558) V; med-1(ok804) X*	(Maduro et al., 2007)
MS401	*end-3(ok1448) V; med-1(ok804) X*	(Maduro et al., 2007)
MS436	*med-2(cx9744) III; end-3(ok1448) V*	(Maduro et al., 2007)
MS1248	*end-1(ok558) end-3(ok1448) V; irEx568[K10F6, F58E10, pAS152(sur-5::dsRed)]*	(Owraghi et al., 2010)
JR3295	*elt-7(tm840) V; elt-2(ca15) X; irEx404[unc-119::CFP and elt-2(+)]*	(Sommermann et al., 2010)
MS851	*elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	(Sommermann et al., 2010)
RW10479	*unc-119(ed3) III; itIs37[pie-1p::mCherry::H2B::pie-1*	CGC

	3'UTR + unc-119(+)] IV; stIs10116[his-72(promoter)::his-24::mCherry::let-858 3'UTR + unc-119(+)]; stIs10426[med-2::TGF(6.2B3)::GFP::TY1::3xFLAG]	
AZ217	*unc-119(ed3) ruIs37 [myo-2p::GFP + unc-119(+)] III*	CGC
FX840	*elt-7(tm840) V*	CGC
MS123	*med-2(cx9744) II*	CGC
OP56	*unc-119(ed3) III; gaEx290[elt-2::TY1::EGFP::3xFLAG(92C12) + unc-119(+)]*	CGC
RB1331	*end-3(ok1448) V*	CGC
RB930	*med-1(ok804) X*	CGC
VC271	*end-1(ok558) V*	CGC
ERT60	*jyIs13 [act-5p::GFP::ACT-5 + rol-6(su1006)] II*	CGC
OH15876	*pha-4(ot946[pha-4::GFP]) V*	CGC
JR4405	*ruIs37 [myo-2p::GFP + unc-119(+)] III; him-5(e1490) V; elt-2(ca15) X; IrEx404[unc-119::CFP, elt-2(+)]*	This study
JR4384	*ruIs37 [myo-2p::GFP + unc-119(+)] III; elt-7(tm840) V; elt-2(ca15) X; IrEx404[unc-119::CFP, elt-2(+)]*	This study
JR4369	*hum-1(cv21[hum-1::RFP]) I; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; gaEx290[elt-2::TY1::EGFP::3xFLAG(92C12) + unc-119(+)]*	This study
JR4371	*hum-1(cv21[hum-1::RFP]) I; gaEx290[elt-2::TY1::EGFP::3xFLAG(92C12) + unc-119(+)]*	This study
JR4368	*ruIs37 [myo-2p::GFP + unc-119(+)] III; end-1(ok558) end-3(ok1448) V; irEx568[K10F6, F58E10, pAS152(sur-5::dsRed)]*	This study
JR2488	*him-8(e1489) IV; end-1(ox134) V*	This study
JR3341	*elt-7(tm840) end-1(ox134)/ dpy-11(e224) unc-76(e911) V*	This study

JR3342	*elt-7(tm840) end-3(ok1448) V*	This study
JR4037	*elt-2(ca15) X; ijIs12[dpy-7p::GFP::lacZ, unc-76(+)]; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4063	*him-8(e1489) irSi24[opt-2p::mCherry::H2B, Cb-unc-119(+)] IV; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4097	*elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V*	This study
JR4194	*jcIs1[ajm-1p::AJM-1::GFP + unc-29(+) + rol-6(su1006)] IV; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V*	This study
JR4195	*elt-7 (tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4200	*him-8(e1489) irSi24 [opt-2p::mCherry::H2B, Cb-unc-119(+)] IV; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4304	*elt-7 (tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; wEx1751[cdf-1p::CDF-1::GFP + rol-6(su1006)]*	This study
JR4313	*elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; gaEx290[elt-2::TY1::EGFP::3xFLAG(92C12) + unc-119(+)]*	This study
JR4364	*ruIs37 [myo-2p::GFP + unc-119(+)] III; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4367	*ruIs37 [myo-2p::GFP + unc-119(+)] III; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V*	This study
MS880	*dpy-11(e224) end-1(ok558) V; elt-2(ca15) X;*	This study

	irEx404[unc-119::CFP, elt-2(+)]	
JR4407	*jcIs1[ajm-1p::AJM-1::GFP + unc-29(+) + rol-6(su1006)] IV; end-1(ox134) V*	This study
MS900	*end-3(ok1448) V; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4412	*jyIs13 [act-5p::GFP::ACT-5 + rol-6(su1006)] II; elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V*	This study
JR4413	*jyIs13 [act-5p::GFP::ACT-5 + rol-6(su1006)] II; end-1(ox134) V*	This study
JR4439	*him-8(e1489) IV; cuIs1[ceh-22p::GFP + rol-6(su1006] end-1(ox134) V; elt-2(ca15); irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4446	*elt-7(tm840) end-1(ox134)/ elt-7(tm840) tmC12[egl-9(tmIs1197)] V; kcIs6[ifb-2p::IFB-2::CFP]*	This study
JR4198	*end-1(ok558) end-3(ok1448) V; ijIs12[dpy-7p::GFP::lacZ, unc-76(+)]; irEx568[K10F6, F58E10, pAS152(sur-5::dsRed)]*	This study
JR4452	*end-1(ox134) V; elt-2(ca15); ijIs12[dpy-7p::GFP::lacZ, unc-76(+)]; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR3315	*elt-2(ca15) X; wIs84[elt-2p::GFP+ rol-6(su1006)]; irEx404[unc-119::CFP, elt-2(+)]*	This study
JR4463	*pha-4(ot946[pha-4::GFP]) V; elt-2(ca15) X; irEx404[unc-119::CFP, elt-2(+)]*	This study

1020

1021   **Supplementary File 1 (.xlsx)**: Model parameters and outputs of the endoderm GRN.

1022   Expression of each factor is determined by the concentration of its activators multiplied by a

1023   coefficient *a*, representing the strength of the inputs. SKN-1 expression follows a square wave in

1024   EMS blastomere. POP-1 expression follows a square wave in the E cell. Feedback coefficients *f*

1025   become nonzero once their respective factors surpass a certain threshold *Φ*.

1026