Elys deficiency constrains Kras-driven tumour burden by amplifying oncogenic stress

Kimberly J Morgan[1,2], Karen Doggett[1,2], Fan-Suo Geng[1,3], Lachlan Whitehead[2,4], Kelly A Smith[5,6], Benjamin M Hogan[6,7], Cas Simons[6,8], Gregory J Baillie[6], Ramyar Molania[2,9], Anthony T Papenfuss[2,9], Thomas E Hall[6], Elke A Ober[10], Didier Y R Stainier[11], Zhiyuan Gong[12] and Joan K Heath[1,2,*]

[1]Epigenetics and Development Division, Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

[2]Department of Medical Biology, University of Melbourne, Parkville, Victoria 3010, Australia

[3]Genomics and Epigenetics Division, Garvan Institute of Medical Research, Darlinghurst, New South Wales, Australia and St Vincent's Clinical School, Faculty of Medicine, University of New South Wales, New South Wales 2010, Australia

[4]Centre for Dynamic Imaging, Advanced Technology and Biology Division, Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

[5]Department of Physiology, The University of Melbourne, Parkville, Victoria 3010, Australia

[6]Institute for Molecular Biosciences, The University of Queensland, Brisbane, Queensland 4072, Australia

[7]Peter MacCallum Cancer Centre, Melbourne, Victoria 3000, Australia

[8]Murdoch Children's Research Institute, Parkville, Victoria 3052, Australia

[9]Bioinformatics Division, Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

[10]Danish Stem Cell Center, University of Copenhagen, 2200 Copenhagen N, Denmark

[11]Max Planck Institute for Heart and Lung Research, Dept. III – Developmental Genetics

Ludwigstrasse 43, 61231 Bad Nauheim, Germany

[12]Department of Biological Science, National University of Singapore, Singapore

*Corresponding Author: Joan K Heath, Walter and Eliza Hall Institute of Medical Research, 1G Royal

Parade, Parkville, Victoria 3052, Australia. Phone: (+613) 9345-2872

Email: joan.heath@wehi.edu.au

Declaration of interests: The authors declare no potential conflicts of interest

ABSTRACT

The nucleoporin ELYS, encoded by *AHCTF1*, is a large multifunctional protein with essential roles in nuclear pore assembly and mitosis. Using a zebrafish model of hepatocellular carcinoma, in which the expression of an inducible mutant *kras* transgene *(kras^{G12V})* drives hepatocyte-specific hyperplasia and liver enlargement, we show that reducing *ahctf1* gene dosage by 50% markedly shrinks tumour burden, while non-hyperplastic tissues are unaffected. We demonstrate that *ahctf1* heterozygosity impairs nuclear pore formation, mitotic spindle assembly and chromosome segregation, leading to DNA damage and activation of TP53-dependent and independent mechanisms of cell death and cell cycle arrest. This selective vulnerability of cancer cells to mild disruption of Elys function uncovers a novel synthetic lethal interaction between *ahctf1* and oncogenic *kras* that could be exploited therapeutically. Heterozygous expression of both *ahctf1* and *ranbp2*, or treatment of heterozygous *ahctf1* larvae with the nucleocytoplasmic transport inhibitor, Selinexor, completely blocked *kras^{G12V}*-driven hepatocyte hyperplasia, revealing promising avenues for combinatorial treatments.

INTRODUCTION

Synthetic lethality is the term used to describe the death of cells in response to co-existing

disruptions in two different genes, neither of which is lethal alone. The phenomenon has emerged

as a promising tool for cancer drug development[1]. The advantage of the approach lies in its capacity

to induce the death of a vulnerable cell population, such as oncogene-expressing cancer cells, while

healthy cells are unaffected. The approach has been validated in the clinic by the use of

poly(adenosine diphosphate [ADP]-ribose) polymerase (PARP) inhibitors to treat tumours carrying

mutations in the breast cancer susceptibility genes, *BRCA1*/*BRAC2*[2], and its success has fuelled the

search for other clinically relevant pairwise combinations. Particular effort has been directed

towards identifying genes whose partial loss of function confers synthetic lethality in cancer cells

containing hitherto 'undruggable' oncogene targets, such as mutant *KRAS*[3,4]. In this paradigm, the

interacting gene is not mutated in cancer nor oncogenic in its own right; however, its function is

essential to maintain the tumourigenic state, inspiring the concept of non-oncogene addiction[5].

In this paper, we tested whether *AHCTF1* exhibits the properties of a mutant *KRAS* synthetic lethal

interacting gene. *AHCTF1* encodes ELYS, a 252 kDa multidomain nucleoporin that was first

discovered in mouse development where it was shown to be required for the proliferation and

survival of inner mass cells[6]. Since then, ELYS has been studied in many model systems, including in a

zebrafish development mutant (ti262)[7], where we and others showed that homozygous inheritance

of an ENU-induced nonsense mutation in *ahctf1* disrupted nuclear pore formation and caused

catastrophic levels of cell death in highly proliferative cell compartments, such as the intestinal

epithelium, while relatively quiescent tissues survived and remained healthy[8,9].

That ELYS is selectively required by proliferating cells is not unexpected given the multiple roles it

plays in the cell cycle (Fig. 1). As well as being a structural component of nuclear pore complexes

(NPCs), ELYS is essential for their formation at the end of mitosis[10-12]. NPCs are huge (110 MDa)

multi-subunit complexes comprising at least 30 different nucleoporins (Nups) in an octameric

array[13]. They form cylindrical channels in the nuclear envelope (NE) and regulate nucleocytoplasmic

transport and intracellular localisation of large (>40 kDa) molecules. In the absence of ELYS, the NE is

re-built at the end of mitosis but this occurs without NPC formation, precluding mRNAs and large

proteins from moving in and out of the nucleus. By providing a platform for NPC re-assembly at the

end of mitosis (Fig. 1a,b), ELYS restores nucleocytoplasmic trafficking, and thereby facilitates entry

into the nucleus of proteins such as G1 and S phase cyclin-dependent kinases that are required for

DNA replication and cell cycle progression (Fig. 1c). Also in G1/S phase, ELYS regulates transcription

and DNA replication by reversible binding to components of the SWI/SNF 80 chromatin remodeller

complex, PBAP, and the DNA replication origin licensing system, MCM2-7, respectively (Fig. 1d-

f)[11,12,14,15].

At the end of G2, the cell has grown to a critical size and is prepared for M phase (Fig. 1g). The NE

breaks down (NEBD) and NPCs are disassembled (Fig. 1h). Whereas most Nups disperse throughout

the cytoplasm and the endoplasmic reticulum (ER)[16,17], ELYS, together with components of the

Nup107-160 complex and RANBP2, play an important role in prophase by binding to the

kinetochores of recently replicated sister chromatids where they contribute to microtubule

recruitment and spindle assembly (Fig. 1i)[18-20]. At telophase, when the chromatids are fully

separated, the spindle and kinetochores dissolve and ELYS becomes tethered to thousands of sites

on chromatin through its C-terminal chromatin binding region where it promotes chromatin

decondensation[15,21]. In G1, the N-terminal half of ELYS provides a scaffold to bind the Nup107-160

complex and other ER-associated Nups, thereby coupling the initiation of NPC assembly with

euchromatin formation, restoration of nucleocytoplasmic transport and transcription[11,22,23].

Given this plethora of functions throughout the cell cycle, it is not surprising that ELYS is an Achilles'

heel of highly proliferative cells[8,9]. To test whether this vulnerability extrapolates to cancer cells *in*

vivo, we employed a highly tractable zebrafish model of hepatocellular carcinoma (HCC) in which

hepatocyte hyperplasia and liver enlargement are driven by an inducible, hepatocyte-specific *EGFP-*

kras^{G12V} transgene[24]. In this study, we used two-photon microscopy to accurately quantitate liver

volume in the presence and absence of a single nonsense mutation in one allele of the *ahctf1*

locus[8,9]. Remarkably, we observed a pronounced reduction in tumour burden in response to this

mild (50%) reduction in *ahctf1* gene dosage, while the rest of the animal was unaffected. We further

show that this selective vulnerability of oncogenic *kras*[G12V]-expressing liver cancer cells to *ahctf1*

heterozygosity possesses the hallmarks of a synthetic lethal interaction that may be exploited

therapeutically.

RESULTS

ahctf1 **heterozygosity reduces tumour burden in a zebrafish** *kras[G12V]***-driven model of**

hepatocellular carcinoma

In the zebrafish *kras[G12V]*-driven HCC model, the doxycycline-induced expression of a single *EGFP-*

kras[G12V] transgene (genotype denoted *TO(kras[G12V])[T/+]* between 2-7 days post-fertilisation (dpf) leads

to the accumulation of a constitutively active, EGFP-tagged oncogenic form of Kras specifically in

hepatocytes[24] (Fig. 2a). This forced expression of *EGFP-kras[G12V]* causes hepatocyte hyperplasia and a

substantial increase in liver volume that can be quantified by two-photon microscopy. To investigate

the role of Elys in a cancer setting, we introduced a mutant *ahctf1* allele (containing a nonsense

mutation at codon 1319) from a zebrafish development mutant (*flo[ti262]*)[7] into the genome of this

model. This resulted in a 57% reduction in *ahctf1* mRNA expression in *ahctf1[+/-]* larvae at 7 dpf (Fig.

2b), indicating that the premature codon encoded by the nonsense mutation triggers nonsense

mediated decay rather than expression of a truncated Elys protein.

To determine the effect of *ahctf1* heterozygosity (and other treatments) on normal liver cells, we

used a control transgenic line, denoted *2-CLiP* (2-Colour Liver Pancreas), in which hepatocytes

express dsRed fluorescence but no oncogenic transgene[25]. We found that the mean liver volume in 7

dpf wildtype zebrafish larvae is $1.95 \times 10^6 \pm 4.99 \times 10^4$ μm^3 (Fig. 2c,d). *ahctf1* heterozygosity had no

significant impact on liver volume in this model, consistent with previous observations that *ahctf1*

heterozygotes develop normally, reach sexual maturity, and exhibit a normal lifespan[9].

Turning to the *TO(kras[G12V])* model, we found that induced expression of oncogenic Kras[G12V] led to

liver hyperplasia and a striking (4.1-fold) increase in liver volume to $7.97 \times 10^6 \pm 1.21 \times 10^5$ μm^3. This

amounted to an increment in liver volume of approximately 6.02×10^6 μm^3 over the course of 5 d,

demonstrating the potency of the oncogenic *kras[G12V]* signal. Remarkably, the excess liver volume

was reduced by 35% in livers of *ahctf1[+/-];TO(kras[G12V])[T/+]* larvae (down to $5.92 \times 10^6 \pm 8.83 \times 10^4$ μm^3).

These data demonstrate that a mild (50%) reduction in *ahctf1* gene dosage is sufficient to restrict

tumour burden in livers expressing oncogenic $kras^{G12V}$, while having no effect on normal livers. These

results demonstrate that reduced *ahctf1* expression is a selective vulnerability of cancer cells and

identify a novel synthetic lethal interaction between *ahctf1* and $kras^{G12V}$.

Robust and persistent overexpression of RAS oncoproteins is frequently associated with oncogene-

induced stress comprising defects in DNA replication, DNA damage and genome instability. In the

presence of wildtype TP53, this damage is limited by activation of TP53 transcription pathways that

induce cell cycle arrest, senescence and apoptosis. To determine whether expression of $kras^{G12V}$

causes oncogene-induced stress and Tp53 activation in our model, we measured the levels of Tp53

protein in pooled lysates of micro-dissected $TO(kras^{G12V})^{+/+}$ and $TO(kras^{G12V})^{T/+}$ livers and the remains

of the larvae (Fig. 2e,f). No Tp53 signal was obtained from non-$kras^{G12V}$-expressing livers or the body

remains after liver removal. We detected a weak Tp53 signal in extracts of $ahctf1^{+/+};TO(kras^{G12V})^{T/+}$

livers, indicating that expression of the $kras^{G12V}$ oncogene alone elicited a weak Tp53 response. In

contrast, a much stronger Tp53 signal (>3.5-fold) was detected in $ahctf1^{+/-};TO(kras^{G12V})^{T/+}$ livers,

demonstrating a more severe level of stress in $ahctf1^{+/-}$ hepatocytes expressing oncogenic $kras^{G12V}$

than occurs in $ahctf1^{+/+}$ hepatocytes.

Loss of Tp53 function enhances $kras^{G12V}$-driven liver enlargement, which is partially alleviated by

***ahctf1* heterozygosity**

To test whether activation of the Tp53 pathway was involved in reducing the volume of $kras^{G12V}$-

expressing livers on a heterozygous *ahctf1* background, we repeated the experiment in the absence

of Tp53 function. Here, we utilised the zebrafish $tp53^{e7}$ or $tp53^{M214K}$ allele, which affects a conserved

amino acid residue within a region of the Tp53 DNA-binding domain corresponding to a mutational

hotspot in human cancer, producing a transactivation-dead Tp53 variant[26]. Abrogating Tp53 function

by homozygous expression of this allele (which we denoted $tp53^{m/m}$) in $ahctf1^{+/+};TO(kras^{G12V})^{T/+}$

larvae supported a further large increase (82%) in excess liver volume ($1.25 \times 10^7 \pm 1.10 \times 10^5$ μm^3)

compared to livers on the $tp53^{+/+}$ background (Fig. 2g,h) (Supplementary Fig. 1-3). These data

indicate that Tp53 function normally restrains tumour growth and liver enlargement in this model.

Nonetheless, the excess liver volume ($9.82 \times 10^6 \pm 1.34 \times 10^5$ μm^3) on the *ahctf1$^{+/-}$*

;tp53$^{m/m}$;TO(kras^{G12V})$^{T/+}$ background was 35% less than on a wildtype *ahctf1* background, indicating

that mildly disrupted *ahctf1* expression can reduce tumour burden, even in the complete absence of

functional Tp53.

159   *ahctf1* heterozygosity triggers cell death in *TO(kras^{G12V})$^{T/+}$* hepatocytes that is partially Tp53-

160   independent

As expression of Tp53 was upregulated in *ahctf1$^{+/-}$;TO(kras^{G12V})$^{T/+}$* livers, we next determined

whether the reduction in liver volume in heterozygous *ahctf1* larvae was due to the induction of a

Tp53-dependent cell death response. To investigate this, we utilised a transgenic line, Tg(*actb2:SEC-*

Hsa.ANXA5-mKate2,cryaa:mCherry)^{uq24rp} (hereafter denoted *Annexin 5-mKate*) that constitutively

expresses a fusion protein comprising Annexin 5 and the far-red fluorophore mKate[27]. This fusion

protein identifies cells undergoing apoptosis by binding to phosphatidylserine that is normally

inaccessible on the inner leaflet of the plasma membrane but is exposed as the membrane breaks

down. This gives rise to discrete fluorescent puncta, which we captured using confocal imaging.

Annexin 5 fluorescent puncta were sparsely distributed (1.5 puncta per 10^{-5} μm^3) throughout the

livers of *ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* larvae demonstrating that expression of *kras^{G12V}* is

associated with low levels of cell death (Fig. 3a,b). In marked contrast, the number of Annexin 5

fluorescent objects was 3.5-fold higher (5.2 puncta per 10^{-5} μm^3) in the livers of *ahctf1$^{+/-}$*

;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ larvae demonstrating that the reduction in tumour burden observed in

response to *ahctf1* heterozygosity is due, at least in part, to an increase in cell death (Fig. 3a,b left

two columns). These results were recapitulated when the cleaved, (active form) of caspase 3 was

used as an alternative marker of apoptosis (Supplementary Fig. 4). That cell death in hyperplastic

hepatocytes is largely dependent on Tp53 function was demonstrated by negligible apoptosis in the

livers of *ahctf1$^{+/+}$;tp53$^{m/m}$;TO(kras^{G12V})$^{T/+}$* larvae (Fig. 3a,b right two columns). Again, the effect of

Tp53 loss was partially alleviated by *ahctf1* heterozygosity. We observed a 17.6-fold increase in the

abundance of Annexin 5 fluorescent puncta in *tp53$^{m/m}$;TO(kras^{G12V})$^{T/+}$* livers on a heterozygous

*ahctf1$^{+/-}$*background, compared to livers on a *ahctf1$^{+/+}$* background.

We showed that the elevated level of Tp53 protein observed in *ahctf1$^{+/-}$;TO(kras^{G12V})$^{T/+}$* livers,

compared to *ahctf1$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers (Fig. 2e,f), was transcriptionally active by the upregulated

(4.7-fold) mRNA expression of a canonical Tp53 target gene, *mdm2* (Fig. 3c). This prompted us to

quantify the mRNA expression levels of a suite of Bcl2 family genes that regulate intrinsic

(mitochondrial) apoptosis (Fig. 3c). Direct Tp53 transcriptional targets, *pmaip1* (aka *noxa*) and *bbc3*

(aka *puma*), were both upregulated by >6.5-fold in *ahctf1$^{+/-}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers compared to

ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ livers. mRNAs encoding the other BH3-only apoptosis effector

proteins, Bim, Bid, Bik and Bad, and the pro-apoptotic executioner protein, Bax were also increased

in *ahctf1$^{+/-}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers compared to *ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers.

Meanwhile, transcripts encoding Bcl2 and Bclxl pro-survival proteins were significantly

downregulated by *ahctf1* heterozygosity. In the absence of Tp53 function, *ahctf1* heterozygosity still

increased the expression of these pro-apoptotic transcripts, albeit less markedly. All these data are

consistent with induction of Tp53 target genes playing a major role in the cell death response to

ahctf1 heterozygosity in *kras^{G12V}*-expressing livers. Notwithstanding this, *ahctf1* heterozygosity still

promotes moderate levels of cell death concomitant with reduced tumour burden, even in the

complete absence of Tp53 function.

ahctf1 heterozygosity restricts DNA replication in *TO(kras^{G12V})$^{T/+}$* hepatocytes

To explore whether the reduction in liver enlargement we observe in *ahctf1$^{+/-}$;TO(kras^{G12V})$^{T/+}$* larvae

also involves impaired cell cycle progression, we used an EdU incorporation assay to identify cells in

S phase (Fig. 4a). We observed 32% EdU-positive cells in the enlarged livers of

ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ larvae and >20% EdU-positive cells (32% reduction) in the livers of

ahctf1$^{+/-}$; tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ larvae (Fig. 4b,c). In Tp53 deficient larvae, liver volume was markedly

increased as expected (Fig. 4b) and there was a doubling in the percentage of EdU-positive cells

(63% compared to 32%) (Fig. 4c). *ahctf1* heterozygosity reduced liver volume in both Tp53 proficient

and deficient larvae (Fig. 4b), consistent with Fig. 2, and reduced the percentage of EdU-positive

cells in both Tp53 proficient (22% compared to 32%) and Tp53 deficient larvae (53% compared to

63%) (Fig. 4c). These data indicate that the overall reduction in liver tumour burden we observe in

response to *ahctf1* heterozygosity is due to reduced numbers of cycling cells and increased numbers

of apoptotic cells.

To investigate this further, we performed RT-qPCR analysis of the negative cell cycle regulators,

cdkn1a and *cdkn2a/b* in the presence and absence of Tp53. We found that *cdkn1a* and *cdkn2a/*b,

encoding p21 and p16, respectively, were upregulated in *ahctf1$^{+/-}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers

compared to *ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers (Fig. 4d). For *cdkn1a*, this response to *ahctf1*

heterozygosity was partially dependent on intact Tp53 function. However, while the level of

cdkn2a/b mRNA expression was increased in response to *ahctf1* heterozygosity in

tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ livers, in this case there was no response to *ahctf1* heterozygosity in

tp53$^{m/m}$;TO(kras^{G12V})$^{T/+}$ livers. This is consistent with *cdkn2a/b* expression being regulated solely by

Tp53. To understand the basis for the beneficial effects on tumour burden by *ahctf1* heterozygosity

in the absence of Tp53, we investigated the impact of *ahctf1* heterozygosity on the expression of

tp53 family members, *tp63* and *tp73*. We found that *tp63* and *tp73* mRNA expression was up-

regulated by 3.0-fold and 2.3-fold, respectively in *ahctf1$^{+/-}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$* livers, compared to

ahctf1$^{+/+}$;tp53$^{+/+}$;TO(kras^{G12V})$^{T/+}$ livers. Interestingly, the expression of *tp63* and *tp73* was upregulated

by almost the same amount in *ahctf1$^{+/+}$;tp53$^{m/m}$;TO(kras^{G12V})$^{T/+}$* livers and this was further enhanced

by *ahctf1* heterozygosity (Fig. 4e). These data suggest that in response to *ahctf1* heterozygosity,

Tp63 and Tp73, like Tp53, may be capable of promoting *cdkn1a* expression to induce cell cycle arrest

even in the absence of functional Tp53. Taken together, the data shown in Figs. 3 and 4 show that

ahctf1 heterozygosity constrains *kras^{G12V}*-driven hepatocyte hyperplasia and liver enlargement by a

mechanism that results in activation of Tp53 family proteins and initiation of transcriptional

programs that induce cell cycle arrest and cell death.

ahctf1 heterozygosity disrupts the abundance of nuclear pore complexes in _TO(kras_^{G12V}_)_$^{T/+}$

hepatocytes.

To determine how _ahctf1_ heterozygosity leads to Tp53 activation, we examined the integrity of

some of the essential processes in which the Elys protein plays a role. As depicted in Fig. 1, Elys fulfils

many distinct functions during the cell cycle. Starting with post-mitotic nuclear pore assembly (Fig.

1a), we used confocal laser-scanning microscopy with Airyscan to look at the abundance and

distribution of NPCs in thick sections of liver (200 μm) stained with an antibody (mAb414) that

recognises FG-repeat Nups (Nup358, 214, 153 and 62) in mature NPCs. Non-_kras_^{G12V} expressing

hepatocytes exhibited fluorescent puncta corresponding to NPCs at the nuclear rim with negligible

staining in the cytoplasm, a pattern that was unaffected by _ahctf1_ genotype (Fig. 5a; left two

columns). Upon induction of expression of the _TO(kras_^{G12V}_)_$^{T/+}$ transgene, fluorescence intensity was

markedly increased in the presence of wildtype _ahctf1_ (Fig 5a, third column) and there was an

increase in the ratio of nuclear:cytoplasmic staining (Fig 5b). This pattern was different in

hepatocytes on a heterozygous _ahctf1_ background. Notably, fluorescence intensity at the nuclear

rim was partially diminished and this was concomitant with the appearance of puncta in the

cytoplasm (Fig. 5a, fourth column; arrows). This apparent reduction in proper NPC assembly resulted

in a 21% reduction in the ratio of nuclear:cytoplasmic fluorescence intensity in liver sections (Fig.

5b).

To determine the abundance of NPCs, we analysed images acquired at the nuclear surface (Fig. 5c).

The pattern and density of fluorescent puncta observed at the nuclear surface of non-_kras_^{G12V}

expressing hepatocytes was not affected by _ahctf1_ genotype (Fig. 5c; left two columns). We found

59% more fluorescent puncta/NPCs at the nuclear surface of hyperplastic hepatocytes expressing

kras^{G12V} in the presence of wildtype _ahctf1_. This _kras_^{G12V}-driven increase in NPCs was reduced by 20%

in response to *ahctf1* heterozygosity (Fig. 5c, d; right two columns). We also observed that induced

kras^{G12V} expression produced a 28% increase in nuclear volume compared to non-*kras^{G12V}* expressing

cells (Fig. 5e), which was reduced to a 13% increase in the context of *ahctf1* heterozygosity.

Together, these data indicate that hyperplastic hepatocytes expressing the *kras^{G12V}* oncogene

contain more abundant nuclear pores and larger nuclei than their non-*kras^{G12V}* expressing

counterparts. While these properties will facilitate nucleocytoplasmic trafficking and be of great

benefit towards meeting the transcriptional and translational needs of rapidly cycling cancerous

cells, we found that they could not be fully sustained on a *ahctf1^{+/-}* background.

***ahctf1* heterozygosity impairs mitotic spindle assembly and chromosome segregation in**

***TO(kras^{G12V})^{T/+}* hepatocytes**

Next, we examined the impact of reduced Elys expression on spindle formation and chromosome

segregation (Fig. 1b). We assessed these features in cryosections of liver using α-tubulin and DAPI to

stain microtubules and chromatin, respectively. Metaphase cells in *ahctf1^{+/+};TO(kras^{G12V})^{T/+}* livers

exhibited normal bipolar spindle formation followed by complete chromosome segregation during

anaphase (Fig. 6a). In contrast, metaphase cells in *ahctf1^{+/-};TO(kras^{G12V})^{T/+}* hepatocytes displayed

abnormal multipolar spindles and misaligned chromosomes (Fig. 6b). Proper chromosome

segregation was disrupted with multiple anaphase bridges formed. While the distribution of cells

observed at different mitotic stages was similar in *ahctf1^{+/+}* and *ahctf1^{+/-}* hepatocytes (Fig. 6c),

mitotic abnormalities were observed in 50% of *ahctf1^{+/-};TO(kras^{G12V})^{T/+}* hepatocytes during

metaphase and anaphase but not at all in *ahctf1^{+/+};TO(kras^{G12V})^{T/+}* hepatocytes (Fig. 6d). Taken

together, these data suggest that hyperplastic hepatocytes require a full complement of *ahctf1*

expression to maintain proper cell division.

***ahctf1* heterozygosity causes DNA damage in *TO(kras^{G12V})^{T/+}* hepatocytes**

So far, our results demonstrate that a 50% decrease in *ahctf1* expression has multiple impacts in

TO(kras^{G12V})^{T/+} hepatocytes, including disrupted NPC assembly, impaired mitotic spindle assembly

and incomplete chromosome segregation, consistent with oncogene-induced stress. As several of

these defects can converge to produce DNA damage, we looked for evidence of DNA damage by

staining cryosections of liver with DAPI and γ-H2AX to mark DNA double-stranded breaks (Fig. 7a,b).

We found that while only 1% of $ahctf1^{+/+};TO(kras^{G12V})^{T/+}$ hepatocyte nuclei were positive for γ-H2AX

(Fig. 7c), 6% of $ahctf1^{+/+};TO(kras^{G12V})^{T/+}$ hepatocyte nuclei were γ-H2AX-positive. Collectively, these

data show that the proper functioning of diverse cellular processes in hyperplastic hepatocytes

depends on wildtype levels of $ahctf1$ expression. We conclude from this that insufficient Elys protein

in rapidly cycling cells produces multiple disruptive events that combine to produce elevated levels

of oncogene-induced stress and DNA damage, in so doing providing the stimulus to activate Tp53

and transcriptional programs that trigger cell cycle arrest and apoptosis.

Combinatorial approaches can completely restrict $kras^{G12V}$-driven liver hyperplasia

Having determined that heterozygous $ahctf1$ mutation is effective in restricting $kras^{G12V}$-driven

hepatocyte hyperplasia and liver enlargement, we investigated the potential of targeting cancer via

another nucleoporin, Ranbp2, in our zebrafish model of HCC. RANBP2 (also known as NUP358) is a

major component of the cytoplasmic filaments of NPCs, where it functions in numerous transport

pathways. Like ELYS, RANBP2 also plays non-canonical roles outside of the NPC, including in mitotic

progression and the maintenance of genome integrity[28-30]. In colon cancer cells with an activated

BRAF signature, RANBP2 silencing reduces microtubule outgrowth from kinetochores, inducing

spindle perturbations and cell death, the frequency of which can be exacerbated by treatment with

microtubule inhibitors[31].

We identified our zebrafish $ranbp2$ mutant in a transgene assisted ENU-mutagenesis screen for

mutations affecting endodermal organ morphogenesis[32] and showed, using whole genome

sequencing and homozygosity mapping[33,34], that the underlying mutation was a nonsense mutation

in $ranbp2$ (Supplementary Fig. 5). Like several other mutants we identified in this screen[8,35,36],

mutant *ranbp2* larvae exhibit morphological deficiencies in proliferative compartments, including

the developing intestinal epithelium, craniofacial complex and eye (Supplementary Fig. 5).

We found that *ranbp2* heterozygosity alone or in combination with *ahctf1* heterozygosity did not

impact on liver volume in *2-CLiP* larvae (Fig. 8a,b). In *TO(kras^{G12V})$^{T/+}$* larvae, liver volume was

increased 3.8-fold, as observed previously, and the excess liver volume was reduced by 38% and 13%

by the introduction of a single mutation in *ahctf1* or *ranbp2*, respectively (Fig. 8b). Remarkably, we

found a striking synergistic benefit of trans heterozygosity in *ranbp2$^{+/-}$;ahctf1$^{+/-}$;TO(kras^{G12V})$^{T/+}$*

larvae, to the extent that liver volume was no longer distinguishable from that of normal *2-CLiP*

livers. These results show that cancer cells are highly susceptible to combinatorial targeting of Nup

function and that it may be possible to exploit this vulnerability to produce highly beneficial

outcomes without impacting negatively on healthy tissues.

To examine this concept in a therapeutic context, we assessed the effect of Selinexor, a selective

inhibitor of nuclear export (SINE) in our *kras^{G12V}*-driven liver hyperplasia model. Selinexor is an XPO1

(exportin) inhibitor with FDA approval for the treatment of refractory or relapsed multiple myeloma

and diffuse large B-cell lymphoma and is in clinical trials for several other cancers, including HCC[37,38].

While we observed no reduction in liver volume in *2-CLiP* larvae exposed to 0.10-2.00 µM Selinexor

from 5 to 7 dpf (Fig. 8c,d) (Supplementary Fig. 6), *ahctf1$^{+/+}$;TO(kras^{G12V})$^{T/+}$* larvae exhibited a dose-

dependent reduction in liver volume when exposed to Selinexor. Strikingly, liver volume was

reduced even further in the context of *ahctf1* heterozygosity, such that at a 1.00 µM dose of

Selinexor, *kras^{G12V}*-driven hepatocyte hyperplasia was completely blocked and liver volume was

indistinguishable from that of normal, non-hyperplastic *2-CLiP* larvae (Fig. 8c,d). These results

indicate that XPO1 inhibition is an effective suppressor of growth and proliferation in our model of

mutant Kras-driven HCC, and that its effect is even more impressive when combined with a mild

reduction in the expression of the nucleoporin, Elys.

We next examined whether there was any clinical rationale for developing Nup inhibitors for the

purpose of cancer treatment. We analysed the survival data of HCC patients (Fig. 8e) collated in the

cBioPortal for Cancer Genomics[39,40] in the context of Nup gene expression (Supplementary Table 3).

Median overall survival of patients with high mRNA expression of components of the NUP107-160

complex, including ELYS is 21.70 months, whereas the survival of patients with low expression of

mRNAs encoding NUP107-160 components is markedly extended to 81.73 months. To be confident

of the integrity of this striking result, we investigated whether the gene expression values in the

TCGA liver hepatocellular carcinoma (LIHC) dataset were subject to batch effects and unwanted

variation, noting that the 372 samples in the LIHC dataset were profiled across 19 sequencing plates.

To do this, we performed ANOVAs to assess the effects of individual sequencing plates on the

expression of genes encoding components of the NUP107-160 complex (Supplementary Fig. 7).

These analyses revealed that differences in gene expression observed between samples were largely

unaffected by unwanted variation between sequencing plates.

a ELYS binds to chromatin and initiates postmitotic NPC assembly

i ELYS mediates microtubule attachment and spindle formation

h ELYS relocalises to kinetochores upon nuclear envelope breakdown and NPC disassembly

b ELYS coordinates NPC reassembly with chromatin decondensation

c Restoration of nucleocytoplasmic transport enables nuclear import of CDKs permitting cell cycle progression

g ELYS interacts with the genome to promote euchromatin formation and mediate transcription

d ELYS interacts with the genome to promote euchromatin formation and mediate transcription

e ELYS interacts with the replication licensing system enabling S phase progression once replication origin licensing is complete

f DNA synthesis

M

G1

G2

S

Cytoplasm

Nucleus

Elys

Elys

Fig. 1 Elys plays critical roles at several stages of the cell cycle.

Elys is a central player in multiple diverse cellular processes far beyond its canonical role in **a, b.**

nuclear pore re-assembly at the end of mitosis. **c.** Elys also restores nucleocytoplasmic trafficking

allowing import of cyclin-dependent kinases (CDKs) required for cell cycle progression. **d.** Elys plays

an active role in genome regulation through its direct interactions with components of large

molecular machineries including chromatin remodellers such as PBAP that favour transcription and

e, f. the MCM2-7 components of the DNA replication origin licensing system. **g.** After cell growth in

G2, **h.** the nuclear envelope breaks down and NPCs are disassembled upon entry into mitosis. Elys

relocalises to kinetochores where **i.** it contributes to microtubule attachment and spindle formation.

Fig. 2 *ahctf1* heterozygosity reduces liver volume in a zebrafish *kras^G12V*-driven model of

hepatocellular carcinoma in both Tp53 proficient and deficient larvae.

a. Protocol for doxycycline induction of *TO(kras^G12V)^T/+* expression in developing zebrafish larvae. **b.**

RT-qPCR analysis of *ahctf1* mRNA levels in pooled micro-dissected livers (n=3 biological replicates). **c.**

Representative three-dimensional reconstructions of *2-CLiP* and *TO(kras^G12V)^T/+* livers of the indicated

ahctf1 genotype. Scale bar 25 μm. **d.** Impact of *ahctf1* heterozygosity on liver volume in *2-CLiP* and

TO(kras^{G12V})$^{T/+}$ larvae (n≥20). **e.** Representative western blot of Tp53 protein signals in lysates of

TO(kras^{G12V})$^{T/+}$ larvae of the indicated *ahctf1* genotype. **f.** Quantification of Tp53 protein levels

normalised by reference to the Gapdh loading control (n=3 independent experiments). **g.**

Representative three-dimensional reconstructions of *TO(kras^{G12V})$^{T/+}$* livers of the indicated *ahctf1* and

tp53 genotypes. Scale bar 25 μm. **h.** Impact of *ahctf1* heterozygosity and homozygous *tp53* mutation

on liver volume in *2-CLiP* and *TO(kras^{G12V})$^{T/+}$* larvae (n≥20). Data are expressed as mean ± SEM.

Significance was calculated using a Student's t-test or one-way ANOVA with Tukey's multiple

comparisons test.

Fig. 3 *ahctf1* heterozygosity increases cell death in *TO(kras^{G12V})^{T/+}* hepatocytes in both Tp53 proficient and deficient larvae.

a. Representative maximum intensity projection images of Annexin 5-mKate fluorescence (white puncta), indicating cells undergoing apoptosis in *TO(kras^{G12V})^{T/+}* livers of the indicated *ahctf1* and *tp53* genotypes. Scale bar 25 μm. **b.** Quantification of the density of Annexin 5 fluorescent foci in *TO(kras^{G12V})^{T/+}* livers of the indicated *ahctf1* and *tp53* genotypes (n≥11). **c.** RT-qPCR analysis of the specified mRNAs in *TO(kras^{G12V})^{T/+}* micro-dissected livers of the indicated *ahctf1* and *tp53* genotypes

376 (n=3 biological replicates). Data are expressed as mean ± SEM. Significance was calculated using a

377 one-way ANOVA with Tukey's multiple comparisons test.

Fig. 4 *ahctf1* heterozygosity restricts DNA replication in *TO(krasG12V)T/+* hepatocytes.

a. Representative maximum intensity projection images of EdU incorporation (white puncta) into

TO(krasG12V)T/+ livers of the indicated *ahctf1* and *tp53* genotypes. Scale bar 25 μm. **b.** Impact of

ahctf1 heterozygosity and homozygous *tp53* mutation on liver volume in *2-CLiP* and *TO(krasG12V)T/+*

larvae (n≥17). **c.** Quantification of the percentage of EdU positive nuclei per Hoechst 43332 positive

nuclei (n≥25). **d.** RT-qPCR analysis of mRNA expression of the cell cycle regulators, *cdkn1a* and

386 *cdkn2a/b* and **e.** *tp63* and *tp73* in *TO(kras^{G12V})^{T/+}* micro-dissected livers of the indicated *ahctf1* and

387 *tp53* genotypes (n=3 biological replicates). Data are expressed as mean ± SEM. Significance was

388 calculated using a one-way ANOVA with Tukey's multiple comparisons test.

Fig. 5 *ahctf1* heterozygosity disrupts nuclear pore complexes in *TO(kras^{G12V})^{T/+}* hepatocytes.

a. Representative Airyscan imaging of liver sections stained with mAb414 (white) marking FG-Nups,

Hoechst 43332 (cyan) marking DNA, and rhodamine phalloidin (magenta) marking the F-actin

cytoskeleton in non-*TO(kras^{G12V})*-expressing cells or EGFP-Kras^{G12V} (magenta) marking the cell

membrane in *TO(kras^{G12V})*-expressing cells of the indicated *ahctf1* and *TO(kras^{G12V})* genotypes.

Arrows indicate mAb414/FG-nucleoporin staining in the cytoplasm. Scale bar 2 µm. **b.** Quantification

of mean nuclear/cytoplasmic fluorescence intensity of mAb414 staining after 3D segmentation and

morphological filtering of nuclear and cytoplasmic areas (n>18). **c.** Representative Airyscan images of

mAb414 staining at the nuclear surface of sections of the indicated *ahctf1* and *TO(kras^{G12V})*

genotype. Scale bar 1 µm. **d.** Quantification of nuclear pore density (n≥25). **e.** Quantification of

nuclear volume (n≥25). Data are expressed as mean ± SEM. Significance was calculated using a one-

way ANOVA with Tukey's multiple comparisons test.

Fig. 6 *ahctf1* **heterozygosity impairs mitotic spindle assembly and chromosome segregation in**

TO(kras^G12V)^T/+ hepatocytes.

a. Representative Airyscan imaging of liver cryosections stained with α-tubulin antibody (magenta)

marking spindle microtubules and DAPI (cyan) marking DNA in mitotic cells of *TO(kras^G12V)^T/+* larvae

on a wildtype *ahctf1^+/+* background. **b.** Mitotic cells in liver cryosections of *TO(kras^G12V)^T/+* larvae on a

410   heterozygous *ahctf1*^{+/-} background exhibit multiple defects, including multipolar spindles, misaligned

411   chromosomes and anaphase bridges (arrows). Scale bar 2 μm. **c.** Distribution of cells observed at

412   different mitotic stages (n=92 livers, 326 mitotic cells) **d.** Quantification of the percentage of mitotic

413   hepatocytes exhibiting an aberrant phenotype (n=14-57).

Fig. 7 *ahctf1* heterozygosity leads to accumulation of DNA double-strand breaks in *TO(kras^{G12V})^{T/+}*

hepatocytes.

a. Representative Airyscan imaging of cryosections of liver from *ahctf1^{+/+};TO(kras^{G12V})^{T/+}* larvae

stained with γ-H2AX antibody (white) marking DNA double-strand breaks, DAPI (cyan) marking DNA

and EGFP-Kras^{G12V} (magenta) marking the cell membrane. Scale bar 5 µm. **a'.** Inset of γ-H2AX positive

nuclei in *ahctf1^{+/+};TO(kras^{G12V})^{T/+}* hepatocytes. Scale bar 2 µm. **b.** Representative images of

cryosections of liver from *ahctf1^{+/-};TO(kras^{G12V})^{T/+}* larvae. Scale bar 5 µm. **b'.** Inset of γ-H2AX positive

nuclei in cryosections of liver from *ahctf1^{+/-};TO(kras^{G12V})^{T/+}* larvae. Scale bar 2 µm. **c.** Quantification of

the percentage of hepatocytes positive for γ-H2AX (n≥31). Data are expressed as mean ± SEM.

Significance was calculated using a Student's t-test.

Fig. 8 Combined approaches completely restrict *kras*[G12V]-driven liver hyperplasia.

a. Representative three-dimensional reconstructions of livers from *2-CLiP* and *TO(kras*[G12V]*)*[T/+] larvae

of the indicated *ahctf1* and *ranbp2* genotypes. Scale bar 25 μm. **b.** Impact of *ahctf1* heterozygosity

and *ranbp2* heterozygosity on liver volume in *2-CLiP* and *TO(kras*[G12V]*)*[T/+] larvae (n≥30). Significance

was calculated using a one-way ANOVA with Tukey's multiple comparisons test. **c.** Representative

three-dimensional reconstructions of *TO(kras^{G12V})$^{T/+}$* livers of the indicated *ahctf1* genotype treated

with DMSO or 2.00 µM Selinexor from 5-7 dpf. **d.** Dose-dependent impact of Selinexor treatment

(0.10-2.00 µM) on liver volume in *2-CLiP* (grey), *ahctf1$^{+/+}$;TO(kras^{G12V})$^{T/+}$* (blue) and *ahctf1$^{+/-}$*

;TO(kras^{G12V})$^{T/+}$ (purple) larvae (n≥20). Data are expressed as mean ± SEM. Significance was

calculated by linear regression analysis. **e.** Overall survival of HCC patients with mRNA expression z-

scores >2 for one or more Nup107-160 complex components (red line; 63 cases); median overall

survival 21.70 months. Overall survival of HCC patients with mRNA expression z-scores <-2 for one or

more Nup107-160 complex components (blue line; 45 cases); median overall survival 81.73 months.

Data from the TCGA LIHC dataset (total 372 samples) available in the cBioPortal for Cancer

Genomics. **f.** Schematic depiction of two of the processes disrupted by mild depletion of Elys protein

culminating in **g.** DNA damage**,** activation of Tp53 transcription programs and **h.** cell death and cell

cycle arrest of hyperproliferative Kras^{G12V}-expressing cancer cells.

DISCUSSION

In this paper we show that *ahctf1* heterozygosity markedly reduces tumour burden in a genetically

engineered zebrafish model of HCC. In this model of mutant *kras*-driven hepatocyte hyperplasia,

ahctf1 heterozygosity produces a reduction in the density of nuclear pores, aberrant chromatid

separation during mitosis (Fig. 8f) and accumulation of DNA damage (Fig. 8g). We propose that these

events combine to intensify the oncogene-induced stress sensed by cancer cells induced to hyper-

proliferate by expression of the Kras^{G12V} oncoprotein. We also show that the heightened levels of

oncogene-induced stress caused by mild depletion of Elys protein provide a powerful stimulus to

Tp53 activation. We observed increased abundance of Tp53 protein and enhanced transcription of

Tp53 target genes, including *pmaip1* and *bbc3*, encoding two pro-apoptotic BH3-only Bcl2 family

proteins (aka Noxa and Puma, respectively), and *cdkn1a* and *cdkn2a/b*, encoding the cell cycle arrest

proteins, Cdkn1a/p21 and Cdkn2a/b/p16(Ink4a)/p14(Arf) (Fig 8h). Induced expression of these

proteins is typically linked to increased apoptosis and cell cycle arrest. We found no evidence that

mild depletion of Elys protein activated this sequence of events in unstressed, non-*kras^{G12V}*-

expressing cells.

Our results suggest that the promising therapeutic effect of inhibiting Elys function could be

diminished in cancer cells lacking wild-type Tp53 protein. This is a clinically relevant issue since

mutations in *TP53*, or amplification/overexpression of its negative regulators *MDM2*/*MDM4*, occur

in 30% of HCC cases[41]. To investigate this further, we conducted experiments on a homozygous *tp53*

mutant background. We found that *ahctf1* heterozygosity still produced a 24% reduction in tumour

burden and induced cell death in the complete absence of Tp53 function, leading us to consider

whether the loss of Tp53 function was compensated for by increased *tp63* and/or *tp73* expression.

These genes encode Tp53 family proteins that share considerable structural homology with Tp53,

particularly within the DNA binding domain, and can activate common and distinct sets of target

genes to produce cell cycle arrest, senescence and apoptosis[42,43]. For example, in response to DNA

damage, Tp63 induces senescence and apoptosis in the same way as Tp53, via transcriptional

induction of *cdkn1a*, *bbc3* and *pmaip1*[44]. We found that *ahctf1* heterozygosity upregulated *tp63* and

tp73 expression by 3.0-fold and 2.3-fold, respectively, and that this was further enhanced by *tp53*

mutation. These results suggest a mechanism through which *ahctf1* heterozygosity achieves a

reduction in tumour burden in the absence of Tp53 function. Since *TP63* and *TP73* are rarely

mutated in cancer[45], it is plausible that inhibition of ELYS function may still produce a beneficial

effect on tumour growth in patients harbouring *TP53* mutations.

Dysregulation of nucleocytoplasmic transport is a common feature in a broad spectrum of cancers,

usually arising from altered expression of nuclear pore components and/or nuclear transport

receptors[46]. For example, overexpression of POM121, a transmembrane NPC component, enhances

the nuclear import of the pro-proliferation transcription factors, E2F1, MYC and AR, and increases

the therapeutic resistance of prostate cancer[47]. Similarly, the nuclear transport receptor XPO1 is

frequently overexpressed in cancers, including HCC, and in this case leads to mis-localisation and

inactivation of tumour suppressor proteins[38,48]. As a result, numerous selective inhibitors of nuclear

export (SINEs) have been developed, including Selinexor (KPT-330), which has advanced through

clinical trials and received FDA approval for the treatment of relapsed or refractory multiple

myeloma and diffuse large B-cell lymphoma[37]. We found that Selinexor treatment in our zebrafish

HCC model produced a dose-dependent reduction in tumour burden. Moreover, this therapeutic

effect was strongly augmented by *ahctf1* heterozygosity, suggesting that new drugs targeting the

ELYS protein could be combined effectively with drugs that target nucleocytoplasmic transport. We

also exploited the genetic tractability of our zebrafish HCC model to show that trans heterozygosity

of *ahctf1*, *ranbp2* delivers a synergistic reduction in liver volume/tumour burden, bringing it back to

the volume of non-hyperplastic livers. Notably, non-*TO(kras^{G12V})* livers were completely unaffected

by *ahctf1*, *ranbp2* trans heterozygosity, indicating a three-pronged synthetic lethal interaction in

cancer cells that may be possible to recapitulate with combinatorial drug treatments.

There are several reasons why we think our *TO(kras^{G12V})* zebrafish HCC model provides a clinically

relevant *in vivo* platform for the study of human cancer and the discovery of new therapeutic

strategies[24]. Firstly, RAS/RAF/MAPK signalling is almost always hyper-activated in human HCC[49].

Secondly, the progressive accumulation of histopathologic features including irregular nuclei,

cytoplasmic vacuolation and increased vascularisation that are characteristic of tumour progression

in HCC are reproduced in the zebrafish model[50]. Thirdly, comparative transcriptomic analysis reveals

that the gene expression profile exhibited by oncogenic $kras^{G12V}$ hepatocytes isolated from larval

zebrafish strongly resembles that of early-stage human HCC, with elevated expression of

RAS/RAF/MAPK target genes such as *FGFR4, ETV4, EPHA2, DUSP6* and *SPRY* and DNA damage

response genes *GADD45B, CCND1* and *H2AX1*[51].

Our results compliment a growing body of evidence that carcinogenesis places persistently high

demands on essential cellular genes, including those required for proper nuclear pore function,

creating a vulnerability that may be therapeutically targeted without causing adverse effects on

normal tissue[52,53]. For example, Sakuma *et al.* used siRNAs targeting 28 out of 32 Nup genes, to

reduce the abundance of mature NPCs in the human melanoma-derived cell line, A375[52]. Focusing

on two Nups exhibiting a severe reduction in NPC density, this study showed that NUP160 and

NUP93 are essential for NPC assembly and that inhibiting this process results in cell death.

Moreover, they found that cancer cells exhibited heightened susceptibility to inhibition of NPC

assembly compared to human pulmonary fibroblasts and a normal cell line derived from retinal

pigmental epithelium (RPE1), which instead underwent reversible cell cycle arrest. The differential

sensitivity between cancer cells and normal cells to NPC assembly inhibition is consistent with

reports that quiescent cells in normal tissues exist for many years with the same set of assembled

NPCs and generally maintain low levels of Nup expression[54,55].

The dynamic and diverse functions of ELYS require its binding to Nups and many other proteins,

including components of kinetochore and replicative licensing complexes, as well as binding to

chromatin in various states of condensation. The molecular topology of the ELYS protein and the

structural basis for its diverse protein-protein and protein-chromatin interactions are hitherto poorly

defined. Crystal structures of ELYS have shown that the β-propeller and α-helical domains in the N-

terminal half of the ELYS protein bind to NUP160, and cryo-electron microscopy reveals the

association of the ELYS C-terminal peptide with nucleosomes[21,56]. To advance our molecular

understanding of the extent to which discrete domains within the ELYS protein contribute to tumour

suppression, further high-resolution structural characterization is required. Hopefully, this will also

reveal opportunities to inhibit distinct regions on the protein for the purpose of disabling ELYS

function therapeutically. However, even if the topology of ELYS lacked features that would facilitate

high affinity binding with small drug-like molecules, as was suggested for other Nups with scaffolding

properties[52], rapidly emerging technological advances in drug development are likely to provide

alternative strategies to target the ELYS protein. For example, carefully controlled proteasomal

degradation of proteins that participate in the oncogenic process with proteolysis targeting chimeras

(PROTACs) is a rapidly advancing field in cancer therapy[57]. This approach depends on the availability

of a ligand or 'warhead' to direct the PROTAC to the targeted protein, but this need not be a

druggable site of protein-protein or protein-chromatin interaction, as is the case for small molecules

designed to disrupt the function of the protein. In parallel, emerging small molecule RNA-targeting

technology is on course to permit direct modulation of the abundance of specific RNA transcripts for

a variety of clinical purposes[58].

In summary, we assert that our *in vivo* cancer studies provide a strong and feasible rationale for

targeting ELYS for the treatment of a broad spectrum of cancers. Moreover, our observations with

RANBP2, indicate that our findings may be extrapolatable to other multi-functional Nups and

provide a foundation for effective combinatorial treatments.

MATERIALS AND METHODS

Zebrafish maintenance and strains

Zebrafish were maintained at 28°C on a 12 h light/12 h dark cycle. The mutant lines *ahctf1^ti262* and

tp53^M214K/M214K (also known as *tp53^e7/e7* and referred to herein as *tp53^m/m*) have been described

previously[7,26]. Tg(*fabp10:dsRed, ela3l:GFP*)[gz12], hereafter referred to as *2-CLiP*, expresses dsRed in the

liver and GFP in the exocrine pancreas but carries no oncogenic transgenes or mutations[25]. The

Tg(*fabp10:rtTA2s-M2;TRE2:EGFP-kras^G12V*)[gz32] line referred to as TO(*kras^G12V*) and the cell death

reporter line, Tg(*actb2:SEC-Hsa.ANXA5-mKate2,cryaa:mCherry*)[uq24rp] were described previously[24,27].

The *ranbp2^s452* mutant line was generated in the Liver^plus ENU mutagenesis screen[32], and its genetic

and morphological characterization is presented in this paper.

Inducing hepatocyte hyperplasia in transgenic zebrafish

To induce mutant *kras^G12V* expression, *TO(kras^G12V)* embryos were treated with 20 μg/mL doxycycline

(Sigma, #D9891) at 2 dpf in E3 medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl₂, 0.33 mM MgSO₄)

with 0.003% 1-Phenyl-2-thiourea (PTU; Sigma, #P7629) to suppress pigmentation. E3 medium was

changed at 5 dpf and fresh doxycycline (final conc. 20 μg/mL) added. For the Selinexor experiments,

the drug (0.10-2.00 mM) (Karyopharm) or 0.001% DMSO (vehicle control) was added at 5 dpf.

Morphological and molecular analyses were performed at 7 dpf. To quantitate liver volume, larvae

were anaesthetized with benzocaine (200 mg/L) and mounted in 1% agarose. Image acquisition was

performed using an Olympus FVMPE- RS multiphoton microscope with a 25x objective and Olympus

FV30-SW software. Excitation wavelengths for GFP and RFP were 840 nm and 1100 nm, respectively.

For volumetric analysis of whole livers, z-stacks with step-size 2 μm, were imported into ImageJ or

Imaris software.

Genotyping

Genomic DNA (gDNA) was extracted from whole zebrafish larvae by incubation at 95°C in 50 μL of 50

mM sodium hydroxide (NaOH) for 20 min, followed by neutralization with 5 μL of 1 M Tris-HCl (pH

8.0). Primer sequences for genotyping are listed in Supplementary Table 1.

mRNA expression analysis

Total RNA was extracted from independent pools of micro-dissected zebrafish livers using the

RNeasy Micro Kit (QIAGEN, #74004). RNA integrity was assessed by a High Sensitivity RNA

ScreenTape assay (Agilent, #5067-5579) on a 2200 TapeStation. cDNA was generated from 1-10 μg

RNA using the Superscript III First Strand Synthesis System (Invitrogen, #18080051) and oligo(dT)

priming according to the manufacturer's instructions. RT-quantitative PCR (RT-qPCR) was performed

using a SensiMix SYBR kit (Bioline, #QT605-05) on an Applied Biosystems ViiATM7 Real-Time PCR

machine. Expression data were normalized by reference to *hrpt1*, *b2m* and *tbp*. LinRegPCR V11.0

was used for baseline correction, PCR efficiency calculation and transcript quantification analysis[59].

Relative expression levels were calculated by the $2^{-\Delta\Delta Ct}$ method and all results were expressed as the

mean ± SEM of three independent pools of biological replicates. Primer sequences are listed in

Supplementary Table 2.

Western blot analysis

Pooled micro-dissected zebrafish livers were lysed in RIPA buffer (20mM Hepes, pH 7.9, 150mM

NaCl, 1mM MgCl2, 1% NP40, 10mM NaF, 0.2mM Na3VO4, 10mM β-glycerol phosphate)

supplemented with cOmplete Proteinase Inhibitor (Roche, #11836170001) and PhosTOP

phosphatase inhibitors (Roche, #04906837001). Samples were incubated for 30 min on ice and the

extracts cleared by centrifugation at 13,000 rpm for 20 min at 4°C. The protein concentration of

samples was determined by BCA protein assay (Thermo Fisher Scientific, #23227). 25 μg of protein

per lane were resolved on NuPAGE Novex Bis-Tris 4-12% polyacrylamide gels (Invitrogen,

#NP0321BOX) and transferred onto nitrocellulose blotting membranes (Amersham Protran,

#10600003). Membranes were blocked with 5% BSA in PBS for 1 h at RT and incubated with primary

antibodies 1:500 Anti-p53 (9.1) mouse mAb (Abcam, #ab77813) overnight at 4°C and 1:1000 Anti-

GAPDH (14C10) Rabbit mAb (Cell Signalling Technology, #2118) for 1 h at RT. Secondary antibodies:

Goat anti-mouse HRP (Dako, #P0447) and Goat anti-Rabbit HRP (Dako, #P0448), were used at 1:5000

and incubated with membranes for 1 h at RT. Signals were developed using Amersham ECL Western

Blotting Detection Kit (Cytiva, # RPN2108) and imaged on a Chemidoc Touch (Biorad). Relative

protein quantitation was calculated based on normalized integrated intensity.

Cell death analysis

601   To assess apoptosis, 7 dpf *TO(kras^{G12V});Annexin 5-mKate* zebrafish larvae were fixed in 4%

602   paraformaldehyde (PFA) overnight at 4^{o}C and livers isolated by micro-dissection. Image acquisition

603   was performed using a Zeiss LSM 880 microscope with a 20x objective and ZEN software. Excitation

604   wavelengths for mKate and GFP were 560 nm and 900 nm, respectively. Liver volume was quantified

605   and 3D segmentation of Annexin 5-mKate signals were performed in ImageJ.

Cell cycle analysis

EdU incorporation was used to assess the percentage of cells in S phase of the cell cycle. Briefly, live

zebrafish larvae (7 dpf) were incubated at 28^{o}C in 2 mM EdU (Invitrogen, #C10340) in E3 medium for

2 h followed by a further incubation in fresh E3 medium for 1 h. Larvae were euthanised using

benzocaine (1000 mg/L; Sigma, #PHR1158) prior to removal of the liver by micro-dissection. EdU

labelling was carried out using the Click-iT Edu Alexa Fluor 647 (AF647) imaging kit (Invitrogen,

#C10340), according to the manufacturer's instructions. The livers were co-stained with Hoechst

33342 (Thermofisher, #62249). Image acquisition was performed using an Olympus FVMPE-RS

multiphoton microscope with excitation wavelengths of 950 nm for Hoechst 33342 dye and 1160 nm

for AF647 (Thermofisher, #A21235). The number of Hoechst 33342 and EdU positive cells was

quantified using Arivis Vision4D software.

Nuclear pore analysis

Larvae fixed in 4% PFA overnight at 4°C were embedded in 4% low melting temperature agarose and transverse sections collected at 200 μm intervals using a vibrating microtome (Leica VT 1000S). Sections were blocked with 1% BSA in PBS/0.3% Triton X-100 and incubated with a 1:750 dilution of mAb414 (Abcam, #ab24609) at 4°C overnight. Sections were then incubated with 1:500 anti-mouse AF647 and Hoechst 33342 at room temperature for 1 h. Sections were mounted and imaged using a Zeiss LSM880 Fast Airyscan Confocal microscope with a 63x objective. 3D segmentation of cells was performed in ImageJ using EGFP signal for *TO(kras^{G12V})$^{T/+}$* larvae or F-actin stained with 1:200 rhodamine phalloidin for *TO(kras^{G12V})$^{+/+}$* larvae . An outline was drawn around the nuclear periphery as segmented using Hoechst signal. mAb414 fluorescence intensity was calculated for the nuclear periphery and for the cytoplasm. NPC density was calculated by finding maxima for mAb414 at the nuclear surface of 5 nuclei per liver.

Cryosectioning and immunofluorescence microscopy analysis

Dissected livers were fixed in 4% PFA overnight at 4°C and washed with PBS/0.1% Tween 20 before incubation in 30% sucrose in PBS overnight at 4°C. Livers were aligned in a tissue mould, embedded in OCT and frozen on dry ice. The livers were sectioned at 10 μm intervals using a Thermofisher Scientific Microm HM550 cryostat. Sections were washed with PBS before blocking with 10% FCS in PBS/0.3% Triton X-100. Incubation with primary antibodies was performed at 4°C overnight, while incubation with secondary antibodies was performed at room temperature for 1 h. Antibodies used in this work were: 1:2000 α-Tubulin DM1A (CST, #3873), 1:1000 γ-H2AX (gift of James Amatruda), 1:250 cleaved caspase 3 (CST, #9664), 1:500 anti-rabbit AF647 (Thermofisher scientific, #A31573) and 1:500 anti-mouse AF647 (Thermofisher, #A21235). Prolong Diamond Antifade reagent with DAPI (Thermofisher #P36962) was used for slide mounting. A Zeiss LSM880 Fast Airyscan Confocal microscope with a 63x objective was used for image acquisition and ImageJ for image analysis.

Statistical analysis

Data are expressed as mean ± SEM unless indicated otherwise and the number of biological

replicates indicating samples from individual animals/livers, or pools of individual animals/livers for

each experiment are stated in the figure legends. P-values were calculated using Student's t-tests

(two-tailed, followed by Welch's correction) when comparing two groups, and by ANOVA followed

by Tukey's post-hoc test when comparing multiple groups. The effect of Selinexor treatment on liver

volume was analysed by linear regression, regressing liver volume against Selinexor concentration.

All analysis was performed using GraphPad Prism V7.03 (GraphPad software) and p-values ≤ 0.05

were considered statistically significant.

ACKNOWLEDGEMENTS

The authors thank Tyson Blanch, Cameron Mackey, Elizabeth Grgacic and Bryan Ko (zebrafish

husbandry), Ellen Tsui (histology), James Amatruda (rabbit polyclonal antibody to zebrafish γ-H2AX),

Brendon Monahan and Leigh Coultas (insightful discussions).

COMPETING INTERESTS

The authors declare no competing financial interests.

ADDITIONAL INFORMATION

FUNDING

This work was funded by an Australian Government Research Training Program Scholarship and a

WEHI Bridging Fellowship (to KM), the National Health and Medical Research Council of Australia

(Grant 1024878 to JKH), Ludwig Cancer Research, a Victorian State Government Operational

Infrastructure Support grant, and the Australian Government NHMRC Independent Research

Institute Infrastructure Support Scheme.

AUTHORS' CONTRIBUTIONS

KM, KD, JKH conceived and designed the experiments; KM, KD, KS, BH, CS, GB, RM, ATP, TH, EO,

DYRS, ZG, JKH developed the methodology; KM, KD, FG, LW, KS, BH, CS, GB, RM, ATP, JKH acquired

and analysed the data; KM, KD and JKH wrote the manuscript; KD, JKH supervised the study; JKH

acquired the funding.

DATA AVAILABILITY

The datasets generated and/or analysed during the current study are available in the cBioPortal

Cancer Genomics database (http://www.cbioportal.org).

REFERENCES

Gao, S. & Lai, L. Synthetic lethality in drug development: the dawn is coming. *Future Med Chem* **10**, 2129-2132, doi:10.4155/fmc-2018-0227 (2018).

Lord, C. J. & Ashworth, A. PARP inhibitors: Synthetic lethality in the clinic. *Science* **355**, 1152-1158, doi:10.1126/science.aam7344 (2017).

Luo, J. *et al.* A genome-wide RNAi screen identifies multiple synthetic lethal interactions with the Ras oncogene. *Cell* **137**, 835-848, doi:10.1016/j.cell.2009.05.006 (2009).

Wang, T. *et al.* Gene Essentiality Profiling Reveals Gene Networks and Synthetic Lethal Interactions with Oncogenic Ras. *Cell* **168**, 890-903 e815, doi:10.1016/j.cell.2017.01.013 (2017).

Solimini, N. L., Luo, J. & Elledge, S. J. Non-oncogene addiction and the stress phenotype of cancer cells. *Cell* **130**, 986-988, doi:10.1016/j.cell.2007.09.007 (2007).

Okita, K. *et al.* Targeted disruption of the mouse ELYS gene results in embryonic death at peri-implantation development. *Genes Cells* **9**, 1083-1091, doi:10.1111/j.1365-2443.2004.00791.x (2004).

Chen, J. N. *et al.* Mutations affecting the cardiovascular system and other internal organs in zebrafish. *Development* **123**, 293-302 (1996).

de Jong-Curtain, T. A. *et al.* Abnormal nuclear pore formation triggers apoptosis in the intestinal epithelium of elys-deficient zebrafish. *Gastroenterology* **136**, 902-911, doi:10.1053/j.gastro.2008.11.012 (2009).

Davuluri, G. *et al.* Mutation of the zebrafish nucleoporin elys sensitizes tissue progenitors to replication stress. *PLoS Genet* **4**, e1000240, doi:10.1371/journal.pgen.1000240 (2008).

Rasala, B. A., Orjalo, A. V., Shen, Z., Briggs, S. & Forbes, D. J. ELYS is a dual nucleoporin/kinetochore protein required for nuclear pore assembly and proper cell division. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 17801-17806, doi:10.1073/pnas.0608484103 (2006).

11   Gillespie, P. J., Khoudoli, G. A., Stewart, G., Swedlow, J. R. & Blow, J. J. ELYS/MEL-28

chromatin association coordinates nuclear pore complex assembly and replication licensing.

Curr Biol **17**, 1657-1662, doi:10.1016/j.cub.2007.08.041 (2007).

12   Franz, C. *et al.* MEL-28/ELYS is required for the recruitment of nucleoporins to chromatin

and postmitotic nuclear pore complex assembly. *EMBO reports* **8**, 165-172,

doi:10.1038/sj.embor.7400889 (2007).

13   Beck, M. & Hurt, E. The nuclear pore complex: understanding its function through structural

insight. *Nature reviews. Molecular cell biology* **18**, 73-89, doi:10.1038/nrm.2016.147 (2017).

14   Hodgson, B., Li, A., Tada, S. & Blow, J. J. Geminin becomes activated as an inhibitor of

Cdt1/RLF-B following nuclear import. *Curr Biol* **12**, 678-683, doi:10.1016/s0960-

9822(02)00778-9 (2002).

15   Kuhn, T. M., Pascual-Garcia, P., Gozalo, A., Little, S. C. & Capelson, M. Chromatin targeting of

nuclear pore proteins induces chromatin decondensation. *J Cell Biol* **218**, 2945-2961,

doi:10.1083/jcb.201807139 (2019).

16   Guttinger, S., Laurell, E. & Kutay, U. Orchestrating nuclear envelope disassembly and

reassembly during mitosis. *Nature reviews. Molecular cell biology* **10**, 178-191,

doi:10.1038/nrm2641 (2009).

17   Chatel, G. & Fahrenkrog, B. Nucleoporins: leaving the nuclear pore complex for a successful

mitosis. *Cell Signal* **23**, 1555-1562, doi:10.1016/j.cellsig.2011.05.023 (2011).

18   Mishra, R. K., Chakraborty, P., Arnaoutov, A., Fontoura, B. M. & Dasso, M. The Nup107-160

complex and gamma-TuRC regulate microtubule polymerization at kinetochores. *Nat Cell*

Biol **12**, 164-169, doi:10.1038/ncb2016 (2010).

19   Orjalo, A. V. *et al.* The Nup107-160 nucleoporin complex is required for correct bipolar

spindle assembly. *Mol Biol Cell* **17**, 3806-3818, doi:10.1091/mbc.e05-11-1061 (2006).

20   Yokoyama, H. *et al.* The nucleoporin MEL-28 promotes RanGTP-dependent gamma-tubulin

recruitment and microtubule nucleation in mitotic spindle formation. *Nat Commun* **5**, 3270,

doi:10.1038/ncomms4270 (2014).

21   Kobayashi, W. *et al.* Structural and biochemical analyses of the nuclear pore complex

component ELYS identify residues responsible for nucleosome binding. *Commun Biol* **2**, 163,

doi:10.1038/s42003-019-0385-7 (2019).

22   Bilokapic, S. & Schwartz, T. U. Molecular basis for Nup37 and ELY5/ELYS recruitment to the

nuclear pore complex. *Proceedings of the National Academy of Sciences of the United States*

of America **109**, 15241-15246, doi:10.1073/pnas.1205151109 (2012).

23   Rasala, B. A., Ramos, C., Harel, A. & Forbes, D. J. Capture of AT-rich chromatin by ELYS

recruits POM121 and NDC1 to initiate nuclear pore assembly. *Mol Biol Cell* **19**, 3982-3996,

doi:10.1091/mbc.E08-01-0012 (2008).

24   Chew, T. W. *et al.* Crosstalk of Ras and Rho: activation of RhoA abates Kras-induced liver

tumorigenesis in transgenic zebrafish models. *Oncogene* **33**, 2717-2727,

doi:10.1038/onc.2013.240 (2014).

25   Korzh, S. *et al.* Requirement of vasculogenesis and blood circulation in late stages of liver

growth in zebrafish. *BMC Dev Biol* **8**, 84, doi:10.1186/1471-213X-8-84 (2008).

26   Berghmans, S. *et al.* tp53 mutant zebrafish develop malignant peripheral nerve sheath

tumors. *Proceedings of the National Academy of Sciences of the United States of America*

102, 407-412, doi:10.1073/pnas.0406252102 (2005).

27   Hall, T. E. *et al.* Cellular rescue in a zebrafish model of congenital muscular dystrophy type

1A. *NPJ Regen Med* **4**, 21, doi:10.1038/s41536-019-0084-5 (2019).

28   Joseph, J., Liu, S. T., Jablonski, S. A., Yen, T. J. & Dasso, M. The RanGAP1-RanBP2 complex is

essential for microtubule-kinetochore interactions in vivo. *Curr Biol* **14**, 611-617,

doi:10.1016/j.cub.2004.03.031 (2004).

29   Salina, D., Enarson, P., Rattner, J. B. & Burke, B. Nup358 integrates nuclear envelope

breakdown with kinetochore assembly. *J Cell Biol* **162**, 991-1001,

doi:10.1083/jcb.200304080 (2003).

30   Hashizume, C., Kobayashi, A. & Wong, R. W. Down-modulation of nucleoporin

RanBP2/Nup358 impaired chromosomal alignment and induced mitotic catastrophe. *Cell*

Death Dis **4**, e854, doi:10.1038/cddis.2013.370 (2013).

31   Vecchione, L. *et al.* A Vulnerability of a Subset of Colon Cancers with Potential Clinical Utility.

Cell **165**, 317-330, doi:10.1016/j.cell.2016.02.059 (2016).

32   Ober, E. A., Verkade, H., Field, H. A. & Stainier, D. Y. Mesodermal Wnt2b signalling positively

regulates liver specification. *Nature* **442**, 688-691, doi:10.1038/nature04888 (2006).

33   Leshchiner, I. *et al.* Mutation mapping and identification by whole-genome sequencing.

Genome Res **22**, 1541-1548, doi:10.1101/gr.135541.111 (2012).

34   Voz, M. L. *et al.* Fast homozygosity mapping and identification of a zebrafish ENU-induced

mutation by whole-genome sequencing. *PLoS One* **7**, e34671,

doi:10.1371/journal.pone.0034671 (2012).

35   Boglev, Y. *et al.* Autophagy induction is a Tor- and Tp53-independent cell survival response

in a zebrafish model of disrupted ribosome biogenesis. *PLoS Genet* **9**, e1003279,

doi:10.1371/journal.pgen.1003279 (2013).

36   Markmiller, S. *et al.* Minor class splicing shapes the zebrafish transcriptome during

development. *Proceedings of the National Academy of Sciences of the United States of*

America **111**, 3062-3067, doi:10.1073/pnas.1305536111 (2014).

37   Jans, D. A., Martin, A. J. & Wagstaff, K. M. Inhibitors of nuclear transport. *Curr Opin Cell Biol*

58, 50-60, doi:10.1016/j.ceb.2019.01.001 (2019).

38   Zheng, Y. *et al.* KPT-330 inhibitor of XPO1-mediated nuclear export has anti-proliferative

activity in hepatocellular carcinoma. *Cancer Chemother Pharmacol* **74**, 487-495,

doi:10.1007/s00280-014-2495-8 (2014).

39   Cerami, E. *et al.* The cBio cancer genomics portal: an open platform for exploring

multidimensional cancer genomics data. *Cancer Discov* **2**, 401-404, doi:10.1158/2159-

8290.CD-12-0095 (2012).

40   Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles using the

cBioPortal. *Sci Signal* **6**, pl1, doi:10.1126/scisignal.2004088 (2013).

41   Zucman-Rossi, J., Villanueva, A., Nault, J. C. & Llovet, J. M. Genetic Landscape and

Biomarkers of Hepatocellular Carcinoma. *Gastroenterology* **149**, 1226-1239 e1224,

doi:10.1053/j.gastro.2015.05.061 (2015).

42   Levrero, M. *et al.* The p53/p63/p73 family of transcription factors: overlapping and distinct

functions. *J Cell Sci* **113 (Pt 10)**, 1661-1670 (2000).

43   Suh, E. K. *et al.* p63 protects the female germ line during meiotic arrest. *Nature* **444**, 624-

628, doi:10.1038/nature05337 (2006).

44   Kerr, J. B. *et al.* DNA damage-induced primordial follicle oocyte apoptosis and loss of fertility

require TAp63-mediated induction of Puma and Noxa. *Mol Cell* **48**, 343-352,

doi:10.1016/j.molcel.2012.08.017 (2012).

45   Dotsch, V., Bernassola, F., Coutandin, D., Candi, E. & Melino, G. p63 and p73, the ancestors

of p53. *Cold Spring Harb Perspect Biol* **2**, a004887, doi:10.1101/cshperspect.a004887 (2010).

46   Kau, T. R., Way, J. C. & Silver, P. A. Nuclear transport and cancer: from mechanism to

intervention. *Nat Rev Cancer* **4**, 106-117, doi:10.1038/nrc1274 (2004).

47   Rodriguez-Bravo, V. *et al.* Nuclear Pores Promote Lethal Prostate Cancer by Increasing

POM121-Driven E2F1, MYC, and AR Nuclear Import. *Cell* **174**, 1200-1215 e1220,

doi:10.1016/j.cell.2018.07.015 (2018).

48   Hill, R., Cautain, B., de Pedro, N. & Link, W. Targeting nucleocytoplasmic transport in cancer

therapy. *Oncotarget* **5**, 11-28, doi:10.18632/oncotarget.1457 (2014).

49   Calvisi, D. F. *et al.* Ubiquitous activation of Ras and Jak/Stat pathways in human HCC.

Gastroenterology **130**, 1117-1128, doi:10.1053/j.gastro.2006.01.006 (2006).

50   Zheng, W. *et al.* Xmrk, kras and myc transgenic zebrafish liver cancer models share

molecular signatures with subsets of human hepatocellular carcinoma. *PLoS One* **9**, e91179,

doi:10.1371/journal.pone.0091179 (2014).

51   Huo, X. *et al.* Transcriptomic analyses of oncogenic hepatocytes reveal common and

different molecular pathways of hepatocarcinogenesis in different developmental stages

and genders in kras(G12V) transgenic zebrafish. *Biochem Biophys Res Commun* **510**, 558-

564, doi:10.1016/j.bbrc.2019.02.008 (2019).

52   Sakuma, S. *et al.* Inhibition of Nuclear Pore Complex Formation Selectively Induces Cancer

Cell Death. *Cancer Discov* **11**, 176-193, doi:10.1158/2159-8290.CD-20-0581 (2021).

53   Beck, M., Schirmacher, P. & Singer, S. Alterations of the nuclear transport system in

hepatocellular carcinoma - New basis for therapeutic strategies. *J Hepatol* **67**, 1051-1061,

doi:10.1016/j.jhep.2017.06.021 (2017).

54   D'Angelo, M. A., Raices, M., Panowski, S. H. & Hetzer, M. W. Age-dependent deterioration of

nuclear pore complexes causes a loss of nuclear integrity in postmitotic cells. *Cell* **136**, 284-

295, doi:10.1016/j.cell.2008.11.037 (2009).

55   Toyama, B. H. *et al.* Identification of long-lived proteins reveals exceptional stability of

essential cellular structures. *Cell* **154**, 971-982, doi:10.1016/j.cell.2013.07.037 (2013).

56   Bilokapic, S. & Schwartz, T. U. Structural and functional studies of the 252 kDa nucleoporin

ELYS reveal distinct roles for its three tethered domains. *Structure* **21**, 572-580,

doi:10.1016/j.str.2013.02.006 (2013).

57   Ocana, A. & Pandiella, A. Proteolysis targeting chimeras (PROTACs) in cancer therapy. *J Exp

Clin Cancer Res* **39**, 189, doi:10.1186/s13046-020-01672-1 (2020).

58   Mukherjee, H. *et al.* PEARL-seq: A Photoaffinity Platform for the Analysis of Small Molecule-

RNA Interactions. *ACS Chem Biol* **15**, 2374-2381, doi:10.1021/acschembio.0c00357 (2020).

59   Ruijter, J. M. *et al.* Amplification efficiency: linking baseline and bias in the analysis of

quantitative PCR data. *Nucleic Acids Res* **37**, e45, doi:10.1093/nar/gkp045 (2009).