Tensor-decomposition–based unsupervised feature extraction in single-cell multiomics data analysis

Y-h. Taguchi [1,]* and Turki Turki [2]

1 Department of Physics, Chuo University, Tokyo 112-8551, Japan
2 Department of Computer Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia; tturki@kau.edu.sa
* Correspondence:tag@granular.com; Tel.: +81-3-3817-1791

Abstract: Analysis of single-cell multiomics datasets is a novel topic and is considerably challenging because such datasets contain a large number of features with numerous missing values. In this study, we implemented a recently proposed tensor-decomposition (TD)–based unsupervised feature extraction (FE) technique to address this difficult problem. The technique can successfully integrate single-cell multiomics data composed of gene expression, DNA methylation, and accessibility. Although the last two have large dimensions, as many as ten million, containing only a few percentages of non-zero values, TD-based unsupervised FE can integrate three omics datasets without filling missing values. Together with UMAP, which is used frequently when embedding single-cell measurements into two-dimensional space, TD-based unsupervised FE can produce two-dimensional embedding coincident with classification when integrating single-cell omics datasets. Genes selected based on TD-based unsupervised FE were also significantly related to reasonable biological roles.

Keywords: tensor-decomposition; feature extraction; single-cell; multiomics data

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Genes* **2021**, *1*, 0. https://doi.org/

Received:
Accepted:
Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

1. Introduction

Single-cell multiomics data analysis is challenging [1]. There are multiple reasons for this issue. First, it inevitably includes too many missing values. In the usual high throughput sequencing (HTS), the so called depth can compensate for this problem. Nevertheless, because of very limited amount of RNA retrieved from individual cells available, 'depth' cannot resolve this missing value problems. Second, too many missing values result in apparent diversity. The primary purpose of single-cell analysis is to identify the diversity of individual cells that cannot be recognized by the tissue-level HTS. Although missing values are random, apparently very variant profiles appear from a single profile, which can be recognized if there is a large enough number of reads available. This compels researchers to distinguish between true biological diversity and apparent diversity caused by missing values [2].

Finally, single-cell analysis is computationally challenging. Because there are not many samples in the standard HTS, even if the number of features is large, the overall required computational resources decided by the product between the number of features and the number of samples are very limited. Nonetheless, since the number of samples that is the same as that of cells can be huge in single-cell analysis, single-cell analysis can be computationally very challenging.

To resolve these difficulties, we employed tensor-decomposition (TD)–based unsupervised feature extraction (FE) [3]. Prior to applying TD to multiomics datasets, singular value decomposition (SVD) was applied to individual omics profiles such that individual omics profiles have common L singular value vectors. Then, K omics profiles are formatted as an $L \times M \times K$ dimensional tensor, where M is the number of single cells. Then, higher-order singular value decomposition (HOSVD), which is a type of TD,

is applied to the tensor. UMAP applied to singular value vectors attributed to single
cells by HOSVD successfully generated two dimensional embedding, coincident with
known classification of single cells.

2. Materials and Methods

2.1. Gene expression profiles

Two single-cell multiomics datasets were downloaded from GEO using the follow-
ing two GEO IDs:

2.1.1. GSE154762:data set 1

The multiomics dataset [4] retrieved from GEO ID GSE154762, which is denoted as
dataset 1 in this study, is composed of 899 single cells for which gene expression, DNA
methylation, and DNA accessibility were measured. These single cells represent human
oocyte maturation (Table 1). For gene expression, the file "GSE154762_hO_scChaRM_count_matix"
was downloaded from the supplementary file of GEO and was loaded onto R [5] using
`read.table` function in R. For DNA methylation and DNA accessibility, 899 files with
the extensions "WCG.bw" and "HCG.bw" were downloaded from supplementary files
of GEO and were loaded onto R using the `import` function in rtracklayer [6] package in
R.

Table 1. The number of single cells within individual cell types included in data set 1.

FGO	GO1	GO2	Granulosa	Immune	MI	MII	StromaC1	StromaC2
81	40	46	93	20	155	90	189	185

2.1.2. GSE121708:data set 2

The multiomics dataset [7] retrieved from GEO ID GSE154762, which is denoted as
dataset 2 in this study, is composed of 852 single cells for which DNA methylation and
DNA accessibility were measured as well as 758 single cells for which gene expression
was measured. These single cells represent the four time points of the mouse embryo
(Table 2). For gene expression, the file "GSE121650_rna_counts.tsv.gz" was downloaded
from the supplementary file of GEO and was loaded onto R using the `read.table`
function in R. For DNA methylation and DNA accessibility, 852 files with the extensions
"met.tsv.gz" and "acc.tsv.gz" were downloaded from supplementary file of GEO and
were loaded onto R using the `read.table` function in R.

Table 2. The number of single cells at four embryonic time points included in dataset 2. For E7.5,
the gene expression profiles of 296 single cells were measured.

E4.5-5.5	E6.5	E6.75	E7.5
267	98	97	390(296)

2.2. Pre-processing of DNA methylation profiles

First, we collected genomic positions for which at least one measurement was
performed for at least one single cell (i.e., union). Then, for each genomic position, three
integers, -1, 0, and 1, were assigned. When the genomic position was measured in a
single-cell and its state was methylated (non-methylated), we attributed 1 (-1) to the
genomic position of the single cell. Otherwise (i.e., missing observation), we attributed 0
to the genomic transition in a single cell. $x_{ij2} \in \mathbb{R}^{N_2 \times M}$ is stored as a sparse matrix object
using the Matrix [8] package in R because of the large N_2.

2.3. Pre-processing of DNA accessibility

First, we divided the whole genome into 200 nucleotide regions, and DNA accessibil-
ity was summed up within individual regions. These values, which show the summation

of DNA accessibility within individual regions, are regarded as DNA accessibility at the individual 200 nucleotide regions, each of which is supposed to approximately correspond to a single nucleosome that is composed of 140 length DNA that wraps around histones and 80 length of linker DNA. In this study, these 200 nucleotide regions are called as "nucelosome regions". $x_{ij3} \in \mathbb{R}^{N_3 \times M}$ was stored as a sparse matrix object using the Matrix package in R because of the large N_3.

2.4. TD-based unsupervised FE

2.4.1. Reduction of feature dimensions

Here, feature denotes gene expression, DNA methylation, or DNA accessiblity. Because the features of these three datasets differ from one another, we first applied SVD to these features. Suppose $x_{ijk} \in \mathbb{R}^{N_k \times M \times K}$ represents the value of the ith feature (expression of the ith gene, methylation of the ith genomic location, or DNA accessibility of the ith nucleosome region) at the kth single cell of the kth omics datae ($1 \leq k \leq K = 3$, $k = 1$:gene expression, $k = 2$:DNA methylation, and $k = 3$:DNA accessibility). Applying SVD to x_{ijk}, and we get

$$x_{ijk} = \sum_{\ell=1}^{L} \lambda_\ell u_{\ell ik} v_{\ell jk} \tag{1}$$

where λ_ℓ is the ℓth singular value, and $u_{\ell_i k}$ and $v_{\ell jk}$ are the ith and jth components of the ℓth left and right singular value vectors, respectively. Then, x_{ijk} is transformed to $x_{\ell jk} \in \mathbb{R}^{L \times M \times K}$ to have the same (common) feature dimension, L, independent of k, as

$$x_{\ell jk} = \sum_{i=1}^{N_k} u_{\ell ik} x_{ijk} \tag{2}$$

2.4.2. Data normalization

Prior to applying SVD to the individual omics profiles in these two datasets $,x_{ijk}, (k = 2, 3)$, that is, DNA methylation and accessibility, of dataset 1 is normalized such that

$$\sum_{i=1}^{N_k} |x_{ijk}| = N_k \tag{3}$$

whereas x_{ij1}, i.e., gene expression, is normalized such that

$$\sum_{i=1}^{N_K} x_{ij1} = 0 \tag{4}$$

$$\sum_{i=1}^{N_k} x_{ij1}^2 = N_k \tag{5}$$

for datasets 1 and 2. The reason why DNA methylation and accessibility of dataset 2 are not normalized is because $\sum_i |x_{ijk}|, (k = 2, 3)$ is very small in some single cells in dataset 2. Thus, applying normalization adds significant weight to these single cells with fewer observations and drastically skewed outcomes. To avoid this problem, $x_{ijk}, (k = 2, 3)$ of dataset 2 was not normalized.

2.4.3. TD applied to dimension reduced multiomics data sets

HOSVD [3] was applied to the tensor, $x_{\ell jk}$, and we got

$$x_{\ell jk} = \sum_{\ell_1=1}^{L} \sum_{\ell_2=1}^{M} \sum_{\ell_3=1}^{K} G(\ell_1 \ell_2 \ell_3) u_{\ell_1 \ell} u_{\ell_2 j} u_{\ell_3 k} \tag{6}$$

where $G \in \mathbb{R}^{L \times M \times K}$ is the core tensor that represents the contribution of $u_{\ell_1 \ell} u_{\ell_2 j} u_{\ell_3 k}$
to $x_{\ell jk}$. $u_{\ell_1 \ell} \in \mathbb{R}^{L \times L}, u_{\ell_2 j} \in \mathbb{R}^{M \times M}, u_{\ell_3 k} \in \mathbb{R}^{K \times K}$ are singular value matrices and are
orthogonal matrices.

2.5. Categorical regression

For categorical regression to test the coincidence between classification shown in
Tables 1 or 2 and singular value vectors attributed to the *j*th single cells, we performed
categorical regression

$$v_{\ell jk} = a_{\ell ks} \delta_{sj} + b_{\ell k} \tag{7}$$
$$u_{\ell_2 j} = a_{\ell_2 s} \delta_{sj} + b_{\ell_2} \tag{8}$$

where *s* denotes one of the classifications shown in Table 1 or 2, $a_{\ell ks}, b_{\ell k}, a_{\ell_2 ks}, b_{\ell_2 k}$ are
regression coefficients, and δ_{js} takes the value of 1 when the *j*th single cell belongs to the
*s*th classification and 0 otherwise. Categorical regression was performed using the `ls`
function in R. The obtained *P*-values were corrected using the Benjamini-Hochberg (BH)
criterion [3]. ℓs or ℓ_2s associated with adjusted *P*-values less than 0.01 are regarded to be
coincident with classification.

2.6. UMAP

Two-dimensional embedding was performed by UMAP [9]. The `umap` function
implemented in R was used.

2.7. Gene selection

After identifying which $u_{\ell_2 j}$ coincides with the classification, we need to identify
which $u_{\ell_1 \ell}$ is associated with the selected $u_{\ell_2 j}$ by investigating $|G(\ell_1 \ell_2 \ell_3)|$; ℓ_1s with a
larger $|G|$ with the selected ℓ_2 are regarded to be coincident with the classification. Then,
the selected $u_{\ell_1 \ell}$ is converted back to $u_{\ell_1 i_1}$ attributed to genes as

$$u_{\ell_1 i} = \sum_{\ell=1}^{L} u_{\ell_1 \ell} u_{\ell i 1} \tag{9}$$

P-values can be attributed to genes, *i*, assuming $u_{\ell_1 i}$ obeys a multiple Gaussian distribution (null hypothesis) as

$$P_i = P_{\chi^2} \left[> \sum_{\ell_1} \left(\frac{u_{\ell_1 i}}{\sigma_{\ell_1}} \right)^2 \right] \tag{10}$$

where the summation is taken over only the selected ℓ_1s, and $P_{\chi^2}[> x]$ is the cumulative
χ^2 distribution, where the argument is larger than *x*, and σ_{ℓ_1} is the standard deviation.
P_is are corrected by the BH criterion [3], and *i*s associated with adjusted P_i less than 0.01
are selected.

3. Results

3.1. Data set 1

We got $x_{ij1} \in \mathbb{R}^{26500 \times 899}$, $x_{ij2} \in \mathbb{R}^{26438807 \times 899}$, and $x_{ij3} \in \mathbb{R}^{15478375 \times 899}$. SVD was
applied to x_{ijk} with $L = 10$, as in Eq. (1). For $x_{ijk}, k = 2, 3$, SVD was performed using the
`irlba` function in the irlba package [10] in R because of the large $N_k, k = 2, 3$ as many as
10 million. Then, HOSVD was applied to $x_{\ell jk}$, as in eq. (2).
One possible validation to check whether the above procedure works properly is
to check whether $v_{\ell jk}$ and $u_{\ell_2 j}$ are coincident with the classification shown in Table 1.
Because the above procedure is fully unsupervised, it is unlikely that $v_{\ell jk}$ and $u_{\ell_2 j}$ are
accidentally coincident with the classification. To quantitatively validate the coincidence
between the classification and $v_{\ell jk}$ or $u_{\ell_2 j}$, we applied categorical regression (see §2.5) .

Table 3. Number of singular value vectors coincident with classification shown in Table 1.

Adjusted P-value	gene expression	SVD ($v_{\ell jk}$) DNA methylation	DNA accesibility	HOSVD ($u_{\ell_2 j}$) DNA methylation and accesiblity	All
< 0.01	10	7	1	10	18
≥ 0.01	0	3	9	10	12

Table 3 shows the number of singular value vectors coincident with the classification shown in Table 1. When SVD was applied to individual omics data, because $L = 10$, the number of singular value vectors was 10 as well. When K omics datasets are integrated and HOSVD is applied, the number of singular value vectors is $KL = 10K$. Thus, when DNA methylation and accessibility are integrated, 20 singular value vectors are available. When all three omics data are integrated, 30 singular value vectors are available. It is obvious that for all five cases, at least one singular value vector is coincident with the classification. Thus, our strategy is essentially successful.

To further validate the successful integration of singular value vectors, we applied UMAP to 20 or 30 singular value vectors obtained by HOSVD (Fig. 1).

Figure 1. Two-dimensional embedding of singular value vectors, $u_{\ell_2 j}$, computed by HOSVD applied to $x_{\ell j k}$ in dataset 1 (Table 3). Upper: $u_{\ell_2 j}, 1 \le \ell_2 \le 20$ when only DNA methylation and accessibility ($k = 2, 3$) are integrated. Lower: $u_{\ell_2 j}, 1 \le \ell_2 \le 30$ when all three omics data points ($1 \le k \le 3$) are integrated. Default settings other than custom.config\$n_neighbors=100 were used.

It is obvious that the integration of all three omics datasets (lower) is more coincident with classification than that of integration of two omics datasets, DNA methylation, and accessibility (upper). This suggests the usefulness of integrating the three omics datasets. In fact, single omics data cannot provide two-dimensional embedding coinciding with classification (Fig. S1).

We also attempted to validate biological outcomes when all three omics datasets were integrated. We selected 47 genes associated with adjusted P_i less than 0.01, as described in §2.7 using u_{1i} because u_{1i} is associated with the largest

$$\sum_{\ell_2} \sum_{\ell_3=1}^{3} G^2(\ell_1 \ell_2 \ell_3), \tag{11}$$

where the summation of ℓ_2 is taken over only 18 ℓ_2s coincident with the classification (Table 3). The selected 47 genes (Data S1) were uploaded to Enrichr [11].

Forty-seven genes were enriched by H3K36me3 based on "ENCODE Histone Modifications 2015"; H3K36m3 is known to play critical roles during oocyte maturation [12]. Forty-seven genes were also targeted by MYC based on "ENCODE and ChEA Consensus TFs from ChIP-X"; Myc is known to play critical roles in oogenesis [13]. Forty-seven genes were also targeted by TAF7 based on "ENCODE and ChEA Consensus TFs from ChIP-X" and "ENCODE TF ChIP-seq 2015"; TAF7 is known to play critical roles during oocyte growth [14]. Forty-seven genes are also targeted by ATF2 based upon "ENCODE and ChEA Consensus TFs from ChIP-X"; expression of ATF2 is known to be altered during oocyte dvelopment [15]. This suggests that our strategy correctly captures regulation-related parts (full data of enrichment analysis can be available as Data S1).

3.2. Data set 2

To confirm that the success in the previous section is not accidental, we applied the same procedure to dataset 2 as well. We got $x_{ij1} \in \mathbb{R}^{22084 \times 758}$, $x_{ij2} \in \mathbb{R}^{20106507 \times 852}$, and $x_{ij3} \in \mathbb{R}^{13627678 \times 852}$. SVD was applied to x_{ijk} with $L = 10$, as in Eq. (1). For $x_{ijk}, k = 2, 3$, SVD was performed using the `irlba` function in the irlba package [10] in R because of the large $N_k, k = 2, 3$ of as many as ten millions. Then, HOSVD was applied to $x_{\ell jk}$, as in eq. (2). Because $N_1 = 758 < N_2 = N_2 = 852$, when HOSVD is applied to $x_{\ell jk}$ composed of all three omics datasets, only 758 single cells shared with all x_{ijk} are considered. As described in the previous section, we first validated the coincidence between singular value vectors attributed to single cells, that is, $v_{\ell j}$ and $u_{\ell_2 \ell k}$, and classification in Table 2 (Table 4).

Table 4. Number of singular value vectors coincident with classification shown in Table 2.

| Adjusted P-value | gene expression | SVD ($v_{\ell jk}$) | | HOSVD ($u_{\ell_2 j}$) | All |
		DNA methylation	DNA accesibility	DNA methylation and accesiblity	
< 0.01	10	7	5	10	18
≥ 0.01	0	3	5	10	12

The coincidence between the singular value vectors and the classification in Table 4 is even better than those in Table 3. Thus, it is unlikely that the superior outcome in Table 3 is purely accidental. To further validate the successful integration of singular value vectors, we applied UMAP to 20 or 30 singular value vectors obtained by HOSVD (Fig. 2).

Figure 2. Two-dimensional embedding of singular value vectors, $u_{\ell_2 j}$, computed by HOSVD applied to $x_{\ell jk}$ in dataset 2 (Table 4). Upper: $u_{\ell_2 j}, 1 \leq \ell_2 \leq 20$ when only DNA methylation and accessibility ($k = 2, 3$) are integrated. Lower: $u_{\ell_2 j}, 1 \leq \ell_2 \leq 30$ when all three omics data points ($1 \leq k \leq 3$) are integrated. Default settings other than custom.config\$n_neighbors=100 were used.

167   It is obvious that the integration of all three omics datasets (lower) is more coincident
168 with classification than that of integration of two omics datasets, DNA methylation, and
169 DNA accessibility (upper), as can be seen in Fig. 2. This again confirms the usefulness
170 of integrating the three omics datasets. In fact, single omics data cannot provide two-
171 dimensional embedding coinciding with classification (Fig. S2).

      We also attempted to validate biological outcomes when all three omics datasets
were integrated. We selected 175 genes associated with adjusted P_i less than 0.01, as
described in §2.7 using u_{1i} because u_{1i} is associated with the largest

$$\sum_{\ell_2} \sum_{\ell_3=1}^{3} G^2(\ell_1 \ell_2 \ell_3) \tag{12}$$

where the summation of ℓ_2 is taken over only 18 ℓ_2s coincident with the classification (Ta-
ble 4). The selected 175 genes (Data S2) were converted to gene symbols by DAVID [16]
gene ID converter and were uploaded to Enrichr.

One hundred and seventy-five genes were enriched by H3K36me3 based on "EN-
CODE Histone Modifications 2015"; H3K36m3 is known to play critical roles during
gastrulation [17]. One hundred and seventy-five genes were also targeted by MYC
based on "ENCODE and ChEA Consensus TFs from ChIP-X"; Myc is also known to
play critical roles in gastrulation [18]. One hundred and seventy-five genes were also
targeted by TAF7 based on "ENCODE and ChEA Consensus TFs from ChIP-X" and
"ENCODE TF ChIP-seq 2015"; TAF7 is known to play critical roles during gastrula-
tion [19]. One hundred and seventy-five genes were also targeted by ATF2 based on
"ENCODE and ChEA Consensus TFs from ChIP-X"; expression of ATF2 is known to be
maintained during gastrulation [20]. This suggests that our strategy correctly captures
regulation-related parts (full data of enrichment analysis is available as Data S2).

These two examples, application to datasets 1 and 2, demonstrate the usefulness
of the present strategy to integrate single-cell multimics datasets composed of gene
expression, DNA methylation, and accessibility.

189 4. Discussion

In this study, we demonstrated the usefulness of our strategy when it is applied to
the integrated analysis of single-cell multiomics datasets composed of gene expression,
DNA methylation, and DNA accessibility. One might wonder if other more popular
methods can achieve similar performance because our strategy is useless if others can
perform comparatively. There are several advantages of our method, which other
methods do not have.

First, we do not have to fill the missing values with non-zero values. Single-cell
measurements are usually associated with a large number of missing values (Table 5).

Table 5. Number of single cells, features, non-zero components, and its ratios.

numbers	expression	DNA methylation	DNA accessibility
	Data set 1		
single cells	899	899	899
features	26500	26438807	15478375
total components	2.38×10^7	2.38×10^{10}	1.39×10^{10}
non-zero components	6.76×10^6	5.50×10^8	3.85×10^8
the ratio of non-zero components	0.28	0.02	0.03
	Data set 2		
single cells	758	852	852
features	22084	20106507	13627678
total components	1.67×10^7	1.71×10^{10}	1.16×10^{10}
non-zero components	4.87×10^6	6.96×10^8	7.87×10^8
the ratio of non-zero components	0.29	0.04	0.07

Although gene expression profiles are associated with a relatively small number
of missing components, more than 70 % were missing. For DNA methylation and
accessibility, the situation is very difficult to treat. Only a few percentages of components
have values, while the rest are missing values. To address this problem, especially
for DNA methylation and accessibility, heavy pre-processing is usually required. For
example, for dataset 1, statistical tests are applied and regions associated with significant
P-values are selected [4], which reduces the number of features attributed to DNA
methylation and accessibility. Because such a statistical test automatically filters out
regions filled with missing values, the ratio of non-zero components is also reduced as a
result. For dataset 2, the authors restricted the features to only the most variable ones
(typically, $\sim 10^3$) and occasionally filled missing components with Bayesian models [7].
These procedures inevitably introduce arbitrariness of outcomes, as pre-processing data
might affect the outcome. In contrast to these arbitrary procedures, our method is almost
unsupervised. We did not select any features or fill in the missing values. Despite these
fully unsupervised strategies, our results are highly coincident with the classification
(Tables 3 and 4 and Figs. 1 and 2). From this perspective, our strategy is superior to the
other methods.
Second, our method can deal with massive datasets. For example, although in-
tegrated analysis of multiomics data was performed using multiomics factor analysis
(MOFA) [21] in the original studies [4,7] of datasets 1 and 2, MOFA cannot accept x_{ijk}
in this study as inputs because MOFA does not implement sparse matrix architecture.
During the computation of MOFA, zero values must be filled with non-zero values to
evaluate the convergence; this results in a dense matrix that cannot be stored in the com-
puter memory because the number of components of DNA methylation and accessibility
are too large to store them as they are (Table 5). In our computation, we can apply SVD
to these large datasets while keeping them in a sparse matrix format using the irlba
package implemented in R. SVD not only reduces the number of features to L but also
fills missing values. Thus, we can manage a large matrix as it is in our implementation.
Third, our method is free from dividing weight between multiomics data sets; how
to weigh individual omics data must be decided based on some criteria outside the
datasets available. Nevertheless, in our implementation, the weight of individual omics
data is represented by $u_{\ell_3 k}$, which is automatically decided by simply applying HOSVD
to a multiomics dataset. Thus, from this perspective, our strategy is outstanding.
Although we have shown that the integration of all three omics data are superior
to that of integration of DNA methylation and accessibility (Figs. 1 and 2), one might
wonder if integration of gene expression and DNA methylation or DNA accessibility
might be comparative with that of all thee omics data sets. In order to deny this

possibility, we also considered these combinations of two of three omics data sets (Figs. S3 and S4). Although the integration of gene expression and DNA accessibility in dataset 1 (Fig. S3(B)) is comparative with that of all three omics data, neither integration of gene expression and DNA memthylation (Fig. S4(A)) nor that of gene expression and DNA accessibility (Fig. S4(B)) is comparative with that of all three omics data in dataset 2. Thus, it is obvious that only integaration of three omics datasets can give us UMAP embeddinng coincident with classification regardless to the dataset considerd.

As for the comparisons with other methods, as mentioned above, no methods implemented with sparse matrix architecture and applicable to multiomics datasets exist to our knowledge. Thus, we could not compare our performance with other methods.

Prospective uses of our methods are as follows. First of all, it can integrate gene expression profiles, DNA methylation and accessibility in single cell measurements with applying no pre-processing; this enables researchers to obtain the reasonable results without struggling on how to convert raw data into treatable formats. In addition to this, since it can save memories required for analysing single cell multiomics data sets, more researchers who cannot have massive computational facilities can analyze massive single cell measurements.

5. Conclusions

In this study, we propose a method for applying TD-based unsupervised FE to single-cell multiomics datasets composed of gene expression, DNA methylation, and DNA accessibility. Together with UMAP, the proposed method successfully integrated a multiomics dataset and generated two-dimensional embedding of single cells coincident with the classification. The implementation requires neither filling missing values nor massive CPU memory to store multiomics datasets of single cells and can deal with DNA methylation and accessibility with ten million features. The present implementation is very promising and can be a *de facto* standard method to integrate single-cell multiomics datasets composed of gene expression, DNA methylation, and DNA accessibility.

Author Contributions: YHT planned the research, performed analyses. YHT and TT have evaluated the results, discussions, outcomes and wrote and reviewed the manuscript.

Funding: This work was supported by KAKENHI [grant numbers 19H05270, 20H04848, and 20K12067] to YHT.

Data Availability Statement: Data used in this study is available in GEO ID GSE154762 and GSE121708. Sample R source can be found at https://github.com/tagtag/scMultiR

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

BH	Benjamini-Hochberg
FE	feature extraction
HOSVD	higher order singular value decomposition
HTS	high throughput sequencing
MOFA	Multi-Omics Factor208Analysis
SVD	singular value decomposition
TD	tensor decomposition

References

1. Lee, J.; Hyeon, D.Y.; Hwang, D. Single-cell multiomics: technologies and data analysis methods. *Experimental & Molecular Medicine* **2020**, *52*, 1428–1442. doi:10.1038/s12276-020-0420-2.

2. Liu, Q.; Herring, C.A.; Sheng, Q.; Ping, J.; Simmons, A.J.; Chen, B.; Banerjee, A.; Li, W.; Gu, G.; Coffey, R.J.; Shyr, Y.; Lau, K.S. Quantitative assessment of cell population diversity in single-cell landscapes. *PLOS Biology* **2018**, *16*, 1–29. doi:10.1371/journal.pbio.2006687.

3. Taguchi, Y.H. *Unsupervised Feature Extraction Applied to Bioinformatics*; Springer International Publishing, 2020. doi:10.1007/978-3-030-22456-1.

4. Yan, R.; Gu, C.; You, D.; Huang, Z.; Qian, J.; Yang, Q.; Cheng, X.; Zhang, L.; Wang, H.; Wang, P.; Guo, F. Decoding dynamic epigenetic landscapes in human oocytes using single-cell multi-omics sequencing. *Cell Stem Cell* **2021**. doi:https://doi.org/10.1016/j.stem.2021.04.012.

5. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2020.

6. Lawrence, M.; Gentleman, R.; Carey, V. rtracklayer: an R package for interfacing with genome browsers. *Bioinformatics* **2009**, *25*, 1841–1842, [https://academic.oup.com/bioinformatics/article-pdf/25/14/1841/606195/btp328.pdf]. doi:10.1093/bioinformatics/btp328.

7. Argelaguet, R.; Clark, S.J.; Mohammed, H.; Stapel, L.C.; Krueger, C.; Kapourani, C.A.; Imaz-Rosshandler, I.; Lohoff, T.; Xiang, Y.; Hanna, C.W.; Smallwood, S.; Ibarra-Soria, X.; Buettner, F.; Sanguinetti, G.; Xie, W.; Krueger, F.; Göttgens, B.; Rugg-Gunn, P.J.; Kelsey, G.; Dean, W.; Nichols, J.; Stegle, O.; Marioni, J.C.; Reik, W. Multi-omics profiling of mouse gastrulation at single-cell resolution. *Nature* **2019**, *576*, 487–491. doi:10.1038/s41586-019-1825-8.

8. Bates, D.; Maechler, M. *Matrix: Sparse and Dense Matrix Classes and Methods*, 2021. R package version 1.3-4.

9. Becht, E.; McInnes, L.; Healy, J.; Dutertre, C.A.; Kwok, I.W.H.; Ng, L.G.; Ginhoux, F.; Newell, E.W. Dimensionality reduction for visualizing single-cell data using UMAP. *Nature Biotechnology* **2018**, *37*, 38–44. doi:10.1038/nbt.4314.

10. Baglama, J.; Reichel, L.; Lewis, B.W. *irlba: Fast Truncated Singular Value Decomposition and Principal Components Analysis for Large Dense and Sparse Matrices*, 2019. R package version 2.3.3.

11. Kuleshov, M.V.; Jones, M.R.; Rouillard, A.D.; Fernandez, N.F.; Duan, Q.; Wang, Z.; Koplev, S.; Jenkins, S.L.; Jagodnik, K.M.; Lachmann, A.; McDermott, M.G.; Monteiro, C.D.; Gundersen, G.W.; Ma'ayan, A. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Research* **2016**, *44*, W90–W97, [https://academic.oup.com/nar/article-pdf/44/W1/W90/18788036/gkw377.pdf]. doi:10.1093/nar/gkw377.

12. Xu, Q.; Xiang, Y.; Wang, Q.; Wang, L.; Brind'Amour, J.; Bogutz, A.B.; Zhang, Y.; Zhang, B.; Yu, G.; Xia, W.; Du, Z.; Huang, C.; Ma, J.; Zheng, H.; Li, Y.; Liu, C.; Walker, C.L.; Jonasch, E.; Lefebvre, L.; Wu, M.; Lorincz, M.C.; Li, W.; Li, L.; Xie, W. SETD2 regulates the maternal epigenome, genomic imprinting and embryonic development. *Nature Genetics* **2019**, *51*, 844–856. doi:10.1038/s41588-019-0398-7.

13. SUZUKI, T.; ichiro ABE, K.; INOUE, A.; AOKI, F. Expression of c-MYC in Nuclear Speckles During Mouse Oocyte Growth and Preimplantation Development. *Journal of Reproduction and Development* **2009**, *55*, 491–495. doi:10.1262/jrd.09-069A.

14. Yu, C.; Cvetesic, N.; Gupta, K.; Ye, T.; Gazdag, E.; Hisler, V.; Negroni, L.; Hajkova, P.; Lenhard, B.; Müller, F.; Berger, I.; Vincent, S.D.; Tora, L. TBPL2/TFIIA complex overhauls oocyte transcriptome during oocyte growth. *bioRxiv* **2020**, [https://www.biorxiv.org/content/early/2020/06/09/2020.06.08.118984.full.pdf]. doi:10.1101/2020.06.08.118984.

15. Vigneault, C.; McGraw, S.; Sirard, M.A. Spatiotemporal expression of transcriptional regulators in concert with the maternal-to-embryonic transition during bovine in vitro embryogenesis. *REPRODUCTION* **01 Jan. 2009**, *137*, 13 – 21. doi:10.1530/REP-08-0077.

16. Huang, D.W.; Sherman, B.T.; Lempicki, R.A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* **2008**, *4*, 44–57. doi:10.1038/nprot.2008.211.

17. Sitbon, D.; Boyarchuk, E.; Dingli, F.; Loew, D.; Almouzni, G. Histone variant H3.3 residue S31 is essential for Xenopus gastrulation regardless of the deposition pathway. *Nature Communications* **2020**, *11*. doi:10.1038/s41467-020-15084-4.

18. Downs, K.M.; Martin, G.R.; Bishop, J.M. Contrasting patterns of myc and N-myc expression during gastrulation of the mouse embryo. *Genes & Development* **1989**, *3*, 860–869, [http://genesdev.cshlp.org/content/3/6/860.full.pdf+html]. doi:10.1101/gad.3.6.860.

19. Langer, D.; Martianov, I.; Alpern, D.; Rhinn, M.; Keime, C.; Dollé, P.; Mengus, G.; Davidson, I. Essential role of the TFIID subunit TAF4 in murine embryogenesis and embryonic stem cell differentiation. *Nature Communications* **2016**, *7*. doi:10.1038/ncomms11063.

20. Villarreal, X.C.; Richter, J.D. Analysis of ATF2 gene expression during early Xenopus laevis development. *Gene* **1995**, *153*, 225–229. doi:https://doi.org/10.1016/0378-1119(94)00770-S.

21. Argelaguet, R.; Velten, B.; Arnol, D.; Dietrich, S.; Zenz, T.; Marioni, J.C.; Buettner, F.; Huber, W.; Stegle, O. Multi-Omics Factor Analysis-a framework for unsupervised integration of multi-omics data sets. *Molecular Systems Biology* **2018**, *14*, e8124, [https://www.embopress.org/doi/pdf/10.15252/msb.20178124]. doi:https://doi.org/10.15252/msb.20178124.