NF-κB-Inducing Kinase Maintains Mitochondrial Efficiency and Systemic Metabolic Homeostasis

Authors: Kathryn M. Pflug[1,2], Dong W. Lee[1], Justin Keeney[1], Raquel Sitcheran[1,2,*]

Affiliations:

[1] Department of Molecular & Cellular Medicine, Texas A&M University Health Science Center, College Station TX 77845, USA

[2] Interdisciplinary Graduate Program in Genetics, Texas A&M University, College Station TX 77845, USA

* Correspondence: sitcheran@tamu.edu

Conflicts of Interest: The authors declare no conflicts of interest.

Keywords

NF-κB-inducing kinase, Glycolysis, Oxidative Phosphorylation (OXPHOS), Spare Respiratory Capacity, Proton Leak, High-Fat Diet

Abstract

Background: NF-κB-inducing kinase (NIK) is a critical regulator of immunity and inflammation and NIK loss-of-function mutations have recently been described in patients with primary immunodeficiency disease. Based on our previous work showing that NIK regulates adaptive metabolic responses in glucose-starved cancer cells, we investigated whether NIK is required for mitochondrial functions in bioenergetic processes and metabolic responses to nutritional stress in NIK knockout (KO) mice, which recapitulate the clinical presentation of NIK PID patients.

Methods: We performed whole body composition analysis of wild type (WT) and NIK KO mice using EchoMRI and DEXA imaging. Seahorse extracellular flux analyses were used to monitor oxidative phosphorylation and glycolysis through oxygen consumption rates (OCR) and extracellular acidification rates (ECAR) in preadipocyte cells and in ex vivo adipose tissue. NIK regulation of systemic metabolic output was measured by indirect calorimetry using TSE Phenomaster metabolic chambers under basal conditions as well as in response to nutritional stress induced by a prolonged high-fat diet (HFD). Finally, we analyzed a role for NIK in adipocyte differentiation, as well as the contributions of canonical and noncanonical NF-κB signaling to adipose development and metabolic output.

Results: We observed that in adipose cells, NIK is required for maintaining efficient mitochondrial membrane potential and spare respiratory capacity (SRC), indicators of mitochondrial fitness. NIK KO preadipocytes and ex vivo adipose tissue exhibited diminished SRC, increased proton leak, with compensatory upregulation of glycolysis. Systemically, NIK KO mice exhibited increased glucose utilization, increased energy expenditure, and reduced adiposity, which persisted under the stress of HFD. Finally, while NIK controlled adipocyte differentiation through activation of RelB and the noncanonical NF-κB pathway, NIK regulation of metabolism in preadipocytes was NF-κB/RelB-independent.

Conclusion: Our results demonstrate that NIK is required for metabolic homeostasis both locally,

on a cellular and tissue level, as well as systemically, on an organismal level. Collectively, the

data suggest that NIK KO cells upregulate glycolytic metabolism as a compensatory response to

impaired mitochondrial fitness (diminished SRC) and mitochondrial efficiency (increased proton

leak). To meet changes in bioenergetic demands, NIK KO mice undergo metabolic rewiring

through increased glucose utilization and glycolysis, which persists under the stress of

overnutrition with a HFD. Moreover, while NIK regulation of metabolism is RelB-independent, NIK

regulation of adipocyte development requires RelB and activation of the noncanonical NF-κB

pathway. Our findings establish NIK as an important regulator of cellular and systemic metabolic

homeostasis, suggesting that metabolic dysfunction may be an important component of primary

immunodeficiency diseases arising from loss of NIK function.

Abbreviations: NIK- NF-κB-Inducing Kinase, NF-κB- Nuclear factor kappa light chain enhancer of

activated B cells, IKKα- inhibitor of kappa B kinase alpha, PID- primary immunodeficiency, SRC- spare

respiratory capacity, OCR- oxygen consumption rate, ECAR- extracellular acidification rate, ingWAT-

inguinal white adipose tissue, DEXA- dual energy X-ray absorptiometry, gWAT- gonadal white adipose

tissue, RER- respiratory exchange rate, HFD- High-fat diet, PPARγ- peroxisome proliferator activated

receptor, C/EBPα- CCAAT enhancer binding protein alpha, ETYA- eicosatetraynoic acid, TNFα- tumor

necrosis factor alpha, TWEAK- tumor necrosis factor weak like.

1. Introduction

Nuclear Factor-κB (NF-κB)-inducing kinase (NIK), encoded by *MAP3K14*, is a serine,

threonine protein kinase that that is best known for its function as an upstream inducer of NF-κB

signaling to regulate innate and adaptive immunity. While NIK can activate canonical and

noncanonical NF-κB pathways, it is uniquely required for activation of noncanonical NF-κB

signaling through phosphorylation of Inhibitor of κB Kinase α (IKKα), which triggers p100

processing to p52, and generation of transcriptionally active p52-RelB NF-κB complexes [1–4].

NIK has critical roles in B-cell, lymphocyte, and lymph node development, immunoglobulin (Ig)

production and T-cell function. Consequently, Nik^{aly} mutant mice (*aly*; alymphoplasia), which lack

NIK activity, and NIK knockout mice (NIK KO) exhibit lymphopenia, abnormal Peyer's patches,

aberrant splenic and thymic structures, reduced B-cell numbers and Ig serum levels leading to

humoral immunodeficiency [1,5–9]. In humans, NIK loss-of-function mutations were recently

identified in patients with primary immunodeficiency (PID) who exhibit similar immune defects as

Nik^{aly} and NIK KO mice [10,11], demonstrating the relevance of NIK deficient mouse models for

immunodeficiency disease.

We, and others, have elucidated important NF-κB-independent metabolic functions for

NIK, including roles in mitochondrial dynamics, metabolic reprogramming in macrophages and

under nutrient stress in cancer cells, as well as regulation of glycolysis in T-cells [12–15]. In

response to diet-induced obesity, gain-of-function studies have shown that NIK induces

hyperglycemia by increasing glucagon activity, and liver steatosis through induction of fatty acid

oxidation, and loss of NIK was shown to increase glucose and insulin tolerance [16–18]. Our

recent work has established NIK mitochondrial localization and regulation of mitochondrial

respiration and fitness, promoting metabolic adaptation of cancer cells to glucose starvation [15].

However, NIK regulation of mitochondrial metabolism has not been examined in adipose tissue,

or on an organismal level.

Metabolic homeostasis relies on a balance between inputs such as glucose and oxygen

necessary for energetic processes and utilization of catabolic pathways involving glycolysis and

oxidative phosphorylation (OXPHOS) for efficient energy production [19,20]. Utilization of these

major metabolic pathways is dynamic and shifts in response to alterations in cellular states or

stress [21,22]. Glycolysis and OXPHOS are interwoven and dysregulation in one can alter the

flux or rate of the other [23–25]. Although glycolysis yields low amounts of ATP, it is important for

feeding into the TCA cycle and electron transport chain (ETC) for rapid production of ATP,

particularly under low oxygen availability [26–28]. Furthermore, glucose catabolism is preferred

in response to stress as it requires the lowest input of ATP [29–31]. Oxidative phosphorylation, or

aerobic respiration, is more commonly utilized to meet ATP demands. Important for mitochondrial

respiration is its spare respiratory capacity (SRC), a measurable indicator of mitochondrial fitness,

or the ability to upregulate oxygen consumption and ATP production to meet changes in energetic

demand [32–34]. Energy derived from mitochondria is essential for maintaining cellular

homeostasis, as well as in response to stress. Reciprocally, on an organismal level, alterations in

mitochondrial functions impact physiological and behavioral responses, as seen in individuals

with impaired ETC function due to primary mitochondrial disorders (PMDs), who are unable to

adapt to dietary and physiological stressors [35,36]. As such, mitochondrial fitness and efficiency

is essential to organismal homeostasis and adaptation to bioenergetic stress.

Here we investigate a functional and developmental role for NIK in controlling metabolic

homeostasis basally and in response to nutritional stress induced by a high-fat diet (HFD). Our

data demonstrate that NIK deficient preadipocytes and *ex vivo* tissue exhibited loss of

mitochondrial SRC and increased proton leak. This uncoupling of mitochondrial respiration and

inefficient metabolism was accompanied by a compensatory increase in glycolysis and

extracellular acidification rate (ECAR). Furthermore, we show that NIK regulation of mitochondrial

SRC and proton leak is independent of noncanonical RelB/NF-κB signaling. Similar to NIK KO

cells and tissue, NIK deficiency in mice resulted in increased glucose utilization and glycolysis to

meet energetic demands. In NIK KO mice, and *ex vivo* adipose tissue, this increased energy

expenditure resulted in reduced adiposity and resistance to adipose accumulation and weight

gain through development, as well as under a HFD. While NIK regulates mitochondrial functions

independently of RelB/NF-κB, we found that NIK promotes adipocyte differentiation in a

noncanonical NF-κB-dependent manner through transcriptional regulation of key adipogenic

transcription factors. These results are the first to describe NIK regulation of adiposity through

NF-κB-dependent development signaling, as well as NF-κB-independent regulation of local-

tissue metabolism and systemic energy expenditure, suggesting that systemic mitochondrial

metabolic defects are an underlying component of immune dysfunction.

2. Results

2.1 NIK KO mice have reduced adiposity

Whole body composition analyses were performed using dual-energy x-ray

absorptiometry (DEXA) imaging to evaluate potential alterations in systemic body composition

due to NIK deficiency. Mice were utilized between the ages of 2-4 months, that exhibited a healthy

deposition. We observed that NIK KO mice displayed a significant reduction in overall fat mass

and considerably smaller visceral and subcutaneous adipose tissue depots (Figure 1A). Further

assessment of whole body composition using EchoMRI^{TM}, demonstrated that both NIK KO males

and females had a significant reduction in overall fat mass and a higher lean mass ratio relative

to body weight (Figure 1B,C). Visceral gonadal white adipose tissue (gWAT) depots were

substantially reduced in male and female NIK KO mice compared to WT and HET mice (Figure

1D,E, Supp. Figure 1A). Similarly, subcutaneous inguinal white adipose tissue (ingWAT) was

reduced in NIK KO mice (Supp. Figure 1B). While WT mice increased in fat mass as they aged

from 2 to 4 months, NIK KO mice maintained a reduced fat mass during this time period (Figure

1F, Supp. Figure 1C,D), and exhibited no signs of infection. Although NIK KO mice gained less

fat, this was not due to behavioral changes as they consumed more food and water with reduced

physical activity than their WT littermates (Supp. Figure 1C,D). Furthermore, though a reduction

in adiposity was notable even at 2 months age (Supp. Fig. 1F), NIK KO mice did not significantly

differ in weight compared to WT mice until 15 weeks of age in males, possibly when the rate of

fat mass development in WT mice surpassed the rate of body growth and adipose development

in NIK KO mice (Supp. Figure 1G,H). Oil Red O staining of liver sections revealed no differences

in exogenous liver lipid accumulation between WT and NIK KO mice (Supp. Figure 1I), which can

occur due to defects in adipocyte function and lipolysis [37,38], suggesting a role for NIK in the

adipose tissue development.

Figure 1: NIK KO mice have reduced adiposity

(A) DEXA scans of chow fed male WT and NIK KO mice at 2 months, displaying overall fat map

(left) and overall low energy map (right). **(B)** EchoMRI™ measurements of overall fat and **(C)** lean

mass of male and female mice normalized to body weight. WT n= 12 females and 18 males, KO

n=12 females and 11 males. **(D)** Gonadal adipose tissue from male or female mice. **(E)** Weight

of gonadal fat between WT and NIK KO male and female mice normalized to body weight. WT

n= 27 females and 36 males, KO n= 26 females and 18 males. **(F)** Fat mass ratio from body

composition data of female and male WT and NIK KO mice at 2 and 4 months of age. At 2-

months-old WT n=17 females and 17 males, KO n= 9 females and 9 males. At 4-months-old WT

n= 12 females and 18 males, KO n= 12 females and 11 males. **(B-C,E-F)** Data represented as

mean ± SD, Unpaired Student t-test.

167   2.2 NIK regulates mitochondrial respiration and efficiency

We previously demonstrated that NIK localizes to mitochondria, promoting mitochondrial

fission and oxidative metabolism in glioblastoma cells [13]. Similarly, we observed that multipotent

C3H10T1/2 cells, capable of differentiating into adipocytes, also exhibited larger, more fused

mitochondria when NIK expression was ablated. Exogenous expression of human NIK in NIK KO

cells (NIK KO-hNIK) restored mitochondrial morphology similar to that observed in WT cells

(Figure 2A,B). Similarly, 3T3-L1 NIK KO preadipocytes also exhibited a fused mitochondria

phenotype. However, cells lacking the downstream noncanonical NF-κB transcription factor, RelB

(RelB KO cells), did not exhibit a fused mitochondrial morphology similar to NIK KO cells,

suggesting this phenotype in is NIK-dependent (Figure 2C). Given the reduction in adiposity and

changes in mitochondria morphology in NIK KO cells, we sought to determine whether NIK

regulates mitochondrial function in adipocytes. Metabolic analysis of 3T3-L1 cells by Seahorse

extracellular flux assay revealed that NIK KO cells exhibited higher basal oxygen consumption

rate (OCR) but with significantly impaired mitochondrial spare respiratory capacity (SRC), and

increased mitochondrial proton leak, characteristic of inefficient mitochondrial metabolism (Figure

2D). Normal SRC and proton leak phenotypes were restored in NIK KO-hNIK cells (Figure 2D,

183 Supp. Fig 1J), demonstrating that the aberrant mitochondrial metabolism is NIK-dependent. RelB

184 KO cells did not have impaired SRC and proton leak, showing that NIK regulation of mitochondrial

185 efficiency is also noncanonical NF-κB-independent (Figure 2D). NIK KO and NIK KO-hNIK cells

186 also showed an increase in basal extracellular acidification rate (ECAR), a readout for glycolysis,

187 while RelB KO cell ECAR was similar to WT (Figure 2E), consistent with NF-κB-independent roles

188 for NIK in adipocyte metabolism. Basally, NIK KO, NIK KO-hNIK and RelB KO cells had similar

189 basal energetic profiles compared with WT cells. However, NIK KO cells exhibited impaired OCR

190 in response to mitochondrial stressors, which was accompanied by a compensatory increase in

191 glycolysis, whereas RelB KO cell metabolism was unaffected (Figure 2F). Exogenous NIK

192 expression rescued OCR elevation under a stressed state in NIK KO cells, demonstrating a

193 requirement for NIK to meet metabolic demand in a stressed state through OXPHOS (Figure 2G).

Figure 2: NIK regulates mitochondrial respiration and efficiency

(A) Representative imaging of C3H10T1/2, mouse stem cells, transfected with Mito-dsRed. **(B)**

Image J quantification of Mito-dsRed transfected WT, NIK KO, and NIK KO-hNIK C3H10T1/2

cells. Data represented as mean ± SD, One-Way ANOVA **(C)** Immunofluorescence imaging of

WT, NIK KO, NIK KO-hNIK, and RelB KO C3H10T1/2 cells with mitochondria (COXIV, green) and nuclear (Hoechst, blue) staining. **(D)** Seahorse extracellular flux analysis of 3T3-L1 preadipocytes oxidative stress test (mitochondria stress test) displaying oxygen consumption rate (OCR) (statistics compared to WT) and **(E)** extracellular acidification rate (ECAR). **(D-E)** OCR and ECAR line graphs are represented as mean ± SEM. Mitochondria stress test dot plots are represented as mean ± SD, One-Way ANOVA. N= 4 independent, biological replicates with 5-8 technical replicates per independent run. **(F)** Cell energy phenotype of 3T3-L1 cells from seahorse analysis showing basal states of cell (open square) by OCR and ECAR to stressed states (closed square). Data represented as mean ± SD. **(G)** Metabolic potential of 3T3-L1 cells from seahorse analysis with contribution by either OXPHOS (OCR) or by glycolysis (ECAR) after FCCP injection (stressed rates). Data represented as mean ± SD, One-Way ANOVA**. (F-G)**. N= 3 independent, biological replicates with 5-8 technical replicates per independent run.

2.3 Glycolytic dependence increases in response to metabolic demand in NIK KO adipose tissue

To evaluate whether NIK regulates adipocyte metabolism in the context of its native microenvironment, we analyzed OCR and ECAR in *ex vivo* adipose tissue from WT and NIK KO mice. Although basal OCR was not significantly altered, NIK KO ingWAT and gWAT depots exhibited impaired maximal respiration and reduced SRC, increased proton leak and decreased ATP production, similar to adipocyte cells (Figure 3A). We also observed robustly elevated ECAR in both NIK KO adipose depots, consistent with metabolic reprogramming to meet energetic demands and compensate for inefficient OXPHOS (Figure 3B). Both adipose depots were able

to increase glycerol levels in response to isoproterenol, a β-agonist and lipolytic stimulant, suggesting that NIK KO adipose tissues are functional, with gWAT displaying enhanced lipid turnover compared to WT (Figure 3C). Together, these findings suggest that loss of NIK in adipocytes and adipose tissue resulted in mitochondrial dysfunction exhibited by impaired SRC and elevated proton leak, causing an increase in energetic demand that is met by enhanced glycolytic metabolism.

Figure 3: Glycolytic dependence increases in response to metabolic demand in NIK KO mice

(A) Seahorse extracellular flux analysis of *ex vivo* mouse gonadal (gWAT) or inguinal WAT (ingWAT) with an oxidative stress test (Mitochondria Stress Test). **(B)** Seahorse extracellular flux

analysis of *ex vivo* mouse gonadal or inguinal WAT with Glycolysis Stress Test. **(A-B)** Data

represented as mean ± SEM for line graphs and mean ± SD for dot plots. WT n=4 females and 5

males, KO n= 4 females and 3 males. Data normalized by protein. **(C)** Glycerol levels of gonadal

or inguinal WAT with lipolysis stimulated with isoproterenol and measured prior to treatment

(untreated) and after 18-hour incubation. Data represented as mean ± SD, Sidak's Multiple

Comparison Test. Data normalized by tissue weight.

2.4 Sex dimorphic metabolic differences are exhibited in NIK KO mice

We next sought to evaluate whether NIK-dependent metabolism observed in cellular and

adipose tissue impacted systemic metabolism. Using indirect calorimetry metabolic cages, whole

body analyses of WT and NIK KO mice were performed across light and dark cycles to analyze

respiratory rates and energy expenditure, along with feed and water intake as well as activity.

Overall, female NIK KO mice displayed higher oxygen consumption and carbon dioxide

production rate throughout both periods of the day compared to their WT counterparts, while NIK

KO male systemic metabolic rates were not significantly altered (Figure 4 A, B). By measuring

respiration exchange rates (RER), substrate utilization for oxidation can be estimated. Glucose

oxidation has the highest respiration exchange rate, followed by protein, and then lipid oxidation

[29]. Tracking changes in oxidation preference between inactive (light) and active (dark) cycles

by RER measurements revealed that at 2 months of age, both WT and NIK KO mice similarly

utilize lipid oxidation during their inactive states (Supp. Figure 2D). However, as NIK KO mice

age, glucose or protein utilization for oxidation increases during their inactive periods, similar to

RER in their active periods. In contrast, WT mice maintain lower RER in their inactive states,

indicative of lipid oxidation. During active periods, WT and NIK KO RER increase similarly at both

ages (Figure 4C). Moreover, NIK KO mice displayed a higher overall energy expenditure,

particularly in females compared with males (Figure 4D). In addition to increased metabolic

output, NIK KO mice also exhibited higher temperatures at 2 months of age (Figure 4E), and

female mice maintained elevated temperatures at 4 months (Supp. Figure 2A). Consistent with

increased temperature and energy expenditure, we observed higher brown adipose tissue in both

female and male NIK KO mice (Figure 4F). While male and female mice displayed adipose

specific increases in metabolism from *ex vivo* extracellular flux analysis, only female mice

exhibited systemic differences on a chow diet. This trend for an overall increase in whole body

metabolic activity in the NIK KO mice is observable in aging mice (Supp. Figure 2).

Figure 4: Sex dimorphic metabolic differences are exhibited in NIK KO mice

(A-D) Systemic metabolic data collected from 4-month-old, chow fed male and female *Map3k14* mice individually housed in TSE Phenomaster metabolic cages. Morning (light) and night (dark) analysis from a 24-hour time period of **(A)** oxygen consumption, **(B)** CO_2 production, **(C)** respiratory exchange rate (CO_2/O_2), and **(D)** caloric energy expenditure. **(E)** Metabolic cage temperature with mice at 2 months of age. **(F)** Brown adipose tissue weights from male and female, chow fed WT and NIK KO mice at 4 months of age. Data represented as mean ± SD Males; WT n=25 females and 33 males, KO n=22 females and 19 males. Line graphs represented as mean ± SEM, bar graphs represented as mean ± SD. **(A-D)** Females WT n=5 females and 4 males, KO n=6 females and 3 males. **(E)** WT n= 6 females and 5 males, KO n=5 females and 3 males. **(A-F)** Data analyzed by Unpaired Student t-test.

2.5 NIK promotes adiposity in response to overnutrition with a high-fat diet

Next, we investigated whether NIK plays a role in regulating systemic metabolism in response to nutritional stress. After 2-3 months on a high-fat diet (HFD), both male and female NIK KO mice exhibited significantly less weight gain than their WT counterparts (Figure 5A, B, C). Whole body composition analysis by EchoMRI demonstrated that NIK KO mice maintained lower fat mass and a higher lean mass ratio by body weight compared to WT mice (Figure 5D, E). Overall, NIK KO mice gained about 3x less fat than WT mice on a HFD compared with standard chow (Figure 5F). Analysis of individual adipose depots revealed that NIK KO mice on a HFD exhibited a 50% decrease in gWAT tissue, along with a significant reduction in the subcutaneous depot, ingWAT, compared to WT mice (Figure 5G,H). In addition to exhibiting reduced adiposity, NIK KO mice were more efficient at clearing glucose from the blood in glucose tolerance tests

compared to WT littermates, suggesting an increase in insulin sensitivity (Figure 5I). Insulin

tolerance testing confirmed this, demonstrating that NIK KO mice also cleared glucose more

rapidly than WT mice after an administration of insulin (Figure 5J). Analysis of adipogenic gene

expression from the inguinal WAT of HFD mice also revealed a reduction in fatty acid binding

protein 4 (*FABP4*; Ap2), and an increase in GLUT4 and *GPD1* (Supp. Figure 3A,B), which is

consistent with both an increase in glucose clearance of NIK KO mice and higher glycolytic

demands of NIK KO adipose tissue. Moreover, glycogen synthase expression in the livers of NIK

KO mice is decreased, suggesting reduced glucose storage compared to WT (Supp. Figure

3C,D). NIK KO mice on a HFD also exhibited reduced lipid levels in the liver compared to WT

mice accompanied by lower gene expression of fatty acid binding protein (*FABP1*) and fatty acid

oxidation genes (*MCAD, CPT1α*) (Supp. Figure 3D,E). Overall, these data demonstrated the

critical role NIK has in facilitating adipose expansion and glucose homeostasis with overnutrition

under a HFD.

Figure 5: NIK promotes adiposity in response to overnutrition with a high-fat diet

(A) Images of male WT and NIK KO mice on HFD. **(B)** Weights of WT and NIK KO male and female mice exposed to a HFD for about 3 months (mice exposed to a HFD in utero up until weaning and then 8-10 weeks after weaning). Data represented as mean ± SEM, Unpaired Student t-test. **(C)** DEXA images of WT and NIK KO male mice on a HFD at about 3 months of age, displaying overall fat map (left) and overall low energy map (right). **(D)** EchoMRI^{TM} data of fat and **(E)** lean mass from female and male WT and NIK KO mice on a HFD normalized to weight. Data represented as mean ± SD, Unpaired Student t-test. **(F)** Fat mass data from EchoMRI^{TM} analysis comparing fat mass ratios from a chow diet to a HFD between WT and NIK KO mice. **(G)** Gonadal white adipose tissue from male mice on a HFD. **(H)** Weights of gonadal and inguinal

white adipose tissue from male and female mice normalized to body weight. Data represented as

mean ± SD, Unpaired Student t-test. **(I)** Glucose tolerance test on WT and NIK KO mice on a

HFD. Data represented at mean ± SEM, Sidak's Multiple Comparisons Test. WT n=8, NIK KO

n=12. **(J)** Insulin tolerance test on WT, HET and NIK KO mice on a HFD. Data represented at

mean ± SEM, Sidak's Multiple Comparisons Test. WT/HET n=10, NIK KO n=6.

2.6 NIK is required for metabolic homeostasis in response to chronic dietary stress

Extracellular flux analyses were then performed on *ex vivo* adipose tissue from mice

subjected to the chronic overnutrition with a HFD in utero through the age of 3 months. Adipose

tissue from NIK KO mice on a HFD showed a striking increase in basal OCR and ATP

production (Figure 6A). NIK KO ingWAT had significantly decreased SRC, while proton leak

was consistently increased in both NIK KO depots (Figure 6A), suggesting elevated, but

inefficient, mitochondrial metabolism, similar to results observed in NIK KO preadipocytes and

ex vivo NIK KO adipose tissue from chow-fed mice. Analysis of glycolytic metabolism also

demonstrated similar results to chow-fed mice, with NIK KO adipose tissue exhibiting

significantly higher basal ECAR rates and higher levels of glycolysis (Figure 6B). We also

observed elevated systemic metabolism in NIK KO mice on a HFD. Although we observed

some sex dimorphic metabolic dysregulation in NIK KO mice under a chow diet, with the chronic

stress of a HFD, both male and female NIK KO mice displayed elevated oxygen consumption

and CO_2 production (Supp. Fig. 4A,B) along with higher respiratory exchange rates, particularly

during the light cycle, or inactive period (Figure 6C, Supp. Fig 4C). Notably, robust increases in

energy expenditure and cage temperature were observed in NIK KO mice, demonstrating

considerably elevated metabolic output compared to WT mice at all times of the day with HFD

overnutrition (Figure 6D, Supp. Fig. 4D,E). Notably, leaner NIK KO mice did not exhibit

increased physical activity (Supp. Fig. 4F), or reduced water or feed intake (Supp. Fig. 4G,H).

Overall, our findings revealed that under either basal or nutrient stressed conditions NIK KO

mice exhibited elevated metabolism with higher energy expenditure but have inefficient

mitochondrial metabolism through impaired SRC and increased proton leak. Furthermore,

increased glycolysis and glucose utilization persisted to meet energetic demands under the

stress of a HFD, demonstrating an important role for NIK in metabolic homeostasis.

Figure 6: NIK is required for metabolic homeostasis in response to chronic dietary stress

(A) Seahorse extracellular flux analysis of *ex vivo* mouse gonadal or inguinal WAT with either

oxidative stress test (Mitochondria Stress Test) or with **(B)** Glycolysis Stress Test. **(A,B)** Data

represented as mean ± SEM for line graphs and mean ± SD for dot plots, Unpaired Student t-

test. WT n=4 females and 5 males , KO n=6 females and 7 males. **(C,D)** Indirect calorimetry

data collected from TSE Phenomaster cages of individually housed male and female *Map3k14*

mice on a HFD at about 3 months of age. Morning and night analysis from a 24-hour time period

of **(C)** respiratory exchange rate (CO_2/ O_2), and **(D)** caloric energy expenditure. WT n= 7

females and 5 males, KO n=8 females and 5 males. Data represented as mean ± SD, Unpaired

Student t-test.

2.7 A role for noncanonical NF-κB-dependent signaling in promoting adiposity

Given previous studies suggesting a role for NF-κB in adipocyte differentiation and

development, we investigated whether NIK-dependent noncanonical NF-κB activity was involved

in promoting adiposity. Compared to control cells, NIK KO C3H10T1/2 and 3T3-L1 cells were

severely impaired in their ability to form mature adipocytes, as demonstrated by staining of lipid

droplets. Expression of a wild-type human NIK construct in NIK KO cells (NIK KO-hNIK) restored

adipogenesis to the same extent as control cells (Figure 7A, Supp. Figure 5A,B). We also

observed that primary bone marrow-derived stem cells (MSCs) isolated from NIK KO mice were

impaired in their ability to differentiate into mature adipocytes compared to MSCs isolated from

WT mice (Supp. Figure 5C). Furthermore, treatment of WT C3H10T1/2 cells with a NIK-specific

inhibitor, B022, impeded adipocyte differentiation similar to cells lacking NIK (Supp. Figure 5D,E)

[39]. These results support a crucial, cell-intrinsic role for NIK in adipocyte development.

Next, we evaluated if NIK regulation of adipogenesis required downstream NF-κB

signaling. We observed that RelB KO 3T3-L1 cells exhibited impaired adipocyte differentiation

and overexpression of IKKα and RelB in NIK KO cells (NIK KO-IKKα, NIK KO-RelB) restored

adipocyte differentiation similar to NIK (NIK KO-hNIK). In contrast, overexpression of the

canonical NF-κB transcription factor RelA in NIK KO or RelB KO cells was unable to restore

adipocyte differentiation and RelA KO cells did not exhibit impaired adipogenesis (Figure 7B,

Supp. Figure 6A). Additionally, RelA phosphorylation at Ser536 (p-RelA-S536), a marker of its

transcriptional activation, decreased significantly 4-6 days after induction of adipocyte

differentiation. Conversely, expression of the noncanonical NF-κB protein RelB, increased

throughout adipocyte differentiation (Supp. Figure 6B). Furthermore, activation of the canonical

NF-κB pathway with TNFα inhibited adipocyte differentiation in WT C3H10T1/2 cells, whereas

preferential activation of the noncanonical NF-κB pathway with TWEAK did not inhibit

adipogenesis (Supp. Figure 6C,D).

Next, we observed that NIK and RelB were required for the induction of the essential

adipogenic transcription factors PPARγ and C/EBPα [40] (Figure 7C, Supp. Figure 6E). Ectopic

expression of NIK or RelB in NIK KO cells rescued their ability to respond to the PPARγ agonist

ETYA (eicosatetraynoic acid), and increased PPARγ and C/EBPα expression (Figure 7D, Supp.

Figure 6F). PPARγ was also observed to be decreased in NIK KO liver extracts (Supp. Figure

6G). Additionally, expression of the PPARγ-regulated, adipocyte-specific genes adiponectin

(*ADIPOQ*), glucose transporter 4 (GLUT4, *SLC2A4*), and glycerol-3-phosphate dehydrogenase

(*GPD1*), were significantly impaired in NIK KO and RelB KO cells, basally and in response to a

PPARγ agonist. Expression of PPARγ and adipogenic genes, as well as functional adipocyte

differentiation was restored in NIK KO cells by re-expression of NIK or RelB (Figure 7E).

Additionally, the overexpression of PPARγ or C/EBPα rescued adipocyte differentiation in NIK

KO C3H10T1/2 and 3T3-L1 cells, demonstrating that NIK KO cells fail to undergo adipogenesis

due to their inability to increase expression of these key adipogenic transcription factors (Figure

7F). Taken together, these findings indicate that NIK promotes adiposity through noncanonical

NF-κB/RelB-dependent regulation of PPARγ and C/EBPα expression.

Figure 7: A role for noncanonical NF-κB-dependent signaling in promoting adiposity

(A) Fluorescent BODIPY lipid staining of C3H10T1/2 and 3T3-L1 cells showing a significant

reduction in adipogenesis with the loss of NIK which is rescued in NIK KO-hNIK cells. **(B)** Oil Red

O staining of differentiated 3T3-L1 cells lacking or over expressing various

noncanonical/canonical NF-κB proteins. **(C)** PPARγ and C/EBPα expression in undifferentiated

vs adipogenic treated in NIK KO and RelB KO 3T3-L1 cells compared to WT control cells. **(D)**

PPARγ and C/EBPα expression in undifferentiated vs adipogenic treated 3T3-L1 cells with re-

expression of human NIK or overexpression of RelB in NIK KO cells (NIK KO-hNIK, NIK KO-

RelB). **(E)** qPCR analysis of adipocyte genes (*ADIPOQ, SLC2A4, GPD1*) in ETYA/insulin treated

C3H10T1/2 cells. Data represented as mean ± SD, Tukey's Multiple Comparison Test. **(F)** Oil

Red O staining of differentiated NIK KO or NIK KO with overexpression of PPARγ or C/EBPα in

C3H10T1/2 and 3T3-L1 cells.

3. Discussion

Maintenance of energy balance is critical for normal physiological processes and overall

health status. In this study, we demonstrate that NIK is required to for metabolic homeostasis

through control of mitochondrial respiration and fitness, establishing a previously unrecognized

role for NIK in controlling cellular, tissue, and systemic metabolism in response to nutritional

stress. Consistent with *in vivo* findings, *in vitro* and *ex vivo* analysis revealed that NIK KO

preadipocytes and tissue exhibited impaired mitochondrial spare respiratory capacity and

elevated proton leak. Furthermore, we observed a compensatory upregulation of glycolysis in the

absence of NIK to meet energy demands (see Figs. 2, 3 & 6). This metabolic reprogramming has

parallels to the Warburg effect in cancer cells [41], which facilitates rapid generation of ATP using

glucose as a carbon source for anabolic pathways while controlling ROS production and redox

homeostasis [42]. Our results are also consistent with the use of glucose and glycolysis to reduce

ATP input for substrate catabolism, especially in a stressed state [29–31].

A key finding of our study is our observation that NIK-deficient exhibit higher energy

expenditure resulting in decreased adiposity, even under the chronic stress of overnutrition

caused by a HFD, which has been shown to induce oxidative stress and promote mitochondrial

dysfunction. We note that NIK KO mice were housed similarly to other immunodeficient animals

and analyzed at 2-4 months of age when they appeared healthy and exhibited no sign of infection

or dermatitis (see Figure 5A). Our data are consistent with a previous report demonstrating that

NIK activity is increased in the livers of genetic (*ob/ob)* or HFD-fed obese mice, and liver-specific

loss of NIK increases glucose metabolism [16,17]. Similarly, another group demonstrated that

NIK overexpression in pancreatic islet cells decreases insulin secretion and glucose homeostasis

[43], consistent with the increased glucose and insulin tolerance found in our NIK KO mice. In this

study, we analyzed adipose tissue as a key metabolic tissue that impacts local and systemic

metabolism. While male and female NIK KO mice exhibited reduced adiposity and elevated

metabolic demand, only females exhibited systemic metabolic changes under a chow-fed diet,

including increased energy expenditure and temperature. Elevated thermogenesis is consistent

with increased mitochondrial proton uncoupling due to higher proton leak observed in NIK KO

cells, as well as the increased brown adipose deposition in NIK KO mice (see Fig 4). Notably,

when challenged with a HFD, both genders exhibited a significant increase in energy expenditure

and higher glucose oxidation than WT mice, demonstrating that with overnutrition, glycolytic

adaptation is fueled to support the increase in energetic demand. Dimorphic sex differences in

energy expenditure might be attributed to distinct hormone-dependent metabolic or immune

changes between males and females. For example, although females are more prone to adipose

deposition than males, estrogen increases insulin sensitivity thereby protecting female mice from

diet-induced obesity [44–46], and estrogen also enhances mitochondrial function in adipocytes

[47]. Furthermore, a recent study demonstrated that RANKL/NIK signaling induced

postmenopausal obesity in ovariectomized Nik[aly/aly] mice, which exhibited reduced adipocyte

hypertrophy and reduced lipid accumulation, consistent with our findings that NIK promotes

adiposity and adipose development [48].

Interestingly, our findings revealed that NIK regulates adiposity through both NF-κB-

independent and NF-κB-dependent mechanisms. Notably, NIK regulates mitochondrial dynamics

and metabolism independently of downstream NF-κB RelB (see Figure 2). However, it promotes

adipocyte differentiation in noncanonical NF-κB-dependent manner, whereas canonical NF-κB

activity suppresses adipogenesis (see Figure 7). These observations are consistent with previous

data showing that activation or overexpression of the canonical NF-κB/RelA pathway inhibits

adipogenesis, and that RelA itself is inhibited by adipogenic-promoting PPARγ agonists *in vitro*

and *in vivo* [49,50]. Furthermore, one study demonstrated an increase of RelB and p52 during

adipocyte differentiation [51], while a separate study demonstrated that mice with adipose-specific

knockout of Tank-binding kinase 1 (TBK1), a repressor of NIK stability, exhibited a reduction in

adipose tissue and adipocyte size under a prolonged HFD [52]. However, a cooperative role for

the canonical and noncanonical NF-κB pathways in promoting adipocyte differentiation was

previously reported in a study of adipose-specific RelB knockout mice in response to lymphotoxin-

β-receptor (LTβR) activation [53]. Our results suggest that there are likely signal-specific roles for

RelB in regulating PPARγ expression and adipogenesis. Moreover, NIK may have RelB-

independent, or non-cell autonomous effects on adipogenesis and adipose development. In the

context of these findings, the signals responsible for activating NIK during adipocyte development

and expansion under a HFD warrant further study.

Further investigation is needed to understand if NIK has adipose cell-intrinsic effects *in*

vivo, as well as assess how other tissue types, such as skeletal muscle which is a major glucose

sink [54], regulate metabolic homeostasis *in vivo*. For example, NIK may impact systemic

metabolism indirectly through its regulation of immune functions, as changes in bacterial flora or

viral infection, which activate NIK/NF-κB signaling, can alter host metabolism to support

replication [55,56]. Furthermore, as a regulator of lymphoid development, NIK deficiency might

affect the metabolic status and recruitment of adipose tissue immune cells that can play critical

roles in regulating tissue homeostasis [14,57]. We note that *in vitro* preadipocytes exhibited

metabolic phenotypes similar to *ex vivo* adipose tissue (see Figs. 2 & 3), demonstrating that NIK

has similar adipocyte cell-intrinsic and tissue-specific metabolic functions.

Our analysis of NIK knockout mice provides new insight into the role of metabolic

dysfunction in the profoundly debilitating phenotypes of primary immunodeficiency diseases

(PIDs). Several genetic defects in the NF-κB signaling pathway are associated with PIDs [58–62],

including recently described patients with loss-of-function mutations in NIK [10,11]. Patients with

PIDs have been observed to have growth defects, mainly in children [63,64]. Risk for infection

have also been linked to factors including body mass index and adipose deposition due to the

immunomodulatory effects of the tissue, but less is understood of the metabolic dynamics in

patients with PIDs [65,66]. An intimate association between primary immunodeficiencies and

primary metabolic diseases is supported by the observation that mitochondrial disease patients

manifest significant immunological defects, [67,68], and some studies have linked inborn errors

of metabolism (IEM) as mimicking or exacerbating immune defects [69]. Overall, our findings are

consistent with the increasing appreciation that mitochondrial functions are important for

integrating metabolic cues and maintaining systemic metabolic health. Moreover, our work

highlights an important role for NIK in regulating metabolic homeostatic and adaptive adipose

remodeling mechanisms in response to a variety of disease contexts, including chronic stress due

to aging, immunodeficiency or overnutrition.

Author Contributions: Conceptualization: K.M.P., D.W.L., R.S.; Data Acquisition: K.M.P.,

D.W.L., J.K.; Writing—original draft preparation, K.M.P.; Writing—review and editing, K.M.P. and

R.S.; Visualization: K.M.P., R.S.; Funding acquisition: R.S.; Supervision: R.S. All authors have

read and agreed to the publication of the manuscript.

Funding: NIH-1R01NS082554 to R.S.

Acknowledgments: We thank Dr. Joseph M. Rutkowski for 3T3-L1 cells, Dr. Larry Suva and Dr.

Kirby for assistance with DEXA imaging, Dr. David Threadgill and his lab, including Dr. Alexandra

Trott and Orion Hicks at the Texas A&M Rodent Preclinical Phenotyping Core, for assistance with

the EchoMRI^{TM} machine and TSE Phenomaster^{TM} metabolic cages, and Texas A&M Health

Science Center COM-CAF core for use of the Seahorse machine.

Conflicts of Interest: The authors declare no conflicts of interest.

4. Materials and Methods

4.1 Animal Procedures & *Ex Vivo* **Work**

All animal experiments were done in accordance with animal use protocol (2019-0102) with

approved IACUC guidelines. *Map3k14* mice were purchased from Jackson lab (B6N.129-

Map3k14tm1Rds/J) and maintained by heterozygous breeding. Mice are housed on

hypoallergenic, alpha dry bedding with weekly cage changes to minimize dermatitis. Chow diet

contained 4% fat while HFD contained 45% fat (Lab Supply 58125). Mice analyzed on a HFD

were weaned from adult females on a HFD and then maintained on a HFD for at least 2 months.

4.2 Cell Culture

Mesenchymal stem cell line and preadipocyte cell line, C3H10T1/2 and 3T3-L1 were cultured in

DMEM supplemented with 10% FBS, 100 U/ml penicillin and 0.1mg/ml streptomycin (Thermo

Fisher Scientific, Waltham, MA). 293-T cells were obtained from ATCC (www.atcc.org) and

cultured in DMEM supplemented with 10% FBS, 100 U/ml penicillin and 0.1mg/ml streptomycin.

Bone marrow MSC cells were isolated from the femurs of mouse and cultured in DMEM

supplemented with 10% FBS, 100 U/ml penicillin and 0.1mg/ml streptomycin.

4.3 Bone Marrow Cell Isolation

Mouse femurs are collected and sterilized in 70% ethanol and then were crushed in sterile PBS

in a mortar with a pestle. From the crushed femurs, 5mL of DMEM (with 10% FBS and 1%

penicillin/streptomycin) is then used to collected cells into a conical tube through a 70μM filter.

Cells are then centrifuged at 1500rpm for 8 minutes. Media is removed and the cell pellet is

resuspended in 3mL of ACK lysis buffer (Lonza 10-548E) for 2 minutes. ACK cell suspension is

diluted with 10mL of media and pelleted again. The cell pellet is resuspended in 10mL of media

and filtered through a 40μM filter.

4.4 CRISPR-Cas9 Gene Knockout

Oligos encoding guide RNAs for murine NIK and RelB are found in Supplemental Table 1. Each

gRNAs were cloned into Lenti-CrispR-v2 (Addgene, Cambridge, MA) respectively. C3H10T1/2

and 3T3-L1 cells were transduced with a mixture of lentiviruses (described below) carrying the

three murine gRNAs. Puromycin resistant single clonal cells were isolated by serial dilution and

experiments were repeated with at least two clones. Loss of NIK or RelB expression was

confirmed by immunoblot analysis. For controls, cells were transduced with empty LentiCrispR-

V2.

4.5 Lentivirus Production

Lentiviral construct of over-expression constructs (NIK, IKKα, RelB, RelA, PPARγ and C/EBPα)

was obtained from DNASU (Tempe, AZ). 24μg of lentiviral plasmids and 72μg of

polyethyleneimine (Sigma-Aldrich) were used to transfect 293T cells. After 3 days of transfection,

viral supernatant was concentrated 20 fold, to 500μl using Lenti-X Concentrator (Clontech,

Mountain View, CA), and 100µl of concentrated virus was used to infect cells. Stably transduced

cells were selected for 72hr in medium containing 0.6µg/ml puromycin or 6µg/ml blasticidin

(Thermo Fisher).

4.6 Seahorse Extracellular Flux Analysis

4.6.1 3T3-L1 Cells

Metabolic activity was analyzed using a Seahorse XFe96 Analyzer (Agilent, Santa Clara, CA).

For 3T3-L1 cells, forty-thousand cells per well were plated in the collagen-coated (40ng/µL)

Agilent Seahorse XFe96 microplates day before analysis. Mitochondrial Stress Tests were

conducted according to the manufacturer's guidelines. Base media was supplemented with 25mM

glucose (Sigma, G7021, St. Louis, MO), 2mM glutamine (Sigma, G85420, St. Louis, MO), and

1mM pyruvate (Gibco). Inhibitors were used at the following final concentrations (10x inhibitor

was added to injection ports to reach final concentration): 1µM oligomycin A (Sigma, 75351, St.

Louis, MO), 2µM FCCP (Sigma, C2920, St. Louis, MO), and mixture of 0.5µM rotenone (Enzo

Life Sciences, Farmingdale, NY, ALX-350-360) and 0.5µM antimycin A (Sigma, A8674, St. Louis,

MO). For Glycolysis Stress Tests, cells were glucose-starved for 1hr in Seahorse DMEM base

media with 2mM glutamine before analysis. Reagents for injections were used at the following

final concentrations (10x reagent was added to injection ports to reach final concentration): 10mM

glucose, 10uM oligomycin A, 50mM 2-DG (Thermo Fischer Scientific, 50-519-066). After

Seahorse analysis, DNA content was measured using DRAQ5 staining (Thermo Fisher Scientific,

50-712-282) for normalization. Assay was ran in 3 biological replicates with samples ran in 5-8

replicates per assay. Analyses were conducted using Seahorse Wave Controller Software v2.6

and XF Report Generators (Agilent Technologies).

4.6.2 *Ex Vivo* Adipose Tissue

Analysis of metabolic activity of adipose tissue was adapted from [70]. 96 well Seahorse plates

were coated twice with Cell-Tak (50µg/mL, Corning® Cell-Tak™ Cell and Tissue Adhesive, Cat.

No. 354240). Tissue was excised fresh, rinsed well in PBS, and cut into small, >1mg, pieces of

similar size and plated into the center of the wells. For Mitochondria Stress Test, base media was

supplemented with 25 mM glucose, 2 mM glutamine, and 1 mM pyruvate. Inhibitors were used at

the following final concentrations (10x inhibitor was added to injection ports to reach final

concentration): 20µM oligomycin A, 20µM FCCP, and mixture of 20µM rotenone and 20µM

antimycin A. Tissue samples are ran in 5-10 replicates per tissue type per assay. Seahorse

Analyses were done following Agilent guidelines for the mitochondria stress test [71] and energy

phenotyping [72].

For Glycolysis Stress Test base media was supplemented with 2mM glutamine, and reagents for

injections were used at the following final concentrations (10x reagent was added to injection

ports to reach final concentration): 25mM glucose, 20µM oligomycin A, 100mM 2-DG. Run time

for tissue was increased to 30 minutes per injection with four measurements per compound (4

cycles of 3 min mix, 1.5 min wait, 3 min measure). Analyses of glycolysis stress test was done

following Agilent guidelines [73]. Afterwards, tissue was homogenized, lysed, and normalized by

protein.

4.7 Immunofluorescence

For immunofluorescence of C3H10T1/2 mitochondria staining, cells were transfected with 1µg

DsRed2-Mito-7 (Mito-dsRed) (Addgene Plasmid #55838) by lipofectamine (Invitrogen L3000008)

for 48-72hrs or stained with COXIV 1:200 overnight. Cells were then fixed with 4% PFA for 20

minutes at 37°C. After fixation, cells were washed with PBS and then stained with a fluorescent

secondary and/or Hoechst (1:1000) diluted in 1% BSA, .1% Triton-X in PBS. Cells were imaged

by confocal (Nikon TI A1R inverted confocal microscope).

4.8 BODIPY Staining

For immunofluorescence imaging, BODIPY™ 490/509 (4,4-Difluoro-1,3,5,7,8-Pentamethyl-4-Bora-3a,4a-Diaza-s-Indacene) (Sigma Aldrich 790389) was used. 5mM of BODIPY dye was diluted 1:2500 in PBS and incubate on live cells for 10 minutes at 37°C. Afterwards, cells were washed twice with PBS and then fixed in 4% PFA for 20 minutes at 37°C. After fixation, cells were washed twice with PBS and then stained with Hoechst (1:1000) diluted in 1% BSA, .1% Triton-X in PBS for 10 minutes. Cells were imaged by confocal (Nikon TI A1R inverted confocal microscope).

4.9 Lipolysis

About 2-5mg was placed in a 12 well plate in duplicate or triplicate. Tissue was left to sit in 300µL DMEM without FBS for an hour in the incubator. Media was removed for t=0 and isoproterenol was added for a final concentration of 100µM. Tissue was incubated with the isoproterenol for an hour and then media was removed for t=1 timepoint, final samples were taken after overnight incubation. 10µL of each sample at the different points were mixed with 160µL of glycerol reagent (Sigma, F6428) and read at an absorbance of 540nm for glycerol levels.

4.10 Adipocyte Differentiation

Adipocyte differentiation was induced by treating confluent cells for 2 days with 5 mg/ml insulin (Thermo Fisher Scientific) and 10µM ETYA (Santa Cruz Biotech, Dallas, TX) in Dulbecco's modified Eagle's medium containing 10% FBS, and then every 2 days for 6 Days with insulin (5 mg./ml). Treatment with NIK inhibitor B022 was purchased from MEDCHEM EXPRESS (Cat. No. HY-120501) and used at a concentration of 5µM, TNFα (ProSpec CYT-223) and TWEAK (PeproTech 31006) were used at 10ng/mL.

4.11 Oil Red O Staining

For the Oil Red O staining, Oil Red O staining solution (0.5% Oil-Red O in isopropyl alcohol

solution-distilled water [60:40]) was filtered through the Whatman no. 1 filter paper. Cells were

fixed with 10% formaldehyde solution for 30 min at 37°C, then washed with 60% isopropyl alcohol

followed by staining with the filtered Oil Red O solution for 20 min and then washed with distilled

water three times. To measure Oil Red O staining of adipocyte differentiated cells, dye is eluted

with 100% isopropanol for 10 minutes. Eluted dye is transferred to 96 well plate and optical density

measurements were done on Perkin Elmer plate reader at 500nm, .5 sec.

4.12 Immunoblot

Cells were lysed in RIPA lysis buffer (Thermo Fisher Scientific) with protease/phosphatase

inhibitor cocktail (Thermo Fisher Scientific). Protein was mixed with NuPage 4X LDS sample

buffer (Thermo Fisher Scientific) containing .1M DTT and denatured at 100°C for 7 min. Proteins

were separated on 8% ~ 12% SDS-PAGE and transferred to nitrocellulose membranes

(Amersham). The membranes were blocked for 1h with 5% non-fat dry milk in 0.1% Tween-

20/TBS (TBST) and incubated with primary antibodies diluted in blocking buffer at 4°C overnight.

After washing in TBST, membranes were incubated with secondary in BSA for 1 hour at room

temperature. The blots were washed with TBST and developed using Chemiluminescent HRP

Substrate (EMD Millipore) on ChemiDoc MP Imaging System (Bio-Rad) for detection of HRP or

an Odyssey Infrared Imaging system (LI-COR Biosciences) for detection of IRDye fluorescent

dyes.

4.13 Antibodies

Following antibodies were used C/EBPα (CST8178) (Cell Signaling Technology), COXIV

(CST11967s), GPD1 (sc-376219) (Santa Cruz Biotechnology, Dallas, TX), IKKα (CST2682),

NFKB2 (CST4882), NIK (CST4994), p-RelA (p-p65) (CST3033), RelA (p65) (sc-8008), PPARγ

(CST2443), RelB (CST4992), GAPDH (sc137179), and β-actin (sc69879).

4.14 RNA Isolation, cDNA Synthesis, and Quantitative-RT-PCR

Primers used for qPCR are in Supplemental Table 2.

4.14.1 Cells: Total RNA was isolated from cells by Purelink™ RNA Mini Kit (Life Technologies). cDNA was synthesized from 1 μg total RNA using iScript reverse transcription supermix (Bio-Rad, Hercules, CA) following the manufacturer's protocol. Quantitative RT-PCR was performed using iTaq Universal SYBR Green Supermix (Bio-Rad) with StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA).

4.14.2 Tissue: Mouse tissue was frozen with liquid nitrogen and ground to a powder with a pestle in a mortar (a minimum of 50mg of tissue is needed, for WAT 1g may be needed). Tissue was then lysed with Trizol (1mL per gram of tissue), followed by addition of chloroform to centrifuged supernatant (200μL chloroform: 1mL Trizol). Equal amounts of 70% ethanol was added to equal parts of aqueous layer from sample in a new tube and then RNA was then collected in subsequent steps using Invitrogen Purelink RNA mini kit. Zymo DNAse was used with 40-80μL per sample. cDNA was synthesized from 2μg of RNA using iScript reverse transcriptase and buffer mix. Samples were ran in triplicate.

4.15 EchoMRI^TM

EchoMRI^TM 100H machine was used to analyze fat and lean mass of chow and HFD mice at 2 or 4 months of age. Mice were weighed before imaging and imaged in MRI tubes.

4.16 DEXA Imaging

Dual-energy x-ray absorptiometry was used on anesthetize mice to visualize low and high-density tissue. Mice from maintained on either a chow or HFD were used.

4.17 Metabolic Cages

Metabolic analysis on the mice was conducted in TSE Phenomaster™ metabolic cages through

Rodent Preclinical Phenotyping Core at TAMU (https://genomics.tamu.edu/preclinical-

phenotyping/). Mice were weighed the day of and housed individually for 48hrs. Data was

analyzed based on last 24hrs to compensate for mouse adjustment to the new environment.

4.18 Glucose Tolerance Testing

For glucose tolerance testing mice were morning fasted for 6hrs in clean cages. Mice were

weighed, and from tail clip, initial glucose levels are recorded using a glucose meter. The mice

are then intraperitoneally injected with 20% D-glucose at 2g/kg (µL= 10 x BW). Blood glucose (in

mg/dL) is then recorded at t=15, t=30, t=60, t=120 minutes post injection.

4.19 Insulin Tolerance Testing

For insulin tolerance testing mice were morning fasted for 4hrs in clean cages. Mice were weighed

and from tail clip initial glucose levels are recorded using a glucose meter. The mice are then

intraperitoneally injected with .5 U/kg Insulin (µL= 5 x BW of .1 U/mL insulin). Blood glucose (in

mg/dL) is then recorded at t=15, t=30, t=45, t=60, t=90 minutes post injection.

4.20 Statistical Analysis

Statistical Analysis was done using GraphPad PRISM software, specifics on data representation

and tests used for analysis can be found in figure legends. *p ≤.05, ** p ≤.01, *** p ≤.001, ****

p≤.0001. Unpaired student t-tests were ran as two-tailed. All statistically significant analyses were

ran based on a 95% confidence interval.

References

[1] Yin, L., Wu, L., Wesche, H., Arthur, C.D., White, J.M., Goeddel, D.V., et al., 2001. Defective lymphotoxin-beta receptor-induced NF-kappaB transcriptional activity in NIK-deficient mice. Science (New York, N.Y.) 291(5511): 2162–5, Doi: 10.1126/science.1058453.

[2] Xiao, G., Fong, A., Sun, S.C., 2004. Induction of p100 processing by NF-kappaB-inducing kinase involves docking IkappaB kinase alpha (IKKalpha) to p100 and IKKalpha-mediated phosphorylation. J Biol Chem 279(29): 30099–105.

[3] Akemi Matsushima, T.K., 2001. Essential role of nuclear factor (NF)-κB-inducing kinase and inhibitor of κB (IκB) kinase α in NF-κB activation through lymphotoxin β receptor, but not through tumor necrosis factor receptor I. Journal of Experimental Medicine 193: 631–6.

[4] Sun, S.C., 2011. Non-canonical NF-kappaB signaling pathway. Cell Res 21(1): 71–85.

[5] Shinkura, R., Kitada, K., Matsuda, F., Tashiro, K., Ikuta, K., Suzuki, M., et al., 1999. Alymphoplasia is caused by a point mutation in the mouse gene encoding Nf-κb-inducing kinase. Nature Genetics 22(1): 74–7, Doi: 10.1038/8780.

[6] Brightbill, H.D., Jackman, J.K., Suto, E., Kennedy, H., Jones, C., Chalasani, S., et al., 2015. Conditional Deletion of NF-kappaB-Inducing Kinase (NIK) in Adult Mice Disrupts Mature B Cell Survival and Activation. J Immunol 195(3): 953–64.

[7] Li, Y., Wang, H., Zhou, X., Xie, X., Chen, X., Jie, Z., et al., 2016. Cell intrinsic role of NF-kappaB-inducing kinase in regulating T cell-mediated immune and autoimmune responses. Sci Rep 6: 22115.

[8] Lacher, S.M., Thurm, C., Distler, U., Mohebiany, A.N., Israel, N., Kitic, M., et al., 2018. NF-kappaB inducing kinase (NIK) is an essential post-transcriptional regulator of T-cell activation affecting F-actin dynamics and TCR signaling. J Autoimmun 94: 110–21.

[9] Hamdan, T.A., Bhat, H., Cham, L.B., Adomati, T., Lang, J., Li, F., et al., 2020. Map3k14 as a Regulator of Innate and Adaptive Immune Response during Acute Viral Infection. Pathogens (Basel, Switzerland) 9(2), Doi: 10.3390/pathogens9020096.

[10] Willmann, K.L., Klaver, S., Dogu, F., Santos-Valente, E., Garncarz, W., Bilic, I., et al., 2014. Biallelic loss-of-function mutation in NIK causes a primary immunodeficiency with multifaceted aberrant lymphoid immunity. Nat Commun 5: 5360.

[11] Schlechter, N., Glanzmann, B., Hoal, E.G., Schoeman, M., Petersen, B.S., Franke, A., et al., 2017. Exome Sequencing Identifies a Novel MAP3K14 Mutation in Recessive Atypical Combined Immunodeficiency. Front Immunol 8: 1624.

[12] Gu, M., Zhou, X., Sohn, J.H., Zhu, L., Jie, Z., Yang, J.-Y., et al., 2021. NF-κB-inducing kinase maintains T cell metabolic fitness in antitumor immunity. Nature Immunology 22(2): 193–204, Doi: 10.1038/s41590-020-00829-6.

[13] Jung, J.U., Ravi, S., Lee, D.W., McFadden, K., Kamradt, M.L., Toussaint, L.G., et al., 2016. NIK/MAP3K14 Regulates Mitochondrial Dynamics and Trafficking to Promote Cell Invasion. Curr Biol 26(24): 3288–302.

[14] Keeney, J.N., Winters, A., Sitcheran, R., West, A.P., 2022. NF-κB-Inducing Kinase (NIK) Governs the Mitochondrial Respiratory Capacity, Differentiation, and Inflammatory Status of Innate Immune Cells. Immunology.

[15] Kamradt, M.L., Jung, J.-U., Pflug, K.M., Lee, D.W., Fanniel, V., Sitcheran, R., 2021. NIK promotes metabolic adaptation of glioblastoma cells to bioenergetic stress. Cell Death & Disease 12(3): 1–18, Doi: 10.1038/s41419-020-03383-z.

[16] Sheng, L., Zhou, Y., Chen, Z., Ren, D., Cho, K.W., Jiang, L., et al., 2012. NF-kappaB-inducing kinase
(NIK) promotes hyperglycemia and glucose intolerance in obesity by augmenting glucagon action.
Nat Med 18(6): 943–9.
[17] Liu, Y., Sheng, L., Xiong, Y., Shen, H., Liu, Y., Rui, L., 2017. Liver NF-kappaB-Inducing Kinase
Promotes Liver Steatosis and Glucose Counterregulation in Male Mice With Obesity. Endocrinology
158(5): 1207–16.
[18] Li, Y., Chen, M., Zhou, Y., Tang, C., Zhang, W., Zhong, Y., et al., 2020. NIK links inflammation to
hepatic steatosis by suppressing PPARalpha in alcoholic liver disease. Theranostics 10(8): 3579–93.
[19] Pang, G., Xie, J., Chen, Q., Hu, Z., 2014. Energy intake, metabolic homeostasis, and human health.
Food Science and Human Wellness 3(3): 89–103, Doi: 10.1016/j.fshw.2015.01.001.
[20] O'Brien, C.M., Mulukutla, B.C., Mashek, D.G., Hu, W.-S., 2020. Regulation of Metabolic
Homeostasis in Cell Culture Bioprocesses. Trends in Biotechnology 38(10): 1113–27, Doi:
10.1016/j.tibtech.2020.02.005.
[21] Ma, Y., Li, J., 2015. Metabolic Shifts during Aging and Pathology. Comprehensive Physiology 5(2):
667–86, Doi: 10.1002/cphy.c140041.
[22] Picard, M., McEwen, B.S., Epel, E.S., Sandi, C., 2018. An energetic view of stress: Focus on
mitochondria. Frontiers in Neuroendocrinology 49: 72–85, Doi: 10.1016/j.yfrne.2018.01.001.
[23] Li, Y., Jia, A., Wang, Y., Dong, L., Wang, Y., He, Y., et al., 2019. Immune effects of glycolysis or
oxidative phosphorylation metabolic pathway in protecting against bacterial infection. Journal of
Cellular Physiology 234(11): 20298–309, Doi: https://doi.org/10.1002/jcp.28630.
[24] Yuan, X., Liu, Y., Bijonowski, B.M., Tsai, A.-C., Fu, Q., Logan, T.M., et al., 2020. NAD + /NADH redox
alterations reconfigure metabolism and rejuvenate senescent human mesenchymal stem cells in
vitro. Communications Biology 3(1): 1–15, Doi: 10.1038/s42003-020-01514-y.
[25] Duan, K., Liu, Z.-J., Hu, S.-Q., Huo, H.-Y., Xu, Z.-R., Ruan, J.-F., et al., 2018. Lactic acid induces lactate
transport and glycolysis/OXPHOS interconversion in glioblastoma. Biochemical and Biophysical
Research Communications 503(2): 888–94, Doi: 10.1016/j.bbrc.2018.06.092.
[26] Bar-Even, A., Flamholz, A., Noor, E., Milo, R., 2012. Rethinking glycolysis: on the biochemical logic
of metabolic pathways. Nature Chemical Biology 8(6): 509–17, Doi: 10.1038/nchembio.971.
[27] Rodic, S., Vincent, M.D., 2018. Reactive oxygen species (ROS) are a key determinant of cancer's
metabolic phenotype. International Journal of Cancer 142(3): 440–8, Doi:
https://doi.org/10.1002/ijc.31069.
[28] Li, X., Jiang, Y., Meisenhelder, J., Yang, W., Hawke, D.H., Zheng, Y., et al., 2016. Mitochondria-
Translocated PGK1 Functions as a Protein Kinase to Coordinate Glycolysis and the TCA Cycle in
Tumorigenesis. Molecular Cell 61(5): 705–19, Doi: 10.1016/j.molcel.2016.02.009.
[29] Even, P.C., Nadkarni, N.A., 2012. Indirect calorimetry in laboratory mice and rats: principles,
practical considerations, interpretation and perspectives. American Journal of Physiology-
Regulatory, Integrative and Comparative Physiology 303(5): R459–76, Doi:
10.1152/ajpregu.00137.2012.
[30] Mizock, B.A., 1995. Alterations in carbohydrate metabolism during stress: A review of the
literature. The American Journal of Medicine 98(1): 75–84, Doi: 10.1016/S0002-9343(99)80083-7.
[31] Livesey, G., Elia, M., 1988. Estimation of energy expenditure, net carbohydrate utilization, and net
fat oxidation and synthesis by indirect calorimetry: evaluation of errors with special reference to
the detailed composition of fuels. The American Journal of Clinical Nutrition 47(4): 608–28, Doi:
10.1093/ajcn/47.4.608.
[32] Marchetti, P., Fovez, Q., Germain, N., Khamari, R., Kluza, J., 2020. Mitochondrial spare respiratory
capacity: Mechanisms, regulation, and significance in non-transformed and cancer cells. The FASEB
Journal 34(10): 13106–24, Doi: https://doi.org/10.1096/fj.202000767R.

[33] Sriskanthadevan, S., Jeyaraju, D.V., Chung, T.E., Prabha, S., Xu, W., Skrtic, M., et al., 2015. AML cells
have low spare reserve capacity in their respiratory chain that renders them susceptible to
oxidative metabolic stress. Blood 125(13): 2120–30, Doi: 10.1182/blood-2014-08-594408.
[34] van der Windt, G.J.W., Everts, B., Chang, C.-H., Curtis, J.D., Freitas, T.C., Amiel, E., et al., 2012.
Mitochondrial Respiratory Capacity Is a Critical Regulator of CD8+ T Cell Memory Development.
Immunity 36(1): 68–78, Doi: 10.1016/j.immuni.2011.12.007.
[35] Haas, R.H., Parikh, S., Falk, M.J., Saneto, R.P., Wolf, N.I., Darin, N., et al., 2007. Mitochondrial
disease: a practical approach for primary care physicians. Pediatrics 120(6): 1326–33, Doi:
10.1542/peds.2007-0391.
[36] Parikh, S., Goldstein, A., Koenig, M.K., Scaglia, F., Enns, G.M., Saneto, R., et al., 2015. Diagnosis and
management of mitochondrial disease: a consensus statement from the Mitochondrial Medicine
Society. Genetics in Medicine: Official Journal of the American College of Medical Genetics 17(9):
689–701, Doi: 10.1038/gim.2014.177.
[37] Mota, M., Banini, B.A., Cazanave, S.C., Sanyal, A.J., 2016. Molecular mechanisms of lipotoxicity and
glucotoxicity in nonalcoholic fatty liver disease. Metabolism 65(8): 1049–61.
[38] Pan, X., Wang, P., Luo, J., Wang, Z., Song, Y., Ye, J., et al., 2015. Adipogenic changes of hepatocytes
in a high-fat diet-induced fatty liver mice model and non-alcoholic fatty liver disease patients.
Endocrine 48(3): 834–47.
[39] Li, Z., Li, X., Su, M.B., Gao, L.X., Zhou, Y.B., Yuan, B., et al., 2020. Discovery of a Potent and Selective
NF-kappaB-Inducing Kinase (NIK) Inhibitor That Has Anti-inflammatory Effects in Vitro and in Vivo. J
Med Chem 63(8): 4388–407.
[40] Rosen, E.D., 2002. C/EBPalpha induces adipogenesis through PPARgamma : a unified pathway.
Genes & Development 16(1): 22–6, Doi: 10.1101/gad.948702.
[41] Calabrese, C., Iommarini, L., Kurelac, I., Calvaruso, M.A., Capristo, M., Lollini, P.-L., et al., 2013.
Respiratory complex I is essential to induce a Warburg profile in mitochondria-defective tumor
cells. Cancer & Metabolism 1(1): 11, Doi: 10.1186/2049-3002-1-11.
[42] Liberti, M.V., Locasale, J.W., 2016. The Warburg Effect: How Does it Benefit Cancer Cells? Trends in
Biochemical Sciences 41(3): 211–8, Doi: 10.1016/j.tibs.2015.12.001.
[43] Li, X., Jia, L., Chen, X., Dong, Y., Ren, X., Dong, Y., et al., 2018. Islet α-cell Inflammation Induced By
NF-κB inducing kinase (NIK) Leads to Hypoglycemia, Pancreatitis, Growth Retardation, and
Postnatal Death in Mice. Theranostics 8(21): 5960–71, Doi: 10.7150/thno.28960.
[44] Mauvais-Jarvis, F., 2015. Sex differences in metabolic homeostasis, diabetes, and obesity. Biology
of Sex Differences 6, Doi: 10.1186/s13293-015-0033-y.
[45] Riant, E., Waget, A., Cogo, H., Arnal, J.-F., Burcelin, R., Gourdy, P., 2009. Estrogens Protect against
High-Fat Diet-Induced Insulin Resistance and Glucose Intolerance in Mice. Endocrinology 150(5):
2109–17, Doi: 10.1210/en.2008-0971.
[46] Handgraaf, S., Riant, E., Fabre, A., Waget, A., Burcelin, R., Liere, P., et al., 2013. Prevention of
Obesity and Insulin Resistance by Estrogens Requires ER Activation Function-2 (ER AF-2), Whereas
ER AF-1 Is Dispensable. Diabetes 62(12): 4098–108, Doi: 10.2337/db13-0282.
[47] Bauzá-Thorbrügge, M., Rodríguez-Cuenca, S., Vidal-Puig, A., Galmés-Pascual, B.M., Sbert-Roig, M.,
Gianotti, M., et al., 2019. GPER and ERα mediate estradiol enhancement of mitochondrial function
in inflamed adipocytes through a PKA dependent mechanism. The Journal of Steroid Biochemistry
and Molecular Biology 185: 256–67, Doi: 10.1016/j.jsbmb.2018.09.013.
[48] Mori, K., Mizokami, A., Sano, T., Mukai, S., Hiura, F., Ayukawa, Y., et al., 2022. RANKL elevation
activates the NIK/NF-κB pathway, inducing obesity in ovariectomized mice. Journal of
Endocrinology 254(1): 27–36, Doi: 10.1530/JOE-21-0424.

[49] Tang, T., Zhang, J., Yin, J., Staszkiewicz, J., Gawronska-Kozak, B., Jung, D.Y., et al., 2010. Uncoupling
898   of inflammation and insulin resistance by NF-kappaB in transgenic mice through elevated energy
899   expenditure. J Biol Chem 285(7): 4637–44.
900   [50] Pascal Peraldi, M.X., 1997. Thiazolidinediones Block Tumor Necrosis Factor-a-induced Inhibition of
901   insulin signaling. J. Clin. Invest. Volume 100(7): 1863–9.
902   [51] Berg, A.H., Lin, Y., Lisanti, M.P., Scherer, P.E., 2004. Adipocyte differentiation induces dynamic
903   changes in NF-κB expression and activity. American Journal of Physiology-Endocrinology and
904   Metabolism 287(6): E1178–88, Doi: 10.1152/ajpendo.00002.2004.
905   [52] Zhao, P., Wong, K. in., Sun, X., Reilly, S.M., Uhm, M., Liao, Z., et al., 2018. TBK1 at the Crossroads of
906   Inflammation and Energy Homeostasis in Adipose Tissue. Cell 172(4): 731-743.e12, Doi:
907   10.1016/j.cell.2018.01.007.
908   [53] Weidemann, A., Lovas, A., Rauch, A., Andreas, N., von Maltzahn, J., Riemann, M., et al., 2016.
909   Classical and alternative NF-kappaB signaling cooperate in regulating adipocyte differentiation and
910   function. Int J Obes (Lond) 40(3): 452–9.
911   [54] Merz, K.E., Thurmond, D.C., 2020. Role of Skeletal Muscle in Insulin Resistance and Glucose
912   Uptake. Comprehensive Physiology, John Wiley & Sons, Ltd p. 785–809.
913   [55] Eisenreich, W., Rudel, T., Heesemann, J., Goebel, W., 2019. How Viral and Intracellular Bacterial
914   Pathogens Reprogram the Metabolism of Host Cells to Allow Their Intracellular Replication.
915   Frontiers in Cellular and Infection Microbiology 9, Doi: 10.3389/fcimb.2019.00042.
916   [56] Sanchez, E.L., Lagunoff, M., 2015. Viral activation of cellular metabolism. Virology 479–480: 609–
917   18, Doi: 10.1016/j.virol.2015.02.038.
918   [57] DeFuria, J., Belkina, A.C., Jagannathan-Bogdan, M., Snyder-Cappione, J., Carr, J.D., Nersesova, Y.R.,
919   et al., 2013. B cells promote inflammation in obesity and type 2 diabetes through regulation of T-
920   cell function and an inflammatory cytokine profile. Proceedings of the National Academy of
921   Sciences of the United States of America 110(13): 5133–8, Doi: 10.1073/pnas.1215840110.
922   [58] Smith, T., Cunningham-Rundles, C., 2019. Primary B-Cell Immunodeficiencies. Human Immunology
923   80(6): 351–62, Doi: 10.1016/j.humimm.2018.10.015.
924   [59] Wang, H.-Y., Ma, C.A., Zhao, Y., Fan, X., Zhou, Q., Edmonds, P., et al., 2013. Antibody deficiency
925   associated with an inherited autosomal dominant mutation in TWEAK. Proceedings of the National
926   Academy of Sciences of the United States of America 110(13): 5127–32, Doi:
927   10.1073/pnas.1221211110.
928   [60] Losi, C.G., Silini, A., Fiorini, C., Soresina, A., Meini, A., Ferrari, S., et al., 2005. Mutational analysis of
929   human BAFF receptor TNFRSF13C (BAFF-R) in patients with common variable immunodeficiency.
930   Journal of Clinical Immunology 25(5): 496–502, Doi: 10.1007/s10875-005-5637-2.
931   [61] Tuijnenburg, P., Lango Allen, H., Burns, S.O., Greene, D., Jansen, M.H., Staples, E., et al., 2018. Loss-
932   of-function nuclear factor κB subunit 1 (NFKB1) variants are the most common monogenic cause of
933   common variable immunodeficiency in Europeans. The Journal of Allergy and Clinical Immunology
934   142(4): 1285–96, Doi: 10.1016/j.jaci.2018.01.039.
935   [62] Chen, K., Coonrod, E.M., Kumánovics, A., Franks, Z.F., Durtschi, J.D., Margraf, R.L., et al., 2013.
936   Germline mutations in NFKB2 implicate the noncanonical NF-κB pathway in the pathogenesis of
937   common variable immunodeficiency. American Journal of Human Genetics 93(5): 812–24, Doi:
938   10.1016/j.ajhg.2013.09.009.
939   [63] Pieniawska-Śmiech, K., Bar, K., Babicki, M., Śmiech, K., Lewandowicz-Uszyńska, A., 2020.
940   Assessment of weight and height of patients with primary immunodeficiency disorders and group
941   of children with recurrent respiratory tract infections. BMC Immunology 21: 42, Doi:
942   10.1186/s12865-020-00372-x.

[64] Caetano, M.C., Silva, R., Ferreira, C.A., Sarni, R.O.S., Carvalho, B.T.C., 2011. Lipid Profile In Patients
With Primary Immunodeficiency. Journal of Allergy and Clinical Immunology 127(2, Supplement):
AB65, Doi: 10.1016/j.jaci.2010.12.269.
[65] Dobner, J., Kaser, S., 2018. Body mass index and the risk of infection - from underweight to
obesity. Clinical Microbiology and Infection 24(1): 24–8, Doi: 10.1016/j.cmi.2017.02.013.
[66] Ruffner, M.A., Sullivan, K.E., 2016. Body Weight and Infectious Outcomes in Patients with Primary
Immunodeficiency Diseases: Outcomes from within the US Immunodeficiency Network (USIDNET).
Journal of Allergy and Clinical Immunology 137(2): AB179, Doi: 10.1016/j.jaci.2015.12.718.
[67] Parikh, S., Goldstein, A., Karaa, A., Koenig, M.K., Anselm, I., Brunel-Guitton, C., et al., 2017. Patient
care standards for primary mitochondrial disease: a consensus statement from the Mitochondrial
Medicine Society. Genetics in Medicine: Official Journal of the American College of Medical
Genetics 19(12), Doi: 10.1038/gim.2017.107.
[68] Bapat, S.P., Whitty, C., Mowery, C.T., Liang, Y., Yoo, A., Jiang, Z., et al., 2022. Obesity alters
pathology and treatment response in inflammatory disease. Nature 604(7905): 337–42, Doi:
10.1038/s41586-022-04536-0.
[69] Parvaneh, N., Quartier, P., Rostami, P., Casanova, J.-L., de Lonlay, P., 2014. Inborn errors of
metabolism underlying primary immunodeficiencies. Journal of Clinical Immunology 34(7): 753–71,
Doi: 10.1007/s10875-014-0076-6.
[70] Bugge, A., Dib, L., Collins, S., 2014. Measuring respiratory activity of adipocytes and adipose tissues
in real time. Methods in Enzymology 538: 233–47, Doi: 10.1016/B978-0-12-800280-3.00013-X.
[71] Agilent Technologies., Seahorse XFp Cell Mito Stress Test Kit User Guide:
https://www.agilent.com/cs/library/usermanuals/public/XFp_Cell_Mito_Stress_Test_Kit_User_Gui
de.pdf, 18.
[72] Agilent Technologies., Seahorse XF Cell Energy Phenotype Test Kit User Guide:
https://www.agilent.com/cs/library/usermanuals/public/XF_Cell_Energy_Phenotype_Test_Kit_Us
er_Guide.pdf, 20.
[73] Agilent Technologies., Seahorse XF Glycolysis Stress Test Kit User Guide:
https://www.agilent.com/cs/library/usermanuals/public/XF_Glycolysis_Stress_Test_Kit_User_Gui
de.pdf, 22.