Genome-wide local ancestry and direct evidence for mitonuclear co-adaptation in African hybrid cattle populations (*Bos taurus/indicus*)

James A. Ward[1] | Gillian P. McHugo[1] | Michael J. Dover[1] | Thomas J. Hall[1] | Said Ismael Ng'ang'a[2,3] | Tad S. Sonstegard[4] | Daniel G. Bradley[5] | Laurent A. F. Frantz[2,3] | Michael Salter-Townshend[6] | David E. MacHugh[1,7]

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[2] Palaeogenomics Group, Department of Veterinary Sciences, Ludwig Maximilian University, Munich D-80539, Germany.

[3] School of Biological and Chemical Sciences, Queen Mary University of London, London E1 4NS, United Kingdom.

[4] Acceligen, 3388 Mike Collins Drive, Eagan, MN, 55121, USA.

[5] Smurfit Institute of Genetics, Trinity College Dublin, Dublin, D02 VF25, Ireland.

[6] UCD School of Mathematics and Statistics, University College Dublin, Dublin, D04 V1W8, Ireland.

[7] UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Dublin, D04 V1W8, Ireland.

Correspondence

David E. MacHugh, Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

Email: david.machugh@ucd.ie

Abstract

Domestic cattle have a key economic role in African societies, providing an important source of mobile wealth through supply of meat, milk, cowhide, fuel, transport, and traction. The phenotypic diversity of African cattle reflects adaptation to a wide range of agroecological conditions and complex patterns of admixture between the humpless *Bos taurus* (taurine) and humped *Bos indicus* (zebu) subspecies, which share a common ancestor 150-500 thousand years ago. Human migration and trade from Asia have left a peak of zebu nuclear ancestry in East Africa and most cattle populations across the continent have a hybrid genetic composition. Notwithstanding this, all African cattle possess taurine mitochondrial haplotypes, even populations with significant zebu nuclear ancestry. In this regard, the efficient functioning of the mitochondrion relies on a network of biochemical interactions between the products of 37 mitochondrial genes and more than one thousand nuclear genes; therefore, admixed African cattle represent ideal populations for evaluating mitonuclear interactions and mismatch between the nuclear and mitochondrial genomes. Using high-density SNP array data from 18 different cattle populations, including ten African admixed breeds, we find strong evidence for mitonuclear coevolution in hybrid African cattle with significant retention of *Bos taurus* alleles at mitochondrially-targeted nuclear genes, particularly those genes with products that directly interact with mtDNA-encoded protein subunits in OXPHOS and ribosomal complexes, or that have functions in mtDNA replication. We also show that subspecific local ancestry varies substantially across the genomes of admixed populations, with a marked signal of taurine ancestry at the major histocompatibility (MHC) gene cluster, which likely reflects adaptation to infectious disease challenges facing African livestock. Our results demonstrate that African admixed cattle represent an excellent comparative model for studying the phenotypic

47   consequences of mitonuclear mismatch and genomic introgression in humans and other large

48   mammals.

1 | INTRODUCTION

The mitochondrion is an endosymbiotic double membrane-bound eukaryotic organelle that evolved from an α-proteobacterial ancestor as consequence of the transition to an oxygen-rich atmosphere that began more than two billion years ago (Dyall *et al.* 2004; Lyons *et al.* 2014; Archibald 2015; Martin *et al.* 2015). Mitochondria harness energy from compounds such as sugars to produce, via oxidative phosphorylation (OXPHOS), adenosine triphosphate (ATP), the principal molecule used to store and transfer energy in the cell. Mitochondria are directly involved in a myriad of cellular and physiological processes, including, for example, calcium homeostasis, cell growth, cell differentiation, cell death, cellular waste management, neuronal function, and aging (McBride *et al.* 2006; Kasahara & Scorrano 2014; Spinelli & Haigis 2018). In addition, the key role of mitochondria in regulation of innate and adaptive immune responses has become more widely appreciated (Weinberg *et al.* 2015; Mills *et al.* 2017; Tiku *et al.* 2020).

Mitochondria have retained their own genome and in vertebrates this is a circular DNA molecule (mtDNA) of approximately 16 kb that encodes 37 genes and is exclusively transmitted by females to their offspring (Boore 1999; Zardoya 2020). Despite this, mitochondria also rely on more than one thousand co-adapted nuclear genes that encode proteins and protein subunits essential to mitochondrial function (Blier *et al.* 2001; Rand *et al.* 2004; Woodson & Chory 2008; Sloan *et al.* 2018). This intracellular mutualism is exemplified by the vertebrate OXPHOS system, which consists of five protein complexes, the four larger of which are chimeric—assembled using subunits encoded by the both the nuclear and mitochondrial genomes (Rand *et al.* 2004; Allen 2015; Isaac *et al.* 2018). The fundamental nature of these interactions and the importance of mitonuclear coevolution have been proposed to profoundly shape the evolution and ecology of eukaryotes (Burton *et al.* 2013; Wolff *et al.* 2014; Hill 2015; Sunnucks *et al.* 2017). Co-adapted sets of mitochondrial and nuclear genes have also been

proposed to underpin eukaryotic speciation and demarcation of species boundaries—the

mitonuclear compatibility species concept (MCSC)—thereby providing an explanatory

framework for using the mitochondrial cytochrome c oxidase I gene (*MT-CO1*) as an objective

and effective molecular barcode for taxonomic species delineation (Hill 2016; Hill 2017).

Functional mismatch between the mtDNA and nuclear genomes can be observed, for

example, in natural and human-generated hybrid populations, or in cybrids—cytoplasmic

hybrids—generated as cell lines, through somatic cell nuclear transfer (SCNT), or via hybrid

backcrosses (Burton *et al.* 2013; Chou & Leu 2015; Hill 2015; Sloan *et al.* 2017; Sunnucks *et*

al. 2017; Lechuga-Vieco *et al.* 2021). In these systems, mitonuclear incompatibilities between

distinct inter- and intraspecific evolutionary lineages can give rise to deleterious biochemical

effects associated with reduced efficacy of OXPHOS protein complexes (Blier *et al.* 2001;

Ellison & Burton 2006; Ellison *et al.* 2008; Ballard & Melvin 2010), which lead to lower ATP

production (McKenzie *et al.* 2003; McKenzie *et al.* 2004; Ellison & Burton 2008; Ellison *et al.*

2008) and increased levels of oxidative damage (Barreto & Burton 2013; Latorre-Pellicer *et al.*

2016; Du *et al.* 2017; Pichaud *et al.* 2019). At the organismal level, the physiological

consequences may be reduced fitness across a range of biological traits in vertebrates and

invertebrates, including those related to fecundity and reproductive success (Barreto & Burton

2013; Tourmente *et al.* 2017; Rank *et al.* 2020), developmental mortality and longevity (Ellison

et al. 2008; Latorre-Pellicer *et al.* 2016; Pichaud *et al.* 2019), metabolism and nutrition (Derr *et*

al. 2012; Latorre-Pellicer *et al.* 2016; Wang *et al.* 2017; Kong *et al.* 2020), endurance exercise

(Sujkowski *et al.* 2019; Spierer *et al.* 2021), and behaviour and neurobiology (Yu *et al.* 2009;

Gonzalez 2021). In F$_3$ hybrids of the marine copepod *Tigriopus californicus*, it was also

demonstrated that wild-type fitness could be restored through maternal backcrosses that

reunited intact nuclear genomes with co-evolved mitochondrial genomes (Ellison & Burton

2008)

Evidence for mitonuclear mismatches that affect biological fitness in natural vertebrate

populations has accumulated during the last decade and includes observations in the Iberian

hare (*Lepus granatensis*) subject to extensive mtDNA introgression from the mountain hare, *L.*

timidus, a northern boreal species (Seixas *et al.* 2018); the Italian sparrow (*Passer italiae*), a

hybrid between *P. domesticus* and *P. hispaniolensis* (Trier *et al.* 2014); hybrids of two diverged

Australian eastern yellow robin (*Eopsaltria australis*) populations (Morales *et al.* 2018);

Mediterranean chameleons (*Chamaeleo chamaeleon*) inhabiting an ancient Levantine marine

barrier (Bar-Yaacov *et al.* 2015); hybrids of two distinct lineages of long-toed salamander

(*Ambystoma macrodactylum*) in western Canada (Lee-Yaw *et al.* 2014); a hybrid population of

Atlantic killifish (*Fundulus heteroclitus*) with two divergent mtDNA haplotypes (Baris *et al.*

2017); and a hybrid fork-tongued goby (*Chaenogobius annularis*) population on the east coast

of Japan (Hirase *et al.* 2021).

The evidence for selection acting to conserve co-adapted mitonuclear interactions in

human populations is equivocal. Analysis of nuclear and mtDNA SNP data from the Human

Genome Diversity Project (HGDP) detected low levels of mitonuclear linkage disequilibria

(LD), even for nuclear genes encoding protein subunits that translocate to the mitochondrion

and form OXPHOS complexes with mtDNA-encoded proteins or that directly interact with

mitochondrial DNA or RNAs (Sloan *et al.* 2015). Contrary to this, however, a more recent study

of six distinct American admixed populations, demonstrated that mtDNA copy number is

inversely correlated with the level of discordance between nuclear and mtDNA ancestry (Zaidi

& Makova 2019). This observation was proposed to be a consequence of suboptimal regulation

of mtDNA replication in individuals with mismatched nuclear and mitochondrial genes. In addition, there was evidence, at least in some populations, for a signature of mitonuclear compatible ancestry enrichment at nuclear genes functionally associated with the mitochondrion, particularly those encoding proteins that translocate to the organelle and interact with mtDNA-encoded proteins in multi-subunit OXPHOS and ribosomal complexes. A third study examined potential mitonuclear incompatibilities arising due to admixture with Neanderthals (*Homo neanderthalensis*) and Denisovans (*Homo denisova*), which has left signatures in modern human genomes of archaic hominin nuclear ancestry without mtDNA introgression (Sharbrough *et al.* 2017). This analysis showed that Neandertal haplotypes for nuclear gene loci associated with the mitochondrion have, as hypothesised, introgressed less readily into both Asian and European human populations. This pattern, however, was not observed for nuclear genomic introgression from Denisovans, likely a consequence of reduced power to detect Denisovan haplotypes in populations exhibiting admixture with both archaic hominins.

After humans, from a population genetics perspective, the most well-studied large mammals are domestic cattle, which are derived from two morphologically distinct wild subspecies, *Bos primigenius taurus* and *Bos primigenius indicus*, informally shortened to *B. taurus* and *B. indicus* and colloquially referred to as "taurine" and "zebu" (or "indicine") cattle, respectively (Zhang *et al.* 2020). Comprehensive genetic and archaeological evidence has shown that taurine and zebu cattle were most likely domesticated separately: humpless taurine cattle from the *B. p. primigenius* subspecies in the Fertile Crescent region of the Middle East approximately 10 thousand years ago (kya), and humped zebu from the *B. p. namadicus* subspecies two millennia later by the early Neolithic cultures located in present-day Baluchistan, Pakistan (Bradley *et al.* 1998; Fuller 2006; Zeder 2011; Utsunomiya *et al.* 2019).

Recent estimates, based on genome-wide sequence data, place the evolutionary divergence time

between the two cattle subspecies at 150 to 500 kya (Chen *et al.* 2018; Wang *et al.* 2018; Wu

et al. 2018), although some analyses of complete mitogenomes have suggested a deeper split

(Hiendleder *et al.* 2008; Bibi 2013; Pramod *et al.* 2019).

The *B. taurus* and *B. indicus* lineages are interfertile and since domestication, and likely

before, there has been significant gene flow and admixture between humpless and humped

cattle across time and traversing multiple locations and environments in Eurasia and beyond

(Decker *et al.* 2014; Ginja *et al.* 2019; Verdugo *et al.* 2019; Gebrehiwot *et al.* 2020; Senczuk

et al. 2021). The largest and most well studied hybrid zone encompasses most of the African

continent and represents a complex mosaic of taurine and zebu ancestry that has been influenced

by a wide range of factors, including climate, physical geography, human migrations, and

infectious disease challenges (MacHugh *et al.* 1997; Hanotte *et al.* 2002; Kim *et al.* 2017; Kim

et al. 2020). However, even in the face of substantial zebu admixture, it has long been

understood that African cattle populations exclusively retain taurine mtDNA haplotypes, the

majority of which (within the T1 haplogroup) differ from T, T2, T3, and T4 haplotypes

observed in other taurine cattle populations (Loftus *et al.* 1994a; Loftus *et al.* 1994b; Bradley

et al. 1996; Troy *et al.* 2001). Therefore, as summarised in Figure 1, hybrid African cattle

represent excellent models for testing hypotheses concerning mitonuclear coevolution in large

mammals. In the present study, we leverage high-density SNP data and established

computational methods to characterise genome-wide local ancestry and systematically evaluate

mitonuclear interactions, coadaptation, and functional mismatch in ten genetically distinct

admixed African cattle populations.

2 | MATERIALS AND METHODS

Animal sampling and genotyping

High-density genome-wide SNP array data sets (Illumina® BovineHD 777K BeadChip) corresponding to a total of 605 animals were obtained from published studies (Bahbahani *et al.* 2015; Verdugo *et al.* 2019) and the Acceligen cattle genotyping database. For the present study, new BovineHD 777K BeadChip SNP data were generated for 73 additional animals (50 Borgou and 23 N'Dama) that were previously published by our group as part of microsatellite-based surveys of cattle genetic diversity (MacHugh *et al.* 1997; Freeman *et al.* 2004). These new SNP genotype data sets were generated by Weatherbys Scientific (Co. Kildare, Ireland) using standard procedures for Illumina SNP array genotyping. In total, 18 different breeds/populations were represented (Table 1), including two West African taurine breeds (Muturu and N'Dama); two European taurine breeds (Holstein-Friesian and Jersey); ten West and East African admixed zebu-taurine (Adamawa Gudali, Ankole, Borgou, Bunaji, East African Shorthorn Zebu, Karamojong, Keteku, Nganda, Red Bororo, and Sokoto Gudali); and four zebu breeds of South Asian origin (Gir, Hariana, Nelore, and Sahiwal). Table 1 also shows the three- or four-letter codes used to designate each breed from this point onwards in this study.

SNP genotype data pre-processing, quality control, and filtering

BovineHD 777K SNP locations were remapped to the current bovine genome assembly ARS-UCD1.2 (Rosen *et al.* 2020) and SNP genotype data were merged using PLINK v1.9 (Chang *et al.* 2015). Quality control (QC) of the combined SNP genotype data set was also performed using PLINK v1.9 and autosomal SNPs with a call rate < 95% and a minor allele frequency (MAF) of < 0.05 were filtered from the data.

Principal component and population structure analyses

Principal component analysis (PCA) of individual animal SNP genotype data for the African taurine (MUTU and NDAG), the East African admixed (ADAG, ANKO, BORG, BUNA, EASZ, KARA, KETE, NGAN, REDB, and SOKG) and two Asian indicine (GIR and NELO) populations was performed using PLINK v1.9 and the results were plotted using GGPLOT2 v3.3.3 (Wickham 2016) in the R v3.6.2 environment for statistical computation and graphics (R Core Team 2019). The genetic structure of each population was also estimated using FASTSTRUCTURE v1.0 with $K = 3$ modelled ancestries to determine mean African and European *B. taurus* and Asian *B. indicus* contributions (Raj *et al.* 2014).

Mitochondrial DNA haplogroup determination

The BovineHD 777K BeadChip includes 346 SNPs located in the mitochondrial genome, which can be used to construct haplotypes and catalogue and distinguish the mitochondrial haplogroups characteristic of *B. taurus* and *B. indicus* cattle lineages. For this analysis, the European JRSY and HOLS taurine breeds and the Indo-Pakistan HARI and SAHI Asian indicine breeds were also included to ensure good representation of the *B. taurus* and *B. indicus* mtDNA haplogroups—the 'T' and 'I' groups, respectively (Troy *et al.* 2001; Chen *et al.* 2010). The mtDNA SNPs were filtered using PLINK v1.9 (Chang *et al.* 2015) such that SNPs with a MAF of < 0.10, and a call rate of < 95% were removed. Individual animals with a genotype missingness of > 95% were also removed. Following this, the most ancestry informative mtDNA SNPs were identified using INFOCALC (Rosenberg *et al.* 2003; Rosenberg 2005), which provides I_n, a general measure of the informativeness of a SNP for ancestry assignment. The 50 top ranked SNPs, based on I_n, were then used to generate mtDNA haplotypes with the FASTPHASE v1.4 program (Scheet & Stephens 2006). Haplotype networks were constructed using the POPART v1.7 package (Leigh & Bryant 2015).

Local ancestry analysis of admixed populations

Local ancestry across the bovine genome for each African admixed breed (ADAG, ANKO, BORG, BUNA, EASZ, KARA, KETE, NGAN, REDB, and SOKG) was inferred using MOSAIC v1.3.7 (Salter-Townshend & Myers 2019). The MOSAIC algorithm, unlike other methods, does not require donor reference populations to be direct surrogates for the mixing ancestral populations; it fits a two-layer Hidden Markov Model (HMM) that determines how closely related each segment of chromosome in each admixed individual genome is to the segments of chromosomes in individual genomes from potential donor populations and infers a stochastic relationship between donor reference panels and mixing populations. While determining local ancestry along each chromosome, MOSAIC also infers the number of generations since the admixture process started for a particular population. The potential donor populations used for the MOSAIC local ancestry analysis were the two West African *B. taurus* breeds (MUTU and NDAG) and two of the Asian *B. indicus* breeds (GIR and NELO). The MOSAIC algorithm requires phased haplotypes and a recombination rate map; therefore, SHAPEIT v2 (r900) (Delaneau *et al.* 2012) was used to generate phased haplotypes and a published cattle recombination map was employed (Ma *et al.* 2015). Functional enrichment analysis of genes proximal to African taurine and Asian zebu local ancestry peaks was performed using Ingenuity® Pathway Analysis—IPA® (version 1.1, summer 2021 release; Qiagen, Redwood City, CA, USA) (Kramer *et al.* 2014).

Detection of subchromosomal deviations in taurine and indicine genomic ancestry

To determine if there was significant retention of *B. taurus* nuclear genes that encode mitochondrially targeted proteins (*N-mito* genes) in African admixed cattle, we used an approach modified from previously published surveys of mitonuclear incompatibilities in modern admixed human populations (Sloan *et al.* 2015; Zaidi & Makova 2019) and because of

ancient gene flow from archaic hominins (*H. neanderthalensis* and *H. denisova*) (Sharbrough *et al.* 2017). Firstly, the MITOCARTA 2.0 database resource (Calvo *et al.* 2016) was used to obtain an inventory of genes that produce the nuclear-encoded component of the mammalian mitochondrial proteome, i.e., proteins with experimental evidence for localisation in the mitochondrion. Following this, the ENSEMBL BIOMART tool (Yates *et al.* 2020) was used to generate a list of 1158 bovine N-mito genes, which was classified into two functional subsets as defined by Sloan *et al.* (2015) and also used by Sharbrough *et al.* (2017) and Zaidi and Makova (2019). These subsets were denoted as 1) high-confidence "*high-mito*" genes (HMG) encoding proteins that directly interact with mtDNA-encoded protein subunits in OXPHOS and ribosomal complexes, or that have functions in mtDNA replication (138 genes); and 2) lower confidence "*low-mito*" genes (LMG), which encode proteins that localise to the mitochondrion, but are not classified as part of the high-mito subset (701 genes). Finally, a third group of "*non-mito*" genes (NMG) was generated, which includes the bulk of the mammalian proteome that does not localise to the mitochondrion (17,834 genes). Table S1 provides further detail for the functional gene subsets used to detect evidence for mitonuclear incompatibilities in African admixed cattle populations.

The local ancestry estimates generated using MOSAIC for each SNP across the genome were catalogued and the BEDTOOLS v2.18 software suite (Quinlan & Hall 2010) was then used to intersect these SNPs with windows spanning 2.5 Mb upstream and downstream of genes within each of the three functional gene subsets. Following this, and as described by Zaidi and Makova (2019), for each of the three subsets an unweighted block bootstrap approach was used to generate distributions of local ancestry deviation towards more or less *B. taurus* ancestry. The first step in this methodology is subtraction of the mean ancestry fraction across the local ancestry estimate for each SNP (the *expectation*), which produces the deviation in local ancestry

at each SNP locus. For each functional gene subset, the number of windows sampled with

replacement was the same as the number of HMG subset genes ($n = 138$). In each case, the

mean ancestry deviation was estimated and then averaged across all windows. Bootstrap

resampling (1000 replicates) was used to generate a distribution of mean deviations in local

ancestry for each of the three functional gene subsets. The overall statistical significance of

deviations from expected local ancestry (using the background NMG subset) was then assessed

using Welch's t-test in R v3.6.2 (R Core Team 2019).

Transcriptomics and Cattle Gene Atlas analysis

Following the methodology described by Sharbrough *et al.* (2017), under the assumption

that highly expressed genes are more functionally constrained, we evaluated if the local ancestry

estimates for HMG, LMG or NMG subsets were correlated with gene expression across a wide

range of bovine tissues. RNA-seq transcriptomics data in the form of fragments per kilobase

per million mapped reads (FPKM) was obtained from the Cattle Gene Atlas database

(http://cattlegeneatlas.roslin.ed.ac.uk; Fang *et al.* 2020), which consists of 723 RNA-seq data

sets across 84 bovine tissues. For each tissue, the expression values were averaged, resulting in

84 mean FPKM values for all known bovine genes. Following this, a K_{ij} matrix was generated

such that each cell contained mean FPKM values organised into columns (i) representing each

tissue and rows (j) corresponding to the 130 HMG, 638 LMG, and 14,758 NMG genes that

could be mapped to the Cattle Gene Atlas. An expression threshold was created such that each

gene was classified as detectably expressed for a particular tissue if its mean FPKM value for

that tissue was >7.8 (corresponds to 50% of the global mean FPKM value of 15.6) across the

complete data set (all genes and all tissues). A tissue expression metric (TEM) for

systematically examining correlated expression for the HMG, LMG and NMG gene subsets

was then created as follows: the mean expression of a gene across all 84 tissues was multiplied

by the number of cases where mean expression of a gene was greater than the global FKKM

mean (15.6) in any tissue. For example, if a gene exhibited a mean expression value of 10.0

across all 84 samples and its expression was greater than 15.6 in five of these tissues, then the

TEM for that gene would be 50.0. The TEM values for each gene in the HMG, LMG, and NMG

subsets were then plotted against the mean African taurine local ancestry scores for each gene

locus across the 10 African admixed breeds and Pearson correlation coefficients were estimated

for the LMG, HMG, and NMG subsets. The statistical analyses and data graphing for the TEM

and African taurine local ancestry analysis were performed using R v3.6.2 (R Core Team 2019)

and GGPLOT2 v3.3.3 (Wickham 2016).

3 | RESULTS

Genomic structure of African admixed cattle

Following filtering and QC there were 562,635 SNPs and 605 individual animals retained

for subsequent analyses. Figure 2 shows a PCA plot generated from SNP genotype data for

individual African taurine and admixed animals and with the GIR and NELO breeds

representing Asian *B. indicus* cattle and the HOLS and JERS representing the European *B.*

taurus group. The first principal component (PC1), which accounts for 58.4% of the variation,

represents the primary split between the *B. taurus* and *B. indicus* lineages and PC2 (17.9% of

the variation) differentiates the African and European taurine populations. The position of

individual admixed animals is determined by the relative proportions of African taurine and

zebu ancestry, with additional European gene flow evident for some East African cattle—

particularly within the EASZ and NGAN breeds. The results of the genetic structure analysis

with FASTSTRUCTURE and an inferred number of clusters of $K = 3$ are shown in Figure 3 and

illustrates the varying patterns of taurine and zebu ancestry across the different East and West

African admixed cattle populations. For the most part, the levels of admixture are relatively

uniform for animals within each breed; however, there are some cattle that deviate noticeably

from the pattern, particularly in the EASZ and NGAN where recent European taurine gene flow

is most evident. Table 2 provides additional detail on the taurine/zebu admixture results

generated using the FASTSTRUCTURE program. As generally observed in previous studies of

B. taurus/*B. indicus* admixture (MacHugh *et al.* 1997; Hanotte *et al.* 2002; Kim *et al.* 2017;

Gebrehiwot *et al.* 2020; Kim *et al.* 2020), the four East African admixed populations exhibit a

greater level of average zebu ancestry (mean = 0.653) than the six West African admixed

populations (mean = 0.612).

Bos taurus and *Bos indicus* mitochondrial DNA haplotype network analysis: admixed African cattle possess taurine mitochondrial genomes

After filtering of the 346 mtDNA SNPs on the BovineHD 777K BeadChip and

identification of the most ancestry-informative SNPs, a phylogenetic haplotype network was

generated using 39 mtDNA SNPs and a total of 491 cattle (47 African taurine, 82 European

taurine, 156 East African admixed, 136 West African admixed, and 70 Asian zebu). Figure 4

shows this network and demonstrates that all 339 African taurine and admixed cattle surveyed

here possess the taurine mitochondrial genome. This observation, which uses, to the best of our

knowledge, the largest sample of African mtDNA genotypes, agrees with previous studies of

mtDNA diversity in African cattle (Loftus *et al.* 1994a; Loftus *et al.* 1994b; Bradley *et al.* 1996;

Troy *et al.* 2001; Dadi *et al.* 2009; Bonfiglio *et al.* 2012). It has been hypothesised that male-

mediated gene flow accounts for this pattern of mitonuclear discordance, which resulted due to

the geographical barriers between Asia and Africa and historical preferences for trade and

transport of *B. indicus* bulls to disseminate desirable traits (Bradley *et al.* 1994; Loftus *et al.*

1994a). Importantly, from the perspective of the present study, these results unambiguously

demonstrate that East and West admixed African cattle breeds retain taurine mitochondrial

genomes, making them ideal populations to investigate mitonuclear interactions and co-

adaptation.

We also observed taurine mtDNA haplotypes in animals from the morphologically Asian

zebu GIR and NELO breeds; this mitochondrial discontinuity reflects their early origins as

hybrid taurine-indicine founder populations that were backcrossed to *B. indicus* bulls to recover

South Asian zebu traits advantageous for the tropical production environments of Brazil (Dani

et al. 2008). In this regard, present-day GIR and NELO cattle exhibit very low levels of

autosomal taurine admixture, less than 1% when estimated using genome-wide high-density

SNP data (O'Brien *et al.* 2015).

341 Genome-wide local ancestry analysis and deviations in African taurine local ancestry
342 for functional mitochondrial gene subsets

Three coancestry curves (two-way African taurine and Asian zebu admixture) generated

using MOSAIC are shown in Figure 5 for each of the East and West African cattle populations.

The ancestry labels on top of each subplot correspond to the closest African taurine and Asian

zebu donor breed as measured by the Weir and Cockerham $\hat{F}_{st}$ estimator of population

differentiation (Weir & Cockerham 1984). In each case, this was the NDAG *B. taurus* and GIR

B. indicus breeds and the full set of $\hat{F}_{st}$ values for each of the admixed African breeds is

provided in Table S2. In addition, the coancestry curve analyses provides estimates of the time

in generations since admixture commenced and these were converted to years before present

(BP) using the NDAG:GIR coancestry curve values (middle subplots) and a generation interval

range of between 4 and 7 years for managed domestic cattle populations (Mészáros *et al.* 2015;

Abin *et al.* 2016). The estimated historical calendar range (common era – CE) for the start of

the admixture process in each of the African admixed breeds is shown in Table 2. A recent

study using whole-genome sequence (WGS) data, albeit with lower samples sizes, also

estimated the time in generations since the beginning of admixture for one of the East African

breeds used for the present study (ANKO). Kim *et al.* (2020) used WGS data from 10 ANKO

cattle to obtain linkage disequilibrium (LD) and haplotype-sharing coancestry based estimates

of 147.3 and 113.6 generations, respectively—more than twice the result obtained here with

SNP array data (54.1 generations; 1641 – 1804 CE) and pushing back the historical calendar

range for the start of admixture in the ANKO population to between 989 and 1566 CE.

For each of the East and West African admixed populations, Asian zebu, and African

taurine SNP-by-SNP local ancestry values were estimated for each of the 29 bovine autosomes

using MOSAIC. As an example of this process and the resulting output, Figure 6 shows Asian

zebu and African taurine local ancestry values plotted for all SNPs across bovine chromosome

23 (BTA23). Individual SNP local ancestry values were then averaged on a gene-by-gene basis

to determine African taurine local ancestry values for individual genes. These can be used to

systematically evaluate deviations in local ancestry for functional mitochondrial gene subsets

(HMG and LMG) that could indicate mitonuclear incompatibilities and coadaptation in

admixed African cattle. African taurine local ancestry deviations for the three different

functional gene subsets (HMG, LMG and NMG) are displayed as violin plots in Figure 7a. A

positive deviation for a breed/subset indicates that a particular functional gene subset exhibits

retention of taurine gene haplotypes. The HMG, LMG, and NMG African taurine local ancestry

deviation distributions for each breed were evaluated for normality using the Shapiro-Wilk test.

Following this, comparative statistical analysis of the local ancestry deviation distributions was

performed, and the results are shown in Table 3. This analysis demonstrated that all 10 African

admixed cattle breeds exhibited statistically significant and consistently increased African

taurine local ancestry deviation distributions for the HMG subset compared to the background

NMG subset. In addition, three out of four of the East African (ANKO, KARA, and NGAN)

and two out of six of the West African (ADAG and KETE) admixed breeds showed a statistically significant increase in African taurine local ancestry deviation distributions for the LMG subset compared to the background NMG subset. Finally, except for the ADAG breed, all populations showed significantly increased African taurine local ancestry in the HMG subset compared to the LMG subset.

A direct statistical comparison of the HMG African taurine local ancestry distributions for the combined East and West African admixed cattle groups (Figure 7b) demonstrated that the East African admixed group exhibit a significantly higher local ancestry deviation for the HMG subset of nuclear genes that encode mitochondrially targeted proteins. We also tested the hypothesis that mean African taurine ancestry deviation is influenced by the number of generations since the start of admixture, assuming a single pulse of admixture as modelled by the MOSAIC software (Salter-Townshend & Myers 2019). The results of this analysis are shown in Figure 7c where a statistically significant negative correlation ($R = -0.678$; $P = 0.031$) was observed between the mean African taurine local ancestry deviation values for the HMG subset and the generations since the start of admixture. This result is somewhat counterintuitive because it might be expected that African taurine local ancestry deviations for the HMG subset would increase with time as selection acts to purge mitonuclear incompatibilities and maintain co-adaption of the mitochondrial and nuclear genomes. However, several cofounding factors may contribute to this result, including varying levels of genome-wide taurine-zebu ancestry and the presence of European taurine ancestry in certain populations (see Figure 3 and Table 3). In addition, the temporal pattern and mode of admixture may not correspond to a simple scenario of a single admixture event or pulse, as modelled by MOSAIC. However, the most persuasive biological explanation is provided by genetic recombination acting across multiple generations, which over time would erode and break down contiguous taurine haplotypes at the

HMG subset gene loci. Consequently, as HMG gene loci in admixed populations recombine, the mean African taurine local ancestry scores averaged across multiple SNPs would decrease and this would lead to dissipation of apparent positive African taurine local ancestry deviation for the HMG subset compared to the background NMG subset. The gene-by-gene local ancestry approach that we employ here to detect mitonuclear incompatibilities may only function accurately for relatively recently admixed populations, such as the human populations with African, European and Native America ancestry examined by Zaidi and Makova (2019). In other words, the underlying signal of mitonuclear compatibility and co-adaption is likely stronger in long established admixed African cattle breeds (*i.e.*, ADAG, BUNA, and SOKG) compared to more recently admixed populations; however, it is masked by recombination and may require higher-resolution local ancestry methods to detect it; for example, using larger sample sizes with WGS data, and an exon-by-exon, rather than a gene-by-gene approach.

Taken together and bearing in mind that ten different independent admixed cattle populations were examined, the results presented here provide compelling evidence supporting the hypothesis that bovine nuclear genes encoding proteins directly (the HMG subset), or indirectly (the LMG subset) involved with the mitochondrion have been influenced by selective processes favouring mitonuclear compatibility and co-adaptation in admixed African cattle populations. This is particularly evident for the HMG subset where all ten breeds exhibited strong signals of African taurine local ancestry enrichment at nuclear genes that encode mitochondrially targeted proteins.

Genome- and chromosome-wide patterns of ancestry across all African admixed cattle populations

To illustrate genome-wide patterns of taurine and zebu admixture across all ten African admixed cattle populations, Figure 8a shows mean African taurine local ancestry for each of

the bovine autosomes (BTA1–BTA29) across all ten African admixed breeds. The three highest

taurine ancestry peaks were observed on BTA7, BTA12, and BTA23 and the three lowest

taurine ancestry troughs (zebu peaks) were observed on BTA3, BTA5, and BTA6. Figure 8b

shows mean African taurine local ancestry plotted across three of these chromosomes (BTA5,

BTA12, and BTA23). The highest genome-wide signal of mean African taurine ancestry across

the ten admixed breeds was observed as a broad plateau encompassing the bovine major

histocompatibility complex (MHC) gene cluster on BTA23 (indicated as a purple rectangle on

Figure 8b), likely reflecting the key role of the MHC gene cluster in innate and adaptive

immunity and responses to pathogens and parasites (Behl *et al.* 2012; Ellis & Hammond 2014).

To further examine patterns of introgression across the genome, the top 1% of the mean

African taurine and Asian zebu local ancestry peaks across the 10 admixed breeds were

identified separately. Following this, genes 2.5 Mb up- and downstream of each peak were

catalogued and used for functional enrichment analysis with the IPA software tool, using the

default settings. Table S3 shows these input gene sets, which contained 858 and 317 genes for

the top 1% of African taurine and Asian zebu local ancestry peaks, respectively. A false

discovery rate (FDR) method Benjamini and Hochberg (1995) was used to set a $P_{adj.}$ threshold

of 0.05 for statistical significance of enrichment for the canonical pathways and physiological

system development and function categories in IPA. The top five enriched canonical pathways

for genes with high African taurine ancestry were *Differential Regulation of Cytokine*

Production in Macrophages and T Helper Cells by IL-17A and IL-17F, *Differential Regulation*

of Cytokine Production in Intestinal Epithelial Cells by IL-17A and IL-17F, *Glutathione-*

mediated Detoxification, *GABA Receptor Signalling*, and *Gustation Pathway*. The top five

enriched physiological system development and function categories genes with high African

taurine ancestry were *Organismal Survival*, *Immune Cell Trafficking*, *Cardiovascular System*

Development and Function, *Haematological System Development and Function*, and *Tissue Morphology*. The top five enriched canonical pathways for genes with high Asian zebu ancestry were *NRF2-mediated Oxidative Stress Response*, *Mitotic Roles of Polo-Like Kinase*, *-tocopherol Degradation*, *Methylglyoxal Degradation III*, and *Role of Osteoblasts, Osteoclasts and Chondrocytes in Rheumatoid Arthritis*. The top five enriched physiological system development and function categories for genes with high Asian zebu ancestry were *Connective Tissue Development and Function*, *Skeletal and Muscular System Development and Function*, *Nervous System Development and Function*, *Organ Morphology*, and *Organismal Development*.

Gene expression analysis of functional HMG, LMG, and NMG subsets

Figure S1 show tissue expression metric (TEM) values obtained from the Cattle Gene Atlas (Fang *et al.* 2020) plotted against mean gene locus taurine local ancestry values in 10 African admixed breeds for the HMG (130 genes), LMG (638 genes), and NMG (14,758 genes) functional subsets (see Table S4 for the data used to generate these plots). Correlation analysis demonstrated that no significant association was observed for the HMG subset TEM values and African taurine local ancestry (Figure S1a; $P = 0.730$). Conversely, as shown in Figures S1b and S1c, very weak but significant negative correlations between TEM and African taurine local ancestry was observed for both the LMG subset ($R = -0.090$; $P = 0.024$) and the much larger NMG subset ($R = -0.021$; $P = 0.010$). In other words, more highly expressed genes in African admixed breeds tend to exhibit greater levels of *B. indicus* ancestry. If we assume that zebu cattle correspond to the introgressive source population then this is the opposite pattern to that observed for archaic human gene flow, where human genes with higher levels of Neanderthal ancestry tend to exhibit lower average multi-tissue expression levels (Sharbrough *et al.* 2017). It is important to keep in mind, however, that as was the case for Neanderthal gene

476 flow, these effects account for a trivial proportion of the variance ($R^2 < 0.01$ for both the LMG

477 and NMG data sets). In addition, direct comparison of the human and cattle results is

478 confounded by the much lower levels of Neanderthal ancestry in modern human populations

479 compared to zebu admixture in African cattle and the much longer time depth in generations

480 since *H. neanderthalensis* and *H. sapiens* interbred (Bergström *et al.* 2021).

4 | DISCUSSION

During the last three decades, increasingly sophisticated molecular tools and statistical

methods have been used to systematically survey genomic variation in African cattle

populations (Loftus *et al.* 1994a; MacHugh *et al.* 1997; Hanotte *et al.* 2002; Decker *et al.* 2014;

Kim *et al.* 2017; Pitt *et al.* 2019; Gebrehiwot *et al.* 2020; Kim *et al.* 2020). These studies have

revealed a complex demographic and evolutionary history fundamentally shaped by gene flow

and admixture between the original substrate of locally adapted humpless taurine animals (*B.*

taurus) and exogenous humped Asian zebu (*B. indicus*) introduced through human migration

and trade (Epstein & Mason 1971; Gifford-Gonzalez & Hanotte 2011). Intriguingly, unlike

other regions of the world such as Southwest Asia and East Asia where taurine and zebu cattle

populations also overlap and interbreed (Edwards *et al.* 2007; Chen *et al.* 2018; Xia *et al.* 2019),

all African cattle examined to-date retain *B. taurus* mitochondrial haplotypes (Figure 4; Loftus

et al. 1994a; Bradley *et al.* 1996; Troy *et al.* 2001; Álvarez *et al.* 2017; Mauki *et al.* 2021). In

this respect, admixed cattle, and animals with predominantly zebu morphology in Africa

represent an example of "massively discordant mitochondrial introgression (MDMI)", a term

introduced by Bonnet *et al.* (2017). This mitochondrial discordance is likely a result of male-

mediated gene flow and genetic drift through preferential dissemination of *B. indicus* genetic

material by a relatively small number of Asian zebu cattle, most of which were bulls (Bradley

et al. 1994; Loftus *et al.* 1994a). The marked evolutionary divergence between the *B. taurus*

and *B. indicus* nuclear genomes and the MDMI observed in admixed African cattle therefore

provide a unique testbed for exploring genomic introgression and mitonuclear interactions in a

numerous large mammalian species spanning diverse physical geographies and agroecological

zones. In addition, the presence of many discrete admixed cattle breeds across the continent

with varying levels of *B. indicus* ancestry (Figure 2a and Figure 3) means that incompatibilities

and coevolution between nuclear and mitochondrial genes can be detected and evaluated in

multiple independent populations.

Evidence for mitonuclear incompatibilities in admixed African cattle breeds

Although previous studies have examined subchromosomal admixture and local ancestry

in hybrid *B. taurus*/*B. indicus* animals (McTavish & Hillis 2014; Chen *et al.* 2018; Koufariotis

et al. 2018; Barbato *et al.* 2020), we present here the first evidence for mitonuclear

incompatibilities in admixed cattle populations. Using a high-density SNP genotyping array,

ten different breeds were examined with genome-wide zebu ancestries ranging between 0.370

(BORG) and 0.740 (KARA), and estimated dates for the start of admixture in each population

extending from the 14^th to the 20^th century (Table 2). A consistent pattern of mitonuclear

disequilibria was observed for the functional HMG subset in admixed African cattle (Figure 7a

and Table 3): African taurine local ancestry was uniformly higher for nuclear genes encoding

proteins that directly engage with mitochondrial-encoded gene products to form multi-subunit

complexes, or that directly interact with mitochondrial DNA or RNAs; this subset encompasses

genes that encode OXPHOS subunits, ribosomal proteins, tRNA synthetases, and DNA and

RNA polymerases. Although the source population divergence is substantially less in admixed

humans, these results are comparable to those obtained by Zaidi and Makova (2019) that

support the hypothesis that selection in admixed human populations has acted against

mitonuclear incompatibility. They observed significant enrichment of sub-Saharan African

ancestry for HMG subset genes in an African American population with sub-Saharan African and European nuclear ancestry and predominantly sub-Saharan African mtDNA haplotypes. They also observed significant enrichment of Native American ancestry at HMG subset genes in a Puerto Rican population with Native American and European nuclear ancestry and predominantly Native American mtDNA haplotypes. It is also noteworthy that the mean ancestry deviation observed for the HMG gene subset in admixed African cattle populations was approximately 2- to 4-fold larger than that observed in the African American and Puerto Rican populations by Zaidi and Makova (2019). This result may reflect stronger selection against mitonuclear incompatibility in admixed cattle that exhibit substantial *B. indicus* nuclear ancestry in parallel with exclusive retention of *B. taurus* mtDNA haplotypes.

For certain admixed African cattle populations (ANKO, KARA, NGAN, ADAG, and KETE) we also observed statistically significant elevated African taurine local ancestry for the LMG gene subset, and except for the ADAG breed, African taurine local ancestry was consistently higher in the HMG subset compared to the LMG subset (Figure 7a and Table 3). A geographic and temporal aspect to potential mitonuclear incompatibility in admixed African cattle populations was also observed. The East African admixed cattle group exhibit a statistically significant larger mean African taurine local ancestry deviation compared to the West African admixed cattle group ($P = 2.2 \times 10^{-16}$; Figure 7b). In addition, somewhat counterintuitively, we observed a decline in African taurine local ancestry deviation for the HMG subset in conjunction with increasing time depth to the onset of population admixture (Figure 7c). The most parsimonious explanation for this pattern is breakdown of African taurine haplotypes at HMG loci through genetic recombination acting across multiple generations.

The functional HMG and LMG subsets containing 138 and 701 genes, respectively were identified, and catalogued in the present study for the purpose of evaluating mitonuclear incompatibilities in admixed African cattle populations. It is also instructive, however, to examine these genes in the context of recently published high-resolution surveys of African cattle genomic diversity and signatures of selection. For example, the aspartyl-tRNA synthetase 2, mitochondrial gene (*DARS2*), an HMG subset gene on BTA16, is in the region encompassed by selective sweeps detected separately in the EASZ breed and a composite sample of East African zebu cattle (Bahbahani *et al.* 2017; Taye *et al.* 2018). In addition, the mitochondrial ribosomal protein S33 gene (*MRPS33*), another HMG subset gene, was detected within a positively selected region on BTA4 when African cattle were compared to commercial European and Asian breeds (Kim *et al.* 2017) and in analyses of selective sweeps focused on the evolution of thermotolerance in African cattle populations (Taye *et al.* 2017). Table S5 provides further information on HMG and LMG subset genes that have been detected in studies of genomic selective sweeps in African cattle populations.

Patterns of local ancestry across African cattle genomes

The patterns of mean African taurine and Asian zebu local ancestry varied across the genome for the 10 African admixed cattle populations, with notable taurine and zebu peaks on BTA7, BTA12, and BTA23, and BTA3, BTA5, and BTA6, respectively. Analysis of genes proximal to the top 1% of taurine and zebu local ancestry peaks demonstrated that genes with high African taurine ancestry were substantial enriched for pathways and physiological systems associated with immunobiology and antimicrobial host defence. Conversely, the genes with high Asian zebu ancestry were overrepresented in pathways and physiological systems associated with organismal growth and development. These functional differences may reflect natural and human-mediated selection for retention of infectious disease resistance or resilience

traits and incorporation of desirable production traits in hybrid African *B. taurus*/*B. indicus*

populations (Mwai *et al.* 2015; Marshall *et al.* 2019).

A particularly notable outcome of the genome-wide analyses of local ancestry was the

observation of the highest signal of African taurine ancestry encompassing the bovine MHC

gene cluster region on BTA23 (Figure 8b). In this regard, a comparable admixture signal has

been observed in a composite North African and Middle Eastern human sample, which

exhibited elevated sub-Saharan ancestry around the MHC gene cluster on HSA6 (Salter-

Townshend & Myers 2019). Enrichment of sub-Saharan African MHC haplotypes has also been

detected in a range of admixed Latin American populations (Norris *et al.* 2020). It is also

noteworthy that previous analyses of genomic microevolution in African cattle populations

have detected selective sweeps at the MHC gene cluster (Kim *et al.* 2017; Taye *et al.* 2018;

Tijjani *et al.* 2019). The bovine MHC region, in common with humans and other mammals,

contains multiple individual genes and gene clusters associated with innate and adaptive

immune responses to pathogens and parasites, including the tumour necrosis factor (*TNF*),

lymphotoxin alpha (*LTA*), and lymphotoxin beta (*LTB*) cytokine genes; the highly polymorphic

genes encoding MHC class I and II molecules responsible for intracellular and extracellular

antigen presentation, respectively (e.g., *BOLA*, *BOLA-NC1*, *BOLA-DMA*, *BOLA-DOA*, *BOLA-*

DQA1, *BOLA-DQB*, and *BOLA-DRA*); and the MHC class I related protein genes (*MIC1*, *MIC2*,

and *MIC3*) (Behl *et al.* 2012; Ellis & Hammond 2014; Maccari *et al.* 2017). A key process

driving evolution of the vertebrate MHC gene cluster region is pathogen- and parasite-mediated

selection caused by infectious disease, particularly balancing selection to maintain allelic

diversity at MHC class I and class II genes (Ebert & Fields 2020; Radwan *et al.* 2020).

Consequently, a marked signal of retention of taurine MHC haplotypes in admixed breeds may

reflect adaptation to significant local disease challenges such as African animal trypanosomiasis

(AAT) caused by a range of *Trypanosoma* species, theileriosis (East Coast fever) caused by

Theileria parva, cowdriosis caused by *Ehrlichia ruminantium*, and Rift Valley fever (RVF)

caused by the RVF virus (Ristic & McIntyre 1981; Seifert 1996). It is also important to note

that patterns of genomic diversity derived from SNP array data for the bovine MHC gene cluster

region could be confounded by ascertainment bias or inaccurate or incomplete genome

annotation. However, ascertainment bias is unlikely to influence our results because the

Illumina® BovineHD 777K BeadChip was designed using a diverse panel representative of

multiple *B. taurus*, *B. indicus* and admixed *B. taurus/indicus* populations (Rincon *et al.* 2011;

Ventura *et al.* 2020). In addition, the 28 SNPs on the array that capture variation at the bovine

MHC gene cluster are located in the relatively stable and well-annotated MHC Class Ib region

(J.A. Hammond, *pers. comm.*).

Conclusions and outlook

Cattle breeding programmes in Africa are currently poised to leverage genomic selection

as a leapfrog technology to bypass conventional genetic evaluation and improvement of

production, health, and welfare traits (Ibeagha-Awemu *et al.* 2019; Marshall *et al.* 2019; Mrode

et al. 2019). Genome-enabled breeding of livestock can also contribute to mitigation of methane

emissions and adaptation to the consequences of climate change (Hayes *et al.* 2013; McManus

et al. 2020); in this regard, the genomic composition of African cattle populations may be

further impacted by the superior thermotolerance of zebu cattle (Hansen 2004; Dikmen *et al.*

2018). The fundamental role of the mitochondrion in the biochemistry and physiology of all

these traits, implies that the mitonuclear ecology framework (Hill 2015) will become

increasingly important for dissecting human-mediated and natural microevolution of livestock.

This will be particularly relevant for African admixed cattle, which as we demonstrate here,

exhibit strong signatures of mitonuclear coevolution. Finally, these populations can serve as

comparative model systems for understanding the phenotypic consequences of mitonuclear

interactions and adaptive and maladaptive genomic introgression in other mammals, including

humans.

ACKNOWLEDGEMENTS

This research was supported by Science Foundation Ireland (SFI) and

Acceligen/Recombinetics Inc. through the SFI CRT in Genomics Data Science under grant no.

18/CRT/6214. Additional support was provided by an SFI Investigator Award to D.E.M (grant

no. SFI/15/IA/3154).

AUTHOR CONTRIBUTIONS

J.A.W., M.J.D., T.S.S., D.G.B., L.A.F., M.S-T., and D.E.M. conceived and designed the

study; J.A.W., T.S.S., D.G.B., and D.E.M coordinated data and sample acquisition; J.A.W. and

G.P.M. conducted the analyses with help from T.J.H, S.I.I.N, L.A.F., and M.S-T.; and J.A.W.

and D.E.M. wrote the manuscript with critical input and revisions from all authors. All authors

gave final approval for publication and agree to be held accountable for the work performed

therein.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

All Illumina® BovineHD 777K BeadChip SNP data used for this study is available from

the Dryad Digital Repository (https://datadryad.org).

REFERENCES

Abin S., Theron H.E. & van Marle-Koster E. (2016) Population structure and genetic trends for indigenous African beef cattle breeds in South Africa. *S. Afr. J. Anim. Sci.* **46**, 152-6.

Allen J.F. (2015) Why chloroplasts and mitochondria retain their own genomes and genetic systems: Colocation for redox regulation of gene expression. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10231-8.

Álvarez I., Pérez-Pardal L., Traoré A., Koudandé D.O., Fernández I., Soudré A., Diarra S., Sanou M., Boussini H. & Goyache F. (2017) Differences in genetic structure assessed using Y-chromosome and mitochondrial DNA markers do not shape the contributions to diversity in African sires. *J. Anim. Breed. Genet.* **134**, 393-404.

Archibald J.M. (2015) Endosymbiosis and eukaryotic cell evolution. *Curr. Biol.* **25**, R911-21.

Bahbahani H., Clifford H., Wragg D., Mbole-Kariuki M.N., Van Tassell C., Sonstegard T., Woolhouse M. & Hanotte O. (2015) Signatures of positive selection in East African Shorthorn Zebu: A genome-wide single nucleotide polymorphism analysis. *Sci. Rep.* **5**, 11729.

Bahbahani H., Tijjani A., Mukasa C., Wragg D., Almathen F., Nash O., Akpa G.N., Mbole-Kariuki M., Malla S., Woolhouse M., Sonstegard T., Van Tassell C., Blythe M., Huson H. & Hanotte O. (2017) Signatures of selection for environmental adaptation and zebu × taurine hybrid fitness in East African Shorthorn Zebu. *Front. Genet.* **8**.

Ballard J.W. & Melvin R.G. (2010) Linking the mitochondrial genotype to the organismal phenotype. *Mol. Ecol.* **19**, 1523-39.

Bar-Yaacov D., Hadjivasiliou Z., Levin L., Barshad G., Zarivach R., Bouskila A. & Mishmar D. (2015) Mitochondrial involvement in vertebrate speciation? The case of mito-nuclear genetic divergence in chameleons. *Genome Biol. Evol.* **7**, 3322-36.

Barbato M., Hailer F., Upadhyay M., Del Corvo M., Colli L., Negrini R., Kim E.S., Crooijmans R., Sonstegard T. & Ajmone-Marsan P. (2020) Adaptive introgression from indicine cattle into white cattle breeds from Central Italy. *Sci. Rep.* **10**, 1279.

Baris T.Z., Wagner D.N., Dayan D.I., Du X., Blier P.U., Pichaud N., Oleksiak M.F. & Crawford D.L. (2017) Evolved genetic and phenotypic differences due to mitochondrial-nuclear interactions. *PLoS Genet.* **13**, e1006517.

Barreto F.S. & Burton R.S. (2013) Elevated oxidative damage is correlated with reduced fitness in interpopulation hybrids of a marine copepod. *Proc. Biol. Sci.* **280**, 20131521.

Behl J.D., Verma N.K., Tyagi N., Mishra P., Behl R. & Joshi B.K. (2012) The major histocompatibility complex in bovines: a review. *ISRN Vet Sci* **2012**, 872710.

Benjamini Y. & Hochberg Y. (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B Method.* **57**, 289-300.

Benson D.A., Cavanaugh M., Clark K., Karsch-Mizrachi I., Ostell J., Pruitt K.D. & Sayers E.W.
(2018) GenBank. *Nucleic Acids Res.* **46**, D41-d7.

Bergström A., Stringer C., Hajdinjak M., Scerri E.M.L. & Skoglund P. (2021) Origins of
modern human ancestry. *Nature* **590**, 229-37.

Bibi F. (2013) A multi-calibrated mitochondrial phylogeny of extant Bovidae (Artiodactyla,
Ruminantia) and the importance of the fossil record to systematics. *BMC Evol. Biol.* **13**,
166.

Blier P.U., Dufresne F. & Burton R.S. (2001) Natural selection and the evolution of mtDNA-
encoded peptides: evidence for intergenomic co-adaptation. *Trends Genet.* **17**, 400-6.

Bonfiglio S., Ginja C., De Gaetano A., Achilli A., Olivieri A., Colli L., Tesfaye K., Agha S.H.,
Gama L.T., Cattonaro F., Penedo M.C., Ajmone-Marsan P., Torroni A. & Ferretti L.
(2012) Origin and spread of *Bos taurus*: new clues from mitochondrial genomes
belonging to haplogroup T1. *PLoS ONE* **7**, e38601.

Bonnet T., Leblois R., Rousset F. & Crochet P.A. (2017) A reassessment of explanations for
discordant introgressions of mitochondrial and nuclear genomes. *Evolution* **71**, 2140-
58.

Boore J.L. (1999) Animal mitochondrial genomes. *Nucleic Acids Res.* **27**, 1767-80.

Bradley D.G., Loftus R.T., Cunningham P. & MacHugh D.E. (1998) Genetics and domestic
cattle origins. *Evol. Anthropol.* **6**, 79-86.

Bradley D.G., MacHugh D.E., Cunningham P. & Loftus R.T. (1996) Mitochondrial diversity
and the origins of African and European cattle. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 5131-
5.

Bradley D.G., MacHugh D.E., Loftus R.T., Sow R.S., Hoste C.H. & Cunningham E.P. (1994)
Zebu-taurine variation in Y chromosomal DNA: a sensitive assay for genetic
introgression in west African trypanotolerant cattle populations. *Anim. Genet.* **25**, 7-12.

Burton R.S., Pereira R.J. & Barreto F.S. (2013) Cytonuclear genomic interactions and hybrid
breakdown. *Annu. Rev. Ecol. Evol. Syst.* **44**, 281-302.

Calvo S.E., Clauser K.R. & Mootha V.K. (2016) MitoCarta2.0: an updated inventory of
mammalian mitochondrial proteins. *Nucleic Acids Res.* **44**, D1251-7.

Chang C.C., Chow C.C., Tellier L.C., Vattikuti S., Purcell S.M. & Lee J.J. (2015) Second-
generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**,
7.

Chen N., Cai Y., Chen Q., Li R., Wang K., Huang Y., Hu S., Huang S., Zhang H., Zheng Z.,
Song W., Ma Z., Ma Y., Dang R., Zhang Z., Xu L., Jia Y., Liu S., Yue X., Deng W.,
Zhang X., Sun Z., Lan X., Han J., Chen H., Bradley D.G., Jiang Y. & Lei C. (2018)
Whole-genome resequencing reveals world-wide ancestry and adaptive introgression
events of domesticated cattle in East Asia. *Nat. Commun.* **9**, 2337.

Chen S., Lin B.Z., Baig M., Mitra B., Lopes R.J., Santos A.M., Magee D.A., Azevedo M.,
Tarroso P., Sasazaki S., Ostrowski S., Mahgoub O., Chaudhuri T.K., Zhang Y.P., Costa
V., Royo L.J., Goyache F., Luikart G., Boivin N., Fuller D.Q., Mannen H., Bradley D.G.
& Beja-Pereira A. (2010) Zebu cattle are an exclusive legacy of the South Asia neolithic.
Mol. Biol. Evol. **27**, 1-6.

Chou J.Y. & Leu J.Y. (2015) The Red Queen in mitochondria: cyto-nuclear co-evolution,
hybrid breakdown and human disease. *Front. Genet.* **6**, 187.

Dadi H., Tibbo M., Takahashi Y., Nomura K., Hanada H. & Amano T. (2009) Variation in
mitochondrial DNA and maternal genetic ancestry of Ethiopian cattle populations.
Anim. Genet. **40**, 556-9.

Dani M.A., Heinneman M.B. & Dani S.U. (2008) Brazilian Nelore cattle: a melting pot
unfolded by molecular genetics. *Genet. Mol. Res.* **7**, 1127-37.

Decker J.E., McKay S.D., Rolf M.M., Kim J., Molina Alcala A., Sonstegard T.S., Hanotte O.,
Gotherstrom A., Seabury C.M., Praharani L., Babar M.E., Correia de Almeida Regitano
L., Yildiz M.A., Heaton M.P., Liu W.S., Lei C.Z., Reecy J.M., Saif-Ur-Rehman M.,
Schnabel R.D. & Taylor J.F. (2014) Worldwide patterns of ancestry, divergence, and
admixture in domesticated cattle. *PLoS Genet.* **10**, e1004254.

Delaneau O., Marchini J. & Zagury J.F. (2012) A linear complexity phasing method for
thousands of genomes. *Nat. Methods* **9**, 179-81.

Derr J.N., Hedrick P.W., Halbert N.D., Plough L., Dobson L.K., King J., Duncan C., Hunter
D.L., Cohen N.D. & Hedgecock D. (2012) Phenotypic effects of cattle mitochondrial
DNA in American bison. *Conserv Biol* **26**, 1130-6.

Dikmen S., Mateescu R.G., Elzo M.A. & Hansen P.J. (2018) Determination of the optimum
contribution of Brahman genetics in an Angus-Brahman multibreed herd for regulation
of body temperature during hot weather. *J. Anim. Sci.* **96**, 2175-83.

Du S.N.N., Khajali F., Dawson N.J. & Scott G.R. (2017) Hybridization increases mitochondrial
production of reactive oxygen species in sunfish. *Evolution* **71**, 1643-52.

Dyall S.D., Brown M.T. & Johnson P.J. (2004) Ancient invasions: from endosymbionts to
organelles. *Science* **304**, 253-7.

Ebert D. & Fields P.D. (2020) Host-parasite co-evolution and its genomic signature. *Nat. Rev.*
Genet. **21**, 754-68.

Edwards C.J., Baird J.F. & MacHugh D.E. (2007) Taurine and zebu admixture in Near Eastern
cattle: a comparison of mitochondrial, autosomal and Y-chromosomal data. *Anim.*
Genet. **38**, 520-4.

Ellis S.A. & Hammond J.A. (2014) The functional significance of cattle major
histocompatibility complex class I genetic diversity. *Annu. Rev. Anim. Biosci.* **2**, 285-
306.

Ellison C.K. & Burton R.S. (2006) Disruption of mitochondrial function in interpopulation hybrids of *Tigriopus californicus*. *Evolution* **60**, 1382-91.

Ellison C.K. & Burton R.S. (2008) Interpopulation hybrid breakdown maps to the mitochondrial genome. *Evolution* **62**, 631-8.

Ellison C.K., Niehuis O. & Gadau J. (2008) Hybrid breakdown and mitochondrial dysfunction in hybrids of *Nasonia* parasitoid wasps. *J. Evol. Biol.* **21**, 1844-51.

Epstein H. & Mason I.L. (1971) *The Origin of the Domestic Animals of Africa (vol. 1)*. Africana Publishing, London.

Fang L., Cai W., Liu S., Canela-Xandri O., Gao Y., Jiang J., Rawlik K., Li B., Schroeder S.G., Rosen B.D., Li C.J., Sonstegard T.S., Alexander L.J., Van Tassell C.P., VanRaden P.M., Cole J.B., Yu Y., Zhang S., Tenesa A., Ma L. & Liu G.E. (2020) Comprehensive analyses of 723 transcriptomes enhance genetic and biological interpretations for complex traits in cattle. *Genome Res.* **30**, 790-801.

Freeman A.R., Meghen C.M., MacHugh D.E., Loftus R.T., Achukwi M.D., Bado A., Sauveroche B. & Bradley D.G. (2004) Admixture and diversity in West African cattle populations. *Mol. Ecol.* **13**, 3477-87.

Fuller D.Q. (2006) Agricultural origins and frontiers in South Asia: a working synthesis. *J. World Prehist.* **20**, 1-86.

Gebrehiwot N.Z., Strucken E.M., Aliloo H., Marshall K. & Gibson J.P. (2020) The patterns of admixture, divergence, and ancestry of African cattle populations determined from genome-wide SNP data. *BMC Genomics* **21**, 869.

Gifford-Gonzalez D. & Hanotte O. (2011) Domesticating animals in Africa: implications of genetic and archaeological findings. *J. World Prehist.* **24**, 1-23.

Ginja C., Gama L.T., Cortés O., Burriel I.M., Vega-Pla J.L., Penedo C., Sponenberg P., Cañón J., Sanz A., do Egito A.A., Alvarez L.A., Giovambattista G., Agha S., Rogberg-Muñoz A., Lara M.A.C., Delgado J.V. & Martinez A. (2019) The genetic ancestry of American Creole cattle inferred from uniparental and autosomal genetic markers. *Sci. Rep.* **9**, 11486.

Gonzalez S. (2021) The role of mitonuclear incompatibility in bipolar disorder susceptibility and resilience against environmental stressors. *Front. Genet.* **12**, 636294.

Hanotte O., Bradley D.G., Ochieng J.W., Verjee Y., Hill E.W. & Rege J.E. (2002) African pastoralism: genetic imprints of origins and migrations. *Science* **296**, 336-9.

Hansen P.J. (2004) Physiological and cellular adaptations of zebu cattle to thermal stress. *Anim. Reprod. Sci.* **82-83**, 349-60.

Hayes B.J., Lewin H.A. & Goddard M.E. (2013) The future of livestock breeding: genomic selection for efficiency, reduced emissions intensity, and adaptation. *Trends Genet.* **29**, 206-14.

Hiendleder S., Lewalski H. & Janke A. (2008) Complete mitochondrial genomes of *Bos taurus*
and *Bos indicus* provide new insights into intra-species variation, taxonomy and
domestication. *Cytogenet. Genome Res.* **120**, 150-6.

Hill G.E. (2015) Mitonuclear ecology. *Mol. Biol. Evol.* **32**, 1917-27.

Hill G.E. (2016) Mitonuclear coevolution as the genesis of speciation and the mitochondrial
DNA barcode gap. *Ecol Evol* **6**, 5831-42.

Hill G.E. (2017) The mitonuclear compatibility species concept. *The Auk* **134**, 393-409.

Hirase S., Tezuka A., Nagano A.J., Sato M., Hosoya S., Kikuchi K. & Iwasaki W. (2021)
Integrative genomic phylogeography reveals signs of mitonuclear incompatibility in a
natural hybrid goby population. *Evolution* **75**, 176-94.

Ibeagha-Awemu E.M., Peters S.O., Bemji M.N., Adeleke M.A. & Do D.N. (2019) Leveraging
available resources and stakeholder involvement for improved productivity of African
livestock in the era of genomic breeding. *Front. Genet.* **10**, 357.

Isaac R.S., McShane E. & Churchman L.S. (2018) The multiple levels of mitonuclear
coregulation. *Annu. Rev. Genet.* **52**, 511-33.

Kasahara A. & Scorrano L. (2014) Mitochondria: from cell death executioners to regulators of
cell differentiation. *Trends Cell Biol.* **24**, 761-70.

Kim J., Hanotte O., Mwai O.A., Dessie T., Bashir S., Diallo B., Agaba M., Kim K., Kwak W.,
Sung S., Seo M., Jeong H., Kwon T., Taye M., Song K.D., Lim D., Cho S., Lee H.J.,
Yoon D., Oh S.J., Kemp S., Lee H.K. & Kim H. (2017) The genome landscape of
indigenous African cattle. *Genome Biol.* **18**, 34.

Kim K., Kwon T., Dessie T., Yoo D., Mwai O.A., Jang J., Sung S., Lee S., Salim B., Jung J.,
Jeong H., Tarekegn G.M., Tijjani A., Lim D., Cho S., Oh S.J., Lee H.-K., Kim J., Jeong
C., Kemp S., Hanotte O. & Kim H. (2020) The mosaic genome of indigenous African
cattle as a unique genetic resource for African pastoralism. *Nat. Genet.* **52**, 1099-110.

Kong M., Xiang H., Wang J., Liu J., Zhang X. & Zhao X. (2020) Mitochondrial DNA
haplotypes influence energy metabolism across chicken transmitochondrial cybrids.
Genes (Basel) **11**.

Koufariotis L., Hayes B.J., Kelly M., Burns B.M., Lyons R., Stothard P., Chamberlain A.J. &
Moore S. (2018) Sequencing the mosaic genome of Brahman cattle identifies historic
and recent introgression including polled. *Sci. Rep.* **8**, 17761.

Kramer A., Green J., Pollard J., Jr. & Tugendreich S. (2014) Causal analysis approaches in
Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-30.

Latorre-Pellicer A., Moreno-Loshuertos R., Lechuga-Vieco A.V., Sanchez-Cabo F., Torroja C.,
Acin-Perez R., Calvo E., Aix E., Gonzalez-Guerra A., Logan A., Bernad-Miana M.L.,
Romanos E., Cruz R., Cogliati S., Sobrino B., Carracedo A., Perez-Martos A.,
Fernandez-Silva P., Ruiz-Cabello J., Murphy M.P., Flores I., Vazquez J. & Enriquez

J.A. (2016) Mitochondrial and nuclear DNA matching shapes metabolism and healthy
ageing. *Nature* **535**, 561-5.

Lechuga-Vieco A.V., Justo-Méndez R. & Enríquez J.A. (2021) Not all mitochondrial DNAs
are made equal and the nucleus knows it. *IUBMB Life* **73**, 511-29.

Lee-Yaw J.A., Jacobs C.G. & Irwin D.E. (2014) Individual performance in relation to
cytonuclear discordance in a northern contact zone between long-toed salamander
(*Ambystoma macrodactylum*) lineages. *Mol. Ecol.* **23**, 4590-602.

Leigh J.W. & Bryant D. (2015) POPART: full-feature software for haplotype network
construction. *Methods Ecol. Evol.* **6**, 1110-6.

Loftus R.T., MacHugh D.E., Bradley D.G., Sharp P.M. & Cunningham P. (1994a) Evidence
for two independent domestications of cattle. *Proc. Natl. Acad. Sci. U. S. A.* **91**, 2757-
61.

Loftus R.T., MacHugh D.E., Ngere L.O., Balain D.S., Badi A.M., Bradley D.G. & Cunningham
E.P. (1994b) Mitochondrial genetic variation in European, African and Indian cattle
populations. *Anim. Genet.* **25**, 265-71.

Lyons T.W., Reinhard C.T. & Planavsky N.J. (2014) The rise of oxygen in Earth's early ocean
and atmosphere. *Nature* **506**, 307-15.

Ma L., O'Connell J.R., VanRaden P.M., Shen B., Padhi A., Sun C., Bickhart D.M., Cole J.B.,
Null D.J., Liu G.E., Da Y. & Wiggans G.R. (2015) Cattle sex-specific recombination
and genetic control from a large pedigree analysis. *PLoS Genet.* **11**, e1005387.

Maccari G., Robinson J., Ballingall K., Guethlein L.A., Grimholt U., Kaufman J., Ho C.S., de
Groot N.G., Flicek P., Bontrop R.E., Hammond J.A. & Marsh S.G. (2017) IPD-MHC
2.0: an improved inter-species database for the study of the major histocompatibility
complex. *Nucleic Acids Res.* **45**, D860-d4.

MacHugh D.E., Shriver M.D., Loftus R.T., Cunningham P. & Bradley D.G. (1997)
Microsatellite DNA variation and the evolution, domestication and phylogeography of
taurine and zebu cattle (*Bos taurus* and *Bos indicus*). *Genetics* **146**, 1071-86.

Marshall K., Gibson J.P., Mwai O., Mwacharo J.M., Haile A., Getachew T., Mrode R. & Kemp
S.J. (2019) Livestock genomics for developing countries - African examples in practice.
Front. Genet. **10**, 297.

Martin W.F., Garg S. & Zimorski V. (2015) Endosymbiotic theories for eukaryote origin.
Philos. Trans. R. Soc. Lond. B. Biol. Sci. **370**, 20140330.

Mauki D.H., Adeola A.C., Ng'ang'a S.I., Tijjani A., Akanbi I.M., Sanke O.J., Abdussamad
A.M., Olaogun S.C., Ibrahim J., Dawuda P.M., Mangbon G.F., Gwakisa P.S., Yin T.T.,
Peng M.S. & Zhang Y.P. (2021) Genetic variation of Nigerian cattle inferred from
maternal and paternal genetic markers. *PeerJ* **9**, e10607.

McBride H.M., Neuspiel M. & Wasiak S. (2006) Mitochondria: more than just a powerhouse. *Curr. Biol.* **16**, R551-60.

McHugo G.P., Dover M.J. & MacHugh D.E. (2019) Unlocking the origins and biology of domestic animals using ancient DNA and paleogenomics. *BMC Biol.* **17**, 98.

McKenzie M., Chiotis M., Pinkert C.A. & Trounce I.A. (2003) Functional respiratory chain analyses in murid xenomitochondrial cybrids expose coevolutionary constraints of cytochrome b and nuclear subunits of complex III. *Mol. Biol. Evol.* **20**, 1117-24.

McKenzie M., Trounce I.A., Cassar C.A. & Pinkert C.A. (2004) Production of homoplasmic xenomitochondrial mice. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 1685-90.

McManus C.M., Rezende Paiva S. & Faria D. (2020) Genomics and climate change. *Rev. Sci. Tech.* **39**, 481-90.

McTavish E.J. & Hillis D.M. (2014) A genomic approach for distinguishing between recent and ancient admixture as applied to cattle. *J. Hered.*

Mészáros G., Boison S.A., Pérez O'Brien A.M., Ferenčaković M., Curik I., Da Silva M.V., Utsunomiya Y.T., Garcia J.F. & Sölkner J. (2015) Genomic analysis for managing small and endangered populations: a case study in Tyrol Grey cattle. *Front. Genet.* **6**, 173.

Mills E.L., Kelly B. & O'Neill L.A.J. (2017) Mitochondria are the powerhouses of immunity. *Nat. Immunol.* **18**, 488-98.

Morales H.E., Pavlova A., Amos N., Major R., Kilian A., Greening C. & Sunnucks P. (2018) Concordant divergence of mitogenomes and a mitonuclear gene cluster in bird lineages inhabiting different climates. *Nat. Ecol. Evol.* **2**, 1258-67.

Mrode R., Ojango J.M.K., Okeyo A.M. & Mwacharo J.M. (2019) Genomic selection and use of molecular tools in breeding programs for indigenous and crossbred cattle in developing countries: current status and future prospects. *Front. Genet.* **9**, 694.

Mwai O., Hanotte O., Kwon Y.J. & Cho S. (2015) African indigenous cattle: unique genetic resources in a rapidly changing world. *Asian-Australas. J. Anim. Sci.* **28**, 911-21.

Norris E.T., Rishishwar L., Chande A.T., Conley A.B., Ye K., Valderrama-Aguirre A. & Jordan I.K. (2020) Admixture-enabled selection for rapid adaptive evolution in the Americas. *Genome Biol.* **21**, 29.

O'Brien A.M., Holler D., Boison S.A., Milanesi M., Bomba L., Utsunomiya Y.T., Carvalheiro R., Neves H.H., da Silva M.V., VanTassell C.P., Sonstegard T.S., Meszaros G., Ajmone-Marsan P., Garcia F. & Solkner J. (2015) Low levels of taurine introgression in the current Brazilian Nelore and Gir indicine cattle populations. *Genet. Sel. Evol.* **47**, 31.

Pichaud N., Bérubé R., Côté G., Belzile C., Dufresne F., Morrow G., Tanguay R.M., Rand D.M. & Blier P.U. (2019) Age dependent dysfunction of mitochondrial and ROS metabolism induced by mitonuclear mismatch. *Front. Genet.* **10**, 130.

Pitt D., Sevane N., Nicolazzi E.L., MacHugh D.E., Park S.D.E., Colli L., Martinez R., Bruford
M.W. & Orozco-terWengel P. (2019) Domestication of cattle: Two or three events?
Evol. Appl. **12**, 123-36.

Pramod R.K., Velayutham D., P K.S., P S.B., Zachariah A., Zachariah A., B C., S S.S., P G.,
Dhinoth Kumar B., Iype S., Gupta R., Santhosh S. & Thomas G. (2019) Complete
mitogenome reveals genetic divergence and phylogenetic relationships among Indian
cattle (*Bos indicus*) breeds. *Anim. Biotechnol.* **30**, 219-32.

Quinlan A.R. & Hall I.M. (2010) BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841-2.

R Core Team (2019) R: A language and environment for statistical computing. R Foundation
for Statistical Computing. http://www.R-project.org, Vienna, Austria.

Radwan J., Babik W., Kaufman J., Lenz T.L. & Winternitz J. (2020) Advances in the
evolutionary understanding of MHC polymorphism. *Trends Genet.* **36**, 298-311.

Raj A., Stephens M. & Pritchard J.K. (2014) fastSTRUCTURE: variational inference of
population structure in large SNP data sets. *Genetics* **197**, 573-89.

Rand D.M., Haney R.A. & Fry A.J. (2004) Cytonuclear coevolution: the genomics of
cooperation. *Trends Ecol. Evol.* **19**, 645-53.

Rank N.E., Mardulyn P., Heidl S.J., Roberts K.T., Zavala N.A., Smiley J.T. & Dahlhoff E.P.
(2020) Mitonuclear mismatch alters performance and reproductive success in naturally
introgressed populations of a montane leaf beetle. *Evolution* **74**, 1724-40.

Rincon G., Weber K.L., Eenennaam A.L., Golden B.L. & Medrano J.F. (2011) Hot topic:
performance of bovine high-density genotyping platforms in Holsteins and Jerseys. *J.*
Dairy Sci. **94**, 6116-21.

Ristic M. & McIntyre I. (1981) *Diseases of Cattle in the Tropics: Economic and Zoonotic*
Relevance. Martinus Nijhoff Publishers, The Hague, Netherlands.

Rosen B.D., Bickhart D.M., Schnabel R.D., Koren S., Elsik C.G., Tseng E., Rowan T.N., Low
W.Y., Zimin A., Couldrey C., Hall R., Li W., Rhie A., Ghurye J., McKay S.D., Thibaud-
Nissen F., Hoffman J., Murdoch B.M., Snelling W.M., McDaneld T.G., Hammond J.A.,
Schwartz J.C., Nandolo W., Hagen D.E., Dreischer C., Schultheiss S.J., Schroeder S.G.,
Phillippy A.M., Cole J.B., Van Tassell C.P., Liu G., Smith T.P.L. & Medrano J.F.
(2020) De novo assembly of the cattle reference genome with single-molecule
sequencing. *Gigascience* **9**.

Rosenberg N.A. (2005) Algorithms for selecting informative marker panels for population
assignment. *J. Comput. Biol.* **12**, 1183-201.

Rosenberg N.A., Li L.M., Ward R. & Pritchard J.K. (2003) Informativeness of genetic markers
for inference of ancestry. *Am. J. Hum. Genet.* **73**, 1402-22.

Salter-Townshend M. & Myers S. (2019) Fine-scale inference of ancestry segments without
prior knowledge of admixing groups. *Genetics* **212**, 869-89.

Scheet P. & Stephens M. (2006) A fast and flexible statistical model for large-scale population
genotype data: applications to inferring missing genotypes and haplotypic phase. *Am. J.*
Hum. Genet. **78**, 629-44.

Seifert H.S.H. (1996) *Tropical Animal Health*. Kluwer Academic Publishers, Dordrecht,
Netherlands.

Seixas F.A., Boursot P. & Melo-Ferreira J. (2018) The genomic impact of historical
hybridization with massive mitochondrial DNA introgression. *Genome Biol.* **19**, 91.

Senczuk G., Mastrangelo S., Ajmone-Marsan P., Becskei Z., Colangelo P., Colli L., Ferretti L.,
Karsli T., Lancioni H., Lasagna E., Marletta D., Persichilli C., Portolano B., Sarti F.M.,
Ciani E. & Pilla F. (2021) On the origin and diversification of Podolian cattle breeds:
testing scenarios of European colonization using genome-wide SNP data. *Genet. Sel.*
Evol. **53**, 48.

Sharbrough J., Havird J.C., Noe G.R., Warren J.M. & Sloan D.B. (2017) The mitonuclear
dimension of Neanderthal and Denisovan ancestry in modern human genomes. *Genome*
Biol. Evol. **9**, 1567-81.

Sloan D.B., Fields P.D. & Havird J.C. (2015) Mitonuclear linkage disequilibrium in human
populations. *Proc. Biol. Sci.* **282**.

Sloan D.B., Havird J.C. & Sharbrough J. (2017) The on-again, off-again relationship between
mitochondrial genomes and species boundaries. *Mol. Ecol.* **26**, 2212-36.

Sloan D.B., Warren J.M., Williams A.M., Wu Z., Abdel-Ghany S.E., Chicco A.J. & Havird
J.C. (2018) Cytonuclear integration and co-evolution. *Nat. Rev. Genet.* **19**, 635-48.

Spierer A.N., Yoon D., Zhu C.T. & Rand D.M. (2021) FreeClimber: automated quantification
of climbing performance in *Drosophila*. *J. Exp. Biol.* **224**.

Spinelli J.B. & Haigis M.C. (2018) The multifaceted contributions of mitochondria to cellular
metabolism. *Nat. Cell Biol.* **20**, 745-54.

Sujkowski A., Spierer A.N., Rajagopalan T., Bazzell B., Safdar M., Imsirovic D., Arking R.,
Rand D.M. & Wessells R. (2019) Mito-nuclear interactions modify *Drosophila* exercise
performance. *Mitochondrion* **47**, 188-205.

Sunnucks P., Morales H.E., Lamb A.M., Pavlova A. & Greening C. (2017) Integrative
approaches for studying mitochondrial and nuclear genome co-evolution in oxidative
phosphorylation. *Front. Genet.* **8**, 25.

Taye M., Lee W., Caetano-Anolles K., Dessie T., Cho S., Jong Oh S., Lee H.-K. & Kim H.
(2018) Exploring the genomes of East African Indicine cattle breeds reveals signature
of selection for tropical environmental adaptation traits. *Cogent Food. Agric.* **4**,
1552552.

Taye M., Lee W., Caetano-Anolles K., Dessie T., Hanotte O., Mwai O.A., Kemp S., Cho S.,
Oh S.J., Lee H.K. & Kim H. (2017) Whole genome detection of signature of positive
selection in African cattle reveals selection for thermotolerance. *Anim. Sci. J.* **88**, 1889-
901.

Tijjani A., Utsunomiya Y.T., Ezekwe A.G., Nashiru O. & Hanotte O. (2019) Genome sequence
analysis reveals selection signatures in endangered trypanotolerant West African
Muturu cattle. *Front. Genet.* **10**, 442.

Tiku V., Tan M.W. & Dikic I. (2020) Mitochondrial functions in infection and immunity.
Trends Cell Biol. **30**, 263-75.

Tourmente M., Hirose M., Ibrahim S., Dowling D.K., Tompkins D.M., Roldan E.R.S. &
Gemmell N.J. (2017) mtDNA polymorphism and metabolic inhibition affect sperm
performance in conplastic mice. *Reproduction* **154**, 341-54.

Trier C.N., Hermansen J.S., Sætre G.P. & Bailey R.I. (2014) Evidence for mito-nuclear and
sex-linked reproductive barriers between the hybrid Italian sparrow and its parent
species. *PLoS Genet.* **10**, e1004075.

Troy C.S., MacHugh D.E., Bailey J.F., Magee D.A., Loftus R.T., Cunningham P., Chamberlain
A.T., Sykes B.C. & Bradley D.G. (2001) Genetic evidence for Near-Eastern origins of
European cattle. *Nature* **410**, 1088-91.

Utsunomiya Y.T., Milanesi M., Fortes M.R.S., Porto-Neto L.R., Utsunomiya A.T.H., Silva M.,
Garcia J.F. & Ajmone-Marsan P. (2019) Genomic clues of the evolutionary history of
Bos indicus cattle. *Anim. Genet.* **50**, 557-68.

Ventura R.V., Brito L.F., Oliveira G.A., Daetwyler H.D., Schenkel F.S., Sargolzaei M.,
Vandervoort G., Silva F.F.E., Miller S.P., Carvalho M.E., Santana M.H.A., Mattos E.C.,
Fonseca P., Eler J.P. & Ferraz J.B.S. (2020) A comprehensive comparison of high-
density SNP panels and an alternative ultra-high-density panel for genomic analyses in
Nellore cattle. *Anim. Prod. Sci.* **60**, 333-46.

Verdugo M.P., Mullin V.E., Scheu A., Mattiangeli V., Daly K.G., Maisano Delser P., Hare
A.J., Burger J., Collins M.J., Kehati R., Hesse P., Fulton D., Sauer E.W., Mohaseb F.A.,
Davoudi H., Khazaeli R., Lhuillier J., Rapin C., Ebrahimi S., Khasanov M., Vahidi
S.M.F., MacHugh D.E., Ertugrul O., Koukouli-Chrysanthaki C., Sampson A., Kazantzis
G., Kontopoulos I., Bulatovic J., Stojanovic I., Mikdad A., Benecke N., Linstadter J.,
Sablin M., Bendrey R., Gourichon L., Arbuckle B.S., Mashkour M., Orton D., Horwitz
L.K., Teasdale M.D. & Bradley D.G. (2019) Ancient cattle genomics, origins, and rapid
turnover in the Fertile Crescent. *Science* **365**, 173-6.

Wang J., Xiang H., Liu L., Kong M., Yin T. & Zhao X. (2017) Mitochondrial haplotypes
influence metabolic traits across bovine inter- and intra-species cybrids. *Sci. Rep.* **7**,
4179.

Wang K., Lenstra J.A., Liu L., Hu Q., Ma T., Qiu Q. & Liu J. (2018) Incomplete lineage sorting rather than hybridization explains the inconsistent phylogeny of the wisent. *Commun. Biol.* **1**, 169.

Weinberg S.E., Sena L.A. & Chandel N.S. (2015) Mitochondria in the regulation of innate and adaptive immunity. *Immunity* **42**, 406-17.

Weir B.S. & Cockerham C.C. (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution* **38**, 1358-70.

Wickham H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer Publishing Company, New York, USA.

Wolff J.N., Ladoukakis E.D., Enriquez J.A. & Dowling D.K. (2014) Mitonuclear interactions: evolutionary consequences over multiple biological scales. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **369**, 20130443.

Woodson J.D. & Chory J. (2008) Coordination of gene expression between organellar and nuclear genomes. *Nat. Rev. Genet.* **9**, 383-95.

Wu D.D., Ding X.D., Wang S., Wojcik J.M., Zhang Y., Tokarska M., Li Y., Wang M.S., Faruque O., Nielsen R., Zhang Q. & Zhang Y.P. (2018) Pervasive introgression facilitated domestication and adaptation in the *Bos* species complex. *Nat. Ecol. Evol.* **2**, 1139-45.

Xia X., Qu K., Zhang G., Jia Y., Ma Z., Zhao X., Huang Y., Chen H., Huang B. & Lei C. (2019) Comprehensive analysis of the mitochondrial DNA diversity in Chinese cattle. *Anim. Genet.* **50**, 70-3.

Yang Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586-91.

Yates A.D., Achuthan P., Akanni W., Allen J., Allen J., Alvarez-Jarreta J., Amode M.R., Armean I.M., Azov A.G., Bennett R., Bhai J., Billis K., Boddu S., Marugan J.C., Cummins C., Davidson C., Dodiya K., Fatima R., Gall A., Giron C.G., Gil L., Grego T., Haggerty L., Haskell E., Hourlier T., Izuogu O.G., Janacek S.H., Juettemann T., Kay M., Lavidas I., Le T., Lemos D., Martinez J.G., Maurel T., McDowall M., McMahon A., Mohanan S., Moore B., Nuhn M., Oheh D.N., Parker A., Parton A., Patricio M., Sakthivel M.P., Abdul Salam A.I., Schmitt B.M., Schuilenburg H., Sheppard D., Sycheva M., Szuba M., Taylor K., Thormann A., Threadgold G., Vullo A., Walts B., Winterbottom A., Zadissa A., Chakiachvili M., Flint B., Frankish A., Hunt S.E., G I.I., Kostadima M., Langridge N., Loveland J.E., Martin F.J., Morales J., Mudge J.M., Muffato M., Perry E., Ruffier M., Trevanion S.J., Cunningham F., Howe K.L., Zerbino D.R. & Flicek P. (2020) Ensembl 2020. *Nucleic Acids Res.* **48**, D682-D8.

Yu X., Gimsa U., Wester-Rosenlöf L., Kanitz E., Otten W., Kunz M. & Ibrahim S.M. (2009) Dissecting the effects of mtDNA variations on complex traits using mouse conplastic strains. *Genome Res.* **19**, 159-65.

Zaidi A.A. & Makova K.D. (2019) Investigating mitonuclear interactions in human admixed
populations. *Nat. Ecol. Evol.* **3**, 213-22.

Zardoya R. (2020) Recent advances in understanding mitochondrial genome diversity.
F1000Res **9**.

Zeder M.A. (2011) The origins of agriculture in the Near East. *Curr. Anthropol.* **52**, S221-S35.

Zhang K., Lenstra J.A., Zhang S., Liu W. & Liu J. (2020) Evolution and domestication of the
Bovini species. *Anim. Genet.* **51**, 637-57.

1052

 TABLE 1 Cattle breeds/populations, geographical origins, and sources of BovineHD 777K SNP data.

Cattle breed/population	Code	Type/morphology	Country of origin	Source 1[a]	Source 2[b]	Source 3[c]	Source 4[d]	Total (*n*)
Muturu	MUTU	West African taurine	Nigeria	8	---	20	---	28
N'Dama	NDAG	West African taurine	Guinea	24	9	---	23	56
Holstein-Friesian	HOLS	European taurine	Netherlands	59	---	---	---	59
Jersey	JRSY	European taurine	United Kingdom	32	---	---	---	32
Ankole	ANKO	East African admixed	Uganda	25	---	---	---	25
East African Shorthorn Zebu	EASZ	East African admixed	Kenya	92	---	19	---	111
Karamojong	KARA	East African admixed	Uganda	16	---	---	---	16
Nganda	NGAN	East African admixed	Uganda	23	---	4	---	27
Adamawa Gudali	ADAG	West African admixed	Nigeria	23	---	2	---	25
Borgou	BORG	West African admixed	Benin	---	---	---	50	50
Bunaji	BUNA	West African admixed	Nigeria	22	---	5	---	27

Keteku	KETE	West African admixed	Nigeria	---	---	22	---	22
Red Bororo	REDB	West African admixed	Nigeria	22	---	4	---	26
Sokoto Gudali	SOKG	West African admixed	Nigeria	19	---	---	---	19
Gir	GIR	Asian zebu	India	28	---	---	---	28
Hariana	HARI	Asian zebu	India	---	10	---	---	10
Nelore	NELO	Asian zebu	Brazil	34	---	---	---	34
Sahiwal	SAHI	Asian zebu	India	---	10	---	---	10

1054   [a] Bahbahani *et al.* (2015); [b] Verdugo *et al.* (2019); [c] Acceligen cattle genotyping database; [d] genotyped for the present study.
1055

TABLE 2 Ancestry components estimated using FASTSTRUCTURE with estimated times for the start of the admixture process in African admixed

cattle populations generated using MOSAIC.

Code	Type/morphology	Country of origin	African taurine ancestry	European taurine ancestry	Asian zebu ancestry	Generations since start of admixture	Start of admixture
MUTU	West African taurine	Nigeria	0.990 ± 0.019	0.010 ± 0.018	0.000 ± 0.003		----
NDAG	West African taurine	Guinea	1.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000		----
HOLS	European taurine	Netherlands	0.010 ± 0.025	0.990 ± 0.025	0.000 ± 0.000		----
JRSY	European taurine	United Kingdom	0.010 ± 0.022	0.990 ± 0.022	0.000 ± 0.000		----
ANKO	East African admixed	Uganda	0.420 ± 0.010	0.020 ± 0.016	0.550 ± 0.013	54.1	1641 – 1804 CE
EASZ	East African admixed	Kenya	0.250 ± 0.015	0.020 ± 0.065	0.730 ± 0.055	46.2	1697 – 1835 CE
KARA	East African admixed	Uganda	0.260 ± 0.008	0.000 ± 0.001	0.740 ± 0.008	65.1	1564 – 1760 CE
NGAN	East African admixed	Uganda	0.310 ± 0.024	0.100 ± 0.052	0.590 ± 0.039	15.9	1909 – 1956 CE
ADAG	West African admixed	Nigeria	0.320 ± 0.010	0.000 ± 0.001	0.680 ± 0.010	79.6	1463 – 1702 CE
BORG	West African admixed	Benin	0.610 ± 0.047	0.010 ± 0.027	0.370 ± 0.055	20.5	1877 – 1938 CE

BUNA	West African admixed	Nigeria	0.340 ± 0.076	0.000 ± 0.007	0.660 ± 0.083	87.9	1405 – 1668 CE
KETE	West African admixed	Nigeria	0.400 ± 0.093	0.000 ± 0.001	0.600 ± 0.093	26.6	1834 – 1914 CE
REDB	West African admixed	Nigeria	0.320 ± 0.009	0.000 ± 0.001	0.680 ± 0.009	70	1530 – 1740 CE
SOKG	West African admixed	Nigeria	0.320 ± 0.008	0.000 ± 0.000	0.680 ± 0.008	95.9	1349 – 1636 CE
GIR	Asian zebu	India	0.000 ± 0.000	0.000 ± 0.000	1.000 ± 0.000		----
HARI	Asian zebu	India	0.000 ± 0.000	0.000 ± 0.000	1.000 ± 0.000		----
NELO	Asian zebu	Brazil	0.000 ± 0.000	0.000 ± 0.001	1.000 ± 0.001		----
SAHI	Asian zebu	India	0.000 ± 0.000	0.000 ± 0.000	1.000 ± 0.000		----

1058 *Note.* For each admixed population, the generations since admixture started were obtained from the NDAG:GIR coancestry plots (Figure 4). A
1059 generation interval range of 4–7 years for managed domestic cattle was used. CE = Common Era.

1060 **TABLE 3** Statistical test results for comparisons of HMG, LMG, and NMG distributions of
1061 African taurine local ancestry deviations for East and West African admixed cattle.

Code	Type/morphology	*P* value (HMG > NMG)	*P* value (LMG > NMG)	*P* value (HMG > LMG)
ANKO	East African admixed	5.69×10^{-76}	2.73×10^{-20}	4.88×10^{-24}
EASZ	East African admixed	3.09×10^{-206}	0.9567	7.51×10^{-239}
KARA	East African admixed	1.52×10^{-86}	5.10×10^{-5}	2.30×10^{-61}
NGAN	East African admixed	6.41×10^{-315}	1.87×10^{-10}	4.13×10^{-250}
ADAG	West African admixed	2.02×10^{-14}	1.46×10^{-41}	1.0000
BORG	West African admixed	8.74×10^{-41}	0.6345	1.98×10^{-50}
BUNA	West African admixed	2.56×10^{-45}	0.9998	3.35×10^{-81}
KETE	West African admixed	2.41×10^{-206}	1.6×10^{-20}	3.54×10^{-110}
REDB	West African admixed	1.08×10^{-51}	1.0000	6.00×10^{-94}
SOKG	West African admixed	8.13×10^{-22}	1.0000	1.61×10^{-95}

1062

1063

1064

FIGURE 1 Geographical patterns of African *Bos taurus* and Asian *Bos indicus* admixture and a schematic illustrating mitonuclear interactions for hybrid African cattle that retain the taurine mitochondrial genome. (a) Map of Africa showing an interpolated synthetic map illustrating spatial distribution of African *B. taurus* and Asian *B. indicus* admixture. Admixture data was generated from the first principal component (PC1) of a principal component analysis (PCA) of microsatellite genetic variation across African cattle populations (Hanotte *et al.* 2002). Modified from (McHugo *et al.* 2019) under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0). (b) Mitonuclear interactions that can give rise to mitonuclear incompatibilities in crossbred and hybrid African cattle populations. All African cattle surveyed to-date retain the taurine mitochondrial genome (some figure components created with a BioRender.com licence).

1071

1072
FIGURE 2 Results of the principal component analysis (PCA) for 605 animals from 18 different cattle breeds genotyped for 562,635 SNPs. (a) PCA plot showing the coordinates for
1073 each animal based on the first two principal components. Principal component 1 (PC1) differentiates the *Bos taurus* and *Bos indicus* evolutionary lineages, while PC2 separates the
1074 African and European taurine groups. (b) Histogram plot showing the relative variance contributions for the first 10 PCs with PC1 and PC2 accounting for 58.4% and 17.9% of the
1075 total variation for PC1–10, respectively.
1076
1077

FIGURE 3 Unsupervised genetic structure plot for Asian zebu, East and West African admixed cattle, and West African and European taurine breeds. Results for an inferred number of ancestry clusters of *K* = 3 is shown, which corresponds to Asian *Bos indicus* (red), European *Bos taurus* (green), and African *B. taurus* (blue) ancestral components, respectively.

1078
1079
1080
1081

1082

FIGURE 4 Phylogenetic network of 491 cattle mtDNA haplotypes generated using 39 ancestry-informative mtDNA SNPs. This mtDNA haplotype network demonstrates that all surveyed African cattle (47 taurine, 156 East African admixed, and 136 West African admixed) possess and retain *Bos taurus* mitochondrial genomes.

FIGURE 5 Coancestry curve plots generated using MOSAIC for 10 East and West African admixed cattle populations. These curves show the exponential decay of the ratio of probabilities of pairs of local ancestries (y-axis) as a function of genetic distance (x-axis). The pair of ancestries used for each curve is shown on the top of each plot with the estimated number of generations since the start of admixture in brackets. For each plot, the green line represents the fitted curve, the black line shows the across targets ratio, and the grey lines indicate the per target ratio (further information in Salter-Townshend & Myers 2019).

FIGURE 6 Local ancestry plots for bovine chromosome 23 (BTA23) showing Asian *Bos indicus* (red) and African *Bos taurus* (blue) subchromosomal ancestry in four East African and six West African admixed cattle populations. Physical distance along BTA23 is indicated in megabases (Mb).

FIGURE 7 African taurine local ancestry deviations for three different functional gene sets. (a) Violin plots of African taurine local ancestry deviations for the HMG, LMG, and NMG subsets with positive deviations indicating retention of African taurine gene haplotypes. Black data points indicate the median values and horizontal lines represent the 95% confidence interval. (b) Box plot of African taurine local ancestry deviations for the HMG subset in the East African and the West African admixed groups. White lines indicate the median values and yellow and purple boxes indicate the interquartile ranges. (c) Linear regression plot of mean taurine local ancestry deviation for the HMG subset against generations since the start of admixture for four East (yellow) and six West African (purple) admixed cattle breeds. A significant negative Pearson correlation coefficient was observed for these data ($R = -0.678$; $P = 0.031$) and the grey shaded area shows the 95% confidence interval for the regression line.

(a)

(b)

1104
1105

FIGURE 8 Mean African taurine local ancestry across all ten African admixed breeds shown genome-wide and individually for bovine chromosomes 5, 12 and 23 (BTA5, BTA12 and BTA23). (a) Genome-wide mean African taurine local ancestry with a solid black line indicating the global mean, dotted black lines bounding the global 95% confidence interval (CI), and orange lines indicating the global minimum and maximum. The three chromosomes shown in panel (b) are indicated with black rectangles. (b) Chromosome-wide plots for BTA5, BTA12, BTA23 with dark blue lines indicating mean African taurine local ancestry for the ten breeds, blue shading indicating the 95% CI, solid black lines indicating the global mean and dashed blue lines indicating the chromosomal means. The location of the major histocompatibility complex (MHC) gene cluster is shown on BTA23.

1114