Cryptococcus neoformans melanization incorporates multiple catecholamines to produce polytypic melanin

Rosanna P. Baker[1], Christine Chrissian[2], Ruth E. Stark[2], Arturo Casadevall[1],*

[1]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD 21205, USA

[2]Department of Chemistry and Biochemistry, The City College of New York and CUNY Institute for Macromolecular Assemblies, New York, NY 10031, USA

*Correspondence: E-mail: acasade1@jhu.edu; Tel.: 410-955-3457; Fax: 410-955-0105

Running title: *Cryptococcal* melanization incorporates multiple precursors

Keywords: fungi, virulence factor, melanin, catecholamine, dopamine, solid-state NMR, reactive oxygen species

Abstract

Melanin is a major virulence factor in pathogenic fungi that enhances the ability of fungal cells to resist immune clearance. *Cryptococcus neoformans* is an important human pathogenic fungus that synthesizes melanin from exogenous tissue catecholamine precursors during infection, but the type of melanin made in cryptococcal meningoencephalitis is unknown. We analyzed the efficacy of various catecholamines found in brain tissue in supporting melanization using animal brain tissue and synthetic catecholamine mixtures reflecting brain tissue proportions. Solid-state NMR spectra of the melanin pigment produced from such mixtures displayed larger melanin contributions than expected from the most robust dopamine constituent, suggesting uptake of additional catecholamines. Probing the biosynthesis of melanin using radiolabeled catecholamines revealed that *C. neoformans* melanization simultaneously incorporated more than one catecholamine, implying that the pigment was polytypic in nature. Nonetheless, melanin derived from individual or mixed catecholamines had comparable ability to protect *C. neoformans* against ultraviolet light and oxidants. Our results indicate that melanin produced during infection differs depending on the catecholamine composition of tissue and that melanin pigment synthesized *in vivo* is likely to accrue from the polymerization of a mixture of precursors. From a practical standpoint our results strongly suggest that using dopamine as a polymerization precursor is capable of producing melanin pigment comparable to that produced during infection. On a more fundamental level our findings uncover additional structural complexity for natural cryptococcal melanin by demonstrating that pigment produced during human infection is likely to be composed of polymerized moieties derived from chemically different precursors.

Introduction

Cryptococcus neoformans is an encapsulated, free-living fungal organism that can establish opportunistic infections in both plant and animal hosts (1). C. neoformans is broadly prevalent in the environment and colonizes a variety of ecological niches, which notably include soil and bird guano (2). Despite ubiquitous exposure of the human population, C. neoformans rarely causes disease in individuals with intact immunity as the infection is either cleared or becomes asymptomatically latent (3). However, C. neoformans poses a serious threat to those with compromised immunity resulting from disease or medical treatment, especially AIDS patients, and is estimated to cause 180,000 deaths annually (4). Cryptococcal infection commences with the inhalation of desiccated fungal cells or spores that become deposited in the lungs, which can result in cryptococcal pneumonia for immunocompromised individuals (5). Progression of the primary infection can occur through extrapulmonary dissemination as yeast cells dispersed through the bloodstream are able to cross the blood brain barrier and take up residence in brain tissue (6, 7). The ensuing meningoencephalitis that is typical of this aggressive stage of cryptococcosis, if left untreated, is almost always fatal (8). Although concerted efforts are underway to develop an effective vaccine (9), there is currently no reliable means of preventing infection and even the most current treatment strategies that employ a combined anti-fungal regimen only decrease 10-week mortality rates to 24% (10). Thus, an improved understanding of factors that contribute to fungal virulence is vital to our efforts in combating this harmful pathogen.

Among the adaptations that safeguard C. neoformans from harsh conditions both in the environment and during host infection is the deposition of a layer of melanin in the fungal cell wall. Melanins are a diverse group of pigments, typically black or dark brown in color, that are formed by the oxidative polymerization of phenolic or indolic precursors (11). They are notorious for their extreme recalcitrance, structural complexity, and unique biophysical properties, such as the ability to absorb a wide spectrum of UV light. Melanin pigments are widely employed throughout the biosphere, serving such diverse functions as camouflage, thermal modulation, and protection from radiation (12); in pathogenic fungi, they are linked to increased virulence (13) and drug resistance (14). In C. neoformans cells, melanin is synthesized from phenolic compounds in small vesicles called melanosomes that are transported and subsequently

deposited within the cell wall (15). There, the melanin particles assemble into larger granules that become tightly interwoven with the constituents comprising the cell wall, which consequently thickens over time (16). In the environment, melanization is credited with protecting *C. neoformans* from electromagnetic radiation, temperature stresses, and amoeba (17, 18). During infection, melanization helps shield *C. neoformans* cells from engulfment and killing by host macrophages (19–21) and contributes to the severity of infection by promoting neurotropism (22, 23). As melanization is one of two primary virulence factors for *C. neoformans*, the disruption of this process, perhaps by inhibiting the laccase enzyme responsible for catalyzing melanin synthesis, has been widely recognized as a potential therapeutic target.

A defining feature of the *C. neoformans* laccase enzyme is its inability to produce melanin pigments from endogenously synthesized compounds like tyrosine, the precursor most commonly used by other melanotic organisms (24). Instead, *C. neoformans* draws on exogenous sources of substrates, producing melanin pigments from a wide variety of phenolic compounds found in natural habitats and in animal hosts, as evidenced by the isolation of melanized cells from both natural isolates (25) and infected host tissues (26, 27). Indeed, the animal nervous system serves as a rich source of precursors in the form of catecholamines, nitrogen-containing diphenolic compounds which include the neurotransmitters dopamine, epinephrine, and norepinephrine. *C. neoformans* melanization has been studied *in vitro* by culturing cells in media supplemented with each of these compounds and more commonly, with L-Dopa, the immediate biosynthetic precursor of dopamine. Considerable insight into the properties of *C. neoformans* melanin has been gained by applying biophysical techniques like solid-state nuclear magnetic resonance (NMR) and electron paramagnetic resonance (EPR) to analyze melanin containing particles derived from such cultures (28–30). These analyses have indicated that the pigments produced by *C. neoformans* from each of these precursors are indeed melanins. They have also revealed distinct differences in molecular structure and EPR signal intensity but the implications of these differences with respect to virulence remain unclear.

Whereas previous studies of *C. neoformans* melanin have largely employed L-Dopa as the precursor (reviewed in (31)), the low abundance of this compound in host tissues prompted us to question how faithfully L-Dopa melanin produced in culture reflects the structure and function of 'natural' melanin produced during the course of *C. neoformans* infection. The progression of cryptococcal disease ultimately results in brain dissemination, which provides a

rich and varied source of catecholamines, thereby presenting the potential for simultaneous incorporation of multiple precursors. Although acid-resistant melanin particles can be isolated from infected human and animal tissue (27, 32), the limited quantity of melanized material that can be recovered from animal models of infection has precluded extensive biophysical analyses. To circumvent this obstacle, we sought to develop a cell culture system that would approximate the repertoire of catecholamine precursors present in the human brain. Drawing on previous neurochemical studies that quantified catecholamines isolated from postmortem human brains (33, 34), we melanized *C. neoformans* in a mixture of three catecholamines according to their relative brain proportions of 60% dopamine, 33% norepinephrine, and 7% epinephrine. Here we describe the isolation and analysis of melanin produced from this brain catecholamine mixture (BCM) and provide evidence that *C. neoformans* can incorporate multiple precursors simultaneously during melanin synthesis. BCM is found to be comparable to L-Dopa melanin in its ability to protect *C. neoformans* from sources of cellular damage like UV radiation and free oxygen radicals. Despite these functional similarities, the ability of *C. neoformans* to utilize a mixture of chemically distinct precursors during melanin synthesis implies architectural heterogeneity for this polymer depending on the local abundance of precursors and significantly complicates its structural characterization.

Results

***C. neoformans* KN99α variant achieves robust melanization in conditions mimicking the human brain environment**

We sought to approximate the melanization process during infection by growing *C. neoformans* in a mixture of catecholamines representative of human brain proportions (brain catecholamine mixture, BCM). A key consideration in this effort was the selection of a laboratory strain variant that would be capable of robust melanin production from BCM at the human body temperature of 37˚C. Several variants derived from the widely used H99 strain of *C. neoformans* var. *grubii* (H99O), including H99E, H99C, and H99W, display reduced virulence in mice and decreased melanization on L-Dopa agar as a result of laboratory passage (35). Conversely, variants H99S, H99F, and KN99α/a that were derived from passage of H99 through the rabbit

central nervous system model are more virulent and produce more melanin when grown on L-Dopa (35). We compared melanization by H99O and KN99α on agar supplemented with BCM as well as the individual precursors DA, NE, and L-Dopa at 30˚C and 37˚C (Fig. 1A). The well-characterized H99-derived ΔLAC1 strain deficient in Laccase-1, the primary enzyme required for melanization, was included as a negative control. *C. neoformans* laccase production and activity has been shown previously to decrease with increasing temperature (36, 37). Accordingly, we observed a slower rate of melanin production and a diminished amount of overall pigmentation for both H99O and KN99α grown at 37˚C compared to 30˚C, regardless of which precursors were provided. Whereas H99O melanization was delayed compared to that of KN99α at 30˚C, similar degrees of pigmentation were observed for the two strains after incubation for 48 h (Fig. 1A, left panel). However, H99O cells showed significantly less pigmentation than KN99α after 48 h at 37˚C (Fig. 1A, right panel; Fig. 1B) and only achieved equivalent melanization levels after an additional two days of growth (data not shown). Melanization by KN99α on BCM-supplemented agar was comparable to the pigmentation observed with each of the individual catecholamines at 30˚C (Fig. 1C, left graph) and was more efficient than NE alone at 37˚C (Fig. 1C, right graph). Given the increased melanin production of KN99α compared to H99O when grown in the presence of BCM at the human body temperature of 37˚C, the KN99α variant of *C. neoformans* was chosen for all subsequent analyses.

Melanin-containing 'ghosts' can be isolated from BCM-supplemented cultures and brain catecholamine extract plates

　　　To further characterize the melanin produced by *C. neoformans* when provided with a mixture of catecholamine precursors, KN99α was melanized in liquid culture for 10 d at 30˚C in MM supplemented with BCM (0.6 mM DA / 0.33 mM NE / 0.07 mM E) or either 1 mM DA or 1 mM NE. Pigmentation intensity increased more rapidly for cells grown in BCM compared to DA initially but both cultures reached maximal melanization levels after 5 d (Fig. 2A). The culture grown in NE melanized more slowly but eventually reached an intensity comparable to the other two cultures after 10 d (Fig. 2A). The robust pigment production achieved in the BCM-supplemented culture supports the utility of this large-scale culture system for acquiring melanin

that approximates that produced in human brain tissue but in sufficient quantities to permit structural characterization.

To obtain melanin for biophysical analysis, cells harvested from the three cultures were subjected to enzymatic cell wall digestion, protein denaturation and hydrolysis, lipid extraction, and finally boiling in concentrated HCl. In each case, acid-resistant material was recovered and examination by light microscopy verified that the isolated particles were dark, yeast-shaped silhouettes referred to as melanin 'ghosts'. However, differences in the physical appearance of particles derived from each culture were noted. The melanin ghost particles isolated from the DA control culture were darkly pigmented (Fig. 2B) and similar in appearance to those isolated previously from cultures melanized in L-Dopa (38). Fewer intact particles were recovered from the NE control culture and they tended to be small and pale (Fig. 2B); there was an overall lower yield of acid resistant material (70 mg/L) compared to that of DA (130 mg/L). The BCM melanin ghosts were intermediate between the two controls both in appearance, with a mixture of intact dark particles and more fragile, pale particles (Fig. 2B), and in terms of the yield of material (100 mg/L). Following the same protocol used to isolate BCM ghost particles, comparable acid-resistant particles were recovered from KN99α cells grown on agar plates containing brain tissue extract (Fig. 2C), although the yield of intact particles was very small (<1 in 10^6 of the total number of cells).

Solid-state NMR suggests that cells provided BCM incorporate more than one precursor into the melanized cell wall architecture

Given the strong resemblance of the particles derived from *C. neoformans* grown in BCM to melanin ghosts, we sought to characterize the molecular profile of this pigmented material. Solid-state NMR (ssNMR) analysis was implemented to examine its molecular architecture and compare it to that of melanin ghost material isolated from cells grown with either of the two major constituent precursors, DA or NE. The 1D ^{13}C cross-polarization (CPMAS) spectra of each of these three samples clearly displayed all of the spectral features that are characteristic of melanin ghosts (Fig. 3) (28–30), notably the group of broad, overlapping peaks located between ~110 and 160 ppm. These peaks correspond to the various aromatic carbons that comprise the inherently heterogeneous melanin polymer: although many of the carbons are chemically similar, they differ in both their bonding patterns and local chemical environments, and consequently

resonate at slightly different frequencies. This results in a multitude of signals that overlap in the aromatic-carbon spectral region, forming what visually appears as a single, broad peak and serves as a telltale indicator of the presence of melanin pigments. Other characteristic features of the cellular components of melanin ghosts were also observed in all three sample spectra: the spectral regions between each ~10-40 ppm and ~50-110 ppm displayed groups of peaks which are attributable to long-chain fatty acids within triacylglycerols and polysaccharide-ring carbons, respectively (39).

Our ssNMR analysis also provided insight into whether *C. neoformans* is able to incorporate more than one melanization precursor into the pigment architecture when provided multiple substrates. The heterogenous nature of the melanin polymer which results in multiple broad, overlapping peaks precludes the identification of specific signals that are unique to a melanin synthesized from DA or from NE. Through visual inspection alone it was evident that the DA, NE and BCM melanin ghost samples all share common spectral features but the relative intensities of the groups of peaks corresponding to pigments, lipids and polysaccharides clearly differed. Notably, the aromatic-carbon signals attributed to the pigment contributed substantially less to the total signal intensity in the NE-only melanin ghost spectrum in comparison to the DA-only spectrum, even though both samples were isolated from cell cultures that contained 1 mM concentrations of the respective precursors. This finding was anticipated since prior studies had demonstrated that precursors such as DA or L-Dopa result in more robust melanin production in comparison to others such as NE or E (28–30). This, in turn, results in melanin ghosts that have a greater pigment content relative to the other cellular remnants present such as polysaccharide and lipids. The spectrum of the BCM melanin ghosts, which were isolated from a cell culture containing 0.6 mM DA, 0.33 mM NE and 0.07 mM E, exhibited pigment contributions with respect to other cellular constituents that exceeded expectations based on the concentration of the robust DA precursor, suggesting incorporation of the additional catecholamines. Quantitatively reliable direct polarization (DPMAS) spectra reinforced the trend of higher than expected pigment contributions in the melanin ghosts (Fig. S1). Thus, even without detailed discrimination of the pigment structures synthesized from different precursors, our ssNMR analysis suggested that *C. neoformans* can use at least two different melanization precursors when they are provided simultaneously.

Radiolabeled norepinephrine and dopamine are incorporated into C. neoformans melanin

As our structural data provided highly suggestive evidence for the incorporation of multiple precursors, we extended our analysis by employing radiolabeled catecholamines as a sensitive and quantitative means of determining whether both DA and NE can be incorporated into the melanin pigment at the same time. Individual *C. neoformans* KN99α cultures were grown in the presence of a low concentration (0.0001 mM) of high specific activity ^{3}H-labeled NE with and without the addition of unlabeled NE or DA. A control culture deficient in the primary laccase enzyme (ΔLAC1) was also supplemented with the same concentration of ^{3}H-NE as a negative control. Even though the molar concentration of NE in the culture was very low, we detected significantly more radioactivity retained in the cell pellet from the KN99α cell culture grown without any additional unlabeled precursors compared to that of the ΔLAC1 control culture (Fig. 4A, graph). The addition of excess unlabeled NE at a concentration of either 0.1 or 1 mM resulted in a noticeable, concentration-dependent increase in pigmentation (Fig. 4A, lower panels) and a concomitant increase in the incorporation efficiency of ^{3}H-NE (Fig. 4A, graph). We attribute these increases to the established dose dependence of melanization for DA and NE precursors, possibly resulting from induction of the laccase enzyme (40). Notably, a comparable increase in ^{3}H-NE incorporation was also detected when cultures were supplemented with excess (1 mM) of unlabeled DA (Fig. 4A, graph), indicating that *C. neoformans* can utilize more than one type of precursor simultaneously.

To confirm that the radioactivity associated with *C. neoformans* cell pellets resulted from the incorporation of the radiolabeled precursors into melanin, we heated cells that had been melanized in mixtures of radiolabeled and unlabeled precursors in concentrated hydrochloric acid to generate melanin ghost particles. Approximately 30-40% of the radioactivity was retained in melanin from cells labeled with either ^{3}H-NE or ^{3}H-DA (Fig. 4B). The efficiency of incorporation of either radioactively-labeled precursor into melanin was not significantly different when melanization was supported by an excess of a single unlabeled catecholamine or a 1:1 mixture of NE and DA (Fig. 4B) since in either case, a substantial proportion of the ^{3}H radioactivity was retained after HCl treatment.

Melanin derived from different precursors provide similar protection from UV radiation and oxidative damage

Melanization of *C. neoformans* in L-Dopa was previously demonstrated to afford protection from the fungicidal effects of UV radiation (41). To investigate the protective capacity of the brain mixture of catecholamines, KN99α cells were melanized for 7 d in BCM, DA, NE, or L-Dopa and compared to non-melanized control cells for survival after UV exposure. Cells melanized under all conditions showed significantly higher percentages of survival compared to non-melanized cells, although melanization in NE was less protective than with the other three precursors (Fig. 5A). Melanization in BCM showed no statistical difference from DA or L-Dopa in shielding cells from UV radiation (P values of 0.8107 and 0.6413 respectively).

Since melanin is a key virulence factor that contributes to *C. neoformans* survival inside human host cells, we sought to explore the ability of 'natural' melanin to safeguard against damaging free radicals generated inside macrophages. An *in vitro* oxidative system that produces oxygen-derived free radicals (42) was used to treat KN99α cells that were non-melanized or had been melanized for 7 d in the BCM, DA, NE, or L-Dopa. Melanization in the presence of each of these precursors, with the exception of NE, significantly increased the percentage of cells that survived exposure to oxidative damage compared to non-melanized cells (Fig. 5B).

Discussion

Melanization by *C. neoformans* is a prominent virulence factor that facilitates evasion from the host immune system. Melanin produced by *C. neoformans* grown in defined culture media supplemented with the phenolic precursor, L-Dopa, has been well characterized, but much less is known about the protective layer of pigment deposited during infection. In the brain, L-Dopa is readily converted to dopamine, which is in turn a precursor of norepinephrine and epinephrine. Hence, *C. neoformans* cells that disseminate to brain tissue are expected to utilize one or more of these three catecholamines for melanin synthesis. We cultured *C. neoformans* in a mixture of dopamine, norepinephrine and epinephrine reflective of their relative proportions in the human brain and found that the resulting melanin was similar to L-Dopa melanin in its ability to protect cells from UV radiation and oxidative damage. Structural analysis using solid-state NMR revealed a spectrum consistent with the simultaneous incorporation of more than one precursor.

Melanization experiments employing radiolabeled catecholamines provided additional evidence that *C. neoformans* can use multiple precursors simultaneously and suggested that melanin made during infection is likely synthesized from a mixture of available substrates.

Initial evidence for *C. neoformans* melanization during infection was provided by the isolation of acid-resistant melanin ghost particles from infected animal and human tissue and from cells grown on tissue homogenate agar plates (27, 32). We isolated similar particles from cells grown on agar plates infused with a lamb brain catecholamine extract but only a very small fraction of the total cells were recovered as melanin ghosts. Neurotransmitters such as dopamine in brain tissue are contained within vesicles and degraded by monoamine oxidase, an enzyme found in glial cells (43). Consequently, several factors could have contributed to the small yield when using brain tissue as the source of substrate, including the relatively lower concentrations of catecholamines in brain tissue compared to the concentrations achievable in culture, low bioavailability of the precursors sequestered in neurotransmitter vesicles, and degradation of catecholamines during the extraction procedure when cells are lysed. In light of these limitations, we opted for culturing *C. neoformans* in a mixture of catecholamines proportional to their relative concentrations present in the human brain and achieved robust melanization on both solid agar substrates and in liquid cultures. We also observed more efficient melanization by the KN99α variant compared to its parental H99O strain, especially at the human body temperature of 37°C, and this is consistent with the correlation between melanization and virulence (35).

The melanin produced by *C. neoformans* when provided with a mixture of catecholamines in the proportions found in the human brain was subjected to structural analysis using solid-state NMR. The relative percentages of 60% dopamine, 33% norepinephrine, and 7% epinephrine were calculated from the global averages measured across all regions of the brain (33, 34). Melanin ghost particles isolated from a KN99α culture supplemented with the 'brain mixture' of catecholamines had relatively more pigment than expected based on the DA concentration. Structural analysis by ssNMR revealed comparable ^{13}C-CPMAS spectra for melanin ghost particles derived from the three cultures with all of the hallmark features of *C. neoformans* melanized cell walls, including remnants of the polysaccharide scaffold and trapped neutral lipids in addition to the purified fungal pigments. The presence of signals corresponding to each of these components confirmed that the black particles recovered from *C. neoformans* cell cultures

supplemented with BCM are indeed melanin ghosts. Quantitative analysis of the relative melanin content in BCM-melanin by DPMAS also indicated a larger amount of pigment than would be expected from the incorporation of DA alone and suggested the use of more than one precursor in the mixture. The increased melanin pigment signal for DA compared to NE melanin ghosts suggests that *C. neoformans* laccase may incorporate DA more efficiently than other catecholamines. Consistent with this notion is our observation that pigmentation increased more rapidly for cells grown on solid or in liquid media supplemented with DA than with NE (Figs. 1 and 2A).

We wondered if the preference for DA incorporation could be substantial enough to preclude the use of other catecholamines when provided in combination. To explore this possibility, we cultured *C. neoformans* in the presence of radiolabeled catecholamines to follow their incorporation into melanin. A significantly higher amount of radioactivity was detected in the cell pellet of the wild-type KN99α strain compared to a laccase-deficient mutant, ΔLAC1, which suggests that the labeled precursor was used to support melanization. However, the efficiency of incorporation by KN99α was very low at approximately 5%, which is not unexpected given the low molar concentration of 0.0001 mM for the labeled precursor. *C. neoformans* melanization in culture was previously shown to have a dose-dependence for L-Dopa and norepinephrine substrates, with an optimal concentration of 1 mM (40). In this instance, supplementing cultures of KN99α with 0.1 and 1 mM unlabeled NE in combination with 0.0001 mM ^{3}H-NE resulted in a concentration-dependent increase in both visible pigmentation and incorporation of ^{3}H-NE. Remarkably, melanization with 1 mM unlabeled DA also significantly increased the incorporation of ^{3}H-NE, although not to the same extent as 1 mM NE, possibly reflecting a slight incorporation bias for DA. However, the increased efficiency of ^{3}H-NE incorporation in the presence of a 10,000-fold molar excess of DA supports the conclusion that *C. neoformans* does not use DA to the exclusion of NE. To confirm that the radiolabeled precursors were being incorporated into melanin pigment and not just peripherally associated with cells, we isolated acid-resistant melanin ghosts from radiolabeled cultures. Using either ^{3}H-NE or ^{3}H-DA, we were able to isolate radiolabeled melanin when cells were melanized in either individual unlabeled catecholamines or an equimolar mixture of NE and DA. Together, these results suggest that the most readily abundant substrate is used to establish a melanin scaffold that is then built upon using any and all available precursors, thereby generating local structural diversity for melanin particles.

The study of *C. neoformans* melanization has used predominantly L-Dopa as a substrate but this precursor would not be the major substrate available in host tissues. *C. neoformans* infections are initially established in the lungs where they encounter alveolar macrophages. As these and other immune cells are known to produce and release catecholamines (44), it is possible that melanization supported by lung resident catecholamines could enhance virulence and promote brain dissemination. The ability of *C. neoformans* to incorporate multiple catecholamines simultaneously suggests that melanin synthesized in brain tissue may differ in distinct anatomical regions as the relative proportions of these neurotransmitters have been found to vary widely from one area of the brain to another (33, 34). Intriguingly, *C. neoformans* infections are often found to be concentrated in the basal ganglia region of the brain where DA levels are highest (45, 46), a finding that could reflect resistance of melanized cells to clearance by host immune mechanisms. Basal ganglia preference is also consistent with the notion that melanin produced by *C. neoformans* in the brain is primarily dopamine-derived. Consequently, we suggest that future studies of the role of melanin in *C. neoformans* pathogenesis are more likely to reflect brain infection conditions if dopamine is used as the melanization substrate. However, L-Dopa may be a more suitable substrate to study the role of cryptococcal melanization in the environment where L-Dopa is a common precursor of plant compounds and accumulates in soils (47). We note that although the type of melanin produced by *C. neoformans* is dependent on the available precursor, our prior ssNMR studies revealed considerable structural similarity between different melanins (28–30).

An improved understanding of the composition and structure of 'natural' melanin produced by *C. neoformans* during infection will help elucidate its function in virulence. We have shown that *C. neoformans* can incorporate multiple precursors, even when their molar ratios differ by several orders of magnitude. Given that melanin synthesis is driven by the accretion of highly reactive intermediates into a solid material, it is expected that *C. neoformans* provided with a mixture of precursors produces polytypic melanin. Furthermore, depending on the relative availability of specific catecholamines, structurally diverse forms of melanin may be produced in different anatomical regions during the course of infection. Despite this added level of complexity, melanins produced in culture from single or mixed substrates were found to be functionally comparable in their capacity to protect *C. neoformans* from UV radiation and oxidative damage, but whether specific forms of melanin could imbue *C. neoformans* with a

greater survival advantage during infection remains unanswered. While the present study has shed light on initial polymerization events, the resulting melanin compound is a highly complex, amorphous substance for which detailed structural elucidation remains beyond our analytical horizon.

Experimental Procedures

Yeast strains

Cryptococcus neoformans serotype A strains H99O (American Type Culture Collection 208821), H99-ΔLAC1 (Fungal Genetics Stock Center, 2007 Lodge library), and KN99α (Fungal Genetics Stock Center, 2007 Lodge library) were recovered from frozen 50% glycerol stocks by growth in YPD broth (Difco 242820) for 48 h at 30°C. The hygromycin resistance cassette in H99-ΔLAC1 was retained by supplementing culture media with 200 U/ml Hygromycin B from *Streptomyces sp.* (Calbiochem 400051). Cells were washed twice with sterile phosphate-buffered saline before sub-culturing into defined media.

Culture media with relative proportions of catecholamines in the human brain

To study melanization under conditions representative of the human brain, we used the total cortical and subcortical measurements of the three brain catecholamines reported previously (33, 34) to calculate the relative proportions of approximately 60% dopamine (DA), 33% norepinephrine (NE), and 7% epinephrine (E). *C. neoformans* variant KN99α was grown in minimal media (MM) consisting of 29.4 mM K_2HPO_4, 10 mM $MgSO_4$, 13 mM glycine, 15 mM D-glucose, and 3 μM thiamine at pH 5.5 that was supplemented with the brain catecholamine mixture (BCM) consisting of 0.6 mM Dopamine hydrochloride (MilliporeSigma H8502), 0.33 mM (-)-Norepinephrine (MilliporeSigma A7257), and 0.07 mM (-)-Epinephrine (MilliporeSigma E4250). Control cultures were also grown in MM supplemented with 1.0 mM DA or 1.0 mM NE.

Melanization assay on agar plates

To prepare melanization assay plates, 4X MM with no additives or with either 1.0 mM DA, 1.0 mM NE, 1.0 mM 3,4-Dihydroxy-L-phenylalanine (L-Dopa, MilliporeSigma D9628), or BCM was mixed 1:3 with cooled 2% agar (Difco 214530) and poured into individual wells of 24-well plates

(2 ml per well). Plates were allowed to cool at RT for 5 h, then 1×10^6 PBS-washed KN99α, H99O, or H99-ΔLAC1 cells were pipetted into each well and plates were incubated at either 30˚C or 37˚C. After 24 and 48 h, the plates that had been incubated at each temperature were photographed in a single field using a 12-megapixel camera. Images were converted to grayscale in Adobe Photoshop (version 22.0.1) and intensities were quantified using Image Studio Lite software (version 5.2.5, Li-Cor Biosciences). The signal measured for the MM control was applied as the image background and intensity measurements for the H99-ΔLAC1 negative control were subtracted from those of KN99α and H99O to derive pigmentation levels above background.

Isolation of melanin particles

Cultures of KN99α were grown in 0.5 L MM supplemented with either BCM or 1.0 mM DA, and in 1.0 L MM with 1.0 mM NE. Melanization of the cultures was monitored by photographing and quantifying pigment intensities (as described above) for 2 mL samples that had been transferred to the wells of a transparent 12-well plate (Celltreat, 229112). After incubation for 10 d at 30˚C, acid-resistant melanin 'ghost' particles were isolated from cultures as described in detail previously (28, 32). Briefly, cells were treated with a 10 mg/ml solution of *Trichoderma harzianum* lysing enzymes (MilliporeSigma L1412) in 1.0 M sorbitol and 0.1 M sodium citrate for 24 h at 30˚C to remove the fungal cell wall. The resulting protoplasts were subjected to protein denaturation with 4.0 M guanidinium thiocyanate for 18 h at RT and then proteolysis with a 20 mg/ml solution of proteinase K (MilliporeSigma 03115801001) in 10 mM Tris-HCl (pH 8), 5.0 mM CaCl$_2$, and 0.5% sodium dodecyl sulfate for 4 h at 65˚C. Following lipid removal by three successive extractions with a 4:8:3 ratio of methanol:chloroform:saline (48) samples were heated at 90˚C in 6 M HCl for 1 h. Melanin-containing particles were recovered by centrifugation and dialyzed for several days against pure H$_2$O prior to lyophilization. Dry material was weighed using an analytical balance (Mettler Toledo AT261 DeltaRange) to determine yield.

Melanization in brain tissue extracts

The tissue from three lamb brains (227 g) was homogenized in ~1 ml/g ice cold buffer comprised of 0.1 M perchloric acid, 1.0 mM sodium metabisulfite, and 0.1 mM EDTA using a 30 ml Potter-Elvehjem tissue grinder (Wheaton 358011) attached to a drill press. Following centrifugation at

5000 rpm for 20 min at 4°C, the pH of the supernatant was adjusted to ~5 by dropwise addition of sodium hydroxide and then passed through a 0.22 μm filter (Millipore SCGPS02RE). The filtrate was rapidly frozen on dry ice and then lyophilized for ~72 h. The freeze-dried extract was resuspended with 75 ml 2X MM containing 1% Penicillin-Streptomycin, mixed with 75 ml cooled 3% agar and poured into six 100 ml petri dishes. After ~24 h at RT, plates were spotted with PBS-washed KN99α and incubated at 30°C for 26 d. Melanin ghost particles were isolated from cells melanized in liquid cultures and on plates as described previously.

Solid-state NMR of melanin particles

Solid-state NMR measurements were carried out on a Varian (Agilent) DirectDrive2 (DD2) spectrometer operating at a ^{1}H frequency of 600 MHz. All data were acquired on ~8-10 mg of lyophilized material using a 1.6-mm T3 HXY fastMAS probe (Agilent Technologies, Santa Clara, CA) with a magic-angle spinning (MAS) rate of 15.00 +\- 0.02 kHz and a spectrometer-set temperature of 25 °C. The 1D ^{13}C cross-polarization (CPMAS) experiments were conducted with 90° pulse lengths of 1.5 and 1.7 μs for ^{1}H and ^{13}C, respectively, a 10% linearly ramped amplitude for ^{1}H and a 1 ms contact time for ^{1}H-^{13}C polarization transfer. The 1D ^{13}C direct polarization (DPMAS) experiments were conducted with 90° pulse lengths of 1.2 and 1.4 μs for ^{1}H and ^{13}C, respectively, and long recycle delays (50 s) were used in order to obtain quantitatively reliable signal intensities. Heteronuclear decoupling was applied during signal acquisition using the small phase incremental alternation pulse sequence (SPINAL) (49) and a radio-frequency field strength of 104 kHz.

Radiolabeled norepinephrine incorporation analysis

KN99α was grown in MM for 2 d at 30°C, then cells were washed and resuspended in fresh media at a cell density of 2×10^8 cells/ml. Cultures were labeled with 2 μCi/ml (0.0001 mM) norepinephrine DL-[7-^{3}H] hydrochloride (^{3}H-NE, American Radiolabeled Chemicals, Inc. ART-0172) for 7 d at 30°C in the absence or presence of 0.1 or 1.0 mM unlabeled NE or 0.1 or 1.0 mM unlabeled DA. H99-ΔLAC1 was also labeled with ^{3}H-NE as a negative control. Color photographs were taken of each culture to document relative pigmentation intensities. Labeled cells were collected by centrifugation at 16,000 $\times$ g for 5 min, washed twice with MM, then resuspended in MM. Samples of media, wash buffer, and resuspended pellets were mixed with

UniverSol-ES liquid scintillation cocktail (MP Biomedicals 01882480) and decays per minute (DPM) were quantified using an LS 6500 liquid scintillation counter (Beckman, Indianapolis, IN). Percent incorporation was calculated by dividing the DPM measured for the pellet sample by the total DPM of the pellet, media, and wash samples.

Isolation of radiolabeled melanin particles

KN99α was grown as described above in MM supplemented with 2 μCi/ml ^{3}H-NE and 1.0 mM unlabeled NE or a mixture of 1.0 mM unlabeled NE and DA or with 4 μCi/mL (0.0001 mM) Dihydroxyphenylethylamine 3,4-[Ring-2,5,6-^{3}H] (^{3}H-DA, Perkin Elmer NET673250UC) and 1.0 mM unlabeled DA or 1.0 mM unlabeled NE and DA for 10 d at 30°C. Labeled cells were collected by centrifugation at 16,000 $\times$ g for 10 min then treated with 6 M hydrochloric acid in a 90°C water bath for 1 h. Acid resistant melanin ghost material was collected by centrifugation at 16,000 $\times$ g for 10 min. DPM were measured as described above for samples taken before and after acid treatment and the percent radioactivity retained in the melanin ghost particles compared to pre-treatment samples was calculated.

Exposure to ultraviolet radiation

Equal numbers of non-melanized and 7-d melanized KN99α were plated on Sabouraud dextrose agar (Difco 210950) and then exposed to 15,000 μJ/cm^2 UV radiation (254 nm) using a Stratalinker 1800 (Stratagene, La Jolla, CA). Percent survival was calculated from CFU counts of treated compared to untreated control plates after incubation at 30°C for 3 d.

Exposure to oxygen-derived oxidants

Cultures of KN99α were grown at 30°C for 7 d in MM (control) or MM supplemented with BCM, 1.0 mM DA, 1.0 mM NE, or 1.0 mM L-Dopa. After washing with PBS, 2.5 $\times$ 10^4 cells were treated for 5 h at 37°C with an oxidative system that uses epinephrine as an electron donor, H_2O_2 as an electron acceptor, and Fe^{3+} as a transition metal catalyst (42). A fresh stock solution of 5.0 mM Fe^{3+} was prepared by mixing 10 mM ferric ammonium sulfate, $NH_4Fe(SO_4)_2$, 1:1 with 1.0 mM sulfuric acid. The oxidative treatment solution consisted of 0.5 mM Fe^{3+}, 1.0 mM H_2O_2, and 1.0 mM epinephrine bitartrate (MilliporeSigma E4375) in a buffer comprised of 50 mM sodium

acetate (pH 5.5) and 1.0 mM MgSO$_4$. Percent survival was calculated from counts of colony forming units (CFUs) for each condition compared to untreated controls.

Statistical Analysis

Biological replicates were performed for experiments as indicated and statistical significance was determined using an ordinary one-way analysis of variance (ANOVA) with Sidak's multiple comparisons test (GraphPad Prism 9 software).

Acknowledgments

We are grateful to the Hardwick lab for providing laboratory space for radioisotope experiments and to the Jacobs-Lorena lab for providing us with a UV Stratalinker 1800. This work was supported by the National Institutes of Health, grant numbers R01-AI052733 (A.C and R.E.S.)

REFERENCES

1. Springer, D. J., Mohan, R., and Heitman, J. (2017) Plants promote mating and dispersal of the human pathogenic fungus Cryptococcus. *PLoS One*. **12**, e0171695

2. Lin, X., and Heitman, J. (2006) The biology of the Cryptococcus neoformans species complex. *Annu. Rev. Microbiol.* **60**, 69–105

3. Coelho, C., Bocca, A. L., and Casadevall, A. (2014) The intracellular life of Cryptococcus neoformans. *Annu. Rev. Pathol.* **9**, 219–238

4. Rajasingham, R., Smith, R. M., Park, B. J., Jarvis, J. N., Govender, N. P., Chiller, T. M., Denning, D. W., Loyse, A., and Boulware, D. R. (2017) Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet. Infect. Dis.* **17**, 873–881

5. May, R. C., Stone, N. R. H., Wiesner, D. L., Bicanic, T., and Nielsen, K. (2016) Cryptococcus: from environmental saprophyte to global pathogen. *Nat. Rev. Microbiol.* **14**, 106–117

6. Chrétien, F., Lortholary, O., Kansau, I., Neuville, S., Gray, F., and Dromer, F. (2002) Pathogenesis of cerebral Cryptococcus neoformans infection after fungemia. *J. Infect. Dis.* **186**, 522–530

7. Chang, Y. C., Stins, M. F., McCaffery, M. J., Miller, G. F., Pare, D. R., Dam, T., Paul-Satyaseela, M., Kim, K. S., and Kwon-Chung, K. J. (2004) Cryptococcal yeast cells invade the central nervous system via transcellular penetration of the blood-brain barrier. *Infect. Immun.* **72**, 4985–4995

8. Perfect, J. R., Dismukes, W. E., Dromer, F., Goldman, D. L., Graybill, J. R., Hamill, R. J., Harrison, T. S., Larsen, R. A., Lortholary, O., Nguyen, M.-H., Pappas, P. G., Powderly, W. G., Singh, N., Sobel, J. D., and Sorrell, T. C. (2010) Clinical Practice Guidelines for the Management of Cryptococcal Disease: 2010 Update by the Infectious Diseases Society of America. *Clin. Infect. Dis.* **50**, 291–322

9. Datta, K., and Pirofski, L. (2006) Towards a vaccine for Cryptococcus neoformans: principles and caveats. *FEMS Yeast Res.* **6**, 525–536

10. Iyer, K. R., Revie, N. M., Fu, C., Robbins, N., and Cowen, L. E. (2021) Treatment strategies for cryptococcal infection: challenges, advances and future outlook. *Nat. Rev. Microbiol.* **19**, 454–466

11. Eisenman, H. C., and Casadevall, A. (2012) Synthesis and assembly of fungal melanin. *Appl. Microbiol. Biotechnol.* **93**, 931–940

12. Hill, H. Z. (1992) The function of melanin or six blind people examine an elephant. *Bioessays.* **14**, 49–56

13. Casadevall, A., Rosas, A. L., and Nosanchuk, J. D. (2000) Melanin and virulence in Cryptococcus neoformans. *Curr. Opin. Microbiol.* **3**, 354–358

14. Nosanchuk, J. D., and Casadevall, A. (2006) Impact of melanin on microbial virulence and clinical resistance to antimicrobial compounds. *Antimicrob. Agents Chemother.* **50**, 3519–3528

15. Eisenman, H. C., Frases, S., Nicola, A. M., Rodrigues, M. L., and Casadevall, A. (2009) Vesicle-associated melanization in Cryptococcus neoformans. *Microbiology.* **155**, 3860–3867

16. Eisenman, H. C., Nosanchuk, J. D., Webber, J. B. W., Emerson, R. J., Camesano, T. A., and Casadevall, A. (2005) Microstructure of cell wall-associated melanin in the human

pathogenic fungus Cryptococcus neoformans. *Biochemistry*. **44**, 3683–3693

17. Cordero, R. J. B., and Casadevall, A. (2017) Functions of fungal melanin beyond virulence. *Fungal Biol. Rev.* **31**, 99–112

18. Fu, M. S., Liporagi-Lopes, L. C., dos Santos Júnior, S. R., Tenor, J. L., Perfect, J. R., Cuomo, C. A., and Casadevall, A. (2020) Amoeba predation of Cryptococcus neoformans results in pleiotropic changes to traits associated with virulence. *bioRxiv*. 10.1101/2020.08.07.241190

19. Wang, Y., Aisen, P., and Casadevall, A. (1995) Cryptococcus neoformans melanin and virulence: mechanism of action. *Infect. Immun.* **63**, 3131–3136

20. Blasi, E., Barluzzi, R., Mazzolla, R., Tancini, B., Saleppico, S., Puliti, M., Pitzurra, L., and Bistoni, F. (1995) Role of nitric oxide and melanogenesis in the accomplishment of anticryptococcal activity by the BV-2 microglial cell line. *J. Neuroimmunol.* **58**, 111–116

21. Liu, L., Tewari, R. P., and Williamson, P. R. (1999) Laccase protects Cryptococcus neoformans from antifungal activity of alveolar macrophages. *Infect. Immun.* **67**, 6034–6039

22. Polacheck, I., Hearing, V. J., and Kwon-Chung, K. J. (1982) Biochemical studies of phenoloxidase and utilization of catecholamines in Cryptococcus neoformans. *J. Bacteriol.* **150**, 1212–1220

23. Lee, S. C., Dickson, D. W., and Casadevall, A. (1996) Pathology of cryptococcal meningoencephalitis: analysis of 27 patients with pathogenetic implications. *Hum. Pathol.* **27**, 839–847

24. Williamson, P. R. (1994) Biochemical and molecular characterization of the diphenol oxidase of Cryptococcus neoformans: identification as a laccase. *J. Bacteriol.* **176**, 656–664

25. Nosanchuk, J. D., Rudolph, J., Rosas, A. L., and Casadevall, A. (1999) Evidence That Cryptococcus neoformans Is Melanized in Pigeon Excreta: Implications for Pathogenesis. *Infect. Immun.* **67**, 5477 LP – 5479

26. Nosanchuk, J. D., Valadon, P., Feldmesser, M., and Casadevall, A. (1999) Melanization of Cryptococcus neoformans in murine infection. *Mol. Cell. Biol.* **19**, 745–750

27. Nosanchuk, J. D., Rosas, A. L., Lee, S. C., and Casadevall, A. (2000) Melanisation of Cryptococcus neoformans in human brain tissue. *Lancet*. **355**, 2049–2050

28. Chatterjee, S., Prados-Rosales, R., Frases, S., Itin, B., Casadevall, A., and Stark, R. E. (2012) Using solid-state NMR to monitor the molecular consequences of Cryptococcus neoformans melanization with different catecholamine precursors. *Biochemistry*. **51**, 6080–6088

29. Chatterjee, S., Prados-Rosales, R., Itin, B., Casadevall, A., and Stark, R. E. (2015) Solid-state NMR Reveals the Carbon-based Molecular Architecture of Cryptococcus neoformans Fungal Eumelanins in the Cell Wall. *J. Biol. Chem.* **290**, 13779–13790

30. Chatterjee, S., Prados-Rosales, R., Tan, S., Phan, V. C., Chrissian, C., Itin, B., Wang, H., Khajo, A., Magliozzo, R. S., Casadevall, A., and Stark, R. E. (2018) The melanization road more traveled by: Precursor substrate effects on melanin synthesis in cell-free and fungal cell systems. *J. Biol. Chem.* **293**, 20157–20168

31. Nosanchuk, J. D., Stark, R. E., and Casadevall, A. (2015) Fungal Melanin: What do We Know About Structure? *Front. Microbiol.* **6**, 1463

32. Rosas, A. L., Nosanchuk, J. D., Feldmesser, M., Cox, G. M., McDade, H. C., and Casadevall, A. (2000) Synthesis of polymerized melanin by Cryptococcus neoformans in infected rodents. *Infect. Immun.* **68**, 2845–2853

33. Herregodts, P., Michotte, Y., and Ebinger, G. (1989) Regional differences in the distribution of norepinephrine and epinephrine in human cerebral cortex: A neurochemical study using HPLC and electrochemical detection. *Neurosci. Lett.* **98**, 321–326

34. Herregodts, P., Ebinger, G., and Michotte, Y. (1991) Distribution of monoamines in human brain: evidence for neurochemical heterogeneity in subcortical as well as in cortical areas. *Brain Res.* **542**, 300–306

35. Janbon, G., Ormerod, K. L., Paulet, D., Byrnes, E. J. 3rd, Yadav, V., Chatterjee, G., Mullapudi, N., Hon, C.-C., Billmyre, R. B., Brunel, F., Bahn, Y.-S., Chen, W., Chen, Y., Chow, E. W. L., Coppée, J.-Y., Floyd-Averette, A., Gaillardin, C., Gerik, K. J., Goldberg, J., Gonzalez-Hilarion, S., Gujja, S., Hamlin, J. L., Hsueh, Y.-P., Ianiri, G., Jones, S., Kodira, C. D., Kozubowski, L., Lam, W., Marra, M., Mesner, L. D., Mieczkowski, P. A., Moyrand, F., Nielsen, K., Proux, C., Rossignol, T., Schein, J. E., Sun, S., Wollschlaeger, C., Wood, I. A., Zeng, Q., Neuvéglise, C., Newlon, C. S., Perfect, J. R., Lodge, J. K., Idnurm, A., Stajich, J. E., Kronstad, J. W., Sanyal, K., Heitman, J., Fraser, J. A., Cuomo,

C. A., and Dietrich, F. S. (2014) Analysis of the genome and transcriptome of Cryptococcus neoformans var. grubii reveals complex RNA expression and microevolution leading to virulence attenuation. *PLoS Genet.* **10**, e1004261

36. Jacobson, E. S., and Emery, H. S. (1991) Temperature regulation of the cryptococcal phenoloxidase. *J. Med. Vet. Mycol. bi-monthly Publ. Int. Soc. Hum. Anim. Mycol.* **29**, 121–124

37. Jacobson, E. S., and Compton, G. M. (1996) Discordant regulation of phenoloxidase and capsular polysaccharide in Cryptococcus neoformans. *J. Med. Vet. Mycol.* **34**, 289–291

38. Garcia-Rivera, J., Eisenman, H. C., Nosanchuk, J. D., Aisen, P., Zaragoza, O., Moadel, T., Dadachova, E., and Casadevall, A. (2005) Comparative analysis of Cryptococcus neoformans acid-resistant particles generated from pigmented cells grown in different laccase substrates. *Fungal Genet. Biol.* **42**, 989–998

39. Chrissian, C., Camacho, E., Fu, M. S., Prados-Rosales, R., Chatterjee, S., Cordero, R. J. B., Lodge, J. K., Casadevall, A., and Stark, R. E. (2020) Melanin deposition in two Cryptococcus species depends on cell-wall composition and flexibility. *J. Biol. Chem.* **295**, 1815–1828

40. Eisenman, H. C., Chow, S.-K., Tsé, K. K., McClelland, E. E., and Casadevall, A. (2011) The effect of L-DOPA on Cryptococcus neoformans growth and gene expression. *Virulence.* **2**, 329–336

41. Wang, Y., and Casadevall, A. (1994) Decreased susceptibility of melanized Cryptococcus neoformans to UV light. *Appl. Environ. Microbiol.* **60**, 3864–3866

42. Polacheck, I., Platt, Y., and Aronovitch, J. (1990) Catecholamines and virulence of Cryptococcus neoformans. *Infect. Immun.* **58**, 2919–2922

43. Meiser, J., Weindl, D., and Hiller, K. (2013) Complexity of dopamine metabolism. *Cell Commun. Signal.* **11**, 34

44. Flierl, M. A., Rittirsch, D., Nadeau, B. A., Chen, A. J., Sarma, J. V., Zetoune, F. S., McGuire, S. R., List, R. P., Day, D. E., Hoesel, L. M., Gao, H., Van Rooijen, N., Huber-Lang, M. S., Neubig, R. R., and Ward, P. A. (2007) Phagocyte-derived catecholamines enhance acute inflammatory injury. *Nature.* **449**, 721–725

45. Lee, S. C., Casadevall, A., and Dickson, D. W. (1996) Immunohistochemical localization of capsular polysaccharide antigen in the central nervous system cells in cryptococcal

meningoencephalitis. *Am. J. Pathol.* **148**, 1267–1274

46. Lee, S. C., and Casadevall, A. (1996) Polysaccharide antigen in brain tissue of AIDS patients with cryptococcal meningitis. *Clin. Infect. Dis.* **23**, 194–195

47. Soares, A. R., Marchiosi, R., Siqueira-Soares, R. de C., Barbosa de Lima, R., Dantas dos Santos, W., and Ferrarese-Filho, O. (2014) The role of L-DOPA in plants. *Plant Signal. Behav.* **9**, e28275

48. Folch, J., Lees, M., and Sloane Stanley, G. H. (1957) A simple method for the isolation and purification of total lipides from animal tissues. *J. Biol. Chem.* **226**, 497–509

49. Fung, B. M., Khitrin, A. K., and Ermolaev, K. (2000) An improved broadband decoupling sequence for liquid crystals and solids. *J. Magn. Reson.* **142**, 97–101

Figure 2. Melanin ghosts from KN99α melanized in pure or tissue-extracted catecholamines. A, Color photographs (left) of 2 ml samples taken from KN99α cultures grown at 30°C for the indicated number of days in minimal media supplemented with a mixture of 0.6 mM dopamine, 0.33 mM norepinephrine, and 0.07 mM epinephrine (BCM), 1 mM dopamine (DA), or 1 mM norepinephrine (NE). Quantification (right) of relative pigment levels of the three cultures showing increasing melanization over time. Brightfield images of acid-resistant melanin 'ghost' particles recovered from each culture (B) and from cells melanized on agar plates supplemented with lamb brain catecholamine extract (C). Scale bars are 5 μm.

Figure 3. Solid-state NMR spectra of melanin 'ghosts' from *C. neoformans* cells grown with DA, NE, or BCM. 1D ^{13}C cross-polarization (CPMAS) ssNMR spectra of melanin ghosts isolated from *C. neoformans* KN99α cultures grown for 10 days at 30°C in minimal media containing either 1 mM dopamine (DA, top), 1 mM norepinephrine (NE, bottom) or in a 'brain catecholamine mixture' of 0.6 mM dopamine, 0.33 mM norepinephrine, and 0.07 mM epinephrine (BCM, middle). The spectral region shaded in red (10-40 ppm) displays peaks primarily corresponding to the aliphatic carbons of lipids, the region in blue (50-110 ppm) to polysaccharide-ring carbons, and the region in yellow (110-160 ppm) to the aromatic carbons of the melanin pigment. The unshaded region (165-190 ppm) displays several overlapping peaks corresponding to the carbonyl carbons in all three constituent types.

Figure 4. Incorporation of radiolabeled precursors into melanin. A, Scatter plot graph (upper panel) of the amount of radioactivity retained in cell pellets of KN99α and ΔLAC1 cultures supplemented with 2 μCi ^{3}H-NE (0.0001 mM) in the absence or presence of the indicated concentration of unlabeled catecholamines as expressed relative to cultures with ^{3}H-NE only (n = 2). Panel below graph displays representative regions taken from color photographs of each culture. B, Radioactivity measured in acid-resistant melanin ghost particles as a percentage of the radioactivity in the samples before acid treatment. Cultures were melanized in minimal media supplemented with 0.0001 mM ^{3}H-NE or ^{3}H-DA and with 1 mM unlabeled catecholamines as indicated (n = 2).

Figure 5. Comparison of melanization conditions for resistance to cellular damage. A, Percent survival relative to untreated controls for melanized and non-melanized KN99α plated on SAB agar plates and then exposed to 15,000 µJ/cm^2 UV radiation (n = 5). B, Survival of KN99α melanized with the indicated catecholamines or a non-melanized control was determined as the number colony forming units (CFU) for cultures exposed to oxygen-derived radicals for 5 hours at 37°C as a percentage of the CFU for untreated controls (n = 3).

^{13}C DPMAS spectra

Figure S1. Quantitatively-reliable solid-state NMR spectra of melanin 'ghosts' from *C. neoformans* cells grown with DA, NE, or BCM. 1D ^{13}C direct-polarization (DPMAS) ssNMR spectra of melanin ghosts isolated from *C. neoformans* KN99α cultures grown for 10 days at 30°C in minimal media containing either 1 mM dopamine (DA, top), 1 mM norepinephrine (NE, bottom) or in a 'brain catecholamine mixture' of 0.6 mM dopamine, 0.33 mM norepinephrine, and 0.07 mM epinephrine (BCM, middle). The experiments were performed with long (50-s) recycle delays to obtain quantitatively reliable signal intensities that allowed the integration of defined spectral regions using the GNU image manipulation program (GIMP). The relative pigment content was estimated by dividing the integrated area corresponding to the aromatic carbons of the melanin pigment (110-160 ppm) by the total ^{13}C NMR signal across the spectrum.