1   **Balancing growth amidst salinity stress – lifestyle perspectives from the extremophyte model**
2   ***Schrenkiella parvula***
3

4   Kieu-Nga Tran[1†], Pramod Pantha[1†], Guannan Wang[1†], Narender Kumar[1†], Chathura
5   Wijesinghege[1], Dong-Ha Oh[1], Nick Duppen[2], Junyi He[2], Hongfei Li[3], Hyewon Hong[4], John C.
6   Johnson[1], Ross Kelt[1], Megan G. Matherne[1], Ashley Clement[1], David Tran[5], Colt Crain[1,6], Prava
7   Adhikari[1], Yanxia Zhang[3], Maryam Foroozani[1], Guido Sessa[7], John C. Larkin[1], Aaron P. Smith[1],
8   David Longstreth[1], Patrick Finnegan[8], Christa Testerink[3], Simon Barak[9], Maheshi Dassanayake[1*]
9

[1]Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

[2]Albert Katz International School for Desert Studies, Ben-Gurion University of the Negev, Sde

Boqer Campus, 8499000, Israel

[3]Laboratory of Plant Physiology, Plant Sciences Group, Wageningen University and Research,

6708PB Wageningen, The Netherlands

[4]Department of Plant Biology, University of Illinois, Urbana-Champaign, IL 61801, USA

[5]Department of Biochemistry & Department of Psychology, University of Miami, Coral Gables,

FL 33146, USA

[6]Louisiana School for Math, Science and the Arts, Natchitoches, LA 71457, USA

[7]School of Plant Sciences and Food Security, The George S. Wise Faculty of Life Sciences, Tel

Aviv University, Tel Aviv, Israel

[8]School of Biological Sciences, University of Western Australia, Perth, 6009, Australia

[9]French Associates' Institute for Agriculture and Biotechnology of Drylands, Jacob Blaustein

Institutes for Desert Research, Ben-Gurion University of the Negev, Sde Boqer Campus,

8499000, Israel

[†]Equal contribution

*Address correspondence to: maheshid@lsu.edu

Keywords: Extremophyte, abiotic stress adaptations, tradeoffs in growth and stress tolerance,

genomics, *Schrenkiella parvula*, salt stress

One sentence summary: *Schrenkiella parvula* salt-resilient growth is facilitated by

uncompromised primary root growth, expansion of xylem vessels, maintenance of leaf water

status and photosynthesis, and early flowering.

Abstract

Schrenkiella parvula, a leading extremophyte model in Brassicaceae, can grow and complete its

life cycle under multiple environmental stresses, including high salinity. While foundational

genomic resources have been created for *S. parvula*, a comprehensive physiological or

structural characterization of its salt stress responses is absent. We aimed to identify the

influential traits that lead to stress-resilient growth of this species. We examined salt-induced

changes in the physiology and anatomy of *S. parvula* throughout its lifecycle across multiple

tissues. We found that *S. parvula* maintains or even exhibits enhanced growth during various

developmental stages at salt stress levels known to inhibit growth in Arabidopsis and most

crops. The resilient growth of *S. parvula* was associated with key traits that synergistically allow

continued primary root growth, expansion of xylem vessels across the root-shoot continuum,

and a high capacity to maintain tissue water levels by developing larger and thicker leaves while

facilitating continued photosynthesis during salt stress. These traits at the vegetative phase

were followed by a successful transition to the reproductive phase via early flowering,

development of larger siliques, and production of viable seeds during salt stress. Additionally,

the success of self-fertilization during early flowering stages was dependent on salt-induced

filament elongation in flowers that aborted in the absence of salt. Our results suggest that the

maintenance of leaf water status and enhancement of selfing in early flowers to ensure

reproductive success, are among the most influential traits that contribute to the extremophyte

lifestyle of *S. parvula* in its natural habitat.

Introduction

Soil characteristics, water availability, and light and temperature regimes set boundaries

for plant growth and dictate how plants complete their life cycles. Balancing plant growth with

environmental stress responses is a constant challenge faced by all plants. This balance is

epitomized by extremophytes, which are plants adapted to thrive in extreme environmental

conditions than mesophytes. Therefore, extremophytes provide insights on evolutionarily-

tested lifestyle strategies that have proven successful in extreme environments (Lloyd and

Oreskes, 2018; Rodell et al., 2018; Schlenker and Auffhammer, 2018; Solis et al., 2020).

Knowledge transfer from extremophyte-focused research to design innovative crops that can

show resilient growth under stress can be a fruitful strategy when envisioning global food

security and expansion of agricultural lands to marginal lands (Oh et al., 2012; Bechtold, 2018;

Kazachkova et al., 2018; Zandalinas et al., 2021). The recent genome explorations in a wider

range of plants makes this an ideal time for mining representative extremophyte traits.

Salt tolerance is a complex trait that requires integration of physiological, anatomical,

and metabolic responses (Cheeseman, 2013; Barros et al., 2021). These multifaceted traits

cannot be achieved in stress-sensitive crops simply through the acclamatory response of a few

genes (Roy et al., 2014). The complexity of these interactions is implied by the fact that

breeding has had limited success in conferring salinity tolerance despite numerous attempts

(Yamaguchi and Blumwald, 2005; Cheeseman, 2013; Roy et al., 2014; Barros et al., 2021). An

understanding of the extent of tradeoffs between plant growth and stress tolerance is required

to assess how selected traits, genes, or pathways affect short- or long-term plant growth and

fitness in extremophytes. Such foundational studies are lagging for all the leading

extremophyte models compared to the breadth of tools and molecular resources available to

identify candidate genes in these species.

Schrenkiella parvula (Schrenk) D.A.German & Al-Shehbaz (previously known as

Thellungiella parvula or *Eutrema parvulum*) is a leading extremophyte model in the

Brassicaceae family (Zhu, 2015; Ali and Yun, 2017; Kazachkova et al., 2018; Krämer, 2018). *S.*

parvula shares many traits with the model plant, *A. thaliana*, that make it an excellent model,

including comparable genome size (Dassanayake et al., 2011), similar lifecycle duration, self-

pollination, and prolific seed production (Table 1 and Figure 1). In the ten years following the

release of its genome, *S. parvula* has been developed as a model system equipped with primary

molecular tools to explore genetic mechanisms underlying plant abiotic stress tolerance

(Dassanayake et al., 2011; Oh et al., 2014; Wang et al., 2018; Pantha et al., 2021; Wang et al.,

2021). *Schrenkiella parvula* is remarkably tolerant to high salinity compared to closely related

Brassicaceae species (Orsini et al., 2010) and is categorized as a halophyte based on its capacity

to complete its life cycle when grown at 200 mM NaCl or higher (Flowers and Colmer, 2008). In

addition to high sodium, *S. parvula* is also uniquely adapted to cope with multiple edaphic

factors found in its native habitat, including boron, potassium, and lithium, among other salts,

that exist at levels toxic for most crops (Helvaci et al., 2004; Oh et al., 2014; Hajiboland et al.,

2018; Tug et al., 2019). Its preferential distribution near saline lakes in the Irano-Turanian

region makes it a robust extremophyte model to investigate multiple environmental stresses

that are often found as compound stresses in many marginal agricultural landscapes.

Despite a growing body of genomic resources for *S. parvula* (Jarvis et al., 2014; Ali et al.,

2018; Wang et al., 2021), a systematic study that describes its life history, and the physiological,

and structural features that are associated with adaptations to environmental stresses,

especially salt stress, is absent. In this study, we evaluated *S. parvula* stress adaptations at both

the structural and physiological levels. We assessed adaptations that are likely to be influential

in balancing growth with stress tolerance and identified inducible adaptive traits that were the

most prominent features in a multivariate space across developmental stages, tissue types, and

response types. We propose these inducible traits to be hallmarks of a stress-adapted lifestyle

for a fast-growing annual in saline soils demonstrating the utility of *S. parvula* as an

extremophyte model plant.

Results

***Schrenkiella parvula* adjusts root growth, structure, and form under high salinity**

Schrenkiella parvula roots, in contrast to those of *A. thaliana*, maintained uninterrupted

root growth in response to salt stress compared to control conditions (Figure 2A). Remarkably,

S. parvula primary root length did not display any growth retardation even after longer

durations or higher salt concentrations tested, while *A. thaliana* showed growth inhibition at

100 mM NaCl (Figure 2B). The same trend was observed for lateral root growth, although the

emergence of lateral roots was delayed in *S. parvula* even at control conditions compared to *A.*

thaliana (Figure 2C). The higher tolerance to salinity shown by *S. parvula* roots compared to *A.*

thaliana roots was consistent when plants were grown under a longer photoperiod and at

higher nutrient levels (Figure S1). Under these conditions, a 125 mM NaCl treatment caused a

growth stimulating effect on *S. parvula* primary root length, but inhibited root growth in *A.*

thaliana (Figure S1B). Stronger inhibition of both average lateral root length and number was

imposed by 175 and 225 mM NaCl, which could explain the reduced total lateral root length

(Figure S1C to E). Taken together, salt stress imposed by 125 mM NaCl promoted primary root

growth and average lateral root length, while a higher salt concentration of 175 mM NaCl

imposed less inhibition on main root growth and lateral root density in *S. parvula* compared to

A. thaliana. Compared to primary and lateral roots, root hair development in *S. parvula* was

sensitive to high salinity; root hairs showed a consistent decrease in length at higher salinities

or during longer exposure times to salt stress (Figures 2D and S2). The ability of *S. parvula* to

maintain or enhance primary root growth under high salinity remained as a consistent trait

shown by mature plants over longer durations and higher salinities tested using 250 mM NaCl

in a hydroponic growth medium (Figure S3).

Arabidopsis thaliana roots showed halotropism by growing away from salt (Figure S4)

(Galvan-Ampudia et al., 2013). Interestingly, under comparable growth conditions, *S. parvula*

showed salt-insensitive primary root growth where it grew towards higher salt concentration

without changing its course, further exemplifying the higher tolerance to high salinity (Figure

S4).

To investigate the potential effect of salt stress on root anatomy, we examined the

tissue level structural responses in roots of 8-week-old *S. parvula* plants that had been treated

for 4 weeks with 150 mM NaCl (Figure 3A). These traits were catalogued from root tips to

mature roots (Figure S5). In young roots, the length of the root tip (measured as the distance

from the tip to the emergence site of the first root hair) was longer under high salinity

conditions compared to control conditions (Figure 3B). However, we did not observe any other

anatomical trait adjustments to salinity in young roots of *S. parvula* (Figure 3C). Interestingly, *S.*

parvula roots develop an extra cell layer between the cortex and endodermis compared to *A.*

thaliana (Brady et al., 2007), formed early on in its root development in both control and salt-

treated samples (Figure S5A and C). In response to high salinity, xylem tissues in mature roots

significantly increased in area under high salinity (Figure 3D). However, this did not affect the

overall mature root area. The xylem tissue expansion was primarily caused by increase in the

area of vessels. The extra space taken up by the xylem tissue was compensated for by reduction

of the cortical air spaces (unstructured aerenchyma) in mature roots (Figure 3D).

***Schrenkiella parvula* shoots change in form and function to response to salt stress**

Structural adjustments in response to long-term salt stress extended to shoot tissues in

S. parvula (Figures 4 and S6). Long-term salt stress leads to larger vessels in *S. parvula* shoots

(Figure 4A) to maintain the root-shoot continuum with root vessel elements that increased in

size when exposed to high salinity (Figure 3D). The expansion in vessel element size in shoots

was matched by a reduction in cambial and cortical zones that kept the average shoot area

constant between control and salt-treated plants (Figure 4A), while other cell and tissue layers

in the shoot remained unchanged (Figure S6B).

Leaves that developed during long-term salt treatments in *S. parvula* exhibited

increased succulence, as indicated by increase in leaf thickness across the leaf including the

midrib (Figures 4B and S7). Increased endoreplication induced by salinity can cause cell size

expansion and succulence in the extremophyte, *Mesembryanthemum crystallinum* (Barkla et

al., 2018). We therefore tested whether the increased succulence in *S. parvula* leaves in

response to salt coincided with endoreplication by quantifying ploidy of leaf cells. Indeed, the

proportion of leaf cells with higher ploidy increased with increased salinity, causing a significant

rise in the endoreplication index under 250 mM NaCl compared to the control condition (Figure

4C). *Schrenkiella parvula* leaves showed both structural and functional adjustments to salt

treatments that correlated with its capacity to maintain relative water content (Figure 5), but

the total leaf number per plant remained similar between control and salt-treated plants

(Figure 5A). Long-term salt treatments not only increased leaf succulence, but also leaf area

(Figure 5A), suggesting a growth-promoting effect at least with the long-term 150 mM NaCl

treatment. *Schrenkiella parvula* was photosynthetically active and maintained growth during

long-term salt treatments at high salinities (Figure 4), but there was a tradeoff in increasing salt

on shoot development at salinities surpassing 150 mM (Figure S8).

Leaf shape, angle and phyllotaxis, stomatal distribution, and boundary layer thickness

caused by thick cuticles or trichomes largely contribute to efficient use of water by plants (Yoo

et al., 2009). *Schrenkiella parvula* leaves are amphistomatous (i.e. stomata exist on both abaxial

and adaxial surfaces), linear-lanceolate with a narrower base, and trichomeless (Table 1, Figure

5A and 5B). In *S. parvula*, stomatal density did not change in fully-mature leaves (Figure 5B).

However, stomatal regulation was adjusted with increasing salinities as indicated by decreased

stomatal conductance (Figure 5C) and decreased CO_2 assimilation, notably at much higher

salinities for *S. parvula* than observed for *A. thaliana* (Figure 5D). While 50 mM salt treatment

was sufficient to cause a decrease in stomatal conductance and photosynthesis in *A. thaliana*, *S.*

parvula treated with 100 mM NaCl was able to maintain stomatal conductance and CO_2

assimilation at levels indistinguishable from those measured in plants grown under control

conditions. Concomitantly, *S. parvula* maintained lower average leaf temperatures than *A.*

thaliana (Figure 5E). Remarkably, *S. parvula* maintained relative leaf water content at all tested

salinities up to 250 mM NaCl (Figure 5F).

Utilizing the Qubit/PSI PlantScreen^{TM} Compact high-throughput phenomics system, we

undertook a comparative analysis of *A. thaliana* and *S. parvula* shoot morphometrics (RGB

digital camera), leaf water content (infra-red camera), and photochemistry (PAM chlorophyll

fluorescence). Figure S9A demonstrates that the phenomics system was able to simultaneously

capture morphological characteristics specific to each species. *A. thaliana* has a rosette

structure (i.e. less protrusions from a perfect circle), and therefore exhibited higher

compactness). In contrast *S. parvula* leaves on a spiral phyllotaxy (Figure 1 and Table 1) that

increasingly protrude as the plant grows, reflected a lower compactness in older plants

regardless of salt concentrations. Salt stress led to a statistically significant reduction in *A.*

thaliana leaf area but had no effect on *S. parvula* leaves (Figure S9B). *S. parvula* was also able to

maintain leaf water content and maximum quantum efficiency of PSII under salt stress (Figure

S9C and D, respectively) whereas *A. thaliana* exhibited a salt stress-mediated reduction in both

parameters. Non-Photochemical Quenching (NPQ), a mechanism for dissipating excess light

energy (Müller et al., 2001), increased in *A. thaliana* under salt stress suggesting that less

energy is being utilized for productive purposes (Figure S9E). On the other hand, salt stress had

no effect on NPQ in *S. parvula*. The resilience of *S. parvula* photochemistry under salt stress is

consistent with this species' ability to maintain CO_2 assimilation in saline conditions (Figure 5D).

Taken together, the phenomics data confirm similar growth and physiological tolerance of *S.*

parvula to salt stress regardless of growth medium.

Minimizing non-stomatal transpiration by enhanced epidermal boundary layer

resistance is equally important as stomatal regulation in preventing water loss under water

stress (Blum, 2009). Indeed, previous findings have demonstrated that *S. parvula* leaves possess

a significantly thicker leaf cuticle supported by a higher level of total wax content compared to

A. thaliana leaves (Teusink et al., 2002) (Figure S10A). Interestingly, when we compared the

basal expression of key genes in the wax biosynthesis pathway (Bernard and Joubès, 2013) from

comparable shoot tissues grown under control conditions, we observed consistently higher

constitutive expression in *S. parvula* compared to expression of their *A. thaliana* orthologs

(Figure S10B).

Changes to leaf shape and arrangement is not a salt-dependent, inducible trait in *S.*

parvula. However, its narrow leaves with elongated petioles arranged spirally on erect stems

with elongated internodes (Figure S11A) are ideally suited to rapid growth in saline habitats

with warm temperatures where efficient transpirational cooling imparts selective advantages

(Lin et al., 2017). Additionally, an erect growth habit is more desirable than a rosette to

minimize leaf contact with soil, given that *S. parvula* is found near saline lakes with topsoils

often enriched in salt crystals (Hajiboland et al., 2018; Tug et al., 2019). Key traits resulting in

narrow leaves, elongated petioles and internodes are influenced by the phytochrome family of

genes (Somers et al., 1991; Li et al., 2011), and we noted that *S. parvula* plants closely resemble

the morphology of *A. thaliana phyB phyD* double mutants (Devlin et al., 1996; Li et al., 2011).

This morphological similarity led us to search for genome-level cues for loss-of-function or

altered function in *S. parvula PHY* genes that may support such a phenotype being selected as

the only documented growth form of *S. parvula* (Figure 1 and Table 1) (German and Al-Shehbaz,

2010). We observed that *PHYB* was conserved as a single copy gene in six closely-related

Brassicaceae genomes that we examined, with a translocation event between lineage I and II

species. However, *PHYD* appeared to have undergone a gene loss specifically in *S. parvula*

(Figure S11B).

Salt stress induces early flowering and silique formation in *Schrenkiella parvula*

We examined long-term salt-stress effects on reproductive traits to investigate how

excess Na$^+$ affected *S. parvula* fitness (Figure 6). Salt treatment induced early flowering

irrespective of whether flowering time was measured as the number of days after planting to

detection of the first floral buds or the number of leaves developed before flowering (Figure 6A

to C). Additionally, early flowering induced by salt was observed under both long-day and short-

day photoperiods (Figure 6C). The initial ~10 flowers produced under control conditions

aborted without developing into siliques, while these flowers produced mature siliques in salt-

treated plants (Figure 6A and B). *Schrenkiella parvula* flowers are self-fertilizing, a process

facilitated by the elongation of filaments to the level of the stigma. However, early flowers

under control conditions developed shorter filaments than in the salt-treated plants (Figure

6D). In control plants, these shorter filaments appeared to hinder successful fertilization and

delayed subsequent silique formation (Figure 6A and E). Notably, the salt-induced early

flowering and salt-dependent elongation of filaments were not specific to NaCl treatment and

were also initiated by KCl at similar concentrations (Figure S12). At later reproductive stages,

the continuous growth of multiple floral meristems led to fertile flowers under control

conditions. Despite the early success of silique formation in salt-treated plants, the cumulative

number of flowers and siliques produced by control plants surpassed that of salt-treated plants

if allowed to grow past two months. For example, at 74 days after planting, the control plants

had a greater number of flowers per plant than salt-treated plants (Figure 6E). The experiment

was discontinued when salt-treated plants stopped producing new inflorescences and were

senescing (approximately 74 days after planting). Therefore, *S. parvula* does not require salt to

increase fitness, and salt treatments appear to only provide a transient advantage during early

reproductive stages. Even if fitness is higher for control plants compared to salt-treated plants

when evaluated based solely on the total number of siliques produced per plant within a

longest possible growing season, there may be additional benefits when seeds develop on

plants exposed to long-term high salinity, especially if seasonal changes in its natural habitats

reward a strategy of salt-accelerated reproduction. The average size of siliques and seeds of

salt-treated plants were significantly greater than in control plants (Figure 6G, H, and I). This

suggested that more resources were available for seeds when produced by salt-treated plants

compared to control plants. The number of seeds per silique was not different between control

and treated plants and the seeds from both groups were indistinguishable in the rate of

germination in the subsequent generation when germinated on media without added NaCl.

***Schrenkiella parvula* seed germination is delayed by high salinity and is sensitive to specific**

salts

Many plants including salt-adapted plants delay germination in saline media even if

their seedling stages can tolerate high salinity (Ungar, 1978; Kazachkova et al., 2016). Both *S.*

parvula and *A. thaliana* exhibited decreased seed germination as NaCl concentrations increased

compared to control conditions (Figure 7A). However, unlike the more salt-tolerant traits *S.*

parvula displayed compared to *A. thaliana* during seedling and mature developmental stages

(Figures 2, 4, and S9), *S. parvula* seed germination was more sensitive to NaCl than *A. thaliana*

seed germination (Figure 7A). When we sowed *S. parvula* seeds directly onto salt plates prior to

stratification, we observed more severe salt-mediated inhibition of germination. This was

demonstrated via detailed germination curves that revealed a progressive reduction in both

germination rate and final percent germination with increasing NaCl concentrations (Figure 7B).

This salt-influenced trait did not lead to a binary outcome between germinated vs

ungerminated seeds. The smaller fraction of *S. parvula* seeds that formed a radicle compared to

A. thaliana continued to develop in a high salinity medium similarly to those seeds germinated

under control conditions (Figure S13A). In contrast, *A. thaliana* seeds that formed radicles in a

high saline medium did not survive to the seedling stage. Up to 98% of *S. parvula* seeds that

failed to germinate (i.e. no visible radicle emergence) on high-salt growth medium germinated

and developed into seedlings when transferred to control medium without added NaCl (Figures

7C and S13B). Moreover, seeds that were incubated for a month on saline medium still

exhibited 98% germination (Figure 7C). Interestingly, adding salt and increasing the duration of

salt-treatment, led to faster germination when salt-inhibited seeds were transferred back to the control medium. These results suggest that NaCl introduces a reversible barrier to *S. parvula* seed germination.

We also tested whether other salts, including K, Li, and B, found in the native soils of *S. parvula* ecotype Lake Tuz caused a reversible inhibition of *S. parvula* seed germination. KCl induced a similar inhibitory response as observed for NaCl (Figures 7D and S13), but neither LiCl nor H_3BO_3 led to inhibition of *S. parvula* seed germination (Figure 7E and F). In contrast, *A. thaliana* seed germination was highly sensitive to high LiCl and H_3BO_3, and KCl at 150 or 250 mM was lethal to *A. thaliana* seeds that formed a radicle (Figures 7D, E, F, and S13A). These data suggest that *S. parvula* is adapted to sense the soil salt level as well as the salt type before radicle emergence, and if seeds germinate, they are more likely to continue development despite toxic levels of salts in the growth medium that are lethal to *A. thaliana*.

Salt-dependent germination of *S. parvula* suggests that it is an adaptive trait to time seed germination for when topsoil salinity decreases during the summer months from June to September. This finding was supported by field observations made in the Lake Tuz region where *S. parvula* grows as a seasonal annual during the summer months that receive the highest rainfall and temperature with the longest day length (Figures S14A to C). The saline soils surrounding Lake Tuz are often covered with a salt crust towards fall, which becomes a dominant feature in the landscape extending to spring of the following year despite having a high saline water table throughout the year (Tug et al., 2019). Seed germination may range widely within the expected growing season and seedlings are more vulnerable to extreme environmental conditions than mature plants. Finally, we compared *S. parvula* and *A. thaliana* seedlings for their tolerance to heat, chilling, and freezing stresses to assess how *S. parvula* may fit within the expected temperature tolerance range required to survive in its native habitat (Figures S14D to F). *Schrenkiella parvula* seedlings displayed higher tolerance to heat stress at 38 °C than *A. thaliana* (Figure S14F), while chilling and freezing tolerance in both species was indistinguishable (Figure S14D and E).

Discussion

Plants are known for their remarkable capacity to tolerate environmental stresses and

ability to modulate growth even to the extent of pausing growth entirely. This plasticity in

response to various environmental stresses is observed in all plants, but as shown in the current

study, extremophytes such as *S. parvula* exhibit a much higher tolerance to stress than

mesophytes, which include most crops and *A. thaliana* (Flowers and Colmer, 2008; Kazachkova

et al., 2018). Halophytes represent only about 0.4% of all flowering plants and about 40% of all

halophytes including *S. parvula* can withstand salt stress at concentrations similar to seawater

(Kotula et al., 2020). At high salinities, most plants respond to salt stress by inhibiting growth to

prioritize survival as a tradeoff (Santiago-Rosario et al., 2021). Nevertheless, the extremophyte

model *S. parvula* provides a genetic system to discover adaptive traits that do not show a

growth compromise at salt concentrations that are known to be lethal to most crops and *A.*

thaliana (Oh et al., 2014). The defining traits of *S. parvula* as an extremophyte should reflect

the environmental constraints that have shaped its phenotype and collectively made it distinct

from other more stress-sensitive annuals in Brassicaceae, including *A. thaliana*. When we

summarized *S. parvula* physiological and structural traits that changed at a developmental

stage-specific and tissue-specific scale in response to salt stress, several traits emerged as

significant features induced under salt stress compared to their non-stressed control group

(Figure 8A). Among them, the higher number of siliques produced during salt-induced early

flowering, expansion of xylem vessel elements, and increased leaf thickness provide the

greatest contribution to the total plastic or inducible trait space (Figure 8B and C).

Salt stress accelerates reproduction thereby contributing to maximizing fitness of *S. parvula*

Salinity delays flowering in most plants (Blits and Gallagher, 1991; Lutts et al., 1995;

Apse et al., 1999; Moriuchi et al., 2016; Cho et al., 2017). For example, high salinity (≥100 mM

NaCl) delays or inhibits the transition from vegetative to reproductive growth in *A. thaliana*

(Achard et al., 2006; Li et al., 2007; Ryu et al., 2014). Delayed flowering ensures survival under

salt stress supported by multiple genetic mechanisms. These pathways include transcription

factors that inhibit flowering and are also induced by salt (Cho et al., 2017). However, a few

plants possess alternative strategies whereby flowering is accelerated in response to salt stress

(Adams et al., 1998; Ventura et al., 2014). Additionally, molecular pathways operating

antagonistically to salt-induced delayed flowering have been described for *A. thaliana* (Yu et al.,

2018). This apparent strategy would facilitate reproduction and improve fitness by allowing

escape from harsh environments, especially for ruderal or annual plants growing in habitats

that become warmer, drier, and more saline (due to increased surface evaporation) towards

the end of their growing season. Salt-induced flowering observed for *S. parvula* suggests that

such an alternative strategy was selected as the dominant trait to maximize its fitness in its

native habitat exemplified by the Lake Tuz ecotype used in our current study (Figures 6 and

S12) over the more common trait of salt-induced delay in flowering observed for many other

plants (Apse et al., 1999; Cho et al., 2017).

Compared to favorable growth conditions, environmental stresses increase the

opportunity for selection. Previous studies have demonstrated that phenotypic selection favors

stress-avoidance traits, including earlier flowering, in addition to stress-tolerance traits (Eshel et

al., 2021). This results in an evolutionary shift toward earlier flowering despite the prevalent

trait observed for model and crop plants to delay flowering when exposed to salt (Stanton et

al., 2000; Ventura et al., 2014; Caño et al., 2016; Moriuchi et al., 2016). Stanton et al., (2000)

further showed that, among multiple environmental stresses tested for their effects on

evolutionary selection in wild mustard, *Sinapis arvensis* (a ruderal annual in Brassicaceae), the

potential for selection was greatest for high salt stress and low light. *Schrenkiella parvula* is

native to the variable shoreline habitats of saline lakes of the Irano-Turanian region where it is

often a ruderal plant growing in frequently inundated saline soils (Ozfidan-Konakci et al., 2015;

Hajiboland et al., 2018; Tug et al., 2019). Therefore, salt stress is the norm for *S. parvula* and

over a multi-generational time scale, salt stress is expected to provide a constant selection

pressure.

When we examined salt-responsive traits throughout the lifecycle of *S. parvula*, an

increased number of siliques produced due to salt-induced early flowering had the largest fold-

change among other salt-induced traits that differentiated the response between control and

salt-treated growth (Figure 8C). Early flowering established an earlier seed set and the

opportunity to produce viable seeds at an earlier developmental age in the salt treated plants

compared to control plants (Figure 6). However, this was a plastic trait not necessarily

imparting higher fitness when assessed based on overall fecundity during a longer growth

period . When control and salt-treated plants were grown for over 2 months, the control plants

displayed longer vegetative growth phases subsequently leading to higher fecundity in *S.*

parvula. These results suggest that salt-induced flowering has evolved as a mechanism of stress

avoidance rather than stress tolerance in a habitat where the severity of environmental

stresses increase as the growing season continues (Figure S14).

A key morphological feature contributing to the transient advantage of salt-mediated

early flowering in *S. parvula,* was the plastic response observed for filament elongation that

was enhanced by salt (Figures 6D and S12). Filament elongation is an important trait for

successful fertilization in self-pollinating plants such as *S. parvula* and *A. thaliana*. However,

whereas salt stress terminally inhibits *A. thaliana* gamete formation and leads to abortion of

flowers and seeds (Sun et al., 2005; Figure 6), filament elongation and production of viable

seeds were not inhibited by salt in *S. parvula*. This finding indicates that *S. parvula* has evolved

to decouple the regulatory pathways that sense salt and inhibit stamen growth.

Salt stress-induced xylem vessel expansion across the root-shoot continuum could contribute

to the salt resilient growth of *Schrenkiella parvula*

Selection on vessel diameter favors narrower vessels to minimize the risk of cavitation

independently of ancestry or habitat for a wide range of angiosperms from rainforests to

deserts (Olson and Rosell, 2013). Furthermore, the effect of salt stress on xylem development

within a species often leads to narrower vessels (Junghans et al., 2006; Ge et al., 2017; Cruz et

al., 2019; Sarker and Oba, 2020). Our results suggest an alternative strategy for *S. parvula*

where we observed an increase in both shoot and root xylem vessel area in plants exposed to

long-term salt stress (given via 150 mM NaCl) (Figures 3 and 4) (Li et al., 2021). The salt-

mediated expansion of xylem tissue in *S. parvula* was the second-most highly induced trait that

responded by a higher fold change between control and salt-treated plants (Figure 8). This

adaptive feature likely allows a higher bulk flow through the xylem to help *S. parvula* leaves

maintain a cooler temperature while allowing uncompromised gas exchange compared to *A.*

thaliana grown under saline conditions (Figure 5), despite the tradeoff of a higher risk of

cavitation. Salt-induced increases to xylem diameter correlated to maintaining growth under

salt stress has been reported for a few extremophytes (e.g. *Nitraria retusa* and *Atriplex halimus*)

adapted to extreme salinities above seawater strength, but is not a common trait associated

with halophytes (Boughalleb et al., 2009; Parida et al., 2016). The expansion of xylem area

without the expansion of root or stem area that we observed as a salt responsive trait in *S.*

parvula (Figure 3) is an atypical adaptation among plants resilient to salt stress (Olson and

Rosell, 2013; Nassar et al., 2020). *S. parvula* seems to optimize effective use of water instead of

simply reducing water loss at high salinities (Figures 5F and S9). Prioritizing effective water use

has been proposed to be a better strategy for plant growth during water deficit stress than

maximizing water conservation at the expense of photosynthetic capacity (Blum, 2009). This

finding has implications for agriculture. If the availability of water can be ensured in agricultural

systems even if the water source is brackish, the crops that are able to maintain relative water

content (as observed for *S. parvula* in Figure 5), while allowing uncompromised transpiration

and gas exchange, will be more resilient to environmental stresses.

Adjustments to shoot architecture modulation follows root responses to cope with salt stress

Schrenkiella parvula root growth and development was less affected by increasing salt

concentrations than in *A. thaliana*. *Schrenkiella parvula* primary root length increased while

root fresh weight was only maintained under high salt conditions (Figures 2, S1 and S3),

suggesting that salt causes a reallocation of resources toward deeper roots. This is also

supported by the reduction in root hair length, total lateral root length and number (Figures 2

and S1). Given the natural environment in which the *S. parvula* ecotype Lake Tuz grows, the top

layers of soil will dry first, causing the soil salt concentration to form a high to low gradient from

topsoil to deeper layers in the rhizosphere (Tug et al., 2019). Thus, *S. parvula* can avoid

relatively higher salinities by growing deeper to lower salt concentrations. The growth strategy

exemplified by *S. parvula* is correlated with uncompromised growth dependent on modulating

root to shoot vasculature and seemed to be efficiently coupled with leaf traits in *S. parvula*.

Even though saline soils directly affect roots, its effects are also observed in shoots. If roots

grow continuously in saline media, salts are often deposited in leaves. As plants age, salt-

adapted plants either need to store salts in leaves and shed those whose capacity to hold

excess salt is reached, or extrude excess salt via salt glands (Jennings, 1968; Dassanayake and

Larkin, 2017). *Schrenkiella parvula* uses the first strategy by developing succulent leaves that

are equipped with larger vacuoles which support salt sequestration. This trait appears to be

facilitated by shifting the leaf cell population from a dominant diploid state to higher ploidy

levels during prolonged salt stress (Figure 4). Indeed, the increase in leaf thickness, along with

larger leaf area, was among the top traits with highest salt-induced fold-changes in *S. parvula*

(Figure 8). Developing larger cells that can store excess salts within larger vacuoles enabled by

endoreduplication has been reported for several other salt adapted extremophytes. It was

proposed to be a key mechanism for surviving environmental stress (De Rocher et al., 1990;

Barkla et al., 2018), while leaf succulence is one of the most common traits observed in

halophytes (Jennings, 1968; Flowers and Colmer, 2008).

To maximize storage capacity via increasing the leaf surface area to volume, leaves are

generally more terete in many extremophytes that have fully developed succulent leaves. This

trait is often accompanied by the loss of leaf abaxial identity whereby the leaves become

adaxialized (Ogburn and Edwards, 2013). Multiple genetic networks regulated by adaxially

expressed HD-ZIPIII transcription factors have been shown to control leaf polarity (Du et al.,

2018). *Schrenkiella parvula* leaves tend to acquire a more terete state with prolonged exposure

to salt (Figure S7B). This phenotypic change is coincident with *S. parvula* leaves exhibiting

reduced differentiation between adaxial and abaxial surfaces as leaf thickness increases with

higher salt concentrations and a clear palisade layer at the adaxial surface is not present.

However, this structural alteration does not affect the relative water content nor cause leaf

curling or other visibly malformed vascular structures, as seen in adaxialized leaves of *A.*

thaliana mutants (Figures 4, 5, and S7) (Du et al., 2018).

Schrenkiella parvula leaves are amphistomatous (possess stomata on the upper and

lower leaf surfaces), in contrast to the *A. thaliana* leaves, which have more stomata on the

abaxial surface (hyposomatous) (Figure 5B). Amphistomatous leaves are relatively rare in

angiosperms (Nadeau and Sack, 2002; Drake et al., 2019) but in such leaves, the distance

between stomata and the mesophyll is reduced allowing greater water-use efficiency due to

increased CO_2 conductance in the mesophyll thereby facilitating a higher relative

photosynthesis rate (de Boer et al., 2016). Therefore, amphistomatous leaves are found in

highly productive fast-growing herbs. However, when this trait is found in extremophytes such

as *S. parvula* with trichomeless leaves arranged mostly with vertical leaf angles (Fig 1), it adds a

tradeoff for water loss through transpiration as well as higher exposure to solar heat (Drake et

al., 2019). Drake et al. (2019) hypothesized that to avoid desiccation in thick amphistomatous

leaves, vein length per unit leaf area must be larger than in hypostomatous leaves of the same

thickness. This anatomy would need to be accompanied by additional traits to facilitate an

increase in transpirational flow via xylem development from root to shoot if leaf thickness

increases as a response to salt stress. Additionally, having mostly vertical leaf angles with thick

amphistomatous leaves is considered a key adaptive trait to reduce the effect of thermal

radiation around midday while maximizing photosynthesis in the morning and late afternoon

thereby lowering the risks of desiccation (King, 1997). Further, traits known to have evolved

under shade avoidance such as internode elongation seems to be exapted in *S. parvula*, most

likely to enhance transpirational cooling selected under environments that demand rapid

growth amidst extreme environmental constraints (Pierik and Testerink, 2014).

Moderate to high salt is a persistent edaphic factor in the natural habitat of *S. parvula*

(Helvaci et al., 2004; Tug et al., 2019). Such an environment would have selected for multiple

traits that act synergistically to create an efficient lifestyle to survive stress. However, unlike

many slow-growing extremophytes, *S. parvula* shows fast growth achieved with short life

cycles, while enduring multiple environmental stresses. Often environmental stress resilience

comes at the cost of yield reduction in crops and the drive for increasing yields has been

prioritized over the need to develop resilient crops (Pardo and VanBuren, 2021). Increasing

threats to global agriculture due to climate change necessitate a change in our priorities for

crop development (IPCC, 2021). *Schrenkiella parvula* provides an excellent genetic model

system that illustrates growth optimization over growth inhibition to cope with environmental

stresses that we can explore to find innovative genetic architectures suitable and transferable

to develop resilient crops.

Materials and Methods

Plant growth conditions

Schrenkiella parvula (Lake Tuz ecotype) and *A. thaliana* (Col-0 ecotype) were grown on

soil, plates or in a hydroponic system as described in Wang et al., (2019), Pantha et al.,(2021),

and Conn et al., (2013). Soil grown plants were used for reproductive trait assessments. Plate

grown plants were used for root growth assessments and thermal tolerance assays.

Hydroponically grown plants were used for shoot physiological assessments and anatomical

characterization. See supplementary method S1 for details.

Root growth assays

Five-day-old seedlings germinated on plates were transferred to 1/4x MS supplemented

with 100 and 150 mM NaCl and kept at a photoperiod of 12 hr light/12 hr dark for 13 days.

Additionally, 4-day-old seedlings of *A. thaliana* and *S. parvula* grown at a photoperiod of 16 hr

light/8hr dark and on 1/2x MS agar plates were transferred to 1/2x MS plates supplemented

with 125, 175, and 225 mM NaCl and monitored for 6 days. These plates were imaged for

further processing using ImageJ (Ferreira and Rasband, 2012) to quantify primary root length,

lateral root number and density, total and average lateral root length, and root hair length

(Figures 2 and S2). See supplementary method S1 for details.

Halotropism assay

Seedlings were grown on 1/2x MS medium solidified with 1.2% (w/v) Phyto agar plates

held at an 85° incline as described in Galvan-Ampudia et al., (2013). After 5 days, the agar

medium 1 cm below the root tips, was removed and replaced by 1/2x MS agar supplemented

with 200 mM NaCl and seedlings were allowed to grow for another 5 days (Figure S4).

Germination assay

Sterilized stratified seeds were germinated on 1/4x MS (Figures 7A, D, E and F) or 1/2x

MS (Figures 7B and C) supplemented with different concentrations of NaCl, KCl, LiCl, or H_3BO_3.

Final percent germination was recorded 3 days and germination curves recorded for up to 12

days after stratification. Seeds were counted as germinated if radicle emergence was observed.

To check the viability of ungerminated seeds, *S. parvula* seeds that failed to germinate on high

NaCl or KCl plates were transferred back to 1/4x MS or 1/2x MS and monitored for up to two

weeks.

Thermal stress tolerance assays

Five-day-old seedlings were subjected to heat stress at 38 °C for 6, 18 and 24 hr; chilling

stress at 4 °C for 24 hr; or freezing stress at 0 °C for 12 hr followed by a recovery time of 5 days

at 23 °C. Seedlings were considered to have survived if roots continued to grow without visible

chlorosis of the cotyledons after the recovery period monitored for another 5 days.

Quantification of leaf traits

Stomatal conductance and CO_2 assimilation rate were measured using the sixth or

seventh leaf from the shoot tip of hydroponically grown plants in LiCor 6400-40 leaf chamber

attached to a 6400XT gas analyzer (LiCor Inc, Lincoln Nebraska). Measurements were taken

after acclimation to 400 μmol photons m^{-2} s^{-1}, 400 μL L^{-1} CO_2 at 23 °C and an air flow rate of 200

μmol s^{-1} until a steady-state photosynthesis rate was attained. Each leaf was measured three

times and three plants were used for each treatment.

Four-week-old hydroponically grown plants subjected to 150 mM NaCl or control

treatments were used for the following leaf measurements. To quantify total leaf area, all

leaves from selected plants were scanned (Perfection V600 scanner, Epson, Suwa, Nagano,

Japan) and analyzed with ImageJ (Ferreira and Rasband, 2012). To obtain the relative water

content (RWC), newly harvested leaves were weighed before and after submerging in water for

24 hr (to get fresh and turgid weights respectively) followed by drying until dry weights were

obtained. RWC was calculated as (fresh weight - dry weight)/(turgid weight - dry weight). Leaf

temperature was measured on intact leaves of 4-week-old plants between ZT3 to ZT4

(Zeitgeber Time) every day for 7 days using thermal imaging (E6, FLIR, Wilsonville, Oregon). Leaf

temperatures were averaged across all leaves for each plant.

High-throughput phenotyping of morphometric and physiological traits

A. thaliana and *S. parvula* soil grown plants were phenotyped for compactness, leaf

water status, leaf area, and chlorophyll fluorescence using a PSI PlantScreen™ Compact system

(Qubit Phenomics, PSI). See supplementary method S1 for details.

Anatomical and flow cytometry analyses

Sections obtained from young roots (1 cm from the tip), mature roots (2 to 5 cm from

the root and shoot junction), stems (fourth, fifth, and sixth internodes from the shoot tip) and

leaves (fourth, fifth, and sixth from the shoot tip) from 8-week-old plants were used for

anatomical trait characterization. The fifth and sixth leaves from the shoot tip were used for

flow cytometry analysis as described in Galbraith et al., (1983). We counted the number of

stomata stained with propidium iodine on both hydroponically- and soil-grown plants and on

adaxial and abaxial surfaces at the same developmental stage and treatment duration. See

supplemental method S1 for details.

Reproductive trait quantification

Three-week-old soil grown plants were treated with an incremental increase of 50 mM

NaCl every 2 days until the final concentration was reached to 150 mM NaCl treatments and

maintained at 150 mM NaCl for the remainder of the experiment under either a long-day (16 hr

light/ 8 hr dark) or short-day (12 hr light/12 hr dark) photoperiod. Flowering time was recorded

when the first open flower was observed. Total numbers of flowering events, including all

developing, mature, and aborted flowers, as well as fully developed siliques were counted

separately for each plant throughout its lifetime. Fully developed siliques were counted and

harvested for imaging from 5-week-old plants that were given 150 mM NaCl or control

treatments for an additional 2 weeks.

Computational analysis

The data used to compare the basal expression of wax biosynthesizing genes between

A. thaliana and *S. parvula* were retrieved from NCBI-SRA database, accession# SRX877979 and

SRX877980, respectively (Oh et al., 2014). The expression profiles were visualized using

Integrated Genomics Viewer (v2.6) (Thorvaldsdóttir et al., 2013). The copy numbers and

transposition states of PhyB/D orthologs were determined using CLfinder-OrthNet pipeline as

described by Oh and Dassanayake, (2019). For principal component analysis (PCA) given in Fig 8,

as traits were measured in different units and scales, the measurements for each trait were first

normalized by either subtracting the mean (for anatomical traits) or dividing by the mean (for

physiological traits). The normalized data were subsequently analyzed with prcomp in R.

Acknowledgement

This work was supported by the US National Science Foundation/US-Israel Binational Science

Foundation award NSF-BSF-IOS-EDGE 1923589/2019610, NSF-MCB-1616827, US Department of

Energy BER-DE-SC0020358, Next-Generation BioGreen21 Program of Republic of Korea

(PJ01317301), and the Goldinger Trust Jewish Fund for the Future awards. Graduate students

K.T., G.W., P.P., and C.W. were supported by an Economic Development Assistantship and

undergraduate students J.C.G. and M.G.M. were supported by the President's Future Leaders in

Research Program at Louisiana State University (LSU). H.L. was sponsored by the China

Scholarship Council (CSC) through a Sino-Dutch Bilateral Exchange Scholarship. C.T. and Y.Z.

were funded by the European Research Council (ERC) under the EU Horizon 2020 Research and

Innovation programme (grant #724321). We acknowledge the productive discussions led by

Drs. Hans Bohnert and John Cheeseman at the University of Illinois at Urbana-

Champaign (UIUC) that prompted us to initiate this study. We thank the LSU High Performance

Computing facility for providing computational resources; Dr. Ying Xiao in the Shared

Instrumentation Facility at LSU for assistance with microscopy imaging; undergraduate

students Stephanie Presedo at LSU and Rebekah Munaretto, Michael Pettineo, and Aditya

Ravindra at UIUC for assisting with plant phenotyping; Jasper Lamers from WUR-PPH for

providing the script for automated root edge detection; and Aliza Finkler and Guilia Meshulam

at Tel Aviv University for providing technical help with the phenomics study.

Author contributions: K.T., P.P., G.W., and N.K. conducted the main experiments and analyzed

data as major contributors to the overall project; C.W., D-H.O., N.D., J.H., H.L., H.H., and P.A.

conducted additional experiments and analyzed data; undergraduate students J.C.J., R.K.,

M.G.M., A.C., D.T., and high school student C.C. conducted complementary experiments

supervised by K.T., P.P., G.W., D-H.O. and C.W.. M.D., C.T., and S.B. designed independent

complementary experiments; supervised students in their labs; and contributed to data

interpretation. D.L., Y.Z., and G.S. provided additional student supervision and assisted with

data generation for selected experiments. K.T., P.P., G.W., S.B., and M.D. wrote the manuscript.

K.T., P.P., G.W., N.K., D-H.O., C.W., P.A., M.F., J.C.L., A.S., D.L., P.F., C.T., S.B., and M.D. provided

critical reviews. M.D. conceptualized and supervised the overall project.

Main figure and table captions

Figure 1. Life cycle of *Schrenkiella parvula* from seeds to siliques. Scale bars are 1 cm unless

indicated in the figure.

Table 1. Morphological comparison between *Arabidopsis thaliana* and *Schrenkiella parvula*.

Figure 2. Effects of NaCl stress on root growth in *Schrenkiella parvula* and *Arabidopsis thaliana*

seedlings. [A] 12-day-old seedlings of *A. thaliana* and *S. parvula* were grown for 5 days on 1/4x

Murashige and Skoog media, and 7 days on the indicated concentrations of NaCl. Plates were

scanned 7 days after treatment (DAT). [B] Primary root growth and average root growth rate,

[C] number of lateral roots and lateral root density, and [D] average length of 10 longest root

hairs. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their

respective control group, determined by Student's *t*-test. Data are mean ± SD (n =3, at least 3

plants per replicate). Open circles are individual measurements.

Figure 3. Effects of NaCl stress on *Schrenkiella parvula* root anatomy. Eight-week-old *S. parvula*

plants were grown under control or 150 mM NaCl conditions for an additional 4 weeks. [A] The

position of transverse sections of *S. parvula* roots, [B] the root tip, [C] the young root, and [D]

the mature root from 12-week-old hydroponically-grown *S. parvula* plants. A minimum of 20

sections from 4 to 13 plants were examined for each root region to quantify each parameter for

each condition. Asterisks indicate significant differences ($p \leq 0.05$) between the treated samples

and their respective control group, determined by Student's *t*-test. Data are means ± SD. Data

points represent individual cross-sections and colors represent individual plants. Representative

cross-sections were obtained from the control plants. Scale bar represents 100 μm.

Figure 4. Effects of NaCl stress on *Schrenkiella parvula* shoot anatomy. Eight-week-old *S.*

parvula plants were grown under control or 150 mM NaCl conditions for an additional 4 weeks.

Transverse section of [A] stem and [B] leaf of 12-week-old hydroponically-grown *S. parvula*

plants. A total of 23 sections from 7 control plants and 32 sections from 9 treated plants were

used for leaf measurements, and a minimum of 20 sections from 5 to 11 plants were used for

stem measurements. Data points represent individual cross-sections and colors represent

individual plants. [C] Nuclear DNA content, distribution of leaf cell ploidy, and

endoreduplication index of leaf cells from 10^{th} and 11^{th} leaves from the shoot meristem in

control and 250 mM NaCl-treated 8-week-old *S. parvula* plants. Data are mean ± SD (n= 4).

Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their

respective control group, determined by Student's *t*-test. Representative cross-sections were

obtained from the control plants Scale bars represent 100 μm.

Figure 5. Effects of NaCl stress on *Schrenkiella parvula* leaf traits. All experiments were

performed with 4-week-old hydroponically-grown plants that were treated for an additional 4

weeks with the indicated NaCl concentrations. [A] Total leaf area, [B] Stomatal density, [C]

Stomatal conductance, [D] Photosynthesis rate, [E] Leaf relative surface temperature, and [F]

Leaf relative water content. Asterisks indicate significant difference ($p \leq 0.05$) between the

treated samples and their respective control samples, determined by Student's *t*-test. Data are

means ± SD (n ≥ 3). Open circles indicate the measurement from each plant (in B and F) and

leaves (in C and D) for each experiment. DAT, days after treatment.

Figure 6. Effects of NaCl on reproductive traits of *Schrenkiella parvula*. [A] Plants grown under

control conditions generally flower at the ~17th leaf stage and the first few flowers are

subsequently aborted. [B] Salt treated plants flower earlier at the ~14th leaf stage and the first

flowers develop into mature siliques. [C] Days from planting to the first observed flower of

plants grown under a long day (16 hr light/8 hr dark) or short day (12 hr light/12 hr dark)

photoperiod with and without the indicated salt treatment. Center line - median; box -

interquartile range (IQR); notch - 1.58 × IQR/sqrt(n); whiskers - 1.5 × IQR. [D] Flowers obtained

from control and treated plants described in panel A and B. [E] Number of flowers and siliques

per plant under control and salt treatment. Data are mean ± SD. Open circles represent

individual measurements from four biological replicates. [F] Ratio between the number of

flowers and siliques observed for control (C) and salt-treated (T) plants described in E. [G]

Siliques from plants under control and NaCl treated conditions. [H] Size of siliques from G. Open

circles represent individual silique counts. [I] Comparison of the area of seeds from plants under

control and NaCl treated conditions. Data are mean ± SD (n = 50). Seeds per condition were

selected from the seed pool of 15 to 50 plants. Open circles represent individual seeds. For

panel [B], salt treatment started 21 DAP, initially at 50 mM NaCl given every other day. The salt

concentration was increased by 50 mM every four days until it reached 200 mM; for panels [C,

E, F], salt treatment was applied to 3-week-old plants until the end of the experiment; for

panels [G, H, I], salt treatment was applied to 5-week-old plants for an additional 2 weeks.

Different letters or asterisks represent significant differences (*, $p < 0.05$) compared to control,

determined by either one-way ANOVA followed by Tukey's post-hoc test [C] or Student's *t*-test

[E, H, I]. DAP, days after planting.

Figure 7. Effects of salt stress on the germination of *Arabidopsis thaliana* and *Schrenkiella*

parvula seeds. [A] Final percent germination recorded at 3 days and [B] germination curves of

seeds treated with different concentrations of NaCl recorded for 12 days after stratification. [C]

Germination curves of NaCl-treated seeds transferred to control (no NaCl) media, after 2 days

or 30 days on salt plates. The germination curves were performed by direct sowing of seeds

onto salt plates prior to stratification. Black arrows point to the day when seeds were

transferred to non-NaCl plates. [D, E, F] Final percent germination of seeds treated with KCl [D],

LiCl [E] or H_3BO_3 [F]. Data are mean ± SD (n = 3-4, each replicate contains ca. 50 seeds). Open

circles indicate individual measurements. Different letters indicate significant difference ($p \leq$

0.05) determined by one-way ANOVA with post-hoc Tukey's test across all conditions in [A, D, E,

F], or across treatment groups at 3 and 12, and 7 days after stratification in [B] and [C]

respectively.

Figure 8. Summary of salt-induced structural and physiological traits in *Schrenkiella parvula*. [**A**]

Salt- induced changes in traits quantified in the current study. Changes in each trait were

calculated as the fold change between treatment and control measurements. Traits highlighted

in the red boxes showed significant differences ($p < 0.05$) under salt treatments compared to

the control determined in previous assays (Fig 2-7). [**B**] and [**C**] PCA biplot of traits quantified

for *S. parvula* under control and salt-treated conditions. Arrows indicate directions of loadings

for each trait and are color-coded by contribution to the variations in PC1 and PC2. [B]

Anatomical traits quantified: LT, leaf thickness; MT, midrib thickness; XS, stem xylem to stem

area; SVA, average area per vessel in stems; CbS, cambium to stem area; CtS, cortex to stem

area; XR, root xylem to root area; RVA, average area per vessel in roots; VN, number of vessels

across the root diameter; AS, air space to root area; YRA, young root area; CT, cortical

thickness; YRD, root diameter in young roots; DRH, distance from root tip to first root hair. [C]

Physiological traits quantified: AbS, abaxial stomatal density; adS, Adaxial stomatal density; LA,

leaf area; SLA, silique area; SA, seed area; PRL, primary root length; LRD, lateral root density;

RWC, relative water content; CA, CO_2 assimilation; SC, stomatal conductance; RHL, root hair

length; NF, number of flowers; NS, number of siliques. DAT, days after treatment; DAP, days

after planting.

Supplement figure and table captions

Figure S1. Effects of NaCl stress on root growth of *Schrenkiella parvula* and *Arabidopsis thaliana*

under long days and higher nutrient conditions. [A] Root growth of 10-day-old seedlings under

indicated concentrations of NaCl. Quantification of [B] primary root growth, [C] number of

lateral roots, [D] average lateral root length, [E] total lateral root length, and [F] lateral root

density. Asterisks indicate significant difference (p-adj ≤ 0.05) between the treated samples and

their respective control samples, determined by two-way ANOVA. Data are represented as

mean ± SD (n = 15 to 20). Open circles indicate the number of plants used for each experiment.

Figure S2. Effects of NaCl stress on root hair growth in *Schrenkiella parvula* and *Arabidopsis*

thaliana. White arrows indicate root tip positions when the 5-day-old seedlings were

transferred to the indicated salt concentrations. Note that the white arrows were absent in *A.*

thaliana and *S. parvula* control because the position of the root tip at the time of transferring

were out of frame when we imaged the root tip region. Scale bars are 0.5 mm. DAT, Days after

treatment.

Figure S3. Long term effects of NaCl stress on the root growth of *Schrenkiella parvula*. [A] Root

fresh weight and [B] primary root length of 12-week-old hydroponically grown *S. parvula* plants.

Four-week-old plants were subjected to 150 mM NaCl for an additional 4 weeks and the

treatment was continued or increased to 250 mM NaCl for another 4 weeks. The plants were

grown at 100-150 µmol m^{-2} s^{-1} photosynthetic photon flux density with a 12 hr light/12 hr dark

photoperiod. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples

and their respective control samples, determined by Student's t-test. Data are represented as

the mean ± SD (n = 9). For each plot, an open circle indicates the measurement from each plant.

Figure S4. Halotropism assay at 200 mM NaCl for *Schrenkiella parvula* and *Arabidopsis thaliana*.

The blue lines indicate the position of seeds sown. The yellow line marks the separation

between 1/4x MS media with 0 mM (top) and 200 mM NaCl (bottom). The bent root angle is

indicated by curved white lines. g, gravity axis.

Figure S5. Root anatomy of *Schrenkiella parvula*. [A] Transverse section of the young root taken

at 1-3 cm from the root tip. [1] Young root area; [2] root diameter; [3] epidermis thickness; [4]

cortical layer thickness; [5] endodermis thickness. [B] Transverse section of the mature root

taken between 2-5 cm below the root-shoot junction. [1] Mature root area; [2] xylem/root

area; [3] Area per vessel; [4] # vessels/root diameter; [5] Air space/root area. [C] Longitudinal

section of young root tip from 7-day-old seedlings. The cortex layer is highlighted in yellow. The

red arrow indicates the additional cell layer.

Figure S6. Stem anatomy of *Schrenkiella parvula*. [A] Transverse section of *S. parvula* stem

(between 4th, 5th, and 6th internodes from the shoot meristem) with cell layers marked in

yellow lines for trait quantification. [B] Measurements of stem anatomical features indicated in

[A]. A minimum of 20 sections from 5-11 plants were used for each measurement. Asterisks

indicate significant difference ($p \leq 0.05$) between the treated samples and its respective control

samples, determined by Student's t-test. Data points represent individual cross-sections and

colors represent individual plants.

Figure S7. Leaf structures of *Schrenkiella parvula*. [A] Transverse section of *S. parvula* leaves. [1]

leaf thickness and [2] midrib thickness. [B] Transverse sections of *S. parvula* leaves treated with

indicated concentration of salts. All sections represent the fifth or sixth leaf from the root-shoot

junction in 8-week-old plants that were treated for 4 weeks.

Figure S8. Effects of salt on growth of 4-week-old *Arabidopsis thaliana* and *Schrenkiella parvula*

treated for an additional 2 weeks at 22 °C under a 12 hr light/12 hr dark photoperiod in soil.

Figure S9. Effects of NaCl stress on *Arabidopsis thaliana* and *Schrenkiella parvula* morphometric

and physiological traits*.* Soil-grown, 8-day-old *A. thaliana* and 12-day-old *S. parvula* were

exposed to the indicated salt concentrations at 21 °C, 16 h light/8 h dark photoperiod. The

Measurements were taken on day 8, 12, and 15 after the start of the salt treatments. [A]

Compactness (the ratio between the rosette (or leaf) area and the rosette (or leaf) convex hull

area); [B] Leaf area; [C] Leaf water content; [D] Maximum quantum efficiency of PSII; and [E]

Non-photochemical quenching were measured/estimated using the Qubit/PSI PlantScreen™

Compact System. For leaf area and leaf water content, values are percent of control (0 mM

NaCl) for each time point. Letters denote significant changes at p-adj ≤ 0.05 (one-way ANOVA

with post-hoc Tukey's test). Data are mean ± SD (n = 10). Open circles indicate the individual

measurements obtained from each plant. DAT, days after treatment.

Figure S10. Leaf surface and the basal expression of genes involved in wax biosynthesis in

Arabidopsis thaliana and *Schrenkiella parvula*. [A] Scanning electron micrographs contrasting *A.*

thaliana and *S. parvula* leaf surfaces. [B] Major wax biosynthesis pathway and [C] wax

biosynthesis genes that exhibited significantly different basal expression between *A. thaliana*

and *S. parvula*.

Figure S11. The potential influence of phytochrome family of genes on *Schrenkiella parvula* leaf

and growth form. [A] Elongated internode and leaf petiole in *S. parvula* compared to *A.*

thaliana. [B] *PHYD* is absent in the *S. parvula* genome as illustrated using OrthNet representing

evolutionary histories of orthologous gene groups derived from six Brassicaceae genomes:

Arabidopsis lyrata (Aly, version 1.0), *Arabidopsis thaliana* (Ath, v. 'TAIR10'), *Capsella rubella*

(Cru, v. 1.0), and *Eutrema salsugineum* (Esa, v. 1.0), *Sisymbrium irio* (Sir, v. 0.2), and *Schrenkiella*

parvula (v. 2.0). Nodes are color-coded according to the species. Edges show either co-linear

(cl) or transposed (tr) properties.

Figure S12. Effects of KCl on reproductive traits of *Schrenkiella parvula*. [A] Plants grown under

control conditions generally flower late and the first few flowers are subsequently aborted. KCl-

treated plants flower earlier and the first flowers develop into mature siliques. [B] Early flowers

in KCl treated plants develop longer filaments compared to control plants. DAP, days after

planting.

Figure S13. Effect of salt on *Arabidopsis thaliana* and *Schrenkiella parvula* inducing barriers for

seed germination and seedling establishment. [A] *S. parvula* (*Sp*) and *A. thaliana* (*At*) seed

germination and growth on NaCl- and KCl-supplemented MS medium. Scale bars are 2 cm. [B]

Growth of ungerminated *S. parvula* seeds treated with different concentrations of NaCl (as

shown in Figure 7) after transferring to control MS medium.

Figure S14. Survival of *Schrenkiella parvula* in the climate conditions found in Lake Tuz region.

[A] Lake Tuz location from Google Earth; [B] Precipitation and temperature recorded in Konya,

Turkey (NOAA, 2019). *S. parvula* growing season is from April/May to August/September (Tug

et al., 2019). Bars indicate precipitation and lines represent temperature. [C] Average day

length recorded in Konya per month. Red line indicates the day length used for typical

laboratory growth assays for *S. parvula*. [D-F] Survival rate of *A. thaliana* and *S. parvula*

seedlings under [D] freezing stress, [E] chilling stress, and [F] heat stress. Five-day-old seedlings

were subjected to temperature treatments and data were recorded 5 days after recovery.

Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their

respective control samples (student's *t*-test). Data are mean ± SD (n = 4). Each replicate plate

contained 3-6 plants. Open circles indicate the biological replicates.

References

Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, Moritz T, Van Der Straeten D, Peng

J, Harberd NP (2006) Integration of plant responses to environmentally activated

phytohormonal signals. Science (80-) **311**: 91–94

Adams P, Nelson DE, Yamada S, Wendy C, Jensen RG, Bohnert HJ, Griffiths H (1998) Tansley

Review No. 97 Growth and development of *Mesembryanthemum crystallinum* (Aizoaceae).

Genet Breed 171–190

Ali A, Khan IU, Jan M, Khan HA, Hussain S, Nisar M, Chung WS, Yun D-J (2018) The High-

Affinity Potassium Transporter EpHKT1;2 From the Extremophile *Eutrema parvula*

Mediates Salt Tolerance. Front Plant Sci. doi: 10.3389/fpls.2018.01108

Ali A, Yun DJ (2017) Salt stress tolerance; what do we learn from halophytes? J Plant Biol **60**:

431–439

Apse MP, Aharon GS, Snedden WA, Blumwald E (1999) Salt Tolerance Conferred by

Overexpression of a Vacuolar Na^+/H^+ Antiport in *Arabidopsis*. Science (80-) **285**: 1256–

1258

Barkla BJ, Rhodes T, Tran KNT, Wijesinghege C, Larkin JC, Dassanayake M (2018) Making

epidermal bladder cells bigger: Developmental-and salinity-induced endopolyploidy in a

model halophyte. Plant Physiol **177**: 615–632

Barros NLF, Marques DN, Tadaiesky LBA, de Souza CRB (2021) Halophytes and other molecular

strategies for the generation of salt-tolerant crops. Plant Physiol Biochem. doi:

10.1016/j.plaphy.2021.03.028

Bechtold U (2018) Plant Life in Extreme Environments : How Do You Improve Drought

Tolerance ? **9**: 1–8

Bernard A, Joubès J (2013) *Arabidopsis* cuticular waxes: Advances in synthesis, export and

regulation. Prog Lipid Res. doi: 10.1016/j.plipres.2012.10.002

Blits KC, Gallagher JL (1991) Morphological and physiological responses to increased salinity in

marsh and dune ecotypes of *Sporobolus virginicus* (L.) Kunth. Oecologia **87**: 330–335

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the target of

crop yield improvement under drought stress. F Crop Res **112**: 119–123

de Boer HJ, Drake PL, Wendt E, Price CA, Schulze ED, Turner NC, Nicolle D, Veneklaas EJ (2016)

Apparent overinvestment in leaf venation relaxes leaf morphological constraints on

photosynthesis in arid habitats. Plant Physiol **172**: 2286–2299

Boughalleb F, Denden M, Tiba B Ben (2009) Anatomical changes induced by increasing NaCl

salinity in three fodder shrubs, *Nitraria retusa*, *Atriplex halimus* and *Medicago arborea*.

Acta Physiol Plant **31**: 947–960

Brady SM, Orlando DA, Lee JY, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN

(2007) A high-resolution root spatiotemporal map reveals dominant expression patterns.

Science (80-). doi: 10.1126/science.1146265

Caño L, Fuertes-Mendizabal T, García-Baquero G, Herrera M, Begoña González-Moro M (2016)

Plasticity to salinity and transgenerational effects in the nonnative shrub *Baccharis*

halimifolia: Insights into an estuarine invasion. Am J Bot **103**: 808–820

Cheeseman JM (2013) The integration of activity in saline environments: Problems and

perspectives. Funct Plant Biol. doi: 10.1071/FP12285

Cho LH, Yoon J, An G (2017) The control of flowering time by environmental factors. Plant J **90**:

708–719

Conn SJ, Hocking B, Dayod M, Xu B, Athman A, Henderson S, Aukett L, Conn V, Shearer MK,

Fuentes S, et al (2013) Protocol: Optimising hydroponic growth systems for nutritional and

physiological analysis of *Arabidopsis thaliana* and other plants. Plant Methods **9**: 1–11

Cruz MV, Mori GM, Signori-Müller C, da Silva CC, Oh DH, Dassanayake M, Zucchi MI, Oliveira

RS, de Souza AP (2019) Local adaptation of a dominant coastal tree to freshwater

availability and solar radiation suggested by genomic and ecophysiological approaches. Sci

Rep **9**: 1–15

Dassanayake M, Larkin JC (2017) Making Plants Break a Sweat: the Structure, Function, and

Evolution of Plant Salt Glands. Front Plant Sci **08**: 406

Dassanayake M, Oh D-H, Haas JSJS, Hernandez A, Hong H, Ali S, Yun DJD-J, Bressan RARA, Zhu

JKJ-K, Bohnert HJHJ, et al (2011) The genome of the extremophile crucifer *Thellungiella*

parvula. Nat Genet **43**: 913–918

Devlin PF, Halliday KJ, Harberd NP, Whitelam GC (1996) The rosette habit of *Arabidopsis*

thaliana is dependent upon phytochrome action: novel phytochromes control internode

elongation and flowering time. Plant J **10**: 1127–1134

Drake PL, de Boer HJ, Schymanski SJ, Veneklaas EJ (2019) Two sides to every leaf: water and

CO_2 transport in hypostomatous and amphistomatous leaves. New Phytol **222**: 1179–1187

Du F, Guan C, Jiao Y (2018) Molecular Mechanisms of Leaf Morphogenesis. Mol Plant **11**: 1117–

1134

Eshel G, Duppen N, Wang G, Oh D-H, Kazachkova Y, Herzyk P, Amtmann A, Gordon M, Chalifa-

Caspi V, Arland Oscar M, et al (2021) Positive Selection and Heat-Response

Transcriptomes Reveal Adaptive Features of the Arabidopsis Desert Relative, *Anastatica*

hierochuntica. bioRxiv

Ferreira T, Rasband W (2012) ImageJ User Guide User Guide ImageJ. Image J user Guid. doi:

10.1038/nmeth.2019

Flowers TJ, Colmer TD (2008) Salinity tolerance in halophytes. New Phytol 945–963

Galvan-Ampudia CS, Julkowska MM, Darwish E, Gandullo J, Korver RA, Brunoud G, Haring

MA, Munnik T, Vernoux T, Testerink C (2013) Halotropism is a response of plant roots to

avoid a saline environment. Curr Biol **23**: 2044–2050

Ge W, Zhang Y, Sun Z, Li J, Liu G, Ma Y, Gao J (2017) Physiological and anatomical responses of

Phyllostachys vivax and *Arundinaria fortunei* (Gramineae) under salt stress. Rev Bras Bot

40: 79–91

German DA, Al-Shehbaz IA (2010) Nomenclatural novelties in miscellaneous Asian Brassicaceae

(Cruciferae). Nord J Bot **28**: 646–651

Hajiboland R, Bahrami-Rad S, Akhani H, Poschenrieder C, Bahrami S, Hossein R, Charlotte A

(2018) Salt tolerance mechanisms in three Irano-Turanian Brassicaceae halophytes

relatives of *Arabidopsis thaliana*. J Plant Res **0**: 0

Helvaci C, Mordogan H, Çolak M, Gündogan I (2004) Presence and Distribution of Lithium in

Borate Deposits and Some Recent Lake Waters of West-Central Turkey. Int Geol Rev **46**:

177–190

IPCC (2021) Climate Change 2021: The Physical Science Basis. Contribution of Working Group I

to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press

IPNI (2021) International Plant Names Index. Published on the Internet http://www.ipni.org,

The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian

National Botanic Gardens. https://www.ipni.org/

Jarvis DE, Ryu CH, Beilstein MA, Schumaker KS (2014) Distinct roles for SOS1 in the convergent

evolution of salt tolerance in *Eutrema salsugineum* and *Schrenkiella parvula*. Mol Biol Evol

31: 2094–2107

Jennings DH (1968) Halophytes, Succulence and Sodium in Plants—a Unified Theory. New

Phytol **67**: 899–911

Junghans U, Polle A, Düchting P, Weiler E, Kuhlman B, Gruber F, Teichmann T (2006)

Adaptation to high salinity in poplar involves changes in xylem anatomy and auxin

physiology. Plant, Cell Environ **29**: 1519–1531

Kazachkova Y, Eshel G, Pantha P, Cheeseman JM, Dassanayake M (2018) Halophytism : What

Have We Learnt From *Arabidopsis thaliana* Relative Model Systems ? **178**: 972–988

Kazachkova Y, Khan A, Acuña T, López-Díaz I, Carrera E, Khozin-Goldberg I, Fait A, Barak S

(2016) Salt Induces Features of a Dormancy-Like State in Seeds of *Eutrema* (*Thellungiella*)

salsugineum, a Halophytic Relative of *Arabidopsis*. Front Plant Sci **7**: 1–18

King DA (1997) The functional significance of leaf angle in *Eucalyptus*. Aust J Bot **45**: 619–639

Kotula L, Garcia Caparros P, Zörb C, Colmer TD, Flowers TJ (2020) Improving crop salt tolerance

using transgenic approaches: An update and physiological analysis. Plant Cell Environ **43**:

2932–2956

Krämer U (2018) Conceptualizing plant systems evolution. Curr Opin Plant Biol **42**: 66–75

Li AJ, Li G, Wang H, Deng XW (2011) Phytochrome Signaling Mechanisms. Arab B. doi:

10.1199/tab.0148

Li H, Testerink C, Zhang Y (2021) How roots and shoots communicate through stressful times.

Trends Plant Sci. doi: 10.1016/j.tplants.2021.03.005

Li K, Wang Y, Han C, Zhang W, Jia H, Li X (2007) GA signaling and CO/FT regulatory module

mediate salt-induced late flowering in *Arabidopsis thaliana*. Plant Growth Regul **53**: 195–

206

Lloyd EA, Oreskes N (2018) Climate Change Attribution: When Is It Appropriate to Accept New

Methods? Earth's Futur **6**: 311–325

Lutts S, Kinet JM, Bouharmont J (1995) Changes in plant response to NaCl during development

of rice (*Oryza sativa* L.) varieties differing in salinity resistance. J Exp Bot **46**: 1843–1852

Moriuchi KS, Friesen ML, Cordeiro MA, Badri M, Vu WT, Main BJ, Aouani ME, Nuzhdin S V.,

Strauss SY, Von Wettberg EJB (2016) Salinity adaptation and the contribution of parental

environmental effects in *Medicago truncatula*. PLoS One **11**: 1–19

Müller P, Li XP, Niyogi KK (2001) Non-photochemical quenching. A response to excess light

energy. Plant Physiol. doi: 10.1104/pp.125.4.1558

Nadeau JA, Sack FD (2002) Stomatal Development in *Arabidopsis*. Arab B **1**: e0066

Nassar RMA, Kamel HA, Ghoniem AE, Alarcón JJ, Sekara A, Ulrichs C, Abdelhamid MT (2020)

Physiological and anatomical mechanisms in wheat to cope with salt stress induced by

seawater. Plants **9**: 1–15

Ogburn RM, Edwards EJ (2013) Repeated origin of three-dimensional leaf venation releases

constraints on the evolution of succulence in plants. Curr Biol **23**: 722–726

Oh D-H, Dassanayake M, Bohnert HJ, Cheeseman JM (2012) Life at the extreme: lessons from

the genome. Genome Biol **13**: 241

Oh D-HHD-H, Hong H, Lee SY, Yun D-JD-J, Bohnert HJ, Dassanayake M (2014) Genome

Structures and Transcriptomes Signify Niche Adaptation for the Multiple-Ion-Tolerant

Extremophyte *Schrenkiella parvula*. Plant Physiol **164**: 2123–2138

Oh DH, Dassanayake M (2019) Landscape of gene transposition-duplication within the

Brassicaceae family. DNA Res. doi: 10.1093/dnares/dsy035

Olson ME, Rosell JA (2013) Vessel diameter-stem diameter scaling across woody angiosperms

and the ecological causes of xylem vessel diameter variation. New Phytol **197**: 1204–1213

Orsini F, D'Urzo MP, Inan G, Serra S, Oh D-HH, Mickelbart M V., Consiglio F, Li X, Jeong JC, Yun

D-JJ, et al (2010) A comparative study of salt tolerance parameters in 11 wild relatives of

Arabidopsis thaliana. J Exp Bot **61**: 3787–3798

Ozfidan-Konakci C, Uzilday B, Ozgur R, Yildiztugay E, Sekmen AH, Turkan I (2015) Halophytes

as a source of salt tolerance genes and mechanisms : a case study for the Salt Lake area ,

Turkey.

Pantha P, Oh D-H, Longstreth D, Dassanayake M (2021) Living with high potassium: an asset or

a hindrance. bioRxiv 2021.07.01.450778

Pardo J, VanBuren R (2021) Evolutionary innovations driving abiotic stress tolerance in C4

grasses and cereals. Plant Cell

Parida AK, Veerabathini SK, Kumari A, Agarwal PK (2016) Physiological, anatomical and

metabolic implications of salt tolerance in the halophyte *Salvadora persica* under

hydroponic culture condition. Front Plant Sci **7**: 1–18

Pierik R, Testerink C (2014) The art of being flexible: How to escape from shade, Salt, And

drought. Plant Physiol **166**: 5–22

De Rocher EJ, Harkins KR, Galbraith DW, Bohnert HJ (1990) Developmentally regulated

systemic endopolyploidy in succulents with small genomes. Science (80-) **250**: 99–101

Rodell M, Famiglietti JS, Wiese DN, Reager JT, Beaudoing HK (2018) Emerging trends in global
freshwater availability. doi: 10.1038/s41586-018-0123-1

Roy SJ, Negrão S, Tester M (2014) Salt resistant crop plants. Curr Opin Biotechnol **26**: 115–124

Ryu JY, Lee HJ, Seo PJ, Jung JH, Ahn JH, Park CM (2014) The *Arabidopsis* floral repressor BFT
delays flowering by competing with FT for FD binding under high salinity. Mol Plant **7**: 377–
387

Santiago-Rosario L, Kyle H, Elderd B, Hart P, Dassanayake M (2021) No Escape: The Influence
of Substrate Sodium on Plant Growth and Tissue Sodium Responses. Ecol Evol. doi:
10.22541/au.162081817.72637329/v1

Sarker U, Oba S (2020) The Response of Salinity Stress-Induced A. tricolor to Growth, Anatomy,
Physiology, Non-Enzymatic and Enzymatic Antioxidants. Front Plant Sci **11**: 1–14

Schlenker W, Auffhammer M (2018) The cost of a warming climate.

Solis CA, Yong MT, Vinarao R, Jena K, Holford P, Hill C, Chen Z (2020) Back to the Wild : On a
Quest for Donors Toward Salinity Tolerant Rice. **11**: 1–14

Somers DE, Sharrock RA, Tepperman JM, Quail PH (1991) The hy3 long hypocotyl mutant of
Arabidopsis is deficient in phytochrome B. Plant Cell **3**: 1263–1274

Stanton ML, Roy BA, Thiede DA (2000) Evolution in Stressful Environments . I . Phenotypic
Variability , Phenotypic Selection , and Response to Selection in Five Distinct
Environmental Stresses Author (s): M . L . Stanton , B . A . Roy and D . A . Thiede Published
by : Society for the Study. Evolution (N Y) **54**: 93–111

Sun K, Cui Y, Hauser BA (2005) Environmental stress alters genes expression and induces ovule
abortion: Reactive oxygen species appear as ovules commit to abort. Planta **222**: 632–642

Teusink RS, Rahman M, Bressan RA, Jenks MA (2002) Cuticular Waxes on *Arabidopsis thaliana*
Close Relatives *Thellungiella halophila* and *Thellungiella parvula*. Int J Plant Sci **163**: 309–
315

Thorvaldsdóttir H, Robinson JT, Mesirov JP (2013) Integrative Genomics Viewer (IGV): High-
performance genomics data visualization and exploration. Brief Bioinform. doi:
10.1093/bib/bbs017

Tug GN, Yaprak AE, Vural M (2019) The Floristical, Ecological, and Syntaxonomical

Characteristics of Salt Marshes and Salt Steppes in Turkey. Tasks Veg. Sci. pp 413–446

Ungar IA (1978) Halophyte seed germination. Bot Rev **44**: 233–264

Ventura Y, Myrzabayeva M, Alikulov Z, Omarov R, Khozin-Goldberg I, Sagi M (2014) Effects of

salinity on flowering, morphology, biomass accumulation and leaf metabolites in an edible

halophyte. AoB Plants. doi: 10.1093/aobpla/plu053

Wang G, Ditusa SF, Oh D, Herrmann AD, Mendoza-cozatl DG, Neill MAO, Smith AP,

Dassanayake M (2021) Cross species multi-omics reveals cell wall sequestration and

elevated global transcript abundance as mechanisms of boron tolerance in plants. 1985–

2000

Wang G, Pantha P, Tran KN, Oh DH, Dassanayake M (2019) Plant growth and agrobacterium-

mediated floral-dip transformation of the extremophyte *Schrenkiella parvula*. J Vis Exp

2019: 1–8

Wang XJ, Hu QJ, Guo XY, Wang K, Ru DF, German DA, Weretilnyk EA, Abbott RJ, Lascoux M,

Liu J quan (2018) Demographic expansion and genetic load of the halophyte model plant

Eutrema salsugineum. Mol Ecol **27**: 2943–2955

Yamaguchi T, Blumwald E (2005) Developing salt-tolerant crop plants: Challenges and

opportunities. Trends Plant Sci **10**: 615–620

Yoo CY, Pence HE, Hasegawa PM, Mickelbart M V. (2009) Regulation of transpiration to

improve crop water use. CRC Crit Rev Plant Sci **28**: 410–431

Yu Y, Wang L, Chen J, Liu Z, Park CM, Xiang F (2018) WRKY71 Acts Antagonistically Against Salt-

Delayed Flowering in *Arabidopsis thaliana*. Plant Cell Physiol **59**: 414–422

Zandalinas SI, Fritschi FB, Mittler R (2021) Global Warming , Climate Change , and

Environmental Pollution : Recipe for a Multifactorial Stress Combination Disaster Trends in

Plant Science. Trends Plant Sci **26**: 588–599

Zhu JK (2015) The Next Top Models. Cell **163**: 18–20

Figure and table captions

Figure 1. Life cycle of *Schrenkiella parvula* from seeds to siliques. Scale bars are 1 cm unless indicated in the figure.

Table 1. Morphological comparison between *Arabidopsis thaliana* and *Schrenkiella parvula*.

Morphology	A. thaliana	S. parvula
Growth habit	Determinate	Indeterminate
Leaf arrangement	Rosette	Spiral phyllotaxy
Leaf shape	Ovate	Linear-lanceolate
Trichomes	Present	Absent
Petals	4	Absent
Number of stamens	6 (2 short and 4 long)	6 (equal size)
Siliques/plant[a]	74.9±18.69	61.6±14.40
Silique length (cm)	1.16±0.09	1.48±0.06
Seeds/silique	23.91±13.60	21.58±5.57
Seed weight[b] (g)	0.020±0.003	0.063±0.013
Seed area (mm^2)	0.12±0.02	0.31±0.04

[a] Quantification from 10-12-week-old plants grown under 14-hr light/10-hr dark photoperiod, 22 °C – 24 °C temperature with a light intensity of 130 µmol m^{-2} s^{-1}.
[b] Average weight of 500 seeds.
Data are represented as mean of 3 independent replicates ± SD.

Figure 2. Effects of NaCl stress on root growth in *Schrenkiella parvula* and *Arabidopsis thaliana* seedlings. [A] 12-day-old seedlings of *A. thaliana* and *S. parvula* were grown for 5 days on 1/4x Murashige and Skoog media, and 7 days on the indicated concentrations of NaCl. Plates were scanned 7 days after treatment (DAT). [B] Primary root growth and average root growth rate, [C] number of lateral roots and lateral root density, and [D] average length of 10 longest root hairs. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their respective control group, determined by Student's t-test. Data are mean ± SD (n =3, at least 3 plants per replicate). Open circles are individual measurements.

Figure S1. Effects of NaCl stress on root growth of *Schrenkiella parvula* and *Arabidopsis thaliana* under long days and higher nutrient conditions. [A] Root growth of 10-day-old seedlings under indicated concentrations of NaCl. Quantification of [B] primary root growth, [C] number of lateral roots, [D] average lateral root length, [E] total lateral root length, and [F] lateral root density. Asterisks indicate significant difference (p-adj ≤ 0.05) between the treated samples and their respective control samples, determined by two-way ANOVA. Data are represented as mean ± SD (n = 15 to 20). Open circles indicate the number of plants used for each experiment.

Figure S2. Effects of NaCl stress on root hair growth in *Schrenkiella parvula* and *Arabidopsis thaliana*. White arrows indicate root tip positions when the 5-day-old seedlings were transferred to the indicated salt concentrations. Note that the white arrows were absent in *A. thaliana* and *S. parvula* control because the position of the root tip at the time of transferring were out of frame when we imaged the root tip region. Scale bars are 0.5 mm. DAT, Days after treatment.

Figure S3. Long term effects of NaCl stress on the root growth of *Schrenkiella parvula*. [A] Root fresh weight and [B] primary root length of 12-week-old hydroponically grown *S. parvula* plants. Four-week-old plants were subjected to 150 mM NaCl for an additional 4 weeks and the treatment was continued or increased to 250 mM NaCl for another 4 weeks. The plants were grown at 100-150 μmol m^{-2} s^{-1} photosynthetic photon flux density with a 12 hr light/12 hr dark photoperiod. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their respective control samples, determined by Student's *t*-test. Data are represented as the mean ± SD (n = 9). For each plot, an open circle indicates the measurement from each plant.

Figure S4. Halotropism assay at 200 mM NaCl for *Schrenkiella parvula* and *Arabidopsis thaliana*. The blue lines indicate the position of seeds sown. The yellow line marks the separation between 1/4x MS media with 0 mM (top) and 200 mM NaCl (bottom). The bent root angle is indicated by curved white lines. g, gravity axis.

Figure 3. Effects of NaCl stress on *Schrenkiella parvula* root anatomy. Eight-week-old *S. parvula* plants were grown under control or 150 mM NaCl conditions for an additional 4 weeks. [A] The position of transverse sections of *S. parvula* roots, [B] the root tip, [C] the young root, and [D] the mature root from 12-week-old hydroponically-grown *S. parvula* plants. A minimum of 20 sections from 4 to 13 plants were examined for each root region to quantify each parameter for each condition. Asterisks indicate significant differences ($p \leq$ 0.05) between the treated samples and their respective control group, determined by Student's *t*-test. Data are means ± SD. Data points represent individual cross-sections and colors represent individual plants. Representative cross-sections were obtained from the control plants. Scale bar represents 100 μm.

Figure S5. Root anatomy of *Schrenkiella parvula*. [A] Transverse section of the young root taken at 1-3 cm from the root tip. [1] Young root area; [2] root diameter; [3] epidermis thickness; [4] cortical layer thickness; [5] endodermis thickness. [B] Transverse section of the mature root taken between 2-5 cm below the root-shoot junction. [1] Mature root area; [2] xylem/root area; [3] Area per vessel; [4] # vessels/root diameter; [5] Air space/root area. [C] Longitudinal section of young root tip from 7-day-old seedlings. The cortex layer is highlighted in yellow. The red arrow indicates the additional cell layer.

Figure 4. Effects of NaCl stress on *Schrenkiella parvula* shoot anatomy. Eight-week-old *S. parvula* plants were grown under control or 150 mM NaCl conditions for an additional 4 weeks. Transverse section of [A] stem and [B] leaf of 12-week-old hydroponically-grown *S. parvula* plants. A total of 23 sections from 7 control plants and 32 sections from 9 treated plants were used for leaf measurements, and a minimum of 20 sections from 5 to 11 plants were used for stem measurements. Data points represent individual cross-sections and colors represent individual plants. [C] Nuclear DNA content, distribution of leaf cell ploidy, and endoreduplication index of leaf cells from 10th and 11th leaves from the shoot meristem in control and 250 mM NaCl-treated 8-week-old *S. parvula* plants. Data are mean ± SD (n= 4). Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their respective control group, determined by Student's *t*-test. Representative cross-sections were obtained from the control plants Scale bars represent 100 μm.

Figure S6. Stem anatomy of *Schrenkiella parvula*. [A] Transverse section of *S. parvula* stem (between 4th, 5th, and 6th internodes from the shoot meristem) with cell layers marked in yellow lines for trait quantification. [B] Measurements of stem anatomical features indicated in [A]. A minimum of 20 sections from 5-11 plants were used for each measurement. Asterisks indicate significant difference ($p \le 0.05$) between the treated samples and its respective control samples, determined by Student's *t*-test. Data points represent individual cross-sections and colors represent individual plants.

Figure S7. Leaf structures of *Schrenkiella parvula*. [A] Transverse section of *S. parvula* leaves. [1] leaf thickness and [2] midrib thickness. [B] Transverse sections of *S. parvula* leaves treated with indicated concentration of salts. All sections represent the fifth or sixth leaf from the root-shoot junction in 8-week-old plants that were treated for 4 weeks.

Figure 5. Effects of NaCl stress on *Schrenkiella parvula* leaf traits. All experiments were performed with 4-week-old hydroponically-grown plants that were treated for an additional 4 weeks with the indicated NaCl concentrations. [A] Total leaf area, [B] Stomatal density, [C] Stomatal conductance, [D] Photosynthesis rate, [E] Leaf relative surface temperature, and [F] Leaf relative water content. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their respective control samples, determined by Student's *t*-test. Data are means ± SD (n ≥ 3). Open circles indicate the measurement from each plant (in B and F) and leaves (in C and D) for each experiment. DAT, days after treatment.

Figure S8. Effects of salt on growth of 4-week-old *Arabidopsis thaliana* and *Schrenkiella parvula* treated for an additional 2 weeks at 22 °C under a 12 hr light/12 hr dark photoperiod in soil.

Figure S9. Effects of NaCl stress on *Arabidopsis thaliana* and *Schrenkiella parvula* morphometric and physiological traits. Soil-grown, 8-day-old *A. thaliana* and 12-day-old *S. parvula* were exposed to the indicated salt concentrations at 21 °C, 16 h light/8 h dark photoperiod. The Measurements were taken on day 8, 12, and 15 after the start of the salt treatments. [A] Compactness (the ratio between the rosette (or leaf) area and the rosette (or leaf) convex hull area); [B] Leaf area; [C] Leaf water content; [D] Maximum quantum efficiency of PSII; and [E] Non-photochemical quenching were measured/estimated using the Qubit/PSI PlantScreen™ Compact System. For leaf area and leaf water content, values are percent of control (0 mM NaCl) for each time point. Letters denote significant changes at *p*-adj ≤ 0.05 (one-way ANOVA with post-hoc Tukey's test). Data are mean ± SD (n = 10). Open circles indicate the individual measurements obtained from each plant. DAT, days after treatment.

Figure S10. Leaf surface and the basal expression of genes involved in wax biosynthesis in *Arabidopsis thaliana* and *Schrenkiella parvula*. [A] Scanning electron micrographs contrasting *A. thaliana* and *S. parvula* leaf surfaces. [B] Major wax biosynthesis pathway and [C] wax biosynthesis genes that exhibited significantly different basal expression between *A. thaliana* and *S. parvula*.

Figure S11. The potential influence of phytochrome family of genes on *Schrenkiella parvula* leaf and growth form. [A] Elongated internode and leaf petiole in *S. parvula* compared to *A. thaliana*. [B] *PHYD* is absent in the *S. parvula* genome as illustrated using OrthNet representing evolutionary histories of orthologous gene groups derived from six Brassicaceae genomes: *Arabidopsis lyrata* (Aly, version 1.0), *Arabidopsis thaliana* (Ath, v. 'TAIR10'), *Capsella rubella* (Cru, v. 1.0), and *Eutrema salsugineum* (Esa, v. 1.0), *Sisymbrium irio* (Sir, v. 0.2), and *Schrenkiella parvula* (v. 2.0). Nodes are color-coded according to the species. Edges show either co-linear (cl) or transposed (tr) properties.

Figure 6.

Figure 6. Effects of NaCl on reproductive traits of *Schrenkiella parvula*. [A] Plants grown under control conditions generally flower at the ~17th leaf stage and the first few flowers are subsequently aborted. [B] Salt treated plants flower earlier at the ~14th leaf stage and the first flowers develop into mature siliques. [C] Days from planting to the first observed flower of plants grown under a long day (16 hr light/8 hr dark) or short day (12 hr light/12 hr dark) photoperiod with and without the indicated salt treatment. Center line - median; box - interquartile range (IQR); notch - 1.58 × IQR/sqrt(n); whiskers - 1.5 × IQR. [D] Flowers obtained from control and treated plants described in panel A and B. [E] Number of flowers and siliques per plant under control and salt treatment. Data are mean ± SD. Open circles represent individual measurements from four biological replicates. [F] Ratio between the number of flowers and siliques observed for control (C) and salt-treated (T) plants described in E. [G] Siliques from plants under control and NaCl treated conditions. [H] Size of siliques from G. Open circles represent individual silique counts. [I] Comparison of the area of seeds from plants under control and NaCl treated conditions. Data are mean ± SD (n = 50). Seeds per condition were selected from the seed pool of 15 to 50 plants. Open circles represent individual seeds. For panel [B], salt treatment started 21 DAP, initially at 50 mM NaCl given every other day. The salt concentration was increased by 50 mM every four days until it reached 200 mM; for panels [C, E, F], salt treatment was applied to 3-week-old plants until the end of the experiment; for panels [G, H, I], salt treatment was applied to 5-week-old plants for an additional 2 weeks. Different letters or asterisks represent significant differences (*, $p <$ 0.05) compared to control, determined by either one-way ANOVA followed by Tukey's post-hoc test [C] or Student's *t*-test [E, H, I]. DAP, days after planting.

Figure S12. Effects of KCl on reproductive traits of *Schrenkiella parvula*. [A] Plants grown under control conditions generally flower late and the first few flowers are subsequently aborted. KCl-treated plants flower earlier and the first flowers develop into mature siliques. [B] Early flowers in KCl treated plants develop longer filaments compared to control plants. DAP, days after planting.

Figure 7. Effects of salt stress on the germination of *Arabidopsis thaliana* and *Schrenkiella parvula* seeds. [A] Final percent germination recorded at 3 days and [B] germination curves of seeds treated with different concentrations of NaCl recorded for 12 days after stratification. [C] Germination curves of NaCl-treated seeds transferred to control (no NaCl) media, after 2 days or 30 days on salt plates. The germination curves were performed by direct sowing of seeds onto salt plates prior to stratification. Black arrows point to the day when seeds were transferred to non-NaCl plates. [D, E, F] Final percent germination of seeds treated with KCl [D], LiCl [E] or H_3BO_3 [F]. Data are mean ± SD (n = 3-4, each replicate contains ca. 50 seeds). Open circles indicate individual measurements. Different letters indicate significant difference ($p \leq 0.05$) determined by one-way ANOVA with post-hoc Tukey's test across all conditions in [A, D, E, F], or across treatment groups at 3 and 12, and 7 days after stratification in [B] and [C] respectively.

Figure S13. Effect of salt on *Arabidopsis thaliana* and *Schrenkiella parvula* inducing barriers for seed germination and seedling establishment. [A] *S. parvula* (*Sp*) and *A. thaliana* (*At*) seed germination and growth on NaCl- and KCl-supplemented MS medium. Scale bars are 2 cm. [B] Growth of ungerminated *S. parvula* seeds treated with different concentrations of NaCl (as shown in Figure 7) after transferring to control MS medium.

Figure S14. Survival of *Schrenkiella parvula* in the climate conditions found in Lake Tuz region. [A] Lake Tuz location from Google Earth; [B] Precipitation and temperature recorded in Konya, Turkey (NOAA, 2019). *S. parvula* growing season is from April/May to August/September (Tug et al., 2019). Bars indicate precipitation and lines represent temperature. [C] Average day length recorded in Konya per month. Red line indicates the day length used for typical laboratory growth assays for *S. parvula*. [D-F] Survival rate of *A. thaliana* and *S. parvula* seedlings under [D] freezing stress, [E] chilling stress, and [F] heat stress. Five-day-old seedlings were subjected to temperature treatments and data were recorded 5 days after recovery. Asterisks indicate significant difference ($p \leq 0.05$) between the treated samples and their respective control samples (student's *t*-test). Data are mean ± SD (n = 4). Each replicate plate contained 3-6 plants. Open circles indicate the biological replicates.

Figure 8. Summary of salt-induced structural and physiological traits in *Schrenkiella parvula*. [**A**] Salt- induced changes in traits quantified in the current study. Changes in each trait were calculated as the fold change between treatment and control measurements. Traits highlighted in the red boxes showed significant differences ($p < 0.05$) under salt treatments compared to the control determined in previous assays (Fig 2-7). [**B**] and [**C**] PCA biplot of traits quantified for *S. parvula* under control and salt-treated conditions. Arrows indicate directions of loadings for each trait and are color-coded by contribution to the variations in PC1 and PC2. [B] Anatomical traits quantified: LT, leaf thickness; MT, midrib thickness; XS, stem xylem to stem area; SVA, average area per vessel in stems; CbS, cambium to stem area; CtS, cortex to stem area; XR, root xylem to root area; RVA, average area per vessel in roots; VN, number of vessels across the root diameter; AS, air space to root area; YRA, young root area; CT, cortical thickness; YRD, root diameter in young roots; DRH, distance from root tip to first root hair. [C] Physiological traits quantified: AbS, abaxial stomatal density; adS, Adaxial stomatal density; LA, leaf area; SLA, silique area; SA, seed area; PRL, primary root length; LRD, lateral root density; RWC, relative water content; CA, CO2 assimilation; SC, stomatal conductance; RHL, root hair length; NF, number of flowers; NS, number of siliques. DAT, days after treatment; DAP, days after planting.