1 **Title:** Temporal scaling of dopamine neuron firing and dopamine release by distinct ion channels
2 shape behavior
3

Authors
Barbara Juarez[1,2,+], Mi-Seon Kong[1], Yong S. Jo[1,§], Jordan E. Elum[1], Joshua X. Yee[1],

Scott Ng-Evans[1], Marcella Cline[1], Avery C. Hunker[2,†], Meagan A. Quinlan[1,†], Madison A.

Baird[2,‡], Abigail J. Elerding[2], Mia Johnson[1], Derek Ban[1], Adriana Mendez[1%], Nastacia L.

Goodwin[3], Marta E. Soden[2], Larry S. Zweifel*[1, 2].

Affiliations

[1]Department of Psychiatry and Behavioral Sciences, University of Washington; Seattle, WA,

United States.

[2] Department of Pharmacology, University of Washington; Seattle, WA, United States.

[3] Department of Biostructure, University of Washington; Seattle, WA, United States.

+ Current affiliation: University of Maryland, Baltimore; Baltimore, MD. United States.

§ Current affiliation: School of Psychology, Korea University; Seoul, Republic of Korea.

†Current affiliation: Allen Brain Institute; Seattle, WA, United States.

‡ Current affiliation: Cajal Neuroscience; Seattle, WA, United States

% Current affiliation: Icahn School of Medicine at Mount Sinai, New York, NY, United States

*Corresponding author. Email: larryz@uw.edu

20
21

Abstract

Despite the widely known role of dopamine in reinforcement learning, how the patterns of dopamine release that are critical to the acquisition, performance, and extinction of conditioned responses are generated is poorly resolved. Here, we demonstrate that the coordinated actions of two ion channels, Kv4.3 and BKCa1.1, control the pattern of dopamine neuron firing and dopamine release on different time scales to regulate separate phases of reinforced behavior and extinction in mice. Inactivation of Kv4.3 in VTA dopamine neurons increases *ex vivo* pacemaker activity and excitability that is associated with decreased rates of bursting and increased *in vivo* ramping dynamics prior to lever press in a learned instrumental response paradigm. Loss of Kv4.3 enhances performance of the learned response and facilitates extinction. In contrast, loss of BKCa1.1 increases duration of burst firing and phasic dopamine release that enhances learning of an instrumental response. Inactivation of BKCa1.1 also enhances extinction burst lever pressing in early extinction training that is associated with a greater change in activity between reinforced and unreinforced actions. These data demonstrate that disruption of intrinsic regulators of neuronal activity differentially impacts dopamine dynamics observed during reinforcement learning and extinction behaviors.

Teaser

Ion channels in midbrain dopamine neurons are critical for patterning action potential firing, dopamine release, and learning.

MAIN TEXT

Introduction

Reinforcement and extinction learning are essential processes by which an organism links environmental stimuli and actions with outcomes. Dopamine-producing neurons in the ventral

tegmental area (VTA) of the midbrain facilitate reinforcement learning to promote the

performance of goal-oriented behavior through modulation of information processing in the

forebrain, most notably in the nucleus accumbens (NAc) (*1*). Distinct patterns of dopamine release,

such as tonic, phasic (burst), or ramping patters, that occur on different time scales during the

acquisition, performance, and extinction of a conditioned response are thought to underlie

dopamine function within these contexts(*2–6*). Despite the evidence that dopamine is critical for

reinforcement learning(*1–3*), how the intrinsic properties of the cell contribute to patterning action

potential firing and neurotransmitter release during distinct phases of reinforcement learning is not

well resolved. Moreover, recent findings suggest that dopamine release dynamics can occur

independently of the activity patterns at the cell body(*5*) raising the possibility that the firing

properties of dopamine neurons at the level of the cell body may be dispensable for some aspects

of reinforcement learning.

VTA dopamine neurons display spontaneous action potential firing in the absence of

afferent inputs(*7*) that is regulated by a suite of voltage-gated ion channels (*8–10*). Potassium

currents shape the action potential waveform to regulate the frequency and pattern of firing (*8–12*).

Kv4.3 subunits contribute to A-type potassium currents (I_A) in dopamine neurons that facilitate

repolarization and control the frequency of firing (*9*). BKCa1.1 subunits are the essential subunits

of voltage-sensitive, calcium-activated big conductance potassium (BK) channels, which conduct

BK currents (I_{BK}). These subunits are expressed in dopamine neurons and conduct currents that

mediate the initial phase of the action potential afterhyperpolarization to control the pattern of

firing (*10*). Based on the recruitment of these channels during different phases of the action

potential repolarization and afterhyperpolarization, we hypothesized that targeted disruption of

their function in VTA dopamine neurons would differentially impact action potential firing

patterns and neurotransmitter release. Specifically, loss of Kv4.3 would induce more pacemaker-

like activity and loss of BKCa1.1 would induce more irregular/burst patterns of activity. Thus,

inactivation of these channels provides a means to further resolve whether intrinsic regulators of

dopamine neuron action potentials influence activity patterns at the level of the cell body and

neurotransmitter release dynamics in terminal fields during reinforcement learning.

Using CRISPR/Cas9 gene-editing technologies, we selectively targeted mutagenesis of

Kv4.3 (gene name: *Kcnd3*) and BKCa1.1 (gene name: *Kcnma1*) in dopamine releasing neurons of

the VTA. We found that inactivation of Kv4.3 and BKCa1.1 differentially affected the action

potential waveform, neuronal excitability, and patterns of action potential firing. We further show

that loss of function (LOF) of these two channels differentially impacted the acquisition,

maintenance, and extinction of a reinforced behavioral response that was associated with distinct

effects on the *in vivo* patterns of cell body activity and dopamine release during these behaviors.

Specifically, loss of BKCa1.1 ion channel subunits in VTA dopamine neurons enhanced burst

firing and phasic dopamine release that was associated with enhanced acquisition of instrumental

behavior and a heightened extinction burst behaviors. In contrast, loss of Kv4.3 increased

dopamine neuron excitability, decreased burst rates, and increased ramping that was associated

with increased performance of a learned response. Further, loss of Kv4.3 also led to facilitated

extinction of an instrumental response that was associated with a lack of differential dopamine

neuron responses between actions associated with rewarded or omitted outcomes. These findings

demonstrate that intrinsic regulators influence patterns of dopamine cell body activity and terminal

release dynamics *in vivo* to shape control the behavior during reinforcement and extinction

learning.

Results

CRISPR/Cas9 inactivation of Kv4.3 and BKCa1.1

To determine the distribution of *Kcnd3* and *Kcnma1* expression in the VTA, we performed

in situ hybridization (RNAscope) to quantify the mRNA levels for these ion channel subunits and

the rate-limiting enzyme in dopamine production, tyrosine hydroxylase (*Th)*. We find that Kv4.3

and BKCa1.1 subunits are expressed in most dopamine neurons along the rostral-caudal axis of

the VTA (Fig. 1, A-C, fig. S1, A-E).

To selectively inactivate these two channels, we used a viral-mediated, Cre-inducible

CRISPR/SaCas9 system (*13*). Adeno-associated viruses (AAV) containing single guide RNA

directed to either of the two loci (AAV1-FLEX-SaCas9-sg*Kcnd3* or AAV1-FLEX-SaCas9-

sg*Kcnma1*) were generated (Fig. 1, D and H) and injected into the VTA of separate groups of adult

Slc6a3(DAT)-IRES-Cre mice (fig. S2, A; fig. S3-S4). Control DAT-IRES-Cre mice received

injections of AAV1-FLEX-eYFP into the VTA. We genetically validated gene mutagenesis of

targeted sequences by co-injection with AAV-FLEX-KASH-GFP into the VTA and harvesting

tissue after four weeks. We found that the AAV mediated Cre-inducible CRISPR/SaCa9 system

yielded high efficiency indel formation (Fig. 1, E, I; fig S2 B-C).

Next, we used patch-clamp electrophysiology to determine impact of CRISPR/SaCas9

induced Kv4.3 or BKCa1.1 gene mutagenesis on channel activity in VTA dopamine neurons. To

visualize cells, we co-injected the LOF mice with AAV-FLEX-eYFP. First, we quantified I_A in

VTA dopamine neurons in control mice and mice that received AAV1-FLEX-SaCas9-sg*Kcnd3*

(Fig. 1, F-G). In control mice, a depolarization protocol induced robust I_A, quantified as the

difference between the instant and sustained parts of the generated current; this was nearly

abolished after bath application of the I_A inhibitor, AmmTx3, suggesting the presence of I_A (Fig.

1, F-G). In mice that received AAV1-FLEX-SaCas9-sg*Kcnd3* in VTA dopamine neurons*,* we

observed almost no I_A in baseline conditions and insensitivity to AmmTx3 (Fig. 1, F-G),

suggesting Kv4.3 LOF. We next quantified I_{BK}, a major component of I_K generated after

depolarization step, in VTA dopamine neurons of in control mice and mice that received AAV1-

FLEX-SaCas9-sg*Kcnma1* (Fig. 1, J-K). In control mice, iberiotoxin, a BK channel-specific

inhibitor, significantly reduced maximal peak of I_K when compared to I_K before iberiotoxin bath

application, suggesting pharmacological blockade of the BK channel current component in

dopamine neurons (Fig. 1, J-K). However, in VTA dopamine neurons from AAV1-FLEX-SaCas9-

sg*Kcnma1* injected mice, there was already a significant reduction in I_K under baseline conditions

that was not changed after iberiotoxin application, suggesting that there is no iberiotoxin-sensitive

BK channel current component following CRISPR injection (Fig. 1, J-K). Using current

subtraction methods, we further quantified the specific I_{BK} and found that AAV1-FLEX-SaCas9-

sg*Kcnma1* significantly reduced I_{BK} (fig. S4, A-C).

It is possible that CRISPR/SaCas9-induced LOF of Kv4.3 or BKCa1.1 channels could

impact the expression of related potassium channel subunits to compensate for their loss. *Kcnd2*

(Kv4.2), another A-type potassium channel subunit, is also expressed in VTA dopamine neurons

in mice (*14*). Mutagenesis of *Kcnd3* did not alter midbrain expression of this subunit (fig. S2, D).

Previous studies have found no other pore forming alpha subunits for BK channels in the VTA.

However, small-conducting calcium activated potassium channels (SK) are expressed in these

cells and contribute to the after hyperpolarization (*15*). We found no differences in *Kcnn3* (SK3)

expression in the midbrain following *Kcnma1* mutagenesis (fig. S2, E). These data suggest that

LOF of either Kv4.3 and BKCa1.1 has significant and specific impacts on their respective currents

without inducing compensatory changes in other related channels.

Kv4.3 and BKCa1.1 regulate *ex vivo* and *in vivo* action potential firing patterns but not *ex vivo* dopamine release from terminals

Next, we profiled the impacts of ion channel subunit LOF on *ex vivo* activity characteristics of VTA dopamine neurons. As predicted, CRISPR/Cas9 induced Kv4.3 LOF altered the dopamine neuron action potential waveform by increasing the action potential half width and slowing repolarization relative to control of spontaneously active action potentials in VTA slices (Fig. 2, A-D). BKCa1.1 LOF cells showed significantly more depolarized afterhyperpolarization compared to control cells (Fig. 2, A-B, E). Dopamine neurons from BKCa1.1 LOF also showed a more depolarized resting membrane potential relative to control cells (Fig. 2, A-B, F). Both Kv4.3 LOF and BKCa1.1 LOF in VTA dopamine neurons significantly depolarized the action potential threshold when compared to control neurons, but neither manipulation affected the peak of the action potential (fig. S4, H-I). These cell-specific CRISPR/Cas9 ion channel subunit LOF changes were different compared to the less specific pharmacological blockade of Kv4 and BK channels, supporting the use of CRISPR for this study (fig. S4, E-G). To determine if the changes we observed in action potential shape from spontaneously action potentials in Kv4.3 LOF and BKCa1.1 LOF dopamine neurons were impacted by the depolarized resting membrane potential, we clamped the neurons at -60 mV and injected current to induce action potential firing (fig S4, J). We found similar effects on action potential shape, although notably no significant change in the anti-peak of the afterhyperpolarization in BKCa1.1 LOF neurons was observed, though the shape of the afterhyperpolarization was affected (fig S4, J-N).

Given the changes in action potential waveform dynamics associated with Kv4.3 LOF and BKCa1.1 LOF, we sought to establish whether these perturbations affect neuronal excitability and patterns of action potential firing. Input-output analysis (excitability) of action potential firing

following depolarizing current injection from cells held at -60 mV (Fig. 2, G-H) and cells at resting

membrane potential (fig. S4, O) revealed that Kv4.3 LOF in VTA dopamine neurons increased

neuronal excitability when compared to control cells. Further, the frequency of spontaneous action

potential firing in Kv4.3 LOF cells is greater relative to control cells (Fig. 2, I-J; fig. S4, P). In

contrast, BKCa1.1 LOF did not affect these parameters (Fig. 2 G-J; S4, O). However, BKCa1.1

LOF did shift the pacemaker-like *ex vivo* activity observed in control and Kv4.3 LOF cells to a

more irregular pattern, as evidenced by an increase in the coefficient of variation in the inter-spike

interval (CV-ISI) (Fig. 2, I-K; fig. S4, P).

Increased CV-ISI in VTA dopamine neurons of BKCa1.1 LOF mice is consistent with a

shift to more burst-like activity (*11*); however, this pattern of firing by dopamine neurons is most

accurately assessed by recording neural activity *in vivo*. To identify putative dopamine neurons

during *in vivo* recording under baseline conditions or during behavior from mice that received

control virus or CRISPR viruses for Kv4.3 LOF or BKCa1.1 LOF, we combined CRISPR/Cas9

mutagenesis with cell-type specific expression of channelrhodopsin (ChR2), *in vivo* single-unit

electrophysiology using optetrode microdrives implanted in the VTA (fig. S5, A-C) (*11,16*). To

optogenetically identify putative dopamine neurons, ChR2 was optically stimulated using 10

pulses of blue light at a frequency of 20 Hz repeated 20 times at the end of a recording session; if

action potential activity was induced in a time-locked manner, it was categorized as an optically-

sensitive putative dopamine neuron (*11*, *16*). We first analyzed the baseline properties of putative

dopamine neurons to understand how VTA dopamine activity is altered after CRISPR/Cas9 LOF.

Overall firing rate did not differ between the cell groups (fig. S5, D). The number of burst events

per second (burst rate) and the percentage of spikes fired in bursts was similar between control and

BKCa1.1 LOF cells but was significantly reduced in Kv4.3 LOF cells relative to controls (Fig. 2L;

fig. S5, E). In contrast, the duration of the burst events and the number of spikes fired in a burst were significantly increased in BKCa1.1 LOF mice relative to controls but were unaffected in Kv4.3 LOF cells (Fig. 2, M; fig. S5, F). Together, these findings indicate that Kv4.3 LOF in VTA dopamine neurons reduces burst activity while BKCa1.1 LOF increases burst duration when burst activity is induced.

In addition to regulating action potentials at the cell body, potassium channels also regulate neurotransmitter release (17, 18). To address whether Kv4.3 or BKCa1.1 LOF function impacts terminally evoked dopamine release dynamics independent from cell body firing, we injected CRISPR/SaCas9 viruses into the VTA along with the redlight-shifted opsin Chrimson (19) and injected the genetically encoded dopamine sensor dLight1.3b (5) into the NAc (Fig. 2, N). After at least 4 weeks, we performed *ex vivo* 2-photon slice imaging of optically evoked dopamine release in acute brain slices containing the NAc, where VTA dopamine cell bodies are not present. This *ex vivo* preparation helped us determine if there are any terminal-specific channel subunit LOF effects on dopamine release, dissociating it from impacts of cell body LOF. VTA dopamine terminals present in the NAc that express Chrimson were optically stimulated (1 s) at different frequencies (1, 5, 10, and 20 Hz) while imaging dLight fluorescence (Fig. 2, N). We did not detect differences between control and Kv4.3 LOF or BKCa1.1 LOF mice (Fig. 2, O-P). These findings suggest that CRISPR induced Kv4.3 LOF and BKCa1.1 LOF have primary effects at the cell body.

Kv4.3 and BKCa1.1 differentially regulate instrumental behavior

To assess whether Kv4.3 and BKCa1.1 LOF impacts reinforcement learning, we first trained mice in an appetitive Pavlovian conditioning paradigm where a lever extension followed by retraction (10 s, CS) signals reward (Rew) delivery (Fig. 3A). We did not observe behavioral differences between Kv4.3 LOF and BKCa1.1 LOF mice relative to controls during the CS

presentation or ITI period (Fig. 3, B-C). Learning, as assessed by the conditioned approach (CS

head entry rate minus ITI head entry rate) was also not different between controls and LOF groups

(Fig. 3, D).

Following Pavlovian conditioning, mice were transitioned to a fixed-ratio (FR1) schedule

of operant reinforcement learning in which lever retraction and reward delivery were contingent

on a lever press action (LP) (Fig. 3, E). BKCa1.1 LOF learned the contingency faster than control

mice on day 1 of training (Fig. 3, F, G left). Control and Kv4.3 LOF mice increased responding

between the first and second day that remained elevated on the third day, with no differences in

total lever presses between them (Fig. 3, F, G left). However, Kv4.3 LOF mice had a higher initial

rate of responding than control mice on the third day, once the task was acquired (Fig. 3, G, right).

Analysis of lever press ethograms during the early (first 30 minutes) or late (last 30 minutes) halves

of the session shows the known distribution of individual variability in LP behavior in control mice

on the first day (Fig. 3, H, J). We also determined inter-press interval timing across groups (Fig.

3, I, K). This all revealed differences in the pattern of lever pressing between the control and LOF

groups (Fig. 3, H-K). BKCa1.1 LOF mice started pressing more quickly with a shorter inter-press

interval on the first day when compared to controls, consistent with the faster learning of the

contingency (Fig. 3, H-I). Once the contingency was learned, control and BKCa1.1 LOF mice had

a relatively stable level of responding on the third day; however, Kv4.3 LOF mice showed a high

level of responding in the first half of the session that diminished in the second half when compared

to controls (Fig. 3, G right, J-K), suggesting an enhanced invigorated performance of the task.

Differences in reinforcement learning were not associated with changes in overall

locomotor activity (fig. S6, A-B). To determine if Kv4.3 LOF or BKCa1.1 LOF can result in

alterations in baseline anxiety, we performed an open field test and elevated plus maze and found

no differences between controls and CRISPR LOF groups (fig. S6, C-F). To assess whether the

differences in operant responding reflected altered motivation, we assayed mice in a progressive

ratio schedule of reinforcement. We did not observe differences between the groups in this task

(fig. S6, G).

Kv4.3 and BKCa1.1 differentially regulate behaviorally-evoked firing patterns and

dopamine release

Differential regulation of action potential waveforms, intrinsic excitability, *in vivo* cell

body activity patterns at baseline, and patterns of instrumental behavior suggest that Kv4.3 LOF

and BKCa1.1 LOF can cause distinct changes in the encoding of reward-related behaviors. To test

this, we performed *in vivo* electrophysiology during the instrumental conditioning task with optical

and pharmacological identification of VTA dopamine neurons (Fig. 4, A-B; fig. S5, C) following

Pavlovian conditioning as above. Due to the restrictive nature of the tethering for recording, all

tethered mice underwent 5 days of FR1 to reach similar behavioral performances as untethered

mice and cell body activity was measured on day 1 and day 5 (fig. S6, H-J). Recordings were

performed throughout the entire behavioral session which was followed by delivery of blue light

for optical identification and injection of quinpirole, a dopamine D2 receptor agonist, to identify

putative dopamine neurons. Although it is difficult to glean much information regarding the action

potential waveform from extracellular recordings because of the unknown distance from the

electrode and the recorded cells, we did observe qualitative differences in the waveforms of light-

sensitive cells from Kv4.3 LOF and BKCa1.1 LOF mice relative to cells from controls (fig. S5,

G) that were similar to those observed in whole cell recordings. The overall firing rates of

individual cells surrounding the lever press and reward delivery period were variable, but clear

differences in the patterns of activity were evident (fig. S7, A-D). In lever press responsive cells,

the mean firing rate during the baseline period prior to lever press did not differ on the first day of

FR1 reinforced learning (fig. S7, A) but was significantly elevated in Kv4.3 LOF mice relative to

controls on the fifth day (fig. S7, D). To account for the overall variance in firing rate, cell activity

was normalized using a Z-score (Fig. 4, C). Analysis of the area under the curve (AUC) of the Z-

score following lever press revealed significantly larger responses to the lever press in BKCa1.1

LOF cells relative to controls on both days (Fig. 4, D). In contrast, we observed significantly

reduced responses in Kv4.3 LOF mice relative to controls on day 5 following reward delivery (Fig.

4, E).

To determine whether changes in behaviorally-evoked action potential firing at the cell

body associated with ion channel LOF is related to dopamine release dynamics downstream, we

expressed dLight1.3b in the NAc of Kv4.3 LOF, BKCa1.1 LOF, and control mice (fig. S8) and

performed fiber photometry during FR1 following Pavlovian conditioning as above. Again,

tethered mice underwent 5 days of FR1 instead of 3 days (fig. S6 H-J) and dopamine signals were

measured on day 1 and day 5. Control mice showed phasic increases in dopamine time-locked to

LP and Rew on day 1 (Fig. 5, A top row) and day 5 (Fig. 5, A middle row, bottom row). Kv4.3

LOF mice showed LP and Rew dopamine signals on the first day (Fig. 5, A top row) and day 5

(Fig. 5, A middle row, bottom row heat maps). BKCa1.1 LOF mice had large dopamine release

events to LP and Rew on day 1 (Fig. 5, A top row) and day 5 (Fig. 5, A middle row, bottom row).

Similar to cell body firing, BKCa1.1 LOF mice had larger dLight1.3b signals during LP on day 1

when compared to controls (Fig. 5, B). Rew-evoked signals were also significantly higher in

BKCa1.1 LOF mice on both days compared to controls (Fig. 5, C). Although phasic increases were

more prominent in BKCa1.1 LOF mice, we found ramping of the dLight1.3b signal prior to the

lever press on day 5 was increased in Kv4.3 LOF mice relative to controls. (Fig. 5, D-E). This was

also evident in the firing rates of Kv4.3 LOF mice relative to controls (fig. S7, E).

Kv4.3 and BKCa1.1 LOF differentially regulates extinction of an operant response

Following Pavlovian and FR1 conditioning, a subset of mice underwent extinction training

with reward omission (Fig. 6, A). Control mice exhibited an extinction burst in lever pressing on

the first day of extinction (higher pressing on day 1 of extinction versus the last day of reinforced

FR1 conditioning). This extinction burst was elevated in BKCa1.1 LOF mice relative to control

mice (Fig. 6, B-C; fig. S9, A-C). In contrast, Kv4.3 LOF mice displayed potentiated extinction, as

evidenced by the significantly reduced lever presses during the first day of extinction training

relative to control mice (Fig. 6, B-C; fig. S9, A-C). These differences in extinction burst behaviors

are also represented using a ratio of lever presses on the first day of extinction compared to the last

day of reinforced FR1; ratios greater than indicate an induction of an extinction burst, as observed

in control mice and BKCa1.1 mice, to a greater degree (fig. S9, G). Notably, Kv4.3 LOF mice do

not have this difference (fig. S9, G). The number of lever presses in BKCa1.1 LOF and control

mice decreased during the extinction session but remained relatively stable in Kv4.3 LOF mice

(Fig. 6, C; fig S9, A-B). All mice performed similarly on the last day of extinction (Fig. 6, B; fig.

S9, D-F).

We were intrigued by the observation that BKCa1.1 mice showed an elevated extinction

burst and that Kv4.3 mice showed a reduced extinction burst relative to controls. Because BKCa1.1

mice have faster acquisition of the FR1 instrumental response, we asked whether there is a

relationship between responding on day 1 of acquisition and day 1 of extinction. We found a

significant positive correlation between the level of responding on acquisition day 1 and extinction

day 1, suggesting that mice that learn the task faster have the largest extinction burst responses with reward is omitted (fig. S9, H).

The altered extinction behaviors observed in Kv4.3 LOF and BKCa1.1 LOF mice may be due to differences in dopamine neuron activity and release between actions of reinforced and omitted outcomes. To explore this further, we compared the normalized action potential firing (Z-score) and dopamine release patterns during the lever press period of the mice during the first day of learning in the FR1 reinforced conditioning presented in Fig. 4 and Fig. 5 and the first day of FR1 extinction (reward omitted) (Fig. 6, D). All groups showed general reductions in putative dopamine neuron action potential firing and dlight1.3b signals during extinction. In control mice normalized firing was reduced, but we observed no significant differences in cell body activity (Fig. 6, E left). However, dLigh1.3b signals in the NAc of control mice on Extinction day 1 were significantly reduced when compared to signals from reinforced FR1 learning (Fig. 6, E right). Kv4.3 LOF mice, which display control-like acquisition of the reinforced task and reduced lever pressing during, showed no differences between days in VTA dopamine cell body activity and dLight1.3b signals in the NAc (Fig. 6, F). Of note, we found that the overall firing rate of lever press responsive cells in the Kv4.3 LOF mice during the pre-lever press period was elevated relative to control mice (fig. S10, A-B), and the overall pattern of activity was remarkably similar to what we observed on the first day of conditioning (compare fig. S7 to fig. S10). BKCa1.1 LOF mice, which have potentiated learning of reinforced task and a larger extinction burst, displayed significant reductions in both VTA dopamine cell body activity and dLight1.3b signals in the NAc on the first day of extinction training when compared to FR1 reinforced Day 1 (Fig. 6, G).

Discussion

Our results show that LOF of ion channels that contribute to the generation of the action

potentials alters the patterns of dopamine neuron firing and dopamine release to influence

reinforcement learning, performance, and extinction. Using viral delivery of CRISPR/SaCas9 we

were able to generate high-efficiency mutagenesis that allowed us to determine the cell-type

specific function of Kv4.3 and BKCa1.1 subunits in VTA dopamine neurons of the adult brain.

We found that CRISPR/SaCas9 mutagenesis of *Kcnd3* (Kv4.3 LOF) effectively reduced the

AmmTx-sensitive A-type potassium current and mutagenesis of *Kcnma1* (BKCa1.1 LOF)

significantly attenuated the iberiotoxin-sensitive BK current. Loss of these channels altered the

action potential waveform by broadening the action potential half-width (Kv4.3 LOF) and

reducing the AHP (BKCa1.1 LOF). Reduced AHP in VTA dopamine neurons following loss of

BKCa1.1 is distinct from the effects of inhibition of BK current in substantia nigra pars

compacta (SNc) dopamine neurons where a paradoxical increase in afterhyperpolarization was

reported (*10*). This could be due in part to the higher expression of the small conductance

potassium channel, SK3 in SNc dopamine neurons (*20*) that also contributes to the

afterhyperpolarization (*15*), or the reciprocal interactions between Kv2 channels and BK

channels in SNc dopamine neurons (*10*).

Kv4.3 LOF resulted in an increase in VTA dopamine neuron excitability and pacemaker

activity, as would be predicted based on the recruitment of A-type potassium channel

conductance during the rapid depolarization phase of the action potential to generate rapid

feedback and allow for slow pacemaker activity (*9*). Loss of BKCa1.1 resulted in irregular action

potential firing in VTA dopamine neurons *ex vivo* and prolonged burst events in these cells *in*

vivo. This *ex vivo* result again contrasts to the effects observed in SNc dopamine neurons, likely

relating to the differential effects on the AHP. Reduced AHP following BKCa1.1 LOF reported

here is similar to that observed following inhibition of BK in cerebellar Purkinje cells that also

increased burst firing (*21*). Based on these observations, we propose that BKCa1.1 channels

terminate burst events through regulation of the action potential afterhyperpolarization in VTA

dopamine neurons, and in their absence, phasic dopamine is enhanced once a burst is initiated,

which assigns a higher value to action. This facilitates the acquisition of the conditioned

response and establishes a greater differential response during extinction thus, generating a larger

extinction burst (*2*, *3*, *22*, *23*).

We find that Kv4.3 channels regulate firing frequency in acute brain slice preparations,

likely through the regulation of membrane repolarization (*9*). However, we did not observe

changes in baseline firing rate *in vivo*, likely due to the influence of GABAergic inhibition in the

intact VTA. Intriguingly, on the last day of FR1 conditioning and the first day of extinction

training, we observed significantly higher baseline firing rates in lever press responsive cells.

We hypothesize that this increased activity is a reflection of a reduction in the inhibitory tone on

these cells following learning, such as through disinhibition (*24*), that elevates tonic activity.

Indeed, it has previously been shown that when tonic signals such as ramping are elevated

performance of a learned response is enhanced (*3–6*). In contrast to previous reports that ramping

of dopamine release occurs independent of activity at the cell body (*5*), our data suggest that the

intrinsic excitability of the cell, regulated by Kv4.3, contributes at least partially to this process

in reinforced trials. During the first day of extinction training, Kv4.3 LOF mice had a lower

extinction behavior burst than control or BKCa1.1 LOF mice. We also observed that Kv4.3 LOF

did not display significant changes in dopamine release of action potential firing between the

first day of acquisition and the first day of extinction. In contrast, BKCa1.1 mice showed the

largest difference between these two days. This suggests that the level of extinction burst is

likely related to the perceived value of the action. BKCa1.1 mice had the largest phasic increase

to the lever press during acquisition and the smallest response during extinction; thus, generating

the largest difference between the reinforced and unreinforced actions. In contrast, Kv4.3 LOF

mice showed prominent ramps at the cell body and in terminal function but reduced phasic

responses that resulted in a loss of significant differences between the responses during

reinforced and unreinforced trials.

A limitation of the current study relating to the loss of function of Kv4.3 and BKCa1.1 is

the potential impact on conductance associated with other ion channels and extrinsic signaling.

Indeed, as mentioned above, BK currents can influence Kv2 currents, and vice versa in SNc

dopamine neurons (9). Because Kv4 channels are activated during rapid ramping of the

depolarization they help to constrain the inward sodium current; thus, their loss would be

predicted to result in a larger sodium conductance and higher frequency firing (9). In VTA

dopamine neurons a small subthreshold sodium current has also been shown to contribute to

pacemaker activity (23), reducing the resting potential to varying degrees following Kv4.3 or

BKCa1.1 mutagenesis could disrupt the subthreshold sodium conductance and alter the

excitability of VTA dopamine neurons or their pacemaker activity. These studies highlight the

critical interactions between ion channels that work collectively to regulate action potential

firing. Further, these ion channels and their respective currents have been previously identified to

have cooperative or opposing roles beyond voltage-gated intrinsic regulation. For example, I_A is

a critical effector of 2-AG modulation of excitability and I_{BK} has been demonstrated to couple to

ligand-gated glutamate receptors to regulate synaptic signaling (34, 35). Regardless of the

mechanism(s) that underlies the impact of Kv4.3 and BKCa1.1 LOF on VTA dopamine neuron

physiology, we find significant effects on the action potential waveform, action potential firing

both *ex vivo* and *in vivo*, and changes in dopamine release dynamics that supports the importance

of their cell autonomous function in these cells.

Although we focused on Kv4.3 and BKCa1.1 in dopamine neurons, these channels are

expressed widely throughout the brain where they likely serve similar functions in patterning

transmitter release. Further, there are numerous other ion channels that contribute to the action

potential. The emergence of many of these ion channels as hotspots of missense mutation in

neurodevelopmental disorders (*15*) reinforces the importance of understanding how these

channels influence cellular and systems function.

Materials and Methods
400
Experimental Design:

Mice: All procedures were approved and conducted in accordance with the guidelines of the

University of Washington's Institutional Animal Care and Use Committee (Protocol ID: 4249-

01). Mice were housed on a 12:12 light: dark cycle with *ad libitum* access to food and water,

except when undergoing food restriction for Pavlovian and operant behavioral conditioning.

Approximately equal numbers of male and female mice were used, starting at 6-10 weeks of age.

Mice were group-housed (2-5 mice per cage), except during night-day locomotion assays. Either

C57BL/6J (Jackson Laboratory for *in situ* hybridization with RNAscope) or *Slc6a3*-IRES-Cre

(DAT-IRES-Cre (Jackson Laboratory, Jax 006660, backcrossed to C57BL/6J) mice were used.

Viruses: All adeno-associated viruses (AAV) were produced in-house with titers of $1\text{-}3\mathrm{X}10^{12}$

particles mL^{-1}, as described (*25*). Cre-inducible CRISPR viruses: AAV1-CMV-FLEX-SaCas9-

HA-U6-sgKcnd3, AAV1- CMV-FLEX-SaCas9-HA-U6-sgKcnma1. Cre-inducible optogenetic

viruses: AAV1- EF1α-FLEX-ChR2-eYFP and AAV1- EF1α-FLEX-Chrimson-TdTomato.

Dopamine sensor virus: AAV1-CAG-dLight1.3B. Cre-inducible control and fluorescent virus:

AAV1-EF1α-FLEX-eYFP. Cre-inducible FACS virus: AAV1-hSyn1-FLEX-KASH-GFP.

CRISPR/Cas9 design: sgRNAs were designed using methods previously described (*13*, *26*). 21

bp sgRNAs for *Kcnd3* (Forward: GAGCGCTATCCCGACACCTTG; Reverse:

CAAGGTGTCGGGATAGCGCTC) and *Kcnma1* (Forward:

GTCTAGGCTGAGATGGTTCGC; Reverse: GCGAACCATCTCAGCCTAGAC) were

generated (Invitrogen) and cloned into pAAV-CMV-FLEX-SaCas9-U6sgRNA for generation of

AAVs in house.

Surgeries: DAT-IRES-Cre mice were 6-8 weeks of age at time of surgery, except for mice used

for *in vivo* electrophysiology which were at least 10 weeks of age at time of surgery. Mice were

anesthetized with isoflurane in a chamber and then placed and secured into the stereotax with

isoflurane delivery through a nose cone. For all implants, an anchoring screw was driven into the

skull away from the implant site. Mice received eYFP (control), CRISPR/eYFP (*in vitro*

electrophysiology or behavioral groups), CRISPR/KASH-GFP (sequencing), or CRISPR/ChR2

(*in vivo* electrophysiology) bilateral injections into the VTA (in mm, relative to bregma: A/P: -

3.3; M/L $\pm$ 0.5; D/V: -4.9 $\rightarrow$ -4.4), total volume 0.5 μL into each side. A/P coordinates were

adjusted for Bregma-Lambda distances using a correction factor of 4.21 mm. For D/V, the

syringe was lowered to the deepest indicated coordinate and then raised at the start of the

injection. For *ex vivo* two-photon dLight experiments, all mice received a bilateral injection of

dLight1.3b into the NAc (in mm, relative to bregma: A/P: +1.42; M/L $\pm$ 0.9; D/V: -4.5$\rightarrow$-4.1)

followed by bilateral VTA injections of CRISPR and Chrimson (0.5 µL each side; control mice

received Chrimson viruses diluted to the same ratio). For *in vivo* fiber photometry of dLight,

mice received control or CRISPR viruses bilaterally into the VTA and then were injected

unilaterally with dLight1.3b into the NAc (in mm, relative to bregma: A/P: -+1.43; M/L - 0.9;

D/V: -4.5→-4.1), total volume 0.5 µL; a photometry fiber purchased from Doric (400 µm optical

fiber, 1.25 mm metal ferrule) was lowered to D/V -4.1 mm. Fibers were secured to the skull with

Metabond (Parkell) and dental cement (Lang Dental). For *in vivo* electrophysiology, microdrives

were constructed in-house with a Neuralynx EIB (EIB-16) mounted on a 3D printed L-frame

(Applied Rapid Technologies). Each microdrive contained in house constructed optetrodes of an

in house constructed optic fiber (200 µm fiber bore and 1.25 mm ceramic ferrule, Precision Fiber

Products) and an in-house constructed tetrode bundle of tungsten wire (each tetrode was 4 wires

25 mm diameter spun together and heat-sealed, California Fine Wire). After virus injection for

CRISPR/ChR2 mixture, a window was created in the skull and the microdrive was lowered to

above the VTA (-3.8 mm). Optetrode microdrives were secured to the skull with Metabond and

dental cement.

***In situ* hybridization using RNAscope:** Wildtype C57BL/6 male and female mice (8-9 weeks

old) were used to map ion channel expression in the VTA using RNAscope (ACDBio) (*27*).

Brains were flash frozen in 2-methylbutane, and stored at -80°C. VTA sections were collected

and processed for RNAscope (20 um, coronal). Representative sections that spanned the VTA

were selected for hybridization (A/P in mm: rostral: -3.1; intermediate: -3.5; caudal: -3.8;

according to the Paxinos atlas). Sections were prepared for hybridization per manufacturer's

(Advanced Cell Diagnostics, Inc) instructions using probes for *Th* (Mm-*Th*), *Kcnd3* (Mm-*Kcnd3*-

C3), and *Kcnma1* (Mm-*Kcnma1*-C2). Slides were coverslipped with Fluoromount with DAPI

(Southern Biotech) and imaged using a confocal fluorescent microscope (University of

Washington Keck Center (NIH grant S10 OD016240), Leica SP8X confocal). Images for each

RNAscope experiment were collected using the same acquisition settings for each slide. Analysis

was performed on Cell Profiler 4.1.3 (https://github.com/CellProfiler/CellProfiler) using a

modified image analysis pipeline for RNAscope (*28*).

Immunohistochemistry for target validation: Following completion of experiments, all mice

were checked for viral targeting or implant placement. Mice were deeply anesthetized with

Buthenesia and transcardially perfused with cold 1X PBS followed by cold 4%

paraformaldehyde (PFA, Sigma-Aldrich). Implants were removed from mice used in *in vivo*

electrophysiology or fiber photometry experiments. Brains were post fixed for 24 hours in PFA

at 4°C, then moved to 30% sucrose in PBS until cryosectioning occurred. All cryosections were

stored in 1X PBS with 0.3% Sodium Azide (Sigma-Aldrich) until immunohistochemistry. For

colocalization validation and behavioral experiments, the VTA was coronally sectioned at 30

μm. For *in vivo* electrophysiology experiments, the VTA was coronally sectioned at 60 μm. For

in vivo dLight fiber photometry experiments, coronal sections of the NAc (60 μm) and the VTA

(30 μm) were obtained. For *ex vivo* dLight slice experiments, the part of the brain containing the

VTA was blocked at the same time as obtaining NAc slices for slice imaging. The block was

post-fixed overnight in 4% PFA at 4°C. They were then transferred to 30% sucrose in PBS for at

least 24 hours and then VTA slices were cryosectioned at 30 μm. NAc slices (250 μm) used in

dLight slice two-photon imaging were immediately transferred to 4% PFA after recording and

were fixed overnight at 4°C. Slices were then transferred to 1X PBS until the day of

immunohistochemistry. Free floating immunohistochemistry experiments were performed using

the following antibodies as needed: rabbit anti-HA (Sigma, cat#: H6908), mouse anti-TH

(Millipore, cat#: MAB318), mouse anti-dsred (Takara, cat#: 632496), rabbit anti-GFP

(Proteintech, cat#: 50430-2-AP), and chicken anti-GFP (Aves, cat#: GFP-1020) to target

CRISPR, TH, Chrimson-TdTomato, Chr2-eYFP, or dLight1.3-GFP viruses followed by

secondary antibody application (JacksonImmuno, cat#s: 703-545-155, 711-165-152, 711-165-

152, 711-165-152). Sections were placed on slides, coverslipped with Fluoromount with DAPI,

and kept in the dark at 4°C until the day of imaging. Images were collected at the UW Keck

Center or on a Keyence Fluorescence Microscope (Keyence). Quantification of co-labeled cells

for immunohistochemistry was performed using ImageJ 1.53 Cell Counter/Multi-point tool.

Fluorescence-activated cell sorting and targeted deep sequencing: Procedures were

performed as previously described (*13*, *26*). AAV1-hSyn1-FLEX-EGFP-KASH and CRISPR

viruses (AAV1-CMV-FLEX-SaCas9-U6-sgRNA) targeting either *Kcnd3* or *Kcnma1* were

stereotaxically co-injected into the VTA of DAT-IRES-Cre mice. Four weeks after surgery, the

VTA of each mouse was extracted, flash frozen in liquid nitrogen, and stored at -80°C. On day of

FACS, VTA tissue from 5 mice per experimental group were combined into a single sample and

homogenized in buffer containing (in mM): 320 Sucrose (sterile filtered), 5 CaCl (sterile

filtered), 3 Mg(Ac)2 (sterile filtered), 10 Tris pH 7.8 (sterile filtered), 0.1 EDTA pH 8 (sterile

filtered), 0.1% NP40, 0.1 Protease Inhibitor Cocktail (PIC, Sigma), 1 β-mercaptoethanol. Nuclei

were isolated by centrifugation in 29% iso-osmolar Optiprep solution. GFP+ and GFP- nuclei

were sorted using a BD AriaFACS III (University of Washington Pathology Flow Cytometry

Core) into a PCR tube strip containing 3 µL of REPLI-g Advanced Single Cell Storage buffer

(Qiagen). Whole genome amplification (WGA) was performed directly following FACS using

the REPLI-g Advanced DNA Single Cell Kit (Qiagen) according to manufacturer's instructions.

For generation of amplicons, samples were amplified (PCR 1) with Phusion High Fidelity

Polymerase (Thermo Fisher) followed by reamplification (PCR 2) with a second set of primers.

The amplicons were gel extracted using the MinElute gel extraction kit (QIAGEN) and sent to

Genewiz for Sanger and amplicon sequencing (Amplicon-EZ). The following primer sets were

used to generate <300bp amplicons: PCR 1: *Kcnd3* forward: CATTGGATGGATGCCAGT;

Kcnd3 reverse: GAGGATGCCATAGAAGGC; *Kcnma1* forward:

GCAACATGGCTGTTGATG; *Kcnma1* reverse: ACCTCCATGGTCACCGGT. PCR 2: *Kcnd3*

forward: TAGCTCCAGCGGACAAGA; *Kcnd3* reverse: CAGAGATGCATTCATAGC;

Kcnma1 forward: CCATTGCTAGCTATGGCA; *Kcnma1* reverse:

ATGAGCGCATCCATCTTG.

Quantitative PCR (qPCR): qPCR was utilized to determine mRNA expression of relevant

potassium channel subunits in the VTA of control, Kv4.3 LOF and BKCa1.1 LOF mice. Four

weeks after CRISPR injection, tissue punches containing the VTA (bilateral) were collected and

immediately frozen on dry ice, prior to storage at -80°C. RNA isolations were conducted from

tissue samples using Qiagen RNeasy Micro Kit (Qiagen, Catalog no. 74004) according to the

manufacturer's instructions. The RNA quality and concentration was assessed for each sample

spectrophotometrically (A260/A280 & gt;1.9) by a NanoDrop instrument. Then 100 ng of RNA

was reverse transcribed by PCR using SuperScript IV VILO (Invitrogen, Catalog no. 11755050).

qPCR was conducted on the cDNA using a Taqman Gene Expression assay consisting of the

Taqman Gene Expression Master Mix, and the appropriate Taqman probes in FAM channel

(Master Max: Thermo Fisher Scientific, Applied Biosystems, catalog no. 4369016, Taqman

probes: *GAPDH* (Assay ID: Mm99999915_g1; Cat. number: 4448489), *En1* (Assay ID:

Mm00438709_m1_g1; Cat. number: 4448892), *Th* (Assay ID: Mm00447557_m1_g1; Cat.

number: 4331182), *Kcnn3* (Assay ID: Mm00446516_m1; Cat. number: 4448892) and *Kcnd2*

(Assay ID: Mm01161732_m1; Cat. number: 4448892). All qPCRs were performed in triplicates

and run on an Agilent Technologies MX3005P qPCR machine at 50°C for 2 mins, 95°C for 20 s,

then 95°C for 3 s and 60°C for 30 s for 40 cycles. The threshold cycle (Ct value) of each gene

was then normalized to GAPDH. ΔCT values were calculated by subtracting the geometric mean

of CT values for *Gapdh*, *En1*, and *Th* from the CT values of the gene of interest (*Kcnd2* or

Kcnn3). Control ΔCT values for the gene of interest were subtracted from the ΔCT values of the

experimental group to generate ΔΔCTs. Fold change was calculated as $2^{-(\Delta\Delta CT)}$.

Slice preparation for *ex vivo* physiology: Mice injected with CRISPR/eYFP or

CRISPR/Chrimson/dLight1.3b were allowed 4-5 weeks recovery after surgery to allow for viral

expression, mutagenesis, and residual protein turnover. All solutions were continuously bubbled

with O_2/CO_2. Coronal brain slices containing the VTA (200 μm for VTA physiology) or NAc

(250 μm for dLight slice two-photon imaging) were prepared in a slush NMDG cutting solution

(in mM: 92 NMDG, 2.5 KCl, 1.25 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5

Na-ascorbate, 3 Na-pyruvate, 0.5 $CaCl_2$, 10 $MgSO_4$, pH 7.3–7.4 (*29*). Slices recovered for ~12

min in the same solution in a 32°C water bath. Slices were then transferred to a room

temperature HEPES-aCSF solution (in mM: 92 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 30 $NaHCO_3$, 20

HEPES, 25 glucose, 2 thiourea, 5 Na-ascorbate, 3 Na-pyruvate, 2 $CaCl_2$, 2 $MgSO_4$). Slices

recovered for an additional 60 min.

Slice electrophysiology: Whole-cell patch-clamp recordings were made using an Axopatch

700B amplifier (Molecular Devices) with sampling at 10 kHz and filtering at 1 kHz. eYFP+ cells

were visually identified via fluorescence for patching with electrodes at 3–5 MΩ. Series

resistance was monitored during all recordings with changes in resistance $\pm$ 20% qualifying the

cell for exclusion. Recordings were made in aCSF (in mM: 126 NaCl, 2.5 KCl, 1.2 NaH_2PO_4,

1.2 $MgCl_2$ 11 D-glucose, 18 $NaHCO_3$, 2.4 $CaCl_2$) at 32°C continually perfused over slices at a

rate of ~2 ml/min. VTA dopamine neurons were identified by fluorescence. Electrodes were

filled with an internal solution containing (in mM): 130 potassium gluconate, 10 HEPES, 5

NaCl, 1 EGTA), 5 Mg2+/ATP, 0.5 Na+-GTP, pH 7.3, 280 mOsmol.

Spontaneous activity, excitability, and action potential shape measurements: Spontaneous

activity was recording in I=0 mode for 1 minute and I/V curves (0-80 pA, 10 pA steps, 1 s) for

excitability were immediately measured in I-Clamp mode (done both at cell's resting membrane

potential in fig. S4, O and when holding the cell at -60 mV in Fig 2, G-H). All data were

analyzed offline using Clampfit (Molecular Devices). Frequency of spontaneous activity was

calculated as number of events/s and CV-ISI was measured using the standard deviation of the

interspike interval (ISI) from spontaneous activity. Excitability was measured as the total number

of events during each current step. Action potentials of spontaneous activity were detected using

the Event Detection function in Clampfit (restricted window parameters: between 50 ms before

action potential peak and 150 ms after action potential peak) and used to determine firing

frequency and CV-ISI, and action potential shape analysis from baseline resting membrane

potential. For action potential shape analysis in a clamped cell, the first action potential of

activity elicited from a 20 pA current injection while the cell was being held at -60 mV was

detected using the Event Detection function in Clampfit with identical restricted window

parameters. Event statistics from the Event Detection function were extracted, including action

potential peak, half-width, decay tau, and afterhyperpolarization (the anti-peak Event Statistics

characteristic corrected for resting membrane potential measure 50 ms before spike peak). For

each cell, detected action potentials were averaged to visualize action potential waveforms and

phase plane plots. Phase plane plots were calculated as the dV/dt and plotted as the dV/dt over

voltage. Action potential threshold was defined as the point at which dV/dt of the AP waveform

reaches 4% of its maximum (*23*). Resting membrane potential was recorded as the membrane

potential at the beginning of the action potential waveform from the Event Detection window (50

ms before peak). For action potential shape analysis of control VTA dopamine neurons following

pharmacological inhibition of A-Type or BK potassium currents, AmmTx3 (10 nM, Tocris) and

Iberiotoxin, (150 nM, Cayman Chemical) were perfused onto the slice in oxygenated aCSF and

spontaneous activity was recorded. Shape analyses were performed as described above.

Potassium current measurements: For A-Type and BK potassium channel current, I_A and I_{BK}

respectively, recordings in V-Clamp mode, aCSF was supplemented with 500 nM TTX to

eliminate action potential activity and cells were held at -60 mV. I_A was activated using voltage

steps from -108mV to -28 mV to -48mV then back to holding potential every 5 s, 5 times.

AmmTx3 (10 nM, Tocris) was used to pharmacologically block Kv4 channels. I_{BK} was measured

using subtraction before and after pharmacological inhibition of BK channels (iberiotoxin, 150

nM (Cayman Chemical)) during depolarizing voltage steps that activated I_K from -63mV to

+2mV to -48 mV to -63 mV every 5 s, 5 times. Data were analyzed offline using Clampfit. For

A-type potassium currents, differences between the peaks at the -28mV and the -48 mV step

were determined and calculated. For quantifying BK currents, we first present in Fig. 1, J-K

maximal peaks of raw I_K in Fig. 1, J-K before (-, I_K^{Pre}) and after (+, I_K^{Post}) application of the BK

channel specific inhibitor iberiotoxin from mice that received control virus or sgKcnma1 virus to

knockdown *Kcnma1* (BKCa1.1). A significant difference from I_K^{Pre} either indicates an

iberiotoxin specific effect indicative of BK channel blockade or a CRISPR/cas9 LOF effect

indicative of BKCa1.1 knockdown. Notably, no difference between sgKcnma1 I_K^{Pre} and I_K^{Post}

indicates that there is no iberiotoxin-sensitive current (I_{BK}). We also present the subtracted I_{BK} in

fig. S4, A-C. which was calculated by subtracting I_K^{Post} from I_K^{Pre}. We then measured the

maximal peak generated from the depolarization and the total area under the curve to quantify

I_{BK} .

***In vivo* single-unit electrophysiology:** Recordings began at least 4 weeks after surgery in a

recording cage with corncob bedding or during behavior using Neuralynx acquisition hardware

and software (Digital Lynx 4SX, Cheetah Data Acquisition). For *in vivo* electrophysiology

recordings during behavior (behaviors described below), mice were recorded in operant

chambers (MedAssociates) that were interfaced with the Data Acquisition software. For all

experiments, the ceramic ferrule of the optic fiber in the microdrive was attached to a patch cord

that was connected to a 473 nm blue laser (Laserglow technologies). A preamplifier was

connected to the microdrive's EIB. Recordings were acquired either in the home cage or during

behavior through Cheetah Data Acquisition. Optical stimulations were synced through Cheetah

with a function generator (Agilent). Data were continuously sampled, amplified, and filtered

between 100 and 9000 Hz, and digitized at 32 kHz. If the spike exceeded a predetermined

threshold, discretely sampled data were recorded and filtered between 100 and 6000 Hz. At the

end of the recording, 10 blue light pulses were delivered 20 times through the optic fiber of the

microdrive to activate neurons that expressed ChR2 (5 ms long at 20 Hz, putative dopamine

neurons). The light intensity (5-15 mW/mm2) was adjusted so that light-evoked spike

waveforms were similar to spontaneous ones. This was followed by quinpirole injections

(described below). All optetrodes were lowered in 80 μm increments at the end of the day to

identify a new cell field the next day. Data were analyzed offline using Plexon Offline Sorter,

SpikeSort3D, Neuroexplorer, and Matlab, as previously described (*30*). Briefly, units were

identified using cluster analysis in the Offline Sorter or SpikeSort3D (KlustaKwik) and saved.

Frequency, burst analysis, and blue-light sensitive (ChR2 expressing putative dopamine neurons)

identification were performed in Neuroexplorer and Matlab. Putative dopamine neurons were

identified as either sensitive to ChR2 stimulation or sensitive to pharmacological identification:

light/optically sensitive putative dopamine neurons were identified as those that had a spike

latency of ≤ 5 s and 0.6 probability during blue light activation of ChR2; pharmacologically-

sensitive putative dopamine neurons were identified as those that had a reduction in firing rate to

30% or less of baseline after an i.p. injection of 0.2 mg/kg of quinpirole chloride (example

histogram in fig. S5, C), a dopamine receptor D2 agonist expressed on a significant

subpopulation of dopamine neurons *(11)*. The onset of a burst was identified as two consecutive

spikes with an inter-spike interval of <80 ms and its termination was defined as an inter-spike

interval of >160 ms *(30)*.

The latency to spike and probability of spiking after ChR2 stimulation under baseline and

behavioral conditions are reported in fig. S5, C. Optically-sensitive putative dopamine units from

mice that underwent recordings only in the home cage and optically-sensitive putative dopamine

units that were recorded during operant conditioning that were not responsive to a lever press

(LP) during behavior were included in the analysis of baseline firing properties reported in Fig.

2, L-M and fig. S5, D-F. Cells that were identified to be responsive to LP events were not

included in analysis of baseline properties due to potential confounds of peri-event activity but

their mean firing rates are reported in fig. S7 and S10. For analysis of *in vivo* firing of putative

dopamine neurons during a LP, 20 s window ($\pm$ 10 s) peri-event windows surrounding the LP

during FR1 reinforced day 1 and day 5, and extinction day 1 were extracted and z-scores

calculated using the first 3 s (-10 to -7 s) of the peri-event window as the peri-event baseline. To

determine which units were behaviorally responsive to the LP, optically- and pharmacologically-

sensitive units that had a $\pm$ 3 STDs shift from the median baseline (-10 to -7 s prior to LP) within

$\pm$ 1 s of the LP event were extracted. AUCs were then determined for these events: LP (0 to 0.7 s

post LP) and Rew (3 s post LP to 6 s post LP).

Overnight locomotor activity: 4 weeks after surgery, baseline locomotion was measured using

locomotion chambers (Columbus instruments) that use infrared beam breaks to calculate

ambulatory activity. Mice were singly housed and provided with *ad libitum* access to food and

water. Locomotion was monitored for 60 consecutive hours (Friday-Monday: 3 nights, 2 days).

Behavioral tracking failed for one BKCa1.1LOF mouse, however this mouse was still included

in operant studies.

Open field test: At least 4 weeks after surgery, mice were tested on an open field test to

determine alterations in exploration and anxiety. Here, mice we placed into a large circular arena

(120 cm diameter) and activity was recorded for a period of 10 min. behavior was analyzed

offline using Ethovision software (Noldus). Here, zones arena edge and arena center zones were

generated. Time in center (a marker of exploratory/anxiolytic behavior) and total distance were

calculated.

Elevated plus maze test: At least 4 weeks after surgery, mice were tested on an elevated plus

maze (Med Associates) to determine alterations in anxiety. The plus-shaped maze has two

opposing closed-walled arms and two opposing open-ledge (1 cm tall) arms with a center zone.

The floors of the maze are white. A mouse was placed into the center zone and allowed to freely

explore the maze for 10 min. Video was recorded and analyzed offline using Ethovision. Total

distance traveled and time spent in the center, closed, and open arms were determined. Percent

time spent in arms was calculated (a marker of anxiolytic behavior).

Pavlovian conditioning: At least 4 weeks after surgery, mice were food restricted to 85% of

their body weight and were maintained at this bodyweight for all conditioning days.

Conditioning sessions took place in mouse conditioning boxes outfitted with a house light,

ultrasensitive levers, a food hopper with a head entry detector, nose poke holes, and barred floors

(MedAssociates). Pavlovian training sessions consisted of 25 trials in which 10 s lever extension

(an audiovisual CS). After 10 s lever presentation, levers were retracted and a sucrose pellet

reward (Rew, 20 mg sucrose pellet, Bioserv) was delivered in the food hopper. Each trial was

separated by a random inter-trial interval averaging 70 s and animals underwent 7 days of

training. The house light remained on during the entire session. Conditioned approach scores

were calculated as the rate of head entries during the CS - rate of head entries during the ITI.

Mice used for *in vivo* electrophysiology or dLight experiments first underwent 3 days of 20 min

acclimation sessions to patch cord tethering in the conditioning boxes followed by 10 days of

Pavlovian conditioning while tethered.

Operant conditioning: After Pavlovian conditioning, mice began an FR1 schedule of reinforced

operant conditioning. Each session lasted for 60 min. Here, levers were extended and remained

extended until a LP. Upon a lever press, levers were retracted, and a sucrose pellet was

immediately delivered into the food hopper (Rew). For *in vivo* electrophysiology or dLight

experiments during FR1 reinforced training, a temporal delay of 3 s between LP and Rew was

included to dissociate dynamics of LP and Rew. The levers did not extend again until the mouse

entered the food hopper to retrieve the sucrose pellet. The house light remained on during the

entire session. Reinforced FR1 sessions lasted for 3 days (5 days for tethered mice). A subset of

mice then underwent a progressive ratio test where the number of lever presses necessary for

sucrose pellet delivery increases non-arithmetically (i.e., 1, 2, 4, 6, 9, 13…) over the course of

the session. The progressive ratio session automatically ended after 3 consecutive min of no

lever presses or after 3 hours. Another subset of mice that had undergone FR1 reinforced training

underwent FR1 extinction for 60 min each session for five days. Here, levers extend and retract

similarly to the FR1 reinforced paradigm, yet a sucrose pellet reward is omitted after a LP. Mice

underwent extinction for 5 days. *In vivo* electrophysiology and dLight mice also underwent 5

days of extinction training following completion of FR1 reinforced training.

dLight recordings using fiber photometry during operant conditioning: Fiber photometry

recordings lasted the entirety of the behavioral conditioning sessions (first and last days of

reinforced FR1 and extinction FR1). dLight1.3b was used to measure dopamine release in the

NAc during these behaviors of control or CRISPR mice. Prior to every session, the implanted

optic fiber was attached to a previously photobleached patchcord (Doric Lenses) using a ceramic

sleeve (Doric Lenses). Recordings were acquired using an RZ5 BioAmp Processor from Tucker

Davis Technologies (TDT). A 465 nm LED (531-Hz, sinusoidal, Doric Lenses) was used to

excite dLight1.3b and a 405 nm LED was used as an isosbestic control (211-Hz, sinusoidal,

Doric Lenses) using an LED driver (Doric Lenses) controlled via the RZ5. Prior to each session,

patch cords had been photobleached for at least one hour. Laser intensities were measured at the

tip of the optic fiber to reach levels of 30-45 uW before being bandpass filtered (525 ± 25 nm,

Doric, FMC4), transduced by a femtowatt silicon photoreceiver (Doric Lenses), and recorded by

a real-time processor (TDT, RZ5). The 531-Hz and 211-Hz signals were extracted in real-time

by the TDT program Synapse at a sampling rate of 1017.25 Hz. MedAssociate signals of

behavioral events such as LP and Rew were synced to the recordings for further analysis offline.

Photometry analysis of perievents: A custom software GUI was generated to extract,

downsample and z-score the photometry signals surrounding behavioral events for each mouse

(20 s window: $\pm$ 10 s behavioral event). Data were first downsampled to 100 Hz and DF/F was

calculated using the custom software GUI. Next, Z-score was calculated relative to the mean and

standard deviation of the first 4 s of the 20 s time window (baseline perievent time). Mouse data

were further processed in custom MatLab script. Z-scores for each mouse were then

downsampled to 50 Hz and then averaged. Data were exported to GraphPad Prism for statistical

analysis, area under the curve (AUC) analysis, peak analysis and trace export. For reinforced

FR1 training and omission extinction training, AUC time bins were calculated for a set time

following the event: LP (3 s post), Rew (5 s post).

Two(2)-photon slice imaging of dLight: NAc slices were transferred to a heated recording in

aCSF (in mM: 126 NaCl, 2.5 KCl, 1.2 NaH_2PO_4, 1.2 $MgCl_2$ 11 D-glucose, 18 $NaHCO_3$, 2.4

$CaCl_2$) at 32°C continually perfused over slices at a rate of ~2 ml/min on an Olympus Fluoview

FVMPE-RS 2-photon microscope. The NAc was identified using anatomical markers and

epifluorescence and expression of dLight GFP and VTA dopamine terminals expressing

Chrimson-TdTomato was quickly confirmed before recording. Previous reports have found that

this preparation does not induce stimulation of Chrimson (*36*). We used a resonant scanner (30

Hz frame rate acquisition) and performed an online averaging of 6 times to get an effective frame

rate of 5 Hz. A GaAsP-PMT with adjustable voltage, gain and offset was used, along with a

green filter cube. We used a 25x immersion objective (Olympus, XLPLN25XWMP, 1.05 NA, 2

mm WD) and imaged with a 920 nm laser (SpectraPhysics, ~100 fs pulse width) with automated

wavelength-specific alignment. Chrimson fibers in the NAc were stimulated at increasing

frequencies for 1 s following a 10 s baseline using full field 615 nm LED stimulation through the

objective (CoolLED, PE-100). Data were analyzed in FiJi ImageJ using the Time Series

Analyzer plug in. Z-score of fluorescence was calculated from a 10 s baseline preceding the

Chrimson stimulation. AUC calculations were performed in GraphPad Prism.

Statistics: Data were analyzed for statistical significance using GraphPad Prism. Data

distribution was assumed to be normal but this was not formally tested. No statistical methods

were used to predetermine sample sizes, but our sample sizes are similar to those reported in

previous publications (*11-13, 30*). Littermates were randomly assigned to experimental groups and

animals were tested in a random order across all biological replicates. All behavioral assays were

repeated in a minimum of three cohorts with similar replication of results. Data for *in situ*

hybridization, genetic validation of mutagenesis, 2-photon imaging, and fiber photometry were

pooled from at least three individual animals, collected from at least two rounds of experiments

with similar replication of results. VTAs for qPCR were pooled from at least two mice per group.

Electrophysiology data are from individual cells collected from at least two individual animals

and collected from at least two rounds of experiments with similar replication of results. Data

were analyzed by an investigator blinded to experimental condition; the experimenter was not

blinded during data collection. Animals with missed viral injections or significant viral spread

outside the targeted region were excluded from analyses. These criteria and other

inclusion/exclusion criteria described above were preestablished. All statistical tests were two-

sided, except for ratio data in fig. S9, G and correlational data in fig. S9, H, and when

appropriate corrected for multiple comparisons to controls (Fisher's or Dunnet's), except for

current data in Fig. 1 G and K where a Sidak's test was used for pharmacological blockade

analysis. Detailed statistical analyses can be found in the figure legends.

773
774

Acknowledgments

We would like to thank members of the Zweifel, Palmiter, and Chavkin laboratories at

the University of Washington for their helpful discussions. We would also like to thank the staff

of the University of Washington's Molecular Genetics Research Core, Imaging and Neural

Circuits Core, Comparative Medicine Animal Facilities, the University of Washington's Keck

Imaging Center, and the University of Washington's Pathology Flow Cytometry Core.

Funding:

This study was supported by:

K99-DA054265 (B.J.)

R00-DA054265 (B.J.)

T32-DA727825 (B.J.)

F32 MH127801 (M-S.K)

F31 DA053724 (J.E.E)

F31-MH116549 (A.C.H.)

T32-GM007270 (A.C.H.)

F31 MH126489 (M.C.)

T32-GM007270 (M.C.)

R03TR003307 (M.E.S.)

R01-MH104450 (L.S.Z.)

R01-DA044315 (L.S.Z.)

University of Washington's Addictions, Drugs and Alcohol Institute:

ADAI-1019-17 (B.J.)

National Research Foundation of Korea:

MEST 2020R1A2C2102134 (Y.S.J)

Core grants:

S10OD016240 (University of Washington Keck Center)

P30-DA048736 (University of Washington Center of Excellence in Opioid Addiction

Research)

B.J. was funded by the postdoctoral Enrichment Program Award from the Burroughs Wellcome

Fund.

Author contributions:
Conceptualization: BJ, LSZ
Methodology: BJ, MES, SN-E, LSZ
Investigation and analysis: BJ, M-SK, YSJ, JEE, JXY, MC, ACH, MAQ, MAB, MJ, DB
AM, NLG, LSZ
Visualization: BJ, ACH, LSZ
Supervision: BJ, LSZ
Writing—original draft: BJ, LSZ
Writing—review & editing: all authors reviewed and approved this manuscript. BJ and
LSZ edited the manuscript following co-author review.

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: Data, materials, and code can be made available upon request.

Figure legends:

Fig. 1 Targeted mutagenesis of *Kcnd3* and *Kcnma1* in dopamine neurons.

(**A-B**) RNAscope *in situ* hybridization for *Kcnd3*, *Kcnma1*, and *Th*. White arrows: representative

Th/Kcnd3/Kcnma1 overlap. Scale bar, 100 μm (**A**) and 50 μm (**B**). (**C**) Quantification of overlap

for *Kcnd3, Kcnma1,* and *Th* in the VTA (N=3 mice). (**D**) Schematic of AAV1-FLEX-SaCas9-

sgKcnd3 targeting virus vector. (**E**) Proportion of targeted deep sequencing reads with indel

mutations following sgKcnd3 targeting (N=5 mice). (**F**) Average pre- and post-AmmTx A-Type

currents (I_A) from control (left) and sgKcnd3 (right) -targeted mice (N=4 cells/group; 3

mice/group) and corresponding depolarization step to elicit I_A. (**G**) Average peak I_A from (**F**) in

control and sgKcnd3-targeted mice (Two-way RM ANOVA, Interaction effect $F_{(1,6)}$=18.96,

P<0.01; Group effect $F_{(1,6)}$=22.6, P<0.01; Drug effect $F_{(1,6)}$=16.1, P<0.01; Sidak's Multiple

Comparison test, ***p<0.001; $N_{control}$=4 cells, 3 mice, $N_{sgKcnd3}$=4 cells, 3 mice). (**H**) Schematic of

AAV1-FLEX-SaCas9-sgKcnma1 targeting virus vector. (**I**) Proportion of targeted deep

sequencing reads with indel mutations following sgKcnma1 targeting (N=5 mice). (**J**) Average

pre- and post-iberiotoxin potassium currents (I_K) from control (left) and sgKcnma1 (right) -

targeted mice ($N_{control}$=11 cells, 4 mice; $N_{sgKcnma1}$=11 cells, 3 mice) and corresponding

depolarization step to elicit I_K. (**K**) Peak I_K from (**J**) in control and sgKcnma1-targeted mice before

(-) and after (+) iberiotoxin (Two-way RM ANOVA, Interaction effect $F_{(1,19)}$=18.45, P<0.001;

Group effect $F_{(1,19)}$=6.553, P<0.05; Drug effect $F_{(1,19)}$=69.84, P<0.0001; Sidak's Multiple

Comparison test, *p<0.05, ***p<0.001; $N_{control}$=11 cells, 4 mice; $N_{sgKcnma1}$=11 cells, 3 mice). Box

and whisker graphs are presented as min-to-max.

Fig. 2 Kv4.3 and BKCa1.1 LOF alter the dopamine action potential waveform and activity.

(**A**) Average action potential waveform from spontaneously active VTA dopamine neurons ($N_{control}$=21 cells, 7 mice, $N_{Kv4.3LOF}$=16 cells, 6 mice, $N_{BKCa1.1}$=19 cells, 7 mice). (**B**) Representative phase plane plot of action potential dynamics. (**C**) Action potential half width in control, Kv4.3 LOF, and BKCa1.1 LOF cells (One-Way ANOVA: $F_{(2,53)}$=9.109, P<0.001, Dunnet's Multiple Comparison's test to controls,t ***p<0.001). (**D**) Decay kinetics of the action potential (One-Way ANOVA: $F_{(2,53)}$=10.67, P<0.0001, Dunnet's Multiple Comparison's test to controls, ***p<0.001). (**E**) Afterhyperpolarization of the action potential (One-Way ANOVA: $F_{(2,53)}$=5.314, P<0.01, Dunnet's Multiple Comparison's test to controls, *p<0.05). (**F**) Resting membrane potential (One-Way ANOVA: $F_{(2,53)}$=6.026, P<0.01, Dunnet's Multiple Comparison's test to controls, **p<0.01). (**G**) Representative evoked excitability traces. (**H**) Current-voltage plot (Two-Way ANOVA, Interaction effect: $F_{(16, 408)}$ = 1.143, P=0.3126; Group effect: $F_{(2, 51)}$ =4.495 P<0.05; Current effect: $F_{(8, 408)}$ =69.36 P<0.0001; Dunnet's Multiple Comparison's test to controls: *p<0.05, **p<0.01; $N_{control}$=19 cells, $N_{Kv4.3 LOF}$=15 cells, $N_{BKCa1.1LOF}$=20 cells). (**I**) Representative spontaneous activity traces. (**J**) Firing rate of spontaneous activity (One-Way ANOVA: $F_{(2,53)}$=3.379, P<0.05, Dunnet's Multiple Comparison's test to controls, *p<0.05). (**K**) CV-ISI of spontaneous activity (One-Way ANOVA: $F_{(2,53)}$=3.606, P<0.05, Dunnet's Multiple Comparison's test to controls, *p<0.05). (**C-F** and **J-K**; $N_{control}$=21 cells, 7 mice, $N_{Kv4.3LOF}$=16 cells, 6 mice, $N_{BKCa1.1}$=19 cells, 7 mice). (**L**) Burst rate in VTA dopamine neurons *in vivo* (One-Way ANOVA: $F_{(2,117)}$=5.673, P<0.01, Dunnet's Multiple Comparison's test to controls **p<0.001). (**M**) Burst duration in VTA dopamine neurons *in vivo* (One-Way ANOVA: $F_{(2,117)}$=5.366, P<0.01, Dunnet's Multiple Comparison's test to controls, *p<0.05). (**L-M** $N_{control}$=28 cells, 7 mice, $N_{Kv4.3 LOF}$=49 cells, 10 mice, and $N_{BKCa1.1}$=41 cells, 9 mice). (**N**) Schematic of 2-photon slice imaging of dLight1.3b with optically evoked

dopamine release of VTA terminals expressing Chrimson + sgRNA in the NAc. (**O**) Average

dLight1.3b signal following 1 s stimulation at different frequencies in control, Kv4.3 LOF and

BKCa1.1 LOF mice. (**P**) Average AUC for responses for data presented in panel **O** (Mixed effects

analysis, Interaction effect $F_{(6, 77)}$=0.6984, P=0.6516, Group effect $F_{(2, 26)}$=0.05105, P=0.9503,

Frequency stimulation effect $F_{(1.725, 44.28)}$=25.99, P<0.0001). (**P-O**, $N_{control}$=6 slices, 5 mice;

$N_{Kv4.3LOF}$=8 slices, 4 mice; $N_{BKCa1.1LOF}$=15 slices, 3 mice; except for 5 Hz where $N_{Kv4.3LOF}$=7 slices

because of brief interruption in data capture that was not detected until post-hoc analysis,). Trace

averages are presented as mean + SEM. Box and whisker graphs are presented as min-to-max.

Fig. 3 Pavlovian and instrumental behavior in control, Kv4.3, and BKCa1.1 LOF mice.

(**A**) Schematic of Pavlovian conditioning paradigm. (**B**) Conditioned approach score (head entries during CS+ - head entries during ITI) across days is not different between groups, (Two-Way RM ANOVA, Interaction effect: $F_{(12, 378)}=1.557$, P=0.1019; Time effect: $F_{(5, 378)}=31.97$, P<0.0001; Group effect: $F_{(2, 63)}=1.093$, P=0.3416) (**C-D**) Head entry rate during CS (C) and ITI period (D) across days used to calculate conditioned approach is not different between groups, (Two-Way RM ANOVA <u>Control ITI vs Cue comparison:</u> Interaction effect $F_{(6, 276)}=3.86$, P<0.01; Time effect $F_{(3.403, 156.5)}=249$, P<0.0001; ITI vs Cue effect $F_{(1, 46)}=6.433$, P<0.05; <u>Kv4.3 LOF ITI vs Cue comparison:</u> Interaction effect $F_{(6, 288)}=2.555$, P<0.05; Time effect $F_{(3.403, 156.5)}=249$, P<0.0001; ITI vs Cue effect $F_{(1, 46)}=6.433$, P<0.05; <u>BKCa1.1 LOF ITI vs Cue comparison:</u> Interaction effect $F_{(6, 204)}=4.286$, P<0.0001; Time effect $F_{(4.053, 137.8)}=36.36$, P<0.0001; ITI vs Cue effect $F_{(1, 34)}=6.989$, P<0.05; Sidak's Multiple Comparison's test, Cue vs ITI, *p<0.05). (**B-D** $N_{control}=23$; $N_{Kv4.3LOF}=25$; $N_{BKCa1.1LOF}=18$). (**E**) Schematic of instrumental FR1 conditioning paradigm. (**F**) Lever presses during days 1-3 of FR1 conditioning (Two-Way RM ANOVA Interaction effect $F_{(4, 106)}=2.781$, P<0.05; Time effect $F_{(1.631, 86.42)}=18.26$, P<0.0001; Group effect $F_{(2, 53)}=0.9557$, P=0.3911; Subject effect $F_{(53, 106)}=2.018$, P<0.01; Dunnet's Multiple Comparison's test to controls, *p<0.05). (**G**) Cumulative lever presses during day 1 (left) and day 3 (right) of FR1 conditioning (Two-Way RM ANOVA, <u>Day 1:</u> Interaction effect $F_{(118, 3127)}=3.769$, ****P<0.0001; Time effect $F_{(1.807, 95.78)}=143.0$, P<0.0001; Group effect $F_{(2, 53)}=4.832$, P<0.05; Subject effect $F_{(53, 3127)}=167.2$, P<0.0001; Dunnet's Multiple Comparison's test to controls, Control vs BKCa1.1LOF: p<0.05 for mins 35-42; p<0.01 for mins 43-60; <u>Day 3:</u> Interaction effect $F_{(118, 3127)}=1.455$, **P<0.01; Time effect $F_{(1.288, 68.26)}=228.8$, P<0.0001; Group effect $F_{(2, 53)}=4.299$, P<0.05; Subject effect $F_{(53, 3127)}=152.3$, P<0.0001; Dunnet's Multiple Comparison's test to controls, Control vs Kv4.3 LOF:

p<0.05 for mins 2, 6-26; p<0.01 for mins 3-5;). (**H**) Lever press ethograms during day 1 of FR1

conditioning. (**I**) Frequency distribution of lever inter-press intervals during the first and second

half of the conditioning session for day 1 (Two-Way RM ANOVA, <u>first half:</u> Interaction effect

$F_{(72, 1908)}$=2.706, ****P<0.0001; Time effect $F_{(2.888, 153.1)}$=20.67, P<0.0001; Group effect $F_{(2, 53)}$=3.458, P<0.05; Subject effect $F_{(53, 1908)}$=3.834; Dunnet's Multiple Comparison's test to controls,

30 s: p<0.05 for control vs Kv4.3 LOF; <u>second half:</u> Interaction effect $F_{(72, 1908)}$=1.012, P=0.4510;

Time effect $F_{(4.049, 214.6)}$=21.56, P<0.0001; Group effect $F_{(2, 53)}$=1.529, P<0.2263; Subject effect

$F_{(53, 1908)}$=2.389, P<0.0001). (**J**) Lever press ethograms during day 3 of FR1 conditioning. (**K**)

Frequency distribution of lever inter-press intervals during the first and second half of the

conditioning session for day 3 day 3 (Two-Way RM ANOVA, <u>first half:</u> Interaction effect $F_{(72, 1908)}$=1.923, P<0.0001; Time effect $F_{(3.798, 201.3)}$=29.10, P<0.0001; Group effect $F_{(2, 53)}$=4.482,

P<0.05; Subject effect $F_{(53, 1908)}$=1.470, P<0.05, Dunnet's Multiple Comparison's test to controls,

20 s: p<.0.05 Kv4.3LOF vs BKCa1.1LOF; 25 s: p<0.01 for Kv4.3LOF vs BKCa1.1LOF; 125 s:

p<0.05 control vs Kv4.3LOF; <u>Second half:</u> Interaction effect $F_{(72, 1908)}$=1.037, P=0.3950; Time

effect $F_{(2.581, 136.8)}$=19.91, P<0.0001; Group effect $F_{(2, 53)}$=0.6340, P<0.5344; Subject effect $F_{(53, 1908)}$=3.668, P<0.0001). (**F-I:** $N_{control}$=19; $N_{Kv4.3LOF}$=24; $N_{BKCa1.1LOF}$=13). **B-D** and **F** data are

presented as mean $\pm$ SEM. Cumulative lever press trace data presented as mean $\pm$ SEM and IPI

data presented as mean + SEM.

Figure 4 Differential action potential firing in putative dopamine neurons from control, Kv4.3 LOF, and BKCa1.1 LOF mice during instrumental conditioning. (**A**) Schematic of instrumental conditioning paradigm. (**B**) Illustration of optetrode Microdrive placement in the VTA and injections of AAV1-FLEX-ChR2-eYFP + AAV1-FLEX-SaCas9-sgRNA injection in DAT-IRES-Cre mice. (**C**) Average peri-event Z-score for putative dopamine neurons responsive to LP in the VTA for day 1 (top row) and day 5 (middle row and bottom row) (day 1: $N_{control}$=17 cells, 2 mice; $N_{Kv4.3LOF}$=39 cells, 5 mice; and $N_{BKCa1.1LOF}$=13 cells, 3 mice; day 5: $N_{control}$=8 cells, 3 mice; $N_{Kv4.3LOF}$=16 cells, 3 mice; and $N_{BKCa1.1LOF}$=15 cells, 3 mice). (**D**) Average area under the curve (AUC) for the peri-event Z-score action potential firing post-LP period on day 1 (One-way ANOVA, $F_{(2,66)}$= 6.29, P=0.0031, Fischer's direct comparison to controls, **p<0.01; $N_{control}$=17, 2 mice; $N_{Kv4.3LOF}$=39 cell, 5 mice; $N_{BKCa1.1LOF}$=13 cells, 3 mice) and day 5 (One-way ANOVA, $F_{(2,44)}$= 25.56, P<0.001, Fisher's direct comparison to controls, **p<0.05, $N_{control}$=8 cells, 2 mice; $N_{Kv4.3LOF}$=16 cells, 3 mice; $N_{BKCa1.1LOF}$=15 cells, 3 mice). (**E**) Average AUC for the Z-score of action potential firing for post-Rew period on day 1 and day 5 (day 1, One-way ANOVA, $F_{(2,66)}$= 4.07, P=0.022; Day 1: $N_{control}$=17 cells, 2 mice; $N_{Kv4.3LOF}$=39 cells, 5 mice; $N_{BKCa1.1LOF}$=13 cells, 3 mice; day 5: One-way ANOVA, $F_{(2,44)}$=3.875, P=0.028, Fischer's direct comparison to controls, **p<0.01; $N_{control}$=8 cells, 3 mice; $N_{Kv4.3LOF}$=16 cells, 3 mice; $N_{BKCa1.1LOF}$=15 cells, 3 mice). Averaged traces are presented as mean + SEM. Box and whisker graphs are presented as min-to-max.

Figure 5 Differential dopamine dynamics in the NAc of control, Kv4.3 LOF, and BKCa1.1 LOF mice during instrumental conditioning. (**A**) Average peri-event Z-score for dLight1.3b signals in the NAc for day 1 (top row) and day 5 (middle row and bottom row). (**B**) Average AUC for the perievent Z-score dLight1.3b signals post-LP period (3 s) (day 1, one-way ANOVA $F_{(2, 14)}$ =4.456, P<0.05; day 5, one-way ANOVA $F_{(2, 14)}$=2.867, P=0.09, Fisher's direct comparison to controls, *p<0.05). (**C**) Average AUC for the perievent Z-score dLight1.3b signals post-Rew period (day 1, one-way ANOVA $F_{(2, 14)}$ =6.701, P<0.01; day 5, one-way ANOVA $F_{(2, 14)}$=2.468, P=0.12, Fisher's direct comparison to controls, *p<0.05). (**D**) Zoomed average peri-event Z-score on day 5 for dLight1.3b in the NAc from panel A showing ramping in the pre-LP period. (**E**) Average ramp to peak Z-score from dLight1.3b signals in the NAc for pre-LP period (One-way ANOVA, $F_{(2,14)}$=3.848, p<0.05, Fisher's direct comparisons to controls *p<0.05). (**A-E**: $N_{control}$=6 mice; $N_{Kv4.3LOF}$=6 mice; $N_{BKCa1.1LOF}$=5 mice). Averaged traces are presented as mean + SEM. Box and whisker graphs are presented as min-to-max.

Figure 6 Extinction behavior and dopamine signal diametrically changes in Kv4.3 and BKCa1.1 LOF mice.

(**A**) Schematic of instrumental FR1 extinction paradigm. (**B**) Lever presses during days 1-5 of extinction training (Two-way RM ANOVA, Interaction effect $F_{(8, 136)}$=7.35, P<0.0001; Time effect $F_{(4, 136)}$=90.43, P<0.0001; Group effect $F_{(2, 34)}$=6.30, P=0.0047; Subject effect $F_{(34, 136)}$=2.911, P<0.0001; Dunnet's Multiple comparison's test to controls, ***p<0.001, ****p<0.0001, vs. control). (**C**) Frequency distribution of lever presses during the first and second half of the conditioning session for day 1 of extinction (Two-way RM ANOVA, <u>first half:</u> Interaction effect $F_{(72, 1152)}$=10.31, ****P<0.0001; Time effect $F_{(1.141, 36.51)}$=57.07, P<0.0001; Group effect $F_{(2, 32)}$=12.42, P<0.0001; Dunnet's multiple comparison's test to controls, 5 s: p<0.01 for control vs Kv4.3LOF; 45 s: p<0.05 for control vs BKCa1.1LOF; <u>second half:</u> Interaction effect $F_{(72, 1152)}$=5.047, P<0.0001; Time effect $F_{(1.459, 46.60)}$=48.96, P<0.0001; Group effect $F_{(2, 32)}$=5.851, P<0.01, Dunnet's multiple comparison's test to controls, 5 s: p<0.05 for control vs Kv4.3LOF). (**B-C**, $N_{control}$=13, $N_{Kv4.3LOF}$=10, and $N_{BKCa1.1LOF}$=12;) (**D**) Average Z-score for putative dopamine neurons in the VTA during FR1 day 1 following LP (top) compared to average Z-score following LP on day 1 of extinction (FR1 day 1, $N_{control}$=17 cells, 2 mice; $N_{Kv4.3LOF}$=39 cells, 5 mice; and $N_{BKCa1.1LOF}$=13 cells, 3 mice; Ext D1, $N_{control}$=10 cells, 3 mice; $N_{Kv4.3LOF}$=27 cells, 5 mice; and $N_{BKCa1.1LOF}$=23 cells, 3 mice) and average Z-score for dLight1.3b signals in the NAc during FR1 day 1 (top) and the first day of extinction (bottom) ($N_{control}$=6; $N_{Kv4.3LOF}$=6; and $N_{BKCa1.1LOF}$=5 mice). Comparison of AUC between Day 1 and Day 5 for Z-score action potential firing and dLight1.3b recordings for (**E**) Control (optetrode unpaired Student's *t*-test t_{25}=1.96, P=0.061; dLight1.3b paired Student's *t*-test t_5=4.17, **P<0.01), (**F**) Kv4.3 LOF (optetrode unpaired Student's *t*-test t_{65}=1.83, P=0.072; dLight1.3b paired Student's *t*-test t_5=1.398, P= 0.22) and (**G**)

980   BKCa1.1 LOF (optetrode unpaired Student's *t*-test t_{37}=4.28, **P<0.001, dLight1.3b paired

981   Student's *t*-test t_4=3.34, *P<0.05). Data are presented as mean $\pm$ SEM (**B-E**), or Box and Whisker

982   min-to-max graphs (**E-G**).

983
984
985   **Supplementary Materials**

986
987   Supplementary figures.

References:

1. R. A. Wise, Dopamine, learning and motivation. *Nat Rev Neurosci*. **5**, 483–494 (2004).
2. W. Schultz, Multiple Dopamine Functions at Different Time Courses. *Annu Rev Neurosci*. **30**, 259–288 (2007).
3. H. R. Kim, A. N. Malik, J. G. Mikhael, P. Bech, I. Tsutsui-Kimura, F. Sun, Y. Zhang, Y. Li, M. Watabe-Uchida, S. J. Gershman, N. Uchida, A Unified Framework for Dopamine Signals across Timescales. *Cell*. **183**, 1600-1616.e25 (2020).
4. M. W. Howe, P. L. Tierney, S. G. Sandberg, P. E. M. Phillips, A. M. Graybiel, Prolonged dopamine signaling in striatum signals proximity and value of distant rewards. *Nature*. **500**, 575–579 (2013).
5. A. Mohebi, J. R. Pettibone, A. A. Hamid, J.-M. T. Wong, L. T. Vinson, T. Patriarchi, L. Tian, R. T. Kennedy, J. D. Berke, Dissociable dopamine dynamics for learning and motivation. *Nature*. **570**, 65–70 (2019).
6. K. Farrell, A. Lak, A. B. Saleem, Midbrain dopamine neurons provide teaching signals for goal-directed navigation. *bioRxiv*. **22** (2021).
7. S. W. Johnson, R. A. North, Two types of neurone in the rat ventral tegmental area and their synaptic inputs. *J Physiol* (1992), doi:10.1113/jphysiol.1992.sp019136.
8. F. White, X.- Hu, M. Marinelli, C. Rudick, Excitability of Dopamine Neurons: Modulation and Physiological Consequences. *CNS Neurol Disord Drug Targets*. **5**, 79–97 (2006).
9. Z. M. Khaliq, B. P. Bean, Dynamic, nonlinear feedback regulation of slow pacemaking by A-type potassium current in ventral tegmental area neurons. *Journal of Neuroscience*. **28** (2008), doi:10.1523/JNEUROSCI.2237-08.2008.
10. T. Kimm, Z. M. Khaliq, B. P. Bean, Differential regulation of action potential shape and burst-frequency firing by BK and Kv2 channels in substantia nigra dopaminergic neurons. *Journal of Neuroscience*. **35** (2015), doi:10.1523/JNEUROSCI.5291-14.2015.
11. M. E. Soden, G. L. Jones, C. A. Sanford, A. S. Chung, A. D. Güler, C. Chavkin, R. Luján, L. S. Zweifel, Disruption of Dopamine Neuron Activity Pattern Regulation through Selective Expression of a Human KCNN3 Mutation. *Neuron*. **80** (2013), doi:10.1016/j.neuron.2013.07.044.
12. J. X. Yee, A. Rastani, M. E. Soden, The potassium channel auxiliary subunit Kvβ2 (*Kcnab2*) regulates Kv1 channels and dopamine neuron firing. *J Neurophysiol*. **128**, 62–72 (2022).
13. A. C. Hunker, M. E. Soden, D. Krayushkina, G. Heymann, R. Awatramani, L. S. Zweifel, Conditional Single Vector CRISPR/SaCas9 Viruses for Efficient Mutagenesis in the Adult Mouse Nervous System. *Cell Rep* (2020), doi:10.1016/j.celrep.2020.02.092.
14. A. S. Chung, S. M. Miller, Y. Sun, X. Xu, L. S. Zweifel, Sexual congruency in the connectome and translatome of VTA dopamine neurons. *Sci Rep*. **7** (2017), doi:10.1038/s41598-017-11478-5.
15. P. D. Shepard, B. S. Bunney, Repetitive firing properties of putative dopamine-containing neurons in vitro: regulation by an apamin-sensitive Ca2+-activated K+ conductance. *Exp Brain Res*. **86** (1991), doi:10.1007/BF00231048.
16. S. Q. Lima, T. Hromádka, P. Znamenskiy, A. M. Zador, PINP: A new method of tagging neuronal populations for identification during in vivo electrophysiological recording. *PLoS One*. **4** (2009), doi:10.1371/journal.pone.0006099.
17. R. Rahamimoff, A. Butkevich, D. Duridanova, R. Ahdut, E. Harari, S. G. Kachalsky, Multitude of ion channels in regulation of transmitter release. *Philos Trans R Soc Lond B Biol Sci*. **354**, 281–288 (1999).
18. K. J. Burke, K. J. Bender, Modulation of Ion channels in the Axon: Mechanisms and function. *Front Cell Neurosci*. **13** (2019), , doi:10.3389/fncel.2019.00221.
19. N. C. Klapoetke, Y. Murata, S. S. Kim, S. R. Pulver, A. Birdsey-Benson, Y. K. Cho, T. K. Morimoto, A. S. Chuong, E. J. Carpenter, Z. Tian, J. Wang, Y. Xie, Z. Yan, Y. Zhang, B. Y. Chow, B. Surek, M. Melkonian, V. Jayaraman, M. Constantine-Paton, G. K. S. Wong, E. S. Boyden, Independent optical excitation of distinct neural populations. *Nat Methods*. **11** (2014), doi:10.1038/nmeth.2836.
20. J. Wolfart, H. Neuhoff, O. Franz, J. Roeper, Differential expression of the small-conductance, calcium-activated potassium channel SK3 is critical for pacemaker control in dopaminergic midbrain neurons. *Journal of Neuroscience*. **21** (2001), doi:10.1523/jneurosci.21-10-03443.2001.
21. Z. Niday, B. P. Bean, BK channel regulation of afterpotentials and burst firing in cerebellar purkinje neurons. *Journal of Neuroscience*. **41** (2021), doi:10.1523/JNEUROSCI.0192-20.2021.
22. J. Clatot, N. Neyroud, R. Cox, C. Souil, J. Huang, P. Guicheney, C. Antzelevitch, Inter-regulation of kv 4.3 and voltage-gated sodium channels underlies predisposition to cardiac and neuronal channelopathies. *Int J Mol Sci*. **21** (2020), doi:10.3390/ijms21145057.

23. Z. M. Khaliq, B. P. Bean, Pacemaking in dopaminergic ventral tegmental area neurons: Depolarizing drive from
background and voltage-dependent sodium conductances. *Journal of Neuroscience*. **30** (2010),
doi:10.1523/JNEUROSCI.0143-10.2010.
24. M. E. Soden, A. S. Chung, B. Cuevas, J. M. Resnick, R. Awatramani, L. S. Zweifel, Anatomic resolution of
neurotransmitter-specific projections to the VTA reveals diversity of GABAergic inputs. *Nat Neurosci* (2020),
doi:10.1038/s41593-020-0657-z.
25. B. B. Gore, M. E. Soden, L. S. Zweifel, Manipulating gene expression in projection-specific neuronal
populations using combinatorial viral approaches. *Curr Protoc Neurosci* (2013),
doi:10.1002/0471142301.ns0435s65.
26. A. C. Hunker, L. S. Zweifel, Protocol to Design, Clone, and Validate sgRNAs for In Vivo Reverse Genetic
Studies. *STAR Protoc* (2020), doi:10.1016/j.xpro.2020.100070.
27. F. Wang, J. Flanagan, N. Su, L. C. Wang, S. Bui, A. Nielson, X. Wu, H. T. Vo, X. J. Ma, Y. Luo, RNAscope: A
novel in situ RNA analysis platform for formalin-fixed, paraffin-embedded tissues. *Journal of Molecular
Diagnostics*. **14** (2012), doi:10.1016/j.jmoldx.2011.08.002.
28. L. Erben, A. Buonanno, Detection and Quantification of Multiple RNA Sequences Using Emerging
Ultrasensitive Fluorescent In Situ Hybridization Techniques. *Curr Protoc Neurosci*. **87** (2019),
doi:10.1002/cpns.63.
29. J. T. Ting, T. L. Daigle, Q. Chen, G. Feng, Acute brain slice methods for adult and aging animals: Application
of targeted patch clamp analysis and optogenetics. *Methods in Molecular Biology*. **1183** (2014),
doi:10.1007/978-1-4939-1096-0_14.
30. Y. S. Jo, G. Heymann, L. S. Zweifel, Dopamine Neurons Reflect the Uncertainty in Fear Generalization.
Neuron (2018), doi:10.1016/j.neuron.2018.09.028.
31. J. M. Ehrich *et al.*, Kappa Opioid Receptor-Induced Aversion Requires p38 MAPK Activation in VTA
Dopamine Neurons. *J Neurosci* **35**, 12917-12931 (2015).
32. R. Spanagel, A. Herz, T. S. Shippenberg, The effects of opioid peptides on dopamine release in the nucleus
accumbens: an in vivo microdialysis study. *J Neurochem* **55**, 1734-1740 (1990).
33. R. Al-Hasani *et al.*, Distinct Subpopulations of Nucleus Accumbens Dynorphin Neurons Drive Aversion and
Reward. *Neuron* **87**, 1063-1077 (2015).
34. S.C. Gantz and B.P. Bean. Cell-Autonomous Excitation of Midbrain Dopamine Neurons by
Endocannabinoid-Dependent Lipid Signaling. *Neuron*. 93, 1375-1387. (2017).
35. R. Gomez, L.F. Maglio, A.J. Gonzalez-Hernandez, B. Rivero-Piréz, D. Bartolomé-Martin, T. Giraldez.
NMDA receptor–BK channel coupling regulates synaptic plasticity in the barrel cortex. *PNAS*. 118 (35) (2021).
36. N.C. Klapoetke *et al*., Independent Optical Excitation of Distinct Neural Populations. *Nature Methods* **3**, 338-
46 (2014).
1077

FIGURE 1

1079
1080

FIGURE 3

FIGURE 4

1090   **FIGURE 5**

dLight fiber photometry in NAc during FR1 reinforcement learning

A Control / Kv4.3 LOF / BKCa1.1 LOF — Day 1, Day 5 traces and Day 5 Trial # heatmaps

B LP — Day 1, Day 5 — AUC (z-score) — Control, Kv4.3 LOF, BKCa1.1 LOF

C Rew — Day 1, Day 5 — AUC — Control, Kv4.3 LOF, BKCa1.1 LOF

Pre-press ramp day 5 of FR1 reinforcement learning

D Control / Kv4.3 LOF / BKCa1.1 LOF

E Peak z-score — Control, Kv4.3 LOF, BKCa1.1 LOF

1091
1092

FIGURE 6