**Endothelial cell signature in muscle stem cells validated by VEGFA-FLT1-AKT1 axis**

**promoting survival of muscle stem cell**

Authors/Affiliations

Mayank Verma[a,b,c,d], Yoko Asakura[b,c,d], Xuerui Wang[b,c,d], Kasey Zhou[b,c,d], Mahmut Ünverdi[b,c,d],

Allison P. Kann[e,f], Robert S. Krauss[e,f] and  Atsushi Asakura[b,c,d]

Department of Pediatrics & Neurology, Division of Pediatric Neurology, The University of

Texas Southwestern Medical Center, TX, USA[a].

Stem Cell Institute[b], Paul & Sheila Wellstone Muscular Dystrophy Center[c], Department of

Neurology[d], University of Minnesota Medical School, MN, USA

Department of Cell, Developmental, and Regenerative Biology[e], and Graduate School of

Biomedical Sciences[f], Icahn School of Medicine at Mount Sinai, NY, USA.

Conflict of Interest

MV and AA are listed as inventors on a patent for anti-FLT1 antibody mediated therapy for

DMD. AA received consulting fees from ShireHGT.

**Abstract**

Endothelial and skeletal muscle lineages arise from common embryonic progenitors. Despite

their shared developmental origin, adult endothelial cells (ECs) and muscle stem cells (MuSCs)

(satellite cells) have been thought to possess distinct gene signatures and signaling pathways.

Here we shift this paradigm by uncovering how adult MuSC behavior is affected by the

expression of a subset of EC transcripts. We used several computational analyses including

single-cell RNAseq to show that MuSCs express low levels of canonical EC markers. We

demonstrate that MuSC survival is regulated by one such prototypic

endothelial signaling pathway (VEGFA-FLT1). Using pharmacological and genetic gain- and

loss-of-function studies, we identify the FLT1-AKT1 axis as the key effector underlying

VEGFA-mediated regulation of MuSC survival. All together, our data support that the VEGFA-

FLT1-AKT1 pathway promotes MuSC survival during muscle regeneration, and highlights how

the minor expression of select transcripts is sufficient for affecting cell behavior.

## Introduction

Skeletal muscle and endothelial cells (ECs) and their progenitors from the trunk and limbs are derived from the somites during early developments. Previous works demonstrated the existence of bipotent progenitors which express both Pax3 and FLK1 (Eichmann et al., 1993; Ema et al., 2006; Esner et al., 2006; Kardon et al., 2002; Tozer et al., 2007). These bipotent progenitors migrate into trunk and limb buds from ventrolateral region of the somites to generate MyoD(+) myogenic cells followed by skeletal muscle and PECAM1(+) ECs followed by vasculatures (Hutcheson and Kardon, 2009; Kardon et al., 2002; Lagha et al., 2009; Mayeuf-Louchart et al., 2014; Mayeuf-Louchart et al., 2016). In addition, FLK1(+) cells give rise to myogenic cells during development and oncologic transformation (Drummond and Hatley, 2018; Mayeuf-Louchart et al., 2014; Motoike et al., 2003). Lastly, multipotent mesoangioblasts, vessel-associated stem cells, have been identified in embryonic dorsal aorta (Minasi et al., 2002). These cells are able to differentiate into several types of mesodermal tissues including skeletal muscle and ECs (Roobrouck et al., 2011). Interestingly, these myogenic cells show the same morphology as muscle satellite cells (MuSCs), stem cell populations for skeletal muscle, and express a number of myogenic and EC markers such as MyoD, M-cadherin, FLK1 and VE-cadherin (De Angelis et al., 1999). However, it is not clear whether adult MuSCs derived from these bipotent progenitors still maintain canonical EC signals. Curiously, blood vessel-associated myoendothelial cell progenitors that express both myogenic and EC markers, and are able to differentiate into myogenic cells following transplantation have been identified in the interstitial spaces of both murine and human adult skeletal muscle (Tamaki et al., 2002; Zheng et al., 2007; Huang et al., 2014). However, the relationship between these myoendothelial cell progenitors and MuSCs remains unclear.

Vascular endothelial growth factor (VEGF), specifically VEGFA modulates many biological aspects including angiogenesis through its two receptors, FLT1 and FLK1. Although FLK1 possesses stronger signaling capability and the major signaling receptor tyrosine kinase (RTK) for VEGFA, FLT1 has considerably higher affinity for VEGF but weaker cytoplasmic signaling capability. In normal tissue, FLT1 acts as a decoy receptor and a sink trap for VEGF thereby preventing excessive normal and pathological angiogenesis. In addition, there are two co-receptors for VEGFA (NRP1 and NRP2) that function with FLK1 to modulate VEGFA signaling. While VEGF signaling has been extensively studied for its role in development, proliferation, and survival of endothelial cells (ECs), its role in non-vascular systems such as neuron and bone has only recently been appreciated (Liu et al., 2012; Okabe et al., 2014; Poesen et al., 2008). Skeletal muscle tissue is the most abundant producer of VEGFA in the body. It has already been extensively studied in the skeletal muscle fibers in models of *Vegfa* knockout mice (Olfert et al., 2009; Tang et al., 2004; Wagner et al., 2006) as well as *Vegfa* overexpression (Arsic et al., 2004; Bouchentouf et al., 2008; Messina et al., 2007; Yan et al., 2005).

Adult skeletal muscle also contains the tissue resident muscle stem cell population, termed MuSCs, which mediate postnatal muscle growth and muscle regeneration (Motohashi and Asakura, 2014). After muscle injury, quiescent MuSCs initiate proliferation to produce myogenic precursor cells, or myoblasts. The myoblasts undergo multiple rounds of cell division before terminal differentiation and formation of multinucleated myotubes by cell fusion. Importantly, the MuSC-derived myoblasts also express VEGFA, which has been shown to increase the proliferation of myoblasts (Christov et al., 2007). Our data obtained from genetical model mice demonstrated that MuSCs express abundant VEGFA, which recruits ECs to establish vascular

niche for MuSC self-renewal and maintenance (Verma et al., 2018). In addition, VEGFA and its

receptors are expressed in the myoblast cell line, C2C12 cells, and the signaling can induce cell

migration and protect apoptotic cell during myogenic differentiation *in vitro* (Bryan et al., 2008;

Germani et al., 2003; Mercatelli et al., 2010). However, it is not clear whether MuSCs also

express VEGF receptors and if cell-autonomous VEGFA signaling plays an essential roles in

MuSC fucction during muscle regeneration *in vivo*.

We have previously shown that *Flt1* heterozygous gene knockout and conditional

deletion of *Flt1* in ECs display increased capillary density in skeletal muscle, indicating the

essential roles for *Flt1* in adult skeletal muscle. More importantly, when crossed with the

Duchenne muscular dystrophy (DMD) model *mdx* mice, these mice show both histological and

functional improvements of their dystrophic phenotypes. This was due to the effect of increased

ECs leading to an increase in MuSCs (Verma et al., 2010; Verma et al., 2019). However, the

effect of VEGFA on MuSC *in vivo* remained unknown. We found that MuSCs express low levels

of canonical EC markers including VEGF receptors using single cell transcriptomics. Therefore,

we examined the effects of VEGFA on MuSCs and show that it has a drastic effect on cell

survival in the via its receptor FLT1 by signaling through AKT1.

**Results**

**EC gene signal including *Vegf receptors* in MuSCs**

EC signatures in MuSCs has been seen in several gene expression data sets (Figure S1A-D,

Table S1) (Fukada et al., 2007; Ryall et al., 2015; van Velthoven et al., 2017; Yue et al., 2020).

However, with the lack of EC control, we questioned whether these were true expression or

artifact. To isolate EC and MuSC populations, we first crossed the *Flk1$^{+/GFP}$* mice to label the

ECs of the vasculature (Ema et al., 2006) and the $Pax7^{+/CreERT2}:ROSA26^{+/Loxp-stop-Loxp-tdTomato}$

($Pax7^{+/CreERT2}:R26R^{+/tdT}$) mice to mark the MuSC lineage (Murphy et al., 2011; Verma et al.,

2018). We performed bulk RNA sequencing (RNAseq) on FACS sorted ECs and MuSCs as well

as freshly isolated single muscle fibers (Figure 1A, S1E-G). We found that single muscle fibers

routinely have ECs fragments attached to the fiber (Figure S1G) and so we removed such fibers

based of Flk1$^{GFP}$ expression from the samples collected for sequencing. We surveyed for

canonical genes for each cell type (Figure 1B) and found minimal but reliable expression of

canonical ECs genes such as *Pecam1*, *Cdh5*, *Kdr*, and *Flt1* in MuSCs.

It is possible that these EC signatures detecting in MuSCs were not due to small amounts

of contaminating ECs with very high expression of the canonical EC genes skewing the average

expression in MuSC RNA samples. To rule out this possibility, we performed single cell

RNAseq (scRNAseq) on MuSCs and ECs isolated from mouse hind limb muscle from both basal

condition and 3-days post injury to look at both quiescent and activated MuSCs from the reporter

mice specified above (Figure 1A). We could reliably delineate injured and activated MuSCs via

side and forward scatter (Figure S1H, S1I). We FACS isolated cells from both days separately

and spiked in 20% of the ECs into the MuSCs, and performed scRNAseq for each time point

(Figure 1A). We performed sequencing with ~300K read/cell compared with the commonly used

sequencing with 60K reads/cell, in order to maximize the possibility of detecting low abundance

transcripts (Zhang et al., 2020). In the aggregated dataset, the MuSCs showed low overlap

between D0 and D3 owing to the different stages of the myogenic differentiation cycle, while the

ECs clusters showed near perfect overlap (Figure 1C, 1D). While drastic morphological changes

in ECs have been shown during muscle regeneration (Hardy et al., 2016), transcriptomic changes

are much more tapered, especially compared with MuSCs (Latroche et al., 2017). We were able

to deconvolve the quiescent MuSCs from the activated and differentiating MuSCs, ECs, and

other cell types from gene signatures. (Figure S1J). We also introduced an artificial chromosome

loci in our sequencing reference genome to allow for mapping of custom genes such as *eGFP-*

*SV40* and *tdTomato-WPRE-BGHPolyA* transgenes and were able to confirm high expression of

these genes to their respective clusters (Figure 1E). Importantly, data from scRNAseq were able

to recapitulate the minimal expression of canonical EC genes in the MuSC clusters such as *Cdh5*

(Figure 1E) as seen in our Bulk RNAseq results (Figure 1B). These included the *Vegf receptors*

*Flk1* (*Kdr*) and *Flt1* (Figure 1E).

As a quality control measure, we mapped parts of the three transgene genes, *eGFP* from

*Flk1$^{+/GFP}$*, and *tdTomato* and *CreERT2* from *Pax7$^{+/CreERT2}$:R26R$^{+/tdT}$* that can be detected in ECs

and MuSCs, respectively, as expected (Figure S1L). Surprisingly, we also found *eGFP* in the

MuSCs and *tdTomato* in EC fraction, while the *CreERT2* expression remained mainly restricted

to the MuSCs (Figure S1K). FACS analysis and FACS-sorted cells confirmed that GFP(+) and

tdTomato(+) cells are exclusively restricted as ECs and MuSCs, respectively (Figure S1E, S1F).

Therefore, we hypothesized that this was due to the ambient free mRNA from the digested cells

that is intrinsic to any droplet based single cell sequencing platform. By using SoupX (Young

and Behjati, 2020), we performed careful background subtraction using genes expressed

exclusively in myofibers as our negative control and genes validated by *in situ* hybridization as a

positive control (Figure S1K) (Kann and Krauss, 2019). We observed decreased but sustained

*eGFP* expression in the MuSC fraction and *tdTomato* expression in the EC fraction after SoupX

subtraction (Figure S1K). In addition, the EC signatures such as *Cdh5* expression in the MuSC

fraction was also sustained. These results conclude that MuSCs contain mRNAs from canonical

ECs genes. We showed that the canonical EC genes were broadly expressed in the myogenic

cells in our dataset (Figure 1E).

Since detection of rare subpopulation in single cell dataset is a factor of cell numbers, we

re-analyzed previously published dataset with 2,232 myogenic cells across different states (Torre

et al., 2018; De Micheli et al., 2020). We were able to classify cell as quiescent, proliferative vs.

differentiating states based on the expression of *Calcr*, *Cdk1* and *Myog*, respectively (Figure

S1L). We noticed that EC prototypic markers such as *Flt1* are broadly expressed with small

amounts in MuSCs. Complementary data from different laboratories showed the clear expression

of EC prototypic markers such as *Cdh5*, *Flt1* and *Kdr*, using microarrays and Bulk-RNAseq

(Figure S1A, S1B; Fukada et al., 2007; Ryall et al., 2015). Recently, RNAseq data from fixed

quiescent, early activated and late activated MuSCs show that *Flt1* may be transiently

upregulated during the early activation process (Figure S1C) (Yue et al., 2020). To confirm

whether the EC gene mRNAs were transcribed from MuSCs, we utilized previously published

MuSC nascent RNA transcriptome from TU-tagged samples (Gay et al., 2013; van Velthoven et

al., 2017). As expected, *Myh1* was represented in the whole muscle but was absent in the TU-

tagged MuSCs (Figure 1F), indicating that the nascent MuSCs were devoid of cellular

contamination from other cells in the muscle. Inversely, the nascent MuSC transcript was over-

represented for MuSC related genes such as *Calcr* and *Sdc4*. Interestingly, we were able to detect

EC genes such as *Kdr* and *Pecam1* in the TU-tagged MuSC samples indicating that they were

actively transcribed by MuSCs (Figures 1F, S1D).

We also verified the expression of *Vegfr* genes (*Kdr*, *Flt1*, *Nrp1* and *Nrp2*) in MuSCs

using RT-qPCR (Figure 1G). In addition, we verified the expression of *Flt1* by performing *in-*

*situ* hybridization using RNAScope on MuSC on whole muscle fiber, which we currently believe

to be the gold standard for expression studies (Figure 1H). Finally, in MuSC-derived myoblasts,

NRP1and NRP2 expression was detectable with comparable intensity compared with EC cell

line, while FLT1 expression was detectable with lower intensity compared with EC cell line

(Figure 1I). By contrast, PECAM1, VE-Cadherin and FLK1 expression, which was clearly

detected in EC cell line, was undetectable level in myoblasts. Taken together, these data indicate

that there are both transcripts of these EC canonical genes and EC canonical proteins in MuSCs.

**VEGFA induces proliferation and cell survival but not differentiation in myoblasts**

Since VEGFRs were expressed in MuSCs in small amounts and their ligand, VEGFA, was

highly expressed in MuSCs (Figure 2A; Verma et al., 2018), we wanted to investigate whether

there were any biological effects to induction by VEGFA. We found that treatment with VEGFA

could increase proliferation of MuSC-derived myoblasts at low dose but inhibit proliferation at

high dose of VEGFA, a phenomenon that has been previously described in ECs (Noren et al.,

2016) (Figure S2A). We saw no effect on differentiation by VEGFA as evaluated by myosin

heavy chain (MyHC) staining, fusion index and RT-qPCR (Figure S2B, S2C, S2D). By contrast,

crystal violet staining showed that VEGFA could significantly increase survived cell number of

myoblasts following UV-mediated apoptotic cell death induction (Figure S2E, S2F). To

investigate apoptosis in detail, we optimized Annexin V assay following thapsigargin-mediated

endoplasmic reticulum (ER)-stress (Hirai et al., 2010) so that we could study deviation at ~ED50

while still performing experiments to remove the confounding variable to proliferation from the

experimental setup (Figure S2G and S2H). We had previously shown that MuSCs are the

predominant cells that secrete VEGFA in skeletal muscle (Figure 2A; Verma et al., 2018) and

while adding exogenous VEGFA did not improve cell survival, blocking VEGFA via a soluble

form of FLT1-FC increased the number of apoptotic and necrotic myoblasts *in vitro* (Figure 2C-

E).

**VEGFA-facilitated cell survival in MuSC-derived myoblasts is mediated through FLT1**

To characterize the VEGF receptor responsible for the anti-apoptotic effect of VEGFA on

MuSC-derived myoblasts, we used pharmacological inhibitors of the VEGF receptors (Figure

2F). We used blocking antibody for the VEGF receptors FLT1 (anti-FLT1 antibody), small

molecule inhibitors for FLK1 (SU4502 and ZM306416) and the FLK1 co-receptor NRP1

(EG00229) following thapsigargin induction (Figure 2D). Surprisingly, inhibiting FLK1, the

major signaling RTK for VEGFA, had no effect on myoblasts survival following thapsigargin

induction (Figure 3E). By contrast, blocking FLT1 via blocking antibody greatly decreased the

survival of myoblasts following thapsigargin induction (Figure 3E). To confirm this interesting

result using genetic tools, we obtained myoblasts with *Pax7-CreER*-inducible deletion of *Flt1*

mice (*Pax7$^{+/CreER}$:Flt1$^{Loxp/Loxp}$* or *MuSC-Flt1$^{\Delta/\Delta}$*) and the control mice (*Pax7$^{+/+}$:Flt1$^{Loxp/Loxp}$*). *In*

*vitro* 4-OHT-mediated genetic deletion of *Flt1* (MuSC-*Flt1$^{\Delta/\Delta}$*) resulted in down-regulation of

*Flt1* RNA and FLT1 protein expression (Figure S2I, S2J), and increased spontaneous apoptotic

cell death even without induction of apoptosis (Figure 2F). By contrast, *Flt1* deletion did not

affect cell proliferation assessed by EdU staining or myogenic differentiation assessed by MyHC

staining (Figure S2K-M). When thapsigargin-induced apoptosis was induced, the *MuSC-Flt1$^{\Delta/\Delta}$*

myoblasts had increased apoptosis that was not responsive to exogenous VEGFA (Figure 2G).

**AKT signaling is involved in apoptosis of muscle stem cells.**

VEGFA signaling is mediated through Extracellular signal-Regulated Kinase (ERK), p38

Mitogen-Activated Protein Kinase (MAPK), and Protein kinase B (AKT). In ECs, VEGFA is

known to protect cells from apoptosis via AKT (Domigan et al., 2015; Lee et al., 2007).

However, it is not known whether VEGFA can similarly activate AKT in MuSC-derived

myoblasts. While the role of AKT has been explored in proliferation and differentiation in

myoblasts, its role in apoptosis has not been well characterized (Loiben et al., 2017). We

assessed for AKT activation via phosphorylated AKT (pAKT) in MuSC-derived myoblasts. We

found that exogenous VEGFA could induce AKT phosphorylation (pAKT) (Figure 2H, 2I). This

response was blunted in $MuSC$-$Flt1^{\Delta/\Delta}$ myoblasts and was no longer responsive to VEGFA

(Figure 2H, 2I). Lastly, we wanted to confirm that AKT activation could improve myoblast

survival. We infected lentiviral $E4ORF1$ or $MyrAKT$ vectors in myoblasts (Figure 2J), both of

which gene products have been shown to specifically activate AKT without activating ERK or

p38 (Kobayashi et al., 2010). We found that overexpression of either of these genes improved

cell survival compared with the control $in$ $vitro$ following induction of apoptosis via thapsigargin

(Figure 2J). These data establish FLT1-AKT as the cascade linking VEGFA to apoptosis in

MuSC-derived myoblasts during muscle regeneration (Figure 2K).

**VEGFA-FLT1 pathway protects MuSCs from apoptosis $in$ $vivo$**

Endogenous and exogenous VEGFA have been shown to regulate cell survival and protect ECs

from apoptosis $in$ $vitro$ (Gerber et al., 1998; Lee et al., 2007). To assess whether additional

VEGFA had an effect of MuSC behaviors $in$ $vivo$, we used mice carrying the $VEGFA^{+/Hyper}$ allele

for injury-mediated tibialis anterior (TA) muscle regeneration following $BaCl_2$ injection (Figure

3A, 3B) (Miquerol et al., 1999). MuSC-derived myoblasts from $Pax7^{+/tdT}$:$VEGFA^{+/Hyper}$ mice

showed around 2.8 fold increased expression of *Vegfa* but not the *Vegfrs* compared with

myoblasts from wild-type mice (Figure S3A). Interestingly, while treatment with VEGFA alone

had no effect on apoptosis *in vitro*, the MuSCs from *Pax7+/tdT:VEGFA+/Hyper* mice showed

decreased cell death in regenerating muscle by 1 day following BaCl$_2$ injection (Figure 3C).

Consequently, single muscle fibers from *Pax7+/tdT:VEGFA+/Hyper* mice showed increased number

of MuSCs, compared with those from *Pax7+/tdT:VEGFA+/+* mice by 28 days following BaCl$_2$

injection (Figure 3D). In addition, muscle regeneration was promoted in *VEGFA+/Hyper* mice in

the early and late muscle repair processes as judged by fiber diameter (Figures 3B, 3E, 3F, S3B-

D).

We then performed the reciprocal experiment to investigate the consequence of *Vegfa*

loss in MuSCs *in vivo*, and utilized MuSC-specific *Vegfa* knockout mice

(*Pax7+/CreER:VEGFALoxp/Loxp*). We have previously shown that vasculature in the MuSC-

*VEGFAΔ/Δ* mouse muscle is perturbed and the proximity between the MuSC and EC is increased

(Verma et al., 2018). However, the functional consequences of this remained unknown. We

confirmed that clear downregulatin of VEGFA protein in MuSC-derived myoblasts isolated from

MuSC-*VEGFAΔ/Δ* mice (Figure S3E). We noticed that deletion of *Vegfa* in MuSCs in the *MuSC-*

*VEGFAΔ/Δ* mouse muscle led to an increase in the proportion of dead MuSCs following BaCl$_2$

injection (Figure 3G). Consequently, the number of MuSCs in the MuSC-*VEGFAΔ/Δ* muscle were

significantly reduced following recovery after injury (Figure 3H). There was no difference in the

MuSC numbers in MuSC-*VEGFAΔ/Δ* muscle at homeostasis. In addition, the muscle had a

regenerative defect as indicated by the shift in fiber size distribution and increased adipose

following muscle injury (Figure 3B, 3I, 3J, S3F-H). While a limitation of this experiment is that

the MuSC fusion into the fiber also deletes *Vegfa* from the fiber themselves, muscle fiber

specific deletion of *Vegfa* has not shown to have an effect on fiber size (Delavar et al., 2014).

These data indicate that cell intrinsic VEGFA improves cell survival of MuSCs and that loss of

MuSC-derived VEGFA results in reduced muscle regeneration.

Since FLT1 but not FLK1 was detected in MuSCs and MuSC-derived myoblasts, we

asked whether the *Flt1* had an effect on MuSC survival *in vivo,* we evaluated cell death in

MuSCs from *MuSC-Flt1$^{\Delta/\Delta}$* mouse muscle. We induced muscle regeneration using $BaCl_2$ for 1

day and assessed for cell death in MuSCs. As seen *in vitro*, we found that loss of *Flt1* in MuSCs

(Figure S2I, S2J) resulted in increased cell death during early regeneration (Figure 3K).

Consequently, single muscle fibers from *MuSC-Flt1$^{\Delta/\Delta}$* mice showed decreased number of

MuSCs, compared with those from *MuSC-Flt1$^{+/+}$* mice by 28 days following $BaCl_2$ injection

(Figure 3L). We also examined the long-term *in vivo* consequence of deleting *Flt1* from MuSC.

There was no significant muscle phenotype in MuSC-*Flt1$^{\Delta/\Delta}$* muscle at homeostasis (Figures 3B,

3M, 3N). However, following injury, the MuSC-*Flt1$^{\Delta/\Delta}$* muscle had a modest regenerative defect

as indicated by the shift in fiber size distribution following muscle injury (Figures 3B, 3M, 3N,

S3J-N).

**VEGFA-FLT1 pathway regulates muscle pathology in DMD model mice**

While angiogenic defects have been reported in the *mdx* mice as well as in golden retrieval

muscular dystrophy (GRMD; canine model of DMD) (Latroche et al., 2015; Verma et al., 2019,

2010; Kodippili et al., 2021; Podkalicka et al., 2021), it is not clear whether VEGF family and its

receptors are implicated in human dystrophinopathies. We probed the VEGF ligands and

receptors in microarrays (Table S1) from skeletal muscles and MuSCs from *mdx* mice

(Pallafacchina et al., 2010; Tseng et al., 2002) and skeletal muscles from the *GRMD* (Vieira et al.,

2015). *Vegfa* was downregulated in both models (Figure S4A). *Flt1* was also downregulated in

*GRMD* but not *mdx* muscles. To examine whether VEGF signaling is altered in DMD patients,

we performed gene expression analysis on previously available data from microarrays from

patients with DMD (Chen et al., 2000). We also aggregated and probed microarray data from

muscle biopsies of patients with various neuromuscular diseases or of healthy individuals after

exercise (Bakay et al., 2006). In the microarray data, *Vegfa* expression was increased after an

acute bout of exercise, and *Vegfa* expression was reduced in ALS muscle, BMD muscle, as well

as both early and late phases of DMD muscle (Figure S4A). These data indicate that *Vegfa*

expression is decreased in dystrophinopathy, and thus may benefit people with DMD by

increasing VEGFA as a therapeutic target.

Therefore, we crossed the *MuSC-Flt1^{Δ/Δ}* mice with the chronically regenerating DMD

model mice (*mdx*) to generate *mdx:MuSC-Flt1^{Δ/Δ}* mice, and analyzed long term effects of *Flt1*

deletion (Figure 4A, S4B). Importantly, we found a significant decrease in fiber diameter and

increased fibrosis (Figures 4B-D, S4C) in TA muscle accompanied by a physiological decease in

muscle perfusion as shown by laser Doppler flow at 12 months age (Figure 4E) as well as a

functional decline in muscle strength as judged by grip strength both acutely and chronically

(Figure 4F).

By contrast, when we crossed the *VEGFA^{+/Hyper}* mice with *mdx* mice (Figure 4A, S4D),

we noticed a significant increase in fiber diameter and decreased fibrosis (Figures 4G-J, S4E) in

both TA and diaphragm muscle of *mdx:VEGFA^{+/Hyper}* mice accompanied by a physiological

increase in muscle perfusion as shown by laser Doppler flow at 12 months age (Figure 4K) as

well as a functional increase in muscle strength as judged by grip strength (Figure 4L). These

data indicate that VEGFA-FLT1 axis is a therapeutic target for the pathology seen in the *mdx*

mice

**Discussion**

In this report, we performed bulk and single cell RNA sequencing on MuSCs and ECs. Since

deep reads can significantly reduce the effect of the technical noise in scRNAseq, it can improve

estimation of minor transcriptional state of a given cell (Zhang, 2020). Unexpectedly, we found

that MuSCs broadly express EC prototypic markers in small amounts and used multiple different

bioinformatics techniques to validate the results. While similar phenomenon in myogenic cells

during development and existence of blood vessel-associated myoendothelial cells in the adult

skeletal muscle have been previously described, no functional follow up as been performed

leading to the questions whether these minor expression profiles were artifacts or functional (De

Angelis et al., 1999; Minasi, 2002; Tamaki et al., 2002; Zheng et al., 2007; Roobrouck, 2011;

Huang et al., 2014; Charville et al., 2015; Goel et al., 2017; Giordani et al., 2018). Our goal was

to see whether this small expression pattern had biological consequences. We ultimately decided

to use *Flt1* for further investigations and used RNAscope and immunostaining to validate its

expression in MuSCs. We found that *Flt1* indeed exerts a biological function even at a low

expression. Signaling through VEGFA-FLT1-AKT can improve cell survival in MuSCs both *in*

*vivo* and *in vitro.*

On a grander scale, our finding of EC prototype markers expressed in MuSC calls into

two questions 1) the genes that we used to specify cellular identities and 2) the cellular identity

of MuSCs and ECs. The former is important as when we experimentally label, induce or perform

Cre-mediated gene knockout experiments based on our assumptions of different gene expression

results which may be confounded for these low expressing genes. For example, we have

previously investigated both *Flt1* and *Kdr* in mouse muscle using three different reporters and

found them to be negative in MuSCs, thereby disregarding their cell-autonomous effect when

evaluating global knockouts (Verma et al., 2018, 2010). It is also possible that EC mRNAs are

results of transcription from the cell or a result of mRNA transfer from neighboring cells

(Desrochers et al., 2016). Of note, the transmission of *tdTomato* mRNA and protein from

$Pax7^{+/CreERT2}:R26R^{+/tdT}$ mice used in this study has been recently shown via exosome, opening

up the possibility of transmission of other mRNA from MuSC to ECs (Murach et al., 2020). The

later is an interesting phenomenon form a developmental point of view. MuSCs and ECs arise

from a bipotent progenitor originated from somites during early development (Kardon et al.,

2002; Hutcheson, 2009; Lagha, 2009; Mayeuf-Louchart, 2014; Mayeuf-Louchart, 2016).

Therefore, it is possible that there is a permissive chromatin state that allows for expression of

reciprocal genes in the two populations. Along the lines of these observations, FLK1(+) or VE-

cadherin(+) cells can contribute to myogenic cells *in vitro* and after cell transplantation (Tamaki

et al., 2002; Le Grand et al., 2004; Zheng et al., 2007; Huang et al., 2014;), and during

development (Drummond and Hatley, 2018; Mayeuf-Louchart et al., 2014; Motoike et al., 2003).

Important notion is that the PDGFRα(-)FLK1(+) population exhibited restricted potential to

differentiate into the MuSCs in injured muscle (Sakurai et al., 2008). Interestingly, in the

zebrafish, exogenous expression of *Etv2* in the fast muscle can lead to transdifferentiation of

muscle fibers into functional vessels so there is evidence of cell fate flexibility (Veldman et al.,

2013). The potential of EC transdifferentiation was also examined by ETV2 overexpression in

five human cell types, skeletal muscle cells, adipose-derived mesenchymal stem cells, umbilical

cord-derived mesenchymal stem cells, embryonic lung fibroblast cells and skin fibroblast cells.

Among them, human skeletal muscle cells showed the highest amenability for this EC induction

following infection with ETV2 lentivirus vector (Yan et al., 2019). Conversely, *Etv2*-deficient

vascular progenitors can differentiate  into skeletal muscle cells (Chestnut et al., 2020). It would

be interesting to see whether other EC gene signatures also have functional consequences in the

MuSC or muscle at large.

We decided to focus on function of *Flt1* among several EC genes expressed in MuSCs

for further investigations on MuSC biology. Our pharmacological and genetic analyses

demonstrate that MuSC-derived VEGFA has a drastic effect on cell survival in the via its

receptor FLT1 by signaling through AKT1. While VEGFA binds to both FLT1 and FLK1,

VEGFB and PGF only bind to FLT1. This creates a scenario where PGF and VEGFB binding

can sequester FLT1, increasing free VEGFA availability for VEGFA-FLK1 binding which is the

major VEGF signaling pathway for many cell types (Vempati et al., 2014). While PGF is not

normally expressed in adult tissues, VEGFB is expressed in the MuSCs and muscle fiber (data

not shown). Importantly, the VEGFB-FLT1 axis has also been shown to inhibit apoptosis in

retina and brain cells in mouse models of ocular neurodegeneration and stroke (Li et al., 2008).

While our results cannot rule out the involvement of VEGFB in protection of MuSC apoptosis,

we provide evidence from both pharmacological and genetic data to indicate that VEGFA is

involved.

Despite drastic effect of VEGFA-FLT1 on apoptosis *in vitro*, the long-term consequences

of *in vivo* deletion of *Flt1* in the MuSC compartment were modest compared with deletion of

*Vegfa* in the MuSCs unless crossing with *mdx* mice. Although *Vegfa* is required for both MuSC

survival and recruitment of vascular niche (Verma et al., 2018), in the steady state, the MuSC

turnover may be low enough that the apoptotic stress burden is low. VEGFA improves cell

survival during the proliferative stage following injury, however this transient improvement in

survival has only a modest impact on the final regenerative process as shown in

*mdx:VEGFA$^{+/Hyper}$* mice.

VEGFA and FLT1 targeted therapies are being explored as both pro- and anti-angiogenic

therapies for several indications including retinal degeneration, cancer, pre-eclampsia and

neuromuscular diseases (Bae et al., 2005; Keifer et al., 2014; Mac Gabhann et al., 2011; Verma

et al., 2010; Verma et al., 2019; Bosco et al., 2021; Xin et al., 2021). As these therapies mature,

it will be important to ascertain the MuSC-specific effects of VEGFA and FLT1.

## Materials and Methods

### Mice

$Flt1^{LoxP/LoxP}$ were obtained from Gua-Hua Fong (Ho et al., 2012). $B6.Cg\text{-}Pax7^{tm1(cre/ERT2)Gaka/J}$ ($Pax7^{+/CreERT2}$; JAX stock# 017763; Murphy et al., 2011), $B6.Cg\text{-}Gt(ROSA)^{26Sortm9(CAG\text{-}tdTomato)Hze/J}$ ($Ai9$; JAX stock # 007909; Madisen et al., 2010), $VEGFA^{+/Hyper}$ ($Vegfatm1.1Nagy/J$; JAX stock# 027314; Miquerol et al., 1999) and $B6Ros.Cg\text{-}Dmd^{mdx\text{-}5cv}/J$ ($mdx^{5cv}$; JAX stock #002379; Chapman et al., 1989) were obtained from Jackson Laboratory. $Kdr^{tm2.1Jrt/J}$ ($Flk1^{+/GFP}$) were obtained from Masatsugu Ema (Ema et al., 2006). $B6.Cg\text{-}Pax7^{tm1(cre/ERT2)Gaka/J}$ ($Pax7^{+/CreERT2}$) mice were crossed with the $B6.Cg\text{-}Gt(ROSA)^{26Sortm9(CAG\text{-}tdTomato)Hze/J}$ ($Ai9$) to yield the $Pax7^{+/CreERT2}\text{:}R26R^{tdT}$($Pax7^{tdT}$) mice. $Pax7^{tdT}$ mice were bred with the $VEGFA^{+/Hyper}$ and $Flk1^{+/GFP}$ to obtain $Pax7^{+/tdT}\text{:}VEGFA^{+/Hyper}$ and $Pax7^{+/tdT}\text{:}Flk1^{+/GFP}$ mice. $VEGFA^{LoxP/LoxP}$ mice obtained from Napoleone Ferrara (Gerber et al., 1999) were crossed with $Pax7^{+/CreERT2}$ to yield the $Pax7^{+/CreERT}\text{:}VEGFA^{LoxP/LoxP}$ mice. $Flt1^{LoxP/LoxP}$ mice obtained from Guo-Hua Fong (Ho et al., 2012) were crossed with $Pax7^{+/CreERT2}$ to yield the $Pax7^{+/CreERT}\text{:}Flt1^{LoxP/LoxP}$ mice. Colonies for all the mice were established in the laboratory. Cre recombination was induced using tamoxifen (Sigma-Aldrich, T5648) dosed as 75 mg/kg body weight x 3 times over one week at 3-6 weeks of age. Mice carrying the wild-type $CreERT2$ allele were used for control experiments. TA muscle regeneration was induced by intramuscular injection of 20 µl of 1% $BaCl_2$ (Sigma-Aldrich, 342920) or 20 µl of 10 µM Cardiotoxin (CTX) (Sigma-Aldrich, V9125). Mice used for this study is summeried in Table S2.

Genotyping to detect the transgenic and mutant alleles was performed by PCR using the primers described on the web site of Jackson Laboratory shown in Table S3. All primers were synthesized as custom DNA oligos from Integrated DNA technologies (IDT). Genotyping to

detect the mutated allele of *mdx$^{5cv}$* was performed by PCR using the primers (0981 and 0982)

shown in Table S1. The PCR product DNA was digested with *Dra*III restriction enzyme (New

England Biolabs, R3510S). Wild-type allele generated 180 bp and mutant allele generated 50

and 130 bp bands.

The animals were housed in an SPF environment and were monitored by the Research

Animal Resources (RAR) of the University of Minnesota. All protocols were approved by the

Institutional Animal Care and Usage Committee (IACUC) of the University of Minnesota and

complied with the NIH guidelines for the use of animals in research.

**Cell isolation by FACS**

*Pax7$^{tdT}$:Flk1$^{GFP}$* mice were utilized for FACS-mediated MuSC and EC isolation as previously

described (Asakura et al, 2002; Verma, 2018). We performed extensive validation of the

fluorescent reporter mice as previously described (Figure S1A-C; Verma, 2018). Briefly,

quiescnet MuSCs and ECs were isolated from the hind limb skeletal muscle of 1–2-mo-old

*Pax7$^{tdT}$:Flk1$^{GFP}$* mice after digestion with collagenase type II. FACS was performed on an FACS

sorter (BD FACSAria) and data were analyzed using FlowJo (BD Biosciences). Sorting gates,

tdTomato(+) for MuSCs and GFP(+) for ECs, were strictly defined based on control cells

isolated from wild-type mice and the forward scatter and side scatter gating. Sorted cells were

immediately characterized by immunostaining on slide glasses, utilzed for RNA preparation or

cultured on collagen-coated plates in the myoblast growth medium as below to obtain MuSC-

derived myoblasts and ECs. FACS analysis was performed as previously described (Turaç et al.,

2013). Cells were either trypsinized (cultured cells) or a single cell suspension was obtained

following enzymatic digestion as whole muscle-derived cells (Asakura, 2002). Cells were then

washed with FACS buffer (2% BSA and 1 mM EDTA in PBS) followed by live/dead staining

using ZombieNIR (Biolegends, 423105). Cells were washed, then immunostained for cell

surface markers. Blocking cells was performed with 1% BSA/PBS, and cells were incubated in

fluorescently-conjugated antibody. FACS was performed on a Fortessa X-20 (BD Biosciences)

with a 355 nm, 405 nm, 488 nm, 561 nm, and 640 nm lasers.

**Cell culture**

MuSC-derived myoblast isolation from adult mice was performed as previously described

(Asakura et al, 2001; Motohashi et al., 2014). Briefly, after collagenase type II (Worthington,

CLS-2) treatment, dissociated cells from mouse hindlimb muscle were incubated with anti-

CD31-PE (eBiosciences, 12-0311), anti-CD45-PE (eBiosciences, 30-F11), anti-Sca1-PE

(eBiosciences, Dec-81) and anti-Integrin α7 (MBL International, ABIN487462), followed by

anti-PE microbeads (Miltenyi Biotec, 130-048-801), and then performed LD column (Miltenyi

Biotec, 130-042-901) separation. Negative cell populations will be incubated with anti-Mouse

IgG beads (Miltenyi Biotec, 130-048-402), and then MS coulumn (Miltenyi Biotec, 130-042-

201) separation was performed to isolate Integrin α7(+) MuSCs. MuSC-derived myoblasts were

maintained in culture on collagen coated plates in myoblast medium containing 20% FBS and 20

ng/ml bFGF (Invitrogen, PHG0263) in HAM's-F10 medium.  Cell cultures were maintained in a

humidified incubator at 37°C with 5% $CO_2$ and 5% $O_2$. 4-hydroxy tamoxifen (4-OHT, Sigma-

Aldrich, H6278) treatment (1 µM in EtOH) was used to induce *Flt1* deletion in myoblasts

isolated from *Flt1[LoxP/LoxP]:Pax7[CreERT2]* mice. For cell survival assay, 1 x $10^5$ cells were allowed to

adhere for one day and starved overnight in 0.1% FBS in HAM's F10 medium. Then, cells were

exposed to 1 µM EdU along with or withpout 2-100 ng/ml recombinant VEGFA (R&D Systems,

493-MV) for 8 hours before being fixed and stained by the Click-iT EdU Alexa Fluor 488

Imaging Kit (Thermo Scientific, C10337). For induction of apoptosis in myoblasts, (1-2 x $10^5$)

cells were allowed to adhere to the plates for 16 hours. Thapsigargin-mediated apoptosis was

induced by 1 µM of thapsigargin (Sigma-Aldrich, T9033) dissolved in EtOH with or without

VEGFA, 100 ng/ml recombinant FLT1-FC (R&D Systems, 7756-FL), 1 µg/ml anti-FLT1

monoclonal antibody (Angio-Proteomie, MAB7072), inhibitors of FLK1, 3 µM ZM306416

(R&D Systems, 2499/1) and 10 µM of SU5402 (R&D Systems, 3300/1) and an inhibitor of

NRP1, 30 µM of EG00229 (R&D Systems, 6986/10) for 24 hours. UV light-mediated apoptosis

was induced by exposing the cells to UV light in cell culture hood for 45 seconds without

medium. After UV exposure, cell survival was assessed 24 hours following culture in 0.1% FBS

in HAM's F10 medium with or without VEGFA using the Crystal violet Assay Kit (Abcam,

ab232855) and quantated the Crystal violet dye after solubilization by absorbance at 570 nm. To

induce differentiation of myoblasts, the myoblast medium was replaced with differentiation

medium that contained DMEM supplemented with 5% horse serum with or withour VEGFA or

bFGF for 1 or 3 days followed by anti-sarcomeric myosin heavy chain antibody (Developmental

Study Hybridoma Bank, MF-20).

**AKT induction**

The lentiviral pCCL-E4ORF1 and pCCL-myrAkt1 constructs were a kind gift from Dr. Jason

Butler (Kobayashi et al., 2010). 293FT cells (Thermo Fisher Scientific, R70007) were seeded in

DMEM with 10% FBS and transfected with the lentivirus vectors along with pCMV-VSV-G

(Addgene, 8454), pRSV-Rev (Addgene, 12253), and pMDLg/pRRE (Addgene, 12251) using

PolyJet transfection reagent (Signagen Laboratories, SL100688). The culture supernatant of the

transfected 293FT cells was added to MuSC-derived myoblast culture with 0.8 μg/ml polybrene

(MilliporeSigma, H9268). pAKT1(+) cells were stainined with anti-pAKT antibody (Cell

Signaling, 4060).

**Apoptosis assay**

Apoptosis was measured using measured using Annexin V-Biotin Apoptosis Detection Kit

(eBioscience™, BMS500BT-100) as per the manufacture's instruction. Streptavidin-conjugated

AlexaFluro-488 was used for detection. Propidium Iodide (PI) was used in all assays except

when Pax7tdT(+) cells were utilized or when ZombieNIR (Biolegends, 423105) was used. FACS

was performed on a Fortessa X-20 (BD Biosciences) equipped with a 355 nm, 405 nm, 488 nm,

561 nm, and 640 nm lasers.

**Immunostaining of cells**

Immunostaining for PECAM1, VE-Cadherin, VEGFA, VEGFRs was performed on collagen

coated coverslips. Other immunostaining was performed on 30 mm tissue culture plates. Cells

were fixed with 2% PFA for 5 minutes and immunostained as previously described (Verma et al.,

2010). For membrane receptor staining, cells were permeabilized with 0.01% saponin

(ThermoFisher Scientific, ICN10285525) which was kept in the staining solution until the

primary antibodies were washed off. At which time, 0.01% Triton-X was added to all the buffers.

The antibodies used for this study are listed in Table S4.

**Single muscle fiber isolation and staining**

Extensor digitorum longus (EDL) muscle was dissected and digested with 0.2% collagenase type

I (Sigma-Aldrich, C0130) for single muscle fiber isolation as previously described (Verma et al.,

2010). Single muscle fibers were fixed with 2% PFA/PBS, permeabilized with 0.2% Triton-

X100 and counterstained with DAPI. Anti-Pax7 antibody(+) or tdTomato(+) MuSCs per single

muscle fiber were counted manually.

**RNAscope**

RNAscope for *Flt1* transcripts was performed as previously described (Kann and Krauss, 2019)

on single muscle fibers from *Pax7^{tdT}* mice using the RNAscope Probe - Mm-Flt1 (C1) (ACDBio,

415541). Briefly, isolated EDL fibers are fixed in 4% PFA, washed with PBS, and dehydrated in

100% methanol. Subsequently, fibers are rehydrated in a stepwise gradient of decreasing

methanol concentrations in PBS/0.1% Tween-20. Fibers are treated with a proteinase for 10

minutes, followed by hybridization, amplification, and fluorophore conjugation steps.

**Histology**

The mouse tibialis anterior (TA) muscle was used for all histological analysis. Tissues were

frozen fresh using $LiN_2$ chilled isopentane and stored at -80ºC. Eight μm thick transverse

cryosections were used for all histological analysis. Hematoxylin & Eosin (HE) staining were

performed as previously described (Verma et al., 2010). Sirius red (Direct Red 80, Sigma-

Aldrich, 365548) staining was performed for muscle sections for fibrosis as previously described

(Shimizu-Motohashi et al., 2015). Muscle sections were stained in Oil Red O solution (Sigma-

Aldrich, O1391-250ML) as previously described (Wang et al., 2017). Microscopic images were

captured by a DP-1 digital camera attached to BX51 fluorescence microscope with 10x or 40×

UPlanFLN objectives with cellSens Entry 1.11 (all from Olympus). Photoshop (Adobe) and Fiji

(NIH) were used for image processing and manually enumerating the fiber diameter (Schindelin

et al., 2012).

**Grip strength test**

Forelimb grip strength test was performed following a previously published procedure (Aartsma-

Rus and van Putten, 2014). Briefly, mice were gently pulled by the tail after fore limb-grasping a

metal bar attached to a force transducer (Grip Strength Meter, Columbus Instruments, 1027CSM-

D52). Grip strength tests were performed by the same blinded examiner. Five consecutive grip

strength tests were recorded, and then mice were returned to the cage for a resting period of 20

minutes. Then, three series of pulls were performed each followed by 20 min resting period. The

average of the three highest values out of the 15 values collected was normalized to the body

weight for comparison.

**Muscle perfusion**

RBC flux was evaluated using the moorLabTM laser Doppler flow meter as previously described

(Verma, 2010) with the MP7a probe that allows for collecting light from a deeper tissue level

than standard probes according to the manufacturer's instructions (Moor Instruments). The fur

from the right hind leg was removed using a chemical depilatory. Readings were taken using the

probe from at least 10 different spots on the TA muscle. The AU was determined as the average

AU value during a plateau phase of each measurement.

**RNA and genomic DNA isolation and qPCR**

Cultured cells were washed with ice cold PBS and lysed on the place with Trizol$^{TM}$. RNA was

isolated using the DirectZol$^{TM}$ RNA Microprep Kit (Zymo Research, R2062) with on-column

DNase digestion followed by cDNA synthesis using the Transcriptor First Strand cDNA

synthesis kit (Roche Molecular Diagnostics, 04379012001) using random primers. Genomic

DNA for genotyping was isolated from mouse tail snips with lysis buffer containing Proteinase

K (Sigma-Aldrich, P2308).  qPCR was performed using GoTaq qPCR Master Mix (Promega,

A6002). The input RNA amount was normalized across all samples and *18S rRNA* or *HtatsF1*

was used for normalization of qPCR across samples. Primer sequences are listed in Table S3. All

primers were synthesized as custom DNA oligos from Integrated DNA technologies (IDT).

**Single cell RNAsequencing and analysis**

Cells for single cell RNAseq were obtained from hind limb muscles of 2-3 moth-old

*Pax7$^{tdT}$:Flk1$^{GFP}$* mice following enzymatic digestion as previously described (Liu et al., 2015).

Dead cells were excluded from the analysis using ZombieNIR (Biolegends, 423105).

TdTomato(+) and GFP(+) cells were sorted individually and then 20% of GFP(+) cells were

spiked into 80% tdTomato(+). We loaded ~5,000 cells into 1 channel of the Chromium system

for each of these samples and prepared libraries according to the manufacturer's protocol using

version 2.0 chemistry (10x Genomics). Following capture and lysis, we synthesized cDNA and

amplified for 12 cycles as per the manufacturer's protocol (10x Genomics). The amplified cDNA

was used to construct Illumina sequencing libraries that were each sequenced with ~300K

read/cell on one lane of an Illumina HiSeq 2500 machine. We used Cell Ranger 3.1 (10X

Genomics) to process raw sequencing data. For A custom genome was constructed to include

*eGFP-SV40*, *tdTomato-WPRE-BGHPolyA* and *Pax7-IRES-CreERT2* transgenes. Detailed step-

by-step instructions can be found at https://github.com/verma014/10XCustomRef. We carried

out analyses of the filtered data using Seurat suite version 3.0 (Stuart et al., 2019) in R studio

(RStudio Team, 2020). For cell imputation, we utilized ALRA through the Seurat wrapper with

default settings(Linderman et al., 2018). Additional scRNAseq datasets were obtained from GEO

and analyzed using the same method as listed above. A myogenic score was calculated based on

the expression of *Myog*, *Pax7*, *Myod1* and *Myf5*. Step-by-step instructions for the analysis can be

found on https://github.com/verma014/10XCustomRef.

**Background Subtraction**

10x Genomics scRNAseq platform uses many more droplets than cells and so following a run,

there are many droplets that do not have cells. These droplets still get sequenced with some of

the RNA that is in the solution. This floating RNA can be used to estimate the "background" in

each droplet. A better description of this can be found by the developers of 'SoupX' (Young and

Behjati, 2020). Since *Cdh5* expression has previously been verified in MuSCs using RNAscope,

we were able to use it as a positive control to remove the background or "soup" from our data. If

*Cdh5* is absent from MuSC, we know that the background subtraction was too aggressive and

that subtracting the Soup is not reliable in our case. In addition, we know certain genes that are

considered to be specific for MuSCs, muscle ECs or muscle fibers based on the bulk RNAseq

(Verma et al., 2018). The top 5 genes that are specific to these population (and also detected by

10x) were selected and used to show the background in our data set was 14.40% and 13.89% for

the D0 and D3 dataset, respectively. The step-by-step instructions can be found on

https://github.com/verma014/10XCustomRef.

**Bulk RNAseq and Microarray Analysis**

FASTQ files were downloaded from SRA using SRA-toolkit. Sequences were trimmed using trimmomatic to remove adapter contamination and low-quality reads. Trimmed sequences were mapped to mouse mm10 using Hisat2 (Pertea et al., 2016). Transcript assembly was performed using StringTie (Pertea et al., 2016). Cell type specificity was determined as previously described (Verma et al., 2018). Microarray analysis was performed using the Affymetrix Transcriptome Analysis Console (TAC). Samples in each experiment were RNA normalized and the expression was acquired using the Affeymetrix Expression analysis console with gene level expression. Heatmaps were generated in Prism 9 (Graphpad, La Jolla, CA). The code for generating each graph is listed in the following table, along with the link to the data in tabular format.

**Quantification and Statistical Analysis**

Statistical analysis was performed using Prism 9 (Graphpad, La Jolla, CA) or RStudio (RStudio Team, 2020). For comparison between two groups, an unpaired T-test was used. For comparison between multiple groups, a one-way ANOVA was used with multiple comparisons to the control. Distributions were compared using a chi-squared test. Graphing of the data was performed using Prism 9. Vector diagrams were modified using Graphic (Autodesk). All values are means $\pm$SEM unless noted otherwise. * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$.

**Data and Software Availability**

All the data was obtained from NCBI GEO. Microarrays of mouse MuSCs were obtained from GSE3483 (Fukada et al., 2007). scRNAseq of MuSCs and muscle ECs was performed in this

study (GSE129057). scRNAseq of whole muscle was obtained from GSE143437 (De Micheli et

al., 2020). Bulk RNAseq of MuSCs, ECs and single muscle fibers was obtained from

GSE108739 (Verma et al., 2018) and GSE64379 (Ryall et al., 2015). Bulk RNAseq of TU-

tagged RNA of MuSCs was obtianed from GSE97399 (van Velthoven et al., 2017). Bulk

RNAseq of fixed and unfixed MuSCs was obtained from GSE113631 (Yue et al., 2020).

Exercise, ALS, DMD, BMD, FSHD GSE3307, Early DMD GSE465, *mdx* GSE466, GRMD

GSE69040, Satellite cells GSE15155. All arrays were normalized to their respective controls. All

arrays and RNAseq data are listed in Supplemental Table 1 (Table S1).

**Acknowledgements**

We thank Minnesota Supercomputing Institute, University of Minnesota Imaging Center,

University of Minnesota FACS Facility, and University of Minnesota Genomics Center for

providing data for this paper. We also thank Jake Trask for critical reading of this paper. We

thank Drs. Yosuke Mukouyama (National Institute of Health), Napoleone Ferrara (Genentech),

Guo-Hua Fong (University of Connecticut) and Masatsugu Ema (Siga University of Medical

Science) for providing *VEGFA^{LoxP/LoxP}*, *Flt1^{LoxP/LoxP}* and *Flk1-GFP* mice, respectively. This work

was supported by NIHT32-GM008244 and NIHF30AR066454 to MV, NIAMS grant AR070231

to RSK, a fellowship of the Training Program in Stem Cell Research from the New York State

Department of Health to A.P.K. (NYSTEM-C32561GG) and NIHR01AR062142,

NIHR21AR070319, MDA Research Grant, and Regenerative Medicine Minnesota (RMM) Grant

to AA.

**Competing interests**

The authors declare no competing interests.

**Reference**

Aartsma-Rus A, van Putten M. 2014. Assessing Functional Performance in the Mdx Mouse

Model. *J Vis Exp*. 51303. doi:10.3791/51303

Arsic N, Zacchigna S, Zentilin L, Ramirez-Correa G, Pattarini L, Salvi A, Sinagra G, Giacca M.

2004. Vascular endothelial growth factor stimulates skeletal muscle regeneration in vivo.

*Mol Ther* **10**:844–854. doi: 10.1016/j.ymthe.2004.08.007

Asakura A, Komaki M, Rudnicki M. 2001. Muscle satellite cells are multipotential stem cells

that exhibit myogenic, osteogenic, and adipogenic differentiation. *Differentiation* **68**:245–

253. doi: 10.1046/j.1432-0436.2001.680412.x.

Asakura A, Seale P, Girgis-Gabardo A, Rudnicki MA. 2002. Myogenic specification of side

population cells in skeletal muscle. *J Cell Biol* **159**:123–134. doi: 10.1083/jcb.200202092.

Bae DG, Kim TD, Li G, Yoon WH, Chae CB. 2005. Anti-flt1 peptide, a vascular endothelial

growth factor receptor 1-specific hexapeptide, inhibits tumor growth and metastasis. *Clin*

*Cancer Res* **11**:2651–2661. doi: 10.1158/1078-0432.CCR-04-1564

Bakay M, Wang Z, Melcon G, Schiltz L, Xuan J, Zhao P, Sartorelli V, Seo J, Pegoraro E,

Angelini C, Shneiderman B, Escolar D, Chen Y-W, Winokur ST, Pachman LM, Fan C,

Mandler R, Nevo Y, Gordon E, Zhu Y, Dong Y, Wang Y, Hoffman EP. 2006. Nuclear

envelope dystrophies show a transcriptional fingerprint suggesting disruption of Rb-MyoD

pathways in muscle regeneration. *Brain* **129**:996–1013. doi:10.1093/brain/awl023

Bosco J, Zhou Z, Gabriëls S, Verma M, Liu N, Miller BK, Gu S, Lundberg DM, Huang Y,

Brown E, Josiah S, Meiyappan M, Traylor MJ, Chen N, Asakura A, De Jonge N, Blanchetot

C, de Haard H, Duffy HS, Keefe D. 2021. VEGFR-1/Flt-1 inhibition increases

angiogenesis and improves muscle function in a mouse model of Duchenne muscular

dystrophy. *Mol Ther Methods Clin Dev* **23**:21:369-381. doi: 10.1016/j.omtm.2021.03.013

Bouchentouf M, Benabdallah BF, Bigey P, Yau TM, Scherman D, Tremblay JP. 2008. Vascular

endothelial growth factor reduced hypoxia-induced death of human myoblasts and

improved their engraftment in mouse muscles. *Gene Ther* **15**:404–414.

doi: 10.1038/sj.gt.3303059

Bryan BA, Walshe TE, Mitchell DC, Havumaki JS, Saint-Geniez M, Maharaj AS, Maldonado

AE, D'Amore PA, Bryan BA, Walshe TE, Mitchell DC, Havumaki JS, Saint-Geniez M,

Maharaj AS, Maldonado AE, D'Amore PA. 2008. Coordinated vascular endothelial growth

factor expression and signaling during skeletal myogenic differentiation. *Mol Biol Cell*

**19**:994–1006. doi:10.1091/mbc.E07-09-0856

Chapman VM, Miller DR, Armstrong D, Caskey CT. 1989. Recovery of induced mutations for

X chromosome-linked muscular dystrophy in mice. *Proc Natl Acad Sci U S A* **86**:1292-6.

doi: 10.1073/pnas.86.4.1292

Charville GW, Cheung TH, Yoo B, Santos PJ, Lee GK, Shrager JB, Rando TA. 2015. Ex Vivo

Expansion and In Vivo Self-Renewal of Human Muscle Stem Cells. *Stem cell Reports*

**5**:621–32. doi:10.1016/j.stemcr.2015.08.004

Chen YW, Zhao P, Borup R, Hoffman EP. 2000. Expression profiling in the muscular

dystrophies: identification of novel aspects of molecular pathophysiology. *J Cell Biol*

**151**:1321–36. doi:10.1083/jcb.151.6.1321

Chestnut B, Casie Chetty S, Koenig AL, Sumanas S. 2020. Single-cell transcriptomic analysis

identifies the conversion of zebrafish Etv2-deficient vascular progenitors into skeletal

muscle. *Nat Commun* **11**:2796. doi:10.1038/s41467-020-16515-y

Christov C, Chretien F, Abou-Khalil R, Bassez G, Vallet G, Authier FJ, Bassaglia Y, Shinin V,

Tajbakhsh S, Chazaud B, Gherardi RK. 2007. Muscle satellite cells and endothelial cells:

close    neighbors    and    privileged    partners.    *Mol    Biol    Cell*    **18**:1397–1409.

doi: 10.1091/mbc.e06-08-0693

De Angelis L, Berghella L, Coletta M, Lattanzi L, Zanchi M, Cusella-De Angelis MG, Ponzetto

C, Cossu G. 1999. Skeletal myogenic progenitors originating from embryonic dorsal aorta

coexpress endothelial and myogenic markers and contribute to postnatal muscle growth and

regeneration. *J Cell Biol* **147**:869–877. doi:10.1083/jcb.147.4.869

De Micheli AJ, Laurilliard EJ, Heinke CL, Ravichandran H, Fraczek P, Soueid-Baumgarten S,

De Vlaminck I, Elemento O, Cosgrove BD. 2020. Single-Cell Analysis of the Muscle Stem

Cell Hierarchy Identifies Heterotypic Communication Signals Involved in Skeletal Muscle

Regeneration. *Cell Rep* **30**:3583-3595.e5. doi:10.1016/j.celrep.2020.02.067

Delavar H, Nogueira L, Wagner PD, Hogan MC, Metzger D, Breen EC. 2014. Skeletal myofiber

VEGF is essential for the exercise training response in adult mice. *AJP Regul Integr Comp*

*Physiol* **306**:R586–R595. doi:10.1152/ajpregu.00522.2013

Desrochers LM, Antonyak MA, Cerione RA. 2016. Extracellular Vesicles: Satellites of

Information Transfer in Cancer and Stem Cell Biology. *Dev Cell* **37**:301-309. doi:

10.1016/j.devcel.2016.04.019.

Domigan CK, Warren CM, Antanesian V, Happel K, Ziyad S, Lee S, Krall A, Duan L, Torres-

Collado AX, Castellani LW, Elashoff D, Christofk HR, van der Bliek AM, Potente M,

Iruela-Arispe ML. 2015. Autocrine VEGF maintains endothelial survival through regulation

of metabolism and autophagy. *J Cell Sci* **128**:2236–48. doi:10.1242/jcs.163774

Drummond CJ, Hatley ME. 2018. A Case of mistaken identity: Rhabdomyosarcoma

development from endothelial progenitor cells. *Mol Cell Oncol* **5**:e1448246.

doi:10.1080/23723556.2018.1448246

Eichmann A, Marcelle C, Bréant C, Le Douarin NM. 1993. Two molecules related to the VEGF

receptor are expressed in early endothelial cells during avian embryonic development. *Mech*

*Dev* **42**:33–48. doi:10.1016/0925-4773(93)90096-g

Ema M, Takahashi S, Rossant J. 2006. Deletion of the selection cassette, but not cis-acting

elements, in targeted Flk1-lacZ allele reveals Flk1 expression in multipotent mesodermal

progenitors. *Blood* **107**:111–7. doi:10.1182/blood-2005-05-1970

Esner M, Meilhac SM, Relaix F, Nicolas J-F, Cossu G, Buckingham ME. 2006. Smooth muscle

of the dorsal aorta shares a common clonal origin with skeletal muscle of the myotome.

*Development* **133**:737–49. doi:10.1242/dev.02226

Fukada S, Uezumi A, Ikemoto M, Masuda S, Segawa M, Tanimura N, Yamamoto H, Miyagoe-

Suzuki Y, Takeda S. 2007. Molecular signature of quiescent satellite cells in adult skeletal

muscle. *Stem Cells* **25**:2448–59. doi:10.1634/stemcells.2007-0019

Gay L, Miller MR, Ventura PB, Devasthali V, Vue Z, Thompson HL, Temple S, Zong H, Cleary

MD, Stankunas K, Doe CQ. 2013. Mouse TU tagging: a chemical/genetic intersectional

method for purifying cell type-specific nascent RNA. *Genes Dev* **27**:98–115.

doi:10.1101/gad.205278.112

Gerber HP, McMurtrey A, Kowalski J, Yan M, Keyt B a, Dixit V, Ferrara N. 1998. Vascular

endothelial    growth    factor    regulates    endothelial    cell    survival    through    the

phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-

1/KDR activation. *J Biol Chem* **273**:30336–43. doi : 10.1074/jbc.273.46.30336

Germani A, Di Carlo A, Mangoni A, Straino S, Giacinti C, Turrini P, Biglioli P, Capogrossi MC.

2003. Vascular endothelial growth factor modulates skeletal myoblast function. *Am J*

*Pathol* **163**:1417–1428. doi: 10.1016/S0002-9440(10)63499-2

Giordani L, He GJ, Negroni E, Sakai H, Law JYC, Siu MM, Wan R, Corneau A, Tajbakhsh S,

Cheung TH, Le Grand F. 2018. High-Dimensional Single-Cell Cartography Reveals Novel

Skeletal Muscle-Resident Cell Populations. *Mol Cell* **74**:609-621.e6. doi:

10.1016/j.molcel.2019.02.026.

Goel AJ, Rieder MK, Arnold HH, Radice GL, Krauss RS. 2017. Niche Cadherins Control the

Quiescence-to-Activation Transition in Muscle Stem Cells. *Cell Rep* **21**:2236–2250.

doi:10.1016/j.celrep.2017.10.102

Hardy D, Besnard A, Latil M, Jouvion G, Briand D, Thépenier C, Pascal Q, Guguin A, Gayraud-

Morel B, Cavaillon J-M, Tajbakhsh S, Rocheteau P, Chrétien F. 2016. Comparative Study

of Injury Models for Studying Muscle Regeneration in Mice. *PLoS One* **11**:e0147198.

doi:10.1371/journal.pone.0147198

Hirai H, Verma M, Watanabe S, Tastad C, Asakura Y, Asakura A. 2010. MyoD regulates

apoptosis of myoblasts through microRNA-mediated down-regulation of Pax3. *J Cell Biol*

**191**:347–65. doi:10.1083/jcb.201006025

Ho VC, Duan L-J, Cronin C, Liang BT, Fong G-H. 2012. Elevated vascular endothelial growth

factor receptor-2 abundance contributes to increased angiogenesis in vascular endothelial

growth factor receptor-1-deficient mice. *Circulation* **126**:741–52.

doi:10.1161/CIRCULATIONAHA.112.091603

Huang P, Schulz TJ, Beauvais A, Tseng YH, Gussoni E. 2014.

Intramuscular adipogenesis is inhibited by myo-endothelial progenitors with functioning

Bmpr1a signalling. *Nat Commun* **5**:4063. doi: 10.1038/ncomms5063.

Hutcheson DA, Kardon G. 2009. Genetic manipulations reveal dynamic cell and gene functions:

Cre-ating a new view of myogenesis. *Cell Cycle* **8**:3675–8. doi:10.4161/cc.8.22.9992

Kann AP, Krauss RS. 2019. Multiplexed RNAscope and immunofluorescence on whole-mount

skeletal myofibers and their associated stem cells. *Development* **146**:dev179259

doi:10.1242/dev.179259

Kardon G, Campbell JK, Tabin CJ. 2002. Local extrinsic signals determine muscle and

endothelial cell fate and patterning in the vertebrate limb. *Dev Cell* **3**:533–45.

doi:10.1016/s1534-5807(02)00291-5

Keifer OP, O'Connor DM, Boulis NM. 2014. Gene and protein therapies utilizing VEGF for

ALS. *Pharmacol Ther* **141**:261–71. doi:10.1016/j.pharmthera.2013.10.009

Kobayashi H, Butler JM, O'Donnell R, Kobayashi M, Ding B-S, Bonner B, Chiu VK, Nolan DJ,

Shido K, Benjamin L, Rafii S. 2010. Angiocrine factors from Akt-activated endothelial cells

balance self-renewal and differentiation of haematopoietic stem cells. *Nat Cell Biol*

**12**:1046–56. doi:10.1038/ncb2108

Kodippili K, Thorne PK, Laughlin MH, Duan D. 2021. Dystrophin deficiency impairs vascular

structure and function in the canine model of Duchenne muscular dystrophy. *J Pathol*

**254**:589-605. doi:10.1002/path.5704

Lagha M, Brunelli S, Messina G, Cumano A, Kume T, Relaix F, Buckingham ME. 2009.

Pax3:Foxc2 reciprocal repression in the somite modulates muscular versus vascular cell fate

choice in multipotent progenitors. *Dev Cell* **17**:892–9. doi:10.1016/j.devcel.2009.10.021

Latroche C, Matot B, Martins-Bach A, Briand D, Chazaud B, Wary C, Carlier PG, Chrétien F,

Jouvion G. 2015. Structural and Functional Alterations of Skeletal Muscle Microvasculature

in Dystrophin-Deficient mdx Mice. *Am J Pathol* **185**:2482-94.

doi:10.1016/j.ajpath.2015.05.009

Latroche C, Weiss-Gayet M, Muller L, Gitiaux C, Leblanc P, Liot S, Ben-Larbi S, Abou-Khalil

R, Verger N, Bardot P, Magnan M, Chrétien F, Mounier R, Germain S, Chazaud B. 2017.

Coupling between Myogenesis and Angiogenesis during Skeletal Muscle Regeneration Is

Stimulated by Restorative Macrophages. *Stem Cell Reports* **9**:2018-2033.

doi:10.1016/j.stemcr.2017.10.027

Le Grand F, Auda-Boucher G, Levitsky D, Rouaud T, Fontaine-Pérus J, Gardahaut M-F. 2004.

Endothelial cells within embryonic skeletal muscles: a potential source of myogenic

progenitors. *Exp Cell Res* **301**:232–41. doi:10.1016/j.yexcr.2004.07.028

Lee S, Chen TT, Barber CL, Jordan MC, Murdock J, Desai S, Ferrara N, Nagy A, Roos KP,

Iruela-Arispe ML. 2007. Autocrine VEGF signaling is required for vascular homeostasis.

*Cell* **130**:691–703. doi:10.1016/j.cell.2007.06.054

Li Y, Zhang F, Nagai N, Tang Z, Zhang S, Scotney P, Lennartsson J, Zhu C, Qu Y, Fang C, Hua

J, Matsuo O, Fong GH, Ding H, Cao Y, Becker KG, Nash A, Heldin CH, Li X. 2008.

VEGF-B inhibits apoptosis via VEGFR-1-mediated suppression of the expression of BH3-

only protein genes in mice and rats. *J Clin Invest* **118**:913–923. doi:10.1172/JCI33673

Linderman GC, Zhao J, Kluger Y. 2018. Zero-preserving imputation of scRNA-seq data using

low-rank approximation. *bioRxiv* 397588. doi**:** https://doi.org/10.1101/397588

Liu L, Cheung TH, Charville GW, Rando T a. 2015. Isolation of skeletal muscle stem cells by

fluorescence-activated cell sorting. *Nat Protoc* **10**:1612–1624. doi:10.1038/nprot.2015.110

Liu Y, Berendsen AD, Jia S, Lotinun S, Baron R, Ferrara N, Olsen BR. 2012. Intracellular

VEGF regulates the balance between osteoblast and adipocyte differentiation. *J Clin Invest*

**122**:3101–3113. doi:10.1172/JCI61209

Loiben AM, Soueid-Baumgarten S, Kopyto RF, Bhattacharya D, Kim JC, Cosgrove BD. 2017.

Data-Modeling Identifies Conflicting Signaling Axes Governing Myoblast Proliferation and

Differentiation Responses to Diverse Ligand Stimuli. *Cell Mol Bioeng* **10**:433–450.

doi:10.1007/s12195-017-0508-5

Mac Gabhann F, Qutub AA, Annex BH, Popel AS. 2011. Systems biology of pro-angiogenic

therapies targeting the VEGF system. *Wiley Interdiscip Rev Syst Biol Med* **2**:694–707.

doi:10.1002/wsbm.92

Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD,

Hawrylycz MJ, Jones AR, Lein ES, Zeng H. 2010. A robust and high-

throughput Cre reporting and characterization system for the whole mouse brain. *Nat*

*Neurosci* **13**:133-40. doi: 10.1038/nn.2467. Epub 2009 Dec 20

Mayeuf-Louchart A, Lagha M, Danckaert A, Rocancourt D, Relaix F, Vincent SD, Buckingham

M. 2014. Notch regulation of myogenic versus endothelial fates of cells that migrate from

the somite to the limb. *Proc Natl Acad Sci U S A* **111**:8844–8849.

doi:10.1073/pnas.1407606111

Mayeuf-Louchart A, Montarras D, Bodin C, Kume T, Vincent SD, Buckingham M. 2016.

Endothelial cell specification in the somite is compromised in Pax3-positive progenitors of

Foxc1/2 conditional mutants, with loss of forelimb myogenesis. *Development* **143**:872–9.

doi:10.1242/dev.128017

Mercatelli N, Dimauro I, Ciafré SA, Farace MG, Caporossi D. 2010. AlphaB-crystallin is

involved in oxidative stress protection determined by VEGF in skeletal myoblasts. *Free*

*Radic Biol Med* **49**:374–82. doi:10.1016/j.freeradbiomed.2010.04.027

Messina S, Mazzeo A, Bitto A, Aguennouz M, Migliorato A, De Pasquale MG, Minutoli L,

Altavilla D, Zentilin L, Giacca M, Squadrito F, Vita G. 2007. VEGF overexpression via

adeno-associated virus gene transfer promotes skeletal muscle regeneration and enhances

muscle function in mdx mice. *FASEB J* **21**:3737–3746. doi : 10.1096/fj.07-8459com

Minasi MG, Riminucci M, De Angelis L, Borello U, Berarducci B, Innocenzi A, Caprioli A,

Sirabella D, Baiocchi M, De Maria R, Boratto R, Jaffredo T, Broccoli V, Bianco P, Cossu G.

2002. The meso-angioblast: a multipotent, self-renewing cell that originates from the dorsal

aorta and differentiates into most mesodermal tissues. *Development* **129**:2773–83. doi:

10.1242/dev.129.11.2773

Miquerol L, Gertsenstein M, Harpal K, Rossant J, Nagy  a. 1999. Multiple developmental roles

of VEGF suggested by a LacZ-tagged allele. *Dev Biol* **212**:307–322.

doi:10.1006/dbio.1999.9355

Motohashi N, Asakura A. 2014. Muscle satellite cell heterogeneity and self-renewal. *Front Cell*

*Dev Biol* **2**:1–11. doi:10.3389/fcell.2014.00001

Motohashi N, Asakura Y, Asakura A. 2014. Isolation, culture, and transplantation of muscle

satellite cells. *J Vis Exp* 50846. doi:10.3791/50846

Motoike T, Markham DW, Rossant J, Sato TN. 2003. Evidence for novel fate of Flk1+

progenitor: contribution to muscle lineage. *Genesis* **35**:153–9. doi:10.1002/gene.10175

Murach KA, Vechetti IJ, Van Pelt DW, Crow SE, Dungan CM, Figueiredo VC, Kosmac K, Fu X,

Richards CI, Fry CS, McCarthy JJ, Peterson CA. 2020. Fusion-Independent Satellite Cell

Communication to Muscle Fibers During Load-Induced Hypertrophy. *Function* **1**:1–15.

doi:10.1093/function/zqaa009

Murphy MM, Lawson J a, Mathew SJ, Hutcheson D a, Kardon G. 2011. Satellite cells,

connective tissue fibroblasts and their interactions are crucial for muscle regeneration. *J*

*Cell Sci* **124**:e1–e1. doi:10.1242/jcs098228

848 Noren DP, Chou WH, Lee SH, Qutub AA, Warmflash A, Wagner DS, Popel AS, Levchenko A.

849  2016. Endothelial cells decode VEGF-mediated Ca2+ signaling patterns to produce distinct

850  functional responses. *Sci Signal* **9**:ra20–ra20. doi:10.1126/scisignal.aad3188

851 Okabe K, Kobayashi S, Yamada T, Kurihara T, Tai-nagara I. 2014. Neurons Limit Angiogenesis

852  by Titrating VEGF in Retina. *Cell* **159**:584–596. doi:10.1016/j.cell.2014.09.025

853 Olfert IM, Howlett RA, Tang K, Dalton ND, Gu Y, Peterson KL, Wagner PD, Breen EC. 2009.

854  Muscle-specific VEGF deficiency greatly reduces exercise endurance in mice. *J Physiol*

855  **587**:1755–1767. doi: : 10.1113/jphysiol.2008.164384

856 Pallafacchina G, François S, Regnault B, Czarny B, Dive V, Cumano A, Montarras D,

857  Buckingham M. 2010. An adult tissue-specific stem cell in its niche: a gene profiling

858  analysis of in vivo quiescent and activated muscle satellite cells. *Stem Cell Res* **4**:77–91.

859  doi:10.1016/j.scr.2009.10.003

860 Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis

861  of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**:1650–1667.

862  doi:10.1038/nprot.2016.095

863 Podkalicka P, Mucha O, Kaziród K, Bronisz-Budzyńska I, Ostrowska-Paton S, Tomczyk M,

864  Andrysiak K, Stępniewski J, Dulak J, Łoboda A. 2021. Age-Dependent Dysregulation of

865  Muscle Vasculature and Blood Flow Recovery after Hindlimb Ischemia in the mdx Model

866  of Duchenne Muscular Dystrophy. *Biomedicines* **9**:481. doi: 10.3390/biomedicines9050481.

867 Poesen K, Lambrechts D, Van Damme P, Dhondt J, Bender F, Frank N, Bogaert E, Claes B,

868  Heylen L, Verheyen A, Raes K, Tjwa M, Eriksson U, Shibuya M, Nuydens R, Van Den

869  Bosch L, Meert T, D'Hooge R, Sendtner M, Robberecht W, Carmeliet P. 2008. Novel Role

870  for Vascular Endothelial Growth Factor (VEGF) Receptor-1 and Its Ligand VEGF-B in

Motor Neuron Degeneration. *J Neurosci* **28**:10451–10459. doi:10.1523/JNEUROSCI.1092-

08.2008

Roobrouck VD, Clavel C, Jacobs SA, Ulloa-Montoya F, Crippa S, Sohni A, Roberts SJ, Luyten

FP, Van Gool SW, Sampaolesi M, Delforge M, Luttun A, Verfaillie CM. 2011.

Differentiation potential of human postnatal mesenchymal stem cells, mesoangioblasts, and

multipotent adult progenitor cells reflected in their transcriptome and partially influenced by

the culture conditions. *Stem Cells* **29**:871–82. doi:10.1002/stem.633

RStudio Team. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.

http://www.rstudio.com/

Ryall JG, Dell'Orso S, Derfoul A, Juan A, Zare H, Feng X, Clermont D, Koulnis M, Gutierrez-

Cruz G, Fulco M, Sartorelli V. 2015. The NAD+-dependent sirt1 deacetylase translates a

metabolic switch into regulatory epigenetics in skeletal muscle stem cells. *Cell Stem Cell*

**16**:171–183. doi:10.1016/j.stem.2014.12.004

Sakurai H, Okawa Y, Inami Y, Nishio N, Isobe K. 2008. Paraxial mesodermal progenitors

derived from mouse embryonic stem cells contribute to muscle regeneration via

differentiation into muscle satellite cells. *Stem Cells* **26**:1865–73.

doi:10.1634/stemcells.2008-0173

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.

*Nat Methods* **9**:676–82. doi:10.1038/nmeth.2019

Shimizu-Motohashi Y, Asakura Y, Motohashi N, Belur NR, Baumrucker MG, Asakura A. 2015.

Pregnancy-induced amelioration of muscular dystrophy phenotype in mdx mice via muscle

membrane stabilization effect of glucocorticoid. *PLoS One*. **10**:e0120325. doi:

10.1371/journal.pone.0120325.

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M,

Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. *Cell* **177**:1888-

1902.e21. doi:10.1016/j.cell.2019.05.031

Tamaki T, Akatsuka A, Ando K, Nakamura Y, Matsuzawa H, Hotta T, Roy RR, Edgerton VR.

2002. Identification of myogenic-endothelial progenitor cells in the interstitial spaces of

skeletal muscle. *J Cell Biol* **157**:571–7. doi:10.1083/jcb.200112106

Tang K, Breen EC, Gerber H-P, Ferrara NM a, Wagner PD. 2004. Capillary regression in

vascular endothelial growth factor-deficient skeletal muscle. *Physiol Genomics* **18**:63–69.

doi:10.1152/physiolgenomics.00023.2004

Torre E, Dueck H, Shaffer S, Gospocic J, Gupte R, Bonasio R, Kim J, Murray J, Raj A. 2018.

Rare Cell Detection by Single-Cell RNA Sequencing as Guided by Single-Molecule RNA

FISH. *Cell Syst* **6**:171-179.e5. doi:10.1016/j.cels.2018.01.014

Tozer S, Bonnin M-A, Relaix F, Di Savino S, García-Villalba P, Coumailleau P, Duprez D. 2007.

Involvement of vessels and PDGFB in muscle splitting during chick limb development.

*Development* **134**:2579–91. doi:10.1242/dev.02867

Tseng BS, Zhao P, Pattison JS, Gordon SE, Granchelli JA, Madsen RW, Folk LC, Hoffman EP,

Booth FW. 2002. Regenerated mdx mouse skeletal muscle shows differential mRNA

expression. *J Appl Physiol* **93**:537–45. doi:10.1152/japplphysiol.00202.2002

Turaç G, Hindley CJ, Thomas R, Davis JA, Deleidi M, Karaöz E, Pruszak J. 2013. Combined

Flow Cytometric Analysis of Surface and Intracellular Antigens Reveals Surface Molecule

Markers of Human Neuropoiesis. *PLoS One* **8**:1–14. doi:10.1371/journal.pone.0068519

van Velthoven CTJ, de Morree A, Egner IM, Brett JO, Rando TA. 2017. Transcriptional

Profiling of Quiescent Muscle Stem Cells In Vivo. *Cell Rep* **21**:1994–2004.

doi:10.1016/j.celrep.2017.10.037

Veldman MB, Zhao C, Gomez G a, Lindgren AG, Huang H, Yang H, Yao S, Martin BL,

Kimelman D, Lin S. 2013. Transdifferentiation of fast skeletal muscle into functional

endothelium in vivo by transcription factor etv2. *PLoS Biol* **11**:e1001590.

doi:10.1371/journal.pbio.1001590

Vempati P, Popel AS, Mac Gabhann F. 2014. Extracellular regulation of VEGF: Isoforms,

proteolysis, and vascular patterning. *Cytokine Growth Factor Rev* **25**:1–19.

doi:10.1016/j.cytogfr.2013.11.002

Verma M, Asakura Y, Hirai H, Watanabe S, Tastad C, Fong G-H, Ema M, Call JA, Lowe DA,

Asakura A. 2010. Flt-1 haploinsufficiency ameliorates muscular dystrophy phenotype by

developmentally increased vasculature in mdx mice. *Hum Mol Genet* **19**:4145–59.

doi:10.1093/hmg/ddq334

Verma M, Asakura Y, Murakonda BSR, Pengo T, Latroche C, Chazaud B, McLoon LK,

Asakura A. 2018. Muscle Satellite Cell Cross-Talk with a Vascular Niche Maintains

Quiescence via VEGF and Notch Signaling. *Cell Stem Cell* **23**:530-543.e9.

doi:10.1016/j.stem.2018.09.007

Verma M, Shimizu-Motohashi Y, Asakura Y, Ennen JP, Bosco J, Zhou Z, Fong G, Josiah S,

Keefe D, Asakura A. 2019. Inhibition of FLT1 ameliorates muscular dystrophy phenotype

by increased vasculature in a mouse model of Duchenne muscular dystrophy. *PLOS Genet*

**15**:e1008468. doi:10.1371/journal.pgen.1008468

Vieira NM, Elvers I, Alexander MS, Moreira YB, Eran A, Gomes JP, Marshall JL, Karlsson EK,

Verjovski-Almeida S, Lindblad-Toh K, Kunkel LM, Zatz M. 2015. Jagged 1 Rescues the

Duchenne Muscular Dystrophy Phenotype. *Cell* **163**:1204–1213.

doi:10.1016/j.cell.2015.10.049

Wagner PD, Olfert IM, Tang K, Breen EC. 2006. Muscle-targeted deletion of VEGF and

exercise capacity in mice. *Respir Physiol Neurobiol* **151**:159–166.

doi:10.1016/j.resp.2005.09.007

Wang C, Liu W, Nie Y, Qaher M, Horton HE, Yue F, Asakura A, Kuang S. 2017. Loss of MyoD

Promotes Fate Transdifferentiation of Myoblasts Into Brown Adipocytes. *EBioMedicine*

**16**:212-223. doi: 10.1016/j.ebiom.2017.01.015.

Xin C, Chu X, Wei W, Kuang B, Wang Y, Tang Y, Chen J, You H, Li C, Wang B. 2021.

Combined gene therapy via VEGF and mini-dystrophin synergistically improves

pathologies in temporalis muscle of dystrophin/utrophin double knockout mice. *Hum Mol*

*Genet* **30**:1349-1359. doi: 10.1093/hmg/ddab120.

Yan G, Yan R, Chen C, Chen C, Zhao Y, Qin W, Veldman MB, Li S, Lin S. 2019. Engineering

vascularized skeletal muscle tissue with transcriptional factor ETV2-induced autologous

endothelial cells. *Protein Cell* **10**:217–222. doi:10.1007/s13238-018-0542-7

Yan H, Guo Y, Zhang P, Zu L, Dong X, Chen L, Tian J, Fan X, Wang N, Wu X, Gao W. 2005.

Superior neovascularization and muscle regeneration in ischemic skeletal muscles following

VEGF gene transfer by rAAV1 pseudotyped vectors. *Biochem Biophys Res Commun*

**336**:287–298. doi: 10.1016/j.bbrc.2005.08.066

Young MD, Behjati S. 2020. SoupX removes ambient RNA contamination from droplet-based

single-cell RNA sequencing data. *Gigascience* **9**:giaa151. doi: 10.1093/gigascience/giaa151.

Yue L, Wan R, Luan S, Zeng W, Cheung TH. 2020. Dek Modulates Global Intron Retention

during Muscle Stem Cells Quiescence Exit. *Dev Cell* **53**:661-676.e6.

doi:10.1016/j.devcel.2020.05.006

Zhang MJ, Ntranos V, Tse D. 2020. Determining sequencing depth in a single-cell RNA-seq

experiment. *Nat Commun* **11**:1–11. doi:10.1038/s41467-020-14482-y

Zheng B, Cao B, Crisan M, Sun B, Li G, Logar A, Yap S, Pollett JB, Drowley L, Cassino T,

Gharaibeh B, Deasy BM, Huard J, Péault B. 2007. Prospective identification of myogenic

endothelial cells in human skeletal muscle. *Nat Biotechnol* **25**:1025-34. doi:

10.1038/nbt1334

     **Figure Legends**

**Figure 1. EC gene signal including VEGF receptor genes in MuSCs.**

A) Experimental schema for bulk and single cell sequencing from the

$Pax7^{CreERT2}:R26R^{tdT}:Flk1^{GFP}$ mice. Bulk sequencing performed on MuSCs, ECs and

single muscle fibers (SMFs) from uninjured muscle. FACS sorted MuSCs and ECs from

uninjured and regenerating TA muscle (3 days following CTX) were run separately on

the 10X single cell platform and aggregated.

B) Bulk RNAseq showing EC signature in MuSCs. Subset dividing genes that are

commonly used to delineate cell identity for MuSCs, ECs and SMFs. Last column shows

genes that define macrophages (Mφ), which should not be highly expressed in any on our

cell types. Red dots indicate MuSCs, green dots indicate ECs and blue dots indicate

SMFs.

C) UMAP from aggregated single cell RNAseq shows expression of different phases of

MuSCs (quiescent MuSCs, activated MuSCs and myoblasts), ECs (tip ECs and ECs) and

from likely contaminant cells such as macrophages (Mφ) and smooth muscle cells

(SMC).

D) UMAP from aggregated data visualized by sample day showing MuSCs segregated by

the sample day but overlap in the EC population. Red dots indicate intact (day 0) and

blue dots indicate 3 days following CTX.

E) Expression of quality control genes such as *eGFP*, *tdTomato*, *CreERT2* and EC genes

such as *Cdh5*, *Kdr* and *Flt1*.

F) Genome browser tracks of whole muscle and TU-tagged MuSC nascent RNA (GSE97399, van Velthoven et al., 2017). *Kdr* and *Pecam1* expression can be found in the MuSC fraction. As control, *Myh1* is highly expressed in the whole muscle preparation but largely absent in the MuSC fraction. *Sdc4* and *Calcr* are highly expressed in MuSC and less so in the whole muscle fraction.

G) qPCR for *Kdr, Flt1, Nrp1* and *Nrp2* in EC lines (B.end3 and C166), muscle cell line (C2C12) and MuSC-derived myoblasts in growth and differentiation medium (DM) shows low level expression of VEGFRs and VEGF co-receptors.

H) RNAScope of *Flt1* on freshly isolated single muscle fibers from *Pax7^{tdT}* mice shows *Flt1* expression (green) and tdTomato (red) in MuSCs. Nuclei were counterstained with DAPI (blue). Scale bar indicates 5 µm.

I) Immunostaining for PECAM1, VE-cadherin (VE-Cad), VEGFA co-receptors (NRP1 and NRP2) and VEGFA receptors (FLT1 and FLK1) in B.End3 EC cell line and MuSC-derived myoblasts (MB). Nuclei were counterstained with DAPI (blue). Scale bar indicates 20 µm.

**Figure 2. VEGFA-FLT1-AKT1 axis controls apoptosis in MuSC *in vitro*.**

A) Immunostaining for VEGFA (green) in MuSC-derived myoblasts. Nuclei were counterstained with DAPI (blue). Scale bar indicates 20 µm.

B) Experimental scheme for assessing apoptosis following thapsigargin induction in myoblast culture.

C) Decreased cell survival in myoblast *in vitro* as VEGFA is blocked using FLT1-FC (a VEFGA trap) following thapsigargin induction. This phenotype is partially rescued with exogenous VEGFA (50 ng/ml).

D) Graphical representation of the VEGF pathway inhibitors used in panel E.

E) Following thapsigargin induction, apoptotic and necrotic cells are increased with inhibition of FLT1 via FLT1-FC or anti-FLT1 antibody (anti-FLT1 mAb) but not FLK1 (SU5402 and ZM306416) or NRP1-FLK1 inhibition (EG00229) following exogenous VEGFA (50 ng/ml).

F) 4-OHT induced deletion of *Flt1* in *Pax7$^{+/CreER}$:Flt1$^{Loxp/Loxp}$* myoblasts is sufficient to reduce cell survival in myoblast without induction of apoptosis.

G) Cell survival is decreased *in vitro* in myoblast with thapsigargin induction following 4-OHT mediated deletion of *Flt1* in *Pax7$^{+/CreER}$:Flt1$^{Loxp/Loxp}$* myoblast that is not rescued by exogenous VEGFA. Blue indicates *MuSC-Flt1$^{+/+}$*, red indicates *MuSC-Flt1$^{+/+}$* with 50 ng/ml VEGFA, green indicates *MuSC-Flt1$^{Δ/Δ}$* and purple indicates *MuSC-Flt1$^{Δ/Δ}$* with 50 ng/ml VEGFA.

H) Representative images of pAKT (red) in myoblast stained by MyoD (green) in *MuSC-Flt1$^{+/+}$* and *MuSC-Flt1$^{Δ/Δ}$* myoblasts induced with exogenous VEGFA. Nuclei were counterstained with DAPI (blue). Scale bar indicates 50 μm.

I) Quantification of pAKT in myoblasts stained by MyoD in *MuSC-Flt1$^{+/+}$* and *MuSC-Flt1$^{Δ/Δ}$* myoblast induced w/wo exogenous VEGFA. VEGFA induction increases pAKT in *MuSC-Flt1$^{+/+}$* myoblasts but this response is lost in *MuSC-Flt1$^{Δ/Δ}$* myoblasts.

1037  J) Annexin V quantification of myoblasts transfected with myr-AKT and E4ORF1 to

1038    activate AKT1 showed increased cell survival of myoblasts following thapsigargin

1039    induction.

1040  K) Representative model for VEGFA-FLT1-AKT axis-mediated MuSC survival.

1041

1042  **Figure 3. MuSC-derived *VEGFA* and *Flt1* requires proper skeletal muscle.**

1043  A) Experimental schema detailing the experiments performed in this figure. The

1044    *Pax7$^{+/CreER}$:R26R$^{tdT}$:VEGFA$^{+/Hyper}$* (*VEGFA$^{+/Hyper}$*) *Pax7$^{+/CreER}$:R26R$^{tdT}$:VEGFA$^{Loxp/Loxp}$*

1045    *for MuSC-VEGFA$^{Δ/Δ}$* and *Pax7$^{tdT}$:Flt1$^{Loxp/Loxp}$* for *MuSC-Flt1$^{Δ/Δ}$* lines were pulsed with

1046    tamoxifen (TMX) prior to investigation.

1047  B) Representative H&E-stained images for intact and on 14-day post injury TA muscle from

1048    *MuSC-VEGFA$^{+/Hyper}$, MuSC-Flt1$^{Δ/Δ}$ and MuSC-VEGFA$^{Δ/Δ}$* mice and their representative

1049    controls. Scale bar indicates 50 μm.

1050  C) Annexin V staining show less necrotic cells in MuSC from *VEGFA$^{+/Hyper}$* mice compared

1051    with the control one day following injury.

1052  D) Quantification of MuSCs from single muscle fibers show increased MuSCs in

1053    *VEGFA$^{+/Hyper}$* EDL muscle compared with the control mice at base line and 14 days post

1054    injury.

1055  E) Fiber size distribution and F) mean feret's diameter of uninjured and regenerating muscle

1056    14 days post injury from *VEGFA$^{+/Hyper}$* and control mice show no difference at baseline

1057    but an increase in fiber diameter following injury.

1058  G) Annexin V staining show increased dead cells in MuSCs from *MuSC-VEGFA$^{Δ/Δ}$* mice

1059    one day following CTX compared with the control *MuSC-VEGFA$^{+/+}$* mice.

H) Quantification of MuSCs from single muscle fiber at base line and 14 days post injury.

I) Fiber size distribution and J) mean feret's diameter of uninjured and regenerating muscle 14 days post injury from $MuSC\text{-}VEGFA^{\Delta/\Delta}$ and $MuSC\text{-}VEGFA^{+/+}$ mice show no difference at baseline but a decrease in fiber diameter following injury.

K) Annexin V staining show increased apoptosis in MuSCs from $MuSC\text{-}Flt1^{\Delta/\Delta}$ mice one day following injury compared with the control $MuSC\text{-}Flt1^{+/+}$ mice.

L) Quantification of MuSCs from single muscle fiber show decreased MuSCs in $MuSC\text{-}Flt1^{\Delta/\Delta}$ EDL muscle at base line and 14 days post injury compared with the control $MuSC\text{-}Flt1^{+/+}$ mice.

M) Fiber size distribution and N) mean feret's diameter of uninjured and regenerating muscle from $MuSC\text{-}Flt1^{\Delta/\Delta}$ and compared with the control $MuSC\text{-}Flt1^{+/+}$ mice show no difference at baseline but a decrease in fiber diameter following injury.

**Figure 4. VEGFA-FLT1 pathway in MuSCs regulates muscle pathology in DMD model mice.**

A) Experimental schema detailing the experiments performed in this figure. The $mdx{:}Pax7^{tdT}{:}Flt1^{Loxp/Loxp}$ was pulsed with tamoxifen (TMX) to generate $mdx{:}MuSC\text{-}Flt1^{\Delta/\Delta}$ mice prior to investigation. $mdx{:}VEGFA^{+/Hyper}$ mouse line was used without any induction.

B) Representative H&E and Sirus red stain from $mdx{:}MuSC\text{-}Flt1^{+/+}$ and $mdx{:}MuSC\text{-}Flt1^{\Delta/\Delta}$ mouse TA muscle at 3 months of age. Scale bar indicates 50 µm.

C) Smaller average fiber size in $mdx{:}MuSC\text{-}Flt1^{\Delta/\Delta}$ compared with the control $mdx{:}MuSC\text{-}Flt1^{+/+}$ mouse TA muscle.

D) Increased fibrotic area in *mdx:MuSC-Flt1$^{\Delta/\Delta}$* compared with the control *mdx:MuSC-*

*Flt1$^{+/+}$* mouse TA muscle.

E) Decreased muscle perfusion in *mdx:MuSC-Flt1$^{\Delta/\Delta}$* compared with the control *mdx:MuSC-*

*Flt1$^{+/+}$* mouse TA muscle.

F) Decreased grip strength normalized to body weight in *mdx:MuSC-Flt1$^{\Delta/\Delta}$* compared with

the control *mdx:MuSC-Flt1$^{+/+}$* mouse TA muscle at both 3 and 12 months of age.

G) Representative H&E and Sirus red stain from *mdx:VEGFA$^{+/Hyper}$* and *mdx:VEGFA$^{+/+}$*

mouse TA muscle at 3 months from TA muscle (top four panels) and diaphragm muscle

(bottom four panels).

H) Increased average fiber size in *mdx:VEGFA$^{+/Hyper}$* compared with the control

*mdx:VEGFA$^{+/+}$* mouse TA and diaphragm (DM) muscle. Scale bar indicates 50 μm.

I) Decreased fibrosis in *mdx:VEGFA$^{+/Hyper}$* compared with the control *mdx:VEGFA$^{+/+}$*

mouse TA muscle.

J) Decreased fibrosis in *mdx:VEGFA$^{+/Hyper}$* compared with the control *mdx:VEGFA$^{+/+}$*

mouse diaphragm (DM) muscle.

K) Muscle perfusion is increased in *mdx:VEGFA$^{+/Hyper}$* compared with the control

*mdx:VEGFA$^{+/+}$* mouse TA muscle.

L) Grip strength normalized to body weight is increased in *mdx:VEGFA$^{+/Hyper}$* compared

with the control *mdx:VEGFA$^{+/+}$* mice.

# Figure 1

GSE97399; van Velthoven et al., 2017

# Figure 2

# Figure 3

# Figure 4