Title: Comparing the efficiency of monofilament and traditional nets for capturing bats

Authors: Silvia Chaves-Ramírez¹, Christian Castillo-Salazar¹, Mariela Sánchez-Chavarría², Hellen Solís-Hernández², Gloriana Chaverri^{2,3*}

ORCID iDs: Christian Castillo-Salazar 0000-0003-0468-983X, Gloriana Chaverri 0000-0002-1155-432X

Affiliations: 1 Escuela de Biología, Universidad de Costa Rica, San Pedro, Costa Rica; 2 Sede del Sur, Universidad de Costa Rica, Golfito, Costa Rica; 3 Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá

*Author for correspondence. E-mail: gloriana.chaverri@ucr.ac.cr

18   **Abstract**

19   Traditional mist nets used for capturing bats have several drawbacks, particularly that they are
 20   inefficient at sampling many insectivorous species. One possible alternative is to use monofilament
 21   nets, whose netting is made of single strands of yarn instead of several as regular nets, making them
 22   less detectable. To date, no study has quantified the capture efficiency of monofilament nets
 23   compared to regular mist nets for the study of bats. Here we compare capture efficiency of
 24   monofilament and regular mist nets, focusing on bat abundance and species diversity at a lowland
 25   tropical forest in southwestern Costa Rica. During our sampling period, we captured 90 individuals
 26   and 14 species in regular nets and 125 individuals and 20 species in monofilament nets. The use of
 27   monofilament nets increased overall capture rates, but most notably for insectivorous species.
 28   Species accumulation curves indicate that samples based on regular nets are significantly
 29   underestimating species diversity, most notably as these nets fail at sampling rare species. We show
 30   that incorporating monofilament nets into bat studies offers an opportunity to expand records of
 31   different guilds and rare bat species and to improve our understanding of poorly-known bat
 32   assemblages while using a popular, relatively cheap and portable sampling method.

33

34   Key words: bats, Chiroptera, Costa Rica, mist nets.

35

36

37 **Introduction**

38 In animal research studies it is often necessary to capture the study organism, and the sample of
 39 individuals that are trapped should ideally be representative of the target species or assemblage.
 40 However, there are many problems associated with trapping methods that hamper the quality of data
 41 obtained, most notably differences in capturing success among individuals or species. A well-known
 42 phenomenon, for example, is the difference in trap-shy or trap-happy species, or those where
 43 individuals consistently avoid traps or those which consistently seek them, respectively, which
 44 creates serious sampling biases [1]. Also, some individuals within a species are easily trapped while
 45 others are not [2], and some sampling methods are consistently more effective than others, even
 46 within the same taxon [3,4]. These biases may significantly affect a study's results and their
 47 interpretation, which thus deteriorates the decisions involving the taxon under study or its habitats.
 48 For example, capture methods may bias estimates of density and population structure [2], population
 49 trends [5], and estimates of species richness and capture rates [6]. It is still possible to account for
 50 capture probabilities as a way to correct for some of these biases [2], yet this requires thorough
 51 knowledge of a trapping method's effect on all target individuals and species [5,7], data which are
 52 commonly not available for the majority of species. Therefore, selecting an appropriate and
 53 representative trapping method is critical for making the correct inferences and appropriate decisions
 54 concerning the focal organism.

55 Bats are the most widely distributed terrestrial mammals on Earth and constitute almost one-
 56 fifth of mammalian biodiversity [8]. However, their ability to fly and nocturnal habits make them a
 57 difficult group to study [9]; thus, efficient sampling methods are essential for their capture and
 58 identification. Despite their large diversity and wide distribution, information on many species is still
 59 deficient, especially in areas that harbor the greatest diversity [10]. A wide variety of methods exists
 60 for the study of bats, which differ in effectiveness and practicality depending on the goal of the

61 study. Among the most commonly known methods for capturing and handling bats are mist nets,
62 harp traps, hand nets, and direct captures at their roosts [11]. On the other hand, indirect recording of
63 species has increased with the use of camera traps, thermal cameras and acoustic recording
64 equipment [12].

65 Bat capture methods, such as mist nets, are considered more invasive as they increase the
66 stress associated with the capture process and require substantial previous experience, especially
67 during the process of extracting individuals from the net [13,14]; however, they are essential for
68 collecting information on morphometrics, acoustics, sensitivity to ectoparasites, species diversity,
69 and population health, among others. An additional problem with the use of mist nets is that some
70 species are very difficult to capture, either because they fly extremely high or because their sensory
71 abilities allow them to detect and therefore avoid the nets [15,16]. Therefore, other trapping methods
72 have been developed, such as harp traps, which tend to reflect fewer high-frequency echoes
73 compared to traditional mist nets, making it easier to trap species that use high-frequency calls,
74 especially those that feed on insects [17–19]. But harp traps also have limitations; for example, their
75 sampling area is small, making it necessary to increase the number of traps and to have prior
76 knowledge of flight routes to increase capture efficiency. On the other hand, indirect methods such
77 as acoustic monitoring may be advantageous since they do not cause stress to the animals and may
78 be able to record individuals that fly very high and/or are difficult to capture. Acoustic monitoring
79 studies, however, depend on the availability of sufficiently large and representative acoustic libraries,
80 which are not yet available for many sites and species [20]. In addition, the use of acoustic recording
81 equipment imposes higher economic costs than other existing capture methods.

82 Despite the difficulties described above, mist nets remain an essential, practical and
83 accessible method for capturing bats, facilitating research and monitoring of species worldwide.
84 Monofilament nets were developed as a novel tool for bat trapping, and these could have great

potential to minimize some of the detectability limitations of using mist nets, while still benefiting from the already well-described advantages of this trapping method. Monofilament nets are designed to be less detectable by bats, as their netting is made of single strands of yarn, unlike traditional nets whose netting is created by twisting several individual strands. However, to date no study has quantified their capture efficiency compared to traditional mist nets. Therefore, the goal of our study is to quantify the efficiency of monofilament mist nets for capturing bats when compared to traditional nylon mist nets. Due to the characteristics of monofilament mist nets, we expect that they will capture i) a greater number of individuals, and ii) a greater diversity of species compared to traditional mist nets. We also expect to capture iii) a greater number of insectivorous bats in monofilament nets, given that they are known to effectively detect traditional mist nets.

Materials and methods

The study was conducted from March 21-28, 2021 at La Cherenga Field Station located at Km 23, Guaycará district, Golfito, Costa Rica (8.639719 N, 83.074489 W). The station is located at 50 m above sea level, and its habitat corresponds to a tropical broad-leaved evergreen lowland forest with average temperatures ranging from 24 to 27°C and annual rainfall of approximately 5000 mm [21]. A large portion of the property is covered by forest in various stages of succession, in addition to an oil palm (*Elais guineensis*) plantation and grasslands. Our sampling was conducted for 8 nights total and was alternated between habitat types, such that in total we sampled 4 nights in areas surrounded by forest (late secondary or primary) and 4 nights in the oil palm plantation.

We captured bats in mist nets (Ecotone, Poland) arranged in a block design; this block included a monofilament net and a regular nylon net (Figure 1) placed next to each other in two possible configurations, depending on space availability, either in a straight line or in an L shape. The position of each type of net within the block was randomly selected. The nets ranged in length

109 from 9 to 12 m; we always tried to place similarly-sized nets in the same block, but an exact match
 110 was not always possible given the net sizes available to us. The monofilament nets we used were
 111 made with 0.08 mm single strand nylon, 14 mm mesh size, 4 shelves and were 2.4 m high. The
 112 regular nets were also made with nylon but of multiple strands, and they had a 19 mm mesh size, 5
 113 shelves and were 3 m high. Most nets were placed at ground level, but during the last 4 nights we
 114 placed a single block at approximately 5 m above the ground.

115

116 Figure 1. A monofilament net (upper panel) compared to a regular net (lower panel).

117

118 We opened nets between sunset at 18:00 until 22:00. Each block of 2 nets was monitored by a
 119 single person, and was checked every 2 minutes to determine the presence of bats in them. If any
 120 individual was captured, it was promptly removed from the net to identify the species [22]. For each
 121 individual, the type of net in which it was captured, the species, sex, age and reproductive status
 122 were noted. Once all data were collected, the animal was released. All sampling protocols followed

guidelines approved by the American Society of Mammalogists for capture, handling and care of mammals [23].

With the data collected we estimated the number of bats captured per net per night. We implemented a correction of the capture effort by estimating the number of individuals captured per square meter per hour, as regular and monofilament nets did not have exactly the same size. We estimated species diversity based on Hill numbers in the package iNext [24], where $q = 0$ represents species richness, $q = 1$ represents the Shannon diversity index, and $q = 2$ represents the Simpson diversity index. Increasing Hill numbers indicate a decreasing emphasis in the contribution of rare species to estimates of diversity [25]. Finally, we determined the diet for bats sampled based on the most commonly known food item consumed by each species or genus [26].

We used R (R Core Team 2018) and lme4 [28] to perform a linear mixed effects analysis of the relationship between the number of bats captured and type of net. As fixed effects we entered type of net and habitat (without interaction term) into the model. As random effects we included block and night. Visual inspection of residual plots did not reveal any obvious deviations from homoscedasticity or normality. P-values were obtained by likelihood ratio tests of the full model with the effect of net type and habitat against the model without the effect of net type, only including the effect of habitat. We also performed a chi-squared test to determine if there was a significant difference in the number of regular or monofilament nets that were able to capture insectivorous and frugivorous bats. Finally, we compared species accumulation curves (interpolated and extrapolated to double the number of captures for the net that captured the most bats) and sample coverage based on Hill numbers for both types of nets used.

Results

We placed a total of 46 nets throughout our 8-night sampling period, 23 regular and 23 monofilament, for a total of 738 and 585.6 m² sampled by regular and monofilament nets, respectively. Both types of nets were opened for 99.5 hours, and in this period, we captured a total of 215 bats from 24 species (table 1); 90 individuals were captured in regular nets while 125 individuals were sampled using monofilament nets. Fourteen species were captured in regular nets and 20 species in monofilament nets. Also, 10 species were only captured in monofilament nets, such as several nectar-feeding bats (i.e., *Hylonycteris underwoodi* and *Lonchophylla concava*), several insectivorous species including many in the genus *Micronycteris* and others such as *Saccopteryx leptura* and *Thyroptera tricolor*. Only 4 species, *Chiroderma villosum*, *Desmodus rotundus*, *Myotis riparius*, and *Trinycteris nicefori*, were solely captured in regular nets. The majority (11 out of 16) of rare species sampled, i.e., those that were captured in 4 or fewer occasions, are considered insectivorous (table 1).

167 Table 1. Number of individuals captured per species in regular and monofilament nets.

	Individuals captured		
Species	Regular	Monofilament	Feeding guild
<i>Artibeus jamaicensis</i>	10	26	Frugivorous
<i>Artibeus lituratus</i>	3	3	Frugivorous
<i>Carollia castanea</i>	8	13	Frugivorous
<i>Carollia perspicillata</i>	27	23	Frugivorous
<i>Carollia sowelli</i>	9	6	Frugivorous
<i>Chiroderma villosum</i>	3	0	Frugivorous
<i>Dermanura watsoni</i>	17	21	Frugivorous
<i>Desmodus rotundus</i>	1	0	Sanguinivorous
<i>Glossophaga soricina</i>	1	7	Nectarivorous
<i>Hylonycteris underwoodi</i>	0	1	Nectarivorous
<i>Lonchophylla concava</i>	0	1	Nectarivorous
<i>Lophostoma brasiliense</i>	0	2	Insectivorous
<i>Micronycteris hirsuta</i>	0	2	Insectivorous
<i>Micronycteris minuta</i>	0	1	Insectivorous
<i>Micronycteris microtis</i>	1	2	Insectivorous
<i>Micronycteris schmidtorum</i>	0	1	Insectivorous
<i>Myotis riparius</i>	3	0	Insectivorous
<i>Peropteryx kappleri</i>	0	1	Insectivorous
<i>Platyrrhinus helleri</i>	0	4	Frugivorous
<i>Saccopteryx bilineata</i>	1	3	Insectivorous
<i>Saccopteryx leptura</i>	0	2	Insectivorous
<i>Thyroptera tricolor</i>	0	1	Insectivorous
<i>Trinycteris nicefori</i>	1	0	Insectivorous
<i>Uroderma bilobatum</i>	5	5	Frugivorous
Total	90	125	

We found that using monofilament nets affected the number of individuals captured ($\chi^2(1) = 6.47, p = 0.01$), increasing capture rates by about 0.04 bats per m^2 per hour compared to regular nets (Figure 2). A significantly larger portion of insectivorous species (9 out of 11; table 1) was captured in monofilament nets compared to those captured in regular nets (4 out of 11; $\chi^2(1) = 4.70, p = 0.03$). Both types of nets were equally efficient at capturing frugivorous species (8 out of 9 species). Monofilament nets also appeared more efficient at capturing rare species than regular nets; 12 out of 16 rare species were captured in monofilament nets, whereas only 6 out of 16 rare species were sampled using regular nets ($\chi^2(1) = 4.57, p = 0.03$).

Figure 2. Violin plots showing the number of bats captured (corrected by sampling effort = bats per m^2 per hour) according to the type of net used and the habitat sampled.

In regular nets, species richness ($q = 0$) was estimated at 11 (s.e. = 4.45), while species richness estimated from captures made in monofilament nets was 20 (s.e. = 4.76). Similar differences

182 were observed using other diversity indexes, including Shannon and Simpson ($q = 1$ & 2,
183 respectively), although the difference in species diversity between net types decreased with
184 increasing values in Hill numbers (Figure 3). Species accumulation curves indicate that samples
185 based on regular nets are significantly underestimating species diversity, even when sample sizes
186 (i.e., individuals captured) are extrapolated to double the study's sampling effort. Also, the calculated
187 sample coverage for both nets is equally large, suggesting that an inflated estimate of sample
188 coverage is created by regular nets even though they seem to be underestimating species diversity
189 (Figure 3).

190
191 Figure 3. Species accumulation curves (upper panels) and estimated sample coverage (lower panels)
192 for the two types of nets used.

193

194 Discussion

195 In this study we compared bat capture efficiency using regular and monofilament mist nets for
 196 sampling Neotropical bat communities. Our results demonstrate that monofilament nets were able to
 197 capture a larger number of individuals and species compared to regular nets. We found that bats of all
 198 trophic guilds were sampled by monofilament nets, yet these nets were particularly more effective at
 199 sampling insectivorous species compared to regular nets. We also show that monofilament nets were
 200 more likely to capture rare or elusive species and that this difference in capture rates was not affected
 201 by habitat type. Thus, net design, particularly with regards to the number of strands used for
 202 constructing the netting, is an important factor to consider when the main interest of the study is to
 203 gauge species diversity or when certain trophic guilds, especially insectivorous, are targeted in bat
 204 surveys.

205 Conspicuousness of mist nets is considered to be a major factor contributing to lower capture
 206 rates in birds; thus, placing nets in the shade or against a dark background significantly increases their
 207 success [29,30]. While no studies to date have assessed how net placement affects capture rates in
 208 bats, our results of an overall increase in the number of sampled individuals strongly suggest that using
 209 monofilament nets may be an alternative and effective way to decrease their detection by bats. This
 210 decrease in detectability of monofilament nets may also be largely responsible for an increased
 211 probability of capturing insectivorous species, which are known to be skilled at detecting and avoiding
 212 regular mist nets [16]. Since insectivorous bats comprised the majority of rare species sampled in our
 213 study, then it is expected that monofilament nets also captured a larger portion of rare species. With
 214 our data, however, we cannot confirm that these seldom-captured species are in fact locally rare, or if
 215 fewer individuals were captured because they remain skilled at avoiding even the monofilament nets.
 216 Despite the latter, monofilament nets were effective at detecting many rare species, particularly those
 217 in the genus *Micronycteris*, which may be missed in bat surveys that use acoustic detectors in addition
 218 to regular nets, as the former may not be able to record low-intensity calls [31,32].

While we found no studies to date that have explicitly compared the efficiency in capture rates between regular and monofilament nets using a paired design like ours, some previous results indicate greater capture rates in the latter. For example, a study by Chaverri et al. (2016) shows that capture rates in regular, polyester, nets were 0.04 bats per net-hour, while capture rates in monofilament nets were 0.13 bats per net-hour. Other studies in birds suggest that both mesh size, color and the number of strands that form the netting influence capture rates of nets placed over fish ponds, with monofilament nets imposing greater risks [34]. The previous results are to be expected, as detectability of capturing devices clearly affects their effectiveness. However, our results further demonstrate that the netting material may affect the species that are sampled through mist-nets.

It is important to emphasize that before deciding to use monofilament nets in a study, all researchers involved must be skillful at removing bats from mist nets, as the strands are often hard to see and they break very easily. Monofilament nets thus require constant revision, since they can be easily and quickly damaged by bats. Therefore, we recommend checking time intervals between 3 to 5 minutes, which represents a shorter time interval compared to what's recommended for regular mist-nets (15 min; (Gannon and Sikes 2007; Kunz et al. 2009). Despite this, we consider that reducing net-checking intervals in general for bat extraction is a practice that should bring significant benefits, particularly regarding animal safety and net capturing efficiency, as an increase in visits could reduce data loss due to escape or predation of bats in the net [35]. We are aware that applying these recommendations may require more field personnel, so researchers should carefully consider the benefits of using monofilament nets, in the form of a significant increase in capturing efficiency, and the costs involved in having additional personnel and an inevitable faster deterioration of their mist nets.

In conclusion, we show that incorporating monofilament nets into bat sampling designs offers not only an opportunity to expand records of different guilds and rare bat species, but ultimately may

help to improve our understanding of poorly-known species and assemblages while still using a relatively cheap and portable method. The use of monofilament nets could help compensate for the known limitations of regular nets (e.g., inefficient at trapping insectivorous bats) and even harp traps (e.g., reduced portability and sampling area), providing an additional tool for the study of bat species. Additional studies are needed to understand the functionality of monofilament nets for other bat assemblages; meanwhile, we advocate for the use of this simple tool and ideally in combination with others, including regular mist nets, but also acoustic and roost surveys.

Ethics statement. All protocols for capturing and handling bats comply with the current laws of the Costa Rican government (permit no. ACOSA-D-R-056-2021).

Data accessibility. All code and raw data have been stored in the GitHub repository (<https://github.com/morceglo/Monofilament-nets-for-bats>).

Author's contributions. All authors participated in the design of the study, collected field data, drafted the manuscript. G.C. conducted statistical analyses. All authors gave final approval for publication.

Competing interests. We have no competing interests.

References

1. Biro PA, Dingemanse NJ. 2009 Sampling bias resulting from animal personality. *Trends Ecol. Evol.* **24**, 66–67. (doi:10.1016/j.tree.2008.11.001)
2. Bisi F, Newey S, Nodari M, Wauters LA, Harrison A, Thirgood S, Martinoli A. 2011 The

- strong and the hungry: bias in capture methods for mountain hares *Lepus timidus*. *Wildlife Biol.* **17**, 311–316. (doi:10.2981/10-133)
3. Williams DF, Braun SE. 1983 Comparison of Pitfall and Conventional Traps for Sampling Small Mammal Populations. *J. Wildl. Manage.* **47**, 841. (doi:10.2307/3808622)
4. Flaquer C, Torre I, Arrizabalaga A. 2007 Comparison of sampling methods for inventory of bat communities. *J. Mammal.* **88**, 526–533.
5. Garshelis DL, Noyce K V. 2006 Discerning Biases in a Large Scale Mark-Recapture Population Estimate for Black Bears. *J. Wildl. Manage.* **70**, 1634–1643.
6. Kolowski JM, Forrester TD. 2017 Camera trap placement and the potential for bias due to trails and other features. *PLoS One* **12**, e0186679. (doi:10.1371/journal.pone.0186679)
7. Baker PJ, Harris S, Robertson CPJ, Saunders G, White PCL. 2001 Differences in the capture rate of cage-trapped red foxes *Vulpes vulpes* and an evaluation of rabies control measures in Britain. *J. Appl. Ecol.* **38**, 823–835. (doi:10.1046/j.1365-2664.2001.00637.x)
8. Burgin CJ, Colella JP, Kahn PL, Upham NS. 2018 How many species of mammals are there? *J. Mammal.* **99**, 1–14. (doi:10.1093/jmammal/gyx147)
9. Voigt CC, Kingston T. 2015 *Bats in the anthropocene: Conservation of bats in a changing world*. (doi:10.1007/978-3-319-25220-9)
10. Frick WF, Kingston T, Flanders J. 2020 A review of the major threats and challenges to global bat conservation. *Ann. N. Y. Acad. Sci.* **1469**, 5–25. (doi:10.1111/nyas.14045)
11. Wilson DE, Cole FR, Nichols JD, Rudran R, Foster MS. 1996 *Measuring and monitoring biological diversity: Standard methods for mammals*. Washington, D.C.: Smithsonian Books. See <http://pubs.er.usgs.gov/publication/5200145>.

- 290 12. Kunz TH, Parsons S. 2009 *Ecological and behavioral methods for the study of bats*.
291 Baltimore, MD: John Hopkins University Press.
- 292 13. Rizo-Aguilar A, Ávila-Torresagaton LG, Fuentes Vargas L, Lara Nuñez AC, Flores Nuñez
293 GI, Albino Miranda S. 2015 Técnicas para el estudio de los murciélagos. In *Manual de*
294 *técnicas del estudio de la fauna* (ed S Gallina Tessaro), Veracruz, México: Instituto de
295 Ecología, A. C.
- 296 14. Kunz TH, Kurta A. 1988 Capture methods and holding devices. In *Ecological and Behavioral*
297 *Methods for the Study of Bats* (ed TH Kunz), pp. 1–29. Washington: Smithsonian Institution
298 Press.
- 299 15. Kalko EK V., Estrada Villegas S, Schmidt M, Wegmann M, Meyer CFJ. 2007 Flying high--
300 assessing the use of the aerosphere by bats. *Integr. Comp. Biol.* **48**, 60–73.
301 (doi:10.1093/icb/icn030)
- 302 16. MacSwiney MC, Clarke FM, Racey PA. 2008 What you see is not what you get: The role of
303 ultrasonic detectors in increasing inventory completeness in Neotropical bat assemblages. *J.*
304 *Appl. Ecol.* **45**, 1364–1371. (doi:10.1111/j.1365-2664.2008.01531.x)
- 305 17. Pech-Canche JM, Estrella E, López-Castillo DL, Hernández-Betancourt SF, Moreno CE. 2011
306 Complementarity and efficiency of bat capture methods in a lowland tropical dry forest of
307 Yucatán, Mexico. *Rev. Mex. Biodivers.* **82**. (doi:10.22201/ib.20078706e.2011.3.683)
- 308 18. Berry N, O’connor W, Holderied MW, Jones G. 2004 Detection and Avoidance of Harp Traps
309 by Echolocating Bats. *Acta Chiropterologica* **6**, 335–346. (doi:10.3161/001.006.0211)
- 310 19. Duffy AM, Lumsden LF, Caddle CR, Chick RR, Newell GR. 2000 The efficacy of AnaBat
311 ultrasonic detectors and harp traps for surveying microchiropterans in south-eastern Australia.
312 *Acta Chiropterologica* **2**, 127–144.

- 313 20. Hayes JP. 2000 Assumptions and practical considerations in the design and interpretation of
314 echolocation-monitoring studies. *Acta Chiropterologica* **2**, 225–236.
- 315 21. Kappelle M, Castro M, Acevedo H, Gonzalez L, Monge H. 2002 *Ecosistemas del Area de*
316 *Conservacion Osa (ACOSA)*. Santo Domingo de Heredia, Costa Rica: Instituto Nacional de
317 Biodiversidad.
- 318 22. York HA, Rodríguez-Herrera B, LaVal RK, Timm RM, Lindsay KE. 2019 Field key to the
319 bats of Costa Rica and Nicaragua. *J. Mammal.* **100**, 1726–1749.
320 (doi:10.1093/jmammal/gyz150)
- 321 23. Sikes RS. 2016 2016 Guidelines of the American Society of Mammalogists for the use of wild
322 mammals in research and education. *J. Mammal.* **97**, 663–688.
323 (doi:10.1093/jmammal/gyw078)
- 324 24. Hsieh TC, Ma KH, Chao A. 2016 iNEXT: An R package for rarefaction and extrapolation of
325 species diversity (Hill numbers). *Methods Ecol. Evol.* **in press**, 1–22.
- 326 25. Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014
327 Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in
328 species diversity studies. *Ecol. Monogr.* **84**, 45–67. (doi:10.1890/13-0133.1)
- 329 26. Kalko EKV, Handley CO, Handley D. 1996 Organization, diversity, and long-term dynamics
330 of a neotropical bat community. In *Long-term Studies in Vertebrate Communities* (eds M
331 Codyand, J Smallwood), pp. 503–553. Los Angeles, California: Academic Press.
- 332 27. R Core T. 2018 R: A language and Environment for Statistical Computing. *Vienna, Austria*
- 333 28. Bates D, Mächler M, Bolker BM, Walker SC. 2015 Fitting Linear Mixed-Effects Models
334 Using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
- 335 29. Jenni L, Leuenberger M, Rampazzi F. 1996 Capture Efficiency of Mist Nets with Comments

on Their Role in the Assessment of Passerine Habitat Use. *J. F. Ornithol.* **67**, 263–274.

30. Keyes BE, Grue CE. 1982 Capturing birds with mist nets: a review. *North Am. Bird Bander* **7**, 2–14.

31. O’Farrell MJ, Gannon WL. 1999 A comparison of acoustic versus capture techniques for the inventory of bats. *J. Mammal.* **80**, 24–30.

32. Portfors C V. *et al.* 2000 Bats from Fazenda Intervales, Southeastern Brazil: species account and comparison between different sampling methods. *Rev. Bras. Zool.* **17**, 533–538. (doi:10.1590/S0101-81752000000200022)

33. Chaverri G, Garin I, Alberdi A, Jimenez L, Castillo-Salazar C, Aihartza J. 2016 Unveiling the hidden bat diversity of a neotropical montane forest. *PLoS One* **11**. (doi:10.1371/journal.pone.0162712)

34. Nemtzov SC, Olsvig-Whittaker L. 2003 The Use of Netting Over Fishponds as a Hazard to Waterbirds. *Waterbirds* **26**, 416–423.

35. Kunz TH, Hodgkison R, Weise CD. 2009 Methods of capturing and handling bats. In *Ecological and Behavioral Methods for the Study of Bats* (eds TH Kunz, S Parsons), pp. 3–35. Baltimore, Maryland: The Johns Hopkins University Press.

36. Gannon WL, Sikes RS. 2007 Guidelines of the American Society of Mammalogists for the Use of Wild Mammals in Research. *J. Mammal.* **88**, 809–823. (doi:10.1644/06-MAMM-F-185R1.1)