Local biodiversity change reflects interactions among changing abundance, evenness and

richness

Shane A. Blowes, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-

Leipzig, Germany; Department of Computer Science, Martin Luther University Halle-

Wittenberg, Halle (Salle), Germany. shane.blowes@idiv.de

Gergana N. Daskalova, School of GeoSciences, University of Edinburgh, Scotland EH9 3FF.

gndaskalova@gmail.com

Maria Dornelas, Centre for Biological Diversity, University of St Andrews, KY16 9TH.

maadd@st-andrews.ac.uk

Thore Engel, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig,

Germany; Department of Computer Science, Martin Luther University Halle-Wittenberg, Halle

(Salle), Germany. thore.engel@idiv.de

Nicholas J. Gotelli, Department of Biology, University of Vermont, Burlington, VT 05405 USA.

Nicholas.Gotelli@uvm.edu

Anne E. Magurran, Centre for Biological Diversity, University of St Andrews, KY16 9TH.

aem1@st-andrews.ac.uk

Inês S. Martins, Leverhulme Centre for Anthropocene Biodiversity and Department of Biology,

University of York, York YO10 5DD, UK; Centre for Biological Diversity, University of St

Andrews, KY16 9TH. iism1@st-andrews.ac.uk

Brian McGill, School of Biology and Ecology and Mitchell Center for Sustainability Solutions,

University of Maine; Orono, ME. mail@brianmcgill.org

Daniel J. McGlinn, Department of Biology, College of Charleston, Charleston, SC.

danmcglinn@gmail.com

Alban Sagouis, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig,

Germany; Department of Computer Science, Martin Luther University Halle-Wittenberg, Halle

(Salle), Germany. alban.sagouis@idiv.de

Hideyasu Shimadzu, Department of Mathematical Sciences, Loughborough University, LE11

3TU, UK; Graduate School of Public Health, Teikyo University, Tokyo, Japan.

H.Shimadzu@lboro.ac.uk

Sarah R. Supp, Data Analytics Program, Denison University, Granville, OH 43023 USA.

supps@denison.edu

Jonathan M. Chase, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-

Leipzig, Germany; Department of Computer Science, Martin Luther University Halle-

Wittenberg, Halle (Salle), Germany. jonathan.chase@idiv.de

Running title: Characterizing multicomponent diversity change

Corresponding author: Shane Blowes, German Centre for Integrative Biodiversity Research

(iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, Leipzig 04103, Germany. shane.blowes@idiv.de

+ 49 341 9733254.

Abstract

Biodiversity metrics often integrate data on the presence and abundance of multiple species. Yet

understanding covariation of changes to the numbers of individuals, the evenness of species'

relative abundances, and the total number of species remains limited. Using individual-based

rarefaction curves, we introduce a conceptual framework to understand how expected positive

relationships among changes in abundance, evenness and richness arise, and how they can break

down. We then examined interdependencies between changes in abundance, evenness and

richness in more than 1100 assemblages sampled either through time or across space. As

predicted, richness changes were greatest when abundance and evenness changed in the same

direction, and countervailing changes in abundance and evenness acted to constrain the

magnitude of changes in species richness. Site-to-site changes in abundance, evenness, and

richness were often decoupled, and pairwise relationships between changes in these components

across assemblages were weak. In contrast, changes in species richness and relative abundance

were strongly correlated for assemblages varying through time. Temporal changes in local

biodiversity showed greater inertia and stronger relationships between the component changes

when compared to site-to-site variation. Local variation in assemblage diversity was rarely due to

a passive sample from a more or less static species abundance distribution. Instead, changing

species relative abundances often dominated local variation in diversity. Moreover, how

changing relative abundances combined with changes to total abundance frequently determined

the magnitude of richness changes. Embracing the interdependencies between changing

abundance, evenness and richness can provide new information for better understanding

biodiversity change in the Anthropocene.

Keywords: biodiversity change, abundance, evenness, species richness, rarefaction

Introduction

Measures of biodiversity and its change are frequently used to determine the magnitude, direction and pace of ecosystem modifications (Diaz et al. 2019). Descriptions of the distribution and abundance of organisms are central to basic ecological research (Krebs 1972, Andrewartha & Birch 1984), and biodiversity (as applied to taxonomic or species diversity) is a multifaceted concept that combines information on the distribution and abundance of multiple species. There have been numerous such aggregate metrics proposed to quantify different aspects of biodiversity (e.g., Hill 1973, Gotelli and Colwell 2001, Magurran & McGill 2011, Chao and Jost 2012), most of which depend on sample effort and scale (Rosenzweig 1995, Whittaker et al. 2001, McGill 2011a, Chase & Knight 2013). This inherent complexity poses significant challenges for quantitative syntheses of temporal and spatial variation in biodiversity.

Here, we argue that in order to understand biodiversity change, it is critical to look beyond aggregate metrics in isolation. Specifically, we need to examine how changes in the key components that lead to aggregate metrics interact and combine with each other. Currently, different components that underlie biodiversity represent important, but are largely viewed as independent lines of evidence for contemporary biodiversity change. For example, one body of work is focused on quantifying total abundances of all species within assemblages, such as recent work documenting declines of birds (e.g., Rosenberg et al. 2019) and insects (Wagner 2020). A related body of work focuses instead on population-level 'winners' and 'losers' within assemblages (McKinney & Lockwood 2000), species with increasing and decreasing trends in occupancy or abundance (e.g., WWF 2020). However, changes in the means of these metrics do not capture the variability and nuance in either the assemblage- and population-level abundance

trends across space and among taxa (e.g., van Klink et al. 2020, Daskalova et al. 2020a, Leung et

al. 2020). Moreover, trends in total assemblage abundance provide only one window into

potentially complex changes that could be occurring in other components of biodiversity, and

while changes to species relative abundances do influence biodiversity metrics, population-level

trends themselves cannot be used to calculate assemblage-level (i.e., biodiversity) metrics (e.g.,

Dornelas et al. 2019). Even when quantifying the same response metric (i.e., species richness),

syntheses of biodiversity change in space due to differential land use conditions have found

declines of species richness (e.g., Newbold et al. 2015, 2018), whereas analyses of time series

have shown that, on average, species richness increases roughly offset decreases (Dornelas et al.

2014, Blowes et al. 2019), despite often significant modifications to climate and habitat (Antão

et al. 2020a, Daskalova et al. 2020b). Finally, the prediction that human activities likely impact

species relative abundances more frequently than species occurrences (Chapin et al. 2000), has

not resulted in a strong focus on assemblage-level evenness in existing syntheses (but see, e.g.,

Crowder et al. 2010, Zhang et al. 2012, Jones & Magurran 2018). Importantly, changes in all

these components -- abundance, evenness, and richness -- contribute to variation in biodiversity.

Yet little is known about how components are changing in combination within assemblages, and

whether certain combinations act to constrain observed biodiversity change.

Species origination (speciation plus colonization) and extinction are the most fundamental

processes for biodiversity dynamics (Storch et al. 2021). These processes combine with

productivity (Currie 1991, Mittelbach et al. 2001), disturbance frequency and intensity (Connell

1978, Miller et al. 2011), historical (Latham & Ricklefs 1993) and biogeographic factors (e.g.,

Kreft et al. 2008), land use modifications (Newbold et al. 2015), and climate change (Antão et al.

2020a) to drive variation in biodiversity. In turn, these processes and drivers combine such that local scale measures of biodiversity estimated from a given (local) sample depends largely on two components (see e.g., He and Legendre 2002, McGill 2011a). First, the total number of individuals (Fisher et al. 1943, Preston 1962), whereby fewer individuals are expected to (non-linearly) lead to fewer species. Second, the total number of species and their relative abundances within the regional species pool (i.e., the set of all potential colonizing species in a region), which we refer to as the Species Abundance Distribution (SAD; McGill et al. 2007). Local samples will have lower species richness when the species pool has relatively few highly abundant species and many rare species (i.e., less even SADs), compared to samples from a species pool of the equivalent size with more equitable species abundances. Moreover, variation or changes in local environmental conditions can alter local patterns of species relative abundances. Whenever two or more samples across space or time differ in the total number of individuals, the shape of the SAD, or both, there will be changes in most metrics of biodiversity. However, changes in abundance and the SAD are not always correlated, and, when decoupled, the magnitude and direction of change in derived biodiversity metrics can differ considerably.

Variation in the total number of individuals is a long-standing, first-order explanation of variation in richness (Fisher et al. 1943, Coleman et al. 1982, Srivastava & Lawton 1998, Gaston 2000, Scheiner & Willig 2005, Storch et al. 2018). In the context of species-area relationships, this has been termed the 'passive sampling hypothesis' (Coleman et al. 1982), and as local assemblages increase in size they are expected to include more species from the regional pool due to sampling processes alone. Larger (Connor & McCoy 1979) or more productive (Wright 1983) areas are also predicted to have more species driven again by increased numbers of

individuals, but in these cases associated with processes other than sampling, such as decreased

likelihood of species extinction with increased population sizes (Preston 1962, Wright 1983,

Srivastava & Lawton 1998), and commonly termed the 'more individuals hypothesis' (Srivastava

& Lawton 1998). Anthropogenic drivers can also influence the number of individuals in

assemblages (e.g., via eutrophication, exploitation, harvesting or land clearing), potentially

impacting biodiversity due to changes to the total number of individuals (Newbold et al. 2015,

Blowes et al. 2020). If biodiversity varies primarily via changes in the numbers of individuals,

positive relationships between altered numbers of individuals and species richness changes are

expected. In such cases, however, other metrics of species diversity that control for variation in

numbers of individuals, such as species richness expected for a given number of individuals—

known as rarefied richness—should be relatively unchanged.

Changes to the shape of the SAD can drive variation in biodiversity through time or space. For

example, co-occurrence and coexistence of species can be altered by changes to resource

diversity (MacArthur 1965), environmental or habitat heterogeneity (Tilman 1982, Shmida &

Wilson 1985), interspecific interactions (e.g., keystone predation; Paine 1974, Menge et al.

2020), biological invasions (Vilà et al. 2011), and external perturbations (Hughes et al. 2007).

Alterations to any of these features can change biodiversity by changing species' relative

abundances and the size of the species pool (via species additions or subtractions).

Anthropogenic factors can also favor some species and disfavor others, potentially altering the

relative abundance of species (e.g., due to selective exploitation; Blowes et al. 2020), or the size

of the species pool (e.g., species with large ranges replacing those with small ranges, Newbold et

al. 2018). In such cases, biodiversity change will be characterized by positive relationships

between species richness change and changes in metrics sensitive to relative abundance, such as

rarefied richness, evenness and diversity metrics sensitive to species relative abundances.

Changing components of biodiversity can covary in different and informative ways. Yet, to date,

there has been little exploration of this covariation in time or space, nor of the theoretical

linkages. For example, whether total abundances and the evenness of species' relative

abundances change in similar or decoupled ways, and how this influences biodiversity change is

largely unknown. However, syntheses of relationships between different biodiversity metrics,

which can reflect different combinations of component changes, have typically found

relationships to be weak. For example, Stirling and Wilsey (2001) showed that although strong

positive correlations between species richness, diversity and evenness metrics were expected

from a neutral model (Caswell 1976), there was considerable variation in the strength, and even

the sign of the relationships in 323 empirical comparisons. Similarly, Soininen et al. (2012)

examined temporal ($n = 212$) and spatial variation ($n = 17$) in aquatic datasets, and again found

considerable heterogeneity in the relationship between richness and evenness. Using data from

91 assemblages, McGill (2011b) concluded that most biodiversity metrics align with three axes

of empirical variation (total abundance, evenness and richness); components subsequently shown

to be relatively uncorrelated across space for a subset of 37 of the 91 assemblages (Chase et al.

2018). Collectively, these studies suggest that static biodiversity estimates are multidimensional,

and that different metrics can covary or be unrelated.

Where ecologists have quantified variation in multiple components of local diversity, the focus

has typically been on averages across assemblages, with each component treated as a separate,

independent response. For example, analyses of the local assemblages documented by the

BioTIME database (Dornelas et al. 2018) show that numbers of individuals, species richness, and

dominance (quantified as the relative abundance of the most numerically dominant species, and

conceptually the complement of evenness) are highly variable among datasets, but on average,

have no directional trend (Dornelas et al. 2014, Jones & Magurran 2018, Blowes et al. 2019). On

the other hand, analyses of the PREDICTS database (Hudson et al. 2017) documenting spatial

contrasts between assemblages in more pristine habitats with those in different land use

categories, show that human-altered habitats frequently have fewer species and often fewer

individuals (Newbold et al. 2015, 2020). However, these results describe average changes across

assemblages estimated independently, whereas, as we describe in more detail below, component

changes are unlikely to be completely independent.

Here, we first provide a conceptual overview of how changes in the main components underlying

local biodiversity (total abundance, evenness and species richness) can combine using

individual-based rarefaction curves. Using simplified scenarios with contrasting component

changes, we show that the signs (or direction) of changes in total abundance and evenness can

combine to determine the magnitude of expected richness changes, and whether positive

pairwise relationships prevail. We then empirically assess interdependencies between abundance,

evenness and richness changes using compilations of ecological assemblage data. In the face of

natural and anthropogenically-driven environmental variation in time and space, we ask whether

changes in the components of local biodiversity show positive relationships (i.e., change in the

same direction). Or, alternatively, are component changes sufficiently heterogeneous that

variation in biodiversity depends on which of the underlying components (numbers of

individuals or the SAD) are changing, and how the different component changes combine?

Conceptual relationships between changes in total abundance, evenness and richness

Individual-Based Rarefaction (IBR) curves (Hurlbert 1971, Gotelli & Colwell 2001) are well

suited for visualizing conceptual relationships among changes in total abundance, evenness and

species richness (Fig. 1a, Cayuela et al. 2015, Chase et al. 2018, McGlinn et al. 2019). The end

point of the IBR depicts the total number of individuals of all species combined, and variation

between assemblages in where the IBR terminates quantifies changes to the number of

individuals (ΔN, Fig.1a) and species richness (ΔS, Fig. 1a). The shape (or curvature) of the IBR

curve reflects species' relative abundances and the size of the species pool (i.e., the SAD). We

use two parts of the curve to characterize changes in the SAD between assemblages. First,

because it is standardized to an equal number of individuals (n), changes in rarefied richness, ΔS_n

(Fig.1a), reflects changes to species' relative abundances only. Second, we use the numbers

equivalent (or effective number of species) transformation of the Probability of Interspecific

Encounter (PIE, Hurlbert 1971). The PIE is equal to the slope at the base of the rarefaction curve

(Olszewski 2004) and represents a metric of evenness that is relatively insensitive to sample

effort (more even communities have a higher PIE, Fig. 1a). Transformation of the PIE to the

numbers equivalent (S_{PIE}) aids comparisons to species richness (i.e., ΔS and ΔS_{PIE} have the same

units; Jost 2006). S_{PIE} is equal to the inverse of Simpson concentration (Jost 2006), and diversity

of order $q = 2$ (Hill 1973, Jost 2007), $D = (\sum_{i=1}^{S} p_i^q)^{1/(1-q)} = \frac{1}{\sum_{i=1}^{S} p_i^2}$, where S is the number of

species and p_i is the proportion of the assemblage represented by species i. As a consequence,

changes in S_{PIE} (ΔS_{PIE}) are most strongly influenced by the number of abundant or common

species in assemblages.

To visualize relationships among changes in total abundance, evenness and richness, we plot

pairwise relationships between changes in four components of the IBR. Specifically, changes in

species richness (ΔS) and total abundance (ΔN; Fig. 1b), and species richness and the two metrics

sensitive to changes in relative abundance – rarefied richness, ΔS_n (Fig. 1c), and ΔS_{PIE}, which we

refer to as evenness (Fig. 1d). Changing components with positive relationships (i.e., the same

sign) fall into the lower left and upper right quadrants, whereas assemblage changes that fall into

the upper left or lower right quadrants of Figs. 1b, c, d reflect a negative relationship between the

respective components.

Altered numbers of individuals, but no change to the SAD, can underpin differences in diversity

between assemblages. Changes only to the number of individuals being passively sampled from

the same underlying SAD (Fig. 2a) result in ΔS and ΔN being positively related with the same

sign (Fig. 2g), whereas ΔS_n (Fig. 2h) and ΔS_{PIE} (Figs. 2i) will be approximately zero (and have a

weak or no relationship with ΔS). This has been variously referred to in the literature as a

sampling effect, the rarefaction effect, and the passive sampling hypothesis (Coleman et al. 1982,

Gotelli & Cowell 2001, Palmer et al. 2000).

Changes in species richness (ΔS) can also be solely associated with changes to relative

abundance (i.e., $\Delta N \approx 0$), which weakens or removes the expectation for a positive relationship

between changes in richness and total abundance. For example, changes in species richness can

be associated with SAD changes due, for e.g., to increased numbers of common species,

increased evenness (He & Legendre 2002), or increases to the size of the species pool (Fig. 2b),

which results in a positive relationship between ΔS and ΔS_{PIE} (Figure 2i). Finally, if total

abundance and relative abundance change in the same direction (e.g., more individuals and

increased evenness, Fig. 2c), then positive pairwise relationships are expected between changes

in abundance, evenness and richness (Figs. 2g, h, i).

In contrast, even if numbers of individuals increase ($\Delta N > 0$), expected gains in species richness

can be constrained by decreased evenness. For example, opposing changes in total abundance

and evenness can potentially result in no change to species richness (Fig. 2d), and no relationship

between ΔS and ΔN (Fig. 2g). Or, if changes to the SAD are sufficiently strong, they can offset

any expected gains due to more individuals (Fig. 2e), and result in a negative relationship

between ΔS and ΔN (Fig. 2g). Alternatively, opposing changes to total numbers of individuals

and evenness could result in a positive relationship between ΔS and ΔN if, for example, the

effects of more individuals on species richness outweighs that of decreased evenness (Figure 2f).

These simplified scenarios illustrate the potential for interdependencies between component

changes. In particular, they show that the signs of changes in total abundance and evenness (i.e.,

ΔN and ΔS_{PIE}) can strongly influence the magnitude of richness changes, and whether expected

positive relationships between changes in abundance, evenness and richness are found. ΔN is

associated with the IBR curve stretching or contracting along the x-axis, and ΔS_{PIE} characterizes

changes that flex the curve up or down from the base along the y-axis (Olszewski 2004). When

ΔN and ΔS_{PIE} have the same sign, assemblages are expected to fall into the lower left and upper

294 right quadrants of Figs. 2g-i (i.e., component changes with the same sign and positive pairwise

295 relationships). In contrast, when ΔN and ΔS_{PIE} have different signs, they can have countervailing

296 effects that constrain richness changes, the strength of their pairwise relationships with ΔS will

297 be diminished and potentially reversed, and the likelihood of assemblages falling into the upper

298 left and lower right quadrants of Figs. 2g-i increases (i.e., opposing signs and negative pairwise

299 relationships).

301 **Empirical relationships among total abundance, evenness and richness**

302 Next, we evaluate empirical relationships by fitting models that allow for correlations between

303 component changes to data from 1125 assemblages. Our goal for the empirical analyses was to

304 examine relationships between changing components in temporal and spatial contexts across a

305 broad range of environmental conditions. We compiled data documenting either temporal or

306 spatial variation of assemblage composition in one of either naturally-varying or perturbed

307 environmental conditions. Temporal variation quantified rates of change (i.e., per year) for each

308 component for an assemblage at a single location through time. Analyses of spatial variation

309 quantified component changes between sites in different land use categories in perturbed

310 environments, or between random pairs of sites in the naturally-varying environment.

312 Based on our conceptual overview, we expect pairwise relationships between abundance,

313 evenness and richness changes to be generally positive. Changes in species richness are also

314 expected to be largest for assemblages where all pairwise relationships are positive. In contrast,

315 opposing changes in total abundance and evenness (i.e., ΔN and ΔS_{PIE} have different signs) are

316 expected to constrain changes in species richness. Additionally, if local diversity changes are

dominated by altered total abundances and species richness, strong positive relationships

between ΔS and ΔN, but weaker relationships between ΔS and ΔS_n, and ΔS and ΔS_{PIE} should

emerge across assemblages. Alternatively, strong relationships between either ΔS and ΔS_n and/or

ΔS and ΔS_{PIE}, accompanied by a weaker relationship between ΔS and ΔN, would indicate that

changes to the SAD are the dominant component of local assemblage change.

Temporal comparisons: natural environmental variation

Temporal changes in natural assemblages at the local scale were quantified using the BioTIME

database (Dornelas et al. 2018). Annual rates of change (change per year) for each metric were

estimated with models fit to data that documents over 45 thousand species in time series with an

average duration of 13 years. Taxonomic groups in our analysis came from surveys in marine,

freshwater and terrestrial ecosystems, and included plants (and other producers), invertebrates,

fish, amphibians, reptiles, birds, and mammals, as well as several surveys that collected data

from multiple taxa. Here, we only used time series that had numerical abundance data available

(i.e., studies that recorded counts of the number of individuals for each species in an

assemblage), and our analysis included 288 studies. Locations sampled in the BioTIME database

document places with varying degrees of anthropogenic environmental change, but do not

include manipulated assemblages or before-after-control-impact studies (Dornelas et al. 2018).

Accordingly, we contrast the environmental variation sampled by BioTIME with assemblage

time series that experienced documented perturbations (see *Temporal comparisons: experimental*

or natural perturbations).

To quantify changes at the local scale within BioTIME, studies with large extents were broken

up into smaller cell-level time series, while still maintaining the integrity of individual studies

(i.e., different studies were not combined, even when samples were collected in the same grid

cell). We used sample-based rarefaction (Gotelli & Colwell 2001) to standardize the number of

samples per year for each time series (see Blowes et al. 2019 for details). For the calculation of

rarefied richness (S_n), the minimum total number of individuals was determined for each time

series, and set as the target n for which expected richness was calculated; cell-level time series

where $n < 5$ were discarded. This process resulted in 42,604 cell-level time series from the 288

studies, and we focus on the study-level estimates of change in our results and discussion.

Temporal comparisons: experimental or natural perturbations

To complement the environmental variation sampled by the BioTIME database, we searched for

time series data with either experimental or natural perturbations. Specifically, we queried the

U.S. LTER network using the Data Portal of the Environmental Data Initiative

(https://portal.edirepository.org/nis/home.jsp) with the search terms 'experiment' and 'time' and

'abundance'. Records returned were checked to confirm that samples documented assemblages

of similar species collected with the same methodology, and following data standardization (i.e.,

minimum of five individuals per sample, and standardization of sample effort through time), our

analysis included 11 studies (see Appendix S1: *Temporal comparisons: experimental or natural*

perturbations for references), and annual rates of change (i.e., per year) were estimated for 63

study-treatment combinations; rates of change for all treatments (including controls) were

quantified in our analyses. Natural and experimental treatments included changes due to

warming, eutrophication, fire, grazing, restoration, severe storms or other disturbances, and kelp

removal. Taxonomic groups included algae, plants, invertebrates, fish, birds, and mammals.

Spatial comparisons: natural environmental variation

We combined two existing compilations of data to examine spatial patterns of biodiversity

change in relatively natural environmental contexts. The CESTES database (Jeliazkov et al.

2019) contains assemblage data from studies that sampled species at multiple sites (it also

includes information on traits and environment that we do not use here); we removed studies

with explicit human impacts identified as an environmental feature, and our analysis included 19

studies that sampled terrestrial, freshwater and marine assemblages from a number of taxonomic

groups (birds, plants, insects, macroinvertebrates, fishes and mammals). McGill (2011b)

compiled datasets with two or more local assemblages containing species abundance data; we

removed studies documenting disturbances and other perturbations, resulting in 32 studies being

retained. From the combined 51 studies, those with many sites were randomly subsampled down

to ten sites so that they did not dominate the results. Within each study, an arbitrary site was

assigned as the 'reference' site, and change was quantified between every site and the reference

within studies; our analysis included a total of 356 spatial comparisons.

Spatial comparisons: anthropogenic perturbations

To quantify spatial differences in biodiversity due to anthropogenic land use, we used the

PREDICTS database (Hudson et al. 2017). We used the 2016 release of the database

(downloaded from https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database on

10th July 2020). We limited our analyses to studies with abundance data for individuals, and

those with known land use categories (primary vegetation, mature secondary vegetation,

intermediate secondary vegetation, plantation forest, cropland, pasture, and urban); studies where

land use was not recorded were omitted. This resulted in 237 combinations of source ID and

study (some sources had multiple studies, denoted SS in the database), and 418 estimates of

change relative to the reference land use (primary vegetation) category.

Statistical models

To estimate changes in the different metrics whilst accounting for expected correlations between

them, we fit multivariate multilevel models to the data. Similar to the way univariate multilevel

(also called hierarchical or mixed-effects) models fit to a single response can allow varying (also

called random) intercepts and slopes to be correlated, this approach estimates changes in all

components simultaneously whilst allowing for (and estimating) correlations between them.

Response distributions for all metrics were chosen to ensure changes were estimated on similar

measurement scales, and because all metrics take only positive values, log response scales were

used for all components.

For the *Temporal comparisons: natural environmental variation* data, total abundance (N) was

fit with a model that assumed a lognormal distribution and identity link function, and Poisson

distributions with log link functions were fit to S_n, S_{PIE} and S; Poisson distributions were chosen

for S_n and S_{PIE} values rounded to integers based on visual assessments that showed lognormal

models fit to raw S_n and S_{PIE} values greatly underpredicted the density of ones in the data. For the

Temporal comparisons: experimental or natural perturbations data, S was no longer an integer

value after standardizing sampling effort, and all metrics were fit with models that assumed

lognormal distributions and identity link functions. Both spatial data sets were fit with models

that assumed lognormal distributions and identity link functions for total abundance (N), rarefied

richness (S_n) and evenness (S_{PIE}), and a Poisson distribution and log-link function for species

richness (S).

The *Temporal comparisons: natural environmental variation* models included non-varying

intercepts and slopes for year, and varying intercepts and slopes for studies and cells for all

responses. To allow for correlations between changes in the different responses, varying study-

and cell-level parameters for all responses were drawn from a single multivariate normal

distribution for each level (i.e., one for studies, one for cells; see Appendix S1: *Temporal*

comparisons: natural environmental variation for equations). The models fit to the *Temporal*

comparisons: experimental or natural perturbations data similarly included non-varying

intercepts and slopes for year, and had varying intercepts for study, site and block fitted

separately for each response. For these data, correlations between changes in the different

responses were modeled by drawing varying intercepts and slopes for each combination of

treatment and study for all responses from a single multivariate normal distribution (see

Appendix S1: *Temporal comparisons: experimental or natural perturbations* for equations).

The models fit to the *Spatial comparisons: natural environmental variation* data included non-

varying intercepts for data source (i.e., CESTES and McGill). Correlations between the different

responses were modeled by assuming varying intercepts and slopes (representing the reference

site and departures for all other sites from the reference, respectively) for each study and

response came from a single multivariate normal distribution; over-dispersion in the richness

response was modeled using an observation-level varying intercept (see Appendix S1: *Spatial*

comparisons: natural environmental variation for equations). Models fit to the *Spatial*

comparisons: anthropogenic perturbations data included non-varying intercepts and slopes

(representing the reference [primary vegetation] category and departures from the reference for

each land use category, respectively), and varying intercepts for sites and blocks were modeled

separately for each response. Correlations between changes in the different responses were

modeled by assuming that varying intercepts and slopes (as per the non-varying intercepts and

slopes) for each combination of source and study and each response came from a single

multivariate normal distribution (see Appendix S1: *Spatial comparisons: anthropogenic*

perturbations for equations).

All statistical models were estimated using the Hamiltonian Monte Carlo (HMC) sampler Stan

(Carpenter et al. 2017), and coded using the 'brms' package (Burkner 2017). Details of all model

specifications, and the iterations and warmup periods are provided in the Appendix, as are the

priors (which were weakly regularizing). Visual inspection of the HMC chains and model

diagnostics (Rhat < 1.05) showed good mixing of chains and convergence, and model adequacy

assessed visually using posterior predictive checks showed that the models were able to make

predictions similar to the empirical data (see Appendix Figure: S1-4). Code for all analyses is

available at https://github.com/sablowes/MulticomponentBioChange, and will be archived

following publication.

Results

Temporal changes in perturbed environments had the highest percentage of assemblages with at

least one component trend (ΔN, ΔS_{n}, ΔS_{PIE}, or ΔS) that differed from zero (44%), followed by

spatial comparisons across land use categories (29%), temporal changes (21%) and spatial

comparisons in naturally varying environments (12%). Component changes that differed from

zero showed broadly similar patterns across datasets, with one exception: trends differing from

zero for multiple components were less common for spatial comparisons between assemblages in

naturally varying environments (Appendix S1: Figure S5).

Pairwise relationships between changing components were typically positive (i.e., had the same

sign), though exceptions to this general pattern were found for all data sources (Figure 3). For

assemblages where ΔN and ΔS_{PIE} had the same sign (though not necessarily differing statistically

from zero), richness changes were typically larger in magnitude (Figure 3). In contrast,

assemblages where ΔN and ΔS_{PIE} had opposing signs typically exhibited changes in richness that

were smaller in magnitude (Figure 3). This tendency for countervailing changes in abundance

and evenness to constrain richness changes was most apparent for spatial comparisons between

different land use categories (Figure 3j-l), and there was a high proportion of assemblages that

were growing in size ($\Delta N > 0$) but with declining species richness ($\Delta S < 0$; Figure 3j), associated

with declining evenness ($\Delta S_{PIE} < 0$).

The strongest relationships were found for components changing through time, and relationships

between richness and changes in the SAD -- rarefied richness (Figs. 3b, e) and evenness (Figs.

3c, f; Figs. 4a, b) -- were stronger than those between changes in richness and total abundance

(Figs. 3a, c; Figs. 4a, b). Spatial comparisons had generally weak relationships overall. No strong

relationships between changing components emerged for comparisons in natural environments

(Figs. 3g-i; Fig. 4c), and only weakly positive relationships between changes in abundance,

evenness and richness were found in comparisons between primary vegetation and different land

use categories (Figure 3j-l, Figure 4d).

Temporal changes in naturally varying assemblages were roughly centered on zero for all

metrics (Figure 3a-c). Across assemblages, altered numbers of individuals and species richness

changes had a moderately positive relationship (Figure 4a), weakened predominantly by

assemblages that had opposing abundance and evenness relationships (Figure 3a). In contrast,

relationships between changes in species richness and rarefied richness, and between richness

changes and evenness changes were strong (Figure 4a). Assemblages in perturbed environments

had slightly positive temporal trends on average in all components (Figure 3d-f). Across

assemblages, ΔS and ΔN (Figure 3d, 4b) and ΔS and ΔS_{PIE} (Figure 3f, 4b) had relatively weak

positive relationships, whereas ΔS and ΔS_n (Figure 3e, 4b) showed a strong positive relationship.

Spatial comparisons in naturally varying environments exhibited highly heterogeneous patterns

of change centered around zero for all metrics (Figure 3g-i). Decoupled component changes

meant that relationships between them were generally absent or weak across assemblages (Figure

4c). Spatial comparisons between assemblages in primary vegetation versus those in different

land use categories were also highly heterogeneous, though there were typically fewer

individuals, less even assemblages and fewer species relative to primary vegetation (Figure 3j-l).

Across assemblages, land use change was typically associated with relatively weak positive

relationships between changes in the components of local diversity (Figure 4d).

Discussion

Our conceptual overview using individual-based rarefaction curves clearly shows how the expectation of positive pairwise relationships between changes in abundance, evenness and richness arises. If curves stretch or contract, we expect positive relationships between changes in total abundance and richness. Similarly, if curves flex upwards or downwards, positive relationships between changes in evenness and richness are expected. Rarefaction curves also show how contrasting signs of changes in abundance and evenness can strongly determine the magnitude of richness changes, and determine whether positive relationships between changes in richness and the other components (abundance and evenness) are likely. Both these predictions were generally well supported by our empirical analyses. Relationships between changes in abundance, evenness and richness were generally positive, and richness changes were typically greater for assemblages with strictly positive pairwise relationships. Countervailing changes in total abundance and evenness, where found, often constrained the magnitude of changes in species richness, and acted to weaken relationships between ΔN and ΔS, and ΔS_{PIE} and ΔS. Spatial comparisons had the most heterogeneous relationships between changes in abundance, evenness, and richness, and in relatively natural environments changes were sufficiently decoupled that no strong relationships emerged across assemblages. In contrast, strong positive correlations between temporal changes in species richness (ΔS) and changes in metrics associated with altered SADs (ΔS_{n}, ΔS_{PIE}) emerged across assemblages. These temporal results show strong support for the prediction that variation in relative abundances can dominate local variation in biodiversity (Chapin et al. 2000), even when human impacts are less direct.

Variation in assemblage size does not dominate local diversity change

Overall, only ~2% of assemblages in this study (22 of 1125) had changes consistent with a

strong 'sampling' effect on changes in species richness (i.e., ΔN & ΔS having the same sign, and

being the only changes different from zero). This finding complements existing evidence

showing that despite many tests, empirical evidence for the more-individuals hypothesis

(Srivastava & Lawton 1998) remains equivocal (Storch et al 2018, Vagle & McCain 2020).

While both (species-level) population variability and variation associated with sampling (Vagle

& McCain 2020) likely contribute to the weak response of species richness to variation in the

total number of individuals, our results are broadly consistent with previous syntheses showing

that broad-scale spatial variation in richness was rarely driven simply by variation in the numbers

of individuals (Currie et al. 2004, Storch et al 2018). Our results indicate that variation in local

diversity, through time or from site-to-site, is not due to changes in assemblage size passively

sampling more or less from a static SAD. Instead, we show that variation in local biodiversity

can be strongly influenced by changes to species' relative abundances. These changes can be

occurring at multiple scales (Hillebrand et al. 2008, Blowes et al. 2020), and could reflect altered

local environmental conditions (e.g., altered resource or habitat availability and diversity,

eutrophication, local harvest or exploitation), or changes at broader scales that alter the species

pool (via species additions or subtractions).

Our general result showing that variation in the total abundance of an assemblage through time

or space is often decoupled from changes in metrics of biodiversity such as species richness also

cautions against making "apples to oranges" comparisons in the context of quantifying

biodiversity change. For example, some estimates of change are based on either population-level

abundance (e.g., Living Planet Index, WWF 2020), or assemblage-level abundance (e.g., insect

declines; Wagner 2000, van Klink et al. 2020), whereas other change estimates are based on

patterns of the number or identity of species present (e.g., Dornelas et al. 2014, Newbold et al.

2015). Our results show that assuming abundance and richness changes are strongly correlated

will often be an oversimplification. Moreover, the importance of altered relative abundances for

local biodiversity variation means that biodiversity change estimates will frequently depend on

whether changes in species relative abundances influence the metrics used (see e.g., Antão et al.

2020b).

Contrasting component relationships between temporal changes versus spatial comparisons of

biodiversity

Relationships between changing components of biodiversity showed strikingly different patterns

between temporal changes and spatial comparisons. Moreover, these differences were generally

greater than those found between naturally-varying and perturbed assemblages, for either

temporal changes or spatial comparisons.

Pairwise relationships between changes in abundance, evenness and richness were typically

weak for spatial comparisons. Decoupling was greatest, and pairwise relationships weakest, for

changes between sites experiencing relatively natural environmental variation. However, given

our simple conceptual framework shows that some degree of interdependence cannot be avoided,

we caution against overinterpreting the relative independence of these component changes, and

further analyses examining component change relationships along continuous spatial gradients

are warranted. Indeed, evenness and richness are never numerically independent (Jost 2010), and

the weak overall relationship between changes in richness and evenness for these data was in

part due to assemblages with countervailing changes in abundance and evenness. Most

importantly, these highly variable component changes further emphasize the need for a holistic

approach to quantifying biodiversity change (Chase et al. 2018, Avolio et al. 2021).

Our prediction that the signs of changes in abundance and evenness can strongly determine the

magnitude of richness changes was most evident in the spatial contrasts between primary

vegetation and other land use categories (Newbold et al. 2015, 2020). Assemblages with the

greatest declines in abundance and evenness had the greatest richness declines. In contrast, when

abundance and evenness changes had opposing signs, richness changes were tempered. Indeed,

countervailing abundance and evenness changes were frequently associated with components

other than species richness (i.e., ΔN, ΔS_n, and/or ΔS_{PIE}) having a trend that differed from zero

across all data sources (Appendix S1: Table S1). This highlights that even apparently decoupled

or weakly correlated component changes have interdependencies that can remain important

determinants of observed changes.

In contrast to assemblage change between sites, there was strong coupling between species

richness and SAD changes through time. In particular, the strength of the relationship between

ΔS_n and ΔS resulted in estimates of change being similar for most assemblage time series in

relatively natural environments (Figure 3b). In some cases, this occurred despite countervailing

changes in total abundance and evenness (Figure 3a, b). For assemblages where abundance and

evenness changed in the same direction, similar estimates of ΔS_n and ΔS indicate that abundance

changes were occurring along a relatively flat region of the individual-based rarefaction curve.

This shows that changes to the total number of individuals need not strongly influence species

richness, even where signs are the same and they have a positive relationship. The strong

association between richness changes and altered relative abundances has important implications

for examining causes and/or consequences of biodiversity change (Hillebrand et al. 2008,

Crowder et al. 2010). Even where the expected positive relationships between abundance,

evenness and richness are found, we can more fully understand assemblage changes when all

component changes are examined simultaneously.

While both approaches, time series and spatial comparisons (or space-for-time substitutions),

have contributed to our understanding of biodiversity change, the relative merits of each for our

understanding of ecological dynamics has not been discussed much (Adler et al. 2020). The

largely decoupled component changes found here for spatial comparisons suggest that too much

focus on average changes across assemblages, such as those in total abundance or in species

richness, risks masking highly heterogeneous changes occurring within assemblages in multiple

components. Moreover, decoupled, heterogeneous component changes complicate using spatial

comparisons to infer temporal changes. The smaller effect sizes found here for time series

indicates greater inertia compared to site-to-site variation. More generally, the strong role of

changes to the SAD for variation in local biodiversity suggests that deepening our understanding

of altered patterns of relative abundance across scales represents an important direction for future

theoretical and empirical work. Here our focus has been on numerical relationships between

component changes, and using process-based models (e.g., Thompson et al. 2020) to examine

how altered metacommunity dynamics impact patterns of relative abundance across scales could

help our understanding how different processes impact component relationships. Similarly,

empirical studies could ask whether local environmental changes are affecting evenness, or if

changes occurring at broader spatial scales are impacting the size of species pool and the

regional SAD?

Conclusions

We found strong correlations between changes in the SAD and species richness changes through

time, whereas relationships between abundance and richness changes for both temporal and

spatial diversity variation were generally weak. Our findings confirm that altered species relative

abundances, and/or changes to the size of the species pool, often strongly influence local

diversity change (Chapin et al. 2010), even where human impacts are less direct. However, our

results also reinforce cautions against examining changes to any one component of biodiversity

change in isolation (e.g., Wilsey et al. 2005, Chase et al. 2018, Avolio et al. 2021).

To be most useful, quantifying (co)variation in the different components of biodiversity needs to

be done coherently. Individual-based rarefaction curves and associated metrics can provide an

intuitive and illustrative characterization of relationships among changing components of local

biodiversity. Whilst ecologists are increasingly looking beyond species richness to quantify

biodiversity change (e.g., Dornelas et al. 2014, Hillebrand et al. 2018), different components of

biodiversity and its change within assemblages are most often analyzed independently, and

frequently with metrics lacking conceptual unity. Conceptually and empirically, our results

emphasize that changes to the most frequently quantified aspects of biodiversity, including

changes to the numbers of individuals, and the relative abundance and total number of species

are highly interdependent. Examining how within-assemblage component changes covary with

potential drivers could reveal insights masked by independent aggregate estimates of change

across assemblages, and provide new information for understanding biodiversity change in the

Anthropocene.

Acknowledgements

SAB thanks the Biodiversity Synthesis group at iDiv for contructive feedback at various stages

of the project. SAB, TE, AS, and JMC were supported by the German Centre for Integrative

Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation

(FZT 118).

References

Antão, L.H., Bates, A.E., Blowes, S.A., Waldock, C., Supp, S.R., Magurran, A.E., Dornelas, M.,

Schipper, A.M., 2020a. Temperature-related biodiversity change across temperate marine and

terrestrial systems. Nature ecology & evolution 4, 927–933.

Antão, L.H., Pöyry, J., Leinonen, R., Roslin, T., 2020b. Contrasting latitudinal patterns in

diversity and stability in a high-latitude species-rich moth community. Global Ecology and

Biogeography 29, 896–907.

Avolio, M.L., Komatsu, K.J., Collins, S.L., Grman, E., Koerner, S.E., Tredennick, A.T., Wilcox,

K.R., Baer, S., Boughton, E.H., Britton, A.J., 2021. Determinants of community

compositional change are equally affected by global change. Ecology Letters 24, 1892–1904.

Blowes, S.A., Chase, J.M., Di Franco, A., Frid, O., Gotelli, N.J., Guidetti, P., Knight, T.M., May,

F., McGlinn, D.J., Micheli, F., 2020. Mediterranean marine protected areas have higher

biodiversity via increased evenness, not abundance. Journal of Applied Ecology 57, 578–589.

Blowes, S.A., Supp, S.R., Antão, L.H., Bates, A., Bruelheide, H., Chase, J.M., Moyes, F.,

Magurran, A., McGill, B., Myers-Smith, I.H., Winter, M., Bjorkman, A.D., Bowler, D.E.,

Byrnes, J.E., Gonzalez, A., Hines, J., Isbell, F., Jones, H.P., Navarro, L.M., Thompson, P.L.,

Vellend, M., Waldock, C., Dornelas, M., 2019. The geography of biodiversity change in

marine and terrestrial assemblages. Science 366, 339–345.

Bolker, B.M., 2008. Ecological models and data in R. Princeton University Press.

Bürkner, P.-C., 2017. brms: An R package for Bayesian multilevel models using Stan. Journal of

Statistical Software 80, 1–28.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker,

M., Guo, J., Li, P., Riddell, A., 2017. Stan: A probabilistic programming language. Journal of

Statistical Software 76.

Caswell, H., 1976. Community structure: a neutral model analysis. Ecological monographs 46,

327–354.

Cayuela, L., Gotelli, N.J., Colwell, R.K., 2015. Ecological and biogeographic null hypotheses for

comparing rarefaction curves. Ecological Monographs 85, 437–455.

Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., Ellison, A.M., n.d.

Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in

species diversity studies. Ecology 84, 45–67.

Chao, A., Jost, L., 2012. Coverage-based rarefaction and extrapolation: standardizing samples by

completeness rather than size. Ecology 93, 2533–2547.

Chapin Iii, F.S., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L.,

Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., 2000. Consequences of changing

biodiversity. Nature 405, 234–242.

Chase, J.M., Blowes, S.A., Knight, T.M., Gerstner, K., May, F., 2020. Ecosystem decay

exacerbates biodiversity loss with habitat loss. Nature 584, 238–243.

Chase, J.M., Knight, T.M., 2013. Scale-dependent effect sizes of ecological drivers on

biodiversity: why standardised sampling is not enough. Ecol Lett 16, 17–26.

Chase, J.M., McGill, B.J., McGlinn, D.J., May, F., Blowes, S.A., Xiao, X., Knight, T.M.,

Purschke, O., Gotelli, N.J., 2018. Embracing scale-dependence to achieve a deeper

understanding of biodiversity and its change across communities. Ecol Lett 21, 1737–1751.

Chase, J.M., McGill, B.J., Thompson, P.L., Antão, L.H., Bates, A.E., Blowes, S.A., Dornelas,

M., Gonzalez, A., Magurran, A.E., Supp, S.R., Winter, M., Bjorkman, A.D., Bruelheide, H.,

Byrnes, J.E., Sarmento Cabral, J., Elahi, R., Gomez, C., Guzman, H.M., Isbell, F., Myers-

Smith, I.H., Jones, H.P., Hines, J., Vellend, M., Waldock, C., O'Connor, M., 2019. Species

richness change across spatial scales. Oikos 1–13.

Coleman, B.D., Mares, M.A., Willig, M.R., Hsieh, Y.-H., 1982. Randomness, area, and species

richness. Ecology 63, 1121–1133.

Connell, J.H., 1978. Diversity in tropical rain forests and coral reefs. Science 199, 1302–1310.

Connor, E.F., McCoy, E.D., 1979. The statistics and biology of the species-area relationship. The

American Naturalist 113, 791–833.

Crowder, D.W., Northfield, T.D., Strand, M.R., Snyder, W.E., 2010. Organic agriculture

promotes evenness and natural pest control. Nature 466, 109–112.

Currie, D.J., 1991. Energy and large-scale patterns of animal-and plant-species richness. Am Nat

27–49.

Currie, D.J., Mittelbach, G.G., Cornell, H.V., Field, R., Guégan, J.-F., Hawkins, B.A., Kaufman,

D.M., Kerr, J.T., Oberdorff, T., O'Brien, E., 2004. Predictions and tests of climate-based

hypotheses of broad-scale variation in taxonomic richness. Ecology letters 7, 1121–1134.

Daskalova, G.N., Myers-Smith, I.H., Bjorkman, A.D., Blowes, S.A., Supp, S.R., Magurran,

A.E., Dornelas, M., 2020a. Landscape-scale forest loss as a catalyst of population and

biodiversity change. Science 368, 1341–1347.

Daskalova, G.N., Myers-Smith, I.H., Godlee, J.L., 2020b. Rare and common vertebrates span a

wide spectrum of population trends. Nature Communications 4394.

Díaz, S., Settele, J., Brondízio, E.S., Ngo, H.T., Agard, J., Arneth, A., Balvanera, P., Brauman,

K.A., Butchart, S.H., Chan, K.M., 2019. Pervasive human-driven decline of life on Earth

points to the need for transformative change. Science 366.

Dornelas, M., Antao, L.H., Moyes, F., Bates, A.E., Magurran, A.E., Adam, D., Akhmetzhanova,

A.A., Appeltans, W., Arcos, J.M., Arnold, H., 2018. BioTIME: A database of biodiversity

time series for the Anthropocene. Glob Ecol Biogeogr 27, 760–786.

Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., Magurran, A.E.,

2014. Assemblage time series reveal biodiversity change but not systematic loss. Science 344,

296–299.

Dornelas, M., Gotelli, N.J., Shimadzu, H., Moyes, F., Magurran, A.E., McGill, B.J., 2019. A

balance of winners and losers in the Anthropocene. Ecol Lett 22, 847–854.

Fisher, R.A., Corbet, A.S., Williams, C.B., 1943. The relation between the number of species

and the number of individuals in a random sample of an animal population. The Journal of

Animal Ecology 42–58.

Gaston, K.J., 2000. Global patterns in biodiversity. Nature 405, 220–227.

726 Gotelli, N.J., Colwell, R.K., 2001. Quantifying biodiversity: procedures and pitfalls in the

727  measurement and comparison of species richness. Ecol Lett 4, 379–391.

728 He, F., Legendre, P., 2002. Species diversity patterns derived from species–area models. Ecology

729  83, 1185–1198.

730 Hill, M.O., 1973. Diversity and evenness: a unifying notation and its consequences. Ecology 54,

731  427–432.

732 Hillebrand, H., Bennett, D.M., Cadotte, M.W., 2008. Consequences of dominance: a review of

733  evenness effects on local and regional ecosystem processes. Ecology 89, 1510–1520.

734 Hillebrand, H., Blasius, B., Borer, E.T., Chase, J.M., Downing, J.A., Eriksson, B.K., Filstrup,

735  C.T., Harpole, W.S., Hodapp, D., Larsen, S., 2018. Biodiversity change is uncoupled from

736  species richness trends: Consequences for conservation and monitoring. Journal of Applied

737  Ecology 55, 169–184.

738 Hudson, L.N., Newbold, T., Contu, S., Hill, S.L., Lysenko, I., De Palma, A., Phillips, H.R.,

739  Alhusseini, T.I., Bedford, F.E., Bennett, D.J., 2017. The database of the Predicts (Projecting

740  responses of ecological diversity in changing terrestrial systems) project. Ecology and

741  Evolution 7, 145–188.

742 Hughes, A.R., Byrnes, J.E., Kimbro, D.L., Stachowicz, J.J., 2007. Reciprocal relationships and

743  potential feedbacks between biodiversity and disturbance. Ecology letters 10, 849–864.

744 Hurlbert, S.H., 1971. The nonconcept of species diversity: a critique and alternative parameters.

745  Ecology 52, 577–586.

746 Jeliazkov, A., Mijatovic, D., Chantepie, S., Andrew, N., Arlettaz, R., Barbaro, L., Barsoum, N.,

747  Bartonova, A., Belskaya, E., Bonada, N., 2020. A global database for metacommunity

748  ecology, integrating species, traits, environment and space. Scientific data 7, 1–15.

Jones, F.A., Magurran, A.E., 2018. Dominance structure of assemblages is regulated over a

period of rapid environmental change. Biology letters 14, 20180187.

Jost, L., 2007. Partitioning diversity into independent alpha and beta components. Ecology 88,

2427–2439.

Jost, L., 2006. Entropy and diversity. Oikos 113, 363–375.

Kreft, H., Jetz, W., Mutke, J., Kier, G., Barthlott, W., 2008. Global diversity of island floras from

a macroecological perspective. Ecology letters 11, 116–127.

Latham, R.E., Ricklefs, R.E., 1993. Global patterns of tree species richness in moist forests:

energy-diversity theory does not account for variation in species richness. Oikos 325–333.

Leung, B., Hargreaves, A.L., Greenberg, D.A., McGill, B., Dornelas, M., Freeman, R., 2020.

Clustered versus catastrophic global vertebrate declines. Nature 588, 267–271.

MacArthur, R.H., 1965. Patterns of species diversity. Biological reviews 40, 510–533.

Magurran, A.E., McGill, B.J., 2011. Biological diversity: frontiers in measurement and

assessment. Oxford University Press.

May, F., Gerstner, K., McGlinn, D.J., Xiao, X., Chase, J.M., 2018. mobsim: An R package for

the simulation and measurement of biodiversity across spatial scales. Methods in Ecology and

Evolution 9, 1401–1408.

McGill, B.J., 2011a. Linking biodiversity patterns by autocorrelated random sampling. Am J Bot

98, 481–502.

McGill, B.J., 2011b. Species abundance distributions, in: Biological Diversity: Frontiers In

Measurement & Assessment (Eds Magurran, A. & McGill, B.). Oxford University Press, pp.

105–122.

McGill, B.J., Etienne, R.S., Gray, J.S., Alonso, D., Anderson, M.J., Benecha, H.K., Dornelas,

M., Enquist, B.J., Green, J.L., He, F., Hurlbert, A.H., Magurran, A.E., Marquet, P.A., Maurer,

B.A., Ostling, A., Soykan, C.U., Ugland, K.I., White, E.P., 2007. Species abundance

distributions: moving beyond single prediction theories to integration within an ecological

framework. Ecol Lett 10, 995–1015.

McGlinn, D.J., Xiao, X., May, F., Gotelli, N.J., Engel, T., Blowes, S.A., Knight, T.M., Purschke,

O., Chase, J.M., McGill, B.J., 2019. Measurement of Biodiversity (MoB): A method to

separate the scale-dependent effects of species abundance distribution, density, and

aggregation on diversity change. Methods in Ecology and Evolution 10, 258–269.

McKinney, M.L., Lockwood, J.L., 1999. Biotic homogenization: a few winners replacing many

losers in the next mass extinction. Trends in ecology & evolution 14, 450–453.

Menge, B.A., Berlow, E.L., Blanchette, C.A., Navarrete, S.A., Yamada, S.B., 1994. The

keystone species concept: variation in interaction strength in a rocky intertidal habitat.

Ecological monographs 64, 249–286.

Miller, A.D., Roxburgh, S.H., Shea, K., 2011. How frequency and intensity shape diversity–

disturbance relationships. Proceedings of the National Academy of sciences 108, 5643–5648.

Mittelbach, G.G., Steiner, C.F., Scheiner, S.M., Gross, K.L., Reynolds, H.L., Waide, R.B.,

Willig, M.R., Dodson, S.I., Gough, L., 2001. What is the observed relationship between

species richness and productivity? Ecology 82, 2381–2396.

Newbold, T., Hudson, L.N., Contu, S., Hill, S.L., Beck, J., Liu, Y., Meyer, C., Phillips, H.R.,

Scharlemann, J.P., Purvis, A., 2018. Widespread winners and narrow-ranged losers: Land use

homogenizes biodiversity in local assemblages worldwide. PLoS biology 16, e2006841.

Newbold, T., Hudson, L.N., Hill, S.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett,

D.J., Choimes, A., Collen, B., 2015. Global effects of land use on local terrestrial biodiversity.

Nature 520, 45–50.

Newbold, T., Oppenheimer, P., Etard, A., Williams, J.J., 2020. Tropical and Mediterranean

biodiversity is disproportionately sensitive to land-use and climate change. Nature Ecology &

Evolution 4, 1630–1638.

Olszewski, T.D., 2004. A unified mathematical framework for the measurement of richness and

evenness within and among multiple communities. Oikos 104, 377–387.

Paine, R.T., 1974. Intertidal community structure. Oecologia 15, 93–120.

Palmer, M.W., Clark, D.B., Clark, D.A., 2000. Is the number of tree species in small tropical

forest plots nonrandom? Community Ecology 1, 95–101.

Preston, F.W., 1962. The canonical distribution of commonness and rarity: Part I. Ecology 43,

185–215.

Rosenberg, K.V., Dokter, A.M., Blancher, P.J., Sauer, J.R., Smith, A.C., Smith, P.A., Stanton,

J.C., Panjabi, A., Helft, L., Parr, M., 2019. Decline of the North American avifauna. Science

366, 120–124.

Rosenzweig, M.L., 1995. Species diversity in space and time. Cambridge University Press.

Scheiner, S.M., Willig, M.R., 2005. Developing unified theories in ecology as exemplified with

diversity gradients. The American Naturalist 166, 458–469.

Shmida, A.V.I., Wilson, M.V., 1985. Biological determinants of species diversity. Journal of

biogeography 1–20.

Soininen, J., Passy, S., Hillebrand, H., 2012. The relationship between species richness and

evenness: a meta-analysis of studies across aquatic ecosystems. Oecologia 169, 803–809.

Srivastava, D.S., Lawton, J.H., 1998. Why more productive sites have more species: an

experimental test of theory using tree-hole communities. The American Naturalist 152, 510–

529.

Stirling, G., Wilsey, B., 2001. Empirical relationships between species richness, evenness, and

proportional diversity. The American Naturalist 158, 286–299.

Storch, D., Bohdalková, E., Okie, J., 2018. The more-individuals hypothesis revisited: the role of

community abundance in species richness regulation and the productivity–diversity

relationship. Ecol Lett 21, 920–937.

Tilman, D., 1982. Resource competition and community structure. Princeton University Press.

Vagle, G.L., McCain, C.M., n.d. Natural population variability may be masking the more-

individuals hypothesis. Ecology 101, e03035.

Van Klink, R., Bowler, D.E., Gongalsky, K.B., Swengel, A.B., Gentile, A., Chase, J.M., 2020.

Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances.

Science 368, 417–420.

Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U.,

Sun, Y., Pyšek, P., 2011. Ecological impacts of invasive alien plants: a meta-analysis of their

effects on species, communities and ecosystems. Ecology letters 14, 702–708.

Wagner, D.L., 2020. Insect declines in the Anthropocene. Annual review of entomology 65,

457–480.

Whittaker, R.J., Willis, K.J., Field, R., 2001. Scale and species richness: towards a general,

hierarchical theory of species diversity. Journal of biogeography 28, 453–470.

Wright, D.H., 1983. Species-energy theory: an extension of species-area theory. Oikos 496–506.

WWF, 2020. Living Planet Report 2020-Bending the curve of biodiversity loss. World Wildlife

Fund, Gland, Switzerland.

Zhang, Y., Chen, H.Y., Reich, P.B., 2012. Forest productivity increases with evenness, species

richness and trait variation: a global meta-analysis. Journal of ecology 100, 742–749.

843 **Figure 1**: (a) Individual-based rarefaction (IBR) curves for two hypothetical assemblages,

844 showing the four components we use to quantify change (N, S, S_n, PIE). We visualize

845 relationships between these four components of the IBR curve by plotting: (b) changes in species

846 richness (ΔS) as a function of altered numbers of individuals (ΔN), (c) changes in species

847 richness (ΔS) as a function of changes in rarefied richness (ΔS_n), and (d) changes in species

848 richness (ΔS) as a function of the numbers equivalent conversion of the Probability of

849 Interspecific Encounter (ΔS_{PIE}). We show ΔPIE on the figure to illustrate changes of the PIE (it is

850 equivalent to the slope at the base of the curve) with the IBR, but use the numbers equivalent

851 transformation (ΔS_{PIE}) in all analyses. Points on panels (b, c, and d) show changes between the

852 two hypothetical assemblages, with the reference assemblage depicted by the grey line.

854 **Figure 2**: Conceptual illustrations of potential pathways of local assemblage diversity change

855 and corresponding relationships between component changes. Starting from a reference

856 assemblage (depicted with grey rarefaction curves), diversity change can be due to (a) more

857 individuals only, (b) changes to the species abundance distribution only (e.g., increased species

858 pool size or increased evenness), or (c) changes in total abundance and the SAD that result in

859 positive pairwise relationships between ΔN, ΔS_n, ΔS_{PIE}, and ΔS. However, if the signs of ΔN and

860 ΔS_{PIE} differ, their relationships with ΔS weaken and species richness can (d) remain static, (e)

861 decrease or (f) increase. We visualize pairwise relationships between component changes for

862 each scenario (i.e., the different shaped symbols) using: (g) changes in species richness as a

863 function of changes to the number of individuals, (h) changes in species richness as a function of

864 changes in rarefied richness, and (i) changes in species richness as a function of changes in

865 evenness.

Figure 3: Empirical relationships between four components of local diversity change. Change in

species richness as a function of changes in the numbers of individuals (left column), rarefied

richness (middle column), and evenness (right column) for (a-c) study-level estimates of

temporal changes in naturally varying environments; (d-f) estimates of temporal change for

combinations of study and treatment in perturbed environments; (g-i) estimates of spatial

changes within studies from an arbitrary reference site along natural environmental gradients;

and, (j-l) estimates of spatial change within studies between primary vegetation and different

land use categories. Colored concentration ellipses show the confidence interval (5 and 95%) of

the posterior distributions. Dotted grey lines are x and y = 0, and x = y. See Appendix S1: Figure

S6 for remaining pairwise relationships. Scale of x- and y-axes vary between panels; one

estimate with $\Delta\log(N) = -1.79$, $\Delta\log(S) = -3.77$, $\Delta\log(S_n) = -3.23$, $\Delta\log(S_{PIE}) = -3.21$, removed

from (j-l) for clarity.

Figure 4: Component correlations among studies within each data source. Density plots for the

posterior distribution of pairwise correlations between component changes for (a) temporal

comparison in naturally varying environments, (b) temporal comparisons in perturbed

environments, (c) spatial comparisons along natural gradients, and (d) spatial comparisons

between different land use categories. Correlations estimated separately for sites and land use

categories relative to the references were combined on (c) and (d).

889   Figure 2

891   Figure 3

894   Figure 4

895
896

Appendix S1
898

899   **Title**: Local biodiversity change reflects interactions among changing abundance, evenness and
900   richness
901

902   **Authors**: Shane A. Blowes, Gergana N. Daskalova, Maria Dornelas, Thore Engel, Nicholas J.
903   Gotelli, Anne E. Magurran, Inês S. Martins, Brian McGill, Daniel J. McGlinn, Alban Sagouis
904   Hideyasu Shimadzu, Sarah R. Supp, Jonathan M. Chase
905

906   **Journal**: Ecology
907

909   **Table S1:** Countervailing changes in abundance (ΔN) and evenness (ΔS_{PIE}) often result in
910   components other than richness changing. Component(s) with high probability of change as per
911   Figure 2 (i.e., 90% CI does not overlap zero); each row shows the number of assemblages with
912   either countervailing [sign(ΔN) x sign(ΔS_{PIE}) = -1] or abundance and evenness changes with the
913   same direction [sign(ΔN) x sign(ΔS_{PIE}) = 1] for the different data sources.
914

Component with high probability of change	sign(ΔN) x sign(ΔS_{PIE})	Temporal change: natural variation — Number of assemblages	Temporal change: perturbed environments — Number of assemblages	Spatial change: natural variation — Number of assemblages	Spatial change: land use change — Number of assemblages
None	-1	97	23	154	107
None	1	131	12	214	191
N only	-1	7	0	7	5
N only	1	2	1	3	5
S_{PIE} only	-1	6	0	1	10
S_{PIE} only	1	2	0	2	12
S_n only	-1	0	1	7	1
S_n only	1	1	1	3	0
S only	-1	1	0	2	1
S only	1	1	0	5	6
S & S_n	-1	0	0	1	1

S & S_n	1	6	2	3	6
S & S_{PIE}	-1	0	0	0	0
S & S_{PIE}	1	1	0	1	3
S & N	-1	1	3	0	3
S & N	1	0	1	3	10
S_n & S_{PIE}	-1	3	4	2	9
S_n & S_{PIE}	1	0	0	0	1
S_n & N	-1	0	0	1	1
S_n & N	1	0	1	0	0
N & S_{PIE}	-1	4	0	0	3
N & S_{PIE}	1	0	0	0	0
S_n, N & S_{PIE}	-1	3	4	0	6
S_n, N & S_{PIE}	1	0	0	0	0
S, N & S_{PIE}	-1	2	0	0	0
S, N & S_{PIE}	1	1	0	0	4
S, S_n, N & S_{PIE}	-1	7	4	0	2
S, S_n, N & S_{PIE}	1	12	6	1	31

 Temporal comparisons: natural environmental variation

922   We fit a model that assumed a lognormal distribution for total abundance (*N*), and
923 poisson distributions (with log link functions) for species richness (S), rarefied richness (*S*n) and
924 evenness (*S*PIE). We ran the model with four chains for 2000 iterations, with 1000 used as
925 warmup. The model took the following form:

926
$$N_{ijt} \sim lognormal\left(\mu_{ijt}^{N}, \sigma\right),$$

927
$$S_{ijt} \sim pois\left(\lambda_{ijt}^{S}\right),$$

928
$$S_{n_{ijt}} \sim pois\left(\lambda_{ijt}^{S_n}\right),$$

929
$$S_{PIE\,ijt} \sim pois\left(\lambda_{ijt}^{S_{PIE}}\right),$$

930
$$\mu_{ijt}^{N} = \beta_0^{N} + \beta_{0i}^{N} + \beta_{0ij}^{N} + \left(\beta_1^{N} + \beta_{1i}^{N} + \beta_{1ij}^{N}\right)year_{ijt},$$

931
$$\log\left(\lambda_{ijt}^{S}\right) = \beta_0^{S} + \beta_{0i}^{S} + \beta_{0ij}^{S} + \left(\beta_1^{S} + \beta_{1i}^{S} + \beta_{1ij}^{S}\right)year_{ijt},$$

932
$$\log\left(\lambda_{ijt}^{S_n}\right) = \beta_0^{S_n} + \beta_{0i}^{S_n} + \beta_{0ij}^{S_n} + \left(\beta_1^{S_n} + \beta_{1i}^{S_n} + \beta_{1ij}^{S_n}\right)year_{ijt},$$

933
$$\log\left(\lambda_{ijt}^{S_{PIE}}\right) = \beta_0^{S_{PIE}} + \beta_{0i}^{S_{PIE}} + \beta_{0ij}^{S_{PIE}} + \left(\beta_1^{S_{PIE}} + \beta_{1i}^{S_{PIE}} + \beta_{1ij}^{S_{PIE}}\right)year_{ijt},$$

934
$$\left[\beta_{0i}^{N}, \beta_{0i}^{S}, \beta_{0i}^{S_n}, \beta_{0i}^{S_{PIE}}, \beta_{1i}^{N}, \beta_{1i}^{S}, \beta_{1i}^{S_n}, \beta_{1i}^{S_{PIE}}\right]' \sim MVNormal(\mathbf{0}, \mathbf{SRS}),$$

935
$$\mathbf{S} = \begin{bmatrix} \sigma_{0i}^{N} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{0i}^{S} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{0i}^{S_n} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{0i}^{S_{PIE}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{1i}^{N} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{1i}^{S} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1i}^{S_n} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1i}^{S_{PIE}} \end{bmatrix},$$

936
$$\mathbf{R} = \begin{bmatrix} 1 & \rho_{N_{0i}S_{0i}} & \rho_{N_{0i}S_{n0i}} & \rho_{N_{0i}SPIE_{0i}} & \rho_{N_{0i}N_{1i}} & \rho_{N_{0i}S_{1i}} & \rho_{N_{0i}S_{n1i}} & \rho_{N_{0i}SPIE_{1i}} \\ \rho_{N_{0i}S_{0i}} & 1 & \rho_{S_{0i}S_{n0i}} & \rho_{S_{0i}SPIE_{0i}} & \rho_{S_{0i}N_{1i}} & \rho_{S_{0i}S_{1i}} & \rho_{S_{0i}S_{n1i}} & \rho_{S_{0i}SPIE_{1i}} \\ \rho_{N_{0i}S_{n0i}} & \rho_{S_{0i}S_{n0i}} & 1 & \rho_{S_{n0i}SPIE_{0i}} & \rho_{S_{n0i}N_{1i}} & \rho_{S_{n0i}S_{1i}} & \rho_{S_{n0i}S_{n1i}} & \rho_{S_{n0i}SPIE_{1i}} \\ \rho_{N_{0i}SPIE_{0i}} & \rho_{S_{0i}SPIE_{0i}} & \rho_{S_{n0i}SPIE_{0i}} & 1 & \rho_{SPIE_{0i}N_{1i}} & \rho_{SPIE_{0i}S_{1i}} & \rho_{SPIE_{0i}S_{n1i}} & \rho_{SPIE_{0i}SPIE_{1i}} \\ \rho_{N_{0i}N_{1i}} & \rho_{S_{0i}N_{1i}} & \rho_{S_{n0i}N_{1i}} & \rho_{SPIE_{0i}N_{1i}} & 1 & \rho_{N_{1i}S_{1i}} & \rho_{N_{1i}S_{n1i}} & \rho_{N_{1i}SPIE_{1i}} \\ \rho_{N_{0i}S_{1i}} & \rho_{S_{0i}S_{1i}} & \rho_{S_{n0i}S_{1i}} & \rho_{SPIE_{0i}S_{1i}} & \rho_{N_{1i}S_{1i}} & 1 & \rho_{S_{1i}S_{n1i}} & \rho_{S_{1i}SPIE_{1i}} \\ \rho_{N_{0i}S_{n1i}} & \rho_{S_{0i}S_{n1i}} & \rho_{S_{n0i}S_{n1i}} & \rho_{SPIE_{0i}S_{n1i}} & \rho_{N_{1i}S_{n1i}} & \rho_{S_{1i}S_{n1i}} & 1 & \rho_{S_{n1i}SPIE_{1i}} \\ \rho_{N_{0i}SPIE_{1i}} & \rho_{S_{0i}SPIE_{1i}} & \rho_{S_{n0i}SPIE_{1i}} & \rho_{SPIE_{0i}SPIE_{1i}} & \rho_{N_{1i}SPIE_{1i}} & \rho_{S_{1i}SPIE_{1i}} & \rho_{S_{n1i}SPIE_{1i}} & 1 \end{bmatrix},$$

937
$$\left[\beta_{0ij}^{N}, \beta_{0ij}^{S}, \beta_{0ij}^{S_n}, \beta_{0ij}^{S_{PIE}}, \beta_{1ij}^{N}, \beta_{1ij}^{S}, \beta_{1ij}^{S_n}, \beta_{1ij}^{S_{PIE}}\right]' \sim MVNormal(\mathbf{0}, \mathbf{SRS}),$$

938
$$S = \begin{bmatrix} \sigma_{0ij}^N & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{0ij}^S & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{0ij}^{Sn} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{0ij}^{SPIE} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{1ij}^N & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{1ij}^S & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1ij}^{Sn} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1ij}^{SPIE} \end{bmatrix},$$

939 R

940
$$= \begin{bmatrix} 1 & \rho_{N_{0ij}S_{0ij}} & \rho_{N_{0ij}Sn_{0ij}} & \rho_{N_{0ij}SPIE_{0ij}} & \rho_{N_{0ij}N_{1ij}} & \rho_{N_{0ij}S_{1ij}} & \rho_{N_{0ij}Sn_{1ij}} & \rho_{N_{0ij}SPIE_{1ij}} \\ \rho_{N_{0ij}S_{0ij}} & 1 & \rho_{S_{0ij}Sn_{0ij}} & \rho_{S_{0ij}SPIE_{0ij}} & \rho_{S_{0ij}N_{1ij}} & \rho_{S_{0ij}S_{1ij}} & \rho_{S_{0ij}Sn_{1ij}} & \rho_{S_{0ij}SPIE_{1ij}} \\ \rho_{N_{0ij}Sn_{0ij}} & \rho_{S_{0ij}Sn_{0ij}} & 1 & \rho_{Sn_{0ij}SPIE_{0ij}} & \rho_{Sn_{0ij}N_{1ij}} & \rho_{Sn_{0ij}S_{1ij}} & \rho_{Sn_{0ij}Sn_{1ij}} & \rho_{Sn_{0ij}SPIE_{1ij}} \\ \rho_{N_{0ij}SPIE_{0ij}} & \rho_{S_{0ij}SPIE_{0ij}} & \rho_{Sn_{0ij}SPIE_{0ij}} & 1 & \rho_{SPIE_{0ij}N_{1ij}} & \rho_{SPIE_{0ij}S_{1ij}} & \rho_{SPIE_{0ij}Sn_{1ij}} & \rho_{SPIE_{0ij}SPIE_{1ij}} \\ \rho_{N_{0ij}N_{1ij}} & \rho_{S_{0ij}N_{1ij}} & \rho_{Sn_{0ij}N_{1ij}} & \rho_{SPIE_{0ij}N_{1ij}} & 1 & \rho_{N_{1ij}S_{1ij}} & \rho_{N_{1ij}Sn_{1ij}} & \rho_{N_{1ij}SPIE_{1ij}} \\ \rho_{N_{0ij}S_{1ij}} & \rho_{S_{0ij}S_{1ij}} & \rho_{Sn_{0ij}S_{1ij}} & \rho_{SPIE_{0ij}S_{1ij}} & \rho_{N_{1ij}S_{1ij}} & 1 & \rho_{S_{1ij}Sn_{1ij}} & \rho_{S_{1ij}SPIE_{1ij}} \\ \rho_{N_{0ij}Sn_{1ij}} & \rho_{S_{0ij}Sn_{1ij}} & \rho_{Sn_{0ij}Sn_{1ij}} & \rho_{SPIE_{0ij}Sn_{1ij}} & \rho_{N_{1ij}Sn_{1ij}} & \rho_{S_{1ij}Sn_{1ij}} & 1 & \rho_{Sn_{1ij}SPIE_{1ij}} \\ \rho_{N_{0ij}SPIE_{1ij}} & \rho_{S_{0ij}SPIE_{1ij}} & \rho_{Sn_{0ij}SPIE_{1ij}} & \rho_{SPIE_{0ij}SPIE_{1ij}} & \rho_{N_{1ij}SPIE_{1ij}} & \rho_{S_{1ij}SPIE_{1ij}} & \rho_{Sn_{1ij}SPIE_{1ij}} & 1 \end{bmatrix},$$

941
$$\beta_0^N, \beta_0^S, \beta_0^{Sn}, \beta_0^{SPIE} \sim N(0,1),$$

942
$$\beta_1^N, \beta_1^S, \beta_1^{Sn}, \beta_1^{SPIE} \sim N(0, 0.2),$$

943
$$\sigma_{0i}^S, \sigma_{0i}^{Sn}, \sigma_{0i}^{SPIE}, \sigma_{1i}^S, \sigma_{1i}^{Sn}, \sigma_{1i}^{SPIE}, \sigma_{0ij}^S, \sigma_{0ij}^{Sn}, \sigma_{0ij}^{SPIE}, \sigma_{1ij}^S, \sigma_{1ij}^{Sn}, \sigma_{1ij}^{SPIE} \sim student(3,0,10),$$

944
$$\sigma, \sigma_{0i}^N, \sigma_{1ij}^N, \sigma_{0ij}^N, \sigma_{1ij}^N \sim student(3,0,16),$$

945
$$R \sim LJK(1),$$

where N_{ijt}, S_{ijt}, $S_{n_{ijt}}$, and $S_{PIE_{ijt}}$ are the values of each of the metrics in the *j*th cell of the *i*th
study in year *t*, and year$_{ijt}$ is the time in years. β_0 and β_1 (with the respective superscripts for each
metric) represent the non-varying intercepts and slopes, respectively. β_{0i} and β_{1i} (with the
respective superscripts for each metric) represent the varying intercepts and slopes for study-
level departures from β_0 and β_1, respectively. β_{0ij} and β_{1ij} (with the respective superscripts for
each metric) represent the varying intercepts and slopes for cell-level departures from β_0 and β_1,
respectively. The covariance matrices of each multivariate normal distribution for varying effects
(one each for the study- and cell-level departures) are parameterised in terms of a correlation
matrix **R** and two matrices **S** with diagonal elements σ (superscripts for metrics, subscripts
denote intercept (0), slopes (1) and level: studies *i* and cells *j*).
956

Figure S1: Posterior predictive checks for the model fit to the temporal data in naturally varying environments. (a) Total
numbers of individuals, (b) species richness, (c) rarefied richness (S_n) and (d) evenness (S_{PIE}). Each panel shows the density of
the data (y) and ten draws of the posterior distribution (yrep).

Temporal comparisons: experimental or natural perturbations

We fit a model that assumed lognormal distributions for all response variables (*N, S, S*n, *S*PIE).
Our focus is on the treatment-level variation in the metrics, and we created a covariate that was
the concatenation of study and treatment to this end. Additionally, as some studies had blocks
within sites, and others did not, we created a new variable that was the concatenation of site and
block, where those without block had no unique identifier. We ran the model with four chains for
3000 iterations, with 1500 used as warmup. The model took the following form:

971
$$N_{ijkt} \sim lognormal(\mu_{ijkt}^N, \sigma^N),$$

972
$$S_{ijkt} \sim lognormal(\mu_{ijkt}^S, \sigma^S),$$

973
$$S_{n\,ijkt} \sim lognormal(\mu_{ijkt}^{Sn}, \sigma^{Sn}),$$

974
$$S_{PIE\,ijkt} \sim lognormal(\mu_{ijkt}^{SPIE}, \sigma^{SPIE}),$$

975
$$\mu_{ijkt}^N = \beta_0^N + \beta_{0i}^N + \beta_{0j}^N + \beta_{0k}^N + (\beta_1^N + \beta_{1k}^N)year_{ijkt},$$

976
$$\mu_{ijkt}^S = \beta_0^S + \beta_{0i}^S + \beta_{0j}^S + \beta_{0k}^S + (\beta_1^S + \beta_{1k}^S)year_{ijkt},$$

977
$$\mu_{ijkt}^{Sn} = \beta_0^{Sn} + \beta_{0i}^{Sn} + \beta_{0j}^{Sn} + \beta_{0k}^{Sn} + (\beta_1^{Sn} + \beta_{1k}^{Sn})year_{ijkt},$$

978
$$\mu_{ijkt}^{SPIE} = \beta_0^{SPIE} + \beta_{0i}^{SPIE} + \beta_{0j}^{SPIE} + \beta_{0k}^{SPIE} + (\beta_1^{SPIE} + \beta_{1k}^{SPIE})year_{ijkt},$$

979
$$[\beta_{0k}^N, \beta_{0k}^S, \beta_{0k}^{Sn}, \beta_{0k}^{SPIE}, \beta_{1k}^N, \beta_{1k}^S, \beta_{1k}^{Sn}, \beta_{1k}^{SPIE}]' \sim MVNormal(\mathbf{0}, \mathbf{SRS}),$$

980
$$\mathbf{S} = \begin{bmatrix} \sigma_{0k}^N & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{0k}^S & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{0k}^{Sn} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{0k}^{SPIE} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{1k}^N & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^S & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^{Sn} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^{SPIE} \end{bmatrix},$$

981
$$\mathbf{R} = \begin{bmatrix} 1 & \rho_{N_{0k}S_{0k}} & \rho_{N_{0k}S_{n0k}} & \rho_{N_{0k}SPIE_{0k}} & \rho_{N_{0k}N_{1k}} & \rho_{N_{0k}S_{1k}} & \rho_{N_{0k}S_{n1k}} & \rho_{N_{0k}SPIE_{1k}} \\ \rho_{N_{0k}S_{0k}} & 1 & \rho_{S_{0k}S_{n0k}} & \rho_{S_{0k}SPIE_{0k}} & \rho_{S_{0k}N_{1k}} & \rho_{S_{0k}S_{1k}} & \rho_{S_{0k}S_{n1k}} & \rho_{S_{0k}SPIE_{1k}} \\ \rho_{N_{0k}S_{n0k}} & \rho_{S_{0k}S_{n0k}} & 1 & \rho_{S_{n0k}SPIE_{0k}} & \rho_{S_{n0k}N_{1k}} & \rho_{S_{n0k}S_{1k}} & \rho_{S_{n0k}S_{n1k}} & \rho_{S_{n0k}SPIE_{1k}} \\ \rho_{N_{0k}SPIE_{0k}} & \rho_{S_{0k}SPIE_{0k}} & \rho_{S_{n0k}SPIE_{0k}} & 1 & \rho_{SPIE_{0k}N_{1k}} & \rho_{SPIE_{0k}S_{1k}} & \rho_{SPIE_{0k}S_{n1k}} & \rho_{SPIE_{0k}SPIE_{1k}} \\ \rho_{N_{0k}N_{1k}} & \rho_{S_{0k}N_{1k}} & \rho_{S_{n0k}N_{1k}} & \rho_{SPIE_{0k}N_{1k}} & 1 & \rho_{N_{1k}S_{1k}} & \rho_{N_{1k}S_{n1k}} & \rho_{N_{1k}SPIE_{1k}} \\ \rho_{N_{0k}S_{1k}} & \rho_{S_{0k}S_{1k}} & \rho_{S_{n0k}S_{1k}} & \rho_{SPIE_{0k}S_{1k}} & \rho_{N_{1k}S_{1k}} & 1 & \rho_{S_{1k}S_{n1k}} & \rho_{S_{1k}SPIE_{1k}} \\ \rho_{N_{0k}S_{n1k}} & \rho_{S_{0k}S_{n1k}} & \rho_{S_{n0k}S_{n1k}} & \rho_{SPIE_{0k}S_{n1k}} & \rho_{N_{1k}S_{n1k}} & \rho_{S_{1k}S_{n1k}} & 1 & \rho_{S_{n1k}SPIE_{1k}} \\ \rho_{N_{0k}SPIE_{1k}} & \rho_{S_{0k}SPIE_{1k}} & \rho_{S_{n0k}SPIE_{1i}} & \rho_{SPIE_{0k}SPIE_{1k}} & \rho_{N_{1k}SPIE_{1k}} & \rho_{S_{1k}SPIE_{1k}} & \rho_{S_{n1k}SPIE_{1k}} & 1 \end{bmatrix},$$

982
$$\beta_0^N, \beta_0^S, \beta_0^{Sn}, \beta_0^{SPIE} \sim N(0, 1),$$

983
$$\beta_1^N, \beta_1^S, \beta_1^{Sn}, \beta_1^{SPIE} \sim N(0, 0.2),$$

984
$$\sigma_{0i}^S, \sigma_{0i}^{Sn}, \sigma_{0i}^{SPIE}, \sigma_{1i}^S, \sigma_{1i}^{Sn}, \sigma_{1i}^{SPIE}, \sigma_{0ij}^S, \sigma_{0ij}^{Sn}, \sigma_{0ij}^{SPIE}, \sigma_{1ij}^S, \sigma_{1ij}^{Sn}, \sigma_{1ij}^{SPIE} \sim normal(0, 1),$$

985
$$\sigma, \sigma_{0i}^N, \sigma_{1ij}^N, \sigma_{0ij}^N, \sigma_{1ij}^N \sim student(3, 0, 16),$$

986
$$\mathbf{R} \sim LJK(1),$$

where N_{ijkt}, S_{ijkt}, $S_{n_{ijkt}}$, and $S_{PIE_{ijkt}}$ are the values of each of the metrics in the kth study-

treatment combination, for the jth study-block combination of the ith study in year t, and year$_{ijkt}$

is the time in years. β_0 and β_1 (with the respective superscripts for each metric) represent the

non-varying intercepts and slopes, respectively. β_{0i} (with the respective superscripts for each

metric) represent the varying intercepts study-level departures from β_0, and β_{0j} (with the

respective superscripts for each metric) represent the varying intercepts study-block level

departures from β_0. β_{0k} and β_{1k} (with the respective superscripts for each metric) represent the

varying intercepts and slopes for study-treatment departures from β_0 and β_1, respectively. The

covariance matrix of the multivariate normal distribution for varying study-treatment effects

were parameterised in terms of a correlation matrix **R** and two matrices **S** with diagonal elements

σ (superscripts for metrics, subscripts denote intercept (0), slopes (1) and study-treatment

combination, k).

Figure S2: Posterior predictive checks for the model fit to the temporal data in perturbed environments. (a) Total numbers of individuals, (b) species richness, (c) rarefied richness (Sn) and (d) evenness (SPIE).). Each panel shows the density of the data (y) and ten draws of the posterior distribution (yrep).

References (for times series with experimental or natural perturbations)

Bateman, H., D. Childers, and P. Warren. 2018. Point-count bird censusing: long-term monitoring of bird abundance and diversity along the Salt River in the greater Phoenix metropolitan area, ongoing since 2013 ver 3.

Carpenter, S., J. Kitchell, J. Cole, and M. Pace. 2017. Cascade Project at North Temperate Lakes LTER Core Data Zooplankton 1984 - 2016 ver 4. Environmental Data Initiative.

Carpenter, S., J. Kitchell, J. Cole, and M. Pace. 2017. Cascade Project at North Temperate Lakes LTER Core Data Phytoplankton 1984 - 2015 ver 4. Environmental Data Initiative.

Ellison, A. and R. Dunn. 2018. Ants Under Climate Change at Harvard Forest and Duke Forest
2009-2015 ver 57. Environmental Data Initiative.
Philip Grime and Jason Fridley. Point Quadrat Vegetation Data, Buxton Climate Change
Experiment, UK, 1994-2009. Knowledge Network for Biocomplexity.
doi:10.5063/AA/fridley.3.2.
Hershey, A. 2016. Total numbers and species of insects taken from rock scrubbings during the
summer of 1984-1988, 1993-1994, 1996-1998, in the Kuparuk River experimental reach near
Toolik Field Station, North Slope Alaska.. ver 3. Environmental Data Initiative.
Joern, A. 2020. CGR02 Sweep Sampling of Grasshoppers on Konza Prairie LTER watersheds
ver 14. Environmental Data Initiative.
Joern, A. 2020. PBG07 Grasshopper species abundances in the Patch-Burn Grazing experiment
at Konza Prairie ver 5. Environmental Data Initiative.
Knops, J. 2018. Arthropod sweep net sampling: Interactive Effects of Deer, Fire and Nitrogen
ver 8. Environmental Data Initiative.
Knops, J. 2018. Small mammal abundance: Interactive Effects of Deer, Fire and Nitrogen ver 8.
Environmental Data Initiative.
Santa Barbara Coastal LTER and D. Reed. 2020. SBC LTER: Reef: Long-term experiment: Kelp
removal: Fish abundance ver 18. Environmental Data Initiative.
1030
1031

Spatial comparisons: natural environmental variation

We fit a model that assumed lognormal distributions for N, S_n, and S_{PIE}, and a Poisson distribution (and log link function) for S. We fit the model with four chains and 4000 iterations, with 2000 as warmup. The model took the following form:

$$N_{ij} \sim lognormal\left(\mu_{ij}^N, \sigma^N\right),$$

$$S_{ij} \sim poisson\left(\lambda_{ij}^S\right),$$

$$S_{n_{ij}} \sim lognormal\left(\mu_{ij}^{S_n}, \sigma^{S_n}\right),$$

$$S_{PIE_{ij}} \sim lognormal\left(\mu_{ij}^{S_{PIE}}, \sigma^{S_{PIE}}\right), \text{(S1.4)}$$

$$\mu_{ij}^N = \beta_0^N + \beta_{0j}^N + (\beta_{1j}^N)site_{ij},$$

$$\log\left(\lambda_{ij}^S\right) = \beta_0^S + \beta_{0i}^S + \beta_{0j}^S + (\beta_{1j}^S)site_{ij},$$

$$\mu_{ij}^{S_n} = \beta_0^{S_n} + \beta_{0j}^{S_n} + (\beta_{1j}^{S_n})site_{ij},$$

$$\mu_{ij}^{S_{PIE}} = \beta_0^{S_{PIE}} + \beta_{0j}^{S_{PIE}} + (\beta_{1j}^{S_{PIE}})site_{ij},$$

$$\left[\beta_{0j}^N, \beta_{0j}^S, \beta_{0j}^{S_n}, \beta_{0j}^{S_{PIE}}, \beta_{1j}^N, \beta_{1j}^S, \beta_{1j}^{S_n}, \beta_{1j}^{S_{PIE}}\right]' \sim MVNormal(\mathbf{0}, \mathbf{SRS}),$$

$$\mathbf{S} = \begin{bmatrix} \sigma_{0j}^N & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{0j}^S & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{0j}^{S_n} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{0j}^{S_{PIE}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{1j}^N & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{1j}^S & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1j}^{S_n} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1j}^{S_{PIE}} \end{bmatrix},$$

$$\mathbf{R} = \begin{bmatrix} 1 & \rho_{N_{0j}S_{0j}} & \rho_{N_{0j}S_{n_{0j}}} & \rho_{N_{0j}S_{PIE_{0j}}} & \rho_{N_{0j}N_{1j}} & \rho_{N_{0j}S_{1j}} & \rho_{N_{0j}S_{n_{1j}}} & \rho_{N_{0j}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{0j}} & 1 & \rho_{S_{0j}S_{n_{0j}}} & \rho_{S_{0j}S_{PIE_{0j}}} & \rho_{S_{0j}N_{1j}} & \rho_{S_{0j}S_{1j}} & \rho_{S_{0j}S_{n_{1j}}} & \rho_{S_{0j}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{n_{0j}}} & \rho_{S_{0j}S_{n_{0j}}} & 1 & \rho_{S_{n_{0j}}S_{PIE_{0j}}} & \rho_{S_{n_{0j}}N_{1j}} & \rho_{S_{n_{0j}}S_{1j}} & \rho_{S_{n_{0j}}S_{n_{1j}}} & \rho_{S_{n_{0j}}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{PIE_{0j}}} & \rho_{S_{0j}S_{PIE_{0j}}} & \rho_{S_{n_{0j}}S_{PIE_{0j}}} & 1 & \rho_{S_{PIE_{0j}}N_{1j}} & \rho_{S_{PIE_{0j}}S_{1j}} & \rho_{S_{PIE_{0j}}S_{n_{1j}}} & \rho_{S_{PIE_{0j}}S_{PIE_{1j}}} \\ \rho_{N_{0j}N_{1j}} & \rho_{S_{0j}N_{1j}} & \rho_{S_{n_{0j}}N_{1j}} & \rho_{S_{PIE_{0j}}N_{1j}} & 1 & \rho_{N_{1j}S_{1j}} & \rho_{N_{1j}S_{n_{1j}}} & \rho_{N_{1j}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{1j}} & \rho_{S_{0j}S_{1j}} & \rho_{S_{n_{0j}}S_{1j}} & \rho_{S_{PIE_{0j}}S_{1j}} & \rho_{N_{1j}S_{1j}} & 1 & \rho_{S_{1j}S_{n_{1j}}} & \rho_{S_{1j}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{n_{1j}}} & \rho_{S_{0j}S_{n_{1j}}} & \rho_{S_{n_{0j}}S_{n_{1j}}} & \rho_{S_{PIE_{0j}}S_{n_{1j}}} & \rho_{N_{1j}S_{n_{1j}}} & \rho_{S_{1j}S_{n_{1j}}} & 1 & \rho_{S_{n_{1j}}S_{PIE_{1j}}} \\ \rho_{N_{0j}S_{PIE_{1j}}} & \rho_{S_{0k}S_{PIE_{1j}}} & \rho_{S_{n_{0j}}S_{PIE_{1j}}} & \rho_{S_{PIE_{0j}}S_{PIE_{1j}}} & \rho_{N_{1j}S_{PIE_{1j}}} & \rho_{S_{1j}S_{PIE_{1j}}} & \rho_{S_{n_{1j}}S_{PIE_{1j}}} & 1 \end{bmatrix},$$

$$\beta_0^N, \beta_0^S, \beta_0^{S_n}, \beta_0^{S_{PIE}} \sim N(0,1),$$

$$\sigma^N, \sigma^{S_n}, \sigma^{S_{PIE}}, \sigma_{0i}^S, \sigma_{0j}^N, \sigma_{0j}^S, \sigma_{0j}^{S_n}, \sigma_{0j}^{S_{PIE}}, \sigma_{1j}^N, \sigma_{1j}^S, \sigma_{1j}^{S_n}, \sigma_{1j}^{S_{PIE}} \sim normal(0,1),$$

$$\mathbf{R} \sim LJK(1),$$

where N_{ij}, S_{ij}, $S_{n_{ij}}$, and $S_{PIE_{ij}}$ are the values of each of the metrics for the ith observation in the jth study-site combination, and site$_{ij}$ is site identifier. β_0 (with the respective superscripts for each metric) represent the non-varying intercepts for each data source (one each for CESTES and

McGill), respectively. β_{0i}^S is an observation level varying intercept to model overdispersion in S.
β_{0j} and β_{1j} (with the respective superscripts for each metric) represent the varying intercepts
(reference sites for each study) and slopes (departures for all other sites). The covariance matrix
of the multivariate normal distribution for varying study-site effects were parameterised in terms
of a correlation matrix **R** and two matrices **S** with diagonal elements σ (superscripts for metrics,
subscripts denote intercept (0), slopes (1)).
1061

1062
1063 *Figure S3: Posterior predictive checks for the model fit to the spatial data in natural environments. (a) Total numbers of*
1064 *individuals, (b) species richness, (c) rarefied richness (S_n) and (d) evenness (S_{PIE}). Each panel shows the density of the data*
1065 *(y) and ten draws of the posterior distribution (yrep).*

1066

Spatial comparisons: anthropogenic perturbations

1068

1069 We fit a model that assumed lognormal distributions for N, S_n, and S_{PIE}, and a Poisson

1070 distribution (and log link function) for S. We fit the model with four chains and 4000 iterations,

1071 with 2000 used as warmup. The model took the following form:

1072

1073
$$N_{ijk} \sim lognormal(\mu_{ijk}^N, \sigma^N),$$

1074
$$S_{ijk} \sim poisson(\lambda_{ijk}^S),$$

1075
$$S_{n_{ijk}} \sim lognormal(\mu_{ijk}^{S_n}, \sigma^{S_n}),$$

1076
$$S_{PIE_{ijk}} \sim lognormal(\mu_{ijk}^{SPIE}, \sigma^{SPIE}),$$

1077
$$\mu_{ijk}^N = \beta_0^N + \beta_{0j}^N + \beta_{0k}^N + (\beta_1^N + \beta_{1k}^N)LU_{ij},$$

1078
$$\log(\lambda_{ijk}^S) = \beta_0^S + \beta_{0i}^S + \beta_{0j}^S + \beta_{0k}^S + (\beta_1^S + \beta_{1k}^S)LU_{ijk},$$

1079
$$\mu_{ijk}^{S_n} = \beta_0^{S_n} + \beta_{0j}^{S_n} + \beta_{0k}^{S_n} + (\beta_1^{S_n} + \beta_{1k}^{S_n})LU_{ijk},$$

1080
$$\mu_{ijk}^{SPIE} = \beta_0^{SPIE} + \beta_{0j}^{SPIE} + \beta_{0k}^{SPIE} + (\beta_1^{SPIE} + \beta_{1k}^{SPIE})LU_{ijk},$$

1081
$$[\beta_{0j}^N, \beta_{0j}^S, \beta_{0j}^{S_n}, \beta_{0j}^{SPIE}, \beta_{1j}^N, \beta_{1j}^S, \beta_{1j}^{S_n}, \beta_{1j}^{SPIE}]' \sim MVNormal(\mathbf{0}, \mathbf{SRS}),$$

1082
$$\mathbf{S} = \begin{bmatrix} \sigma_{0k}^N & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{0k}^S & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{0k}^{S_n} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{0k}^{SPIE} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{1k}^N & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^S & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^{S_n} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sigma_{1k}^{SPIE} \end{bmatrix},$$

1083
$$\mathbf{R} = \begin{bmatrix} 1 & \rho_{N_{0k}S_{0k}} & \rho_{N_{0k}S_{n0k}} & \rho_{N_{0k}SPIE_{0k}} & \rho_{N_{0k}N_{1k}} & \rho_{N_{0k}S_{1k}} & \rho_{N_{0k}S_{n1k}} & \rho_{N_{0k}SPIE_{1k}} \\ \rho_{N_{0k}S_{0k}} & 1 & \rho_{S_{0k}S_{n0k}} & \rho_{S_{0k}SPIE_{0k}} & \rho_{S_{0k}N_{1k}} & \rho_{S_{0k}S_{1k}} & \rho_{S_{0k}S_{n1k}} & \rho_{S_{0k}SPIE_{1k}} \\ \rho_{N_{0k}S_{n0k}} & \rho_{S_{0k}S_{n0k}} & 1 & \rho_{S_{n0k}SPIE_{0k}} & \rho_{S_{n0k}N_{1k}} & \rho_{S_{n0k}S_{1k}} & \rho_{S_{n0k}S_{n1k}} & \rho_{S_{n0k}SPIE_{1k}} \\ \rho_{N_{0k}SPIE_{0k}} & \rho_{S_{0k}SPIE_{0k}} & \rho_{S_{n0k}SPIE_{0k}} & 1 & \rho_{SPIE_{0k}N_{1k}} & \rho_{SPIE_{0k}S_{1k}} & \rho_{SPIE_{0k}S_{n1k}} & \rho_{SPIE_{0k}SPIE_{1k}} \\ \rho_{N_{0k}N_{1k}} & \rho_{S_{0k}N_{1k}} & \rho_{S_{n0k}N_{1k}} & \rho_{SPIE_{0k}N_{1k}} & 1 & \rho_{N_{1k}S_{1k}} & \rho_{N_{1k}S_{n1k}} & \rho_{N_{1k}SPIE_{1k}} \\ \rho_{N_{0k}S_{1k}} & \rho_{S_{0k}S_{1k}} & \rho_{S_{n0k}S_{1k}} & \rho_{SPIE_{0k}S_{1k}} & \rho_{N_{1k}S_{1k}} & 1 & \rho_{S_{1k}S_{n1k}} & \rho_{S_{1k}SPIE_{1k}} \\ \rho_{N_{0k}S_{n1k}} & \rho_{S_{0k}S_{n1k}} & \rho_{S_{n0k}S_{n1k}} & \rho_{SPIE_{0k}S_{n1k}} & \rho_{N_{1k}S_{n1k}} & \rho_{S_{1k}S_{n1k}} & 1 & \rho_{S_{n1k}SPIE_{1k}} \\ \rho_{N_{0k}SPIE_{1k}} & \rho_{S_{0k}SPIE_{1k}} & \rho_{S_{n0k}SPIE_{1i}} & \rho_{SPIE_{0k}SPIE_{1k}} & \rho_{N_{1k}SPIE_{1k}} & \rho_{S_{1k}SPIE_{1k}} & \rho_{S_{n1k}SPIE_{1k}} & 1 \end{bmatrix},$$

1084
$$\beta_0^N, \beta_1^N \sim student(3, 5, 10),$$

1085
$$\beta_0^S, \beta_1^S \sim student(3, 3, 10),$$

1086
$$\beta_0^{S_n}, \beta_0^{SPIE}, \beta_1^{S_n}, \beta_1^{SPIE} \sim student(3, 2, 10),$$

1087
$$\sigma^N, \sigma^{S_n}, \sigma^{SPIE}, \sigma_{0i}^S, \sigma_{0j}^N, \sigma_{0j}^S, \sigma_{0j}^{S_n}, \sigma_{0j}^{SPIE}, \sigma_{0k}^N, \sigma_{0k}^S, \sigma_{0k}^{S_n}, \sigma_{0k}^{SPIE}, \sigma_{1k}^N, \sigma_{1k}^S, \sigma_{1k}^{S_n}, \sigma_{1k}^{SPIE} \sim student(3, 0, 10),$$

1088
$$\mathbf{R} \sim LJK(1),$$

1089

where N_{ijk}, S_{ijk}, $S_{n_{ijk}}$, and $S_{PIE_{ijk}}$ are the values of each of the metrics for the ith concatenation

of source ID, study number, block and site number (denoted SSBS in the database), in the jth

concatenation of source ID, study number, block (denoted SSB in the database), of the kth

combination of source ID and study (denoted SS in the database), and LU is an identifier for

land use types (other than primary vegetation, that was fit as the intercept). β_0 (with the

respective superscripts for each metric) represent the non-varying intercepts for reference land

use category (i.e., primary vegetation). β_{0i}^S is a site-level varying intercept to model

overdispersion in S. β_{0j} (with the respective superscripts for each metric) is a varying intercept

for blocks. β_{0k} and β_{1k} (with the respective superscripts for each metric) represent the varying

intercepts (reference sites for each study) and slopes (departures for all other sites). The

covariance matrix of the multivariate normal distribution for varying study-site effects were

parameterised in terms of a correlation matrix $\mathbf{R}$ and two matrices $\mathbf{S}$ with diagonal elements σ

(superscripts for metrics, subscripts denote intercept (0), slopes (1)).

1103
1104 *Figure S4: Posterior predictive checks for the model fit to the spatial data in anthropogenically perturbed environments. (a)*
1105 *Total numbers of individuals, (b) species richness, (c) rarefied richness (S_n) and (d) evenness (S_{PIE}). Each panel shows the*
1106 *density of the data (y) and ten draws of the posterior distribution (yrep).*

1107
Figure S5: Summary of assemblage diversity components with a high probability of change (90% credible interval
did not overlap zero).(a) temporal changes in naturally varying environments, (b) temporal changes in perturbed
environments, (c) spatial changes relative to an arbitrary reference, (d) spatial changes relative to primary
vegetation. Assemblages with no component changes different from zero are reported as insets for clarity. Metric
abbreviations: total number of individuals (N), expected number of species for n individuals (S_n), numbers
equivalent transformation of the Probability of Interspecific Encounter (S_{PIE}), and total species richness (S). Number
following each bar is the count of assemblages for that category.
1115
1116

Figure S6: Relationships between four components of local diversity change (not shown on Figure 3). Change in rarefied richness as a function of changes in evenness (left), change in rarefied richness as a function of changes in total abundance (middle) and evenness (right) for (a-c) study-level estimates of temporal changes in naturally varying environments; (d-f) estimates of temporal change for combinations of study and treatment in perturbed environments; (c) estimates of spatial changes within studies from an arbitrary reference site along natural

environmental gradients; and, (d) estimates of spatial change within studies between primary vegetation and different land use categories. Coloured concentration ellipses show 10% increments (5 – 95%) of the posterior distributions. Dotted grey lines are x = y = 0, and x = y for visual reference. NB: Scale of x- and y-axes vary between data sources; one estimate with ΔN = -1.79, ΔS = -3.77, ΔSn = -3.23, $\Delta SPIE$ = -3.21, removed from (j-l) for clarity.