# Proteomic phenotyping of stimulated Müller cells uncovers profound pro-inflammatory signaling and antigen-presenting capacity

**Adrian Schmalen[1,2], Lea Lorenz[2], Antje Grosche[3], Diana Pauly[4], Cornelia A. Deeg[2], Stefanie M.**
**Hauck[1*]**

[1]Research Unit Protein Science and Metabolomics and Proteomics Core, Helmholtz Zentrum
München, German Research Center for Environmental Health, D-85764 Neuherberg, Germany

[2]Department of Veterinary Sciences, Ludwig-Maximilians-Universität Munich, D-82152
Martinsried, Germany

[3]Department of Physiological Genomics, Biomedical Center, Ludwig Maximilians University
Munich, D-82152 Martinsried, Germany

[4]Experimental Ophthalmology, Philipps-University Marburg, D-35043 Marburg, Germany

**\* Correspondence:**
Corresponding Author
hauck@helmholtz-muenchen.de

**Keywords: Müller cells, atypical antigen-presenting cell, immune response, diabetic**
**retinopathy, pro-inflammatory, complement system, cytokines.**

**Abstract**

Müller cells are the main macroglial cells of the retina exerting a wealth of functions to maintain
retinal homoeostasis. Upon pathological changes in the retina, they become gliotic with both
protective and detrimental consequences. Accumulating data also provide evidence for a pivotal role
of Müller cells in the pathogenesis of diabetic retinopathy (DR). While microglial cells, the resident
immune cells of the retina are considered as main players in inflammatory processes associated with
DR, the implication of activated Müller cells in chronic retinal inflammation remains to be
elucidated. In order to assess the signaling capacity of Müller cells and their role in retinal
inflammation, we performed in-depth proteomic analysis of Müller cell proteomes and secretomes
after stimulation with INFγ, TNFα, IL-4, IL-6, IL-10, TGFβ1, TGFβ2 and TGFβ3. We used both,
primary porcine Müller cells and the human Müller cell line MIO-M1 for our hypothesis generating
approach. Our results point towards an intense signaling capacity of Müller cells, which reacted in a
highly discriminating manner upon treatment with different cytokines. Stimulation of Müller cells
results in a primarily pro-inflammatory phenotype with secretion of cytokines and components of the
complement system. Furthermore, we observed evidence for mitochondrial dysfunction, implying
oxidative stress after treatment with the various cytokines. Finally, both MIO-M1 cells and primary
porcine Müller cells showed several characteristics of atypical antigen-presenting cells, as they are
capable of inducing MHC class I and MHC class II with co-stimulatory molecules. In line with this,
they express proteins associated with formation and maturation of phagosomes. Thus, our findings
underline the importance of Müller cell signaling in the inflamed retina, indicating an active role in
chronic retinal inflammation underlying the pathogenesis of diabetic retinopathy.

## 1    Introduction

Neurodegenerative diseases of the retina are characterized by progressive photoreceptor damage eventually resulting in vision loss (1). Accumulating evidence over the last years led to the recognition of chronic inflammation as an important part of the pathogenesis underlying this heterogeneous group of retinal diseases (2-4). While the loss of photoreceptors in inherited retinal diseases such as retinitis pigmentosa originates in various genetic mutations (5), the driving force that leads to retinal degeneration and blindness in diabetic retinopathy (DR) is a disturbed metabolism with hyperglycemia and dyslipidemia, resulting in microvascular damage (6). DR is among the most frequent causes of blindness worldwide with a rising prevalence (7). Chronic hyperglycemia in diabetes patients induces the activation of leukocytes in the periphery (8) as well as micro- and macroglial cells in the retina (9-11). This results in the release of pro-inflammatory cytokines and eventually leads to photoreceptor cell death (12, 13). Microglial cells, the resident immune cells of the retina, are acknowledged as the main drivers of retinal immune responses (14). However, growing evidence suggests that the interaction of micro- and macroglial cells essentially shapes retinal inflammation and photoreceptor degeneration (15, 16).

Retinal Müller glial cells constitute the primary macroglial cells of the retina (17). They span the entire width of the retina and are in close contact with the vitreous, the retinal blood-vessels and with all retinal neurons (18). While they maintain retinal homeostasis during steady-state conditions, activation of Müller cells under pathological conditions results in gliosis, a cellular attempt to restore insulted tissue, with both protective and detrimental effects (19). It is known that Müller cells are an important source of neurotrophic factors but also of pro-inflammatory and angiogenetic cytokines (20-23). They play an important role in DR pathogenesis (24, 25) and it was also suggested that they are involved in retinal immune responses (26-30). However, the impact of their protective or detrimental effects on retinal inflammation in DR and other neurodegenerative retinal diseases remains elusive so far. Thus, we performed an in-depth-analysis of the Müller cell proteome and secretome after stimulation with various pro- and anti-inflammatory cytokines as well as growth factors. For our analysis, we used cells of the human Müller glia cell-line MIO-M1 as well as primary porcine retinal Müller Glia (pRMG), as porcine eyes resemble human eyes regarding their anatomy (31). Furthermore, the pig is established as a useful model for research in diabetic retinopathy (32, 33). Comparative quantitative proteomic analysis allows to elucidate key proteins or pathways involved in disease pathogenesis. In addition, this approach enables the discovery of new biomarkers and has therefore proven as a valuable tool for deciphering pathophysiological key mechanisms (34, 35). Our hypothesis-generating study yielded comprehensive data on the capacity of Müller cells to react in a very differentiated manner to varying stimulants. In addition to the secretion of pro-inflammatory cytokines, we observed expression of MHC class I and II molecules and proteins that are associated with the processing of antigens. We therefore propose that Müller cells are critical modulators of the retinal immune response and might exert an antigen-presenting function. Thus, attention should be paid to their implication in chronic inflammation underlying degenerative retinal diseases.

## 2    Materials and methods

### 2.1    Cell preparation and culture

Cells were maintained in Dulbecco's modified eagle medium (DMEM) containing 10% (*v/v*) inactivated fetal calf serum (FCS), 100 U/mL penicillin and 100 μg/mL streptomycin unless

otherwise stated. Cell culture media and reagents were purchased from Gibco (Life Technologies
GmbH, Darmstadt, Germany).

Ten porcine eyes from healthy pigs were kindly provided from a local abattoir. The use of porcine
material from the abattoir was approved for purposes of scientific research by the appropriate board
of the veterinary inspection office, Munich, Germany (registration number DE 09 162 0008-21). No
experimental animals were involved in this study. Within 2h after enucleation, eyes were processed
under a laminar flow hood under sterile conditions as previously described (30, 36). In short,
periocular tissue was removed and the eyeballs were rinsed in 80% ethanol followed by washing with
cold PBS. Afterwards, eyeballs were stored in DMEM until further processing. The eyeballs were
opened circumferentially parallel to the limbus corneae, and anterior parts of the eyes were discarded.
The retina was detached from the posterior eyeballs and transferred into a petri dish containing
DMEM. After removal of vitreous and pigment epithelium residues, major blood-vessels were
excised and the remaining retinal tissue was cut into very small fragments using micro-scissors.
Resulting fragments were washed in Ringer's solution followed by enzymatic digestion at 37 °C with
papain previously activated by incubation with 1.1 µM EDTA, 0.067 µM mercaptoethanol and 5.5
µM cysteine-HCl. Enzymatic digestion was stopped after 12 min by adding serum-containing
DMEM, followed by addition of Desoxyribonuclease I (Sigma-Aldrich Chemie GmbH, Taufkirchen,
Germany) and trituration. After sedimentation of the cells, the supernatant was carefully removed
using Pasteur pipettes. The remaining pellets were resuspended in DMEM, pooled and seeded into 6-
well plates (Sarstedt, Nümbrecht, Germany). The following day, thorough panning of the plates and
removal of the supernatant were performed in order to eliminate non-attached cells, yielding pure
Müller cell cultures as previously described (37, 38). Cells were cultured at 37 °C and 5% $CO_2$ with
regular exchange of medium and repeated microscopic control of cell density and purity according to
previous reports (37, 39).

The human Müller cell line Moorfields/Institute of Ophthalmology-Müller 1 (MIO-M1;
RRID:CVCL_0433) was a kind gift of G. A. Limb (39). They were tested negative for mycoplasma
contamination. Two days before treatment, $1 \times 10^5$ MIO-M1 cells per well were seeded in 6-well
plates and incubated at 37 °C and 5% $CO_2$ until further processing.

## 2.2   Cell stimulation

The human cytokines Interleukin 10 (IL-10), Transforming Growth Factor beta-1 (TGFβ1),
Transforming Growth Factor beta-2 (TGFβ2), Transforming Growth Factor beta-3 (TGFβ3), and
Tumor Necrosis Factor Alpha (TNFα) were purchased from Sigma-Aldrich, Interleukin 4 (IL-4) and
Interferon Gamma (IFNy) from R&D Systems/Bio-Techne (R&D Systems/Bio-Techne, Wiesbaden-
Nordenstadt, Germany), and Interleukin 6 (IL-6) and Vascular Endothelial Growth Factor$_{165}$ (VEGF)
from PeproTech (PeproTech, Winterhude, Germany). Porcine TGFβ3 was purchased from Biozol
(Biozol, Eching, Germany), whereas porcine TGFβ1, TGFβ2, IL-4, IL-6, IL-10, IFNy and TNFα
were from R&D System/Bio-Techne. Since there was no porcine VEGF available, the above
mentioned human VEGF was also used for stimulation of pRMG.

To diminish the influence of cytokines present in FCS, both confluent pRMG and MIO-M1 cells
were rinsed two times with prewarmed serum-free medium, followed by starvation for 1 h at 37 °C
and 5% $CO_2$ with serum-deprived medium. Afterwards, cells were treated over night with IFNy, IL-
4, IL-6, IL-10, TGFβ1, TGFβ2, TGFβ3, TNFα or VEGF$_{165}$, respectively, in a randomized plate
design at a concentration of 5 ng/mL in medium without FCS. Untreated cells cultured in serum-free
medium served as a control.

**2.3    Sample collection and proteolysis**

Supernatants were collected 24 h after treatment, passed through medium equilibrated 0.2 µm
Millex-GP filter units (Merck Chemicals GmbH, Darmstadt, Germany), and transferred into 2 mL
Lo-Bind tubes (Eppendorf AG, Hamburg, Germany). Afterwards, cells were washed once with
DPBS. 200 µL RIPA buffer containing Roche cOmplete Mini Protease Inhibitor Cocktail (Merck
Chemicals GmbH) was applied directly into each well and cells were detached with a cell scraper.
Lysates were transferred into freshly prepared 1.5 mL Lo-Bind tubes (Eppendorf AG). Protein
concentration of the lysates was determined by Pierce BCA assay (Thermo Fisher Scientific). Ten µg
protein per lysate or 400 µL supernatant per sample were digested with Lys-C and trypsin using a
modified FASP procedure as described elsewhere (40, 41).

**2.4    LC-MS/MS and quantitative analysis**

LC-MSMS analysis was performed on a QExactive HF mass spectrometer (Thermo Fisher Scientific)
online coupled to a UItimate 3000 RSLC nano-HPLC (Dionex, Sunnyvale, USA). Samples were
automatically injected and loaded onto a C18 trap column for 5 min. Afterwards, samples were eluted
and separated on a C18 analytical column (Acquity UPLC M-Class HSS T3 Column, 1.8 µm, 75 µm
× 250 mm; Waters, Milford, USA). Samples were separated by a 95 min non-linear acetonitrile
gradient at a flow rate of 250 nl/min. Resolution of the MS spectra was recorded at 60,000 with an
AGC target of $3\times10^6$ and a maximum injection time of 50 ms from 300 to 1,500 m/z. The 10 most
abundant peptide ions were selected from the MS scan and fragmented via HCD. Thereby, the
normalized collision energy was 27 with an isolation window of 1.6 m/z, and a dynamic exclusion of
30 s. MS/MS spectra were recorded at a resolution of 15,000 with an AGC target of $10^5$ and a
maximum injection time of 50 ms. Spectra with unassigned charges, and charges of +1 and >8 were
excluded from the precursor selection.

The four datasets (lysates of pRMG and MIO-M1 cells, and secretomes of pRMG and MIO-M1
cells) were analyzed separately. The Proteome Discoverer 2.4 SP1 software (version 2.4.1.15;
Thermo Fisher Scientific) was used for peptide and protein identification via a database search
(Sequest HT search engine) against the SwissProt Human (MIO-M1) and Ensembl Pig (pRMG).
Database search was performed with full tryptic specificity, allowing for up to one missed tryptic
cleavage site, a precursor mass tolerance of 10 ppm, and fragment mass tolerance of 0.02 Da.
Carbamidomethylation of Cys was set as a static modification. Dynamic modifications included
deamidation of Asn and Gln, oxidation of Met, and a combination of Met loss with acetylation on
protein N-terminus. Peptide spectrum matches and peptides were validated with the Percolator
algorithm (42). Only the top-scoring hits for each spectrum were accepted with a false discovery rate
(FDR) <1% (high confidence). The final list of proteins satisfying the strict parsimony principle
included only protein groups passing an additional protein confidence filter FDR <5% filter
(target/decoy concatenated search validation).

Quantification of proteins, after precursor recalibration, was based on intensity values (at RT apex)
for all unique peptides per protein. Peptide abundances were normalized to the total peptide amount.
The protein abundances were calculated summing the normalised abundance values of corresponding
unique peptides. These protein abundances were used for calculation of enrichment ratios of proteins
in the stimulated samples to the untreated samples, resulting in single ratios for every quantified
protein in every treated sample. Significance of the ratios was tested using a background-based t-test
with correction for multiple testing according to Benjamini-Hochberg (adjusted *P*-value) (43, 44).

**2.5    Data analysis and visualization**

Calculation of the abundance ratio weight requires abundance values for both, the stimulated sample
and the control. However, if a protein is exclusively expressed in one of these samples, the Proteome
Discoverer software fails to calculate a respective abundance ratio weight. Since these extreme
values were of special interest to us, the missing abundance ratio weights were imputed using the R
package mice (version 3.13.0) and the "classification and regression trees" imputation method.

Ingenuity Pathway Analysis (IPA; Qiagen, Hilden, Germany) was used to analyze overrepresentation
of proteins in canonical pathways of the IPA library, as described elsewhere (45). IPA allows
deducing potential physiological effects of the various tested cytokines. Analysis was performed
based on the fold-change of the stimulated samples and the abundance ratio $P$-value. Fisher's exact
test allowed testing for nonrandom associations of proteins in the datasets and the different canonical
pathways (46). Furthermore, the method of Benjamini-Hochberg (B-H $P$-value) corrected for
multiple testing (44).

The euclidean distance for the heatmap analysis was calculated with the open source software Cluster
3.0 and hierarchically clustered by complete linkage clustering (47). The resulting heat map was
visualized with the open source software Java Treeview (version 1.2.0) (48).

**3    Results**

**3.1    Differential secretion of proteins after stimulation of Müller cells with various cytokines**

Müller cells are in close contact to all retinal cells, the vitreous and the blood vessels (17). To
address, whether this privileged position within the retina also translates into extensive signaling
between Müller cells and the surrounding cells, the secretomes of the human Müller cell-derived cell
line MIO-M1 and of pRMG were quantitatively analyzed by mass spectrometry after stimulation for
24h with the cytokines IFNγ, IL-10, IL-4, IL-6, TGFβ1, TGFβ2, TGFβ3, TNFα and VEGF,
respectively. By this means, we quantified 2,032 proteins in the supernatant of MIO-M1 cells
(Supplementary Table 1) and 3,093 proteins in the supernatant of pRMG across all treatment groups
(Supplementary Table 2).

**Figure 1.** Scatterplot of all identified proteins from MIO-M1 secretomes after treatment with
the indicated cytokines for 24 h (**A-I**). Proteins with significant changes in their abundance
(±log$_2$(1.5) fold expression, corrected p-value ≤0.05) were colored, with upregulated proteins
being depicted as yellow dots, while down-regulated proteins are colored cyan. Proteins with
significantly altered abundance in both, MIO-M1 and pRMG secretomes, are labeled with their
gene symbol. Keratins were excluded.

Fig. 1 and Supplementary Fig. S1 summarize changes in the secretome after treatment of Müller cells
with the different cytokines. A log$_2$ fold change of ±0.58 and a corrected p-value of equal or less than
0.05 served as cutoff to define significantly upregulated or downregulated genes, respectively.
Proteins equally regulated in MIO-M1 cells and pRMG were labeled with their gene symbol.
However, this was only possible for proteins with identical gene symbols in the human and the
porcine protein database. After treatment with IFNγ, 107 proteins in the secretome of MIO-M1 cells
and 176 proteins in the secretome of pRMG were significantly more abundant, while 67 proteins of
MIO-M1 cells and 96 proteins of pRMG were significantly less abundant in the supernatants (Fig.
1A; Suppl. Fig. S1A). Intriguingly, MIO-M1 cells and pRMG shared 21 upregulated and one
downregulated protein. Among these shared regulated proteins after treatment with IFNγ were many
with immune system functions, like signaling molecules (e.g. C-X-C Motif Chemokine Ligand 9
(CXCL9), CXCL10, IL-6) and components of the complement system (e.g. C1R, Serpin Family G
Member 1 (SERPING1)). Upon treatment with TNFα, 127 (MIO-M1) or 143 (pRMG) proteins were
more abundant and 57 (MIO-M1) or 87 (pRMG) proteins were less abundant in the supernatant (Fig.
1B, Suppl. Fig. S1B). Within these groups, MIO-M1 cells and pRMG shared 20 upregulated and
three downregulated proteins, again with many pro-inflammatory proteins like C-X3-C Motif

Chemokine Ligand 1 (CX3CL1), CXCL10, C-C Motif Chemokine Ligand 2 (CCL2), IL-6, and C1R
being upregulated. Thus, IFNγ and TNFα resulted in the most conserved changes of the secretome of
Müller cells when comparing between stimulated MIO-M1 cells and pRMG. In contrast, MIO-M1
cells and pRMG only shared ten proteins that were more abundant and three which were less
abundant after treatment with VEGF, with seven proteins being keratins (Fig. 1C, Suppl. Fig. S1C).
Treatment of Müller cells with the three different interleukins had a subtle influence on secreted
proteins with conserved regulation of seven (IL-4), nine (IL-6), and two (IL-10) proteins in the
secretome of MIO-M1 cells and pRMG (Fig. 1D-F, Suppl. Fig. S1D-F).

Finally, TGFβs led to pronounced alterations in the secretome of MIO-M1 cells. TGFβ1 enhanced
the secretion of 125 proteins while simultaneously reducing the abundance of 67 proteins, TGFβ2
increased the abundance of 131 proteins while impairing secretion of 69 proteins, and TGFβ3 raised
abundance of 135 proteins and reduced the abundance of 76 proteins (Fig. 1G-I). Furthermore, eleven
proteins of MIO-M1 cells and pRMG were similarly regulated by TGFβ1, eight upregulated and
three downregulated (Suppl. Fig. S1G). After treatment with TGFβ2, three proteins were more
abundant and four proteins less abundant in the secretome of both cell types (Suppl. Fig. S1H).
Additionally, the secretome of TGFβ3 treated MIO-M1 cells and pRMG shared one downregulated
protein and eight upregulated proteins, with most proteins like Osteonectin (Secreted Protein Acidic
And Rich In Cysteine; SPARC), Matrix Metallopeptidase 1 (MMP1), Biglycan (BGN) and various
keratins being functionally related to extracellular matrix organization (Suppl. Fig. S1I).
Furthermore, stimulation with TGFβ3 evoked upregulation of pro-inflammatory cytokine IFNγ in
both, the MIO-M1 cell line and pRMG. Overall, there are subtle, but intriguing differences in protein
abundances after stimulation of Müller cells with the different TGFβ isoforms. Some of the proteins
differentially induced by the TGFβ isoforms in MIO-M1 cells are pro-inflammatory cytokines, like
IFNγ, TNFα, and CCL2 (Supplementary Table 1).

**Figure 2.** Heatmap of hierarchical cluster analysis of proteins secreted by MIO-M1 cells after
treatment with various cytokines. Identified proteins were filtered for extracellular proteins
with significant changes in expression (±log$_2$(1.5) fold expression, corrected p-value ≤0.05).
Down-regulated proteins are presented in cyan, while up-regulated proteins are depicted yellow
for the respective treatments. Gray squares represent proteins that were neither identified in the

untreated control, nor in the respective treatment. Clusters were defined using the branches of a
dendrogram and shown as close up with the corresponding gene symbols.

Next, we aimed to examine the physiological functions of proteins secreted by Müller cells.
Therefore, we ordered secreted proteins in distinct clusters, grouped by similar expression patterns.
From the identified proteins of the MIO-M1 supernatant, all cytosolic contaminants were removed,
resulting in a list that exclusively contained extracellular proteins. Furthermore, only those proteins
that showed a significant change of abundance (corrected p-value ≤0.05 and ± 1.5-fold abundance) in
at least one treatment were selected for further analysis. As a result, a hierarchical heat map and an
associated dendrogram consisting of 171 proteins was generated using the $\log_2$ abundance ratio of
treated and untreated control cells (Suppl. Fig. S2). Furthermore, clusters were highlighted depending
on their position on the dendrogram (Fig. 2). To avoid single protein clusters, the proteins Polymeric
Immunoglobulin Receptor (PIGR) and Lacritin (LACRT) were assigned to Cluster F, despite being
on different branches of the dendrogram.

Notably, we identified proteins which were secreted by MIO-M1 cells exclusively after stimulation
(cluster A), while another set of proteins was completely absent in MIO-M1 cells (cluster B) treated
with VEGF. The largest cluster comprises proteins upregulated by at least one cytokine (cluster C).
In contrast to cluster D, consisting of proteins downregulated by TGFβs, TGFβs induce secretion of
proteins of cluster E. Finally, secretion of most proteins of cluster F is deterred by at least one
cytokine.

**A**

GO Enrichment: Cluster A

**B**

GO Enrichment: Cluster C

**C**

GO Enrichment: Cluster E

**Figure 3.** A shiny GO enrichment analysis for the previously defined cluster A (**A**), cluster C
(**B**) and cluster E (**C**) was performed. Redundant pathways were reduced to show only one
representative pathway. Depicted is the –log10(FDR) of the top 10 pathways for each cluster.
The red dashed line indicates the significance threshold.

The clusters A, C and E are large clusters with proteins upregulated by at least one cytokine, with
cluster A being the cluster with the most pronounced changes in secretion of its members and cluster
C being the most diffuse one. In order to deduce the physiological functions of the secreted proteins
of each of these clusters, we performed gene ontology (GO) analysis. Thereby, redundant pathways
were condensed to a single representative pathway and the top 10 pathways with the lowest
enrichment FDR were displayed. Intriguingly, the top 10 pathways for cluster A were related to

immune system processes, with "Humoral immune response" and "Immune system process" being
the most significant pathways with an enrichment FDR of $3.6 \times 10^{-9}$ and $6.9 \times 10^{-9}$, respectively (Fig.
3A). Furthermore, cluster C contained the pathway "Immune system process", while most other
pathways in this cluster C were involved in shaping the extracellular environment (Fig. 3B). Besides
further pathways involved in extracellular remodeling, the proteins of cluster E were also associated
to the pathway "AGE-RAGE signaling pathway in diabetic complications" (Fig. 3C).

**3.2    In-depth analysis of the Müller cell proteome after stimulation with a selection of**
282            **cytokines**

Our secretome analysis hints towards extensive signaling between Müller cells and their cellular
environment, differentially induced by various cytokines. To elucidate the underlying cellular
alterations, we also investigated differences in the proteome of MIO-M1 cells and pRMG cells by
mass spectrometry after treatment with these cytokines for 24 h. In total, 5,514 proteins were
quantified in the lysates of MIO-M1 cells (Suppl. Table 3) and 4,187 proteins in the lysates of pRMG
cells (Suppl. Table 4) across all treatment groups.

**Figure 4.** Scatterplot of all identified proteins from MIO-M1 lysates after treatment with the
291            indicated cytokines for 24 h (**A-I**). Proteins with significant changes in their abundance
292            ($\pm\log_2(1.5)$ fold expression, corrected p-value $\leq 0.05$) were colored, with upregulated proteins
293            being depicted as yellow dots, while down-regulated proteins are colored cyan. Proteins with

significantly altered abundance in both, MIO-M1 and pRMG lysates, are labeled with their
gene symbol. Keratins were excluded.

The threshold for significant abundance changes was set using the same cutoff values as for the
secretome, and equally regulated proteins in MIO-M1 cells and pRMG were labeled with their gene
symbol, if they shared the same gene symbol in the human and the porcine database (Fig. 4; Suppl
Fig. S3). Although the porcine protein database contains mostly humanized gene symbols, the swine
leukocyte antigens (SLA) genes show little sequence homology between the human and the porcine
genome and cannot properly be humanized (49). However, Human Leukocyte Antigen-C Alpha
Chain (HLA-C) was part of the porcine protein database.

Treatment of Müller cells with IFNγ resulted in 206 more abundant and 88 less abundant proteins in
MIO-M1 cells and 331 more abundant and 36 less abundant proteins in pRMG lysates (Fig. 4A;
Suppl Fig. S3A). Thereof, 29 proteins showed higher expression levels in both cells types. Among
the overlapping proteins were restriction factors like SAM And HD Domain Containing
Deoxynucleoside Triphosphate Triphosphohydrolase 1 (SAMHD1) and MX Dynamin Like GTPase
1 (MX1), transcription factors like Signal Transducer And Activator Of Transcription 1 (STAT1) and
STAT2, regulators of protein homeostasis like Leucine Aminopeptidase 3 (LAP3) or the Proteasome
20S Subunit Beta 9 (PSMB9), and proteins associated with peptide transport and antigen presentation
like Transporter 2, ATP Binding Cassette Subfamily B Member (TAP2), TAP Binding Protein
(TAPBP), Beta-2-Microglobulin (B2M), as well as HLA-C. After treatment with TNFα, 204 proteins
were more abundant in the proteome of MIO-M1 cells, while 119 proteins were less abundant (Fig.
4B). In pRMG, 207 proteins with higher abundance and 285 proteins with lower abundance were
identified upon treatment with TNFα, with 18 proteins that were differentially regulated in both cell
types (Suppl. Fig. S3B). Among shared proteins that were more abundant after treatment with TNFα
were pro-inflammatory proteins like B2M and Nuclear Factor Kappa B Subunit 2 (NFKB2), or
adhesion molecules like Intercellular Adhesion Molecule 1 (ICAM1) or Vascular Cell Adhesion
Molecule 1 (VCAM1). VEGF led to 143 more and 102 less abundant proteins in MIO-M1 cells or
232 more and 224 less abundant proteins in pRMG, respectively (Fig. 4C; Suppl. Fig. S3C). Thereof,
MIO-M1 cells and pRMG shared nine more abundant proteins, inter alia proteins associated with
reorganization of the cortical cytoskeleton like Alpha-Actin-1 (ACTA1) or HCLS1 Associated
Protein X-1 (HAX1), and two less abundant proteins, Thymosin Beta 10 (TMSB10) and Thymosin
Beta 4 X-Linked (TMSB4X), both inhibitors of actin polymerization. Upon treatment with
interleukins IL-4, IL-6 and IL-10, the Müller cell proteomes mirrored the subtle effects of these
cytokines on the abundance of proteins observed for the Müller cell secretomes (Fig. 4D-F, Suppl.
Fig. S3D-F). Also in line with the secretome data, the overlap between differentially abundant
proteins of the MIO-M1 and pRMG proteome after treatment with the various interleukins contained
only few proteins. In contrast, TGFβ1 increased the abundance of 143 proteins, while decreasing the
abundance of 94 proteins in the proteome of MIO-M1 cells and increased the abundance of 203
proteins, while decreasing the abundance of 103 proteins in the proteome of pRMG (Fig. 4G; Suppl.
Fig. S3G). In comparison to the lower abundant proteins Phosphodiesterase 5A (PDE5A) and
Inhibitor Of Nuclear Factor Kappa B Kinase Subunit Beta (IKBKB), the proteins Collagen Type I
Alpha 1 Chain (COL1A1), Insulin Like Growth Factor Binding Protein 7 (IGFBP7), JunB Proto-
Oncogene (JUNB), and 2-Hydroxyacyl-CoA Lyase 1 (HACL1) were more abundant in both, MIO-
M1 cells and pRMG after treatment with TGFβ1. Following treatment with TGFβ2, 125 proteins of
the proteome of MIO-M1 cells and 266 proteins of the proteome of pRMG were more abundant,
whereas 67 proteins of the MIO-M1 proteome and 229 proteins of the pRMG proteome were less
abundantly expressed (Fig. 4H; Suppl. Fig. S3H). In the case of treatment with TGFβ3, 130 proteins
in the MIO-M1 proteome and 185 in the pRMG proteome showed higher abundances, while 94

proteins in MIO-M1 proteome and 250 in the pRMG proteome were less abundant (Fig. 4I; Suppl.
Fig. S3I). The overlap of MIO-M1 cells and pRMG treated with TGFβ2 comprised three proteins,
and treatment with TGFβ3 resulted in an overlap of seven proteins. Overall, pRMG reacted more
pronounced to treatment with the various cytokines compared to MIO-M1 cells.

**3.3  Canonical pathways enriched in Müller cells upon stimulation**

Treatment with various cytokines partly induced pronounced changes in the secretome and proteome
of Müller cells. In the secretome, these changes primarily included the secretion of pro-inflammatory
cytokines and proteins associated with organization of the extracellular matrix. To elucidate
overrepresented mechanisms and pathways in stimulated Müller cells, we performed Ingenuity
pathway analysis (IPA). We limited the IPA to significantly regulated proteins (p-value $\leq 0.05$)
identified in the MIO-M1 and pRMG lysates. Since IPA cannot handle porcine gene symbols, we
replaced the only canonical SLA gene SLA-1 in our pRMG dataset by the canonical human HLA
gene HLA-A. Hence, an IPA core analysis was performed with 1,543 proteins for pRMG and with
2,262 proteins for the MIO-M1 cells. IPA identified 338 canonical pathways within the proteome of
MIO-M1 cells and 218 canonical pathways in the proteome of pRMG that were significantly
enriched by at least one of the used cytokines (IPA p-value $\leq 0.05$; Suppl. Table 5). Among the
identified canonical pathways were many pathways associated with signaling, cell death, immune
system processes and the cellular redox state (Fig. 5; Suppl. Fig. S4). A selection of canonical
pathways enriched in both pRMG and MIO-M1 cells after treatment with at least one of the tested
cytokines is depicted in Fig. 5.

**Figure 5.** A comparative IPA analysis with the significantly regulated proteins identified in the pRMG lysates after stimulation with the indicated cytokines was performed. Canonical pathways related to signaling, cell death, immune system processes and oxidative stress were selected. Pathways with significant enrichment of genes after stimulation with at least one cytokine are presented. Significance of the gene enrichment for each pathway and treatment is indicated by purple squares in the left array. Thereby, treatments that did not meet the

significance threshold (p-value ≤ 0.05) are marked with a dot. The z-score is indicated in the
right array and represents a prediction of activation (orange) or inhibition (blue) of the
pathway. Gray squares mark treatments where the activation state of a pathway could not be
calculated.

Among the most significantly regulated pathways in MIO-M1 cells and pRMG were the canonical
pathways "Mitochondrial Dysfunction" and "Oxidative Phosphorylation". These pathways were
significantly induced by all examined cytokines in pRMG. The pathways "Ferroptosis Signaling
Pathway", "iNOS Signaling", "NRF2-mediated Oxidative Stress Response", and "Production of
Nitric Oxide and Reactive Oxygen Species in Macrophages" are closely linked to the cellular redox
state and were among the enriched pathways in MIO-M1 cells and pRMG after treatment with
various cytokines. Furthermore, proteins associated with the maturation of phagosomes were
significantly enriched in pRMG. In line with this, "Caveolar-mediated Endocytosis Signaling" was
significantly enriched in pRMG after treatment with IFNγ, TGFβ1, TNFα and VEGF, and "Clathrin-
mediated Endocytosis Signaling" was significantly enriched in pRMG after treatment with all
cytokines except IL-10. Besides these two pathways associated with the recycling of the extracellular
environment, intracellular protein homeostasis and MHC class I peptide generation was facilitated by
enrichment of the "Protein Ubiquitination Pathway" in pRMG after treatment with IFNγ, TGFβ3,
TNFα and VEGF. Similarly, IFNγ, IL-4, TGFβ1, TGFβ3, TNFα and VEGF enriched "Protein
Ubiquitination Signaling" in MIO-M1 cells. "Neuroinflammation Signaling" was induced by IFNγ,
TNFα and VEGF in MIO-M1 cells and by IFNγ, TGFβ1, TGFβ3 and TNFα in pRMG, whereas
TGFβ2 and VEGF led to a slight inhibition of this pathway in pRMG.

**3.4    A deeper look into Müller cell complement secretion upon cytokine stimulation**

Because the enrichment analysis of the secretome yielded highly significant hits such as "humoral
immune response" and "immune system process," we took a closer look at complement proteins in
the secretome and cell lysates. Notably, most complement proteins are secreted as key components of
the humoral immune system. The identified complement components include central complement
proteins, regulators, and receptors. Consistent with their localization in the cell membrane, the latter
(including ITGAM, ITGB2, C5aR1) were detected only in cell lysates, and here specifically in those
of pRMGs. The complement regulators clusterin (CLU), vitronectin (VTN), CD59, and SERPING
were found in most test samples. With regard to the central complement components, the pRMG
secretome took a prominent position and showed results for complement components for all three
different activation pathways (e.g. C1q, FD, MASP1) and the terminal pathway (e.g. C9). The central
complement protein C3 was found in both the MIO-M1 and pRMG secretomes and in the MIO-M1
lysate. Interestingly, cytokine treatment induced changes in complement proteins and regulators but
had no effect on complement receptor expression. We observed that C1q subunits, which initiate the
classical complement pathway by binding to antibodies, were detectable only in pRMG but not in
MIO-M1 cells. C1q levels in cell lysates and the corresponding secretome were consistently reduced
after TNFα treatment but were increased by IFNγ. Moreover, complement proteases C1r and C1s,
which bind to C1q therewith continuing the cascade of classical pathway activation, were enriched in
the supernatants of MIO-M1 and pRMG cells treated with IFNγ (Fig. 2C). In contrast, C1r
concentration was significantly decrease in supernatants of MIO-M1 cells but not pRMG after VEGF
and TGFβ2 application. Notably, C1s and C1r were not detected in cellular lysates. Interestingly, the
abundance of the central complement proteins C3 and C4A were modified by the supplemented
cytokines in MIO-M1 secretomes only and not in any other data set (Fig. 2C). These proteins are
cleaved upon complement activation as for example triggered by the C1q-mediated classical pathway
and result in cleaved products which interact with cellular receptors (e.g. C3a/C3b, C4a). Here,

complement protein C3 is mainly increased following TNFα addition and C4 upon exposure to IFNγ
(Fig. 2C). In fact, IFNγ was also the major player modulating the secreted complement components
in pRMG: C2 and FI were significantly increased while C9, FD and MASP1 were clearly reduced in
its presence. These complement components absent from any other sample.

Regarding the complement regulators factor H (FH), SERPING and CLU are of interest. Secretion of
FH was not observed in untreated MIO-M1 and pRMG, but it was significantly upregulated in MIO-
M1 secretomes following IFNγ, TNFα, TGFβ3, and VEGF treatment (Fig. 2A). Similar results were
obtained for SERPING, whose levels were increased by IFNγ in the MIO-M1 secretome, pRMG cell
lysates and secretome. Remarkably, the MIO-M1 lysate showed decreased values for CLU following
IFNγ, TGFβ1 and TNFα, and similar but not significant trends was observed for the respective
secretome. Finally, while CLU was upregulated in pRMGs lysates upon IL-6 or VEGF treatment, no
significant alterations could be found in corresponding secretomes.

In summary, IFNγ and TNFα seemed to be the most effective cytokines to modulate the Müller cell
complement expression and secretion (Fig. 2).

**3.5   Müller cells as non-professional antigen-presenting cells**

Intriguingly, treatment of pRMG with IFNγ, TGFβ2, TGFβ3 and TNFα significantly enriched
proteins associated with the "Antigen Presentation Pathway". Likewise, the "Antigen Presentation
Pathway" was induced in MIO-M1 cells by treatment with IFNγ, TGFβ1, TNFα and VEGF. Thereby,
antigen presentation is an umbrella term for two distinct processes. MHC class I antigen presentation
is common to all nucleated cells and allows CD8[+] cytotoxic T cells (CTL) to assess whether cells are
infected with an intracellular pathogen (50, 51). In contrast, MHC class II is presented to antigen
specific CD4[+] T cells mainly by professional antigen-presenting cells inducing their activation and
differentiation to T helper cells (52).

**Figure 6.** Heatmap of hierarchical cluster analysis of proteins involved in MHC class I (upper panel) and MHC class II (lower panel) antigen presentation expressed by MIO-M1 cells (**A**) and pRMG cells (**B**) after treatment with various cytokines. Down-regulated proteins are presented in cyan, while up-regulated proteins are depicted yellow for the respective treatments. Gray squares represent proteins that were neither identified in the untreated control, nor in the respective treatment. The heatmap was generated on the basis of the log₂ fold change of the respective proteins.

To investigate the antigen presentation capacity of Müller cells, we constructed a hierarchical heatmap for MIO-M1 cells (Fig. 6A) and pRMG (Fig. 6B) challenged with various cytokines. Proteins linked to antigen presentation were selected and clustered hierarchically. Proteins associated with MHC class I antigen presentation are displayed in the upper panel and proteins correlated to MHC class II antigen presentation are depicted in the lower panel. In MIO-M1 cells, IFNγ induced the majority of proteins linked to both, MHC class I and II antigen presentation, whereas TNFα exerted its inductive effect exclusively on MHC class I antigen presentation. The only protein linked to MHC class II antigen presentation induced by TNFα in MIO-M1 cells was Cathepsin S (CTSS). Furthermore, the other cytokines did not induce proteins related to antigen presentation in MIO-M1 cells. Quite the contrary, TGFβ2 and TGFβ3 reduced the abundance of proteins linked to MHC class

I antigen presentation in these cells. In contrast to MIO-M1 cells, pRMG reacted to all tested
cytokines by induction of components of both MHC class I and II antigen presentation to varying
degrees. IFNγ and TNFα induced proteins of MHC class I and II antigen presentation in pRMG,
among others SLA-DQA1, SLA-DQB1, SLA-DRA and SLA-DRB1. Furthermore, class I and II
antigen presentation was upregulated by TGFβ isoforms 1-3 in pRMG. While TGFβ2 and TGFβ3
induced the components of the MHC class I peptide loading complex TAP2 and TAPBP, TGFβ1 also
increased the abundance of HLA-DMA and HLA-DMB, proteins involved in the peptide loading on
MHC class II. We saw only a subtle induction of proteins related to antigen presentation by IL-10,
IL-4 and IL-6. The smallest impact on antigen presentation proteins was seen after stimulation with
IL-6. Furthermore, IFNγ significantly upregulated the expression of the co-stimulatory molecule
CD40 in pRMG, while TGFβ2, TGFβ3, TNFα and VEGF resulted in lower abundance of CD40
(Suppl. Table 4).

**Figure 7.** IPA for pRMG cells after treatment with IFNγ was performed. Depicted is the
canonical antigen presentation pathway. Yellow proteins are induced. The intensity of the
yellow color indicates the degree of upregulation. Grey proteins are in the dataset but did not

pass the analysis cutoffs of the pathway analysis. Purple circles highlight proteins (single
circle) or complexes (double circles).

Induction of the canonical MHC class I and MHC class II antigen presentation pathway as assessed
by IPA for pRMG after treatment with IFNγ is summarized in Fig. 7. This pathway was enriched in
pRMG cells with a p-value of $4.15 \times 10^{-12}$. Besides MHC class I and MHC class II, also components
of the peptide loading complex of MHC class I (TAP2 and TPN) were upregulated in pRMG after
IFNγ treatment. Furthermore, IFNγ induces the Large Multifunctional Peptidase 2 (LMP2;
synonymous to PSMB9) and Large Multifunctional Peptidase 7 (LMP7; synonymous to PSMB8)
subunits of the immunoproteasome in MIO-M1 cells.

**4    Discussion**

DR is a condition associated with microvascular degeneration, resulting in ocular inflammation and
eventually in complete blindness (53). As the prevalence of diabetes in developed and developing
countries rises, DR became the leading cause of blindness in the working-age population (54).
Microglia are resident innate immune cells in the retina and as such, they are considered to regulate
the inflammation during DR (55). In contrast, the role of Müller cells during the pathology of DR has
been often described to be the metabolic support of neurons (53) . However, growing evidence
suggests excessive signaling between Müller cells and the surrounding tissue including microglia,
with both beneficial and detrimental consequences for the pathogenesis of DR (56). We show here
that stimulation of Müller cells with pro-inflammatory cytokines like IFNγ or TNFα, but also with
growth factors like TGFβ or VEGF resulted in profound, yet distinct response profiles in their
secretomes, thus confirming their central role in cell-to-cell communication within the retina.

Previously, it has been shown that the conditional knock-out of VEGF in Müller cells of mice
reduced the expression of inflammatory markers like TNFα or ICAM1 compared to diabetic control
mice, indicating a pro-inflammatory effect of VEGF on Müller cells (57, 58). Here we could show
that VEGF induced the Müller cell line MIO-M1 to secrete proteins associated with immune effector
processes like IFNγ, B2M, HRNR, and complement-associated proteins like SERPING1 and
complement factor H (FH). Furthermore, ICAM1 was more abundant in our proteome data of MIO-
M1 cells and pRMG treated with VEGF compared to the respective control cells. Thus, we were able
to confirm the pro-inflammatory role of VEGF in Müller cells.

Overall, we observed that MIO-M1 cells secreted TNFα upon stimulation with VEGF, IL-6, and the
TGFβ isoforms 2 and 3. Increased levels of TNFα were previously identified in the vitreous of DR
patients (59). Furthermore, Müller cells have been reported to secrete increased amounts of VEGF
and TNFα under stress conditions like inflammation or hyperglycemia, thereby promoting retinal
inflammation (57, 60-64). Our results show that TNFα induced MIO-M1 cells to secrete
inflammatory proteins but also proteins associated with tissue development like Leukemia Inhibitory
Factor (LIF), Follistation (FST), Neuregulin 1 (NRG1), or Pleiotrophin (PTN), to name but a few.
Thus, our data suggest a pleotropic role of TNFα in Müller cells.

TGFβs regulate early embryogenesis, maintenance and regeneration of mature tissues, and various
disease processes (65). TGFβ1 has been described to be anti-inflammatory, inducing immune
tolerance in the periphery and preventing autoimmunity. Thus, knock out of TGFβ1 in mice induced
autoantibodies and multifocal inflammatory disease in many tissues, eventually causing the mice to
die within 4 weeks of age (66). In our *in vitro* model, treatment of MIO-M1 cells with TGFβ1
induced secretion of CCL2 and therefore might result in the recruitment of leukocytes.

Simultaneously, TGFβ1 enhanced the secretion of anti-inflammatory proteins like TNF Alpha
Induced Protein 6 (TNFAIP6) or LIF in Müller cells. In contrast, TGFβ2 and TGFβ3 also promoted
immune system processes in MIO-M1 cells by inducing secretion of pro-inflammatory cytokines
TNFα, CXCL8, CXCL10 and IFNγ. It has been shown that TGFβ2 acts immunomodulatory in
aqueous humor by decreasing the expression of IL-6, CXCL1, CCL2, G-CSF and IGFBP-5 (67).
However, we demonstrated that treatment of Müller cells with TGFβ2 induced expression of IL-6
and CCL2, suggesting a complex and diverse role of TGFβ2 signaling in the eye. Although TGFβ
isoforms are closely related (sharing 71-79% sequence identity), have similar three-dimensional
structures and signal canonically through the same receptor (68), our secretome analyses provide
evidence that they affect Müller cells differentially. While knock-out of the TGFβ isoforms resulted
in different phenotypes, all null mutations were lethal (69-72). Replacement of the coding sequences
by one of the other isoforms only partially rescued these phenotypes, implying intrinsic differences
of these isoforms (73, 74). These differences might originate from varying flexibilities of an
interfacial helix and thus modify the binding of the TGFβ isoforms to matrix proteins as suggested by
Huang and colleagues (68). Latent TGFβ binds Thrombospondin 1 (THBS1), augmenting its
activation (75-77). Intriguingly, in our study, all three TGFβs elevate the abundance of THBS1
indicating a positive feedback loop. Recently, transcriptome analysis linked the expression of TGFβ
isoforms of mice to activation of different signaling cascades. While TGFβ1 and TGFβ2 evoked the
non-canonical p38MAPK signaling pathway, which has been linked to gliosis, TGFβ3 induced the
SMAD canonical signaling pathway for TGFβs in mice (78, 79). During glaucoma, the aqueous
humor contains elevated levels of TGFβ2, exceeding levels for homeostatic signaling (77, 80).
Thereby, TGFβ2 has been associated with pathological remodeling of the trabecular meshwork and
the optical nerve head (77). Furthermore, it stimulated secretion of extracellular matrix proteins by
astrocytes and cells of the lamina cribrosa (81-84). Our data suggest that Müller cells also contribute
to the remodeling of the extracellular matrix as stimulation with TGFβ1 and TGFβ3 resulted in
enhanced secretion of extracellular matrix proteins like Fibrillin-2 (FBN2), various keratins, and
collagens. Furthermore, they simultaneously enhanced the turnover of extracellular matrix by
inducing the secretion of the Matrix Metalloprotease-1 (MMP1) and MMP2. Previously, MMP2 has
been described to be elevated in DR patients and to promote the pathogenesis of DR by inducing
mitochondrial dysfunction and apoptosis of retinal capillary cells (85). Intriguingly, our IPA also
indicates that the canonical pathway for mitochondrial dysfunction is enriched in Müller cells upon
stimulation with all TGFβ isoforms. However, this is not solely due to the enhanced secretion of
MMP2, since mitochondrial dysfunction is enriched in Müller cells by all tested cytokines, whereas
TGFβs exclusively induce MMP2.

A meta-analysis of eleven studies recently showed significantly increased levels of the anti-
inflammatory cytokine IL-10 in the vitreous humor of DR patients (86). Moreover, IL-10 levels were
higher in patients with proliferative DR compared to patients with non-proliferative DR (86). While
our data show no contribution of Müller cells to elevated IL-10 secretion upon stimulation with the
tested cytokines, IL-10 induced proteins associated with tissue development, cell adhesion, and
angiogenesis in MIO-M1 cells, namely Corneodesmosin (CDSN), LIF, PTN, and various collagens,
to name a few. In line with this, it has been shown that IL-10 is involved in the pathological
angiogenesis during the development by modulating the macrophage response to hypoxia (87). Thus,
we hypothesize that IL-10 might also be involved in the abnormal angiogenesis during DR.

Previously, it has been shown that the canonically anti-inflammatory IL-4 can potentiate cytokine
and chemokine production in macrophages following a pro-inflammatory stimulus (88-90). In line
with this, we observed increased abundances of many pro-inflammatory proteins like IFNγ, S100A9,
S100A7, CXCL10, and Lysozyme C (LYZ) upon stimulation of MIO-M1 cells with IL-4, indicating

that IL-4 does not exert an anti- but a pro-inflammatory influence in these cells. Interestingly, MIO-
M1 cells did not secrete the anti-inflammatory interleukin IL-4 upon treatment with the various tested
cytokines. Secretion of pro-inflammatory IL-6 by Müller cells has been described upon treatment
with IL-1β or LPS (91). Furthermore, elevated levels of IL-6 were found in the vitreous and serum of
DR patients and even further increased in patients suffering from proliferative DR (92). Here we
show that IFNγ, TNFα, TGFβ2, and TGFβ3 also induced the secretion of IL-6 in MIO-M1 cells.
Previously, it was shown that IL1β induced IL-6 through activation of the p38MAPK signaling
pathway (93). The murine TGFβ isoforms 1 and 2 also activated the p38MAPK signaling in Müller
cells of mice (78). Thus, the involvement of p38MAPK signaling in secretion of IL-6 by Müller cells
should be addressed in further studies. Brandon and colleagues demonstrated induction of VEGF by
IL-6 in Müller cells, especially under hyperglycemic conditions, preventing Müller cells from
glucose toxicity (94). In contrast, our analysis revealed no VEGF secretion by MIO-M1 cells upon
IL-6 treatment. High glucose concentrations of 25 mM potentiated the induction of VEGF by IL-6
(94). However, by default the standard culture medium used in our study has a D-glucose
concentration of 25 mM. Thus, an adaption of MIO-M1 cells to these conditions might have occurred
negating the induction of VEGF by IL-6.

Elevated levels of IFNγ in the vitreous of DR patients have been described previously (95, 96). In our
study, we could observe distinctly elevated INFγ secretion only after stimulation with INFγ.
Treatment with IFNγ furthermore induced the secretion of CXCL9, CXCL10, IL-6 and complement
subcomponent C1R in MIO-M1 cells and pRMG.

Interestingly, we observed an induction of CX3CL1 in the whole-cell lysates and in the secretomes.
Specifically, except for TGFβ1 and TGFβ2, all stimulants used in this study induced expression of
CX3CL1 in the cell proteome of MIO-M1 cells, while only IFNγ and TNFα treatment resulted in
significantly higher abundance of CX3CL1 in the secretome as well. CX3CL1 is a membrane-bound
chemokine, which functions as an adhesion molecule for leukocytes, but can also be proteolytically
cleaved, resulting in a soluble form with chemotactic function (97, 98). Several studies already
demonstrated the implication of CX3CL1 in DR pathogenesis as the vitreous of PDR patients
contains elevated levels of CX3CL1 (99) Furthermore, circulating CD11b+ leukocytes that are
involved in leukostasis in DR express higher levels of CX3CR1 in diabetic mice compared to
controls (100). Thus, membrane-bound CX3CL1 on the surface of Müller cells might be involved in
leukostasis in DR. Since TNFα levels are elevated in the eyes of DR patients, this might result in the
upregulation of CX3CL1 in diabetic Müller cells (101). In a previous study, incubation of microglia
with Müller cell supernatant containing secreted CX3CL1 resulted in the upregulation of the
respective receptor CX3CR1 in microglia. The authors proposed that Müller cells might be able to
promote microglial motility via the chemotactic effect of CX3CL1 (102). Thus, secretion of CX3CL1
from Müller cells might contribute to DR pathogenesis by recruiting peripheral inflammatory cells
and microglia. Upregulation of CX3CL1 after treatment with TNFα has been demonstrated in aortic
endothelial cells, but could not be observed in retinal endothelial cells (103). Here, we demonstrate
that stimulation with TNFα elicits secretion of CX3CL1 in pRMG and MIO-M1 cells and expression
of CX3CL1 in MIO-M1 cells. Thus, we lay the ground for further research concerning the expression
and regulation of CX3CL1 in Müller cells during DR.

Another route of communication between microglia/macrophages and Müller cells could be local
changes in complement expression. We recently demonstrated, that Müller cells in the mouse retina
are the main producers of complement components of the classical (C1s, C4), the alternative pathway
(FB) and C3 as the central component to all pathways under homeostatic, but also under ischemic
stress conditions (104). In the retina, it is the microglia that by far express highest levels of

complement receptors including ITGAM (alias CD11b), C3aR, C5aR1 and C5aR2 (104). In our
present study, TNFα and INFγ triggerd the most prominent effects on complement expression
consistently in MIO-M1 and pRMG. Given that *in vivo* microglia and potentially other immune cell
serve as major source of TNFα and INFγ (105) (resource data from Cowan et al., 2020 (106) at
https://data.iob.ch), the strong effect of these cytokines on Müller cell could be central to coordinate
the tissue immune homeostasis in pathologies. In this context, the enhanced expression of activating
complement components by Müller glia could serve as feedback mechanisms towards microglia in
turn modulating their activation profile. Taken together, our results point towards a pro-inflammatory
phenotype of Müller cells, which is in line with a previous study, where we analyzed the surfaceome
of primary equine Müller cells and MIO-M1 cells after stimulation with Lipopolysaccharide (30).
While the surfaceome of Müller cells in this previous study revealed expression of MHC class I and
II as well as costimulatory molecules especially in primary equine Müller cells, we could now further
complement these results with LC-MS/MS-analyses of whole cell lysates and of cell supernatants,
confirming an antigen-presenting phenotype of Müller cells. While stimulation with LPS resulted in
enhanced expression of MHC class II molecules in primary equine Müller cells, no MHC class II
molecules could be identified in MIO-M1 cells upon LPS treatment (30). In contrast, stimulation
with IFNγ in this study induced the expression of proteins that are associated with both MHC-I and
MHC-II antigen presentation in pRMG as well as in MIO-M1 cells. This is in accordance with an
early study that demonstrated the induction of MHC class I and MHC class II molecules in primary
human Müller cells by IFNγ *in vitro* (107). In contrast to MIO-M1 cells, pRMG also showed basal
expression of MHC class II without stimulation. Thus, our data demonstrate that Müller cells exhibit
several criteria for atypical antigen-presenting cells (108). While microglial cells are the predominant
immune cells of the retina, our data confirm that Müller cells are crucially involved in immunological
processes in the retina as well, as they possess an antigen processing and presenting machinery and
secrete pro-inflammatory cytokines (14). We have previously shown that the cultivation of primary
porcine Müller cells under hyperglycemic conditions resulted in higher expression levels of MHC
class II molecules, pointing towards an immunologically activated state of Müller cells in DR (36).

Besides expression of MHC class I and II molecules, we demonstrated that stimulation of Müller
cells with various cytokines resulted in the enrichment of proteins and pathways that are associated
with the formation and maturation of phagosomes. Previously, Müller cells have been described to be
phagocytic cells, capable of phagocytosing cell debris, dead photoreceptor cells and even bacteria
(109-111). Our IPA showed that proteins of phagocytosis pathways in Müller cells are induced upon
stimulation with various cytokines. Furthermore, phagocytosis is not only clathrin- but also caveolar-
mediated. Since our data showed enrichment of phagocytic pathways, as well as the canonical
antigen presentation pathway, it is possible that Müller cells present exogenous peptides on MHC
class II to CD4+ T helper cells. Intriguingly, phagocytosis of dead photoreceptors would also allow
Müller cells to present proteins expressed by photoreceptors on MHC class II, and via cross-
presentation on MHC class I (112, 113). The relationship of HLA alleles and haplotypes to the
progression of diabetic retinopathy is part of an ongoing scientific debate (114-122). However, the
rigorous pro-inflammatory signaling of Müller cells paired to potential activation of CD4+ and CD8+
T cells in the context of DR should be elaborated in further studies. Towards this goal it should be
addressed, whether Müller cells are sufficient to stimulate alloreactive naïve T cells or memory T
cells (108).

Furthermore, our proteomic analysis revealed significantly higher abundance of the costimulatory
molecule CD40 in pRMG after stimulation with IFNγ. Intriguingly, CD40 has been shown to play a
vital role in DR pathogenesis (123-125). A study with transgenic mice, which express CD40 only in
Müller cells, demonstrated that the upregulation of CD40 in Müller cells under diabetic conditions is

sufficient for thriving leukostasis and degeneration of capillaries, thus promoting retinal
inflammation and neurodegeneration in streptozotocin induced DR (124). Expression of CD40 has
also been demonstrated in primary human Müller cells and its ligation with CD154 resulted in
increased expression of ICAM-1 in Müller cells (123). Furthermore, it was shown that activation of
Müller cells through CD40 leads to an enhanced production of TNFα and IL-1β in bystander
microglia (124). Since several studies revealed an association between elevated levels of IFNγ and
DR, and stimulation of CNS astrocytes led to expression of CD40, IFNγ in DR might lead to the
upregulation of CD40 in Müller cells and thus promote chronic inflammation (126-129). Here, we
provide new insight in CD40 signaling in Müller cells, as IFNγ increased the expression of CD40,
whereas TGFβ2, TGFβ3, TNFα and VEGF resulted in lower abundance of CD40 in pRMG. Future
studies are needed to further investigate this issue, as the CD40 signaling pathway constitutes a
possibility for therapeutical intervention (130). In contrast to the porcine dataset, we could not
identify CD40 in the MIO-M1 cells. As CD40 expression has already been shown in primary human
Müller cells, our result might be due to the dedifferentiation of immortalized cells in culture (123).

Oxidative stress and reactive oxygen species (ROS) are known to play a central role during the
pathogenesis of DR (131). Rat-derived Müller cells under hyperglycemic conditions developed
mitochondrial dysfunction and oxidative stress, causing swelling and eventually apoptosis of the cells
(132, 133). Mitochondrial dysfunction can lead to ROS production, which then promotes
inflammatory response by activation of NF-κB and release of pro-inflammatory cytokines (134, 135).
Our analysis revealed that proteins associated with mitochondrial dysfunction were enriched after
treatment of pRMG with all tested cytokines. Furthermore, two significantly enriched pathways in
our data sets are associated with reactive oxygen species, namely "NRF2 mediated Oxidative Stress
Response" and "Production of Nitric Oxide and Reactive Oxygen Species in Macrophages".
Intriguingly, Müller cells have previously been found to regulate the ROS levels via Nrf2 and to be
more resistant to ROS formation compared to photoreceptor cells or bipolar cells (41, 136). In line
with this, we showed that treatment with IL-4, TGFβ2, TGFβ3, TNFα and VEGF inhibited death
receptor signaling pRMG. Phagocytic cells often produce ROS to protect themselves from pathogens
(137). Furthermore, macrophages stabilize cytosolic Nrf2 to be more resistant against ROS (138).
Since Müller cells have been shown to by phagocytic, we propose that induction of ROS in these
cells also serves as a defense mechanism (109-111).

In this study, we used both primary porcine Müller cells and the spontaneously immortalized human
Müller cell line MIO-M1 that was established and described in the Moorfields Institute of
Ophthalmology 2002 (39). Although MIO-M1 cells largely retain the phenotypic and functional
characteristics of Müller cells *in vitro*, previous studies documented some characteristics of neuronal
stem cells and expression of postmitotic neuronal cell markers in MIO-M1 cells (39, 139, 140).
Furthermore, it has been shown that Müller cells tend to adapt to cell culture conditions by reduced
secretion of neurotrophic factors (141-143). Our analysis revealed many proteins commonly
regulated in pRMG and MIO-M1 cells, as well as pathways similarly enriched and regulated in both
cells. However, we also observed effects specific for either MIO-M1 cells or pRMG. Thus, for
example our IPA revealed that MIO-M1 cells decrease the activity of the significantly enriched
canonical pathway oxidative phosphorylation, while it is amongst the most induced pathways of
pRMG following treatment with the majority of tested cytokines. While these differences might
originate from different adaption of MIO-M1 cells and pRMG to the cell culture conditions or the
immortalization of MIO-M1 cells, it might as well be possible that they originate from differences in
human and porcine Müller cells. Further studies should elaborate similarities and differences between
freshly prepared primary human and porcine Müller cells.

Taken together, our in-depth proteomic profiling of MIO-M1 cells and pRMG revealed the capacity
of Müller cells to react in a differentiated manner upon treatment with various growth factors and
cytokines. Furthermore, we demonstrated a primarily pro-inflammatory phenotype of Müller cells, as
they secreted a variety of pro-inflammatory cytokines, complement components and upregulated
proteins associated with antigen processing and presentation, suggesting a function as atypical
antigen-presenting cell in the course of retinal inflammation. Furthermore, we observed enrichment
of proteins connected to mitochondrial dysfunction, as well as proteins related to the formation and
maturation of phagosomes. Thus, Müller cells are capable of modulating immune responses in the
retina, and may significantly contribute to chronic inflammation during DR.

**5    Conflict of Interest**

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

**6    Author Contributions**

Conceptualization, S.M.H. and C.A.D.; Formal analysis, S.M.H.; Funding acquisition, S.M.H. and
C.A.D.; Investigation, A.S., L.L., C.A.D., and S.M.H.; Project administration, S.M.H.; Supervision,
S.M.H.; Visualization, A.S.; Writing - original draft, A.S., L.L., A.G., and D.P.; Writing - review and
editing, C.A.D., and S.M.H. All authors have read and agreed to the published version of the
manuscript.

**7    Funding**

This work was funded by Deutsche Forschungsgemeinschaft in the SPP 2127, grant numbers DFG
DE 719/7-1 to C.A.D., HA 6014/5-1 to S.M.H., GR 4403/5-1 to A.G. and PA 1844/3-1 to D.P.

**8    Acknowledgments**

The authors would like to thank the butcher's shop "Wirtshaus Metzgerei Adolf Sandbiller" in D-
94336 Hunderdorf, Germany for providing porcine eye samples as well as Juliane Merl-Pham for her
valuable advice.

**9    Data Availability Statement**

All data are contained within the article or in supplementary materials.

## 10    References

1.      Duncan JL, Pierce EA, Laster AM, Daiger SP, Birch DG, Ash JD, et al. Inherited Retinal Degenerations: Current Landscape and Knowledge Gaps. Transl Vis Sci Technol. 2018;7(4):6.

2.      Wooff Y, Man SM, Aggio-Bruce R, Natoli R, Fernando N. IL-1 Family Members Mediate Cell Death, Inflammation and Angiogenesis in Retinal Degenerative Diseases. Front Immunol. 2019;10:1618.

3.      Olivares-González L, Velasco S, Campillo I, Rodrigo R. Retinal Inflammation, Cell Death and Inherited Retinal Dystrophies. Int J Mol Sci. 2021;22(4).

4.      Forrester JV, Kuffova L, Delibegovic M. The Role of Inflammation in Diabetic Retinopathy. Front Immunol. 2020;11:583687.

5.      Broadgate S, Yu J, Downes SM, Halford S. Unravelling the genetics of inherited retinal dystrophies: Past, present and future. Prog Retin Eye Res. 2017;59:53-96.

6.      Lechner J, O'Leary OE, Stitt AW. The pathology associated with diabetic retinopathy. Vision Research. 2017;139:7-14.

7.      Collaborators GBaVI. Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight: an analysis for the Global Burden of Disease Study. Lancet Glob Health. 2021;9(2):e144-e60.

8.      Chen M, Obasanmi G, Armstrong D, Lavery NJ, Kissenpfennig A, Lois N, et al. STAT3 activation in circulating myeloid-derived cells contributes to retinal microvascular dysfunction in diabetes. J Neuroinflammation. 2019;16(1):138.

9.      Gerhardinger C, Costa MB, Coulombe MC, Toth I, Hoehn T, Grosu P. Expression of acute-phase response proteins in retinal Müller cells in diabetes. Invest Ophthalmol Vis Sci. 2005;46(1):349-57.

10.     Mizutani M, Gerhardinger C, Lorenzi M. Müller cell changes in human diabetic retinopathy. Diabetes. 1998;47(3):445-9.

11.     Zeng HY, Green WR, Tso MO. Microglial activation in human diabetic retinopathy. Arch Ophthalmol. 2008;126(2):227-32.

12.     Coughlin BA, Feenstra DJ, Mohr S. Müller cells and diabetic retinopathy. Vision Research. 2017;139:93-100.

13.     Kinuthia UM, Wolf A, Langmann T. Microglia and Inflammatory Responses in Diabetic Retinopathy. Front Immunol. 2020;11:564077.

14.     Karlstetter M, Scholz R, Rutar M, Wong WT, Provis JM, Langmann T. Retinal microglia: just bystander or target for therapy? Prog Retin Eye Res. 2015;45:30-57.

15.     Di Pierdomenico J, Martínez-Vacas A, Hernández-Muñoz D, Gómez-Ramírez AM, Valiente-Soriano FJ, Agudo-Barriuso M, et al. Coordinated Intervention of Microglial and Müller Cells in Light-Induced Retinal Degeneration. Invest Ophthalmol Vis Sci. 2020;61(3):47.

16.     Wang M, Ma W, Zhao L, Fariss RN, Wong WT. Adaptive Müller cell responses to microglial activation mediate neuroprotection and coordinate inflammation in the retina. J Neuroinflammation. 2011;8:173.

17.     Reichenbach A, Bringmann A. Glia of the human retina. Glia. 2020;68(4):768-96.

18.   Newman E, Reichenbach A. The Müller cell: a functional element of the retina. Trends
Neurosci. 1996;19(8):307-12.

19.   Bringmann A, Pannicke T, Grosche J, Francke M, Wiedemann P, Skatchkov SN, et al. Müller
cells in the healthy and diseased retina. Prog Retin Eye Res. 2006;25(4):397-424.

20.   Bringmann A, Iandiev I, Pannicke T, Wurm A, Hollborn M, Wiedemann P, et al. Cellular
signaling and factors involved in Müller cell gliosis: Neuroprotective and detrimental effects. Prog
Retin Eye Res. 2009;28(6):423-51.

21.   Ruzafa N, Pereiro X, Lepper MF, Hauck SM, Vecino E. A Proteomics Approach to Identify
Candidate Proteins Secreted by Müller Glia that Protect Ganglion Cells in the Retina.
PROTEOMICS. 2018;18(11):1700321.

22.   von Toerne C, Menzler J, Ly A, Senninger N, Ueffing M, Hauck SM. Identification of a
Novel Neurotrophic Factor from Primary Retinal Müller Cells Using Stable Isotope Labeling by
Amino Acids in Cell Culture (SILAC). Mol Cell Proteomics. 2014;13(9):2371-81.

23.   Hauck SM, Schoeffmann S, Amann B, Stangassinger M, Gerhards H, Ueffing M, et al.
Retinal Mueller glial cells trigger the hallmark inflammatory process in autoimmune uveitis. J
Proteome Res. 2007;6(6):2121-31.

24.   Subirada PV, Paz MC, Ridano ME, Lorenc VE, Vaglienti MV, Barcelona PF, et al. A journey
into the retina: Müller glia commanding survival and death. European Journal of Neuroscience.
2018;47(12):1429-43.

25.   Ghaseminejad F, Kaplan L, Pfaller AM, Hauck SM, Grosche A. The role of Müller cell
glucocorticoid signaling in diabetic retinopathy. Graefes Arch Clin Exp Ophthalmol.
2020;258(2):221-30.

26.   Natoli R, Fernando N, Madigan M, Chu-Tan JA, Valter K, Provis J, et al. Microglia-derived
IL-1β promotes chemokine expression by Müller cells and RPE in focal retinal degeneration. Mol
Neurodegener. 2017;12(1):31.

27.   Rutar M, Natoli R, Chia RX, Valter K, Provis JM. Chemokine-mediated inflammation in the
degenerating retina is coordinated by Müller cells, activated microglia, and retinal pigment
epithelium. J Neuroinflammation. 2015;12(1):8.

28.   Roberge FG, Caspi RR, Nussenblatt RB. Glial retinal Müller cells produce IL-1 activity and
have a dual effect on autoimmune T helper lymphocytes. Antigen presentation manifested after
removal of suppressive activity. J Immunol. 1988;140(7):2193-6.

29.   Lorenz L, Amann B, Hirmer S, Degroote RL, Hauck SM, Deeg CA. NEU1 is more abundant
in uveitic retina with concomitant desialylation of retinal cells. Glycobiology. 2021.

30.   Lorenz L, Hirmer S, Schmalen A, Hauck SM, Deeg CA. Cell Surface Profiling of Retinal
Muller Glial Cells Reveals Association to Immune Pathways after LPS Stimulation. Cells.
2021;10(3).

31.   Middleton S. Porcine ophthalmology. Vet Clin North Am Food Anim Pract. 2010;26(3):557-
72.

32.   Kleinwort KJH, Amann B, Hauck SM, Hirmer S, Blutke A, Renner S, et al. Retinopathy with
central oedema in an INS (C94Y) transgenic pig model of long-term diabetes. Diabetologia.
2017;60(8):1541-9.

33.    Renner S, Blutke A, Clauss S, Deeg CA, Kemter E, Merkus D, et al. Porcine models for
studying complications and organ crosstalk in diabetes mellitus. Cell and Tissue Research. 2020.

34.    Konigsberg IR, Borie R, Walts AD, Cardwell J, Rojas M, Metzger F, et al. Molecular
Signatures of Idiopathic Pulmonary Fibrosis. Am J Respir Cell Mol Biol. 2021.

35.    Weigand M, Hauck SM, Deeg CA, Degroote RL. Deviant proteome profile of equine
granulocytes associates to latent activation status in organ specific autoimmune disease. J
Proteomics. 2021;230:103989.

36.    Sagmeister S, Merl-Pham J, Petrera A, Deeg CA, Hauck SM. High glucose treatment
promotes extracellular matrix proteome remodeling in Mller glial cells. PeerJ. 2021;9:e11316.

37.    Eberhardt C, Amann B, Stangassinger M, Hauck SM, Deeg CA. Isolation, characterization
and establishment of an equine retinal glial cell line: a prerequisite to investigate the physiological
function of Muller cells in the retina. J Anim Physiol Anim Nutr (Berl). 2012;96(2):260-9.

38.    Hauck SM, Suppmann S, Ueffing M. Proteomic profiling of primary retinal Muller glia cells
reveals a shift in expression patterns upon adaptation to in vitro conditions. Glia. 2003;44(3):251-63.

39.    Limb GA, Salt TE, Munro PM, Moss SE, Khaw PT. In vitro characterization of a
spontaneously immortalized human Muller cell line (MIO-M1). Invest Ophthalmol Vis Sci.
2002;43(3):864-9.

40.    Wisniewski JR, Zougman A, Nagaraj N, Mann M. Universal sample preparation method for
proteome analysis. Nat Methods. 2009;6(5):359-62.

41.    Grosche A, Hauser A, Lepper MF, Mayo R, von Toerne C, Merl-Pham J, et al. The Proteome
of Native Adult Muller Glial Cells From Murine Retina. Mol Cell Proteomics. 2016;15(2):462-80.

42.    Kall L, Canterbury JD, Weston J, Noble WS, MacCoss MJ. Semi-supervised learning for
peptide identification from shotgun proteomics datasets. Nat Methods. 2007;4(11):923-5.

43.    Navarro P, Trevisan-Herraz M, Bonzon-Kulichenko E, Nunez E, Martinez-Acedo P, Perez-
Hernandez D, et al. General statistical framework for quantitative proteomics by stable isotope
labeling. J Proteome Res. 2014;13(3):1234-47.

44.    Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. Controlling the false discovery rate in
behavior genetics research. Behav Brain Res. 2001;125(1-2):279-84.

45.    Kramer A, Green J, Pollard J, Jr., Tugendreich S. Causal analysis approaches in Ingenuity
Pathway Analysis. Bioinformatics. 2014;30(4):523-30.

46.    Fisher RA. On the Interpretation of $\chi^2$ from Contingency Tables, and the Calculation of P.
Journal of the Royal Statistical Society. 1922;85(1):87-94.

47.    Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-
wide expression patterns. Proc Natl Acad Sci U S A. 1998;95(25):14863-8.

48.    Saldanha AJ. Java Treeview--extensible visualization of microarray data. Bioinformatics.
2004;20(17):3246-8.

49.    Lunney JK, Ho CS, Wysocki M, Smith DM. Molecular genetics of the swine major
histocompatibility complex, the SLA complex. Dev Comp Immunol. 2009;33(3):362-74.

50.    Sigal LJ. Activation of CD8 T Lymphocytes during Viral Infections. Encyclopedia of
Immunobiology. 2016:286-90.

51.    Hewitt EW. The MHC class I antigen presentation pathway: strategies for viral immune
evasion. Immunology. 2003;110(2):163-9.

52.    Roche PA, Furuta K. The ins and outs of MHC class II-mediated antigen processing and
presentation. Nat Rev Immunol. 2015;15(4):203-16.

53.    Lechner J, O'Leary OE, Stitt AW. The pathology associated with diabetic retinopathy. Vision
Res. 2017;139:7-14.

54.    Zheng Y, He M, Congdon N. The worldwide epidemic of diabetic retinopathy. Indian J
Ophthalmol. 2012;60(5):428-31.

55.    Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, et al. Fate mapping analysis
reveals that adult microglia derive from primitive macrophages. Science. 2010;330(6005):841-5.

56.    Coughlin BA, Feenstra DJ, Mohr S. Muller cells and diabetic retinopathy. Vision Res.
2017;139:93-100.

57.    Wang J, Xu X, Elliott MH, Zhu M, Le YZ. Muller cell-derived VEGF is essential for
diabetes-induced retinal inflammation and vascular leakage. Diabetes. 2010;59(9):2297-305.

58.    Bai Y, Ma JX, Guo J, Wang J, Zhu M, Chen Y, et al. Muller cell-derived VEGF is a
significant contributor to retinal neovascularization. J Pathol. 2009;219(4):446-54.

59.    Demircan N, Safran BG, Soylu M, Ozcan AA, Sizmaz S. Determination of vitreous
interleukin-1 (IL-1) and tumour necrosis factor (TNF) levels in proliferative diabetic retinopathy. Eye
(Lond). 2006;20(12):1366-9.

60.    Kelly K, Wang JJ, Zhang SX. The unfolded protein response signaling and retinal Muller cell
metabolism. Neural Regen Res. 2018;13(11):1861-70.

61.    Zhong Y, Li J, Chen Y, Wang JJ, Ratan R, Zhang SX. Activation of endoplasmic reticulum
stress by hyperglycemia is essential for Muller cell-derived inflammatory cytokine production in
diabetes. Diabetes. 2012;61(2):492-504.

62.    Zhang SX, Wang JJ, Gao G, Shao C, Mott R, Ma JX. Pigment epithelium-derived factor
(PEDF) is an endogenous antiinflammatory factor. FASEB J. 2006;20(2):323-5.

63.    He J, Wang H, Liu Y, Li W, Kim D, Huang H. Blockade of vascular endothelial growth
factor receptor 1 prevents inflammation and vascular leakage in diabetic retinopathy. J Ophthalmol.
2015;2015:605946.

64.    Liu Y, Li L, Pan N, Gu J, Qiu Z, Cao G, et al. TNF-alpha released from retinal Muller cells
aggravates retinal pigment epithelium cell apoptosis by upregulating mitophagy during diabetic
retinopathy. Biochem Biophys Res Commun. 2021;561:143-50.

65.    Gilbert RWD, Vickaryous MK, Viloria-Petit AM. Signalling by Transforming Growth Factor
Beta Isoforms in Wound Healing and Tissue Regeneration. J Dev Biol. 2016;4(2).

66.    Yaswen L, Kulkarni AB, Fredrickson T, Mittleman B, Schiffman R, Payne S, et al.
Autoimmune manifestations in the transforming growth factor-beta 1 knockout mouse. Blood.
1996;87(4):1439-45.

67.    Yamagami S, Yokoo S, Mimura T, Amano S. Effects of TGF-beta2 on immune response-
related gene expression profiles in the human corneal endothelium. Invest Ophthalmol Vis Sci.
2004;45(2):515-21.

68.     Huang T, Schor SL, Hinck AP. Biological activity differences between TGF-beta1 and TGF-
beta3 correlate with differences in the rigidity and arrangement of their component monomers.
Biochemistry. 2014;53(36):5737-49.

69.     Kulkarni AB, Huh CG, Becker D, Geiser A, Lyght M, Flanders KC, et al. Transforming
growth factor beta 1 null mutation in mice causes excessive inflammatory response and early death.
Proc Natl Acad Sci U S A. 1993;90(2):770-4.

70.     Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, et al. Targeted disruption
of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease.
Nature. 1992;359(6397):693-9.

71.     Sanford LP, Ormsby I, Gittenberger-de Groot AC, Sariola H, Friedman R, Boivin GP, et al.
TGFbeta2 knockout mice have multiple developmental defects that are non-overlapping with other
TGFbeta knockout phenotypes. Development. 1997;124(13):2659-70.

72.     Proetzel G, Pawlowski SA, Wiles MV, Yin M, Boivin GP, Howles PN, et al. Transforming
growth factor-beta 3 is required for secondary palate fusion. Nat Genet. 1995;11(4):409-14.

73.     Yang LT, Kaartinen V. Tgfb1 expressed in the Tgfb3 locus partially rescues the cleft palate
phenotype of Tgfb3 null mutants. Dev Biol. 2007;312(1):384-95.

74.     Hall BE, Wankhade UD, Konkel JE, Cherukuri K, Nagineni CN, Flanders KC, et al.
Transforming growth factor-beta3 (TGF-beta3) knock-in ameliorates inflammation due to TGF-beta1
deficiency while promoting glucose tolerance. J Biol Chem. 2013;288(44):32074-92.

75.     Schultz-Cherry S, Lawler J, Murphy-Ullrich JE. The type 1 repeats of thrombospondin 1
activate latent transforming growth factor-beta. J Biol Chem. 1994;269(43):26783-8.

76.     Schultz-Cherry S, Murphy-Ullrich JE. Thrombospondin causes activation of latent
transforming growth factor-beta secreted by endothelial cells by a novel mechanism. J Cell Biol.
1993;122(4):923-32.

77.     Murphy-Ullrich JE, Downs JC. The Thrombospondin1-TGF-beta Pathway and Glaucoma. J
Ocul Pharmacol Ther. 2015;31(7):371-5.

78.     Conedera FM, Quintela Pousa AM, Presby DM, Mercader N, Enzmann V, Tschopp M.
Diverse Signaling by TGFbeta Isoforms in Response to Focal Injury is Associated with Either
Retinal Regeneration or Reactive Gliosis. Cell Mol Neurobiol. 2021;41(1):43-62.

79.     Kaminska B, Gozdz A, Zawadzka M, Ellert-Miklaszewska A, Lipko M. MAPK signal
transduction underlying brain inflammation and gliosis as therapeutic target. Anat Rec (Hoboken).
2009;292(12):1902-13.

80.     Tripathi RC, Li J, Chan WF, Tripathi BJ. Aqueous humor in glaucomatous eyes contains an
increased level of TGF-beta 2. Exp Eye Res. 1994;59(6):723-7.

81.     Fuchshofer R, Birke M, Welge-Lussen U, Kook D, Lutjen-Drecoll E. Transforming growth
factor-beta 2 modulated extracellular matrix component expression in cultured human optic nerve
head astrocytes. Invest Ophthalmol Vis Sci. 2005;46(2):568-78.

82.     Fuchshofer R. The pathogenic role of transforming growth factor-beta2 in glaucomatous
damage to the optic nerve head. Exp Eye Res. 2011;93(2):165-9.

83.     Fuchshofer R, Tamm ER. The role of TGF-beta in the pathogenesis of primary open-angle
glaucoma. Cell Tissue Res. 2012;347(1):279-90.

84.     Zode GS, Sethi A, Brun-Zinkernagel AM, Chang IF, Clark AF, Wordinger RJ. Transforming growth factor-beta2 increases extracellular matrix proteins in optic nerve head cells via activation of the Smad signaling pathway. Mol Vis. 2011;17:1745-58.

85.     Mohammad G, Kowluru RA. Matrix metalloproteinase-2 in the development of diabetic retinopathy and mitochondrial dysfunction. Lab Invest. 2010;90(9):1365-72.

86.     Tan W, Zou JL, Yoshida S, Jiang B, Zhou YD. Increased vitreal levels of interleukin-10 in diabetic retinopathy: a Meta-analysis. Int J Ophthalmol. 2020;13(9):1477-83.

87.     Dace DS, Khan AA, Kelly J, Apte RS. Interleukin-10 promotes pathological angiogenesis by regulating macrophage response to hypoxia during development. PLoS One. 2008;3(10):e3381.

88.     Luzina IG, Keegan AD, Heller NM, Rook GA, Shea-Donohue T, Atamas SP. Regulation of inflammation by interleukin-4: a review of "alternatives". J Leukoc Biol. 2012;92(4):753-64.

89.     Major J, Fletcher JE, Hamilton TA. IL-4 pretreatment selectively enhances cytokine and chemokine production in lipopolysaccharide-stimulated mouse peritoneal macrophages. J Immunol. 2002;168(5):2456-63.

90.     Gadani SP, Cronk JC, Norris GT, Kipnis J. IL-4 in the brain: a cytokine to remember. J Immunol. 2012;189(9):4213-9.

91.     Yoshida S, Sotozono C, Ikeda T, Kinoshita S. Interleukin-6 (IL-6) production by cytokine-stimulated human Muller cells. Curr Eye Res. 2001;22(5):341-7.

92.     Yao Y, Li R, Du J, Long L, Li X, Luo N. Interleukin-6 and Diabetic Retinopathy: A Systematic Review and Meta-Analysis. Curr Eye Res. 2019;44(5):564-74.

93.     Liu X, Ye F, Xiong H, Hu DN, Limb GA, Xie T, et al. IL-1beta induces IL-6 production in retinal Muller cells predominantly through the activation of p38 MAPK/NF-kappaB signaling pathway. Exp Cell Res. 2015;331(1):223-31.

94.     Coughlin BA, Trombley BT, Mohr S. Interleukin-6 (IL-6) mediates protection against glucose toxicity in human Muller cells via activation of VEGF-A signaling. Biochem Biophys Res Commun. 2019;517(2):227-32.

95.     Ucgun NI, Zeki-Fikret C, Yildirim Z. Inflammation and diabetic retinopathy. Mol Vis. 2020;26:718-21.

96.     Wu H, Hwang DK, Song X, Tao Y. Association between Aqueous Cytokines and Diabetic Retinopathy Stage. J Ophthalmol. 2017;2017:9402198.

97.     Bazan JF, Bacon KB, Hardiman G, Wang W, Soo K, Rossi D, et al. A new class of membrane-bound chemokine with a CX3C motif. Nature. 1997;385(6617):640-4.

98.     Imai T, Hieshima K, Haskell C, Baba M, Nagira M, Nishimura M, et al. Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. Cell. 1997;91(4):521-30.

99.     Abu El-Asrar AM, Nawaz MI, Ahmad A, De Zutter A, Siddiquei MM, Blanter M, et al. Evaluation of Proteoforms of the Transmembrane Chemokines CXCL16 and CX3CL1, Their Receptors, and Their Processing Metalloproteinases ADAM10 and ADAM17 in Proliferative Diabetic Retinopathy. Front Immunol. 2020;11:601639.

100.     Serra AM, Waddell J, Manivannan A, Xu H, Cotter M, Forrester JV. CD11b+ bone marrow-derived monocytes are the major leukocyte subset responsible for retinal capillary leukostasis in

experimental diabetes in mouse and express high levels of CCR5 in the circulation. Am J Pathol.
2012;181(2):719-27.

101.    Wu F, Phone A, Lamy R, Ma D, Laotaweerungsawat S, Chen Y, et al. Correlation of
Aqueous, Vitreous, and Plasma Cytokine Levels in Patients With Proliferative Diabetic Retinopathy.
Invest Ophthalmol Vis Sci. 2020;61(2):26.

102.    Zhang S, Zhang S, Gong W, Zhu G, Wang S, Wang Y, et al. Muller Cell Regulated
Microglial Activation and Migration in Rats With N-Methyl-N-Nitrosourea-Induced Retinal
Degeneration. Front Neurosci. 2018;12:890.

103.    Greene JA, Portillo JA, Lopez Corcino Y, Subauste CS. CD40-TRAF Signaling Upregulates
CX3CL1 and TNF-alpha in Human Aortic Endothelial Cells but Not in Retinal Endothelial Cells.
PLoS One. 2015;10(12):e0144133.

104.    Pauly D, Agarwal D, Dana N, Schafer N, Biber J, Wunderlich KA, et al. Cell-Type-Specific
Complement Expression in the Healthy and Diseased Retina. Cell Rep. 2019;29(9):2835-48 e4.

105.    Kaczmarek-Hajek K, Zhang J, Kopp R, Grosche A, Rissiek B, Saul A, et al. Re-evaluation of
neuronal P2X7 expression using novel mouse models and a P2X7-specific nanobody. Elife. 2018;7.

106.    Cowan CS, Renner M, De Gennaro M, Gross-Scherf B, Goldblum D, Hou Y, et al. Cell
Types of the Human Retina and Its Organoids at Single-Cell Resolution. Cell. 2020;182(6):1623-40
e34.

107.    Mano T, Tokuda N, Puro DG. Interferon-gamma induces the expression of major
histocompatibility antigens by human retinal glial cells. Exp Eye Res. 1991;53(5):603-7.

108.    Kambayashi T, Laufer TM. Atypical MHC class II-expressing antigen-presenting cells: can
anything replace a dendritic cell? Nat Rev Immunol. 2014;14(11):719-30.

109.    Bejarano-Escobar R, Sanchez-Calderon H, Otero-Arenas J, Martin-Partido G, Francisco-
Morcillo J. Muller glia and phagocytosis of cell debris in retinal tissue. J Anat. 2017;231(4):471-83.

110.    Sakami S, Imanishi Y, Palczewski K. Muller glia phagocytose dead photoreceptor cells in a
mouse model of retinal degenerative disease. FASEB J. 2019;33(3):3680-92.

111.    Singh PK, Shiha MJ, Kumar A. Antibacterial responses of retinal Muller glia: production of
antimicrobial peptides, oxidative burst and phagocytosis. J Neuroinflammation. 2014;11:33.

112.    Gutierrez-Martinez E, Planes R, Anselmi G, Reynolds M, Menezes S, Adiko AC, et al. Cross-
Presentation of Cell-Associated Antigens by MHC Class I in Dendritic Cell Subsets. Front Immunol.
2015;6:363.

113.    Larsson M, Fonteneau JF, Bhardwaj N. Dendritic cells resurrect antigens from dead cells.
Trends Immunol. 2001;22(3):141-8.

114.    Kastelan S, Tomic M, Salopek-Rabatic J, Pavan J, Lukenda A, Gotovac M, et al. The
association between the HLA system and retinopathy development in patients with type 1 diabetes
mellitus. Coll Antropol. 2013;37 Suppl 1:65-70.

115.    Barbosa J, Ramsay RC, Knobloch WH, Cantrill HL, Noreen H, King R, et al.
Histocompatibility antigen frequencies in diabetic retinopathy. Am J Ophthalmol. 1980;90(2):148-
53.

116.    Dornan TL, Ting A, McPherson CK, Peckar CO, Mann JI, Turner RC, et al. Genetic
susceptibility to the development of retinopathy in insulin-dependent diabetics. Diabetes.
1982;31(3):226-31.

117.    Falck AA, Knip JM, Ilonen JS, Laatikainen LT. Genetic markers in early diabetic retinopathy
of adolescents with type I diabetes. J Diabetes Complications. 1997;11(4):203-7.

118.    Cruickshanks KJ, Vadheim CM, Moss SE, Roth MP, Riley WJ, Maclaren NK, et al. Genetic
marker associations with proliferative retinopathy in persons diagnosed with diabetes before 30 yr of
age. Diabetes. 1992;41(7):879-85.

119.    Birinci A, Birinci H, Abidinoglu R, Durupinar B, Oge I. Diabetic retinopathy and HLA
antigens in type 2 diabetes mellitus. Eur J Ophthalmol. 2002;12(2):89-93.

120.    Agardh D, Gaur LK, Agardh E, Landin-Olsson M, Agardh CD, Lernmark A. HLA-
DQB1*0201/0302 is associated with severe retinopathy in patients with IDDM. Diabetologia.
1996;39(11):1313-7.

121.    Santos DC, Porto LC, Pizarro MH, de Melo LGN, Silva DA, Oliveira RV, et al. Human
Leukocyte Antigens class II (HLA II) gene profile from an admixed population of patients with type
1 diabetes with severe diabetic retinopathy: a nested case-control study in Brazil. Diabetol Metab
Syndr. 2021;13(1):83.

122.    Liu C, Chen G, Bentley AR, Doumatey A, Zhou J, Adeyemo A, et al. Genome-wide
association study for proliferative diabetic retinopathy in Africans. NPJ Genom Med. 2019;4:20.

123.    Portillo JA, Greene JA, Okenka G, Miao Y, Sheibani N, Kern TS, et al. CD40 promotes the
development of early diabetic retinopathy in mice. Diabetologia. 2014;57(10):2222-31.

124.    Portillo JC, Lopez Corcino Y, Miao Y, Tang J, Sheibani N, Kern TS, et al. CD40 in Retinal
Müller Cells Induces P2X7-Dependent Cytokine Expression in Macrophages/Microglia in Diabetic
Mice and Development of Early Experimental Diabetic Retinopathy. Diabetes. 2017;66(2):483-93.

125.    Subauste CS. The CD40-ATP-P2X 7 Receptor Pathway: Cell to Cell Cross-Talk to Promote
Inflammation and Programmed Cell Death of Endothelial Cells. Front Immunol. 2019;10:2958.

126.    Cvitkovic K, Sesar A, Sesar I, Pusic-Sesar A, Pejic R, Kelava T, et al. Concentrations of
Selected Cytokines and Vascular Endothelial Growth Factor in Aqueous Humor and Serum of
Diabetic Patients. Semin Ophthalmol. 2020;35(2):126-33.

127.    Ozay Y, Ozek D, Yildirim F, Yildirim Z. The effect of diabetes on vitreous levels of
adiponectin and inflammatory cytokines in experimental rat model. Adv Clin Exp Med.
2020;29(4):449-52.

128.    Vujosevic S, Micera A, Bini S, Berton M, Esposito G, Midena E. Proteome analysis of retinal
glia cells-related inflammatory cytokines in the aqueous humour of diabetic patients. Acta
Ophthalmol. 2016;94(1):56-64.

129.    Tan L, Gordon KB, Mueller JP, Matis LA, Miller SD. Presentation of proteolipid protein
epitopes and B7-1-dependent activation of encephalitogenic T cells by IFN-gamma-activated SJL/J
astrocytes. J Immunol. 1998;160(9):4271-9.

130.    Portillo JC, Yu JS, Hansen S, Kern TS, Subauste MC, Subauste CS. A cell-penetrating CD40-
TRAF2,3 blocking peptide diminishes inflammation and neuronal loss after ischemia/reperfusion.
FASEB J. 2021;35(3):e21412.

131.    Cecilia OM, Jose Alberto CG, Jose NP, Ernesto German CM, Ana Karen LC, Luis Miguel
RP, et al. Oxidative Stress as the Main Target in Diabetic Retinopathy Pathophysiology. J Diabetes
Res. 2019;2019:8562408.

132.    Tien T, Zhang J, Muto T, Kim D, Sarthy VP, Roy S. High Glucose Induces Mitochondrial
Dysfunction in Retinal Muller Cells: Implications for Diabetic Retinopathy. Invest Ophthalmol Vis
Sci. 2017;58(7):2915-21.

133.    Krugel K, Wurm A, Pannicke T, Hollborn M, Karl A, Wiedemann P, et al. Involvement of
oxidative stress and mitochondrial dysfunction in the osmotic swelling of retinal glial cells from
diabetic rats. Exp Eye Res. 2011;92(1):87-93.

134.    Behl T, Kotwani A. Exploring the various aspects of the pathological role of vascular
endothelial growth factor (VEGF) in diabetic retinopathy. Pharmacol Res. 2015;99:137-48.

135.    Homme RP, Singh M, Majumder A, George AK, Nair K, Sandhu HS, et al. Remodeling of
Retinal Architecture in Diabetic Retinopathy: Disruption of Ocular Physiology and Visual Functions
by Inflammatory Gene Products and Pyroptosis. Front Physiol. 2018;9:1268.

136.    Wang J, Shanmugam A, Markand S, Zorrilla E, Ganapathy V, Smith SB. Sigma 1 receptor
regulates the oxidative stress response in primary retinal Muller glial cells via NRF2 signaling and
system xc(-), the Na(+)-independent glutamate-cystine exchanger. Free Radic Biol Med. 2015;86:25-
36.

137.    Geng J, Sun X, Wang P, Zhang S, Wang X, Wu H, et al. Kinases Mst1 and Mst2 positively
regulate phagocytic induction of reactive oxygen species and bactericidal activity. Nat Immunol.
2015;16(11):1142-52.

138.    Wang P, Geng J, Gao J, Zhao H, Li J, Shi Y, et al. Macrophage achieves self-protection
against oxidative stress-induced ageing through the Mst-Nrf2 axis. Nat Commun. 2019;10(1):755.

139.    Hollborn M, Ulbricht E, Rillich K, Dukic-Stefanovic S, Wurm A, Wagner L, et al. The
human Muller cell line MIO-M1 expresses opsins. Mol Vis. 2011;17:2738-50.

140.    Lawrence JM, Singhal S, Bhatia B, Keegan DJ, Reh TA, Luthert PJ, et al. MIO-M1 cells and
similar muller glial cell lines derived from adult human retina exhibit neural stem cell characteristics.
Stem Cells. 2007;25(8):2033-43.

141.    Merl J, Ueffing M, Hauck SM, von Toerne C. Direct comparison of MS-based label-free and
SILAC quantitative proteome profiling strategies in primary retinal Muller cells. Proteomics.
2012;12(12):1902-11.

142.    Hauck SM, Gloeckner CJ, Harley ME, Schoeffmann S, Boldt K, Ekstrom PA, et al.
Identification of paracrine neuroprotective candidate proteins by a functional assay-driven
proteomics approach. Mol Cell Proteomics. 2008;7(7):1349-61.

143.    von Toerne C, Menzler J, Ly A, Senninger N, Ueffing M, Hauck SM. Identification of a
novel neurotrophic factor from primary retinal Muller cells using stable isotope labeling by amino
acids in cell culture (SILAC). Mol Cell Proteomics. 2014;13(9):2371-81.

1088