Do closely related species interact with similar partners? Testing for phylogenetic signal in bipartite interaction networks

Benoît Perez-Lamarque [1,2], Odile Maliet [1], Marc-André Selosse [2,3], Florent Martos [2], and Hélène Morlon [1]

[1] *Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France*

[2] *Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France*

[3] *Department of Plant Taxonomy and Nature Conservation, University of Gdansk, Wita Stwosza 59, 80-308 Gdansk, Poland*

<u>Corresponding author:</u> Benoît Perez-Lamarque (benoit.perez@ens.psl.eu)

<u>Author contributions:</u> BPL, OM, MAS, FM, and HM designed the study. BPL performed the analyses and FM gathered the data. BPL and HM wrote the first draft of the manuscript and all authors contributed to revisions.

<u>Data accessibility statement:</u> All the R functions used to measure phylogenetic signals in bipartite interaction networks, including (simple, partial, and clade-specific) Mantel tests and PBLM, are available in the R-package RPANDA (Morlon *et al.* 2016) (functions *phylosignal_network* and *phylosignal_sub_network*). A tutorial and the simulated networks can be found at https://github.com/BPerezLamarque/Phylosignal_network. Amended functions of *BipartiteEvol* are also included in RPANDA.

Abstract

Whether interactions between species are conserved on evolutionary time-scales is a

central question in ecology and evolution. It has spurred the development of both

correlative and model-based approaches for testing phylogenetic signal in interspecific

interactions: do closely related species interact with similar sets of partners? Here we

test the statistical performances of some of these approaches using simulations. We

find that one of the most widely used model-based approach, the Phylogenetic

Bipartite Linear Model (PBLM), often detects phylogenetic signal when it should not.

Simple Mantel tests investigating the correlation between phylogenetic distances and

dissimilarities in sets of interacting partners instead have low type-I error rates and

satisfactory statistical power, especially when using weighted interactions and

phylogenetic dissimilarity metrics; however, they often artifactually detect anti-

phylogenetic signals. Partial Mantel tests, which are used to partial out the

phylogenetic signal in the number of partners, actually fail at correcting for this

confounding effect, and we instead propose the sequential use of simple Mantel tests.

We also explore the ability of simple Mantel tests to analyze clade-specific

phylogenetic signal, while current methods only measure an overall signal. We

provide general guidelines and an empirical application on an interaction network

between orchids and mycorrhizal fungi.

Keywords: ecological network, phylogenetic signal, Mantel tests, clade-specific signal,

species interactions, mycorrhizal symbiosis.

Introduction

Species in ecological communities engage in numerous types of interspecific interactions, such as pollination, mycorrhizal symbioses, predation, and parasitism (Bascompte *et al.* 2003; Fontaine *et al.* 2011; Martos *et al.* 2012; Bascompte & Jordano 2013). Understanding the processes that shape these interaction networks, including the role of evolutionary history, *i.e.* the phylogeny, is a major focus of ecology and evolution (Rezende *et al.* 2007; Vázquez *et al.* 2009; Krasnov *et al.* 2012; Elias *et al.* 2013; Rohr & Bascompte 2014). One way to assess the role of evolutionary history in shaping contemporary interactions is to test for phylogenetic signal in species interactions, *i.e.* whether closely related species interact with similar sets of partners (Peralta 2016).

Testing for phylogenetic signal in a unidimensional trait (*i.e.* whether a trait is phylogenetically conserved) for a given clade is mainstream (Felsenstein 1985; Blomberg *et al.* 2003; Münkemüller *et al.* 2012). One approach (the 'correlative' approach) is to perform a Mantel test between phylogenetic and trait distances (Mantel 1967); another approach (the 'model-based' approach) relies on trait evolution models such as Pagel's λ (Pagel 1999) or Blomberg's *K* (Blomberg *et al.* 2003). The model-based approach has a higher ability to detect an existing phylogenetic signal (power) and a lower propensity to infer a phylogenetic signal when it should not (type-I error; Harmon & Glor 2010): The correlative approach should therefore only be used when the model-based approach is not applicable, *e.g.* if the 'trait' data is expressed in terms of pairwise distances.

Testing for phylogenetic signal in species interactions falls in the category of cases where the 'trait' data are pairwise distances, here the between-species dissimilarity in sets of interacting species. Simple Mantel tests have therefore been widely used in this context (*e.g.* Rezende *et al.* 2007; Elias *et al.* 2013; Fontaine & Thébault 2015). Partial Mantel tests have also been used to test whether the

phylogenetic signal reflects more the identity of the interacting partners than their

number, *i.e.* the degree, as similarity in the number of partners can increase the value

of similarity metrics (Rezende *et al.* 2007; Jacquemyn *et al.* 2011; Aizen *et al.* 2016).

Besides such correlative approaches, the Phylogenetic Bipartite Linear Model (PBLM,

Ives & Godfray 2006) has been widely used to test for phylogenetic signal in species

interactions in a variety of empirical networks, *e.g.* in host-parasite, plant-fungus, and

pollination networks (Martos *et al.* 2012; Martín González *et al.* 2015; Xing *et al.* 2020).

Here, using simulations, we perform a comparative analysis of the statistical

performances of the correlative (simple or partial) Mantel tests and of the model-based

PBLM to test for phylogenetic signal in species interactions (Box 1), considering

weighted and unweighted bipartite interaction networks between species from two

guilds A and B (Fig. 1). We also investigate the ability of Mantel tests to detect the

presence of phylogenetic signal in the different clades of a phylogenetic tree, because

current methods only measure the overall signal in the phylogeny. Finally, we provide

general guidelines and apply them to an orchid-fungus mycorrhizal network

identified across the oceanic island of Réunion (Martos *et al.* 2012).

Methods

Simulating bipartite interaction networks with or without phylogenetic signal in species interactions

We used *BipartiteEvol*, an individual-based eco-evolutionary model (Maliet *et al.* 2020), to generate interaction networks with or without phylogenetic signal between two guilds interacting in a mutualistic, antagonistic, or neutral way. Each individual from guild A (resp. B) is characterized by a multidimensional continuous trait and interacts with one individual from guild B (resp. A). The effect of this interaction on the fitness of each individual from guilds A or B is determined by the distance in trait space of the two interacting individuals, according to a classical trait matching expression parametrized by two parameters α_A and α_B (Supplementary Methods 1, Maliet *et al.* 2020). These parameters determine the nature and specificity of the interaction: positive α_A and α_B correspond to mutualistic interactions, negative α_A and positive α_B to antagonistic interactions (with guild A representing hosts/preys and guild B parasites/predators), high $|\alpha|$ values to scenarios with strong fitness effects (*i.e.* highly specialized interactions), and $|\alpha|$ values close to 0 to more neutral scenarios. *BipartiteEvol* simulates individual's deaths and births (proportional to the individual's fitness) and new individuals have a probability μ to mutate, in which case new traits are drawn independently in a normal distribution centered on the parent traits. Here, we amended the species delineation of the original *BipartiteEvol* model (Maliet *et al.* 2020) and considered that each combination of traits forms a new species. Under this process, closely related species tend to interact with similar sets of partners (*i.e.* there is a phylogenetic signal in species interactions) if (and only if): (1) closely related species have similar traits (*i.e.* there is a phylogenetic signal in species traits) and (2) these traits determine who interacts with whom, *i.e.* $\alpha \neq 0$. Similarly, an anti-phylogenetic signal in species interactions (*i.e.* the tendency for closely related species

to associate with dissimilar partners) is expected if there is anti-phylogenetic signal in species traits and $\alpha \neq 0$.

Using the R-package RPANDA (Morlon *et al.* 2016; R Core Team 2020), we simulated a total of 2,400 interaction networks with individuals characterized by a six-dimensional trait. To obtain a wide range of network sizes, we considered a total number of 500, 1,000, 2,000, 3,000, 4,000, or 5,000 pairs of interacting individuals per simulation. For each size, we simulated the evolution of 100 neutral networks (α_A=0 ; α_B=0), 120 mutualistic networks (**i**: α_A=1; α_B=1; **ii**: α_A=0.1; α_B=0.1; **iii**: α_A=0.01; α_B=0.01; **iv**: α_A=1; α_B=0.1; **v**: α_A=1; α_B=0.01; and **vi**: α_A=0.1; α_B=0.01) and 180 antagonistic networks (**i**: α_A=-1; α_B=1; **ii**: α_A=-0.1; α_B=0.1; **iii**: α_A=-0.01; α_B=0.01; **iv**: α_A=-1; α_B=0.1; **v**: α_A=-1; α_B=0.01; **vi**: α_A=-0.1; α_B=1; **vii**: α_A=-0.1; α_B=0.01; **viii**: α_A=-0.01; α_B=1; **ix**: α_A=-0.01; α_B=0.1). We used a mutation rate μ=0.01 and followed the interacting individuals during 50^6 death events. At the end, we extracted for each guild a species tree from its genealogy by randomly selecting one individual per species (Fig. S1) and reconstructed the corresponding weighted interaction network by counting the number of occurrences of each interspecific interaction.

We separated the 2,400 simulated networks between those for which we should expect a phylogenetic signal in species interactions and those for which we should not. We did not expect phylogenetic signal in species interactions in neutral networks and in non-neutral networks with no phylogenetic signal in species traits. Conversely, we expected phylogenetic signal in non-neutral networks with phylogenetic signal in species traits. Phylogenetic signal in species traits was evaluated using Mantel tests (Pearson correlation) between phylogenetic distances and trait distances computed as the Euclidian distances between trait values for each species pair.

Computing phylogenetic signal in species interactions

<u>Mantel tests:</u> We evaluated the phylogenetic signal in species interactions in guilds A and B separately using simple Mantel tests between phylogenetic and ecological (set

of interacting partners) distances. Ecological distances were measured both without

accounting for evolutionary relatedness of the interacting partners, using (weighted or

unweighted) Jaccard, and accounting for relatedness using (weighted or unweighted)

UniFrac distances (Box 1). We used Pearson, Spearman, and Kendall correlations by

extending the *mantel* function in the R-package ecodist (Goslee & Urban 2007); we

evaluated the significance of each correlation using 10,000 permutations, except for the

computationally intensive Kendall correlation (100 permutations only). For each

network, we considered that there was a significant phylogenetic signal (resp. anti-

phylogenetic signal) if the correlation coefficient was higher (resp. lower) than >95%

of the randomized correlations; we computed the p-value of each one-tailed Mantel

test as the fraction of the randomized correlations above (resp. below) the original

value.

PBLM: To estimate phylogenetic signal based on PBLM, we modified the function *pblm*

from the R-package picante (Kembel *et al.* 2010) to more efficiently perform matrix

inversions and handle large interaction networks. We followed Ives & Godfray (2006;

Box 1) by considering that the phylogenetic signal is significant when the mean square

error (MSE) of the model is smaller than that obtained using star phylogenies (MSE_{star});

we also used a more stringent criterion by considering that the signal is significant

when the MSE is at least 5% lower than MSE_{star}. Finally, we applied the bootstrapping

method of Ives & Godfray (2006; Box 1) for the smallest networks (its computational

cost prevented us from applying it on larger networks).

Confounding effect of the phylogenetic signal in the number of partners

To test the performances of the partial Mantel test at measuring phylogenetic signal in

species interactions while controlling for the number of partners (Box 1), we first

performed partial Mantel tests between phylogenetic and ecological distances, while

controlling for the absolute differences in degrees, on the networks simulated with

BipartiteEvol. There is no reason to produce a phylogenetic signal in the number of

partners in *BipartiteEvol* simulations, and we verified this by performing Mantel tests

between phylogenetic distances and degree differences. These analyses were

performed to assess whether partial Mantel tests loose power compared to simple

Mantel tests.

Second, we assessed whether partial Mantel tests successfully correct for phylogenetic

signal in the number of partners using networks simulated under a process that

generate phylogenetic conservatism in the number, but not the identity, of interacting

partners (*i.e.* partial Mantel tests should not be significant when applied to such

networks). To simulate network with only phylogenetic conservatism in the number

of partners in guild A, we first simulated phylogenetic trees for guilds A and B using

pbtree (R-package phytools; Revell 2012) with a number of species uniformly sampled

between 40 and 150 by guild. Next, we simulated the number of partners of the species

from guild A using an Ornstein-Uhlenbeck process with an attraction toward 0, a

variance of 0.1 (noise of the Brownian motion), and a selection strength (a_A) ranging

from 5 (strong stabilizing effect, weak phylogenetic signal) to 0 (Brownian motion,

strong phylogenetic signal). We computed the number of partners per species by

calibrating the simulated values between 1 and the number of species in guild B and

taking the integer part. For each a_A value (5, 1, 0.5, 0.05, or 0), we performed 100

simulations using *mvSIM* (R-package mvMORPH; Clavel *et al.* 2015). Finally, for each

species in A, we attributed the corresponding number of partners in B at random to

obtain binary networks. We checked that our simulations indeed generated a signal in

the number of partners by performing simple Mantel tests between phylogenetic and

degree difference distances. Finally, we performed on each simulated network a

partial Mantel test.

Given the poor performances of partial Mantel tests (see Results), we tested

whether using sequential Mantel tests would provide a good alternative: based on

simple Mantel tests, we consider that there is a phylogenetic signal in the identity of

the partners if there is a phylogenetic signal in species interactions and no phylogenetic

signal in the number of partners. We applied this sequential testing to all our simulated

networks.

Effect of phylogenetic uncertainty, sampling asymmetry, and network

heterogeneity on measures of phylogenetic signal in species interactions

Unlike simulations (such as those provided by *BipartiteEvol*), empirical bipartite

networks suffer from phylogenetic uncertainty (*e.g.* in the microbial partners' tree

when studying host-associated microbiota – which often prevents accounting for

evolutionary relatedness; *i.e.* using UniFrac distances), sampling asymmetry (*i.e.* one

side of the network is more thoroughly sampled than the other), and network

heterogeneity (*i.e.* different sub-clades in the network have different levels of

phylogenetic signal). We performed additional analyses to investigate the effect of

these aspects on phylogenetic signals in species interactions measured using simple

Mantel tests.

First, we tested the effect of phylogenetic uncertainty in the partners' tree on the

measure of phylogenetic signal when evolutionary relatedness is accounted for (*i.e.*

using UniFrac distances). We performed these analyses to assess whether accounting

for the partners' evolutionary relatedness remains advantageous (see Results) when

phylogenetic uncertainty is high. To add some variability in the phylogenetic tree of

guild B (resp. A) used to compute the UniFrac distances between species pairs from

guild A (resp. B), we first simulated, on the original partners tree, the evolution of a

short DNA sequence and then reconstructed the tree from the simulated DNA

alignment using neighbor-joining (*nj* function, R-package APE (Paradis *et al.* 2004)).

We used *simulate_alignment* (R-package HOME; Perez-Lamarque & Morlon 2019) to

simulate sequences of length 75, 150, 300, 600, or 1,200 base-pairs, with 30% of variable

sites, and a substitution rate of 1.5 (shorter fragments should result in noisier

phylogenies).

Second, we tested the influence of sampling asymmetry on measures of

phylogenetic signal. Empirical networks are often an incomplete representation of the

actual interactions between two guilds because they are under-sampled, and

frequently, in an asymmetrical way. For instance, by sampling targeted species from

guild A, observed networks are constituted by few species from guild A which have

the complete set of their partners and by often more species from guild B which have

an incomplete set of their partners (as they likely interact with unsampled species from

guild A). We tested the influence of such sampling asymmetry by selecting only 10%

of the most abundant species from guild A in each simulated network (while retaining

at least 10 species) and computed phylogenetic signal in these asymmetrically-

subsampled networks.

Third, both Mantel tests and PBLM neglect the heterogeneity within networks.

Indeed, a non-significant phylogenetic signal at the level of the entire network can

potentially hide a sub-clade presenting significant phylogenetic signal. Alternatively,

a phylogenetic signal in the entire network may be driven by only two sub-clades of

guilds A and B. To explore the potential heterogeneity of the phylogenetic signal

within one guild, one possibility is to apply Mantel tests to the sub-networks formed

by a given sub-clade (*e.g.* Song *et al.* 2020). For each node of the tree of guild A having

at least 10 descendants, we estimated the clade-specific phylogenetic signal using a

Mantel test investigating whether closely related species from this sub-clade of A tend

to interact with similar partners (and *vice-versa* for guild B). Using UniFrac distances,

we performed the Mantel tests with 100,000 permutations, and introduced a

Bonferroni correction for multiple testing to keep a global alpha-risk of 5%. To test this

approach, we generated synthetic networks with known sub-clade signal by

artificially combining networks simulated under neutrality with networks simulated

with the mutualistic parameters **v** (see Results). We grafted each "mutualistic"

phylogenetic tree from guilds A and B within a "neutral" phylogenetic tree by

randomly selecting a branch, such that it creates a separate module with strong

phylogenetic signal. Such simulations could correspond to the evolution of a different

niche, *e.g.* terrestrial *versus* epiphytic plants associating with different mycorrhizal

fungi (Martos *et al.* 2012). We then performed our clade-specific analysis of

phylogenetic signal and investigated in which nodes we recovered significant

phylogenetic signals.

General guidelines and illustration with application on the orchid-fungus

mycorrhizal network from La Réunion

We used our results and other empirical considerations to provide general

guidelines for testing for phylogenetic signal in interaction networks. We illustrated

these guidelines by applying them in a network between orchids and mycorrhizal

fungi from La Réunion island (Martos *et al.* 2012). This network encompasses 70 orchid

species (either terrestrial or epiphytic species) and 93 molecularly-identified fungal

partners (defined according to 97% sequence similarity; Martos *et al.* 2012). We

gathered the maximum-likelihood plant and fungal phylogenies on TreeBASE (Study

Accession S12721), calibrated the orchid phylogeny using a relaxed clock with *chronos*

(Paradis 2013), and arbitrarily added 10 million-years-old polytomies in unresolved

genera to obtained a species-level orchid phylogeny.

Results

Expected phylogenetic signals in species interactions in *BipartiteEvol* networks

The networks simulated using *BipartiteEvol* gave realistic ranges of sizes (from <50 to >250 species; Fig. S2-3).

We found a significant phylogenetic signal in species traits for most antagonistic and neutral simulations (Fig. S4). In contrast, for many mutualistic simulations, closely related species often did not tend to have similar traits, except when α_B=0.01 (*i.e.* mutualistic parameters **iii**, **v**, and **vi**; Fig. S4). When α_B were higher (*i.e.* mutualistic parameters **i**, **ii**, and **iv**, we suspect stabilizing selection to occur and erase the phylogenetic signal in the traits (Maliet *et al.* 2020): we therefore do not expect phylogenetic signal in species interactions for these simulations. In addition, we found an anti-phylogenetic signal in species traits in less than 1% of the simulations (likely to be false-positives given the alpha-risk of Mantel tests; Fig. S4): these networks were thus removed when evaluating the performance of the different approaches and we therefore do not expect anti-phylogenetic signal in species interactions for the remaining networks.

Tests of phylogenetic signal in species interactions in *BipartiteEvol* networks

Using Mantel tests, as expected, we did not find significant phylogenetic signals in species interactions for most neutral networks or for networks with no signal in species traits (Fig. 2, Figs. S5-6-7): the type-I error rate was below 5%, corresponding to the alpha-risk of the test (Table S1), with one notable exception for small networks when using weighted Jaccard distances and Pearson correlations (~8% type-I error). Conversely, we detected a significant unexpected anti-phylogenetic signal in more

than 10% of the simulated networks, in particular in the small ones (Fig. 2, Figs. S5-6-7).

Many mutualistic or antagonistic networks where we expected a phylogenetic signal in species interactions (*i.e.* non-neutral networks with signal in species traits) presented no significant signal with Mantel tests (Fig. 2, Figs. S5-6-7), in particular those simulated with low α_A and α_B values (*e.g.* antagonism **vii**), where non-neutral effects were weak. In mutualistic networks, phylogenetic signals in species interactions were present only when there was a large asymmetry in the effects of trait matching on the fitnesses of the species from guilds A or B (case **v**: α_A=1; α_B=0.01), *i.e.* when only one guild was specialized. Conversely, in antagonistic networks, phylogenetic signals were found mainly when trait matching had a strong impact on the fitness of guild B (the obligate parasites/predators; $\alpha_B \geq 0.1$). Additionally, when phylogenetic signal was significant in one guild, it was generally also significant in the other; in antagonistic networks, the signal was usually higher in guild A compared to guild B (Figs. S5-6-7).

The statistical power of Mantel tests measuring phylogenetic signal in species interactions seems to be modulated by network size, as phylogenetic signals were less often significant but generally stronger in smaller networks (Figs. S5-6-7). Moreover, Mantel tests based on Pearson correlations had higher power than Spearman and Kendall correlations (Figs. S5-6-7) and weighted UniFrac distances outperformed other ecological distances in terms of power (Figs. S5-6-7; Table S2).

When using mean square errors to evaluate the significance of PBLM, we found a significant phylogenetic signal in most of the simulated networks including when we did not expect any (Fig. 2e). The propensity of PBLM to detect phylogenetic signal decreased in large unweighted networks, but the type-I errors remained >30%,

including when using a more stringent significance cutoff (Figs. S8-9). Similar results

were obtained when bootstrapping to evaluate the significance (Fig. S10).

Confounding effect of phylogenetic signal in the number of partners

As expected, tests of phylogenetic signal in the number of partners were non-

significant in the large majority of the *BipartiteEvol* networks, especially the larger ones

(Fig. S11). We did however observe significant correlations between ecological

distances and degree difference distances (Fig. S12). Partial Mantel tests testing for

phylogenetic signal in species interactions while accounting for phylogenetic signal in

the number of partners had similar type-I error and power as simple Mantel tests (Figs.

S5-13; Table S2). Performing sequential Mantel tests decreased the statistical power by

less than 2% (Table S2).

Networks simulated with phylogenetic conservatism in the number, but not the

identity of partners covered a realistic range of sizes (Fig. S14). As expected, Mantel

tests revealed significant phylogenetic signals in the number of partners in >60% of

these networks, with an increasing percentage of significant tests with decreasing a_A

(*i.e.* increasing conservatism in the number of partners; Fig. S18). We found significant

correlations between degree differences and ecological distances in most of these

simulated networks (Fig. S15). As a result, simple Mantel tests testing for phylogenetic

signal in species interactions without accounting for phylogenetic signal in the number

of partners were frequently significant (>30%; Fig. S16; Table S3). Partial Mantel tests

controlling for degree differences slightly decreased the proportion of false-positives,

but it remained high (type-I error >25%; Fig. S17). In addition, partial Mantel tests

detected a spurious significant anti-phylogenetic signal in species interactions in >15%

of the networks (Fig. S17). Conversely, only few networks with a significant simple

Mantel test in species interactions did not produce a significant simple Mantel test in

the number of partners, such that sequential Mantel tests had only a ~7% type-I error rate (Table S3).

Effect of phylogenetic uncertainty, sampling asymmetry, and network heterogeneity

The statistical power of Mantel tests using UniFrac distances decreased, as expected, when the length of the simulated DNA sequences decreased (*i.e.* when phylogenetic uncertainty increased; Fig. S19). However, even when the simulated DNA sequences were the shortest (75 base-pairs), resulting in very noisy reconstructed partners' tree (Fig. S20), the statistical power of the Mantel tests using UniFrac distances remained larger than when using Jaccard distances (Fig. S19).

Our results on the statistical performance of tests of phylogenetic signal were similar when considering sampling asymmetry (Figs. S21-24): PBLM spuriously detected phylogenetic signal when it should not, and Mantel tests had decent statistical performances, especially when using weighted UniFrac distances. In addition, the correlations of the Mantel tests in guild A were generally higher when significant (Fig. S23).

Our clade-specific tests of phylogenetic signal using Mantel tests while correcting for multiple testing recovered a significant phylogenetic signal in 82% of the nodes where mutualism originated (Fig. S25), as well as in most of the ascending nodes. Conversely, we did not find spurious phylogenetic signals in nodes with only neutrally-evolving lineages (Fig. S25).

General guidelines and illustration with application on the orchid-fungus mycorrhizal network from La Réunion

Box 2 provides general guidelines based on our results and empirical

considerations for accurate tests of phylogenetic signal in interaction networks. We

applied these guidelines on the orchid-fungus mycorrhizal network from La Réunion

(available in Martos et al. (2012); Fig. 3). First (step 1), simple Mantel tests of

phylogenetic signal in species interactions for fungi and orchids revealed a significant

but low phylogenetic signal (R<0.10) on the orchid side using Jaccard distances;

however, the significance disappeared with UniFrac distances (Table S4). Similarly,

marginally not-significant and low phylogenetic signals were detected in the

mycorrhizal fungi side (R<0.04; Table S4). Next (step 2), simple Mantel tests of

phylogenetic signal in the number of partners were not significant (p-values>0.05).

Our investigation of clade-specific phylogenetic signals in species interactions in

orchids (option 1) revealed a significant phylogenetic signal in Angraecinae, a sub-

tribe composed of 34 epiphytic species (sequential Mantel test: R=0.37; Bonferroni-

corrected p-value=0.016; Fig. 4) interacting with 53 fungi, suggesting that closely

related Angraecinae tend to interact with more similar mycorrhizal fungi. When we

checked the robustness of the significant phylogenetic signal detected in Angraecinae

(option 2) by subsampling the Angraecinae clade down to 10 species, we still

recovered significant signal in species interactions in both cases (Fig. S26).

Discussion:

We used simulations to perform a comparative analysis of the statistical performances of Mantel tests and the Phylogenetic bipartite linear model (PBLM; Ives & Godfray 2006) for testing for phylogenetic signal in species interactions. Our results highlight the weaknesses of PBLM and partial Mantel tests, and advocate for the use of simple and sequential Mantel tests.

The Phylogenetic bipartite linear model (PBLM) is widely used to test for phylogenetic signal in species interactions, however we found that it has a very high type-I error rate (>30%). PBLM assumes that the interaction strength between two species is determined by the product of two unobserved traits evolving on the phylogenies of guilds A and B respectively, according to two independent Ornstein-Uhlenbeck processes with the selection strengths d_A and d_B (Box 1). PBLM tests the significance of d_A and d_B, which measure the phylogenetic signal of the unobserved traits. A species with a high trait value will have high interaction strengths with many partner species (*i.e.* it is a generalist species), while a species with a low trait value will have low interaction strengths with most partner species, except with the few species with high trait values (*i.e* it is a specialist species). Therefore, we suspect d_A and d_B to measure phylogenetic signals in the number of partners rather than in species interactions. However, we also found significant d_A and d_B in the absence of phylogenetic signal in the number of partners, suggesting that PBLM is sensitive to model misspecification (it relies on strong hypotheses on how the number of partners evolves). In any case, PBLM should not be used as a routine for measuring phylogenetic signal in species interactions. Other model-based approaches inspired from PBLM propose to infer more parameters describing the phylogenetic structure of interactions networks (Rafferty & Ives 2013; Hadfield *et al.* 2014). Unfortunately their computational intensity prevented their incorporation in our comparative analysis.

We found that simple Mantel tests have a moderate statistical power and a
reasonable type-I error rate when testing for phylogenetic signal in species
interactions. Hence, although these tests might fail at detecting low phylogenetic
signal, we can trust their results when they are significant. On the contrary, we found
a high proportion of false-positives (5-10%) when testing for anti-phylogenetic signal;
therefore, detection of such signal in empirical networks should be interpreted with
caution. In addition, Pearson correlations performed better than Spearman and
Kendall correlations, which is somewhat surprising, as correlations between
phylogenetic and ecological distances are not particularly expected to be linear:
Spearman and Kendall correlations have less stringent hypotheses, as they only
assume monotonicity (Box 1), but they probably loose information. We also reported
that using ecological distances that consider interaction abundances and phylogenetic
relatedness of the partners, such as weighted UniFrac distances, significantly improves
the detection of phylogenetic signal, even when reconstructed partners trees are not
robust. Given that species delineation may be somewhat arbitrary, especially for
microbial interactors, and that Jaccard distances are directly sensitive to species
delineation (Sanders *et al.* 2014), we advocate the use of weighted UniFrac distances.
An exception might be if communities of interactors differ mainly in terms of recently
diverged species; in this case Jaccard distances may perform better, as UniFrac
distances emphasize differences in long branches rather than recent splits (Sanders *et*
al. 2014). Finally, we found that multiple simple Mantel tests combined with a
Bonferroni correction perform rather well to investigate clade-specific phylogenetic
signals. Such an approach can therefore be valuable for measuring local phylogenetic
signal in large "meta-networks", such as those describing host-microbiota
phylosymbiosis (Song *et al.* 2020), which likely have heterogeneous phylogenetic
signals across the network.

While simple Mantel tests have satisfactory statistical performances, these tests do
not control for the potential confounding effect of phylogenetic signal in the number

of partners. Partial Mantel tests are frequently used for investigating phylogenetic

signal in species interactions while controlling for signal in the number of partners;

however, we found that they often detected significant signals in species interactions

when we simulated signals in only the number of partners. Thus, partial Mantel tests

fail at discerning whether evolutionary relatedness strictly affects the identity of

partners, independently of the total number of partners associated with each species

(Rezende *et al.* 2007). This corroborates the poor statistical performances of partial

Mantel tests frequently observed in other contexts (Harmon & Glor 2010; Guillot &

Rousset 2013). We therefore suggest to perform sequential simple Mantel tests, testing

first for phylogenetic signal in species interactions, and if significant, testing for

phylogenetic signal in the number of partners. If there is no signal in the number of

partners but a signal in interactions, then we can safely conclude that evolutionary

relatedness strictly affects the identity of partners. This approach has a low type-I error

rate and a very limited power decrease; however, it does not allow testing if there is a

specific signal in species identity when there is a signal in the number of partners. A

hint at whether signal in species interactions is entirely due to signal in the number of

partners or not can be gained by comparing the correlation coefficients obtained when

correlating phylogenetic distance to ecological distance *versus* degree distance.

By definition, phylogenetic signals in species interactions measure general patterns

that are not informative of the processes at play (Losos 2008). A better understanding

of the ecological and evolutionary processes playing a role in the assembly of

interaction networks (Harmon *et al.* 2019) will require developing integrative process-

based approaches, for instance inference machineries for eco-evolutionary models

such as *BipartiteEvol*. Classical inferences (generalized least-squares or likelihood-

based approaches) might be challenging for such complex models (Hadfield *et al.*

2014), but strategies such as machine learning provide promising alternatives.

In the mycorrhizal network from La Réunion, we found non-significant or weak phylogenetic signals in species interactions at the level of the entire orchid-fungus network, suggesting these interactions are generally poorly conserved over long evolutionary timescales (Jacquemyn *et al.* 2011; Martos *et al.* 2012). Conversely, clade-specific Mantel tests detected a significant phylogenetic signal in the Angraecinae epiphytic clade that is experiencing a radiation in La Réunion island. This signal is likely produced by the different orchids genera in Angraecinae associating with specific fungal clades (Martos *et al.* 2012). Thus, our results corroborate a trend toward mycorrhizal specialization in epiphytic orchids compared with terrestrial species (Xing *et al.* 2019), as the epiphytic habitats might require particular adaptations and stronger dependences toward specific mycorrhizal fungi.

Interaction networks are increasingly being analyzed to unravel the evolutionary processes shaping their structure and to predict their stability. Currently-used tools for measuring phylogenetic signals are clearly misleading. We propose instead an alternative approach based on sequential Mantel tests. By emphasizing the limits of current tests of phylogenetic signal, we also hope to stimulate new developments in the statistical adjustment to empirical data of process-based models for the evolution of interaction networks.

Acknowledgments:

The authors acknowledge M. Elias, E. Thébault, and D. de Vienne for helpful discussions. They also thank I. Overcast, S. Lambert, I. Quintero, C. Fruciano, J. Clavel, and A. Silva for comments on an early version of the manuscript. This work was supported by a doctoral fellowship from the École Normale Supérieure de Paris attributed to BPL and the École Doctorale FIRE – Programme Bettencourt. Funding of the research of FM was from the Agence Nationale de la Recherche (ANR-19-CE02-0002). HM acknowledges support from the European Research Council (grant CoG-PANDA).

References:

References:

Aizen, M.A., Gleiser, G., Sabatino, M., Gilarranz, L.J., Bascompte, J. & Verdú, M.

(2016). The phylogenetic structure of plant-pollinator networks increases with

habitat size and isolation. *Ecol. Lett.*, 19, 29–36.

Bascompte, J. & Jordano, P. (2013). *Mutualistic networks. Monogr. Popul. Biol.*

Princeton. Princeton University Press, Princeton.

Bascompte, J., Jordano, P., Melian, C.J. & Olesen, J.M. (2003). The nested assembly of

plant-animal mutualistic networks. *Proc. Natl. Acad. Sci.*, 100, 9383–9387.

Blomberg, S.P., Garland, T. & Ives, A.R. (2003). Testing for phylogenetic signal in

comparative data: Behavioral traits are more labile. *Evolution (N. Y).*, 57, 717–745.

Clavel, J., Escarguel, G. & Merceron, G. (2015). mvMORPH: An R package for fitting

multivariate evolutionary models to morphometric data. *Methods Ecol. Evol.*, 6,

1311–1319.

Elias, M., Fontaine, C. & Frank Van Veen, F.J. (2013). Evolutionary history and

ecological processes shape a local multilevel antagonistic network. *Curr. Biol.*, 23,

1355–1359.

Felsenstein, J. (1985). Phylogenies and the comparative method. *Am. Nat.*, 125, 1–15.

Fontaine, C., Guimarães, P.R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten,

W.H., *et al.* (2011). The ecological and evolutionary implications of merging

different types of networks. *Ecol. Lett.*, 14, 1170–1181.

Fontaine, C. & Thébault, E. (2015). Comparing the conservatism of ecological

interactions in plant–pollinator and plant–herbivore networks. *Popul. Ecol.*, 57,

29–36.

Goslee, S.C. & Urban, D.L. (2007). The ecodist package for dissimilarity-based

analysis of ecological data. *J. Stat. Softw.*, 22, 1–19.

Guillot, G. & Rousset, F. (2013). Dismantling the Mantel tests. *Methods Ecol. Evol.*, 4,

336–344.

Hadfield, J.D., Krasnov, B.R., Poulin, R. & Nakagawa, S. (2014). A tale of two

phylogenies: Comparative analyses of ecological interactions. *Am. Nat.*, 183, 174–187.

Hansen, T.F. & Martins, E.P. (1996). Translating between microevolutionary process and macroevolutionary patterns: The correlation structure of interspecific data. *Evolution (N. Y).*, 50, 1404–1417.

Harmon, L.J., Andreazzi, C.S., Débarre, F., Drury, J., Goldberg, E.E., Martins, A.B., *et al.* (2019). Detecting the macroevolutionary signal of species interactions. *J. Evol. Biol.*, 32, 769–782.

Harmon, L.J. & Glor, R.E. (2010). Poor statistical performance of the Mantel test in phylogenetic comparative analyses. *Evolution (N. Y).*, 64, 2173–2178.

Ives, A.R. & Godfray, H.C.J. (2006). Phylogenetic analysis of trophic associations. *Am. Nat.*, 168, E1–E14.

Jacquemyn, H., Merckx, V.S.F.T., Brys, R., Tyteca, D., Cammue, B.P.A.A., Honnay, O., *et al.* (2011). Analysis of network architecture reveals phylogenetic constraints on mycorrhizal specificity in the genus Orchis (Orchidaceae). *New Phytol.*, 192, 518–528.

Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., *et al.* (2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*, 26, 1463–1464.

Krasnov, B.R., Fortuna, M.A., Mouillot, D., Khokhlova, I.S., Shenbrot, G.I. & Poulin, R. (2012). Phylogenetic signal in module composition and species connectivity in compartmentalized host-parasite networks. *Am. Nat.*, 179, 501–511.

Losos, J.B. (2008). Phylogenetic niche conservatism, phylogenetic signal and the relationship between phylogenetic relatedness and ecological similarity among species. *Ecol. Lett.*, 11, 995–1003.

Maliet, O., Loeuille, N. & Morlon, H. (2020). An individual based model for the eco-evolutionary emergence of bipartite interaction networks. *Ecol. Lett.*, ele.13592.

Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer Res.*, 27, 209–220.

Martín González, A.M., Dalsgaard, B., Nogués-Bravo, D., Graham, C.H., Schleuning,

M., Maruyama, P.K., *et al.* (2015). The macroecology of phylogenetically

structured hummingbird-plant networks. *Glob. Ecol. Biogeogr.*, 24, 1212–1224.

Martos, F., Munoz, F., Pailler, T., Kottke, I., Gonneau, C. & Selosse, M.-A.A. (2012).

The role of epiphytism in architecture and evolutionary constraint within

mycorrhizal networks of tropical orchids. *Mol. Ecol.*, 21, 5098–5109.

Morlon, H., Lewitus, E., Condamine, F.L., Manceau, M., Clavel, J. & Drury, J. (2016).

RPANDA: An R package for macroevolutionary analyses on phylogenetic trees.

Methods Ecol. Evol., 7, 589–597.

Münkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schiffers, K., *et al.*

(2012). How to measure and test phylogenetic signal. *Methods Ecol. Evol.*, 3, 743–

756.

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*.

Paradis, E. (2013). Molecular dating of phylogenies by likelihood methods: A

comparison of models and a new information criterion. *Mol. Phylogenet. Evol.*, 67,

436–444.

Paradis, E., Claude, J. & Strimmer, K. (2004). APE: Analyses of phylogenetics and

evolution in R language. *Bioinformatics*, 20, 289–290.

Peralta, G. (2016). Merging evolutionary history into species interaction networks.

Funct. Ecol., 30, 1917–1925.

Perez-Lamarque, B. & Morlon, H. (2019). Characterizing symbiont inheritance during

host–microbiota evolution: Application to the great apes gut microbiota. *Mol.*

Ecol. Resour., 19, 1659–1671.

R Core Team. (2020). R: A language and environment for statistical computing.

Rafferty, N.E. & Ives, A.R. (2013). Phylogenetic trait-based analyses of ecological

networks. *Ecology*, 94, 2321–2333.

Revell, L.J. (2012). phytools: An R package for phylogenetic comparative biology

(and other things). *Methods Ecol. Evol.*, 3, 217–223.

Rezende, E.L., Lavabre, J.E., Guimarães, P.R., Jordano, P. & Bascompte, J. (2007).

627 Non-random coextinctions in phylogenetically structured mutualistic networks.

628 *Nature*, 448, 925–928.

629 Rohr, R.P. & Bascompte, J. (2014). Components of phylogenetic signal in antagonistic

630 and mutualistic networks. *Am. Nat.*, 184, 556–564.

631 Sanders, J.G., Powell, S., Kronauer, D.J.C., Vasconcelos, H.L., Frederickson, M.E. &

632 Pierce, N.E. (2014). Stability and phylogenetic correlation in gut microbiota:

633 lessons from ants and apes. *Mol. Ecol.*, 23, 1268–1283.

634 Smouse, P.E., Long, J.C. & Sokal, R.R. (1986). Multiple regression and correlation

635 extensions of the mantel test of matrix correspondence. *Syst. Zool.*, 35, 627–632.

636 Song, S.J., Sanders, J.G., Delsuc, F., Metcalf, J., Amato, K., Taylor, M.W., *et al.* (2020).

637 Comparative analyses of vertebrate gut microbiomes reveal convergence

638 between birds and bats. *MBio*, 11, 1–14.

639 Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009). Evaluating multiple determinants

640 of the structure of plant–animal mutualistic networks. *Ecology*, 90, 2039–2046.

641 Xing, X., Jacquemyn, H., Gai, X., Gao, Y., Liu, Q., Zhao, Z., *et al.* (2019). The impact of

642 life form on the architecture of orchid mycorrhizal networks in tropical forest.

643 *Oikos*, 128, 1254–1264.

644 Xing, X., Liu, Q., Gao, Y., Shao, S., Guo, L., Jacquemyn, H., *et al.* (2020). The

645 architecture of the network of orchid–fungus interactions in nine co-occurring

646 Dendrobium species. *Front. Ecol. Evol.*, 8, 1–10.

 Text boxes:

Box 1: Methods for measuring phylogenetic signal in species interactions:

Mantel tests (Mantel 1967) were introduced to examine the correlation between

two dissimilarity matrices. They have been used to measure the phylogenetic signal in

species interactions by computing the correlation between the matrix of phylogenetic

distances (for species pairs from guild A for example, Figure 1) and the matrix of

'ecological dissimilarities' between the sets of interacting partners (*i.e.* species from

guild B interacting with species pairs from guild A). The correlation (R) with -1<R<1 is

often evaluated using Pearson correlation, that is the mean of the products of the

corresponding elements in the two standardized dissimilarity matrices. Alternatively,

R can be evaluated using Spearman correlation (computed from dissimilarities

transformed into ranks), or Kendall correlation (computed by counting the pairs of

observations that have the same rank). The parametric Pearson correlation is

statistically more powerful, but makes stronger hypotheses (it assumes a linear

relationship) than the non-parametric Spearman and Kendall correlations (which

assume only a monotonic relationship). A positive (resp. negative) correlation

indicates a phylogenetic (resp. anti-phylogenetic) signal in species interactions. Its

significance is evaluated using randomizations by repeatedly permuting one of the

original dissimilarity matrices: one-tailed p-values are obtained by comparing the rank

of the original correlation R to the randomized correlations. Ecological dissimilarities

of interacting partners between two species from guild A can be measured with

various indices. Two classical indices are the Jaccard distance, defined as the number

of their unshared partners from guild B divided by their total number of partners, and

the UniFrac distance, which incorporates phylogenetic relatedness between partners,

computed as the fraction of unshared branch length in the phylogenetic tree of their

partners from guild B. Both indices also have a weighted version that accounts for

interaction strength (e.g. weighted Jaccard and generalized/weighted UniFrac; Chen

et al. 2012).

Partial Mantel tests examine the correlation between two dissimilarities

matrices while accounting for a third dissimilarity matrix (Smouse *et al.* 1986). When

testing for phylogenetic signal in bipartite interaction networks, these tests are useful

for controlling the phylogenetic signal in the number of partners; indeed similarity in

the number of partners can decrease the value of ecological dissimilarity metrics

independently of the identity of the partners, such that a phylogenetic signal in the

number of partners can generate a phylogenetic signal in species interactions that is

not linked to an evolutionary conservatism of interacting partners. Partial Mantel tests

therefore investigate the correlation between phylogenetic and ecological

dissimilarities while controlling for the absolute difference in the degrees (*i.e.* the

number of partners) between species pairs (Rezende *et al.* 2007).

The Phylogenetic bipartite linear model (PBLM; Ives & Godfray 2006) assumes

that interaction strengths between species from guilds A and B are determined by

(unobserved) traits that evolve on the two phylogenies each following a simplified

Ornstein-Uhlenbeck process (Blomberg *et al.* 2003). The strength of interaction between

two species is assumed to be given by the product of their two traits. Under these

assumptions, d_A and d_B can be estimated from the two phylogenies and the matrix of

interaction strengths using generalized least squares (Ives & Godfray 2006). d_A and d_B

are then interpreted as a measure of phylogenetic signal in species interactions. If d=1,

the traits evolve as Brownian processes; if d=0, there is no effect of the phylogeny

(similar as evolution on a star phylogeny); 0<d<1 generates stabilizing selection and

d>1 disruptive selection. Ives & Godfray (2006) proposed two approaches to assess the

significance of the signal. The simplest consists in comparing the mean square errors

(MSE) of the generalized least squares regression to the same MSE obtained using star

phylogenies (MSE_{star}). $MSE<MSE_{star}$ is interpreted as a significant phylogenetic signal

in species interactions. The second approach uses a bootstraping strategy to build 95%

confidence intervals around the estimated d_A and d_B values: the null hypothesis

(absence of phylogenetic signal in guild A, resp. B) is rejected if the confidence interval

around d_A (resp. d_B) does not incude 0. While designed primarily for applications to

bipartite networks characterized by matrices of interaction strengths (*e.g.* net attack

rate of a parasitoid on its hosts), PBLM has been applied to weighted networks

characterized by matrices of interaction abundance (*i.e.* the number of times the

interaction has been observed) and unweighted (binary) networks, using 1 for the

interaction strength when species interact and 0 otherwise (Ives & Godfray 2006;

Vázquez *et al.* 2009; Jacquemyn *et al.* 2011; Martos *et al.* 2012; Xing *et al.* 2020).

Box 2: Recommended guidelines to measure phylogenetic signal in species interactions within bipartite ecological networks.

This guideline is composed of two fixed steps followed by two optional ones and can be applied as soon as a bipartite interaction network (with or without abundances) and at least the phylogenetic tree of guild A are available. The phylogenetic tree does not need to be binary, rooted, or ultrametric.

Step 1: The first step consists in testing for phylogenetic signal in species interactions for guild A (*i.e.* whether closely related species from guild A tend to interact with similar partners from guild B) using a one-tailed simple Mantel test. This step requires to pick an ecological distance (UniFrac distances are recommended compared to Jaccard distances) and a type of correlation (Pearson correlation by default).

Step 2: Next, to assess whether a phylogenetic signal in species interactions really comes from the identity of species interactions, the second step consists in testing whether there is phylogenetic signal in the number of partners of guild A (*i.e.* whether closely related species from guild A tend to interact with the same number of partners from guild B) using a one-tailed simple Mantel test.

Option 1: Clade-specific phylogenetic signal in guild A can be tested using simple Mantel tests while correcting for multiple testing (*e.g.* Bonferroni correction). It can be used to test whether some clades present different intensities of phylogenetic signal (*e.g.* because of higher specificity).

Option 2: The robustness of the findings can be tested by looking at how the conclusions might be affected by phylogenetic uncertainty (e.g. using a Bayesian posterior of tree) or sampling bias. The potential effect of sampling bias can be investigated by subsampling all clades to the same number of species.

744   If a phylogenetic tree for guild B is available, all these steps can be replicated to test

745   for phylogenetic signal in species interaction in guild B.

Figures:

Figure 1: Illustration of the data used to test for phylogenetic signal in species interactions

Toy example of an interaction network between orchids (in green) and mycorrhizal fungi (in brown) with associated phylogenetic trees. The bipartite interaction network between two guilds A (here the orchids) and B (the fungi) is represented by a matrix which elements indicate either whether or not species interact (*i.e.* 1 if they do and 0 otherwise, 'unweighted' or 'binary' network) or the frequency of the interaction ('weighted' network; for example here we indicated the number of times a given pairwise interaction has been observed using shades of gray from white (no interaction) to dark gray (many interactions)). Each guild is also characterized by a rooted phylogenetic tree, used to compute phylogenetic distances between pairs of species.

Figure 2: Statistical performances of the simple Mantel tests and the Phylogenetic bipartite linear model (PBLM; Ives & Godfray, 2006)

For each panel, the simulations are divided between networks where phylogenetic signal in species interactions is expected (*i.e.* networks (i) simulated with an effect of the traits on individual fitness - antagonistic and mutualistic simulations - and (ii) presenting traits that are phylogenetically conserved – see Supplementary Figure 2) and networks where phylogenetic signal in species interactions is not expected (*i.e.* neutral simulations ($\alpha = 0$) or simulated networks where we observed no phylogenetic signal in the traits).

a-d: Phylogenetic signals in species interactions estimated using simple Mantel tests with Pearson correlation (R) in the guilds A (a, c) and B (b, d). The different panels in rows correspond to the 2 tested ecological distances: weighted Jaccard (a, b) or weighted UniFrac (c, d) distances. One-tailed Mantel tests between phylogenetic distances and ecological distances were performed using 10,000 permutations. In each panel, the bars indicate the percentage of simulated networks that present a significant positive correlation (p-value>0.05 for the test of phylogenetic signal), a significant negative correlation (in red; p-value>0.05 for the test of anti-phylogenetic signal), or no significant correlation (in yellow; both p-values>0.05).

e: Phylogenetic signals estimated using PBLM. For a given combination of parameters, the bar indicates the percentage of simulated networks that present no significant (in yellow; $MSE \geq MSE_{star}$) or a significant (green; $MSE < MSE_{star}$) phylogenetic signal. Phylogenetic signals are shaded from light green to dark green according to the strength of the signal (e.g. in dark green if $d_A > 0.15$ or $d_B > 0.15$). PBLM were run on the weighted networks.

In each panel, the first bar indicates the statistical power of the test, whereas the second and third bar indicate the type-I error rate of the test.

Figure 3: Illustration of the recommended guidelines for measuring phylogenetic

signal in species interactions within bipartite interaction networks (Box 2).

At the top, a toy example of an interaction network between orchids (in green) and

mycorrhizal fungi (in brown) is informed with the phylogenetic trees of each guild.

For each step of the guidelines (Box 2), an example of the corresponding function

available in the R-package RPANDA is indicated in grey.

guild A

guild B

Interaction network

Phylogenetic signal in guild A:

Step 1: test the phylogenetic signal in the **species interactions** (simple Mantel test)
(i) choice of ecological distances (Jaccard, UniFrac…)
(ii) with or without interaction abundances

phylosignal_network(network, tree_A, tree_B, method = "GUniFrac", correlation = "Pearson")

Step 2: test the phylogenetic signal in the **degree of generalism** (simple Mantel test)

phylosignal_network(network, tree_A, method = "degree", correlation = "Pearson")

Option 1: investigate clade-specific phylogenetic signals (simple Mantel tests with Bonferroni correction)

phylosignal_sub_network(network, tree_A, tree_B, method = "GUniFrac", correlation = "Pearson")

Option 2: test the robustness of the findings to phylogenetic uncertainty and/or sampling bias

(repeat for guild B)

Figure 4: Empirical application on an orchid-fungus interaction network from La
Réunion island (Martos *et al.,* 2012): the clade-specific analyses of phylogenetic
signal in species interactions revealed a significant phylogenetic signal in the
epiphytic subtribe Angraecinae.

The orchid phylogeny (Martos *et al.,* 2012) is represented with its nodes colored
according to the results of the Mantel test performed on the corresponding sub-
network: in blue if non-significant, in grey when the node has less than 10 descendent
species (the Mantel test was not performed), and in red when the phylogenetic signal
is significant. Each one-tailed simple Mantel test was performed using the Pearson
correlation and 100,000 permutations and its significance was evaluated while
correcting for multiple testing (Bonferroni correction).

For each species, its habitat (terrestrial or epiphytic) is indicated at the tips of the tree
and the main orchid clades are highlighted in colors. Only the genera are indicated at
the tips of the tree (see Supplementary Figure S28 for the species list).