Disentangling the flow of signals between populations of neurons

Evren Gokcen[1], Anna I. Jasper[2], João D. Semedo[1], Amin Zandvakili[2],

Adam Kohn[2,3,4†], Christian K. Machens[5†], and Byron M. Yu[1,6†*]

[†]These authors contributed equally to this work.

[*]Corresponding author.

[1]Dept. of Electrical and Computer Engineering, Carnegie Mellon University, Pittsburgh, PA, USA

[2]Dominick Purpura Dept. of Neuroscience, Albert Einstein College of Medicine, Bronx, New York, NY, USA

[3]Dept. of Ophthalmology and Visual Sciences, Albert Einstein College of Medicine, Bronx, New York, NY, USA

[4]Dept. of Systems and Computational Biology, Albert Einstein College of Medicine, Bronx, New York, NY, USA

[5]Champalimaud Research, Champalimaud Centre for the Unknown, Lisbon, Portugal

[6]Dept. of Biomedical Engineering, Carnegie Mellon University, Pittsburgh, PA, USA

Abstract

Technological advances have granted the ability to record from large populations of neurons across multiple brain areas. These recordings may illuminate how communication between areas contributes to brain function, yet a substantial barrier remains: How do we disentangle the concurrent, bidirectional flow of signals between two populations of neurons? We therefore propose here a novel dimensionality reduction framework: Delayed Latents Across Groups (DLAG). DLAG disentangles signals relayed in both directions; identifies how these signals are represented by each population; and characterizes how they evolve over time and trial-to-trial. We demonstrate that DLAG performs well on synthetic datasets similar in scale to current neurophysiological recordings. Then we study simultaneously recorded populations in primate visual areas V1 and V2, where DLAG reveals signatures of concurrent yet selective communication. Our framework lays the foundation for dissecting the intricate flow of signals across populations of neurons, and how this signaling contributes to cortical computation.

Introduction

Simultaneous recordings from large populations of neurons across multiple brain areas are growing in availability [1–4]. These recordings present opportunities to illuminate how inter-areal communication enables basic brain function [5], but they also present substantial conceptual and statistical challenges. Brain areas involved in sensory [6–9], cognitive [10], and motor functions [11] are often reciprocally connected: signals are relayed not only from one area to the next, but bidirectionally, and likely concurrently. Recorded neural activity, however, provides only a tangled view of this concurrent communication (Fig. 1, top): individual neurons simultaneously reflect an area's inputs, outputs, and ongoing internal computations [12].

Determining the flow of signals between recorded areas is therefore a nontrivial task; but importantly, inter-areal communication is not instantaneous. The physiological properties of axons and synapses introduce delays in signal transmission. These delays provide a working definition of signal flow, which underlies many approaches to studying inter-areal communication. For instance, several studies have compared the timing of the onset of neural responses [13–16] or of the emergence of selectivity attributable to top-down processes [17–22] across areas following the presentation of a stimulus. Other studies, leveraging simultaneous recordings, have measured temporal delays between two areas through pairwise spiking correlations [23–28] and information theoretic measures [29, 30]. Similarly, inter-areal phase delays have been measured through the coherence between local field potentials (LFP) [31–34]. These timing-based approaches have significantly advanced our understanding of how signals propagate across brain areas. However, given that neuronal computations are believed to be carried out by neuronal populations, such approaches—focused largely on pairs of neurons or aggregate measures of neural activity—lack the richness to fully describe the nature of signals relayed between areas.

A full characterization of inter-areal signal flow therefore requires relating the activity of populations of neurons across two or more areas—a challenging high-dimensional problem. Dimensionality reduction techniques capable of identifying low-dimensional patterns of population activity common to two or more recorded areas are thus increasingly prevalent [35, 36]. These techniques

have driven new proposals for population-level mechanisms of gating between motor cortex output and muscle movement [37, 38]; selective communication between cortical areas [39, 40]; enhanced communication of stimulus information with attention [41]; and the robustness of certain local computations to perturbations upstream [42].

The relationship between the population activity patterns identified in these studies and the flow of inter-areal signals, however, remains unclear: is the observed inter-areal communication driven by one area or both, and when does this (potentially bidirectional) communication occur? If communication were to occur in one direction at a time, then existing dimensionality reduction methods could, in principle, provide further insight into these questions. If two areas were to communicate in both directions concurrently, however, then existing methods would only identify the dominant direction of population-level signal flow [43]. Disentangling the concurrent flow of signals between populations remains a substantial barrier in neuroscience (Fig. 1, bottom left).

We therefore propose a novel dimensionality reduction framework: Delayed Latents Across Groups, or DLAG (Fig. 1, bottom right). DLAG disentangles signals relayed in both directions; identifies how these signals are represented by each population; and characterizes how they evolve over time and trial-to-trial. We first demonstrate that DLAG performs well on synthetic datasets similar in scale to current neurophysiological data. Then we study simultaneously recorded populations in primate visual areas V1 and V2, where DLAG reveals that V1-V2 interactions are selective and asymmetric. DLAG therefore unlocks new opportunities to investigate the concurrent flow of signals between populations of neurons and how inter-areal communication contributes to brain function.

Results

Delayed Latents Across Groups (DLAG)

Suppose we observe the activity of two populations of neurons (Fig. 2, left column), measured as, for example, the number of spikes counted within nonoverlapping time bins. Here we will take

these populations as belonging to two different brain areas, A and B; however, in principle, they can belong to any meaningful groups, such as cortical layer or cell type. DLAG functionally dissects the observed population activity in each area into a linear combination (weighted sum) of two types of latent variables (Fig. 2, center column).

The first type of latent variable, *across-area* variables, describes interactions among neurons in both areas (illustrated by the magenta box spanning both areas in Fig. 2). Critically, across-area variables are paired, and associated with a relative time delay (Fig. 2, D_1 between the first pair and D_2 between the second pair). The across-area variables evolve smoothly over time, and different latent variables may vary more rapidly or more slowly, as determined by the neural recordings. If a particular time course is reflected in the population activity of area A, and a similar time course is reflected in the population activity of area B, but after a time delay, then an across-area variable pair can describe the apparent flow of that signal from A to B. And if, concurrently—and along a distinct set of population activity patterns—a time course is first reflected in area B, followed by area A, a second across-area variable pair can also describe the flow of that inter-areal signal. In fact, DLAG can employ potentially many across-area variable pairs, each with a delay of its own sign and magnitude, to capture multiple concurrent streams of signal flow between the two populations at different timescales.

The second type of latent variable, *within-area* variables, describes interactions among neurons within one area that cannot be explained by interactions with neurons in the other area (Fig. 2; blue: within-area A; red: within-area B). Like across-area variables, they evolve smoothly over time, and different latent variables may vary more rapidly or more slowly, as determined by the neural recordings. Within-area variables are also critical to the estimation of across-area variables (see Methods and Supplementary Discussion).

The relationship between within- and across-area latent variables and observed population activity in each area can be represented geometrically with the concept of a population space (Fig. 2, right column). For each area, we can define a high-dimensional population activity space where each axis represents the activity of one neuron. Each point in the space represents the population activity at a particular time; and the points trace out a trajectory over time. DLAG's two types

of latent variables each define the axes (dimensions) of a low-dimensional plane (subspace) in this high-dimensional space (in Fig. 2 we show only the across-area subspaces, for visual clarity). We interpret each dimension of these subspaces as representing a particular spatial pattern of population activity whose time course is described by the corresponding latent variable.

The temporal structure of within- and across-area variables are both described by relating each latent variable at different time points through Gaussian processes (Supplementary Fig. 1). Each Gaussian process is associated with its own characteristic timescale that controls the temporal smoothing of neural activity (Supplementary Fig. 2). DLAG estimates both these timescales and time delays from the neural activity. And finally, after the DLAG model parameters are estimated from the neural activity, the smoothed time courses of within- and across-area latent variables can be studied on a trial-to-trial basis.

Validation on realistic-scale synthetic data

Before applying DLAG to experimental data, we characterized its performance on synthetic datasets similar in scale to state-of-the-art neurophysiological recordings from multiple brain areas, and on additional synthetic datasets covering a wider range of conditions. Informed by our recordings in macaque V1 and V2 [28, 39] (see next section), we simulated independent datasets with representative numbers of neurons (area A: 80; area B: 20); trial counts (100); trial lengths (1,000 ms); and "signal-to-noise" ratios, where "signal" is defined as the shared activity described by latent variables. We set the combined total dimensionality in each area to representative values, but varied the relative numbers of within- and across-area latent variables across datasets (see Methods for additional details). A selection of within- and across-area time courses (black solid traces) from a representative trial and dataset are displayed in Fig. 3a.

Importantly, among these datasets, we included datasets with an absence of across- or within-area structure, to test if our framework could identify such cases. Specifically, we included datasets for which across-area dimensionality was zero: the two areas are independent, and any correlations between neurons are purely within-area. We also included datasets for which within-area

dimensionality was zero: all variance shared among neurons in one area can be attributed to their interactions with neurons in the other area.

Scalable model selection   In practice, the optimal number of within- and across-area latent variables is not known *a priori*, and must be estimated from data. Selecting the number of within- and across-area variables, however, is a challenging problem of scale. For example, performing a grid search over just 10 possibilities for each type of latent variable would result in 1,000 model candidates. We therefore developed a streamlined cross-validation procedure that significantly improves scalability (described in Methods).

Applied to the synthetic datasets described above, our cross-validation procedure proved highly accurate. Across all datasets—including those with no across- or within-area structure—the selected dimensionalities matched the ground truth (Fig. 3b, across-area; Fig. 3c, within-area). Furthermore, model selection remained accurate in additional, more challenging scenarios, where we considered synthetic datasets with significantly lower signal-to-noise ratios (lower than typically encountered in our V1-V2 recordings, below) (Supplementary Fig. 3). And, in the instances where the greater statistical challenge induced imperfect estimates of dimensionality, DLAG's parameter and latent variable estimates remained stable (Supplementary Fig. 4, Supplementary Fig. 5).

Estimation of parameters and latent variables   We next characterized the performance of the optimal DLAG models, selected above. Across all datasets, within- and across-area latent time courses (Fig. 3a); delays (Fig. 3d); across-area Gaussian process (GP) timescales (Fig. 3e); and within-area GP timescales (Fig. 3f, Fig. 3g) were all consistently and accurately estimated. We highlight, in particular, DLAG's ability to estimate time delays between the two areas (Fig. 3d). Delay error was 1.3±0.1 ms (mean and SEM across all delays; max error 7.0 ms), despite observations occurring at 20 ms time steps. This accuracy emphasizes an important feature of the DLAG model that distinguishes it from other time series modeling approaches (see Discussion). Because latent time courses and time delays are continuous-valued, DLAG can leverage the cor-

related activity of the neuronal populations to recover delays with sub-sample period (or spike count bin width, in the case of spiking neural activity) resolution.

The synthetic datasets presented here were generated with a variety of parameters representative of realistic data, but we also verified that DLAG performed predictably well over a wider range of simulated conditions. Specifically, we systematically characterized DLAG's performance as a function of number of trials (Supplementary Fig. 6); number of neurons (Supplementary Fig. 7); and latent timescale (Supplementary Fig. 8). We also characterized the runtime of the DLAG fitting procedure as a function of number of trials, trial length, number of neurons, and latent dimensionality (Supplementary Fig. 9).

Dissecting bidirectional interactions between V1 and V2

We then used DLAG to study interactions between two areas in the early visual system: V1 and V2. V1 and V2 share strong reciprocal connections [44, 45] and strongly interact [26, 28, 39], but the bidirectional nature of their interactions is not yet well understood. We simultaneously recorded the activity of neuronal populations in the superficial (output) layers of V1 (61 to 122 neurons; mean 86.3), and the middle (input) layers of V2 (15 to 32 neurons; mean 19.6) in three anesthetized monkeys (Fig. 4a; data reported previously in [28, 39]). Recording locations were selected to maximize the probability that the recorded V1 and V2 populations directly interacted by ensuring spatial receptive field alignment. We analyzed neuronal responses measured during the 1.28 s presentation of drifting sinusoidal gratings of different orientations, and counted spikes in 20 ms time bins. The temporal period of the drifting gratings (160 ms) is evident in peristimulus time histograms (PSTHs) for an example recording session and stimulus condition (Fig. 4b).

In total, we fit DLAG models separately to 40 "datasets," corresponding to five recording sessions, each with eight different orientations. Additionally, to provide an intra-areal contrast against the DLAG models dissecting V1-V2 interactions, we also applied DLAG to V1 only (Fig. 4c). For each dataset, we randomly split V1 into two equally sized subpopulations ("V1a" and "V1b"). We then applied DLAG to dissect these "V1a-V1b" interactions in a manner identical to V1-V2.

V1-V2 interactions are selective and affect V1 and V2 differently   We first used DLAG to study whether V1 and V2 interact selectively: are there activity patterns in V1 and in V2 that are not shared between the two areas? Representative single-trial latent variable time courses suggest that V1 and V2 do interact selectively (Fig. 5a). There exist significant within-area latent variables (i.e., latent variables that were selected via cross-validation), consistent with dimensionalities across all datasets (Fig. 6a; median dimensionality within-V1: 14; within-V2: 2; across: 3). Furthermore, upon scaling the amplitude of each latent variable (Fig. 5a) by the fraction of shared variance it explains in its corresponding area, the within-area variables in both V1 and V2 appear to explain shared variance comparable in magnitude to that explained by across-area variables. Indeed, across all datasets, across-area activity explained only a portion of the overall shared variance in V1 and in V2 (Fig. 6b; median: 34% in V1; 76% in V2). Collectively, these observations are consistent with the presence of a communication subspace between V1 and V2 [39], through which only a subset of activity patterns are shared between the two areas.

Interestingly, across-area activity was more dominant in V2 than it was in V1 (Fig. 6b, points above the diagonal), a phenomenon that could not be fully attributed to differences in recorded population size or in the total dimensionality of each area (Supplementary Fig. 10). This discrepancy in across-area strength might be a consequence of the cortical layers from which we recorded: much of the activity in the middle layers of V2 is likely driven by V1. The superficial layers of V1, on the other hand, receive input from other areas that do not also project to the middle layers of V2.

V1a and V1b, by contrast, do not appear to interact selectively. The latent variables within-V1a and within-V1b ("within-population" variables) explain little shared variance relative to "across-population" variables (Fig. 5b). V1a-V1b across-population activity was consistently higher-dimensional (Fig. 6c; median dimensionality within-V1a: 2; within-V1b: 1; across: 11), and accounted for nearly all of the overall shared variance in V1a and in V1b (Fig. 6d; median: 96% in V1a; 98% in V1b).

Given the selectivity with which V1 and V2 interact, what is the nature of activity not shared between the two areas? Stereotyped periodic signals, for instance, whose temporal periods match

the period of the drifting grating presented, appear strongly within-V1 (Fig. 5a, "Within 1" and "Within 2", top); and to the extent that they do appear in both areas, they appear only weakly in V2 (Fig. 5a, "Across 3"). Periodic signals also appear strongly in both V1a and V1b (Fig. 5b, "Across 1" and "Across 2") and, unlike V1-V2, are not evident in within-population variables. The prominence of this stimulus-related periodic structure in V1 relative to V2 is consistent with the stimulus response properties of neurons in each area, evident in the neuronal PSTHs (Fig. 4b).

V1-V2 interactions are bidirectional and asymmetric   We next used DLAG to study the bidirectional nature of interactions between V1 and V2. We therefore focus on DLAG's time delays, which identify whether an interaction is feedforward (positive delay: V1 to V2) or feedback (negative delay: V2 to V1). For example, the first representative V1-V2 across-area variable (Fig. 5a, "Across 1") is associated with a -23 ms delay, implying a feedback interaction. In contrast, the visually similar V1a-V1b across-population variable (Fig. 5b, "Across 3") is associated with a 0 ms delay. We developed a statistical procedure to test whether delays such as these significantly deviate from zero (described in Methods). Indeed, the directionality of this latent variable was identified as statistically "ambiguous" (i.e. not significantly different from zero, indicated by the bidirectional gray arrow in Fig. 5b). A V1a-V1b delay at or near zero is expected, given that the V1a and V1b populations belong to the same area, V1, and likely receive common inputs with similar latencies (in contrast to the populations in distinct areas V1 and V2).

Delays across all datasets reflected bidirectional interactions between V1 and V2 (Fig. 6e). Like the strength of across-area activity observed in V1 and in V2, above, the distribution of delays might also reflect the cortical layers from which we recorded. The superficial layers of V1 directly project to the middle layers of V2, consistent with the typically short positive delays [26, 28]; whereas several intermediate connections likely lie between the middle layers of V2 and the superficial layers of V1, consistent with the typically longer negative delays. But notably, the delays between V1 and V2 exhibited a striking asymmetry. The interactions across these areas were predominantly directed from V2 to V1 (Fig. 6e; median over "non-zero" delays: -8 ms; median over all delays: -5 ms). This asymmetry remained even when we subsampled the V1 population to match V2 in size, and re-applied DLAG (Supplementary Fig. 10).

V1a-V1b interactions, in contrast, were symmetric (Fig. 6f; median over "non-zero" delays: -2 ms; median over all delays: 0 ms; neither median significantly different from zero). This centering of the delay distribution around zero is again expected, given that the neurons in V1a and V1b were randomly chosen and belong to the same area. Still, the magnitudes of V1a-V1b delays were not universally zero. These non-zero delays likely reflect aggregate differences in the stimulus response properties of the randomly chosen V1a and V1b subpopulations. For example, inspection of PSTHs (Fig. 4b) suggests that the phase of trial-averaged periodic structure can vary by tens of ms between individual V1 neurons (see Discussion).

Timescales increase from V1 to V2   Finally, how might the timescales of neural activity differ between V1 and V2? Within-V2 Gaussian process (GP) timescales were consistently longer than within-V1 GP timescales (Fig. 6g; median within-V1: 24 ms; within-V2: 74 ms). Within-V1a and within-V1b GP timescales, on the other hand, were nearly the same (Fig. 6h; median within-V1a: 20 ms; within-V1b: 23 ms). Across-area GP timescales were also typically longer for V1-V2 (Fig. 6e; median over non-zero and ambiguous delays: 37 ms) than for V1a-V1b (Fig. 6f; median over non-zero and ambiguous delays: 29 ms). V1-V2 temporal structure again remained intact for analyses with V1 and V2 populations of equal size (Supplementary Fig. 10). Overall, these observations are consistent with previous evidence that timescales increase for areas higher up the cortical hierarchy [46].

Discussion

Two areas in the early visual system, V1 and V2, have served as an excellent testbed for our framework. To our knowledge, our application of DLAG to these areas is the first time observations have been made of persistent, concurrent interactions from spiking activity of neuronal populations. DLAG uncovered signatures of inter- and intra-areal interaction that are largely consistent with previous work, such as the selectivity with which V1 and V2 interact [39], as well as an increase in timescale moving up the cortical hierarchy, from V1 to V2 [46].

However, DLAG provided a novel ability to study the bidirectional nature of interactions between these areas, and characterize these interactions on a trial-to-trial and moment-to-moment basis. DLAG identified population-level interactions in both directions, whose strengths and associated time delays appear to reflect the cortical layers from which we recorded. But surprisingly, it also revealed a marked asymmetry, biased in the feedback (V2 to V1) direction. In contrast, anatomically, feedback inter-cortical connections equal feedforward (V1 to V2) connections in number [44, 45], while feedback signals appear modulatory (rather than driving) in nature [47]. This apparent disparity presents an opportunity for future study.

While we applied DLAG to electrophysiological recordings (i.e., spiking activity) of populations in distinct brain areas, we emphasize that DLAG is general: in principle, DLAG is applicable to any high-dimensional time series data. Hence, it can be applied not only to electrophysiological recordings, but any recording type (e.g., calcium imaging). It can also be applied not only to populations in distinct brain areas, but any meaningfully distinct subpopulations, such as different cortical layers or cell types. In fact, both observed groups need not be neuronal populations: one could use DLAG to study, for example, the relationship between a neuronal population and a dynamic stimulus or behavioral variables.

Relation to previous statistical methods   Through dimensionality reduction, DLAG directly reveals inter-areal interaction structure that would not obviously emerge from other multivariate time series methods, such as Granger causal modeling [48–50], Generalized Linear Models [40, 51, 52], or recurrent neural networks [53]. For instance, the selectivity with which V1 and V2 communicate—evident through DLAG's across- and within-area dimensionalities (Fig. 6a-d)— might have been difficult to characterize otherwise. We were also able to uncover the spatiotemporal structure of activity within V1 and within V2 (i.e., activity not shared across areas) not only across datasets (Fig. 6), but also on a trial-to-trial and moment-to-moment basis (Fig. 5). This characterization would have been difficult with existing time series methods that incorporate dimensionality reduction but discard within-area activity as noise [54, 55].

DLAG also offers unique advantages when characterizing the temporal structure of activity within and across areas. Applied to V1 and V2, DLAG uncovered latent variables with diverse temporal profiles and timescales. The ability to capture diverse dynamical motifs stems from DLAG's definition via Gaussian processes [56]: beyond temporal smoothness, DLAG makes no additional assumptions about the form of dynamics within or across areas. In contrast, multi-area methods proposed by [57] and [58], for instance, assume interactions evolve over time according to particular parametric (e.g., linear) dynamical models. Gaussian processes provide DLAG with another advantage [59]: the ability to discover wide-ranging delays with high precision. Existing multi-area methods (all of which, above, are defined in discrete-time) are limited to delays restricted to be integer multiples of the sample period or spike count bin width of neural activity. To approach the flexibility and precision of DLAG's delay discovery, these methods must overcome additional computational and statistical challenges for both model selection and fitting.

With the conceptual and statistical advantages described above, DLAG is a potentially powerful tool for exploratory data analysis; and it is ultimately complementary with other approaches. For example, given a new dataset, one might use DLAG to generate data-driven hypotheses about plausible dynamical structures within and across areas, and then test these hypotheses using a dynamical system-based approach. Data-constrained recurrent networks, as in [53], may also be particularly powerful tools moving forward: a one-to-one correspondence between artificial and recorded neurons offers a mechanistic description of neuronal interactions within and across areas, and a potential bridge to experiments *in silico*. DLAG could be used to enhance the study of both the recorded and artificial neuronal populations.

Interpretation of DLAG's latent variables and time delays   One might interpret the population activity patterns represented by DLAG's across-area variables as distinct "channels" with which two areas communicate [43]. As with any statistical method, however, interpretation of the features extracted by DLAG is subject to ambiguities, particularly when not all relevant brain areas and neurons are recorded [60]. An across-area latent variable, for instance, could reflect an interaction between areas A and B that is direct or indirect, mediated by a third (unob-

served) area C. Similarly, a within-area latent variable could reflect activity internal to one area, or it could reflect inputs sent from unrecorded neurons to one area but not the other.

The sign and magnitude of DLAG's time delays can, however, narrow the set of hypotheses consistent with the data. We might reasonably suspect, for example, that short positive (V1 to V2) delays identified by DLAG reflect direct interactions from the output layers of V1 to the input layers of V2 (the layers from which we recorded) [26, 28]. Larger negative (V2 to V1) delays might instead indicate indirect interactions, given that several intermediate connections likely lie between the input layers of V2 and the output layers of V1. Many of the V1-V2 interactions were associated with delays statistically indistinguishable from zero (i.e., "ambiguous"), and could reflect a third case: common input from an unobserved source. Additional experimental interventions could further disambiguate these cases.

A few assumptions explicit in the DLAG model definition warrant additional care when interpreting estimated delays. First, DLAG treats time delays as constant parameters. However, interactions between areas might not be constant across different trial epochs or different experimental (e.g., stimulus) conditions. Estimated delays therefore resemble an average over these variations; and we interpret delays as summaries of the persistent direction of interactions associated with a particular activity pattern, throughout the course of a trial. Second, DLAG assumes that each dimension of population activity is associated with one delay, or direction. If an interaction is bidirectional yet one-dimensional (as in a coupled oscillator), then teasing apart directionality might be difficult—albeit for any statistical method, not just DLAG.

Finally, DLAG ties the delays among neurons in each area so that each neuron is assumed to take part in an inter-areal interaction with the same latency. The PSTHs of V1 and V2, however (Fig. 4b), demonstrate that the stimulus response properties of neurons within a population can vary by tens of ms. DLAG's delay estimates must aggregate these disparate response properties, and hence could vary significantly for different subsamples of each population. We therefore interpret, for example, the stereotyped periodic signals evident in V1-V2 across-area latent variables (Fig. 5a; "Across 3") as likely less a reflection of communication between V1 and V2, and

more a reflection of aggregate differences in the stimulus response properties of neurons in V1 and in V2.

Solutions to these interpretational challenges might already be well within reach, if not already available through DLAG's existing machinery. For example, one could simply fit DLAG to subsets of trials, subsets of neurons, or to separate trial epochs to obtain appropriate statistics from DLAG model parameters or posterior estimates. We have already employed some of these strategies here, and could continue to build upon that foundation.

Interactions across more than two brain areas   In this work, we have presented DLAG as a means to disentangle the flow of signals between two neuronal populations. Of course, most sensory [6–9], cognitive [10], and motor functions [11] rely on distributed networks of neuronal populations, and recordings involving three or more distinct populations are becoming more accessible. To study these many-area recordings, statistical methods will need to disambiguate not only within-area versus across-area interactions, but more generally single-area, two-way, three-way, up to global interactions. In addition, disambiguating the direction of signal flow through a network of several areas is a well-known statistical challenge [60]. Future methods, including extensions to our current work, will need to address these challenges while remaining computationally tractable as larger networks of brain areas are considered. Still, DLAG is an important step toward tracking the intricate, multi-dimensional flow of signals across many interacting populations of neurons.

Methods

Code availability

All methods and data analyses described here were implemented and carried out in Matlab (The Mathworks, Inc.). Implementations of DLAG in Matlab and Python will be available by November 2021.

Mathematical notation

To disambiguate each variable or parameter in the DLAG model, we need to keep track of up to four labels that indicate their associated (1) brain area; (2) neuron or latent variable index; (3) time point; or (4) designation as within- or across-area. We indicate the first three labels via subscripts, where brain areas are indexed by $i = 1, 2$; neurons or latent variables are indexed by j (we'll indicate the upper bound as appropriate); and time is indexed by $t = 1, \ldots, T$. For example, we define the observed activity of neuron j (out of q_i) in area i at time t as $y_{i,j,t} \in \mathbb{R}$. To indicate a collection of all variables along a particular index, we replace that index with the ':' symbol. Hence we represent the simultaneous activity of the population of q_i neurons observed in area i at time t as the vector $\mathbf{y}_{i,:,t} \in \mathbb{R}^{q_i}$. For concision, where a particular index is either not applicable or not immediately relevant, we omit it. The identities of the remaining indices should be clear from context. For example, throughout this work we consider only the activity of a full population, and not of single neurons, so we rewrite $\mathbf{y}_{i,:,t}$ as $\mathbf{y}_{i,t}$. Finally, we indicate a latent variable's or parameter's designation as within- or across-area via a superscript, where 'w' indicates within-area, and 'a' indicates across-area. For example, we define across-area latent variable j (out of p^a) in area i at time t as $x^a_{i,j,t} \in \mathbb{R}$, and the collection of all p^a latent variables as the vector $\mathbf{x}^a_{i,:,t} \in \mathbb{R}^{p^a}$. We similarly define within-area latent variable j (out of p^w_i) in area i at time t as $x^w_{i,j,t} \in \mathbb{R}$, and the collection of all p^w_i latent variables as the vector $\mathbf{x}^w_{i,:,t} \in \mathbb{R}^{p^w_i}$.

It is conceptually helpful to understand the above notation for observed ($\mathbf{y}$) and latent ($\mathbf{x}$) variables as taking cross-sections of matrices. For example, observed activity in area i can be grouped

into the matrix $Y_i = [\mathbf{y}_{i,1} \cdots \mathbf{y}_{i,T}] \in \mathbb{R}^{q_i \times T}$. Then, each $\mathbf{y}_{i,t}$ is a column of Y_i. Similarly, across-area latent variables in area i can be grouped into the matrix $X_i^a = [\mathbf{x}_{i,:,1}^a \cdots \mathbf{x}_{i,:,T}^a] \in \mathbb{R}^{p^a \times T}$. Each $\mathbf{x}_{i,:,t}^a$ is a column of X_i^a. Similarly, we represent a row of X_i^a (i.e., the values of a single latent variable j at all time points) as $\mathbf{x}_{i,j,:}^a \in \mathbb{R}^T$. Within-area latent variables can be understood analogously from the matrix $X_i^w = [\mathbf{x}_{i,:,1}^w \cdots \mathbf{x}_{i,:,T}^w] \in \mathbb{R}^{p_i^w \times T}$.

We will explicitly define all other variables and parameters as they appear, but for reference, we list common variables and parameters below:

Observed neural activity

- q_i – number of neurons observed in area i
- Y_i – $q_i \times T$ matrix of observed activity in area i
- $\mathbf{y}_{i,t}$ – $q_i \times 1$ vector of observed activity in area i at time t; the t^{th} column of Y_i

Latent variables

- p^a – number of across-area variables (same for both areas)
- X_i^a – $p^a \times T$ matrix of across-area variables in area i
- $\mathbf{x}_{i,:,t}^a$ – $p^a \times 1$ vector of across-area variables in area i at time t; the t^{th} column of X_i^a
- $\mathbf{x}_{i,j,:}^a$ – $T \times 1$ vector of values of across-area variable j in area i over time; the j^{th} row of X_i^a
- p_i^w – number of within-area variables in area i
- X_i^w – $p_i^w \times T$ matrix of within-area variables in area i
- $\mathbf{x}_{i,:,t}^w$ – $p_i^w \times 1$ vector of within-area variables in area i at time t; the t^{th} column of X_i^w
- $\mathbf{x}_{i,j,:}^w$ – $T \times 1$ vector of values of within-area variable j in area i over time; the j^{th} row of X_i^w

Model parameters

- C_i^a – $q_i \times p^a$ across-area loading matrix for area i
- C_i^w – $q_i \times p_i^w$ within-area loading matrix for area i
- $\mathbf{d}_i$ – $q_i \times 1$ mean parameter for area i
- R_i – $q_i \times q_i$ observation noise covariance matrix for area i
- $D_{i,j}$ – time delay parameter between area i and across-area variable j
- D_j – relative time delay associated with across-area variable j; $D_j = D_{2,j} - D_{1,j}$
- τ_j^a – Gaussian process timescale for across-area variable j
- σ_j^a – Gaussian process noise parameter for across-area variable j

 • $\tau_{i,j}^w$ – Gaussian process timescale for within-area variable j in area i

• $\sigma_{i,j}^w$ – Gaussian process noise parameter for within-area variable j in area i

Gaussian process covariances

• $K_{i_1,i_2,j}^a$ – $T \times T$ covariance matrix for across-area variable j, between areas i_1 and i_2

• $k_{i_1,i_2,j}^a$ – covariance function for across-area variable j, between areas i_1 and i_2

• $K_{i,j}^w$ – $T \times T$ covariance matrix for within-area variable j in area i

• $k_{i,j}^w$ – covariance function for within-area variable j in area i

DLAG observation model

For area i at time t, we define a linear-Gaussian relationship between observed activity, $\mathbf{y}_{i,t}$, and

latent variables, $\mathbf{x}_{i,:,t}^a$ and $\mathbf{x}_{i,:,t}^w$ [61]:

$$\mathbf{y}_{i,t} = C_i^a \mathbf{x}_{i,:,t}^a + C_i^w \mathbf{x}_{i,:,t}^w + \mathbf{d}_i + \boldsymbol{\varepsilon}_i \tag{1}$$

$$\boldsymbol{\varepsilon}_i \sim \mathcal{N}(\mathbf{0}, R_i) \tag{2}$$

where $C_i^a \in \mathbb{R}^{q_i \times p^a}$, $C_i^w \in \mathbb{R}^{q_i \times p_i^w}$, $\mathbf{d}_i \in \mathbb{R}^{q_i}$, and $R_i \in \mathbb{S}^{q_i \times q_i}$ ($\mathbb{S}^{q_i \times q_i}$ is the set of $q_i \times q_i$ symmetric

matrices) are model parameters to be estimated from data. The relationship between observed

and latent variables is illustrated graphically in Supplementary Fig. 1. The loading parameter

matrices C_i^a and C_i^w linearly combine latent variables and map them to observed neural activity.

The parameter $\mathbf{d}_i$ can be thought of as the mean firing rate of each neuron. $\boldsymbol{\varepsilon}_i$ is a zero-mean

Gaussian random variable, where we constrain the covariance matrix R_i to be diagonal, as in

factor analysis (FA) [62] and Gaussian process factor analysis (GPFA) [56], to capture variance

that is independent to each neuron. This constraint encourages the latent variables to explain as

much of the shared variance among neurons as possible.

As we will describe, at time point t, across-area variables $\mathbf{x}_{1,:,t}^a$ and $\mathbf{x}_{2,:,t}^a$ in area 1 and area 2,

respectively, are coupled with each other, and thus each area has the same number of across-

area variables, p^a. Within-area variables are not coupled across areas, on the other hand, and

thus each area i may have a different number of within-area variables, p_i^w. Because we seek a low-dimensional description of neural activity in each area, the combined number of across- and within-area variables is less than the number of neurons, i.e., $p^a + p_i^w < q_i$, where p^a and p_i^w are determined by the data (see below).

The parameters C_i^w and C_i^a have an intuitive geometric interpretation (Fig. 2, right column). If we represent each element of $\mathbf{y}_{1,t}$ and $\mathbf{y}_{2,t}$, the activity of each neuron in each area, as an axis in a high-dimensional space (one set of axes for each area), then the columns of C_i^w and C_i^a define hyperplanes, or subspaces, in these high-dimensional spaces. These subspaces represent patterns of activity shared among neurons within (C_i^w) and across (C_i^a) areas; each dimension corresponds to a distinct latent variable. Additionally, as we will discuss below, since the j^{th} pair of across-area variables ($\mathbf{x}_{1,j,:}^a$, $\mathbf{x}_{2,j,:}^a$) is associated with a direction of population signal flow (Fig. 2, center column), so too are the corresponding columns in C_1^a and C_2^a. The across-area subspace can thus be partitioned further based on the nominal directionality of activity patterns (area 1 to area 2, or area 2 to area 1). Finally, note that the columns of C_i^a and C_i^w (and the subspaces they define) are linearly independent; but they are not, in general, orthogonal.

DLAG state model

The temporal structure of interactions within and across areas are captured entirely through relationships between latent variables. We describe the time course of within- and across-area variables by relating each $\mathbf{x}_{i,:,t}^w$ and $\mathbf{x}_{i,:,t}^a$ in brain area i at different time points through Gaussian processes [63].

Within-area latent variables We first describe within-area temporal structure. In particular, we consider $\mathbf{x}_{i,j,:}^w \in \mathbb{R}^T$, the j^{th} row of X_i^w. For each within-area variable $j = 1, \ldots, p_i^w$ in brain area i, we define a separate Gaussian process (GP) as follows [56]:

$$\mathbf{x}_{i,j,:}^w \sim \mathcal{N}(\mathbf{0}, K_{i,j}^w) \tag{3}$$

where $K_{i,j}^w \in \mathbb{S}^{T \times T}$ is the covariance matrix for within-area variable j of area i. DLAG is compatible with any valid form of GP covariance, but for the present work, we choose the commonly used squared exponential (SE) function. Then, element (t_1, t_2) of $K_{i,j}^w$, the covariance between samples at times t_1 and t_2, can be computed according to:

$$k_{i,j}^w(t_1, t_2) = \left(1 - (\sigma_{i,j}^w)^2\right) \exp\left(-\frac{(\Delta t)^2}{2(\tau_{i,j}^w)^2}\right) + (\sigma_{i,j}^w)^2 \cdot \delta_{\Delta t} \tag{4}$$

$$\Delta t = t_2 - t_1 \tag{5}$$

where the characteristic timescale, $\tau_{i,j}^w \in \mathbb{R}_{>0}$, and GP noise variance, $(\sigma_{i,j}^w)^2 \in (0, 1)$, are model parameters. $\delta_{\Delta t}$ is the kronecker delta, which is 1 for $\Delta t = 0$ (equivalently, $t_1 = t_2$) and 0 otherwise.

Notice that $k_{i,j}^w$ is stationary: the SE function depends only on the time difference $(t_2 - t_1)$ (Supplementary Fig. 2a). This stationarity gives the covariance matrix $K_{i,j}^w$ a characteristic banded structure (Supplementary Fig. 2b). The characteristic timescale, $\tau_{i,j}^w$, dictates the width of $k_{i,j}^w(t_1, t_2)$, or equivalently, how rapidly the latent variable changes over time. Each $\tau_{i,j}^w$ is estimated from data. While the GP noise variance, $(\sigma_{i,j}^w)^2$, can be estimated in the same manner as $\tau_{i,j}^w$, we follow the same conventions as in [56], and fix $(\sigma_{i,j}^w)^2$ to a small value (10^{-3}). Note also that, under this definition, the process is normalized so that $k_{i,j}^w(t_1, t_2) = 1$ for $t_1 = t_2$. Equivalently, the prior distribution of within-area latent variables $\mathbf{x}_{i,:,t}^w$ in area i at each time t follows the standard normal distribution, $\mathcal{N}(\mathbf{0}, I)$. This normalization removes model redundancy in the scaling of X_i^w and C_i^w.

Beyond describing within-area interactions, within-area variables are critical to the interpretability of across-area variables. As we will define below, across-area variables can describe the activity of neurons in both areas; so in principle, it is perfectly valid for them to describe interactions that are shared only among neurons in one area, but not both. If the model does not allow for within-area variables, then, across-area variables are forced to trade off between explaining activity that is truly shared between both areas and explaining activity that is present

in only one area but statistically salient (i.e., high-variance). By capturing within-area interactions, within-area variables allow the across-area variables to isolate activity that is truly shared between both areas. This statistical phenomenon applies to other statistical models, and is not specific to DLAG [36, 57]. See Supplementary Discussion for further mathematical discussion.

Across-area latent variables We next describe across-area temporal structure. As we'll detail, across-area variables are different from within-area variables in two respects: across-area variables are (1) coupled; and (2) time-delayed relative to each other (Fig. 2, center column). Thus in contrast to our definition of within-area variables, in which we considered each area independently, we now consider across-area variables in both areas together: $\mathbf{x}_{1,j,:}^a \in \mathbb{R}^T$ and $\mathbf{x}_{2,j,:}^a \in \mathbb{R}^T$, the j^{th} rows of X_1^a and X_2^a, respectively, for the j^{th} across-area variable. For each across-area variable $j = 1, \ldots, p^a$, we let $\mathbf{x}_{1,j,:}^a$ and $\mathbf{x}_{2,j,:}^a$ covary according to a common Gaussian process:

$$\begin{bmatrix} \mathbf{x}_{1,j,:}^a \\ \mathbf{x}_{2,j,:}^a \end{bmatrix} \sim \mathcal{N}\left(\mathbf{0}, \begin{bmatrix} K_{1,1,j}^a & K_{1,2,j}^a \\ K_{2,1,j}^a & K_{2,2,j}^a \end{bmatrix} \right) \tag{6}$$

where $K_{1,1,j}^a = K_{2,2,j}^a$ describe the autocovariance of each across-area variable, and $K_{1,2,j}^a = K_{2,1,j}^{a\top}$ describe the cross-covariance that couples the two areas. The cross-covariance therefore captures interactions across areas over time and, as we'll detail below, at a relative time delay.

To express the auto- and cross-covariance functions, we introduce additional notation. Specifically, we indicate brain areas with two subscripts, $i_1 = 1, 2$ and $i_2 = 1, 2$. Then, we define $K_{i_1,i_2,j}^a \in \mathbb{R}^{T \times T}$ to be either the auto- or cross-covariance matrix between across-area variable $\mathbf{x}_{i_1,j,:}^a$ in area i_1 and across-area variable $\mathbf{x}_{i_2,j,:}^a$ in area i_2. We again choose to use the SE function for GP covariances. Therefore, element (t_1, t_2) of each $K_{i_1,i_2,j}^a$ can be computed as follows [59]:

$$k_{i_1,i_2,j}^a(t_1, t_2) = \left(1 - (\sigma_j^a)^2\right) \exp\left(-\frac{(\Delta t)^2}{2(\tau_j^a)^2}\right) + (\sigma_j^a)^2 \cdot \delta_{\Delta t} \tag{7}$$

$$\Delta t = (t_2 - D_{i_2,j}) - (t_1 - D_{i_1,j}) \tag{8}$$

where, like the within-area case, the characteristic timescale, $\tau_j^a \in \mathbb{R}_{>0}$, and the GP noise vari-

ance, $(\sigma_j^a)^2 \in (0, 1)$, are model parameters. $\delta_{\Delta t}$ is the kronecker delta, which is 1 for $\Delta t = 0$ and 0

otherwise.

We also introduce two new parameters: the time delay to area i_1, $D_{i_1,j} \in \mathbb{R}$, and the time de-

lay to area i_2, $D_{i_2,j} \in \mathbb{R}$. Notice that, when computing the autocovariance for area i (i.e., $i_1 =$

$i_2 = i$), the time delay parameters $D_{i_1,j}$ and $D_{i_2,j}$ are equal, and so Δt (equation (8)) reduces

simply to the time difference $(t_2 - t_1)$, as in the within-area case (equation (5)). Time delays

are therefore only relevant when computing the cross-covariance between area 1 and area 2. The

time delay to area 1, $D_{1,j}$, and the time delay to area 2, $D_{2,j}$, by themselves have no physically

meaningful interpretation. Their difference $D_j = D_{2,j} - D_{1,j}$, however, represents a well-defined,

continuous-valued time delay from area 1 to area 2. The sign of the relative time delay D_j indi-

cates the directionality of the lead-lag relationship between areas captured by latent variable j

(positive: area 1 leads area 2; negative: area 2 leads area 1), which we interpret as a description

of inter-areal signal flow.

The shape of the auto-covariance function for area i, $k_{i,i,j}^a(t_1, t_2)$, and cross-covariance function

between area 1 and area 2, $k_{1,2,j}^a(t_1, t_2)$, are illustrated in Supplementary Fig. 2c as a function

of the time difference $(t_2 - t_1)$. The characteristic timescale, τ_j^a, dictates the width of both the

auto- and cross-covariances, or equivalently, how rapidly the latent variables in both areas change

over time. The relative delay D_j shifts the peak of the cross-covariance away from zero; or equiv-

alently, effectively shifts (along the common GP) the samples of the across-area variable in area

2, $\mathbf{x}_{2,j,:}^a$, relative to the samples of the across-area variable in reference area 1, $\mathbf{x}_{1,j,:}^a$. The corre-

sponding structure of the auto- and cross-covariance matrices is illustrated in Supplementary

Fig. 2d. Note again the banded structure that results from the SE function; and in particular,

note how the banded structure of the cross-covariance matrices is shifted away from the diagonal because of the relative delay D_j.

Both the characteristic timescales τ_j^a and relative delays D_j are estimated from data. More specifically, to ensure identifiability of time delay parameters, we designate area 1 as the reference area, and fix the delays for area 1 at 0, that is, $D_{1,j} = 0$ for all across-area variables $j = 1, \ldots, p^a$. Then, each relative time delay D_j is simply the time delay parameter to area 2, $D_{2,j}$, which is fit to data as detailed in the next section. As in the within-area case, the across-area GP noise variance, $(\sigma_j^a)^2$, is set to a small value (10^{-3}). Furthermore, the across-area GP is also normalized so that $k_{i_1,i_2,j}^a(t_1, t_2) = 1$ if $\Delta t = 0$, thereby removing model redundancy in the scaling of X_i^a and C_i^a.

DLAG special cases   Finally, we consider some special cases of the DLAG model that illustrate its relationship to other dimensionality reduction methods. First, by fixing all time delays to zero ($D_j = 0$), and by removing within-area latent variables ($p_1^w = p_2^w = 0$), DLAG becomes equivalent to Gaussian process factor analysis (GPFA) [56] applied to both areas jointly. By removing instead the across-area latent variables ($p^a = 0$), and keeping the within-area latent variables intact, DLAG becomes equivalent to GPFA applied to each area independently. And finally, by removing temporal smoothing (i.e., in the limit as all GP noise parameters σ_j^a, $\sigma_{i,j}^w$ approach 1), while keeping both within- and across-area latent variables, DLAG becomes equivalent to a model closely related to probabilistic canonical correlation analysis (pCCA) [61, 64].

Fitting the DLAG model

In this section, we describe how DLAG model parameters are fit using exact Expectation Maximization (EM). Toward that end, we first write the DLAG observation model more compactly as follows. Define the joint activity of neurons in all brain areas by vertically concatenating the observations in each area, $\mathbf{y}_{1,t}$ and $\mathbf{y}_{2,t}$:

$$\mathbf{y}_t = \begin{bmatrix} \mathbf{y}_{1,t} \\ \mathbf{y}_{2,t} \end{bmatrix} \in \mathbb{R}^q \tag{9}$$

where $q = q_1 + q_2$. Next we group together the across- and within-area latent variables for the i^{th} brain area to define $\mathbf{x}_{i,t} = [\mathbf{x}_{i,:,t}^{a\top} \ \mathbf{x}_{i,:,t}^{w\top}]^\top \in \mathbb{R}^{p_i}$, where $p_i = p^a + p_i^w$. We then vertically concatenate the latent variables in each area:

$$\mathbf{x}_t = \begin{bmatrix} \mathbf{x}_{1,t} \\ \mathbf{x}_{2,t} \end{bmatrix} \in \mathbb{R}^p \tag{10}$$

where $p = p_1 + p_2$. We also define the following structured matrices. First define $C_i = [C_i^a \ C_i^w] \in \mathbb{R}^{q_i \times p_i}$ by horizontally concatenating C_i^a and C_i^w. Then, we collect the C_i into a block-diagonal matrix as follows:

$$C = \begin{bmatrix} C_1 & \mathbf{0} \\ \mathbf{0} & C_2 \end{bmatrix} \in \mathbb{R}^{q \times p} \tag{11}$$

Similarly, define

$$R = \begin{bmatrix} R_1 & \mathbf{0} \\ \mathbf{0} & R_2 \end{bmatrix} \in \mathbb{R}^{q \times q}, \tag{12}$$

$$\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix} \in \mathbb{R}^q \tag{13}$$

We can then write the DLAG observation model compactly as follows:

$$\mathbf{y}_t \mid \mathbf{x}_t \sim \mathcal{N}(C\mathbf{x}_t + \mathbf{d}, R) \tag{14}$$

The observation model expressed in equation (14) defines a distribution for neural activity at a single time point, but to properly fit the DLAG model, we must consider the distribution over all time points. Thus we define $\bar{\mathbf{y}} = [\mathbf{y}_1^\top \cdots \mathbf{y}_T^\top]^\top \in \mathbb{R}^{qT}$ and $\bar{\mathbf{x}} = [\mathbf{x}_1^\top \cdots \mathbf{x}_T^\top]^\top \in \mathbb{R}^{pT}$, obtained by vertically concatenating the observed variables $\mathbf{y}_t$ and latent variables $\mathbf{x}_t$, respectively, across all $t = 1, \ldots, T$. Then, we rewrite the state and observation models as follows:

$$\bar{\mathbf{x}} \sim \mathcal{N}(\mathbf{0}, \bar{K}) \tag{15}$$

$$\bar{\mathbf{y}} \mid \bar{\mathbf{x}} \sim \mathcal{N}(\bar{C}\bar{\mathbf{x}} + \bar{\mathbf{d}}, \bar{R}), \tag{16}$$

where $\bar{C} \in \mathbb{R}^{qT \times pT}$ and $\bar{R} \in \mathbb{S}^{qT \times qT}$ are block diagonal matrices comprising T copies of the matrices C and R, respectively. $\bar{\mathbf{d}} \in \mathbb{R}^{qT}$ is constructed by vertically concatenating T copies of $\mathbf{d}$. The elements of $\bar{K} \in \mathbb{R}^{pT \times pT}$ are computed using equations (3)-(8). Then, the joint distribution over observed and latent variables is given by

$$\begin{bmatrix} \bar{\mathbf{x}} \\ \bar{\mathbf{y}} \end{bmatrix} \sim \mathcal{N}\left(\begin{bmatrix} \mathbf{0} \\ \bar{\mathbf{d}} \end{bmatrix}, \begin{bmatrix} \bar{K} & \bar{K}\bar{C}^\top \\ \bar{C}\bar{K} & \bar{C}\bar{K}\bar{C}^\top + \bar{R} \end{bmatrix} \right) \tag{17}$$

E-step In the E-step, our goal is to compute the posterior distribution of the latent variables $\bar{\mathbf{x}}$ given the recorded neural activity $\bar{\mathbf{y}}$, $P(\bar{\mathbf{x}}|\bar{\mathbf{y}})$, using the most recent parameter estimates

$$\theta = \left\{ C, \ \mathbf{d}, \ R, \ \{D_j\}_{j=1}^{p^a}, \ \{\tau_j^a\}_{j=1}^{p^a}, \ \{\tau_{1,j}^w\}_{j=1}^{p_1^w}, \ \{\tau_{2,j}^w\}_{j=1}^{p_2^w} \right\} \tag{18}$$

Using basic results of conditioning for jointly Gaussian random variables, we get

$$\bar{\mathbf{x}} \mid \bar{\mathbf{y}} \sim \mathcal{N}\left(\bar{K}\bar{C}^\top \left(\bar{C}\bar{K}\bar{C}^\top + \bar{R}\right)^{-1} (\bar{\mathbf{y}} - \bar{\mathbf{d}}), \bar{K} - \bar{K}\bar{C}^\top \left(\bar{C}\bar{K}\bar{C}^\top + \bar{R}\right)^{-1} \bar{C}\bar{K} \right) \tag{19}$$

Thus, posterior estimates of latent variables are given by

$$\mathbb{E}\left[\bar{\mathbf{x}}|\bar{\mathbf{y}}\right] = \bar{K}\bar{C}^\top \left(\bar{C}\bar{K}\bar{C}^\top + \bar{R}\right)^{-1} (\bar{\mathbf{y}} - \bar{\mathbf{d}}) \tag{20}$$

The marginal likelihood can be computed as

$$\bar{\mathbf{y}} \sim \mathcal{N}\left(\bar{\mathbf{d}}, \bar{C}\bar{K}\bar{C}^\top + \bar{R}\right) \tag{21}$$

M-step In the M-step, our goal is to maximize $\mathcal{E}(\theta) = \mathbb{E}\left[\log P(\bar{\mathbf{x}}, \bar{\mathbf{y}})|\theta\right]$ with respect to θ, using the latest inference of the latent variables, computed in the E-step. As in [56, 59], we adopt the following notation. Given a vector $\mathbf{v}$,

$$\langle \mathbf{v} \rangle = \mathbb{E}\left[\mathbf{v}|\bar{\mathbf{y}}\right] \tag{22}$$

$$\langle \mathbf{v}\mathbf{v}^\top \rangle = \mathbb{E}\left[\mathbf{v}\mathbf{v}^\top|\bar{\mathbf{y}}\right] \tag{23}$$

The appropriate expectations can be found using equation (19).

Maximizing $\mathcal{E}(\theta)$ with respect to C, $\mathbf{d}$ yields the following closed-form update for the i^{th} brain area:

$$\begin{bmatrix} C_i & \mathbf{d}_i \end{bmatrix} = \left(\sum_{t=1}^{T} \mathbf{y}_{i,t} \cdot \begin{bmatrix} \langle \mathbf{x}_{i,t} \rangle^\top & 1 \end{bmatrix}\right) \left(\sum_{t=1}^{T} \begin{bmatrix} \langle \mathbf{x}_{i,t}\mathbf{x}_{i,t}^\top \rangle & \langle \mathbf{x}_{i,t} \rangle \\ \langle \mathbf{x}_{i,t} \rangle^\top & 1 \end{bmatrix}\right)^{-1} \tag{24}$$

After performing the update for each area separately, we collect all updated values into C and $\mathbf{d}$. Then we update R for both brain areas together, as follows:

$$R = \frac{1}{T}\text{diag}\left\{\sum_{t=1}^{T}\left((\mathbf{y}_t - \mathbf{d})(\mathbf{y}_t - \mathbf{d})^\top - (\mathbf{y}_t - \mathbf{d})\langle \mathbf{x}_t \rangle^\top C^\top - C\langle \mathbf{x}_t \rangle(\mathbf{y}_t - \mathbf{d})^\top + C\langle \mathbf{x}_t\mathbf{x}_t^\top \rangle C^\top\right)\right\} \tag{25}$$

There are no closed-form solutions for the Gaussian process parameter updates, but we can compute gradients and perform gradient ascent. Note that, for this work, we choose not to fit the Gaussian process noise variances, but rather, we set them to small values (10^{-3}), as in [56]. Within-area timescale gradients for the i^{th} brain area and j^{th} within-area latent variable are given by

$$\frac{\partial \mathcal{E}(\theta)}{\partial \tau_{i,j}^w} = \text{tr}\left(\left(\frac{\partial \mathcal{E}(\theta)}{\partial K_{i,j}^w} \right)^{\top} \left(\frac{\partial K_{i,j}^w}{\partial \tau_{i,j}^w} \right) \right) \tag{26}$$

where

$$\frac{\partial \mathcal{E}(\theta)}{\partial K_{i,j}^w} = -\frac{1}{2}(K_{i,j}^w)^{-1} + \frac{1}{2}\left((K_{i,j}^w)^{-1} \langle \mathbf{x}_{i,j,:}^w \mathbf{x}_{i,j,:}^{w\top} \rangle (K_{i,j}^w)^{-1} \right) \tag{27}$$

and element (t_1, t_2) of $\partial K_{i,j}^w / \partial \tau_{i,j}^w$ is given by

$$\frac{\partial k_{i,j}^w(t_1, t_2)}{\partial \tau_{i,j}^w} = \left(1 - (\sigma_{i,j}^w)^2 \right) \frac{(t_2 - t_1)^2}{(\tau_{i,j}^w)^3} \exp\left(-\frac{(t_2 - t_1)^2}{2(\tau_{i,j}^w)^2} \right) \tag{28}$$

To express the across-area timescale and delay parameter gradients, we introduce more compact notation for the variables in equation (6). Let $\mathbf{x}_{j,:}^a = [\mathbf{x}_{1,j,:}^{a\top} \mathbf{x}_{2,j,:}^{a\top}]^{\top} \in \mathbb{R}^{2T}$ for the j^{th} across-area latent variable, and

$$K_j^a = \begin{bmatrix} K_{1,1,j}^a & K_{1,2,j}^a \\ K_{2,1,j}^a & K_{2,2,j}^a \end{bmatrix} \in \mathbb{S}^{2T \times 2T} \tag{29}$$

Then, across-area timescale gradients are given by

$$\frac{\partial \mathcal{E}(\theta)}{\partial \tau_j^a} = \text{tr}\left(\left(\frac{\partial \mathcal{E}(\theta)}{\partial K_j^a} \right)^{\top} \left(\frac{\partial K_j^a}{\partial \tau_j^a} \right) \right) \tag{30}$$

where

$$\frac{\partial \mathcal{E}(\theta)}{\partial K_j^a} = -\frac{1}{2}(K_j^a)^{-1} + \frac{1}{2}\left((K_j^a)^{-1}\langle \mathbf{x}_{j,:}^a \mathbf{x}_{j,:}^{a\top}\rangle (K_j^a)^{-1}\right) \tag{31}$$

and each element of $\partial K_j^a / \partial \tau_j^a$ is given by

$$\frac{\partial k_{i_1,i_2,j}^a(t_1,t_2)}{\partial \tau_j^a} = \left(1 - (\sigma_j^a)^2\right)\frac{(\Delta t)^2}{(\tau_j^a)^3}\exp\left(-\frac{(\Delta t)^2}{2(\tau_j^a)^2}\right) \tag{32}$$

Δt is defined as in equation (8). To optimize the timescales while respecting non-negativity constraints, we perform a change of variables, and then perform unconstrained gradient ascent with respect to $\log \tau_{i,j}^w$ or $\log \tau_j^a$.

Next, delay gradients for brain area i and across-area latent variable j are given by

$$\frac{\partial \mathcal{E}(\theta)}{\partial D_{i,j}} = \text{tr}\left(\left(\frac{\partial \mathcal{E}(\theta)}{\partial K_j^a}\right)^\top \left(\frac{\partial K_j^a}{\partial D_{i,j}}\right)\right) \tag{33}$$

where $\frac{\partial \mathcal{E}(\theta)}{\partial K_j^a}$ is defined as in equation (31), and each element of $\partial K_j^a / \partial \tau_j^a$ is given by

$$\frac{\partial k_{i_1,i_2,j}^a(t_1,t_2)}{\partial D_{i,j}} = \left(1 - (\sigma_j^a)^2\right)\frac{\Delta t}{(\tau_j^a)^2}\exp\left(-\frac{(\Delta t)^2}{2(\tau_j^a)^2}\right)\frac{\partial(\Delta t)}{\partial D_{i,j}} \tag{34}$$

$$\frac{\partial(\Delta t)}{D_{i,j}} = \begin{cases} 1 & \text{if } i = i_2 \\ -1 & \text{if } i = i_1 \end{cases} \tag{35}$$

where Δt, i_1, and i_2 are defined as in equation (8). In practice, all delay parameters for area 1 are fixed at 0 to ensure identifiability. As with the timescales, one might wish to constrain the delays within some physically realistic range, such as the length of an experimental trial, so that $-D_{\max} \leq D_{i,j} \leq D_{\max}$. Toward that end, we make the change of variables $D_{i,j} = D_{\max}\frac{1-e^{-D_{i,j}^*}}{1+e^{-D_{i,j}^*}}$ and perform unconstrained gradient ascent with respect to $D_{i,j}^*$. Here we chose $D_{\max}$ to be half the length of a trial.

Finally, note that all of these EM updates are derived for a single sequence, or trial. It is straight-forward to extend these equations to N independent sequences (each with a potentially different number of time steps, T) by maximizing $\frac{\partial}{\partial \theta} \left[\sum_{n=1}^{N} \mathcal{E}_n(\theta) \right]$.

Parameter initialization To initialize DLAG observation model parameters to reasonable values prior to fitting with the EM algorithm, we first fit a probabilistic canonical correlation analysis (pCCA) [64] model to the data, with the same number of across-area latent variables as the desired DLAG model. pCCA is defined by the following state and observation models:

$$\mathbf{x}_t^a \sim \mathcal{N}(\mathbf{0}, I) \tag{36}$$

$$\mathbf{y}_{i,t} \mid \mathbf{x}_t^a \sim \mathcal{N}(C_i^a \mathbf{x}_t^a + \mathbf{d}_i, R_i) \tag{37}$$

where $C_i^a \in \mathbb{R}^{q_i \times p^a}$ maps the p^a-dimensional across-area latent variables $\mathbf{x}_t^a \in \mathbb{R}^{p^a}$ to area i; $\mathbf{d}_i \in \mathbb{R}^{q_i}$ is a mean parameter; and $R_i \in \mathbb{S}^{q_i \times q_i}$ is the observation noise covariance matrix. R_i is not constrained to be diagonal. The fitted values for C_i^a and $\mathbf{d}_i$ are used as initial values for their DLAG analogues. We take only the diagonal elements of R_i to initialize its DLAG analogue.

pCCA does not incorporate within-area latent variables. Therefore, we initialized each DLAG within-area loading matrix C_i^w so that its columns spanned a subspace uncorrelated with that spanned by the columns of C_i^a, returned by pCCA. Such a subspace can be computed as follows. Let $\Sigma_i \in \mathbb{S}^{q_i \times q_i}$ be the sample covariance matrix of activity in area i. Then define $M_i = C_i^{a\top} \Sigma_i \in \mathbb{R}^{p^a \times q_i}$. The singular value decomposition of M_i is given by $M_i = U_i S_i V_i^\top$, where $U_i \in \mathbb{R}^{p^a \times p^a}$, $S_i \in \mathbb{R}^{p^a \times q_i}$, and $V_i \in \mathbb{R}^{q_i \times q_i}$. The first p^a columns of V_i span the same across-area subspace spanned by the columns of C_i^a. The remaining $q_i - p^a$ columns form an orthonormal basis for the subspace uncorrelated with this across-area subspace. We initialized C_i^w with the first p_i^w of these uncorrelated basis vectors. Finally, we initialized all delays to zero, and all within- and across-area Gaussian process timescales to the same value, equal to twice the bin width or sample period of the data.

Selecting the number of within- and across-area latent variables

DLAG has three hyperparameters: p^a, the number of across-area latent variables; and p_1^w and p_2^w, the number of within-area latent variables for each area. Model selection therefore poses a significant scaling challenge. Grid search over even a small range of within- and across-area dimensionalities can result in a large number of models that need to be fitted and validated. For example, considering just 10 possibilities for each type of latent variable would result in 1,000 model candidates. Thus, exhaustive search for the optimal DLAG model is impractical.

We therefore developed a streamlined cross-validation procedure that significantly improves scalability. In brief, our model selection procedure occurs in two stages. First, we consider each area separately, and—using factor analysis (FA) [62]—we find the number of latent variables needed to explain the shared variance among neurons within each area. We reasoned that, while there is not a direct correspondence between the optimal number of latent variables in DLAG and FA models (because of temporal smoothing and other differences in model structure), it is unlikely that the total number of within- and across-area latent variables extracted by DLAG will exceed the FA dimensionality for an area (such a case would imply that there exists a neuron in, for example, area A that covaries with one or more neurons in area B, but no other neurons in area A). Hence we believe this approach to be reasonable given the significant computational benefits. We then use the FA dimensionality in each area to reduce the space of DLAG model candidates to a practical size.

In greater detail, we first applied FA to each area independently, and identified the optimal FA dimensionality through K-fold cross-validation (here we chose $K = 4$). The FA model with the highest cross-validated data likelihood was taken as optimal. We then used the optimal FA dimensionalities ($p_i^{FA}, i = 1, 2$) to constrain the space of DLAG model candidates. In particular, we consider only DLAG models that satisfy $p^a + p_i^w = p_i^{FA}$, for $i = 1, 2$; and, $p^a \leq \min(p_1^{FA}, p_2^{FA})$. In words, we consider only DLAG models such that the number of within- and across-area latent variables in each area sum to that area's optimal FA dimensionality. Furthermore, the number of across-area latent variables is limited by the area with the smallest optimal FA dimensionality.

Not only does this streamlined cross-validation approach provide an upper limit on the possible number of within- and across-area latent variables, it also effectively collapses the DLAG hyperparameter space from three free hyperparameters to one (across-area dimensionality, p^a), drastically improving scalability. Among the model candidates within this constrained search range, we selected models that exhibited the largest cross-validated data likelihood (computed as in equation (21)), using the same K-fold cross-validation scheme as for FA. To further reduce runtime, we limited the number of EM iterations for cross-validated models to 1,000. The optimal model was then re-fit to full convergence, where the data log-likelihood improved from one iteration to the next by less than a preset tolerance (here we used 10^{-8}).

We also note that throughout this work, we explicitly considered model candidates for which across-area dimensionality was zero ($p^a = 0$): the two areas are independent, and any correlations between neurons are purely within-area. Similarly, we explicitly considered model candidates for which within-area dimensionalities were zero ($p_1^w = 0$ or $p_2^w = 0$): all variance shared among neurons in one area can be attributed to their interactions with neurons in the other area. The case where all dimensionalities are zero ($p^a = p_1^w = p_2^w = 0$) is equivalent to fitting a multivariate Gaussian distribution to the data with diagonal covariance (i.e., all neurons are treated as independent). We similarly considered zero-dimensional FA models ($p_1^{FA} = 0$ or $p_2^{FA} = 0$) during the first stage of our model selection procedure, equivalent to fitting a multivariate Gaussian distribution with diagonal covariance to observations in the respective area. The inclusion of these zero-dimensionality model candidates protects against the identification of spurious interactions across or within areas.

Synthetic data generation

We generated synthetic datasets according to the DLAG generative model, so that we could leverage known ground truth to evaluate the accuracy of estimates and characterize DLAG's performance over a range of simulated conditions. We started by randomly generating the set of model parameters, θ, subject to constraints informed by experimental data. For all datasets, we chose realistic values for the numbers of neurons in each area based on our V1-V2 recordings (area A:

$q_1 = 80$; area B: $q_2 = 20$). We set the combined total dimensionality in each area to representative values (area A: $p^a + p_1^w = 10$; area B: $p^a + p_2^w = 5$), but varied the relative number of within- and across-area latent variables across datasets. Generating 20 datasets at each of six configurations ($p^a = 0, \ldots, 5$; $p_1^w = 5, \ldots, 10$; $p_2^w = 0, \ldots, 5$) resulted in a total of 120 independent datasets.

For each dataset, we drew all timescales ($\{\tau_j^a\}_{j=1}^{p^a}$, $\{\tau_{1,j}^w\}_{j=1}^{p_1^w}$, $\{\tau_{2,j}^w\}_{j=1}^{p_2^w}$) uniformly from $U(\tau_{\min}, \tau_{\max})$, with $\tau_{\min} = 10$ ms and $\tau_{\max} = 150$ ms. We drew all delays ($\{D_1, \ldots, D_{p^a}\}$) uniformly from $U(D_{\min}, D_{\max})$, with $D_{\min} = -30$ ms and $D_{\max} = +30$ ms. All Gaussian process noise variances ($\{(\sigma_j^a)^2\}_{j=1}^{p^a}$, $\{(\sigma_{1,j}^w)^2\}_{j=1}^{p_1^w}$, $\{(\sigma_{2,j}^w)^2\}_{j=1}^{p_2^w}$) were fixed at 10^{-3}. To ensure that synthetic datasets exhibited realistic noise levels, we first evaluated the strength of latent variables relative to the strength of single-neuron variability exhibited in the V1-V2 recordings. Specifically, we computed the "signal-to-noise" ratio (where "signal" is defined as the shared activity described by latent variables), $\mathrm{tr}(C_i C_i^\top)/\mathrm{tr}(R_i)$, for V1 and V2 using the parameters of the optimal DLAG models fit to each V1-V2 dataset. Representative values were 0.3 and 0.2 for V1 and V2, respectively. We then generated our synthetic observation model parameters, C_i and R_i, to satisfy these representative ratios. Finally, we drew the entries of $\mathbf{d}$ from the standard normal distribution $\mathcal{N}(\mathbf{0}, I)$. With all model parameters specified, we then generated $N = 100$ independent and identically distributed trials ($\bar{\mathbf{x}}_n, \bar{\mathbf{y}}_n, n = 1, \ldots, N$) according to equations (15) and (16). Each trial comprised $T = 50$ time points, corresponding to $1,000$ ms sequences sampled with a period of 20 ms, to mimic the 20 ms spike count time bins used to analyze the experimental data.

Synthetic data performance metrics

To quantify DLAG's performance across all synthetic datasets, we employed a variety of metrics. To assess the accuracy of loading matrix estimation (C_i^a, C_i^w; reported in Fig. 3, Supplementary Fig. 6, and Supplementary Fig. 7), we first computed a normalized subspace error [65]:

$$e_{\mathrm{sub}} = \frac{\|(I - \hat{M}(\hat{M}^\top \hat{M})^{-1} \hat{M}^\top)M\|_F}{\|M\|_F} \tag{38}$$

 where M is the appropriate ground truth parameter, $\hat{M}$ is the corresponding estimate, and $\|\cdot\|_F$

is the Frobenius norm. e_{sub} quantifies the magnitude of the projection of the column space of M

onto the null space of $\hat{M}$. A value of 1 indicates that the column space of M lies completely in

the null space of $\hat{M}$, and therefore the estimate captures no component of the ground truth. A

value of 0 indicates that the column space of $\hat{M}$ contains the full column space of M, and there-

fore the estimate captures all components of the ground truth. This metric offers two advantages:

(1) it does not require that the columns of M and $\hat{M}$ are ordered in anyway (the ordering of

DLAG latent variables is arbitrary); and (2) it does not require that M and $\hat{M}$ have the same

number of columns, so it can be used to compare the performance of models with different num-

bers of latent variables. Finally, we report the accuracy of loading matrix estimation as $1 - e_{\text{sub}}$.

As for the loading matrices, quantifying the accuracy of the moment-by-moment estimation of

latent variables requires extra care since the sign and ordering of latent variables is arbitrary

and will not, in general, match between estimates and the ground truth. To align estimates with

the ground truth, we instead computed the error between denoised, or smoothed, observations.

First, let $\widetilde{\mathbf{x}}_i^a = [\mathbf{x}_{i,:,1}^{a\top} \cdots \mathbf{x}_{i,:,T}^{a\top}]^\top \in \mathbb{R}^{p^a T}$ be a collection of all (ground truth) across-area variables

at all time points in area i. Similarly, let $\widetilde{\mathbf{x}}_i^w = [\mathbf{x}_{i,:,1}^{w\top} \cdots \mathbf{x}_{i,:,T}^{w\top}]^\top \in \mathbb{R}^{p_i^w T}$ be a collection of all

(ground truth) within-area variables at all time points in area i. Finally, define $\widetilde{C}_i^a \in \mathbb{R}^{q_i T \times p^a T}$

and $\widetilde{C}_i^w \in \mathbb{R}^{q_i T \times p_i^w T}$ to be block diagonal matrices comprising T copies of the (ground truth) ma-

trices C_i^a and C_i^w, respectively; and define $\widetilde{\mathbf{d}}_i \in \mathbb{R}^{q_i T}$ by vertically concatenating T copies of (the

ground truth) $\mathbf{d}_i$. We'll denote the estimates of each of these values by $\widehat{\widetilde{\mathbf{x}}}_i^a$, $\widehat{\widetilde{\mathbf{x}}}_i^w$, $\widehat{\widetilde{C}}_i^a$, $\widehat{\widetilde{C}}_i^w$, and $\widehat{\widetilde{\mathbf{d}}}_i$.

The estimates $\widehat{\widetilde{\mathbf{x}}}_i^a$ and $\widehat{\widetilde{\mathbf{x}}}_i^w$ are posterior means, computed according to equation (20).

Then, to separate the accuracy of across-area variable estimation from the accuracy of within-

area variable estimation (as reported in Fig. 3, Supplementary Fig. 6, and Supplementary Fig. 7),

we estimated denoised (smoothed) observations, using only across-area or only within-area latent

variable estimates:

$$\widehat{\widetilde{\mathbf{y}}}_i^* = \widehat{\widetilde{C}}_i^* \widehat{\widetilde{\mathbf{x}}}_i^* + \widehat{\widetilde{\mathbf{d}}}_i \tag{39}$$

where $\hat{\tilde{\mathbf{y}}}_i^* = [\hat{\mathbf{y}}_{i,1}^{*\top} \cdots \hat{\mathbf{y}}_{i,T}^{*\top}]^\top \in \mathbb{R}^{q_i T}$. Here, the '$*$' symbol is used to indicate either a or w as a superscript, where observations have been denoised using only across- or within-area variable estimates, respectively. We then collect the denoised sequences on all N trials, $\hat{\tilde{\mathbf{y}}}_{i,n}^*$, $n = 1, \ldots, N$, into the matrix $\hat{Y}_i^* = [\hat{\tilde{\mathbf{y}}}_{i,1}^* \cdots \hat{\tilde{\mathbf{y}}}_{i,N}^*] \in \mathbb{R}^{q_i T \times N}$.

Analogously, define $Y_i^* \in \mathbb{R}^{q_i T \times N}$ to be the set of ground truth sequences generated prior to adding noise (i.e., the noise term $\boldsymbol{\varepsilon}_i$, defined in equation (2)). We then compute the R^2 value between estimated and (noiseless) ground truth sequences:

$$R^2 = 1 - \frac{\|Y_i^* - \hat{Y}_i^*\|_F^2}{\|Y_i^* - \bar{Y}_i^*\|_F^2} \tag{40}$$

where $\bar{Y}_i^* = [\bar{\mathbf{y}}_i^* \cdots \bar{\mathbf{y}}_i^*] \in \mathbb{R}^{q_i T \times N}$ is constructed by horizontally concatenating N copies of the sample mean for each neuron in the ground truth Y_i^*, taken over all time points and trials ($\bar{\mathbf{y}}_i^* \in \mathbb{R}^{q_i}$). Note that, in the multivariate case, $R^2 \in (\infty, 1]$, where a negative value implies that estimates predict the ground truth less accurately than simply the sample mean.

Reporting the accuracy of delay and timescale estimates required explicitly matching estimated latent variables to the ground truth. Given the large number of synthetic datasets presented here, we automated this matching process as follows. First, for each area i, we took the unordered across- and within-area latent variable estimates, $\hat{\mathbf{x}}_i^a$ and $\hat{\mathbf{x}}_i^w$, and computed the pairwise correlation between each estimated latent variable and each ground truth latent variable, $\mathbf{x}_i^a$ and $\mathbf{x}_i^w$, across all time points and trials. We then reordered the estimated latent variables to match the ground truth latent variables with which they showed the highest magnitude of correlation. Because latent variables generated from the DLAG model are independent of one another *a priori*, matches based on correlation were straightforward to identify, in practice. To report delay and timescale estimation performance, we computed the absolute error between ground truth and (matched) estimated parameters, to express the error in units of time (ms).

Finally, to assess the accuracy of estimating $\mathbf{d}$ and R (reported in Supplementary Fig. 6 and Supplementary Fig. 7), we first computed the normalized error

$$e_{\text{vec}} = \frac{\|\mathbf{v} - \hat{\mathbf{v}}\|_2}{\|\mathbf{v}\|_2} \tag{41}$$

729 where $\mathbf{v}$ is either $\mathbf{d}$ or $\text{diag}(R)$, and $\hat{\mathbf{v}}$ is the corresponding estimate. We then report the accuracy

730 of estimates as $1 - e_{\text{vec}}$.

Visual stimuli and neural recordings

Animal procedures and recording details have been described in previous work [28, 66]. Briefly,

animals (macaca fascicularis, young adult males) were anesthetized with ketamine (10 mg/kg)

and maintained on isoflurane (1%-2%) during surgery. Recordings were performed under sufen-

tanil (typically 6-18 mg/kg/hr) anesthesia. Vecuronium bromide (150 mg/kg/hr) was used to

prevent eye movements. The duration of each experiment (which comprised multiple recording

sessions) varied from 5 to 7 days. All procedures were approved by the IACUC of the Albert Ein-

stein College of Medicine.

The data analyzed here are a subset of those reported in [28], and similar but not identical to

the subset used in [39, 43]. Activity in V1 output layers was recorded using a 96 channel Utah

array (400 micron inter-electrode spacing, 1 mm length, inserted to a nominal depth of 600 mi-

crons; Blackrock, UT). We recorded V2 activity using a set of electrodes/tetrodes (interelectrode

spacing 300 microns) whose depth could be controlled independently (Thomas Recording, Ger-

many). These electrodes were lowered through V1, the underlying white matter, and then into

V2. Within V2, we targeted neurons in the input layers. We verified the recordings were per-

formed in the input layers using measurements of the depth in V2 cortex, histological confirma-

tion (in a subset of recordings), and correlation measurements. For complete details see [66] and

[28]. Voltage snippets that exceeded a user-defined threshold were digitized and sorted offline.

The sampled neurons had spatial receptive fields within 2-4°of the fovea, in the lower visual field.

We measured responses evoked by drifting sinusoidal gratings (1 cyc/°; drift rate of 3-6.25 Hz;

2.6-4.9°in diameter; full contrast, defined as Michelson contrast, $(L_{\max} - L_{\min})/(L_{\max} + L_{\min})$,

where $L_{\min}$ is 0 cd/m^2 and $L_{\max}$ is 80 cd/m^2) at 8 different orientations (22.5°steps), on a cal-

ibrated CRT monitor placed 110 cm from the animal (1024 x 768 pixel resolution at a 100 Hz refresh rate; Expo: http://sites.google.com/a/nyu.edu/expo). Each stimulus was presented 400 times for 1.28 s. Each presentation was preceded by an interstimulus interval of 1.5 s during which a gray screen was presented.

We recorded neuronal activity in three animals. In two of the animals, we recorded in two different but nearby locations in V2, providing distinct middle-layer populations, yielding a total of five recording sessions. We treated responses to each of the 8 stimuli in each session separately, yielding a total of 40 datasets.

Data preprocessing

We counted spikes in 20 ms time bins during the 1.28 s stimulus presentation period (64 bins per trial). For all analyses corresponding to each recording session, we excluded neurons that fired fewer than 0.5 spikes/s, on average, across all trials and all stimulus orientations. We subtracted the mean response within each trial from each neuron to remove slow fluctuations in mean firing rate across trials. We then applied DLAG to each dataset separately.

Intra-areal and subsampled population comparisons

To contrast against the DLAG models dissecting V1-V2 interactions, we also applied DLAG to V1 only. For each dataset, we randomly split V1 into two equally sized subpopulations (for datasets with an odd number of V1 neurons, we discarded one neuron at random). Each subpopulation was labeled arbitrarily as either "V1a" or "V1b" (Fig. 4c). We then applied DLAG to dissect these "V1a-V1b" interactions in a manner identical to V1-V2 (Fig. 5, Fig. 6).

We also sought to understand the extent to which the spatiotemporal characteristics reported for V1-V2 analyses were driven by disparities in population size between V1 and V2. For each dataset, we therefore randomly subsampled the V1 population to match the size of the V2 population. We then applied DLAG to each subsampled dataset in the same manner as above.

Variance explained by DLAG latent variables

After fitting a DLAG model to each experimental dataset, we sought to compare the relative strengths of across- or within-area latent variables extracted from the same dataset (as in Fig. 5) and across different datasets (as in Fig. 6b,d). To quantify these comparisons, we computed the variance each latent variable explained, as derived from fitted model parameters. From equation (21), it is straightforward to show that the total variance in area i simplifies to

$$\text{var}_{\text{total}} = \text{tr}\left(C_i^a C_i^{a\top} + C_i^w C_i^{w\top} + R_i\right) \tag{42}$$

By inspection, the total variance decomposes into three separable components: $\text{tr}(C_i^a C_i^{a\top})$, the variance due to across-area activity; $\text{tr}(C_i^w C_i^{w\top})$, the variance due to within-area activity; and $\text{tr}(R_i)$, the variance that is independent to each neuron. In fact, the across-area and within-area components can be decomposed further into contributions by individual latent variables. Let $\mathbf{c}_{i,j}^a \in \mathbb{R}^{q_i}$ be the j^{th} column of C_i^a; and $\mathbf{c}_{i,j}^w \in \mathbb{R}^{q_i}$ be the j^{th} column of C_i^w. Then, $\text{tr}(C_i^a C_i^{a\top}) = \sum_{j=1}^{p^a} \|\mathbf{c}_{i,j}^a\|_2^2$, and $\text{tr}(C_i^w C_i^{w\top}) = \sum_{j=1}^{p_i^w} \|\mathbf{c}_{i,j}^w\|_2^2$.

Because we were interested in variance shared among neurons, rather than independent to each neuron, we focused on the variance components involving C_i^a and C_i^w, rather than R_i. Furthermore, since the total variance of recorded neural activity may vary widely across animals, stimuli, and recording sessions, we computed two normalized metrics to facilitate comparison of these shared variance components across datasets. First, let $\mathbf{c}_{i,j}$ be the j^{th} column of C_i, where $C_i = [C_i^a \ C_i^w]$ is the same as in equation (11). To visualize the relative strength of latent variables in each area (Fig. 5), we computed

$$\alpha_{i,j} = \frac{\|\mathbf{c}_{i,j}\|_2^2}{\text{tr}\left(C_i^a C_i^{a\top} + C_i^w C_i^{w\top}\right)} \tag{43}$$

that is, the fraction of shared variance explained by latent variable j in area i. We then displayed latent time courses multiplied by the appropriate $\alpha_{i,j}$ at each time point. Similarly, to quantify

the strength of across-area activity (relative to within-area activity) in each area (Fig. 6b,d), we computed

$$\alpha_i^a = \frac{\text{tr}\left(C_i^a C_i^{a\top}\right)}{\text{tr}\left(C_i^a C_i^{a\top} + C_i^w C_i^{w\top}\right)} \tag{44}$$

that is, the fraction of shared variance explained by all across-area latent variables in area i.

Uncertainty of estimated delays

DLAG's performance on the synthetic data presented here suggests that time delays are estimated with high accuracy and precision. For our neural recordings, however, where no "ground truth" is accessible, we sought to assess the certainty with which fitted delay parameters were indeed positive or negative—indicating a particular direction of inter-areal signal flow. We therefore developed the following nonparametric bootstrap procedure.

First, assume we have a DLAG model that has been fit to a particular dataset with N trials. We construct a bootstrap sample $b = 1, \ldots, B$ from this dataset by selecting N trials uniformly at random with replacement (here we used $B = 1,000$). Then, let ℓ_b be the data log-likelihood of the DLAG model, evaluated on bootstrap sample b. And let $\ell_{b,j=0}$ be the data log-likelihood— also evaluated on bootstrap sample b—of a model for which D_j, the delay for across-area latent variable j, has been set to zero (all other model parameters remain unaltered).

To compare the performance of this "zero-delay" model to the performance of the original model, we define the following statistic:

$$\Delta \ell_{b,j=0} = \ell_b - \ell_{b,j=0} \tag{45}$$

If the zero-delay model performed at least as well as the original DLAG model (equivalently, $\Delta \ell_{b,j=0} \leq 0$) on 5% or more of the bootstrap samples, then we could not say, with sufficient certainty, that the delay for across-area variable j was strictly positive or strictly negative. Oth-

erwise, we took the magnitude of the delay for across-area variable j to differ significantly from zero.

For each of our V1-V2 datasets, then, this procedure allowed us to label some delays as "ambiguous," where the corresponding population signal could not be confidently categorized as flowing in one direction or the other. Finally, note that the concept of ambiguity defined here is distinct from the concept of a variable's importance in describing observed neural activity: for example, an ambiguous across-area variable could, in principle, still explain a large portion of an area's shared variance.

Acknowledgements   This work was supported by the Dowd Fellowship (E.G.), Simons Collaboration on the Global Brain 542999 (A.K.), 543009 (C.K.M.), 543065 (B.M.Y.), 364994 (A.K., B.M.Y.), NIH R01 EY028626 (A.K.), Irma T. Hirschl Trust (A.K.), NIH U01 NS094288 (C.K.M.), NIH R01 HD071686 (B.M.Y.), NIH CRCNS R01 NS105318 (B.M.Y.), NSF NCS BCS 1533672 and 1734916 (B.M.Y.), NIH CRCNS R01 MH118929 (B.M.Y.), and NIH R01 EB026953 (B.M.Y.).

References

1. Ahrens, M. B. *et al.* Brain-wide neuronal dynamics during motor adaptation in zebrafish. *Nature* **485,** 471–477 (2012).

2. Yang, W. & Yuste, R. In vivo imaging of neural activity. *Nature Methods* **14,** 349–359 (2017).

3. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551,** 232–236 (2017).

4. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and engagement across the mouse brain. *Nature* **576,** 266–273 (2019).

5. Kohn, A. *et al.* Principles of Corticocortical Communication: Proposed Schemes and Design Considerations. *Trends in Neurosciences* **43,** 725–737 (2020).

6. Lamme, V. A., Supèr, H. & Spekreijse, H. Feedforward, horizontal, and feedback processing in the visual cortex. *Current Opinion in Neurobiology* **8,** 529–535 (1998).

7. Angelucci, A. & Bressloff, P. C. Contribution of feedforward, lateral and feedback connections to the classical receptive field center and extra-classical receptive field surround of primate V1 neurons. *Progress in Brain Research* **154,** 93–120 (2006).

8. Gilbert, C. D. & Li, W. Top-down influences on visual processing. *Nature Reviews Neuroscience* **14,** 350–363 (2013).

9. Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503,** 51–58 (2013).

10. Miller, E. K., Lundqvist, M. & Bastos, A. M. Working Memory 2.0. *Neuron* **100,** 463–475 (2018).

11. Shadmehr, R. & Krakauer, J. W. A computational neuroanatomy for motor control. *Experimental Brain Research* **185,** 359–381 (2008).

12. Keemink, S. W. & Machens, C. K. Decoding and encoding (de)mixed population responses. *Current Opinion in Neurobiology* **58,** 112–121 (2019).

13. Schmolesky, M. T. *et al.* Signal timing across the macaque visual system. *Journal of Neurophysiology* **79,** 3272–3278 (1998).

14. de Lafuente, V. & Romo, R. Neural correlate of subjective sensory experience gradually builds up across cortical areas. *Proceedings of the National Academy of Sciences of the United States of America* **103,** 14266–14271 (2006).

15. Hernández, A. *et al.* Decoding a perceptual decision process across cortex. *Neuron* **66,** 300–314 (2010).

16. Siegel, M., Buschman, T. J. & Miller, E. K. Cortical information flow during flexible sensorimotor decisions. *Science* **348,** 1352–1355 (2015).

17. Supèr, H., Spekreijse, H. & Lamme, V. A. F. Two distinct modes of sensory processing observed in monkey primary visual cortex (V1). *Nature Neuroscience* **4,** 304–310 (2001).

18. Freeman, J., Ziemba, C. M., Heeger, D. J., Simoncelli, E. P. & Movshon, J. A. A functional and perceptual signature of the second visual area in primates. *Nature Neuroscience* **16,** 974–981 (2013).

19. Pooresmaeili, A., Poort, J. & Roelfsema, P. R. Simultaneous selection by object-based attention in visual and frontal cortex. *Proceedings of the National Academy of Sciences* **111,** 6467–6472 (2014).

20. Chen, M. *et al.* Incremental integration of global contours through interplay between visual cortical areas. *Neuron* **82,** 682–694 (2014).

21. Schwiedrzik, C. M. & Freiwald, W. A. High-Level Prediction Signals in a Low-Level Area of the Macaque Face-Processing Hierarchy. *Neuron* **96,** 89–97 (2017).

22. Issa, E. B., Cadieu, C. F. & DiCarlo, J. J. Neural dynamics at successive stages of the ventral visual stream are consistent with hierarchical error signals. *eLife* **7,** e42870 (2018).

23. Reid, R. C. & Alonso, J. M. Specificity of monosynaptic connections from thalamus to visual cortex. *Nature* **378,** 281–284 (1995).

24. Roe, A. W. & Ts'o, D. Y. Specificity of Color Connectivity Between Primate V1 and V2. *Journal of Neurophysiology* **82,** 2719–2730 (1999).

25. Nowak, L. G., Munk, M., James, A. C., Girard, P. & Bullier, J. Cross-Correlation Study of the Temporal Interactions Between Areas V1 and V2 of the Macaque Monkey. *Journal of Neurophysiology* **81,** 1057–1074 (1999).

26. Jia, X., Tanabe, S. & Kohn, A. Gamma and the Coordination of Spiking Activity in Early Visual Cortex. *Neuron* **77,** 762–774 (2013).

27. Oemisch, M., Westendorff, S., Everling, S. & Womelsdorf, T. Interareal Spike-Train Correlations of Anterior Cingulate and Dorsal Prefrontal Cortex during Attention Shifts. *Journal of Neuroscience* **35,** 13076–13089 (2015).

28. Zandvakili, A. & Kohn, A. Coordinated Neuronal Activity Enhances Corticocortical Communication. *Neuron* **87,** 827–839 (2015).

29. Campo, A. T. *et al.* Task-driven intra- and interarea communications in primate cerebral cortex. *Proceedings of the National Academy of Sciences* **112,** 4761–4766 (2015).

30. Campo, A. T. *et al.* Feed-forward information and zero-lag synchronization in the sensory thalamocortical circuit are modulated during stimulus perception. *Proceedings of the National Academy of Sciences* **116,** 7513–7522 (2019).

31. Gregoriou, G. G., Gotts, S. J., Zhou, H. & Desimone, R. High-frequency, long-range coupling between prefrontal and visual cortex during attention. *Science* **324,** 1207–1210 (2009).

32. Salazar, R. F., Dotson, N. M., Bressler, S. L. & Gray, C. M. Content-Specific Fronto-Parietal Synchronization During Visual Working Memory. *Science* **338,** 1097–1100 (2012).

33. van Kerkoerle, T. *et al.* Alpha and gamma oscillations characterize feedback and feedforward processing in monkey visual cortex. *Proceedings of the National Academy of Sciences* **111,** 14332–14341 (2014).

34. Bastos, A. M., Vezoli, J. & Fries, P. Communication through coherence with inter-areal delays. *Current Opinion in Neurobiology* **31,** 173–180 (2015).

35. Keeley, S. L., Zoltowski, D. M., Aoi, M. C. & Pillow, J. W. Modeling statistical dependencies in multi-region spike train data. *Current Opinion in Neurobiology* **65,** 194–202 (2020).

36. Semedo, J. D., Gokcen, E., Machens, C. K., Kohn, A. & Yu, B. M. Statistical methods for dissecting interactions between brain areas. *Current Opinion in Neurobiology* **65,** 59–69 (2020).

37. Kaufman, M. T., Churchland, M. M., Ryu, S. I. & Shenoy, K. V. Cortical activity in the null space: permitting preparation without movement. *Nature Neuroscience* **17,** 440–448 (2014).

38. Ames, K. C. & Churchland, M. M. Motor cortex signals for each arm are mixed across hemispheres and neurons yet partitioned within the population response. *eLife* **8,** e46159 (2019).

39. Semedo, J. D., Zandvakili, A., Machens, C. K., Yu, B. M. & Kohn, A. Cortical Areas Interact through a Communication Subspace. *Neuron* **102,** 249–259 (2019).

40. Perich, M. G., Gallego, J. A. & Miller, L. E. A Neural Population Mechanism for Rapid Learning. *Neuron* **100,** 964–976 (2018).

41. Ruff, D. A. & Cohen, M. R. Simultaneous multi-area recordings suggest that attention improves performance by reshaping stimulus representations. *Nature Neuroscience* **22,** 1669–1676 (2019).

42. Veuthey, T. L., Derosier, K., Kondapavulur, S. & Ganguly, K. Single-trial cross-area neural population dynamics during long-term skill learning. *Nature Communications* **11,** 4057 (2020).

43. Semedo, J. D. *et al.* Feedforward and feedback interactions between visual cortical areas use different population activity patterns. Preprint at https://doi.org/10.1101/2021.02.08.430346 (2021).

44. Felleman, D. & Essen, D. V. v. Distributed hierarchical processing in the primate cerebral cortex. *Cerebral cortex* (1991).

45. Markov, N. T. *et al.* Cortical High-Density Counterstream Architectures. *Science* **342** (2013).

46. Murray, J. D. *et al.* A hierarchy of intrinsic timescales across primate cortex. *Nature Neuroscience* **17,** 1661–1663 (2014).

47. Sherman, S. M. & Guillery, R. W. On the actions that one nerve cell can have on another: Distinguishing "drivers" from "modulators". *Proceedings of the National Academy of Sciences* **95,** 7121–7126 (1998).

48. Kamiński, M., Ding, M., Truccolo, W. A. & Bressler, S. L. Evaluating causal relations in neural systems: Granger causality, directed transfer function and statistical assessment of significance. *Biological Cybernetics* **85,** 145–157 (2001).

49. Quinn, C. J., Coleman, T. P., Kiyavash, N. & Hatsopoulos, N. G. Estimating the directed information to infer causal relationships in ensemble neural spike train recordings. *Journal of Computational Neuroscience* **30,** 17–44 (2011).

50. Kim, S., Putrino, D., Ghosh, S. & Brown, E. N. A Granger Causality Measure for Point Process Models of Ensemble Neural Spiking Activity. *PLOS Computational Biology* **7,** e1001110 (2011).

51. Pillow, J. W. *et al.* Spatio-temporal correlations and visual signalling in a complete neuronal population. *Nature* **454,** 995–999 (2008).

52. Truccolo, W., Hochberg, L. R. & Donoghue, J. P. Collective dynamics in human and monkey sensorimotor cortex: predicting single neuron spikes. *Nature Neuroscience* **13,** 105–111 (2010).

53. Perich, M. G. *et al.* Inferring brain-wide interactions using data-constrained recurrent neural network models. Preprint at https://doi.org/10.1101/2020.12.18.423348 (2021).

54. Rodu, J., Klein, N., Brincat, S. L., Miller, E. K. & Kass, R. E. Detecting multivariate cross-correlation between brain regions. *Journal of Neurophysiology* **120,** 1962–1972 (2018).

55. Bong, H. *et al.* Latent Dynamic Factor Analysis of High-Dimensional Neural Recordings. *Advances in Neural Information Processing Systems* **33,** 16446–16456 (2020).

56. Yu, B. M. *et al.* Gaussian-Process Factor Analysis for Low-Dimensional Single-Trial Analysis of Neural Population Activity. *Journal of Neurophysiology* **102,** 614–635 (2009).

57. Semedo, J., Zandvakili, A., Kohn, A., Machens, C. K. & Yu, B. M. Extracting Latent Structure From Multiple Interacting Neural Populations. *Advances in Neural Information Processing Systems* **27,** 2942–2950 (2014).

58. Glaser, J., Whiteway, M., Cunningham, J. P., Paninski, L. & Linderman, S. Recurrent Switching Dynamical Systems Models for Multiple Interacting Neural Populations. *Advances in Neural Information Processing Systems* **33,** 14867–14878 (2020).

59. Lakshmanan, K. C., Sadtler, P. T., Tyler-Kabara, E. C., Batista, A. P. & Yu, B. M. Extracting Low-Dimensional Latent Structure from Time Series in the Presence of Delays. *Neural Computation* **27,** 1825–1856 (2015).

60. Reid, A. T. *et al.* Advancing functional connectivity research from association to causation. *Nature Neuroscience* **22,** 1751–1760 (2019).

61. Archambeau, C. & Bach, F. Sparse probabilistic projections. *Advances in Neural Information Processing Systems* **21** (2008).

62. Everett, B. *An Introduction to Latent Variable Models* (Springer Netherlands, 1984).

63. Rasmussen, C. E. & Williams, C. K. I. *Gaussian processes for machine learning* (MIT Press, Cambridge, MA, 2006).

64. Bach, F. R. & Jordan, M. I. A Probabilistic Interpretation of Canonical Correlation Analysis (2005).

65. Golub, G. H. & Van Loan, C. F. *Matrix computations* Fourth edition (The Johns Hopkins University Press, Baltimore, 2013).

66. Smith, M. A. & Kohn, A. Spatial and Temporal Scales of Neuronal Correlation in Primary Visual Cortex. *Journal of Neuroscience* **28,** 12591–12603 (2008).

Figure 1. Disentangling the flow of signals between populations of neurons. Top: Recorded neural activity provides only a tangled view of the bidirectional, concurrent interactions between brain areas (illustrated by the thick translucent arrows; magenta: interactions directed from area A to area B; orange: interactions directed from area B to area A). Bottom left: Existing dimensionality reduction methods identify undirected population activity patterns shared between areas (each activity pattern is represented by a cluster of multi-colored arrows; four different activity patterns are shown). These undirected population activity patterns likely reflect a mixture of the (multi-dimensional) interactions in both directions. Within each activity pattern, individual arrows represent a directed interaction; color depicts the direction of signal flow (magenta: A to B; orange: B to A), and shading (light vs dark) distinguishes distinct interactions. Bottom right: We propose a novel dimensionality reduction method, Delayed Latents Across Groups (DLAG). DLAG identifies both within- and across-area activity patterns (indicated by color and source/target of arrow; blue: within-A; red: within-B; magenta/orange: across-area). DLAG further disentangles the concurrent flow of population signals between two areas. The color of each across-area activity pattern depicts the direction of signal flow (magenta: A to B; orange: B to A), and shading (light vs dark) distinguishes distinct interactions.

Figure 2. DLAG conceptual illustration. From left to right: neurons, latent variables, and population space representations in two recorded brain areas analyzed by DLAG (top row / blue box: area A; bottom row / red box: area B). Left column: Activity of neurons simultaneously recorded in each area. Only three neurons are shown in each area for clarity. Center column: DLAG expresses the observed neural activity in each area as a linear combination (weighted sum) of two types of (low-dimensional) latent variables: within-area and across-area. Within-area variables are shown in the color corresponding to the area in which they belong (Within A: blue; Within B: red). For clarity, only two within-area variables are shown in each area, but in principle there may be a greater number. Across-area variables are shown in magenta. The magenta box inset overlaps the blue and red boxes for area A and B, respectively, to indicate that across-area variables are shared among neurons in both areas. The location of each across-area variable (i.e., within the bounds of Area A's box or area B's box) indicates which area's activity it reflects. Between area A and area B, across-area variables are vertically paired. The time courses of each pair are related after a time delay (D_1: delay between the left pair; D_2: delay between the right pair). The sign of this delay allows each pair to be associated with a directed interaction (A to B or B to A), which is indicated by gray arrows. Right column: The activity

of each neural population can be represented in a population space, where each axis represents the activity of a single neuron. Each point in population space represents the population activity at a particular time; and the points trace out a trajectory over time (magenta curve). DLAG identifies two linearly independent subspaces in each area: a within-area subspace (not shown, for clarity) and an across-area subspace (magenta-shaded plane). Each dimension of the across-area subspace is associated with a directed interaction, corresponding to those captured by the latent variables ('dim.': dimension).

Figure 3. DLAG accurately estimates within- and across-area time courses and their parameters. (**a**) Select single-trial latent-variable time course estimates on a representative dataset. Top row / blue box: area A; bottom row / red box: area B. For visual clarity, two latent variables of each type are shown (left: across-area; right: within-area). The total number of latent variables for the displayed dataset are as follows: across-area – 2; within-area A – 8; within-area B – 3. Orange dashed traces: DLAG estimates; black solid traces: ground truth. a.u.: arbitrary units. Scale bar: 250ms. Across all datasets for which across- or within-area dimensionality was

non-zero (across: 100; within A: 120; within B: 100), mean accuracy (R^2) of time course estimation was as follows: area A, across – 0.90; area B, across – 0.91; area A, within – 0.88; area B, within – 0.82 (all SEM values less than 0.01). Similarly, mean accuracy of subspace (loading matrix) estimation was as follows: C_1^a – 0.89; C_2^a – 0.93; C_1^w – 0.92; C_2^w – 0.94 (all SEM values less than 0.01). (**b**) Across-area dimensionality estimates versus the ground truth for all 120 synthetic datasets. (**c**) Within-area dimensionality estimates versus the ground truth for all 120 synthetic datasets (blue: within-area A; red: within-area B). For both (b) and (c), data points are integer-valued, but randomly jittered to show points that overlap. (**d**) Delay estimates versus the ground truth. Displayed error ('err.') indicates mean absolute error and SEM reported across 300 across-area variables. (**e**) Across-area Gaussian process (GP) timescale estimates versus the ground truth. Displayed error ('err.') indicates mean absolute error and SEM reported across 300 across-area variables. (**f**) Within-area A GP timescale estimates versus the ground truth. Displayed error ('err.') indicates mean absolute error and SEM reported across 900 within-area variables in area A. (**g**) Within-area B GP timescale estimates versus the ground truth. Displayed error ('err.') indicates mean absolute error and SEM reported across 300 within-area variables in area B.

Figure 4. Simultaneous population recordings in V1 and V2. (**a**) Schematic showing a sagittal section of occipital cortex and the recording setup. V1 population activity was recorded using a 96-channel Utah array. V2 population activity was recorded using a set of movable electrodes and tetrodes. (**b**) Peristimulus time histograms during the stimulus presentation period, for an example session and stimulus condition. For visualization only, neuronal spike trains were first smoothed using a sliding Gaussian window of width 20ms, and then z-scored to produce normalized firing rates. Neurons are ordered from top to bottom (separately for V1 and V2) according to the time at which their peak firing rate occurs. Scale bar: 160ms, the temporal period of the drifting grating stimulus. (**c**) Inter- and intra-areal comparisons. (Left) We applied DLAG to spike counts in V1 (light blue) and V2 (red). (Right) To contrast against DLAG's outputs from the V1-V2 analysis, we also applied DLAG to V1 only: we randomly split V1 into two equally sized populations, and then applied DLAG, as we did with V1 and V2. Each triangle represents a neuron. Box sizes illustrate typical relative population sizes.

Figure 5. Representative DLAG time courses for inter- and intra-areal analyses (same dataset as shown in Fig. 4b). (**a**) Inter-areal (V1-V2) time courses. Left: Across-area time courses. Right: Within-area time courses. Top row / light blue box: V1. Bottom row / red box: V2. Each panel corresponds to a latent variable, plotted vs. time. All time courses are aligned to stimulus onset. Scale bar: 160ms, the temporal period of the drifting grating stimulus. a.u.: arbitrary units. Each black trace corresponds to one trial; for clarity, only 10 of 400 are shown. Note that the

polarity of traces is arbitrary, as long as it is consistent with the polarity of C_i^a or C_i^w, as appropriate. Across-area variables are paired vertically; vertical arrows point in the direction of the lagging area, as determined by the sign of the delay next to each arrow. All delays for the displayed dataset were deemed significantly different from zero (see Methods). For visualization only, latent variables have been scaled and ordered by the fraction of shared variance they explain (across- and within-area variables are sorted separately; across-area variables are sorted according to shared variance explained in V2). All across-area variables and within-V2 variables uncovered by DLAG are shown here. The top 2 of 14 within-V1 variables are displayed, which explain 46% of V1's within-area shared variance. (**b**) Intra-areal (V1a-V1b) time courses. Left: Across-population time courses. Right: Within-population time courses. Top row / light blue box: V1a. Bottom row / dark blue box: V1b. All other conventions the same as in (a). Here, the delay for the third across-population variable (Across 3) was deemed to have an ambiguous sign, indicated by the bidirectional gray arrow. All other delays for the displayed dataset were deemed significantly different from zero, indicated by the unidirectional black arrows. Three of 10 across-population variables uncovered by DLAG are shown here, which explain 23% and 17% of V1a's and V1b's total shared variance, respectively. All uncovered within-V1a variables are shown, and 2 of 5 within-V1b variables are shown. Within-population variables (including those not shown here) explained 5% and 7% of V1a's and V1b's total shared shared variance, respectively.

Figure 6. DLAG reveals that V1-V2 interactions are selective and asymmetric. Top row: Summary of V1-V2 (inter-areal) interactions. Bottom row: Summary of V1a-V1b (intra-areal) interactions. (**a**) V1-V2 within- and across-area dimensionalities (determined via cross-validation). Histograms show distribution of within-V1 (light blue), within-V2 (red), and across-area (magenta) dimensionalities across 40 datasets. Arrows indicate the median of each distribution (within-V1: 14; within-V2: 2; across-area: 3). Within-V2 dimensionality was 0 in 5 of 40 datasets. (**b**) Shared variance explained by across-area activity in V1 and in V2. Black points: across-area strength in V2 vs. across-area strength in V1 on each of 40 datasets. Red point: median across datasets (V1: 0.34; V2: 0.76). Across-area strength is significantly greater in V2 than in V1 (one-sided paired sign test; $p < 0.001$). (**c**) V1a-V1b within- and across-population dimensionalities (determined via cross-validation). Histograms show distribution of within-V1a (light blue), within-V1b (dark blue), and across-population (magenta) dimensionalities across 40 datasets. Arrows indicate the median of each distribution (within-V1a: 2; within-V1b: 1; across-population: 11). Within-V1a dimensionality was 0 in 8 of 40 datasets; within-V1b dimensionality was 0 in 11 of 40 datasets. (**d**) Shared variance explained by across-population activity in V1a and in V1b.

Black points: across-population strength in V1b vs. across-population strength in V1a on each of 40 datasets. Red point: median across datasets (V1a: 0.96; V1b: 0.98). Across-population strength is not significantly greater in one population or the other (two-sided paired sign test; $p = 0.87$). (**e**) V1-V2 across-area Gaussian process (GP) timescale vs. time delay. Black points: across-area latent variables for which the delays were deemed significantly non-zero (see Methods; 95 of 135 across-area variables across all 40 datasets). Gray points: across-area latent variables for which delays were deemed ambiguous (not significantly positive or negative; 40 of 135 across-area variables across all 40 datasets). Red point: median across all significantly non-zero across-area variables (GP timescale: 37ms; Delay: -8ms). Red dashed line indicates the median delay. '***': delays are significantly biased in the negative (feedback; V2 to V1) direction (one-sided one-sample sign test on 'non-zero' delays, $p < 0.001$). (**f**) V1a-V1b across-population GP timescale vs. time delay. Same conventions as in (e). Out of 437 across-population latent variables uncovered across all 40 datasets, 316 were deemed significantly non-zero (black points), while 121 were deemed ambiguous (gray points). Red point: median across all significantly non-zero across-population variables (GP timescale: 27ms; Delay: -2ms). Red dashed line indicates the median delay. 'ns': delays are not significantly biased in the negative direction (one-sided one-sample sign test on 'non-zero' delays, $p = 0.08$). (**g**) GP timescales within V1 and within V2. Normalized histograms show the distribution of within-V1 (light blue) and within-V2 (red) GP timescales across all 40 datasets (total within-V1 latent variables: 562; total within-V2 latent variables: 121). Arrows indicate the median of each distribution (within-V1: 24ms; within-V2: 74ms). '***': within-V2 GP timescales are significantly longer than within-V1 GP timescales (one-sided Wilcoxon rank sum test, $p < 0.001$). (**h**) GP timescales within V1a and within V1b. Normalized histograms show the distribution of within-V1a (light blue) and within-V1b (dark blue) GP timescales across all 40 datasets (total within-V1a latent variables: 100; total within-V1b latent variables: 92). Arrows indicate the median of each distribution (within-V1a: 20ms; within-V1b: 23ms). '*': within-V1b GP timescales are significantly longer than within-V1a GP timescales (one-sided Wilcoxon rank sum test, $p < 0.05$).

Supplementary information

Supplementary Figure 1. DLAG graphical model representation. Filled circles represent observed variables (i.e., observed neural activity in each area), where $\mathbf{y}_{1,t}$ and $\mathbf{y}_{2,t}$ are the observed neural activity in area A and B, respectively, at time t. Unfilled circles represent latent variables, where $\mathbf{x}_t^a$ are across-area variables at time t; $\mathbf{x}_{1,t}^w$ and $\mathbf{x}_{2,t}^w$ are within-area variables in area A and B, respectively, at time t. Rectangles indicate model parameters; D represents the set of relative time delay parameters between the two areas. Color indicates a variable's or parameter's association with area A (blue), area B (red), or both (magenta). Arrows indicate statistical associations between variables. In particular, the arrows point from latent variables to observed neural activity, framing DLAG as a generative model. Thick black lines indicate that variables are related in time via a Gaussian process. Here two time steps are shown ($t-1$ and t), and time evolves from left to right.

Supplementary Figure 2. DLAG state model details and Gaussian process structure. (**a**) Within-area state model. Left column: Within-area time courses (area A: $\mathbf{x}^w_{1,j,:}$, blue points; area B: $\mathbf{x}^w_{2,j,:}$, red points) can be described as a finite number of samples drawn from a Gaussian process (GP). Right column: The temporal structure of each within-area GP is governed by a covariance function (area A: $k^w_{1,1}$; area B: $k^w_{2,1}$). The squared exponential (SE) function, chosen for the present work, is defined by a timescale parameter ($\tau^w_{1,j}$, $\tau^w_{2,j}$), which controls the width the covariance kernel, or equivalently, how quickly the latent variable changes over time. (**b**) An example set of within-area GP covariance matrices (K^w_j). The banded structure emerges from the choice of squared exponential function and stationarity of the GP covariance. Note the complete decoupling of within-area latent variables across areas: each latent variable has its own characteristic timescale, and cross-covariance terms are all zero. (**c**) Across-area state model. Left column: Like within-area time courses, across-area time courses can also be described as a finite number of samples drawn from a GP. Unlike within-area time courses, however, which are

independent, across-area time courses are coupled across areas, drawn from a common process $(\mathbf{x}^a_{j,:})$, but shifted by a relative delay (D_j). The samples mapped to each area (area A: $\mathbf{x}^a_{1,j,:}$, blue points; area B: $x^a_{2,j,:}$, red points) are aligned to the times at which observations occur (indicated by vertical gray dashed lines). Right column: The temporal structure of the common GP is governed by a SE covariance function. The width of the auto- and cross-covariances ($k^a_{i,i,j}$ and $k^a_{1,2,j}$, respectively) is controlled by a timescale parameter (τ^a_j). The center of the cross-covariance is controlled by the delay parameter D_j (positive delays: A leads B; negative delays: B leads A). (**d**) An example across-area GP covariance matrix (K^a). The banded structure emerges from the choice of squared exponential function and stationarity of the GP covariance. Note the non-zero cross-covariance terms in the off-diagonal blocks of K^a: the banded structure is shifted from the diagonal of each block by the delay parameter D_j.

Supplementary Figure 3. Model selection remained accurate in a regime of weak shared variance. To explore a scenario in which our model selection procedure began to misestimate within- or across-area dimensionalities, we synthesized an additional 120 independent datasets in which the strength of activity due to latent variables (i.e., shared variance in each area) was significantly weaker relative to the strength of variance independent to each neuron. Specifically, we generated datasets from the DLAG generative model such that the ratio $\mathrm{tr}(C_i C_i^\top)/\mathrm{tr}(R_i)$ was 0.1 for each area i (compared to 0.3 and 0.2 in area A and area B, respectively, in the original synthetic datasets; see Methods). All other data characteristics remained the same as in the original data. (**a**) Estimated versus ground truth across-area dimensionalities. (**b**) Estimated versus ground truth within-area dimensionalities. For both (a) and (b), data points are integer-valued, but randomly jittered to show points that overlap. Estimated across- and within-area dimensionalities never deviated from the ground truth by more than one, and the datasets with an absence of across- or within-area structure were always correctly identified. On datasets where a mismatch between estimated and ground truth dimensionalities did occur, the primary source of that inaccuracy likely originated from the initial factor analysis (FA) stage of model selection, rather than the second stage involving DLAG. Of the 66 datasets in which FA correctly estimated the total dimensionalities in both area A and area B, 65 datasets were correctly estimated following the second stage of model selection with DLAG models; and, irrespective of the accuracy after the FA stage, across-area dimensionality was correct on 101 of 120 datasets. This phenomenon might be expected, given the mismatch between the FA model and the DLAG generative model used to synthesize these datasets. It also provides the opportunity to explore the impact of imperfect dimensionality estimates—inevitable in real data—on the estimation and interpretation of DLAG's parameters and latent variables following fitting (see Supplementary Fig. 4, Supplementary Fig. 5).

Supplementary Figure 4. DLAG's parameter and latent variable estimates remained stable when dimensionality was underestimated. Here we present a case study from one of the synthetic datasets presented in Supplementary Fig. 3, in which the total dimensionality of area B was underestimated during the initial factor analysis (FA) model selection stage, and consequently so was across-area dimensionality. (**a**) Single-trial latent-variable time course estimates produced by a model with the correct number of within- and across-area latent variables (even though dimensionality was actually misestimated during model selection; see (b)). Top row / blue box: area A; bottom row / red box: area B. Left: across-area; right: within-area. One example trial shown. Orange dashed traces: DLAG estimates; black solid traces: ground truth. a.u.: arbitrary

units. Scale bar: 250ms. Delays reported in ms. Even in the weak-shared variance regime, estimates are qualitatively close to the ground truth. The asterisks ('*') are intended to highlight the third across-area latent variable for each area, which becomes misclassified as a within-area A latent variable when area B's dimensionality is underestimated (see within-area A latent variable 8 in (b)). (**b**) Single-trial latent-variable time course estimates produced by the model chosen through our model selection procedure. Same trial shown as in (a). The estimated number of latent variables in area B was one fewer than the ground truth; and so, too, was the estimated number of across-area variables. Still, qualitatively, time course estimates closely match those of the "matched" model in (a), in which the correct number of within- and across-area variables was used (compare estimated latent variables with the same index across (a) and (b)); and delay estimates are only slightly affected. By inspection, the third across-area latent variable pair (marked by the asterisks in (a)) now appears as the eighth within-area A latent variable (also marked by an asterisk). Note that the ordering of latent variables is not, in general, meaningful; we have ordered the latent variables here to facilitate visual illustration.

Supplementary Figure 5. DLAG's parameter and latent variable estimates remained stable when dimensionality was overestimated. Here we present a case study from one of the synthetic datasets presented in Supplementary Fig. 3, in which the total dimensionality of area B was overestimated during the initial factor analysis (FA) model selection stage, and consequently so was across-area dimensionality. (**a**) Single-trial latent-variable time course estimates produced by a model with the correct number of within- and across-area latent variables (even though dimensionality was actually misestimated during model selection; see (b)). Top row / blue box: area A; bottom row / red box: area B. Left: across-area; right: within-area. One example trial shown. Orange dashed traces: DLAG estimates; black solid traces: ground truth. a.u.: arbitrary

units. Scale bar: 250ms. Delays reported in ms. Even in the weak-shared variance regime, estimates are qualitatively close to the ground truth. The asterisk ('*') is intended to highlight the eighth within-area A latent variable, which becomes misclassified as an across-area variable when area B's dimensionality is overestimated (see across-area variable 3 in (b)). (**b**) Single-trial latent-variable time course estimates produced by the model chosen through our model selection procedure. Same trial shown as in (a). The estimated number of latent variables in area B was one more than the ground truth; and so, too, was the estimated number of across-area variables. Still, qualitatively, time course estimates closely match those of the "matched" model in (a), in which the correct number of within- and across-area variables was used (compare estimated latent variables with the same index across (a) and (b)). By inspection, the eighth within-area A latent variable (marked by the asterisk in (a)) now appears as the third across-area latent variable (also marked by asterisks). This phenomenon is straightforward to diagnose: here, we have additionally scaled each latent variable by the fraction of shared variance it explains within its respective area (see Methods; same convention as in Fig. 5). The third across-area latent variable explains little shared variance in area B, consistent with the ground truth. Note that the ordering of latent variables is not, in general, meaningful; we have ordered the latent variables here to facilitate visual illustration.

Supplementary Figure 6. DLAG performance improves with increasing number of trials. We sought to characterize DLAG's performance as a function of the number of available trials. We therefore synthesized 25 additional datasets (via the DLAG generative model) with the following characteristics: $N = 1000$ trials; $q_1 = q_2 = 50$ neurons per area; 500ms trial lengths with 20ms sample period (for $T = 25$ samples per trial); latent dimensionalities $p^a = p_1^w = p_2^w = 5$; signal-to-noise ratios $\text{tr}(C_1 C_1^\top)/\text{tr}(R_1) = \text{tr}(C_2 C_2^\top)/\text{tr}(R_2) = 0.3$; GP timescales $\tau^a, \tau_1^w, \tau_2^w \in [10, 150]$ms; and delays $D \in [-30, 30]$ms. We took subsets of trials from these datasets, and fit DLAG to increasingly large subsets (sizes equally spaced on a log scale from 10 to 1000 trials). (**a**) Accuracy of observation model parameter estimates versus number of trials. Top: Accuracy of across- and within-area loading matrix estimates increases with number of trials. (C_1^a: solid magenta; C_2^a: dashed magenta; C_1^w: solid blue; C_2^w: dashed red). Bottom: Covariance matrix (R: light gray) and mean parameter (**d**: dark gray) estimates improve with increasing number of trials. (**b**) Absolute error (in ms) of state model parameter estimates versus number of trials. Top: The error in across- and within-area GP timescale estimates decreases as number of trials increases (τ_1^a: magenta; τ_1^w: blue; τ_2^w: red). Error of within-area timescale estimates (τ_2^w) have been omit-

ted for values of 10 trials, where timescale estimates were too large to be meaningful relative to the length of trials. Bottom: The error in delay parameter estimates decreases as number of trials increases. (**c**) Accuracy (R^2) of latent variable time course estimates versus number of trials. Moment-by-moment estimates of latent variable time courses improve as the number of trials increases ($\mathbf{x}_1^a$: solid magenta; $\mathbf{x}_2^a$: dashed magenta; $\mathbf{x}_1^w$: solid blue; $\mathbf{x}_2^w$: dashed red). In (a)-(c) error bars represent SEM across 25 independent simulated datasets.

Supplementary Figure 7. DLAG performance improves with increasing number of neurons (and fixed latent dimensionality). We sought to characterize DLAG's performance as a function of the number of available neurons. We therefore synthesized 25 additional datasets (via the DLAG generative model) with the following characteristics: $N = 100$ trials; $q_1 = q_2 = 100$ neurons per area; 500ms trial lengths with 20ms sample period (for $T = 25$ samples per trial); latent dimensionalities $p^a = p_1^w = p_2^w = 5$; signal-to-noise ratios $\mathrm{tr}(C_1 C_1^\top)/\mathrm{tr}(R_1) = \mathrm{tr}(C_2 C_2^\top)/\mathrm{tr}(R_2) = 0.3$; GP timescales $\tau^a, \tau_1^w, \tau_2^w \in [10, 150]$ms; and delays $D \in [-30, 30]$ms. We took subsets of neurons from these datasets, and fit DLAG to increasingly large subsets (11, 25, 50, 75, and 100 available neurons in each area). (**a**) Accuracy of observation model parameter estimates versus number of neurons per area. Top: Accuracy of across- and within-area loading matrix estimates increases with number of neurons per area. (C_1^a: solid magenta; C_2^a: dashed magenta; C_1^w: solid blue; C_2^w: dashed red). Bottom: Covariance matrix (R: light gray) and mean parameter (**d**: dark gray) estimates improve with increasing number of neurons per area. (**b**) Absolute error (in ms) of state model parameter estimates versus number of neurons per area. Top: The error in across- and within-area GP timescale estimates decreases as number of neurons per area increases (τ_1^a:

magenta; τ_1^w: blue; τ_2^w: red). Error of within-area timescale estimates have been omitted for values of 11 neurons per area, where timescale estimates were too large to be meaningful relative to the length of trials. Bottom: The error in delay parameter estimates decreases as number of neurons per area increases. (**c**) Accuracy (R^2) of latent variable time course estimates versus number of neurons per area. Moment-by-moment estimates of latent variable time courses improve as the number of neurons per area increases ($\mathbf{x}_1^a$: solid magenta; $\mathbf{x}_2^a$: dashed magenta; $\mathbf{x}_1^w$: solid blue; $\mathbf{x}_2^w$: dashed red). In (a)-(c) error bars represent SEM across 25 independent simulated datasets.

Supplementary Figure 8. Uncertainty of DLAG timescale and delay estimates increases with increasing latent timescale. (**a**) Error (in ms; estimate minus ground truth value) of across-area GP timescale (top) and delay (bottom) estimates for each latent variable shown in Fig. 3c,d. The variance of both estimated GP timescales and delays appear to increase as the underlying ground truth GP timescale increases. (**b**) To verify the apparent trend in (a), we systematically characterized the accuracy of GP timescale and delay parameter estimates as a function of underlying GP timescale. We synthesized additional datasets (via the DLAG generative model) with the following fixed characteristics: $N = 100$ trials; $q_1 = q_2 = 50$ neurons per area; 500ms trial lengths with 20ms sample period (for $T = 25$ samples per trial); latent dimensionalities $p^a = p_1^w = p_2^w = 5$; signal-to-noise ratios $\mathrm{tr}(C_1 C_1^\top)/\mathrm{tr}(R_1) = \mathrm{tr}(C_2 C_2^\top)/\mathrm{tr}(R_2) = 0.3$; and delays $D \in [-30, 30]$ms. Each dataset's within- and across-area latent variables were given the same GP timescale; and across 150 datasets, we considered six different timescales (25 datasets synthesized for each timescale), ranging in length from half the sample period to the length of the trial (10ms, 20ms, 50ms, 100ms, 200ms, 500ms). Top: Absolute error (in ms) of across- and within-

area GP timescale estimates increases as underlying GP timescale increases (τ_1^a: magenta; τ_1^w: blue; τ_2^w: red). Bottom: Absolute error (in ms) of delay parameter estimates increases as underlying GP timescale increases. The lowest error is achieved when GP timescales are equal to sample period of observations. Error bars represent SEM across 25 independent simulated datasets.

Supplementary Figure 9. DLAG runtime as a function of number of trials, number of neurons, trial length, and latent dimensionality. (**a**) The average clock time (in seconds) per DLAG EM iteration scales (approximately) linearly with the number of trials. These runtime analyses were carried out on synthetic datasets with $q_1 = q_2 = 50$ neurons in each area; $T = 25$ time steps per trial; and latent dimensionalities $p^a = p_1^w = p_2^w = 5$ (total number of latent dimensions $2p^a + p_1^w + p_2^w = 20$). (**b**) The average clock time (in seconds) per DLAG EM iteration scales (approximately) linearly with the number of neurons per area. These runtime analyses were carried out on synthetic datasets with $N = 100$ trials; $T = 25$ time steps per trial; and latent dimensionalities $p^a = p_1^w = p_2^w = 5$ (total number of latent dimensions $2p^a + p_1^w + p_2^w = 20$). (**c**) The average clock time (in seconds) per DLAG EM iteration scales (approximately) quadratically with the number of time steps per trial. Runtime scales quadratically, rather than linearly (as in (a)), because DLAG describes the temporal structure within each trial via Gaussian processes. These runtime analyses were carried out on synthetic datasets with $N = 100$ trials; $q_1 = q_2 = 50$ neurons in each area; and latent dimensionalities $p^a = p_1^w = p_2^w = 5$ (total number of latent dimensions $2p^a + p_1^w + p_2^w = 20$). (**d**) The average clock time (in seconds) per DLAG EM iteration scales (approximately) quadratically with the total number of latent dimensions ($2p^a + p_1^w + p_2^w$). These runtime analyses were carried out on synthetic datasets with $N = 100$ trials; $q_1 = q_2 = 50$ neurons in each area; and $T = 25$ time steps per trial. In (a)-(d), error bars represent SEM across 25 independent simulated datasets. Results were obtained on a Red Hat Enterprise Linux machine (release 7.9, 64-bit) with 250GB of RAM running Matlab (R2019a), on an Intel Xeon CPU (E5-2695 v3, 2.3 GHz).

Supplementary Figure 10. V1-V2 spatiotemporal structure is preserved when V1 is subsampled to match V2 in population size. We sought to understand the extent to which the spatiotemporal characteristics reported for V1-V2 analyses were driven by disparities in population size between V1 and V2. For each dataset, we randomly subsampled the V1 population ('V1sub') to match the size of the V2 population. We then applied DLAG to each subsampled dataset in the same manner as in Fig. 6. (**a**) V1sub-V2 within- and across-area dimensionalities. Histograms show distribution of within-V1sub (light blue), within-V2 (red), and across-area (magenta) dimensionalities across 40 datasets. Arrows indicate the median of each distribution (within-V1sub: 7.5; within-V2: 3; across-area: 3). Within-V2 dimensionality was 0 in 1 of 40 datasets. (**b**) Shared variance explained by across-area activity in V1sub and in V2. Black points: across-area strength in V2 vs. across-area strength in V1sub on each of 40 datasets. Red point: median across datasets (V1sub: 0.33; V2: 0.70). Across-area strength is significantly greater in V2 than in V1sub (one-sided paired sign test; $p < 0.001$). It is possible that the difference in across-area strength seen in (b) is implied by, and therefore redundant with, the difference in within-area dimensionalities seen in (a). To test this possibility, we recomputed the median across-area strengths for V1 and V2, considering only datasets such that the distributions of within-V1sub and within-V2 dimensionalities were the same. Sixteen datasets remained after this distribution-matching procedure (the 16 datasets for V1 were not necessarily the same 16 datasets as for V2). The medians in V1 and V2 were nearly unchanged (V1sub: 0.33; V2: 0.67). Across-area strengths therefore convey a difference in the structure of V1 and V2 activity that could not be seen from differences in dimensionality alone. (**c**) V1sub-V2 across-area Gaussian process (GP) timescale vs. time delay. Black points: across-area latent variables for which the

delays were deemed significantly non-zero (see Methods; 97 of 136 across-area variables across all 40 datasets). Gray points: across-area latent variables for which delays were deemed ambiguous (not significantly positive or negative; 39 of 136 across-area variables across all 40 datasets). Red point: median across all significantly non-zero across-area variables (GP timescale: 32ms; Delay: -8ms). Red dashed line indicates the median delay. '***': delays are significantly biased in the negative (feedback; V2 to V1sub) direction (one-sided one-sample sign test on 'non-zero' delays, $p < 0.001$). (**d**) GP timescales within V1sub and within V2. Normalized histograms show the distribution of within-V1sub (light blue) and within-V2 (red) GP timescales across all 40 datasets (total within-V1sub latent variables: 307; total within-V2 latent variables: 153). Arrows indicate the median of each distribution (within-V1sub: 21ms; within-V2: 68ms). '***': within-V2 GP timescales are significantly longer than within-V1sub GP timescales (one-sided Wilcoxon rank sum test, $p < 0.001$).

Supplementary Discussion

Statistical tradeoffs between within- and across-area latent variables

Throughout this work, we have described how DLAG decomposes observed neural activity into a linear combination of within- and across-area latent variables. Equivalently, DLAG partitions each area's population space into distinct within- and across-area subspaces, which represent particular patterns of population activity (Fig. 2). Here we investigate more deeply why the within-area latent variables are a necessary model component, even if across-area activity is of primary scientific interest. Toward that end, we will consider an alternative interpretational perspective: namely, that DLAG performs a low-rank decomposition of the covariance matrix of a time-varying sequence. This alternative perspective also illuminates a general statistical phenomenon—not specific to DLAG—that any multi-area time series method must consider.

DLAG performs a low-rank covariance decomposition

Let us first express the DLAG model not only for a single time point, as in equation (14), but for all time points in a sequence. In particular, we will collect observed and latent variables in a manner that highlights group structure (i.e., organized differently than in equations (15) and (16)). Thus we define $\widetilde{\mathbf{y}}_1 = [\mathbf{y}_{1,1}^\top \cdots \mathbf{y}_{1,T}^\top]^\top \in \mathbb{R}^{q_1 T}$ and $\widetilde{\mathbf{y}}_2 = [\mathbf{y}_{2,1}^\top \cdots \mathbf{y}_{2,T}^\top]^\top \in \mathbb{R}^{q_2 T}$, obtained by vertically concatenating the observed neural activity $\mathbf{y}_{1,t}$ and $\mathbf{y}_{2,t}$ in areas 1 and 2, respectively, across all times $t = 1, \ldots, T$. We collect the across- and within-area latent variables for each area similarly. Let $\widetilde{\mathbf{x}}_1^a = [\mathbf{x}_{1,:,1}^{a\top} \cdots \mathbf{x}_{1,:,T}^{a\top}]^\top \in \mathbb{R}^{p^a T}$, $\widetilde{\mathbf{x}}_1^w = [\mathbf{x}_{1,:,1}^{w\top} \cdots \mathbf{x}_{1,:,T}^{w\top}]^\top \in \mathbb{R}^{p_1^w T}$, $\widetilde{\mathbf{x}}_2^a = [\mathbf{x}_{2,:,1}^{a\top} \cdots \mathbf{x}_{2,:,T}^{a\top}]^\top \in \mathbb{R}^{p^a T}$, and $\widetilde{\mathbf{x}}_2^w = [\mathbf{x}_{2,:,1}^{w\top} \cdots \mathbf{x}_{2,:,T}^{w\top}]^\top \in \mathbb{R}^{p_2^w T}$.

Then, we rewrite the state and observation models as follows:

$$
\begin{bmatrix}
\widetilde{\mathbf{x}}_1^a \\
\widetilde{\mathbf{x}}_1^w \\
\widetilde{\mathbf{x}}_2^a \\
\widetilde{\mathbf{x}}_2^w
\end{bmatrix}
\sim \mathcal{N}\left(
\begin{bmatrix}
\mathbf{0} \\
\mathbf{0} \\
\mathbf{0} \\
\mathbf{0}
\end{bmatrix},
\begin{bmatrix}
\widetilde{K}_{1,1}^a & \mathbf{0} & \widetilde{K}_{1,2}^a & \mathbf{0} \\
\mathbf{0} & \widetilde{K}_1^w & \mathbf{0} & \mathbf{0} \\
\widetilde{K}_{2,1}^a & \mathbf{0} & \widetilde{K}_{2,2}^a & \mathbf{0} \\
\mathbf{0} & \mathbf{0} & \mathbf{0} & \widetilde{K}_2^w
\end{bmatrix}
\right)
\tag{46}
$$

$$
\begin{bmatrix}
\widetilde{\mathbf{y}}_1 \\
\widetilde{\mathbf{y}}_2
\end{bmatrix}
\;\Bigg|\;
\begin{bmatrix}
\widetilde{\mathbf{x}}_1^a \\
\widetilde{\mathbf{x}}_1^w \\
\widetilde{\mathbf{x}}_2^a \\
\widetilde{\mathbf{x}}_2^w
\end{bmatrix}
\sim \mathcal{N}\left(
\begin{bmatrix}
\widetilde{C}_1^a & \widetilde{C}_1^w & \mathbf{0} & \mathbf{0} \\
\mathbf{0} & \mathbf{0} & \widetilde{C}_2^a & \widetilde{C}_2^w
\end{bmatrix}
\begin{bmatrix}
\widetilde{\mathbf{x}}_1^a \\
\widetilde{\mathbf{x}}_1^w \\
\widetilde{\mathbf{x}}_2^a \\
\widetilde{\mathbf{x}}_2^w
\end{bmatrix}
+
\begin{bmatrix}
\widetilde{\mathbf{d}}_1 \\
\widetilde{\mathbf{d}}_2
\end{bmatrix},
\begin{bmatrix}
\widetilde{R}_1 & \mathbf{0} \\
\mathbf{0} & \widetilde{R}_2
\end{bmatrix}
\right),
\tag{47}
$$

where $\widetilde{C}_1^a \in \mathbb{R}^{q_1 T \times p^a T}$, $\widetilde{C}_1^w \in \mathbb{R}^{q_1 T \times p_1^w T}$, $\widetilde{C}_2^a \in \mathbb{R}^{q_2 T \times p^a T}$, $\widetilde{C}_2^w \in \mathbb{R}^{q_2 T \times p_2^w T}$, $\widetilde{R}_1 \in \mathbb{S}^{q_1 T \times q_1 T}$, and

$\widetilde{R}_2 \in \mathbb{S}^{q_2 T \times q_2 T}$ are all block diagonal matrices comprising T copies of the loading matrices C_1^a,

C_1^w, C_2^a, and C_2^w; and observation noise covariance matrices R_1, and R_2, respectively. $\widetilde{\mathbf{d}}_1 \in \mathbb{R}^{q_1 T}$

and $\widetilde{\mathbf{d}}_2 \in \mathbb{R}^{q_2 T}$ are constructed by vertically concatenating T copies of mean parameters $\mathbf{d}_1$ and

$\mathbf{d}_2$, respectively. Note that equations (46) and (47) above are equivalent to equations (15) and

(16), but with variables rearranged.

Each within-area covariance matrix $\widetilde{K}_i^w \in \mathbb{S}^{p_i^w T \times p_i^w T}$, for area $i = 1, 2$ has the following block

structure:

$$
\widetilde{K}_i^w =
\begin{bmatrix}
\widetilde{K}_i^w(1,1) & \cdots & \widetilde{K}_i^w(1,T) \\
\vdots & \ddots & \vdots \\
\widetilde{K}_i^w(T,1) & \cdots & \widetilde{K}_i^w(T,T)
\end{bmatrix}
\tag{48}
$$

where each block $\widetilde{K}_i^w(t_1, t_2) = \mathrm{diag}(k_{i,1}^w(t_1,t_2), \ldots, k_{i,p_i^w}^w(t_1,t_2)) \in \mathbb{S}^{p_i^w \times p_i^w}, t_1, t_2 \in \{1,\ldots,T\}$ is

a diagonal matrix whose elements are computed according to the covariance function defined in

equations (4) and (5).

Each across-area auto- or cross-covariance matrix $\widetilde{K}_{i_1,i_2}^a \in \mathbb{R}^{p^a T \times p^a T}$, for areas $i_1, i_2 \in \{1,2\}$ has

analogous structure:

$$\widetilde{K}^a_{i_1,i_2} = \begin{bmatrix} \widetilde{K}^a_{i_1,i_2}(1,1) & \cdots & \widetilde{K}^a_{i_1,i_2}(1,T) \\ \vdots & \ddots & \vdots \\ \widetilde{K}^a_{i_1,i_2}(T,1) & \cdots & \widetilde{K}^a_{i_1,i_2}(T,T) \end{bmatrix} \tag{49}$$

where each block $\widetilde{K}^a_{i_1,i_2}(t_1,t_2) = \text{diag}(k^a_{i_1,i_2,1}(t_1,t_2),\ldots,k^a_{i_1,i_2,p^a}(t_1,t_2)) \in \mathbb{S}^{p^a \times p^a}, t_1, t_2 \in \{1,\ldots,T\}$ is a diagonal matrix whose elements are computed according to the covariance function defined in equations (7) and (8). Note that the cross-covariance matrices are transposes of one another, i.e., $\widetilde{K}^a_{i_1,i_2} = \widetilde{K}^{a\top}_{i_2,i_1}$.

Upon inspection of equation (46), the statistical dependency between latent variables becomes clear. However, the statistical dependency between observed neural activity in each area, $\widetilde{\mathbf{y}}_1$ and $\widetilde{\mathbf{y}}_2$, is not obvious, since the structure of equation (47) suggests that they might be decoupled. The relationship between observed areas becomes clear when we consider their joint distribution, after marginalizing out the latent variables:

$$\begin{bmatrix} \widetilde{\mathbf{y}}_1 \\ \widetilde{\mathbf{y}}_2 \end{bmatrix} \sim \mathcal{N}\left(\begin{bmatrix} \widetilde{\mathbf{d}}_1 \\ \widetilde{\mathbf{d}}_2 \end{bmatrix}, \widetilde{\Sigma} \right) \tag{50}$$

where

$$\widetilde{\Sigma} = \begin{bmatrix} \widetilde{C}^a_1 \widetilde{K}^a_{1,1} \widetilde{C}^{a\top}_1 + \widetilde{C}^w_1 \widetilde{K}^w_1 \widetilde{C}^{w\top}_1 + \widetilde{R}_1 & \widetilde{C}^a_1 \widetilde{K}^a_{1,2} \widetilde{C}^{a\top}_2 \\ \widetilde{C}^a_2 \widetilde{K}^a_{2,1} \widetilde{C}^{a\top}_1 & \widetilde{C}^a_2 \widetilde{K}^a_{2,2} \widetilde{C}^{a\top}_2 + \widetilde{C}^w_2 \widetilde{K}^w_2 \widetilde{C}^{w\top}_2 + \widetilde{R}_2 \end{bmatrix} \tag{51}$$

Equation (51) makes explicit the alternative interpretational perspective of DLAG: DLAG performs a low-rank decomposition of the covariance matrix $\widetilde{\Sigma}$. This decomposition is illustrated visually in Fig. S1a. For simplicity, we illustrate a covariance matrix for areas with three neurons each, over two time points. The shading of blocks of the covariance matrix illustrate which type of DLAG parameter is responsible for explaining that particular portion of covariance (magenta: across-area; blue/red: within-area; gray: independent single-neuron variability). Regions of overlap (i.e., where both blue/magenta or red/magenta shading are present) illustrate portions of covariance that both within- and across-area variables are responsible for explaining. Any regions of white indicate that no model parameters explain that portion of covariance.

The across-area parameters (note the fully magenta-shaded across-area covariance component in Fig. S1a) serve to explain covariance among all neurons, in both areas. Within-area parameters (blue and red shading, for areas 1 and 2, respectively) serve to explain covariance among neurons within each area, but not across areas (note the white across-area blocks for the within-area covariance component). Importantly, the only parameters in the DLAG model capable of explaining covariance across areas are the across-area parameters (only magenta shading is present in the across-area blocks of $\widetilde{\Sigma}$). And interestingly, within-area components fully overlap across-area components in the within-area blocks of $\widetilde{\Sigma}$, suggesting a potential redundancy. However, as we will discuss below, the overall structure of the decomposition shown in Fig. S1a is critical to the interpretation of across-area variables—that they isolate neural interactions *across* areas (and minimally reflect purely within-area interactions).

Within-area latent variables are critical to isolating across-area activity

To build further intuition, let us consider the scenario where within- and across-area covariances are modeled statically—without considering the flow of time (Fig. S1b). Static covariance decompositions result, for example, from the probabilistic canonical correlation analysis (pCCA) model [64], which includes static across-area latent variables and no within-area latent variables (within-area covariance is instead captured in non-diagonal observation noise covariance matrices, R_1 and R_2). The covariance matrix $\widetilde{\Sigma}$ still decomposes into across- and within-area components; however, covariances at non-zero time lags (i.e., the covariance between neural activity at a time point t_1 and a different time point $t_2 \neq t_1$, indicated by the white-shaded blocks of $\widetilde{\Sigma}$ in Fig. S1b) are all zero, by definition. Just like the DLAG case (Fig. S1a), only the across-area parameters can explain across-area covariance; and within-area components fully overlap across-area components in the within-area blocks of $\widetilde{\Sigma}$. Across-area activity is successfully isolated by across-area variables.

The problematic case arises when we update the across-area model to be time-varying, but leave the within-area model static (Fig. S1c). For example, what if we proposed a version of DLAG that did not include within-area latent variables, and instead simply adopted the same observa-

tion model as pCCA (i.e., non-diagonal observation noise covariance matrices, R_1 and R_2)? In this case, while the within-area model components do explain covariance among neurons within each area, they fail to capture any within-area covariance across non-zero time lags, by definition. This "shortcoming" leaves the across-area variables to explain within-area covariance *over time*. Visually, all non-zero time lag within-area blocks of the covariance matrix $\widetilde{\Sigma}$ have solely magenta shading (these blocks are highlighted by the '*' symbols in Fig. S1c). In contrast, the true DLAG model and fully static models avoid this pitfall. These successful models define no within-area blocks of $\widetilde{\Sigma}$ for which across-area parameters are solely responsible—or, visually, no within-area blocks that have solely magenta shading (Fig. S1a,b).

In summary, across-area variables can describe the activity of neurons in both areas; so in principle, it is perfectly valid for them to describe interactions that are actually shared only among neurons in one area, but not both. In the absence of within-area latent variables, then, across-area latent variables are forced to trade off between explaining activity that is truly shared between both areas and explaining activity that is present in only one area but statistically salient (i.e., high-variance). By capturing purely within-area interactions *over time*, within-area latent variables allow the across-area latent variables to isolate activity that is truly shared between both areas (as depicted in Fig. S1a). This statistical phenomenon applies to other statistical models, and is not specific to DLAG [36, 57].

a Time-varying across- and within-area models

b Static across- and within-area models

c Time-varying across-area model, but static within-area model

Figure S1. Full-sequence (trial) covariance matrix decompositions. For simplicity, in (a)-(c), we illustrate a covariance matrix for areas with three neurons each, over two time points. (**a**) DLAG decomposes the covariance of a full sequence (trial) into low-rank components. From left to right, panels represent the full covariance matrix; its across-area component; its within-area component; and its independent, single-neuron component. Across-area parameters (magenta shading) are solely responsible for explaining across-area covariance over time (i.e., there is no overlap of magenta with blue, red, or gray in the off-diagonal blocks of the full covariance matrix, on the left). Covariance among neurons within an area that cannot be explained by across-area covariance is captured by within-area parameters (area A: blue; area B: red). Note the overlap in blue/magenta and red/magenta shading in the within-area blocks of the full covariance matrix. (**b**) Models such as probabilistic canonical correlation analysis (pCCA), for example, similarly de-

compose the full-sequence covariance matrix into across- and within-area components, but make no attempt to model covariance over time, either across or within areas (indicated by blocks with white shading). (**c**) In the absence of a time-varying within-area model, across-area parameters fit to data are forced to explain salient within-area covariance over time. Note the within-area blocks of the full covariance matrix that have only magenta shading (indicated by the '*' symbols).