Structurally complex osteosarcoma genomes exhibit limited heterogeneity within

individual tumors and across evolutionary time

Sanjana Rajan[1,2*], Simone Zaccaria[3,4,5*], Matthew V. Cannon[2*], Maren Cam[2], Amy C.

Gross[2], Benjamin J. Raphael[3,6#], Ryan D. Roberts[2,7,8#]

1. Molecular, Cellular, and Developmental Biology Program, The Ohio State

University, Columbus, Ohio, USA.

2. Center for Childhood Cancers and Blood Diseases, Abigail Wexner Research

Institute at Nationwide Children's Hospital, Ohio, USA.

3. Department of Computer Science, Princeton University, Princeton, NJ, USA.

4. Computational Cancer Genomics Research Group, University College London

Cancer Institute, London, UK

5. Cancer Research UK Lung Cancer Centre of Excellence, University College

London Cancer Institute, London, UK

6. Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, USA.

7. The Ohio State University James Comprehensive Cancer Center, Columbus,

Ohio, USA.

8. Division of Pediatric Hematology, Oncology, and BMT, Nationwide Children's

Hospital, Columbus, Ohio, USA

*Contributed equally

#Contributed equally

Correspondence: Ryan D. Roberts

Email: Ryan.Roberts@nationwidechildrens.org

Running Title: Complex but stable copy-number profiles in osteosarcoma

Author Contributions: S.R., S.Z., M.V.C, B.J.P. and R.D.R. designed research, M.C.,

A.C.G., M.V.C. and S.R. performed research, S.Z. and M.V.C. analyzed the data. S.R.

and S.Z. wrote the paper, S.R., S.Z., M.V.C., B.J.P. and R.D.R. edited the paper.

Competing Interest Statement: The authors declare no potential conflicts of interest.

Keywords: Osteosarcoma, heterogeneity, somatic copy number aberration

Abstract

Osteosarcoma is an aggressive malignancy characterized by high genomic complexity.

Identification of few recurrent mutations in protein coding genes suggests that somatic

copy-number aberrations (SCNAs) are the genetic drivers of disease. Models around

genomic instability conflict - it is unclear if osteosarcomas result from pervasive ongoing

clonal evolution with continuous optimization of the fitness landscape or an early

catastrophic event followed by stable maintenance of an abnormal genome. We address

this question by investigating SCNAs in >12,000 tumor cells obtained from human

osteosarcomas using single cell DNA sequencing, with a degree of precision and

accuracy not possible when inferring single cell states using bulk sequencing. Using the

CHISEL algorithm, we inferred allele- and haplotype-specific SCNAs from this whole-

genome single cell DNA sequencing data. Surprisingly, despite extensive structural

complexity, these tumors exhibit a high degree of cell-cell homogeneity with little sub-

clonal diversification. Longitudinal analysis of patient samples obtained at distant

therapeutic time points (diagnosis, relapse) demonstrated remarkable conservation of

SCNA profiles over tumor evolution. Phylogenetic analysis suggests that the majority of

SCNAs were acquired early in the oncogenic process, with relatively few structure-

altering events arising in response to therapy or during adaptation to growth in

metastatic tissues. These data further support the emerging hypothesis that early

catastrophic events, rather than sustained genomic instability, give rise to structural

complexity, which is then preserved over long periods of tumor developmental time.

Significance Statement

Chromosomally complex tumors are often described as genomically unstable. However,

determining whether complexity arises from remote time-limited events that give rise to

structural alterations or a progressive accumulation of structural events in persistently

unstable tumors has implications for diagnosis, biomarker assessment, mechanisms of

treatment resistance, and represents a conceptual advance in our understanding of

intra-tumoral heterogeneity and tumor evolution.

Introduction

Osteosarcoma is the most common primary bone tumor affecting children and

adolescents[1]. Nearly always high grade and aggressive, this disease exhibits extensive

structural variation (SV) that results in a characteristically chaotic genome[2–4]. With few

recurrent point mutations in protein coding regions, osteosarcoma genomes often exhibit

widespread structural complexity, giving rise to associated somatic copy-number

aberrations (SCNAs), a likely genomic driver of malignant transformation[5]. Indeed,

osteosarcoma is the prototype tumor whose study led to the discovery of

chromothripsis[6,7], a mutational process that causes the shattering of chromosomes

leading to localized genomic rearrangements causing extreme chromosomal

complexity[8]. However, genomic complexity in osteosarcoma often goes beyond

alterations caused by the canonical processes associated with chromothripsis[9,10]. Many

have reasonably interpreted chromosomal complexity to be evidence of sustained

chromosomal instability (CIN), often with supporting evidence from other cancer types[11–

[14]. Indeed, cancer sequencing studies have identified the presence of extensive SCNAs

as a marker for CIN[13].

Two distinct models have been proposed to explain the evolution of chromosomal

structure and copy numbers in cancer genomes. One model suggests that underlying

genomic instability gives rise to populations of cells with diverse phenotypic variations

and that ongoing selection of advantageous phenotypes drives evolution and

adaptation[15,16]. A somewhat competing model argues that discrete periods of genomic

instability, isolated in tumor developmental time, give rise to extreme chromosomal

complexity driven by a small number of impactful catastrophic events[17,18]. In

osteosarcoma, investigators have put forward data that would seem to support both

models. For instance, several groups have used single cell RNA sequencing

experiments, which reveal a high degree of transcriptional heterogeneity, to infer a high

degree of copy number heterogeneity within osteosarcoma tumors[19,20], an observation

which would support a malignant process driven by ongoing instability and gradual

evolution. However, others have shown that SCNA profiles differ little when comparing

primary to metastatic or diagnostic to relapse samples[5,21,22], which would suggest that

ongoing mechanisms of malignancy do not create an environment of chromosomal

instability. Overall, it remains unclear whether the structurally complex genomes

characteristic of osteosarcoma emerge from continuous cycles of diversification and

fitness optimization within a context of ongoing instability and significant intra-tumoral

chromosomal heterogeneity or from an early catastrophic event that gave rise to

widespread structural changes that are then maintained over long periods of tumor

development, with evidence from the literature supporting both potential mechanisms.

One challenge in addressing this question comes from challenges in data interpretation

and deconvolution, as the existing studies describing copy number clonality and

evolution have inferred cell-specific copy number states from bulk tumor sequencing,

often from a single time point[23]. However, investigating ongoing clonal evolution from

bulk sequencing data remains particularly challenging, as each bulk tumor sample is an

unknown mixture of millions of normal and cancer cells[24–27]. The emergence of single

cell genomic DNA sequencing technologies now permits scalable and unbiased whole-

genome single cell DNA sequencing of thousands of individual cells in parallel[24,28],

providing an ideal framework for analyzing intra-tumor genomic heterogeneity and SCNA

evolution. Complementing these technical developments, recent computational

advances – most notably the CHISEL algorithm[27] – enable highly accurate ploidy

estimates and the inference of allele- and haplotype-specific SCNAs in individual cells

and sub-populations of cells from low coverage single cell DNA sequencing. This allows

cell-by-cell assessment of intra-tumoral SCNA heterogeneity, identification of allele-

specific alterations and reconstruction of the evolutionary history of a tumor from

thousands of individual cancer cells obtained at a single or multiple time points during

tumor progression.

Here, we leverage these approaches to determine whether the widespread SCNAs in

osteosarcoma result from ongoing genomic instability, providing a mechanism for tumor

growth and evolution. Using expanded patient tissue samples, our studies revealed

widespread aneuploidy and SCNAs in 12,019 osteosarcoma cells from ten tumor

samples. Using this approach, we found negligible intra-tumor genomic heterogeneity,

with remarkably conserved SCNA profiles when comparing either the individual cells

within a tumor or tumors collected at different therapeutic time points from the same

patient. These findings suggest that the widespread patterns of genomic SVs in

osteosarcoma are likely acquired early in tumorigenesis, and the resulting patterns of

SVs and SCNAs can be preserved within an individual tumor, across treatment time and

through the metastatic bottleneck.

Results

Individual cells within osteosarcoma tumors exhibit extensive SCNAs, but a high

degree of homogeneity

Single cell DNA sequencing was performed on 12,019 tumor cells from expanded

patient tissue samples. These nine patient tissues were obtained from diagnostic

biopsies of localized primary tumors ($n = 3$), from post-chemotherapy resection

procedures ($n = 2$), or from relapsed metastatic lung lesions ($n = 4$), representing a

spectrum of disease progression (Supplemental **Table S1**, Supplemental **Table S2**).

Apart from OS-17, a well-established model of metastatic osteosarcoma[29], all patient

tissues were expanded for a single passage in mice as either subcutaneous flank or

orthotopic bone tumors to obtain fresh tissue to perform single cell DNA sequencing

(300-2,500 single cells per sample; Supplemental **Figure S1A**). Previous studies have

shown that this procedure yields samples with a high degree of fidelity relative to the

diagnostic specimens, especially in early passages, an observation that we also

validated in our own samples[30]. We then used CHISEL[27] to identify allele- and

haplotype-specific SCNAs from the sequencing data.

Consistent with previous reports[6,31], sequencing showed a high degree of aneuploidy

and extensive SCNAs across the entire osteosarcoma genome (Figure 1). If driven by a

process of chromosomal instability and ongoing/continuous clonal evolution, we would

expect to observe multiple subclones with distinct complements of SCNAs within each

same tumor, such as has been shown in recent single cell studies of other cancer

types[24,27,32]. However, in each of the ten samples investigated, we identified one

dominant clone that comprised nearly all cells within each sample, with many samples

composed entirely of a single clone (Supplemental **Figure S1B**). To ensure that our

results were not an artifact caused by the algorithm or the selected thresholds for noise

control, we confirmed that the cells discarded as poor quality/noisy by CHISEL bear

SCNAs similar to the dominant clones identified in each sample – thus no rare clones

with distinct copy number profiles were discarded inappropriately (Supplemental **Figure**

S2). Interestingly, we found that a substantial fraction of the overall copy-number

changes involved allele-specific SCNAs, including copy-neutral LOHs (i.e., allele-specific

copy numbers {2, 0}) that would have been missed by previous analyses of total copy

numbers.

Figure 1. Extensive genomic complexity in ten expanded osteosarcoma patient tissue samples using single cell DNA sequencing. Allele-specific copy numbers (heatmap colors) are inferred by using the CHISEL algorithm[27] from each of ten datasets including 300-2300

single cancer cells from osteosarcoma tumors. In each dataset, cancer cells are grouped into clones (colors in leftmost column) by CHISEL based on the inferred allele-specific copy numbers. Corrected allele-specific copy-numbers are correspondingly obtained by consensus. Note that cells classified as noisy by CHISEL have been excluded. '*' and '#' represent samples obtained from the same patient.

Genome-wide ploidy of single cells showed high variability across samples, ranging from

1.5 to 4, demonstrating a high degree of aneuploidy (Supplemental **Figure S3**).

Consistent with the high levels of aneuploidy, we identified the presence of whole-

genome doubling (WGD, a phenomenon identified with much greater precision in the

single cell data) across nearly all cancer cells of six tumors (NCH-OS-8, OS-17, NCH-

OS-11, SJOS046149_X2, SJOS003939_X2 and SJOS003939_X1; **Figure 1 A-C, H-J**).

One tumor (SJOS003939_X2) shows two subclones that appear to be undergoing whole

genome duplication, with one subclone exhibiting a SCNA pattern that is almost exactly

double that of the other, across the genome.

To further assess tumor stability, we used paired datasets from patients collected at time

points along tumor progression. We observed that whole-genome copy-number profiles

were highly consistent within each patient. The first set includes NCH-OS-4, which was

obtained shortly after diagnosis at the time of resection (after two rounds of neoadjuvant

MAP chemotherapy), and NCH-OS-7, which was obtained at the time of relapse the

following year. Comparing genomic windows where at least one sample had a SCNA in

the primary clone, 77-78% of genomic windows had identical copy number assignments

in both samples, despite variation in tumor purity (**Supplemental Figure S4**). This

contrasts with between 1% and 35% concordance for non-related samples. The

correlation between related samples may be even higher, given inaccuracies expected

from low-coverage single cell SCNA detection.

The second set of paired primary and metastatic lesions (SJOS003939_X1,

SJOS003939_X2) also showed SCNA profiles that were highly similar (58% of SCNAs

identical, **Supplemental Figure S4**), suggesting a high degree of conservation of

genomic aberration profiles over therapeutic time. Overall, we observed a very high

degree of homogeneity within cancer cells sequenced from the same tumor. Even in

tumors where small proportions of cells (5-20%) are classified as part of small

subclones, these sub-clonal cells are only distinguished by modest SCNAs differences

within a few chromosomes. The exception to this general observation arose in

SJOS003939_X2, a second xenograft from a patient with a germline TP53, raising

suspicion for a second malignancy (rather than a relapse). Thus, despite the high levels

of aneuploidy and massive SCNAs identified in all ten samples, these osteosarcoma

cells demonstrated very modest levels of intra-tumor heterogeneity and variation across

therapeutic time.

Osteosarcoma cells harbor extensive SCNAs that mostly correspond to deletions.

The occurrence of WGD events correlates with high levels of aneuploidy and higher

frequency of SCNAs[33]. Recent reports have identified that WGDs serve as a

compensatory mechanism for cells to mitigate the effects of deletions[34]. We investigated

cell ploidy and fraction of genome affected by SCNAs (aberrant), amplifications,

deletions, and sub-clonal CNAs between tumors affected by WGDs (NCH-OS-8, OS-17,

NCH-OS-11, SJOS046149_X2, SJOS003939_X2 and SJOS003939_X1) and tumors not

affected by WGDs (NCH-OS-10, NCH-OS-4 and NCH-OS-7). Osteosarcoma cells in all

analyzed tumors demonstrate extensive SCNAs, affecting more than half of the genome

in every tumor cell. We found that the fraction of genome affected by SCNAs ranged

from 50-70% on average (**Figure 2A**, Supplemental **Figure S5A**). This result might not

be surprising for tumors affected by WGDs, however, we observed that tumors not

affected by WGD had a high fraction of aberrant genome as well (higher than 50% on

average; **Figure 2A**). This aberrant fraction is substantially higher than has been

reported for other cancer types[33].

Figure 2. Osteosarcoma cancer cells exhibit extensive genetic alterations, especially deletions, but a relatively low level of heterogeneity. (A) Ploidy (y-axis) and fraction of aberrant genome (x-axis) of every cell (point) across the ten analyzed datasets (colors). The kernel density of the marginal distributions of each value is reported accordingly in every plot. (B) Fraction of genome affected by deletions (x-axis) vs. fraction of genome affected by amplifications (y-axis) of every cell (point) across the ten analyzed datasets (colors). (C) Fraction of aberrant genome (x-axis) and fraction of sub-clonal SCNAs (i.e. fraction of the genome with SCNAs different than the most common clone for the same region across all cells in the same dataset, y-axis) of every cell (point) across the ten analyzed datasets (colors).

We observed a clear enrichment of deletions among the identified SCNAs across all

cancer cells. The fraction of genome affected by amplifications is 0-40% on average in

every tumor, while the fraction of the genome affected by deletions is 40-100% on

average across all cancer cells in every tumor (**Figure 2B**). This result is not particularly

surprising for tumors with WGD events and is consistent with a recent study of Lopez et

al.[34] that demonstrated a similar correlation in non-small-cell lung cancer patients.

However, in the osteosarcoma tumors analyzed in this study, we found that cancer cells

in non-WGD tumors are similarly affected by a high fraction of deletions (**Figure 2B**).

Importantly, we observed that >80% of the cells in all but two of our samples harbored

LOH events at the TP53 locus (in-line with frequency previously reported[3])

(**Supplemental Figure S5**). This substantiates the correlation between LOH of TP53

and high levels of genomic instability (including the occurrence of WGDs) reported in

recent studies[13,34,35], and suggests that these events might have a critical role in the

maintenance of a highly aberrant genomic state. Notably, CHISEL identified 50% of the

samples to harbor copy-neutral LOH alterations at the TP53 locus that would be missed

by total copy number analyses (**Supplemental Figure S5**).

We found it interesting that sub-clonal SCNAs that likely occurred late in the evolutionary

process (present only in subpopulations of cancer cells) are relatively rare across all

analyzed osteosarcoma tumors, irrespective of WGD status (with a frequency of 0-20%

in most cancer cells; **Figure 2C**, **Supplemental Figure S5**). Note the only exceptions to

this observation correspond to cells in NCH-OS-11, a sample with overall higher noise

and variance, and a subpopulation of cells in two other tumors (SJOS046149_X2 and

SJOS003939_X2) that appear to be cells that have not undergone WGD (**Supplemental**

Figure S5B). Indeed, the average fraction of SCNAs in SJOS046149_X2,

SJOS003939_X2 is lower than 20%. Overall, we observed that osteosarcoma cells

investigated in these ten samples, whether passaged in cell culture over a few

generations (OS-17), treatment naïve or exposed to extensive chemotherapy, bear high

levels of aneuploidy marked with extensive deletions and negligible sub-clonal

diversification, irrespective of WGD status.

Longitudinal single cell sequencing shows modest evolution of SCNA from

diagnosis to relapse

Increased aneuploidy has previously been associated with chromosomal instability (CIN)

and accelerated tumor evolution[13,36], though some have suggested that this observation

specifically applies to tumors that exhibit not only high levels of SCNA, but also high

levels of sub-clonal SCNA[37]. To assess the degree of structural instability exhibited by

these tumors, we examined a pair of samples, NCH-OS-4 and NH-OS-7, collected at

diagnosis and at relapse respectively, from the same patient to determine whether

SCNAs remained stable over therapeutic time or showed signs of significant

instability/evolution. This included an expansion in both the flank and orthotopic tibia

locations to determine whether these environments drove a niche-specific expansion of

a selected clone. Results suggest that expansion in mouse did not lead to evolutionary

disequilibrium.

We used CHISEL to jointly analyze 4,238 cells from these paired tumor samples and to

infer corresponding allele- and haplotype-specific SCNAs (**Figure 3A**). Based on

existing evolutionary models for SCNAs, we reconstructed a phylogenetic tree that

describes the evolutionary history of the different tumor clones identified in these tumors

(**Figure 3B**). The result from this phylogenetic analysis confirmed our findings in two

ways. First, we found that the evolutionary ordering of the different clones in the

phylogenetic tree is concordant with the longitudinal ordering of the corresponding

samples (**Figure 3B**): the tumor clones identified in the early sample (NCH-OS-4)

correspond to ancestors of all the other tumor clones identified in later samples (NCH-

OS-7-tib and NCH-OS-7-flank). Second, we observed that the vast majority of SCNAs

accumulated during tumor evolution are truncal, indicating that these aberrations are

accumulated early during tumor evolution and shared across all the extant cancer cells

(**Figure 3B**). In fact, only three significant events distinguish the most common ancestor

of all cells from this patient (identified in NCH-OS-4) from the cells within the relapse

lesion: gain of chromosome 14, gain of chromosome 16q (resulting in copy-neutral

LOH), and deletion of one allele of chromosome 18 (resulting in LOH). Note that we

cannot be certain of when these clones arose. It is possible these changes occurred

early in tumor formation and were present in the primary tumor but were not present in

the biopsied sample and so we must exercise caution when assessing if there is ongoing

low-level chromosomal instability.

To further assess the effects that environmental stressors might play on the creation

and/or emergence of sub-dominant clones, which could be masked due to extreme

rarity, we expanded samples from the same tumor within two different

microenvironments in mice. Consistent with the diagnosis-relapse sample comparison,

clones identified within tumors grown orthotopically within the tibia (NCH-OS-7-tib) or

within subcutaneous flank tissues (NCH-OS-7-flank) are highly concordant (78% of

genomic windows called with identical SCNA values) and distinguished by few focal

SCNAs (primarily single copy changes). These changes could be either be variance in

SCNA calling from the sequencing data, stochastic differences caused by the presence

of sub-clones within the original tumor sample that was bisected and implanted or

biologically relevant. Without targeted studies it is not possible to confidently define the

biological role of these focal changes, if any. A third comparison of tumors separated in

time and space was possible using another paired set of primary and metastatic lesions

(SJOS003939_X1, SJOS003939_X2), which also demonstrated negligible sub-clonal

diversification (**Supplemental Figure S7**). Indeed, each sample was dominated by one

major clone, which exhibited only subtle differences from the paired sample. While these

results do not suggest the absence of SCNA changes over the course of tumor

evolution, they do suggest a level of stability quite similar to genomically simple tumors

and that the mechanisms giving rise to these limited focal changes are different from

those that gave rise to widespread genomic complexity.

To further explore temporal and spatial consistency of patient tumor samples, we

combined whole genome sequencing data obtained from paired osteosarcoma samples

within the St. Jude database[38–40] with our own whole genome sequencing and performed

SCNA analysis. This combined data yielded between two and six tumor samples for

each patient, in addition to a germline reference sample. In most cases, all samples

taken from a single patient at different timepoints were highly similar and clustered

together (**Figure 4** and **Supplemental Figure S8A-K**). There were, however, five

samples that had more than one distinct clone in separately collected samples which

reduced the overall average. In these instances, the average correlation between clonal

populations within a patient was only 0.28, which was close to the correlation we

observed between samples taken from different patients (mean Pearson cor = 0.18).

Deeper exploration of these samples revealed germline TP53 mutations in some

patients (shown with a red asterisk in **Figure 4**), suggesting an underlying cancer

predisposition and a likelihood that these are tumors arising from distinct oncogenic

events. The correlation within a clone was very high (mean Pearson cor = 0.67), despite

the noise created by the sparse coverage sequencing inherent to this method.

Xenograft-derived samples did not cluster separately from samples derived directly from

patients (**Figure 4**), except for two samples from SJOS030645 which formed a distinct

cluster. The xenograft samples had a high correlation with non-xenograft samples from

the same patient (mean Pearson cor = 0.70), suggesting that the SCNA patterns in

these samples were not dramatically altered by clonal selection within the mouse.

Determination of SCNA-based clonal composition and tumor purity was performed using

the HATCHet algorithm[41], providing additional context for interpretation of results.

HATCHet results show a very high degree of SCNA-based clonal conservation from one

clinical timepoint to the next.

Figure 3. Phylogenetic reconstruction of tumor evolution is consistent with longitudinal ordering of matched tumor samples and reveals conservation of SCNA profiles. (A) Allele-specific copy numbers (heatmap colors) across all autosomes (columns) have been inferred by CHISEL jointly across 4238 cells (rows) in 3 tumor samples from the same patient: 1 pre-treatment sample (NCH-OS-4 tibia) and two post-treatment samples (NCH-OS-7 tibia and NCH-OS-7 flank). CHISEL groups cells into 4 distinct clones (blue, green, red, and purple) characterized by different complements of SCNAs. (B) Phylogenetic tree describes the evolution in terms of SCNAs for the four identified tumor clones. The tree is rooted in normal diploid clone (white root) and is characterized by two unobserved ancestors (white internal nodes). Edges are labelled with the corresponding copy-number events that occurred and transformed the copy-number profile of the parent into the profile of the progeny. The four tumor clones (blue, green, red, and purple) are labelled according to the sample in which they were identified.

Figure 4. CNA correlation between osteosarcoma samples. Pearson R values denoting
correlation of binned copy numbers between samples. Colors on x and y-axes indicate each
sample's patient of origin and type as well as the clones defined from the correlation analysis.
Red asterisks denote samples from patients with germline TP53 mutations. Note that
SJOS003939_X1 is from the same patient as SJOS031478_* samples.

Discussion

Osteosarcoma is one of several malignancies typified by chaotic genomic landscapes

dominated by structural variation and corresponding copy number alterations[3].

Chromosomal complexity in osteosarcoma and other cancers with complex karyotypes

has often been assumed to represent underlying genomic instability, suggesting that

these tumors gradually accumulate structural changes that lead to increasing

complexity, with continual selection of ever more aggressive clones driving malignant

progression. This concept was supported by previous reports demonstrating that, in

some patients, spatially separated tumor samples exhibit slightly divergent SNP and

SCNA patterns [42,43]. These studies also noted that, while there is heterogeneity between

samples, there seem to be clones that are shared across multiple metastatic foci. By

nature, these studies understandably focused on identifying SNP and copy number

differences contained within distinct lesions in these highly aggressive tumors, with the

largest sample sets collected at autopsy. By utilizing single cell DNA sequencing, we

have been able to investigate intra-tumor genomic heterogeneity and tumor evolution in

concrete ways that were previously possible only by estimation and inference using bulk

sequencing methods[44–47]. This allowed us to ask different questions related to the

stability of these complex genomes within a tumor sample. We were surprised to find

that cells within a tumor demonstrated surprisingly little cell-to-cell variability in SCNA

profiles - results that, at first, seemed discordant with previous reports of intra-tumoral

genomic heterogeneity in osteosarcoma[19].

Analyzing longitudinal sets of paired samples, we showed that these particular

osteosarcoma tumors maintained relatively stable SCNA profiles from diagnosis to

relapse, primary tumor to metastasis, and during growth in two distinct environments.

Our phylogenetic analysis suggested that the most recent common ancestor of these

related samples harbored almost all of the observed SCNAs, suggesting that most of the

genomic aberrations arose early in the tumorigenic process within these patients,

followed by a long period of stable clonal expansion (clonal stasis)[18]. Our analysis of

bulk whole genome sequence data from St. Jude supports this observation and

highlights an important observation. Where we observed multiple clones in our single

cell data, each clone was homogeneous in its SCNA patterns across cells within the

clone, but highly distinct from other clones (**Figure 1**). We observed the same

phenomenon in the bulk St. Jude data where a single clone detected in multiple

temporally or anatomically separated samples had highly conserved SCNAs while

distinct clones were highly divergent (Figure 4). This suggests that each of these clones

either derive from a very early event that produced multiple distinct clones, or

independent tumorigenesis events.

An inherent limitation of single cell analysis of biopsy samples is that they are not

representative of the entire tumor and so the homogeneous cell populations we observe

could, in part, derive from the small sample size involved. However, our data include

independent data from multiple biopsies that showed similar clonal patterns. Also, our

analysis of the St. Jude samples includes multiple independent biopsies from patients

and demonstrates the same pattern of SCNA conservation across samplessingle cell. A

potential unexpected advantage of the small sample size inherent to tumor biopsies is

that these samples tended to be clonal in nature in our single cell data. Given this, it may

be feasible to assume that bulk SCNA results are representative of most cells within the

biopsy.

Another limitation of biopsy samples is the potential for normal cell types within the

sample to interfere with the evaluation of SCNAs. If too large of a proportion of normal

cells are present, estimates of copy number will be less accurate. For instance,

Supplemental **Figure S8A** shows sample SJOS031478_D1 which has very low copy

number alteration values, suggesting that this sample may have a large proportion of

normal cells, making detection of SCNA difficult. Deconvolution can improve, but not

fully overcome, this issue[41]. To help compensate for this issue, for **Figure 4** we used

correlation between samples instead of comparison of absolute copy numbers. This

allows sample SJOS031478_D1 to cluster closely with SJOS031478_D2 in **Figure 4**

despite apparent contamination with normal tissue. Samples SJOS031478_D1 and

SJOS031478_D3 had very low correlation in their SCNA patterns. It is notable that these

samples harbor distinct SCNAs including a deletion of a large portion of chromosome

five in SJOS031478_D3 that is absent from SJOS031478_D1 and a large amplification

of chromosome eight present in SJOS031478_D1 but absent from SJOS031478_D3

(Supplemental **Figure S8A**) indicating that these are distinct clones. Similar patterns can

be seen in Supplemental **Figure S8J** where SJOS046149_R2 and SJOS046149_X2 are

distinct from SJOS046149_R1, SJOS046149_X1 and SJOS046149_X1b.

One genomic change that was readily evident within our data was the common

occurrence of WGD. Using the CHISEL algorithm[27], we identified high levels of

aneuploidy and extensive genomic aberrations that were dominated by deletions within

these osteosarcoma tumors. Consistent with previous reports suggesting WGD as a

mechanism to mitigate the effects of widespread deletions[34], we identified extensive

deletions even in tumors that had not undergone duplication. Indeed, some of our

samples showed subclones of cells that differed across the genome by almost exactly

two-fold, which may represent populations of cells that had undergone duplication (with

the duplicated fraction being the dominant clone). These findings support the hypothesis

that duplication is a process that produces a more aggressive clone from cells that are

first affected by widespread deletion.

To expand the analysis addressing the question of stability beyond our single cell WGS

samples, we evaluated SNCAs in bulk WGS data derived from osteosarcoma samples.

We investigated paired tumor samples across 14 patients (and included the associated

patient-derived xenografts, where available) to determine if SCNA patterns were stable

across time. We observed that there were both identical and divergent clones within

single patients. Clones were similar with correlations as high as 0.92. In the few patients

where relapse specimens contained highly divergent clones (**Supplemental Figure**

S8A-K), a deeper analysis revealed germline TP53 mutations in many cases (**Figure 4**).

In these patients harboring a genetic predisposition to developing osteosarcoma, it is

likely that these genetically distinct lesions represent independent oncogenic events and

it is possible that TP53 activity was impaired through alternative means in the other

patients.

Historically, studies in osteosarcoma (and other cancers) have equated a high level of

SCNA with ongoing genomic instability[31,48], and some direct evidence has supported this

concept[43]. However, several recent studies seem to challenge this conclusion, showing

preservation of SCNA profiles in primary vs metastatic and diagnostic vs relapse

samples[5,22,42]. Our findings support the hypothesis that mechanisms leading to

widespread structural alterations are active early in tumorigenesis but resolve and are

followed by long periods of relative stability. These seemingly discordant observations

may both be true. First, there may be different paths to chromosomal complexity in

different tumors—processes that resolve in some tumors, but do not in others. Indeed,

nearly all these publications contain sample sets that seem to support higher and lower

levels of chromosomal (in)stability.

Second, tumor cells may experience time-limited periods of relative instability, resulting

in the phenomenon of punctuated equilibrium, as has been shown in other cancer

types[18]. In a punctuated equilibrium scenario, the timing of the biopsy would change the

likelihood of finding more or less SCNA heterogeneity within the tumor using methods

like single cell WGS.

To synthesize these concepts, there are several potential models for the emergence of

SCNA-defined clones in osteosarcoma (**Figure 5**). A single initiation event giving rise to

a single dominant clone followed by highly stable genomic organization (**Figure 5A**)

would cause all tumor samples from a patient to have consistent SCNA patterns in both

bulk and single cell sequencing. This mechanism, however, would not be supported by

previously published data [42,43] or the results presented here.

Figure 5. Possible models for temporal SCNA stability. (A) After tumor initiation, if chromosomal instability is low, tumors will have identical SCNA patterns across all cells. (B) High tumor instability would result in tumors with highly heterogeneous SCNA patterns across cells which may not be apparent in bulk sequencing. (C) If there are multiple initiation events with low subsequent genomic instability SCNA patterns will be consistent across clones, which will be apparent by both bulk and single cell sequencing. (D) If there are multiple initiation events with ongoing genomic instability, clones derived from each will be similar but with highly variable

434 SCNA patterns within a clone. This heterogeneity would be apparent by single cell sequencing
435 but not bulk sequencing. (E) If a single initiation event is followed by an initial period of genomic
436 instability, divergent clones could emerge. Patterns of stability within a clone would suggest that
437 chromosomal stability is re-established prior to clonal expansion.
438

In an alternative model, instability persists from the oncogenic insult forward (**Figure**

5B). This model would produce samples containing multiple divergent subclones, which

would be evident in bulk sequencing collected in different loci, though not detectable

within a single sample. Single cell analyses would identify several SCNA-defined clones

within each sample. This model seems less likely considering our single cell results.

A third outcome could result if there were multiple independent initiation events,

producing several competing clones within a tumor, followed by a period of stable

chromosomal organization (**Figure 5C**). This mechanism, which is consistent with the

punctuated equilibrium hypothesis[18], could produce multiple samples from a patient with

divergent SCNA patterns by bulk sequencing, however, cells within each sample would

demonstrate highly similar copy number patterns (assuming the sample does not

overlap a boundary between clones). This model agrees with both published

observations[42,43] and the results presented here.

A slight modification of this model would invoke early mechanisms giving rise to multiple

competing clones, followed by a period where an independent mechanism causes

ongoing low-level chromosomal instability within each founder clone (**Figure 5D**). This

mechanism would generate multiple slightly divergent clones within each sample. We

see some evidence to support this model, such as the similar, but distinct, clones

observed in the NCH-OS-7 samples in Figures 1 and 3. These patterns could also be

explained by tissue sampling bias, experimental variance or computational noise caused

by the low sequencing depth inherent to single cell data. A larger study would be needed

to evaluate this.

A final model, which would also be consistent with both our single cell data and the

published record, suggests a single initiation event followed by a period of where

daughter cells exhibit chromosomal instability (**Figure 5E**), creating a diversity of

competing clones. Eventually, clones emerge that exhibit chromosomal stability and

have a competitive advantage. In this scenario, tumor cells within a patient are distinct

from clone to clone, but homogeneous within a clone, with a small subset of shared

SCNAs that were present in the origin cell and maintained through the subsequent

chromosomal instability.

While copy number patterns might be stable after an initial structure-altering event,

SNVs arise through completely different mechanisms and likely have different

evolutionary dynamics. Previous studies looking at osteosarcoma across therapeutic

time show clear sequence related changes dominated by patterns that suggest a

cisplatin induced mutational burden [43]. However, the structural integrity of the

chromosomes does not seem to be affected by treatment[42,43].

A brief (single cycle) expansion of tissue using an animal host proved useful for

generating high-quality single cell suspensions of sufficient quantity while maintaining

high fidelity to the original patient sample. This approach is not intended to model tumor

progression in a murine host, but rather to maximize the data obtained from each of

these incredibly valuable samples. Some have expressed concerns that mouse-specific

evolution selects for sub clonal populations[49]. However, the mouse-specific evolution

that occurs over many passages (such as in the development of a PDX) does not occur

when the mouse is used as a vehicle for brief expansion[30]. Our SCNA analysis

comparing results of these expanded tissues to bulk sequencing performed directly on

the patient samples showed a very high correlation between expanded and primary

samples. Therefore, this approach may represent a productive compromise enabling

multiple lines of research on tissues with limited availability in rare diseases.

Our findings of clonal stasis in osteosarcoma sheds some light on the complex

evolutionary history of this cancer type and could have important implications for tumor

evolution, patient diagnosis and treatment of osteosarcoma. However, a much larger

sample size of patient tissues is needed to capture the full heterogeneity of

osteosarcoma seen in the human disease and describe the prevalence of multiple tumor

sub-clones. Somewhat ironically, one may conclude from this data that bulk sequencing

methods likely produce an adequate assessment of SCNA profiles and heterogeneity in

osteosarcoma, given the lack of heterogeneity found in our analysis. These data likewise

suggest that, in a clinical setting, sequencing analyses based on SCNA likely remain

valid, even into treatment and relapse, assuming separate samples derive from the

same clonal tumor population.

At a biological level, these results support the early catastrophe model as a primary

mechanism of osteosarcoma complexity, suggesting that most structural

rearrangements occur early in the tumorigenic process. While other rearrangements

certainly can occur during malignant progression, subsequent structural events do not

appear to be *necessary* for invasion, metastasis, or therapeutic resistance (though they

certainly may *contribute* to such processes), nor do they appear to be the same

mechanisms that create widespread structural complexity. Ongoing research will

continue to inform our understanding of the contributions that initial catastrophic events

and ongoing mechanisms of genomic evolution have and how they influence clinical

outcomes.

It is important to note that our study is performed in a way that is generally insensitive to

other alterations (such as SNVs) as a source of genomic variation, though few recurrent

mutations have been identified in osteosarcoma, despite extensive genetic analysis[5,48,50].

If both observations hold true, one must conclude that the acquisition of traits that drive

malignant progression arise through epigenetic-based evolutionary processes, which

remain poorly understood. Interestingly, we and others have shown that these same

osteosarcomas demonstrate a high level of intra-tumor transcriptional heterogeneity[19,51].

This heterogeneity of gene expression in cells that are genomically homogeneous

suggests that there may be microenvironmental differences or an underlying epigenetic

heterogeneity, which could be a basis for competition and selection of tumor cells.

Materials and Methods

Experimental model – Expanded patient tissues and murine studies

Expanded patient tissue. Patient samples NCH-OS-4, NCH-OS-7, NCH-OS-8, NCH-OS-

10 and NCH-OS-11 were obtained from patients consented under an Institutional

Review Board (IRB)-approved protocol IRB11-00478 at Nationwide Children's Hospital

(Human Subject Assurance Number 00002860). Germline whole genome sequencing

(WGS) was generated from patient blood collected under IRB approved protocol IRB11-

00478. Patient samples SJOS046149_X1, SJOS046149_X2, SJOS003939_X1 and

SJOS031478_X2, with matched normal WGS were received from St. Jude's Children's

Research Hospital through the Childhood Solid Tumor Network[38–40]. The OS-17 PDX

was established from tissue obtained in a primary femur biopsy performed at St. Jude's

Children's Research Hospital in Memphis and was a gift from Peter Houghton[29].

Murine Studies. Flank tumors. Viable tissue fragments from patient tissue were

expanded in C.B-17/IcrHsd-Prkdc[scid] mice as subcutaneous tumors following approved

IACUC protocols. These tumors were allowed to grow to 300-600 mm^3 before harvest.

Passage 1 expanded tissue was used for all samples, with the exception of OS-17 (p18).

Orthotopic primary tumors. Single cell suspensions of 5x10^5 cells were injected intra-

tibially in C.B-17/IcrHsd-Prkdc^{scid} mice as per IACUC guidelines. These tumors were

harvested once they grew to 800 mm^3 and prepped for single cell DNA-seq.

Single cell suspension and DNA library generation

Tumors harvested from mice were processed using the human tumor dissociation kit

(Miltenyi Biotec, 130-095-929) with a GentleMacs Octo Dissociator with Heaters

(Miltenyi Biotec, 130-096-427). Single cell suspensions in 0.04% BSA-PBS of

dissociated tumor tissues were generated and frozen down using the 10X freezing

protocol for SCNA. The frozen down single cell suspensions were processed using the

Chromium Single Cell DNA Library & Gel Bead Kit (10X genomics #1000040) according

to the manufacturer's protocol with a target capture of 1000-2000 cells. These barcoded

single cell DNA libraries were sequenced using the NovaSeq 6000 System using paired

sequencing with a 100bp (R1), 8bp (i7) and 100bp (R2) configuration and a sequencing

coverage ranging from 0.01X to 0.05X (~0.02X on average) per cell. Germline WGS was

performed on NovaSeq SP 2x150BP.

Single cell SCNA calling using CHISEL

Paired-end reads were processed using the Cell Ranger DNA Pipeline (10X Genomics),

obtaining a barcoded BAM file for every considered single cell sequencing dataset. As

described previously[27], the pipeline consists of barcode processing and sequencing-

reads alignment to a reference genome, for which we used hg19. We applied CHISEL

(v1.0.0) to analyze each generated barcoded BAM file using the default parameters and

by increasing to 0.12 the expected error rate for clone identification in order to account

for the lower sequencing coverage of the analyzed data[27]. In addition, we provided

CHISEL with the available matched-normal germline sample from each patient and

phased germline SNPs according to the recommended pipeline by using Eagle2 through

the Michigan Imputation Server with the Haplotype Reference Consortium (HRC)

reference panel (v.r1.1 2016). CHISEL inferred allele- and haplotype-specific copy

numbers per cell and used these results to group cells into distinct tumor clones, while

excluding outliers and likely noisy cells. To determine fraction of aberrant genome

(genome affected by SCNAs), we defined aberrant as any non-diploid genomic region

(i.e., allele-specific copy numbers different than {1, 1}) in tumors not affected by WGDs

(NCH-OS-10, NCH-OS-4, and NCH-OS-7) or any non-tetraploid genomic region (i.e.,

allele-specific copy numbers different than {2, 2}) in tumors affected by WGDs (NCH-

OS-8, OS-17, NCH-OS-11, SJOS046149_X2, SJOS003939_X2 and SJOS003939_X1).

We defined deletions as previously described in cancer evolutionary studies[26,52–54]. We

say that a genomic region in a cell is affected by a deletion when any of the two allele-

specific copy numbers inferred by CHISEL is lower than the expected allele-specific

copy number (1 for non-WGD tumors or 2 for tumors affected by WGD). Conversely, a

genomic region is amplified when any of the two allele-specific copy numbers is higher

than expected.

Reconstruction of copy-number trees

We reconstructed copy-number trees for tumor samples NCH-OS-4 (tibia), NCH-OS-7

(flank) and NCH-OS-7 (tibia), to describe the phylogenetic relationships between distinct

tumor clones inferred by CHISEL based on SCNAs using the same procedure proposed

in previous studies[27]. Briefly, we reconstructed the trees using the maximum parsimony

model of interval events for SCNAs[52,53] and the copy-number profiles of each inferred

clone. These copy number profiles were obtained as the consensus across the inferred

haplotype-specific copy numbers derived by CHISEL for all the cells in the same clone,

where we also considered the occurrence of WGDs predicted by CHISEL. We classified

copy-number events as deletions (i.e., del), as LOH which are deletions resulting in the

complete loss of all copies of one allele (loh), as copy-neutral LOH which are LOHs in

which the retained allele is simultaneously amplified, and as gains (gain).

SCNA calling on whole genome data

To compare SCNA patterns across multiple tumor samples from the same patients, we

downloaded a total of 47 whole genome sequence datasets from St. Jude's DNAnexus

from 14 patients including germline data and multiple tumor samples (diagnosis, relapse,

metastasis and xenograft). We also included the seven scSCNA datasets we generated

which had matched germline whole genome data in the St. Jude data and treated these

as bulk sequencing data for this analysis. We used samtools[55] to convert the bam files to

fastq and aligned all datasets to a joint hg38/mm10 reference. We filtered out all mouse

sequences and removed PCR duplicates. We then called SCNAs with Varscan[56]. Next,

we combined all SCNA data by calculating the median copy number for 1,000 bp non-

overlapping bins. Correlation between samples was calculated using the cor function in

R and the resulting output was plotted as a heatmap using the pheatmap R package

(https://github.com/raivokolde/pheatmap).

SNP calling on whole genome data

To assess genetic heterogeneity of all samples, we produced phylogenetic trees from

SNP data. We used the bam alignment files produced during the SCNA calling analysis

and called SNPs using bcftools' mpileup function[55]. We removed SNP calls with a quality

below 20 and read depth below 20, and then generated vcf files using bcftools[55]. To

check TP53 status we merged the SNP calls with known SNPs from ClinVar[57] and kept

SNPs with a clinical significance (CLNSIG) of "Pathogenic"[58].

Data and code availability

607   All the processed data, scripts and results from CHISEL are available on GitHub at

608   https://github.com/kidcancerlab/sc-OsteoCNAs. Whole genome sequencing data for

609   pediatric relapse tumor samples used for analysis in this study were obtained from St.

610   Jude Cloud[38–40].

Acknowledgments

We thank our funding sources. This work was generously supported by NIH/NCI

K08CA201638 (R.D.R.), R01CA260178 (R.D.R.) and U24CA248453 (B.J.R.), St.

Baldrick's Foundation Scholar Award (R.D.R.), Hyundai Hope on Wheels Young

Investigator Award (R.D.R.), Cancer Free Kids Foundation (R.D.R.), Steps for Sarcoma

Foundation (R.D.R.), Sarcoma Foundation of America (R.D.R.), a Pelotonia Fellowship

(S.R.), a Nationwide Children's Director's Strategic Development Fund, and an NIH

CTSA Grant UL1TR002733. S.Z. is a Cancer Research UK Career Development Fellow

(Award Reference RCCCDF-Nov21\100005) and is also supported by Rosetrees Trust

(Grant Reference M917) and by a Cancer Research UK UCL Centre Non-Clinical

Training Award (CANTAC721\100022).

References

1.   Casali, P. G. *et al.* Bone sarcomas: ESMO-PaedCan-EURACAN Clinical Practice
Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* **29**, iv79–
iv95 (2018).

2.   Bridge, J. A. *et al.* Cytogenetic findings in 73 osteosarcoma specimens and a
review of the literature. *Cancer Genet Cytogenet* **95**, 74–87 (1997).

3.   Chen, X. *et al.* Recurrent somatic structural variations contribute to tumorigenesis
in pediatric osteosarcoma. *Cell Rep* **7**, 104–112 (2014).

4. Squire, J. A. *et al.* High-resolution mapping of amplifications and deletions in
pediatric osteosarcoma by use of CGH analysis of cDNA microarrays. *Genes*
Chromosomes Cancer **38**, 215–225 (2003).

5. Sayles, L. C. *et al.* Genome-informed targeted therapy for osteosarcoma. *Cancer*
Discov **9**, 46–63 (2019).

6. Stephens, P. J. *et al.* Massive genomic rearrangement acquired in a single
catastrophic event during cancer development. *Cell* **144**, 27–40 (2011).

7. Meyerson, M. & Pellman, D. Cancer genomes evolve by pulverizing single
chromosomes.   *Cell*   vol.   144   9–10   Preprint   at
https://doi.org/10.1016/j.cell.2010.12.025 (2011).

8. Li, Y. *et al.* Constitutional and somatic rearrangement of chromosome 21 in acute
lymphoblastic leukaemia. *Nature* **508**, 98–102 (2014).

9. Behjati, S. *et al.* Recurrent mutation of IGF signalling genes and distinct patterns
of genomic rearrangement in osteosarcoma. *Nat Commun* **8**, 1–8 (2017).

10. Perry, J. A. *et al.* Complementary genomic approaches highlight the PI3K/mTOR
pathway as a common vulnerability in osteosarcoma. *Proc Natl Acad Sci U S A*
111, E5564–E5573 (2014).

11. Zhao, Y. *et al.* Single-cell RNA sequencing reveals the impact of chromosomal
instability on glioblastoma cancer stem cells. *BMC Med Genomics* **12**, 79 (2019).

12. Bakker, B. *et al.* Single-cell sequencing reveals karyotype heterogeneity in murine
and human malignancies. (2016) doi:10.1186/s13059-016-0971-7.

13. Watkins, T. B. K. *et al.* Pervasive chromosomal instability and karyotype order in
tumour evolution. *Nature* **587**, 126–132 (2020).

14. Bach, D. H., Zhang, W. & Sood, A. K. Chromosomal instability in tumor initiation
and development. *Cancer Research* vol. 79 3995–4002 Preprint at
https://doi.org/10.1158/0008-5472.CAN-18-3235 (2019).

15. Fearon, E. R. & Vogelstein, B. A genetic model for colorectal tumorigenesis. *Cell*
61, 759–767 (1990).

16. Hoglund, M. *et al.* Multivariate analyses of genomic imbalances in solid tumors
reveal distinct and converging pathways of karyotypic evolution. *Genes*
Chromosomes Cancer **31**, 156–171 (2001).

17. Umbreit, N. T. *et al.* Mechanisms generating cancer genome complexity from a
single cell division error. *Science (1979)* **368**, (2020).

18. Gao, R. *et al.* Punctuated copy number evolution and clonal stasis in triple-
negative breast cancer. *Nat Genet* **48**, 1119–1130 (2016).

19. Zhou, Y. *et al.* Single-cell RNA landscape of intratumoral heterogeneity and
immunosuppressive microenvironment in advanced osteosarcoma. *Nat Commun*
11, 1–17 (2020).

20.   Liu, Y. *et al.* Single-Cell Transcriptomics Reveals the Complexity of the Tumor
669   Microenvironment of Treatment-Naive Osteosarcoma. *Front Oncol* **0**, 2818
670   (2021).

671   21.   Xu, H. *et al.* Genetic and clonal dissection of osteosarcoma progression and lung
672   metastasis. *Int J Cancer* **143**, 1134–1142 (2018).

673   22.   Negri, G. L. *et al.* Integrative genomic analysis of matched primary and metastatic
674   pediatric osteosarcoma. *J Pathol* **249**, 319–331 (2019).

675   23.   Wang, Y. & Navin, N. E. Advances and Applications of Single Cell Sequencing
676   Technologies. *Mol Cell* **58**, 598 (2015).

677   24.   Laks, E. *et al.* Clonal Decomposition and DNA Replication States Defined by
678   Scaled Single-Cell Genome Sequencing. *Cell* **179**, 1207-1221.e22 (2019).

679   25.   Tarabichi, M. *et al.* A practical guide to cancer subclonal reconstruction from DNA
680   sequencing. *Nature Methods 2021 18:2* **18**, 144–155 (2021).

681   26.   Zaccaria, S. & Raphael, B. J. Accurate quantification of copy-number aberrations
682   and whole-genome duplications in multi-sample tumor sequencing data. *Nature*
683   *Communications 2020 11:1* **11**, 1–13 (2020).

684   27.   Zaccaria, S. & Raphael, B. J. Characterizing allele- and haplotype-specific copy
685   numbers in single cells with CHISEL. *Nat Biotechnol* (2020) doi:10.1038/s41587-
686   020-0661-6.

687   28.   Andor, N. *et al.* Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines
688   reveals rules of in vitro evolution. *NAR Genom Bioinform* **2**, (2020).

689   29.   Houghton, P. J. *et al.* The pediatric preclinical testing program: Description of
690   models   and   early   testing   results.   *Pediatr   Blood   Cancer*   (2007)
691   doi:10.1002/pbc.21078.

692   30.   Woo, X. Y. *et al.* Conservation of copy number profiles during engraftment and
693   passaging of patient-derived cancer xenografts. *Nature Genetics 2021 53:1* **53**,
694   86–99 (2021).

695   31.   Martin, J. W., Squire, J. A. & Zielenska, M. The genetics of osteosarcoma.
696   *Sarcoma* **2012**, 11 (2012).

697   32.   Minussi, D. C. *et al.* Breast tumours maintain a reservoir of subclonal diversity
698   during expansion. *Nature* **592**, 302–308 (2021).

699   33.   Zack, T. I. *et al.* Pan-cancer patterns of somatic copy number alteration. *Nat*
700   *Genet* **45**, 1134–1140 (2013).

701   34.   López, S. *et al.* Interplay between whole-genome doubling and the accumulation
702   of deleterious alterations in cancer evolution. *Nat Genet* **52**, 283–293 (2020).

703   35.   Bielski, C. M. *et al.* Genome doubling shapes the evolution and prognosis of
704   advanced cancers. *Nat Genet* **50**, 1189–1195 (2018).

705   36.   Passerini, V. *et al.* The presence of extra chromosomes leads to genomic
706   instability. *Nat Commun* **7**, 1–12 (2016).

707   37.   Sheltzer, J. M. A transcriptional and metabolic signature of primary aneuploidy is
708   present in chromosomally unstable cancer cells and informs clinical prognosis.
709   *Cancer Res* **73**, 6401–6412 (2013).

710   38.   McLeod, C. *et al.* St. Jude Cloud: A Pediatric Cancer Genomic Data-Sharing
711   Ecosystem. *Cancer Discov* **11**, 1082–1099 (2021).

712   39.   Downing, J. R. *et al.* The Pediatric Cancer Genome Project. *Nat Genet* **44**, 619–
713   622 (2012).

714   40.   Stewart, E. *et al.* Orthotopic patient-derived xenografts of paediatric solid tumours.
715   *Nature* **549**, 96–100 (2017).

716   41.   Zaccaria, S. & Raphael, B. J. Accurate quantification of copy-number aberrations
717   and whole-genome duplications in multi-sample tumor sequencing data. *Nature
718   Communications 2020 11:1* **11**, 1–13 (2020).

719   42.   Wang, D. *et al.* Multiregion sequencing reveals the genetic heterogeneity and
720   evolutionary history of osteosarcoma and matched pulmonary metastases.
721   *Cancer Res* **79**, 7–20 (2019).

722   43.   Brady, S. W. *et al.* The Clonal Evolution of Metastatic Osteosarcoma as Shaped
723   by Cisplatin Treatment. *Mol Cancer Res* **17**, 895–906 (2019).

724   44.   Navin, N. *et al.* Tumour evolution inferred by single-cell sequencing. *Nature* **472**,
725   90–95 (2011).

726   45.   Wang, Y. *et al.* Clonal evolution in breast cancer revealed by single nucleus
727   genome sequencing. *Nature* **512**, 155–160 (2014).

728   46.   Navin, N. E. The first five years of single-cell cancer genomics and beyond.
729   *Genome   Research*   vol.   25   1499–1507   Preprint   at
730   https://doi.org/10.1101/gr.191098.115 (2015).

731   47.   Gawad, C., Koh, W. & Quake, S. R. Single-cell genome sequencing: current state
732   of the science. *Nature Reviews Genetics 2016 17:3* **17**, 175–188 (2016).

733   48.   Gröbner, S. N. *et al.* The landscape of genomic alterations across childhood
734   cancers. *Nature* **555**, 321–327 (2018).

735   49.   Ben-David, U. *et al.* Patient-derived xenografts undergo mouse-specific tumor
736   evolution. *Nature Genetics 2017 49:11* **49**, 1567–1575 (2017).

737   50.   Ma, X. *et al.* Pan-cancer genome and transcriptome analyses of 1,699 paediatric
738   leukaemias and solid tumours. *Nature* **555**, 371–376 (2018).

739   51.   Rajan, S. *et al.* Osteosarcoma tumors maintain intratumoral heterogeneity, even
740   while adapting to environmental pressures that drive clonal selection. *bioRxiv*
741   2020.11.03.367342 Preprint at https://doi.org/10.1101/2020.11.03.367342 (2020).

742   52.   Schwarz, R. F. *et al.* Phylogenetic Quantification of Intra-tumour Heterogeneity.
743   *PLoS Comput Biol* **10**, e1003535 (2014).

744   53.   El-Kebir, M. *et al.* Complexity and algorithms for copy-number evolution problems.
745   *Algorithms for Molecular Biology 2017 12:1* **12**, 1–11 (2017).

746   54.   ZaccariaSimone, El-KebirMohammed, W., K. & J., R. Phylogenetic Copy-Number
747   Factorization of Multiple Tumor Samples. *https://home.liebertpub.com/cmb* **25**,
748   689–708 (2018).

749   55.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4
750   (2021).

751   56.   Koboldt, D. C. *et al.* VarScan 2: Somatic mutation and copy number alteration
752   discovery in cancer by exome sequencing. *Genome Res* **22**, 568–576 (2012).

753   57.   Landrum, M. J. *et al.* ClinVar: improving access to variant interpretations and
754   supporting evidence. *Nucleic Acids Res* **46**, D1062–D1067 (2018).

755   58.   Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics
756   Analysis Version 11. *Mol Biol Evol* **38**, 3022–3027 (2021).

Figure 1

Figure 1. Extensive genomic complexity in ten expanded osteosarcoma patient tissue samples using single cell DNA sequencing. Allele-specific copy numbers (heatmap colors) are inferred by using the CHISEL algorithm27 from each of ten datasets including 300-2300 single cancer cells from osteosarcoma tumors. In each dataset, cancer cells are grouped into clones (colors in leftmost column) by CHISEL based on the inferred allele-specific copy numbers. Corrected allele-specific copy-numbers are correspondingly obtained by consensus. Note that cells classified as noisy by CHISEL have been excluded. '*' and '#' represent samples obtained from the same patient.

Figure 2. Osteosarcoma cancer cells exhibit extensive genetic alterations, especially deletions, but a relatively low level of heterogeneity. (A) Ploidy (y-axis) and fraction of aberrant genome (x-axis) of every cell (point) across the ten analyzed datasets (colors). The kernel density of the marginal distributions of each value is reported accordingly in every plot. (B) Fraction of genome affected by deletions (x-axis) vs. fraction of genome affected by amplifications (y-axis) of every cell (point) across the ten analyzed datasets (colors). (C) Fraction of aberrant genome (x-axis) and fraction of sub-clonal SCNAs (i.e. fraction of the genome with SCNAs different than the most common clone for the same region across all cells in the same dataset, y-axis) of every cell (point) across the ten analyzed datasets (colors).

Figure 3. Phylogenetic reconstruction of tumor evolution is consistent with longitudinal ordering of matched tumor samples and reveals conservation of SCNA profiles. (A) Allele-specific copy numbers (heatmap colors) across all autosomes (columns) have been inferred by CHISEL jointly across 4238 cells (rows) in 3 tumor samples from the same patient: 1 pre-treatment sample (NCH-OS-4 tibia) and two post-treatment samples (NCH-OS-7 tibia and NCH-OS-7 flank). CHISEL groups cells into 4 distinct clones (blue, green, red, and purple) characterized by different complements of SCNAs. (B) Phylogenetic tree describes the evolution in terms of SCNAs for the four identified tumor clones. The tree is rooted in normal diploid clone (white root) and is characterized by two unobserved ancestors (white internal nodes). Edges are labelled with the corresponding copy-number events that occurred and transformed the copy-number profile of the parent into the profile of the progeny. The four tumor clones (blue, green, red, and purple) are labelled according to the sample in which they were identified.

Figure 4

Figure 4. CNA correlation between osteosarcoma samples. Pearson R values denoting correlation of binned copy numbers between samples. Colors on x and y-axes indicate each sample's patient of origin and type as well as the clones defined from the correlation analysis. Red asterisks denote samples from patients with germline TP53 mutations. Note that SJOS003939_X1 is from the same patient as SJOS031478_* samples.

Figure 5

- Single initiation event
- Chromosomal stability after initiation
- CNV/SNP patterns consistent among cells
- Single clone

- Single initiation event
- Unstable chromosomal arrangement
- CNV/SNP patterns divergent between cells
- Shared phylogeny between all cells

- Multiple initiation events
- Chromosomal stability after initiation
- CNV/SNP patterns consistent per clone
- No shared phylogeny between clones

- Multiple initiation events
- Low-level chromosomal instability
- Variation within clones
- No shared phylogeny between clones

- Single, non-competitive initiation event
- Expansion of competitive sub-clones
- CNV/SNP patterns consistent per clone
- Shared phylogeny from distinct progenitor cell

Figure 5. Possible models for temporal SCNA stability. (A) After tumor initiation, if chromosomal instability is low, tumors will have identical SCNA patterns across all cells. (B) High tumor instability would result in tumors with highly heterogeneous SCNA patterns across cells which may not be apparent in bulk sequencing. (C) If there are multiple initiation events with low subsequent genomic instability SCNA patterns will be consistent across clones, which will be apparent by both bulk and single cell sequencing. (D) If there are multiple initiation events with ongoing genomic instability, clones derived from each will be similar but with highly variable SCNA patterns within a clone. This heterogeneity would be apparent by single cell sequencing but not bulk sequencing. (E) If a single initiation event is followed by an initial period of genomic instability, divergent clones could emerge. Patterns of stability within a clone would suggest that chromosomal stability is re-established prior to clonal expansion.