Caspase-mediated nuclear pore complex trimming

in cell differentiation and endoplasmic reticulum stress

Ukrae H. Cho[1, *] and Martin W. Hetzer[1, 2, *]

[1] Molecular and Cell Biology Laboratory, Salk Institute for Biological Studies, La Jolla, CA 92037, USA

[2] Institute of Science and Technology Austria (IST Austria), Klosterneuburg, Austria

*Correspondence should be addressed to
U.H.C (ucho@salk.edu) or M.W.H (martin.hetzer@ist.ac.at)

Abstract

During apoptosis, caspases degrade 8 out of ~30 nucleoporins to irreversibly demolish the

nuclear pore complex. However, for poorly understood reasons, caspases are also activated during cell

differentiation. Here, we show that sublethal activation of caspases during myogenesis results in the

transient proteolysis of four peripheral Nups and one transmembrane Nup. "Trimmed" NPCs become

nuclear export-defective, and we identified in an unbiased manner several classes of cytoplasmic,

plasma-membrane, and mitochondrial proteins that rapidly accumulate in the nucleus. NPC trimming by

non-apoptotic caspases was also observed in neurogenesis and endoplasmic reticulum stress. Our

results suggest that caspases can reversibly modulate nuclear transport activity, which allows them to

function as agents of cell differentiation and adaptation at sublethal levels.

INTRODUCTION

Caspases are a group of apoptotic proteases that target hundreds of proteins (Araya et al., 2021;

Mahrus et al., 2008; Shimbo et al., 2012). The activation of executioner caspases, especially that of

caspase-3, is equated with cell death. However, studies in the last two decades have demonstrated that

it is transiently activated at sublethal levels in differentiating cells (Burgon and Megeney, 2018; Connolly

et al., 2014; Ding et al., 2016; Tang et al., 2015). This phenomenon has been described in myogenic,

neuronal, osteogenic, and many other types of differentiation.

The nuclear pore complex (NPC), the sole gateway between the nucleoplasm and the cytoplasm,

is extensively targeted by caspase-3 (Ferrando-May et al., 2001; Kihlmark et al., 2001; Patre et al., 2006).

A total of 8 nucleoporins (Nups), the NPC subunits, are degraded by this cysteine protease: 2 out of the

8 Nups are on the cytoplasmic side of the NPC (Nup358 and Nup214), 2 form NPC scaffold (Nup96 and

Nup93), 3 reside on the nucleoplasmic side (Nup153, Tpr, and Nup50), and 1 is a transmembrane protein

(Pom121). Hence, all major NPC substructures – cytoplasmic filaments, central scaffold, nuclear basket,

and membrane-bound portion – are destroyed. Caspase-3-mediated NPC disintegration has been

studied extensively in apoptotic cells, but not in differentiating cells. It is unknown if/how the NPC is

modified by caspase-3 in differentiating cells. Are the same set of 8 Nups cleaved? What happens to

nuclear transport and passive permeability barrier? How does NPC proteolysis facilitate cell identity

transition?

We address these questions by examining caspase-mediated NPC proteolysis in C2C12 cells

undergoing myoblast-to-myotube transition where pulse-like, non-apoptotic caspase-3 activation has

been well-characterized (Bloemberg and Quadrilatero, 2014; Boonstra et al., 2018; Fernando et al., 2002;

M et al., 2020; Murray et al., 2008). Transient caspase-3 activation in differentiating myoblasts has been

shown by immunoblotting and fluorogenic probes. Moreover, genetic deletion or pharmacological

inhibition of caspase-3 significantly inhibits cell-cell fusion and expression of key muscle genes. Here, we

show that during myogenesis, caspase-3 molecularly "trims" the NPC in a reversible manner rather than

dismantling the entire structure as in apoptosis. This partial degradation of the NPC transiently impairs

nuclear export, but not nuclear import and passive permeability barrier, and accompanies nuclear

accumulation of focal adhesion proteins, tubulins, Annexin A's, mitochondrial enzymes, etc. Of note, NPC

trimming explains how FAK becomes transiently nuclear, which previously was shown to be required for

myogenin de-repression (Luo et al., 2009). NPC trimming is observed in differentiating primary myoblasts

and neural precursor cells as well, illustrating that it may be a general feature of cell differentiation. We

also observed NPC trimming in ER-stressed cells, suggesting that caspase-mediated modulation of

nuclear transport might represent not only a programmed step in differentiation but also a part of the

cellular stress response system.

RESULTS

5 Nups are completely degraded by caspase-3 during myogenesis

Caspase-3 can degrade 8 Nups (Nup358, Nup214, Nup153, Nup96, Nup93, Nup50, Tpr, and

Pom121) during staurosporine-, actinomycin D-, etoposide-, or TRAIL-induced apoptosis (Ferrando-May

et al., 2001; Patre et al., 2006). Recombinant caspase-3 reproduces *in vivo* Nup fragmentation pattern

when added to permeabilized cells, and its pharmacological inhibition by DEVD-CHO blocks Nup

degradation during apoptosis (Ferrando-May et al., 2001). Nups were identified as caspase-3 targets in

multiple substrate profiling experiments as well (https://wellslab.ucsf.edu/degrabase/). Rendering them

caspase-3-resistant is technically not feasible as most of them are cleaved at multiple sites – for example,

8+ for Nup358 and 4+ for Tpr (Patre et al., 2006). We observe 6+ fragments for Nup153 (see below).

However, human Pom121 has been made uncleavable by D531A mutation (Kihlmark et al., 2004).

We first asked whether caspase-3 proteolyzes Nups during myogenesis. As reported previously,

we detected the active form of caspase-3 as well as that of caspase-9 and -12 in differentiating C2C12

cells (**Figures 1A** and **S1A**) (Fernando et al., 2002; Gomez-Cavazos and Hetzer, 2015; Murray et al.,

2008; Nakanishi et al., 2005). Typically, caspase activity peaks at 16-24 hours after inducing myogenesis,

and is fully quenched by 32-60 hours. (See **Cell culture** in **MATERIALS AND METHODS**. As a reference,

peak caspase-3 activity level in differentiating C2C12 cells is comparable to that in C2C12 cells treated

with 1 μM staurosporine for 4 hours (Murray et al., 2008).) During this time window, 4 peripheral Nups,

Nup358, Nup214, Nup153, and Tpr, were reversibly and completely degraded in a caspase-dependent

manner (**Figure S1B**). Complete loss of full-length proteins demonstrates that this proteolytic event takes

place in the differentiating population, not just in 2-3% of the population that undergo apoptosis during

C2C12 myogenesis. (We washed plates 3 times with PBS before harvesting to minimize their contribution.

We also show the loss of Nup fragments in differentiating cells by immunofluorescence below.) Although

partially, Nup50 was proteolyzed as well (**Figure S1C**). Nup96 and Nup93, which form NPC scaffold and

are cleaved during apoptosis (Patre et al., 2006), remained intact. (For Nup93, a faster-running, second

band arises in a caspase-dependent manner (**Figure S1C**). The molecular nature of this band is unclear

at the moment.) This agrees with our previous report showing the persistence of the same Nup96 and

Nup93 copies through the course of myogenesis (Toyama et al., 2019).

For more comprehensive survey, we tested 8 additional scaffold Nups and 1 transmembrane Nup,

Pom121, which is cleaved by caspase-3 in apoptotic cells (**Figure 1B**). Only Pom121 exhibited caspase-

dependent proteolysis. In summary, caspase-3 degrade 4 peripheral Nups and 1 transmembrane Nup

during myoblast-to-myotube transition. Given that caspase-3-exposed NPCs will lack cytoplasmic

filaments (Nup358 and Nup214), nuclear basket (Nup153 and Tpr), and the C-terminal tail of Pom121

that extends into the cytoplasm, we termed this phenomenon "NPC trimming". Importantly, since the NPC

core is unaffected, functional NPCs can be quickly restored once caspases are quenched. RNA-seq

shows that the transcript levels of all but one Nups remain stable (**Figure S2**). Nup210 is the only exception whose expression level changes dramatically during myotube formation at both transcript and protein levels as we have previously reported (D'Angelo et al., 2012).

Calpains are another class of proteases that are activated in differentiating myoblasts (Louis et al., 2008), and are known to proteolyze Nups in multiple contexts (Bano et al., 2010; Sugiyama et al., 2017; Yamashita et al., 2017). We tested whether calpain inhibition prevents the digestion of Nups (**Figure S1B**). Interestingly, it delayed but did not prevent caspase-3/12 activation and peripheral Nup degradation. This suggests that they participate in caspase activation process but are not responsible for NPC trimming.

Given that central FG Nups, such as Nup98 and Nup62, are not targeted by caspases (Ferrando-May et al., 2001; Patre et al., 2006), we surmised that the nuclear passive permeability barrier would remain intact. Indeed, nuclear exclusion of ≥40-kDa dextrans demonstrates that the passive barrier is functional even when the NPC is partially disintegrated (**Figure S1D**).

Caspase-mediated NPC trimming impairs nuclear export

The loss of peripheral Nups suggested that the engagement of nuclear transport receptors, and thus nuclear transport, might be affected. The removal of Nup358, Nup214, Nup153, or Tpr has been shown to block nuclear export and cause nuclear accumulation of RNAs and NES (nuclear export signal)-containing proteins (Aksenova et al., 2020; Forler et al., 2004; Hutten and Kehlenbach, 2006; Ullman et al., 1999). We thus assessed mRNA export during the first three days of myogenic differentiation by combining nuclear isolation and oligo(dT) bead-based poly(A)$^+$ RNA purification. The nuclear-to-total mRNA ratio increases from 34% to 60% during differentiation, indicating an impairment in mRNA export (**Figures 1C, S3A,** and **S3B**). Of note, the distribution of housekeeping gene RNAs, 18S rRNA and *Gapdh* transcript, remains unchanged during this period as they have extended half-lives (**Figure S3C**) (Lee et al., 2010; Yi et al., 1999). This might explain how cells can continue to synthesize essential proteins and maintain homeostasis even when RNA export is temporarily inhibited.

We then examined Crm1-dependent nuclear export of proteins. The cytoplasmic-to-nuclear ratio

of NES$_{Rev}$-GFP, a widely used export probe, was significantly lower in myoblasts undergoing cell-cell

fusion than in myotubes (**Figures 1D, 1E,** and **S3D**). Similarly, the cytoplasmic-to-nuclear ratio of NES$_{PKI}$-

GFP significantly decreased when myoblasts started to produce "bubbling" blebs and became fusion-

competent (**Figures S3E-G**) (Lian et al., 2020). We also asked if the localization of α-tubulin is affected.

It is normally absent in the nucleus due to the presence of multiple NESs (Schwarzerova et al., 2019).

Strikingly, 1 day post-differentiation when NPCs are partially disintegrated, α-tubulin was detected in the

nuclear lysate, although at a relatively low level (**Figure 1F**). When NPC trimming was blocked by Q-VD-

OPh, a pan-caspase inhibitor, its nuclear translocation was suppressed (**Figure S4A**). We next checked

if focal adhesion proteins behave similarly since they (1) contain NESs as well and (2) are likely to be

released to the cytoplasm by calpains in differentiating C2C12 cells (Louis et al., 2008). Remarkably, all

four focal adhesion proteins that we examined (Hic-5, zyxin, paxillin, and FAK) became transiently

nuclear during myogenesis (**Figure 1G**). In the presence of Q-VD-OPh, their nuclear entrapment was

notably suppressed, albeit not fully (**Figure S4B**). The residual amount can be attributed to forced import

during myogenesis by their binding partners that make use of them as transcription cofactors in the

nucleus. We indeed detect interaction between FAK and MBD2 (Luo et al., 2009) and Hic-5 and androgen

receptor (Alpha et al., 2020) by proximity ligation assay (**Figure S5**).

In summary, differentiation-associated caspase activity impairs nuclear export and leads to the

nuclear retention of NES-containing proteins. (Nuclear import is discussed below.) Given the rapid influx

of NES-harboring proteins, we hypothesized that leptomycin B, a Crm1 inhibitor, could support

myogenesis. Indeed, when differentiation medium is supplemented with a partial dose of leptomycin B,

myogenin expression is enhanced (**Figure S4C**). This agrees with the reported ~2-fold myogenin

enhancement by 6 hours of leptomycin B pre-treatment in growth medium (Bustos et al., 2015). In

contrast, a full dose of Q-VD-OPh in differentiation medium significantly suppresses myogenin

upregulation (**Figure S4D**). Q-VD-OPh significantly reduces cell-cell fusion and myosin heavy chain

expression as well (Murray et al., 2008).

Tpr and Pom121 C-terminal fragments dissociate from the NPC

Since Nups are large proteins that are held in place via multiple interactions, we asked whether the entire proteins are released from the NPC after proteolysis. Using four different antibodies (**Figure S6A**), we first analyzed whether various domains of Nup153 and Tpr are retained in late-stage apoptotic nuclei where caspase levels are significantly higher than in differentiating cells and chromatins are fully condensed. Immunostaining revealed that only the C-terminal epitope of Tpr dissociates from the NPC while proteolyzed Nup153 and Tpr N-terminus remain bound **(Figures 2A** and **B)**. Their persistent association with the NPC is reminiscent of that of enterovirus 2A or rhinovirus 3C protease-cleaved Nup98 (Park et al., 2015; Walker et al., 2013).

Differentiating myoblasts exhibited a proteolytic phenotype that is distinct from that of apoptotic cells. Quantitative analysis suggests that Tpr C-terminal fragment dissociates partially (**Figures 2C** and **D**). Only in 5-10% of the cells were the fragment completely dissociated as in apoptotic cells (dotted boxes in **Figure 2C**). "TprC⁻" population can be clearly distinguished from dying cells by nuclear morphology (**Figure S6B**). Moreover, while the nucleoplasmic pool of Crm1 is lost and it becomes confined to the nuclear periphery in apoptotic cells (**Figure S6C**), Crm1 localization did not change in differentiating cells, even in TprC⁻ cells (**Figure 2C**). This might explain why NPC trimming results in a subdued nuclear export defect compared to a full dose of leptomycin B; NES-containing focal adhesion proteins accumulate in the nucleus to a limited extent during NPC trimming (**Figure 1G**) but become predominantly nuclear upon leptomycin B treatment (Nix et al., 2001; Ossovskaya et al., 2008).

To further validate (in)complete retention of Tpr C-terminal tail and Nup153 fragments, respectively, we isolated the nuclei of differentiating C2C12 cells and fractionated them into "nuclear insoluble" (nuclear envelope, NPC, and chromatin-bound) and "nuclear soluble" (**Figure 2E**; fractionation steps detailed below). Nup153 fragments were found exclusively in the former. In contrast, the C-terminal piece of Tpr, which is expected to be ~37 kDa (**Figure S6A**) (Ferrando-May et al., 2001), was detected in both subfractions. This agrees with the partial-dissociation-upon-cleavage model.

We also analyzed Pom121 fragmentation. Pom121 has a short transmembrane and a long pore-side domain (aminoacid 35-55 and 56-1249 respectively in human; **Figure S6A**). The latter is caspase-

cleaned and is undetectable by immunostaining in apoptotic cells (Kihlmark et al., 2001; Kihlmark et al.,

2004). Although Pom121 is proteolyzed to completion at 24 hours post-differentiation (**Figures 1B**),

immunostaining shows that its pore side domain does not fully dissociate from the NPC (**Figures 2F** and

G). As in the case of Tpr, Pom121 C-terminus immunoreactivity does decrease on average but only a

subset of nuclei exhibits its full loss. To conclude, there is some heterogeneity in the behavior of caspase-

generated Nup fragments while NES-GFP reporter demonstrates that most, if not all, NPCs become

export-defective. Lastly, we confirmed that trimmed NPCs do not cluster like they do in late-stage

apoptosis via maximum intensity Z projection (**Figure S6D**) (Kihlmark et al., 2001), and that the

localization of importins is unaffected by NPC trimming (**Figure S7**).

NPC trimming coincides with a rapid change in the nuclear proteome

Next, we employed quantitative mass spectrometry to study how NPC trimming affects the

nuclear proteome. For better coverage of low-abundance proteins, we fractionated nuclei into "nuclear

insoluble" and "nuclear soluble" by hypotonic lysis (**Figure 3A**). The former contains proteins strongly

bound to the NPC, nuclear envelope, and chromatin, whereas in the latter are ones solubilized in the

nucleoplasm (**Figure 3B**). Using tandem mass tag labeling, we determined how the levels of ~2800

proteins in the two subfractions change during caspase activation and NPC trimming, i.e., before and

after 24 hours of myogenic differentiation. We overlaid RNA-seq data on top of mass spectrometry result

and generated a color-mapped volcano plot for the "nuclear soluble" fraction (**Figures 3C** and **D** and

Supplemental Spreadsheet 1). Our multi-omic analysis well matches previous candidate-based studies.

As shown in **Figure 3C**, alpha-actinin 2 (Actn2) is known to be transcriptionally upregulated and removed

from the nucleus during myogenesis (Lin et al., 2010).

In the "nuclear soluble" fraction, there were 236 proteins that increased >4-fold in the first 24

hours. We grouped them by function/localization (**Figure 4A**). All focal adhesion proteins but tensins

became significantly more nuclear as expected. Several α/β-tubulin isoforms and microtubule regulators

displayed similar behavior. Interestingly, Tppp3 (Tubulin polymerization promoting protein family member

3), which has been reported to localize to nucleolar fibrillar center (the Human Protein Atlas; HPA047629),

was the top hit in this group. Another noticeable group was Annexin A's. Among them is Annexin A2, which is an NES-containing protein that becomes more nuclear in genotoxic condition and mitigates DNA damage. The nuclear level of S100a10, an Annexin A2 interactor, increased sharply as well (**Figure 3D**). We also find that several mitochondrial enzymes, but not ATP synthase or TIM/TOM complex subunits barring Tomm34, accumulate in the nucleus. A subset of the Krebs cycle enzymes has been reported to enter the nucleus during embryogenesis (Nagaraj et al., 2017), and we find the same in myogenic differentiation. Lastly, the nuclear levels of most E3 ligases and proteasome subunits remain relatively stable, but there were a few exceptions such as Itch, Ufl1, Psme1 and Psmb3.

In addition to caspase-mediated NPC trimming, other mechanisms could increase the nuclear level of a protein during myogenesis. Our color-mapped volcano plot shows that most transcriptionally upregulated proteins (red data points in **Figures 3C** and **D**) become more abundant in the nucleus, demonstrating that enhanced synthesis is another major factor. Enhanced nuclear import and/or stability can be involved as well. To determine the contribution of NPC trimming to a given translocation event, one would need to generate 5 caspase-resistant Nups, which for technical reasons is impractical. Our alternative approach was to pharmacologically inhibit caspases during differentiation and test if the influx/retention of non-nuclear proteins is mitigated. Out of 700 proteins that show >2-fold increase in the nuclear level during the first 24 hours (area shaded in red in **Figure 4B**), all but 4 display less increase when caspases are inhibited. This is in line with our view that NPC trimming plays a key role in shaping the nuclear proteome (**Figure 4C**). In sum, our data (1) illustrates that the nuclear proteome changes dramatically and rapidly in response to the myogenic signal, (2) reveals the importance of protein translocation in cell differentiation, and (3) provides a fresh list of proteins that have been overlooked in the context of myotube formation.

NPC trimming occurs after survivin downregulation and before myogenin upregulation

We find that caspase-mediated NPC trimming (**Figure 5A**) coincides with a rapid nuclear proteome change. We next wanted to pinpoint the timing of this event. Cell cycle exit is one of the earliest events in myogenesis. Survivin is a chromosomal passenger complex subunit that is highly expressed in

G2/M phase but rapidly downregulated in early G1 (Li et al., 1998). In differentiating C2C12 cells, survivin downregulation precedes caspase-3 activation, indicating that the latter occurs after the very last cell division and as cells enter G0 (**Figure 5B**). We then determined the timing of caspase-3 activation relative to the upregulation of myogenin, which marks the point-of-no-return. Both immunoblotting and immunstaining show that active caspase-3 arises before myogenin (**Figures 5B** and **S8A**). Interestingly, cells can remain NPC-trimmed for a brief period after myogenin upregulation. We not only identified TprC⁻ Myog⁻ but also TprC⁻ Myog⁺ cells (**Figure S8A**). The presence of the latter signifies that nuclear import is functional even when NPCs are trimmed. The same transitory cell states appear in differentiating primary myoblasts (**Figure S8B**), demonstrating that this is not a C2C12-specific phenomenon. Caspase-3 is quenched by the time myosin heavy chain is expressed, i.e., when cell-cell fusion occurs (**Figure S8C**). When myogenesis is accelerated by GFP-myogenin overexpression, caspase-3 is quenched a day earlier compared to the control condition (**Figure S8C**), signifying that a caspase-quenching pathway exists and is activated once myoblasts pass a certain differentiation stage. **Figure 5C** summarizes the order of mentioned events.

We also checked whether caspase-3 activation, and hence NPC trimming, is affected by key signaling pathways involved in myogenesis: ERK1/2, calcineurin/NFAT, and Notch. ERK1/2 inhibition stimulates myogenesis by promoting cell-cycle exit as well as cell-cell fusion (Eigler et al., 2021). Consistently, we find that SCH772984, an ERK1/2 inhibitor, accelerates survivin downregulation, caspase-3 activation, Nup153 degradation, and myogenin induction (**Figure S8D**). When the nuclear translocation of NFAT was blocked by FK506, a calcineurin inhibitor, cells failed to upregulate myogenin as previously reported (Armand et al., 2008) but still activated caspase-3 (**Figure S8E**). This result illustrates that the calcineurin/NFAT pathway is orthogonal to the caspase-3 activation pathway. Lastly, DAPT, a γ-secretase inhibitor that blocks Notch intracellular domain release, neither affected myogenin upregulation nor caspase-3 activation (**Figure S8F**).

NPC trimming occurs during neurogenesis and ER stress

We next asked whether caspase-mediated NPC trimming is an event that is specific to myogenesis, which results in huge, multinucleated cells, or represents a general phenomenon associated with cell differentiation. To test this, neural precursor cells were differentiated into post-mitotic neurons over four weeks, and caspase-related events were analyzed (**Figure 5D**). As in myogenic differentiation, survivin is highly expressed in precursor cells but undetectable in mature neurons, and the activation of caspase-3 and the proteolysis of Nup153 and PARP occurred concomitantly. This suggests that caspases play similar roles during neuronal differentiation, albeit over a longer duration.

Our results so far suggest that NPC trimming and resultant nuclear proteome change are parts of a well-orchestrated sequence of events during early stages of cell differentiation. Interestingly, however, the calpain/caspase activation cascade in differentiating myoblasts is reminiscent of that in unfolded protein response. We thus asked whether ER stress alone can trigger partial NPC disintegration. C2C12 myotubes and reserve cells (mononucleated population) were treated with 1 μg/mL tunicamycin (**Figure 5E**). Active caspase-3 arose in both cell types at comparable levels; however, in myotubes, Nup153 and Tpr were fully proteolyzed by 6 hours. The discrepancy can be attributed to XIAP. XIAP is the major caspase-inhibiting protein, and its level is significantly lower in myotubes at 0 hour. (XIAP in myotubes migrates faster than that in reserve cells during gel electrophoresis, possibly due to Ser87 dephosphorylation, which stabilizes XIAP (Kato et al., 2011).) ER stress-induced NPC trimming exhibits multiple similarities to that during myogenesis: (1) NES$_{Rev}$-GFP becomes more nuclear (**Figure S8G**), (2) 5-10% of myotubes become TprC$^-$ while retaining Nup153 epitopes (**Figure S8H**), and (3) Nup93 remains unaffected, again illustrating that scaffold Nups are better protected from caspases (**Figure 5E**). Additionally, we tested whether low-level but sustained ER stress induces the same phenotype since tunicamycin treatment at the full dose may not be physiologically relevant. We maintained differentiated C2C12 cells at low doses of tunicamycin (<50 ng/mL) for a week (**Figure 5F**). As in differentiating myoblasts and acutely ER stressed-myotubes, the breakdown of Nup93 was minimal, whereas 4 peripheral Nups were noticeably degraded. In short, chronic ER stress forces the NPC to remain in the trimmed form for a prolonged period before apoptosis is triggered.

DISCUSSION

During apoptosis, the NPC is completely demolished by fully activated caspase-3. In

differentiating cells, caspase-3 activation is sublethal, and it selectively targets NPC projections, i.e.,

cytoplasmic filaments, nuclear basket, and Pom121 pore-side domain (**Figure 6**). Caspase-mediated

NPC trimming is a reversible process that transiently impairs nuclear export. This phenotype is somewhat

reminiscent of that of cells exposed to picornavirus 2A protease, where Nup98 is selectively degraded

from the NPC and mRNA/protein export is inhibited while nuclear import and passive permeability barrier

remain mostly functional (Serganov et al., 2022).

NPC trimming coincides with the nuclear influx/retention of non-nuclear proteins. Why do

differentiating cells need non-nuclear proteins in the nucleus? While it will require multiple follow-up

studies to answer this question, NPC trimming provides an explanation to one key event in myogenesis

whose most upstream trigger remained unknown. The nuclear level of FAK, a focal adhesion protein with

both nuclear export and localization signals, was previously shown to surge during myogenesis to remove

MBD2 – the main component of the gene-repressing NuRD (nucleosome remodeling and deacetylation)

complex – from methylated CpGs within the *Myog* promoter (**Figure S9A**) (Luo et al., 2009). In other

words, nuclear FAK de-represses *Myog*. Nevertheless, how FAK rapidly translocates into the nucleus in

differentiation myoblasts has remained unclear. We here show that caspase-mediated NPC trimming

impairs its export from the nucleus during myogenic differentiation. In line with this model, we find that

the loss of MBD2-genome binding occurs within 24 hours as FAK becomes nuclear and precedes the

degradation of MBD2 protein itself during myogenesis (**Figures S9B and C**). Of note, in mechanically

stressed cells, it is well-established that focal adhesion proteins quickly translocate to the nucleus to

function as transcription co-factors and induce rapid gene expression changes necessary for stress

resolution (Hervy et al., 2006). It is possible that this modus operandi has been adapted for cell

differentiation.

In addition, we find in the nucleus diverse classes of proteins that have been linked to DNA

damage response, such as Annexin A2, E3 ligases (Itch and Ufl1), and proteasome subunits (Psmb3)

(Chang et al., 2019; Jacquemont and Taniguchi, 2007; Madureira et al., 2012; Qin et al., 2019). This is

in line with the fact that DNA damage/repair is required for myogenesis (Al-Khalaf et al., 2016). Multiple

mitochondrial proteins enter the nucleus as well, hinting that nuclear-mitochondrial communication plays

a role myogenic differentiation (Soledad et al., 2019).

NPC trimming might also be an NPC-genome interaction reset mechanism. It is well established

that NPC-genome interaction (1) is cell type-specific, (2) is reconfigured during differentiation, and (3)

regulates cell identity genes (Cho and Hetzer, 2020; Ibarra et al., 2016; Jacinto et al., 2015; Labade et

al., 2021; Liang et al., 2013). Transient proteolysis of the nuclear basket can be an elegant reinitialization

mechanism. This is an interesting hypothesis to test in the future.

In the context of acute ER stress, NPC trimming might be a stress-comping mechanism that

curtails mRNA export and reduces the protein synthesis/folding load. However, if ER stress becomes

chronic and caspases linger for a prolonged period, it can jeopardize cellular homeostasis as nuclear

levels of NES-containing proteins will remain abnormally high. Our quantitative, subcellular fraction mass

spectrometry workflow (**Figure 3A**) can be employed to examine the nuclear proteome change during

chronic ER stress.

The present study establishes caspase-mediated NPC trimming as a key event in cell

differentiation and stress response. It also points to the idea that exhaustive NPC disintegration during

apoptosis may simply be an extreme version of this phenomenon. In fact, some consider apoptosis as a

radical form of caspase-mediated cellular transformation (Burgon and Megeney, 2018; Connolly et al.,

2014), and our results suggest that time is ripe to reassess the death-centric view of caspases.

MATERIALS AND METHODS

Data and code availability. RNA-sequencing data have been uploaded to the Gene Expression Omnibus (GEO) database (NCBI) under accession number GSE183521.

Cell lines. C2C12 cells were obtained from ATCC; HEK293T cells, from the Salk Stem Cell Core Facility; H9 embryonic stem cells, from the Gage lab (the Salk Institute); and mouse primary myoblasts, from the Alessandra Sacco lab (Sanford Burnham Prebys).

Immunoblot techniques. Cultured cells were washed with phosphate-buffered saline (PBS) and harvested by trypsinization. Cell pellet was washed with PBS, and lysed in RIPA buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% Triton X100, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate) containing protease and phosphatase inhibitors (Pierce Protease Inhibitor Mini Tablet, EDTA-free, Thermo Scientific; PhosSTOP, Roche) for 45 minutes at 4ºC. Insoluble material was pelleted at 16,000 x g at 4ºC for 20 minutes. 15-30 μg of total protein per sample were added 6x Laemmli sample buffer, boiled for 4 minutes, and loaded on a Tris or Bis-tris gel for electrophoresis. Proteins were transferred to a nitrocellulose membrane and stained with Ponceau S solution to confirm equal protein loading and successful transfer. After washing with Tris-buffered saline containing 0.05% (w/v) Tween 20 (TBST) several times, the membrane was blocked with 5% non-fat milk power in TBST for an hour at room temperature, and subsequently immunoblotted at 4ºC overnight with primary antibodies listed in **Supplemental Methods**. Chemiluminescent detection was conducted using either SuperSignal West Pico or Femto kits (Thermo Scientific) after 45-minute incubation with secondary antibodies listed in **Supplemental Methods** at room temperature. Western blot images were obtained using KwikQuant Imager (Kindle Biosciences) or Odyssey CLx (LI-COR).

Cloning. In-fusion cloning was performed using the In-Fusion HD EcoDry Cloning Plus kit (Takara), and standard cut-and-paste cloning using T7 DNA ligase (New England Biolabs). NES$_{Rev}$: LPPLERLTL. NES$_{PKI}$: LALKLAGLDI.

Cell culture. C2C12 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 20% fetal bovine serum and penicillin-streptomycin. For myogenic differentiation, they were grown to confluency, washed with PBS twice, and added DMEM with 2% horse serum and the same antibiotics. C2C12 differentiation medium was replenished every 24 or 48 hours. After 120 hours, mature myotubes and reserve cells were obtained. Myotubes were harvested with minimal contamination of reserve cells by mild trypsinization (1:3 or 1:4 dilution of trypsin in PBS). We want to note that the exact timing and duration of caspase activation, and thus NPC trimming, varies depending on cell confluency, culture substrate, growth area-to-medium volume ratio, passage number, etc. Compare **Figures 1A** and **S1B**. Typically, caspase activity peaks at 16-24 hours after inducing myogenesis, and is fully quenched by 32-60 hours in differentiating C2C12 cells. H9 embryonic stem cells were differentiated to neural precursor cells using a previously published method (Marchetto et al., 2017). Purified neural precursor cells were cultured in neurogenic conditions (DMEM/F12 based medium with 1x N2, 1x B27, 20ng/mL GDNF, 20ng/mL BDNF, 1mM cAMP, and 200nM ascorbic acid) for 4 weeks to generate mature post-mitotic neurons.

C2C12 cell fractionation. Cells were harvested by trypsinization, washed with PBS, and chilled on ice. Cells were then lysed in ice-cold 0.1% NP40 in PBS, and rotated at 4ºC for 10 minutes. Nuclei were pelleted by centrifugation at 500 x g at 4ºC for 4 minutes, and the supernatant (cytoplasmic fraction) was transferred to a fresh tube and stored at -80ºC until use. Nuclei were again resuspended in ice-cold 0.1% NP40 in PBS, rotated at 4ºC for 5 minutes, and pelleted at 500 x g at 4ºC for 4 minutes. The supernatant was discarded, and the pellet (nuclear fraction) was stored at -80ºC until use or further fractionated into "nuclear soluble" and "nuclear insoluble". To obtain the former, nuclei were added 3 mM EDTA, 0.2 mM EGTA, and 0.2 mM DTT, and rotated at 4ºC for 45 minutes. Hypotonically lysed nuclei were centrifuged

at 1700 x g at 4ºC for 5 minutes, and the supernatant was transferred to a fresh tube ("nuclear soluble"). Lysed nuclei were washed once with the same buffer, and then further extracted using RIPA buffer at 4ºC for 45 minutes. DNA-wrapped histones were pelleted at 16,000 x g at 4ºC for 20 minutes, and the supernatant was transferred to a fresh tube ("nuclear insoluble").

Lentivirus packaging, infection, and selection. Third-generation lentiviral protocol was followed to produced virus in HEK293T cells. C2C12 cells were infected with viral supernatant at 30-40% confluency in the presence of 6 μg/mL polybrene for 24 hours, and selected 24 hours after infection with 1 mg/mL puromycin.

RNA fluorescence *in situ* hybridization (FISH). C2C12 cells were grown on a No. 1.5 coverslip placed in a 12-well cell culture plate. After removing media, cells were washed with PBS, and fixed in 3.7% paraformaldehyde in PBS for 10 minutes at room temperature. Fixed cells were washed with PBS twice and permeabilized in 70% (vol/vol) ethanol at 4ºC for at least a day. We then followed Stellaris RNA FISH protocol for adherent cells (https://www.biosearchtech.com/support/resources/stellaris-protocols) to fluorescently visualize RNAs. Fluorescent images were obtained using a Leica SP8 confocal microscope equipped with a 63x oil-immersion objective. Images were cropped and pseudocolored using FIJI. See **Supplemental Methods** for RNA FISH probes used in this study.

Immunofluorescence. C2C12 cells and mouse primary myoblasts were grown on a chambered cell culture slide (Ibidi). For the latter, slides were coated with Matrigel prior to seeding. Cells were fixed in PBS containing 2% paraformaldehyde for 10 minutes at room temperature, and washed with PBS 3 times. Fixed cells were permeabilized and blocked in immunofluorescence buffer (PBS containing 0.1% Triton-X, 0.02% sodium dodecyl sulfate, and 10 mg/mL bovine serum albumin) for 30 minutes. The cells were then incubated with primary antibodies diluted in immunofluorescence buffer at room temperature for 2 hours, washed 3 times with immunofluorescence buffer, and incubated with the appropriate secondary antibodies diluted in immunofluorescence buffer at room temperature for 45 minutes. Finally, the cells

were washed with immunofluorescence buffer 3 times, and added VECTASHIELD antifade mounting medium with DAPI (Vector Laboratories). Fluorescent images were obtained using a Leica SP8 confocal microscope equipped with a 63x oil-immersion objective. Images were cropped and pseudocolored using FIJI. See **Supplemental Methods** for antibodies used for immunofluorescence.

Poly(A)$^+$ RNA quantification. Poly(A)$^+$ RNA was isolated from nuclei and whole cells (see C2C12 cell fractionation above) using the Magnetic mRNA Isolation Kit (New England Biolabs). Two rounds of binding, washing, and elution were performed. Eluted RNA was quantified via Qubit RNA HS Assay (Invitrogen).

Proximity ligation assay. Proximity ligation assay was performed using a Duolink *in situ* Red Starter Kit Mouse/Rabbit (Sigma). Following antibodies were used: FAK (BD Biosciences 610087), MBD2 (Sigma M7318), Hic-5 (BD Biosciences 611164), androgen receptor (Novus Biologicals NBP2-67497).

Dextran exclusion assay. 20, 40, and 65-85 kDa dextrans (Sigma, FD20S/FD40S/T1162) were dissolved in "Transport Buffer (TB)" (20 mM HEPES (pH 7.3), 110 mM potassium acetate, 5 mM sodium acetate, 1 mM EGTA, and 2 mM DTT) at 10 mg/ml. Degradation products were removed using 10 and 30 kDa-cutoff size exclusion columns (Millipore UFC501096/UFC503096). Dextran stock solutions were diluted in TB 1:50 to 1:200 prior to use. C2C12 cells seeded in a chambered cell culture slides were incubated with TB supplemented with 20 μg/ml digitonin for 5 min at room temperature, washed with TB (without digitonin) three times, and added dextrans. After 7.5 min at 37ºC, images were acquired using a Leica SP8 confocal microscope equipped with a 63x oil-immersion objective. Images were cropped and pseudocolored using FIJI.

mRNA-sequencing. 1-2 million C2C12 cells were lysed in 1 ml TRIzol. 0.4 ml chloroform was added and vigorously shaken for RNA extraction. The aqueous phase was transferred to a fresh tube, and one volume of 70% ethanol was added dropwise while vortexing at the lowest speed at room temperature.

The mixture was purified using RNeasy Mini kit (Qiagen) to yield several hundred ng/μL total RNA. mRNA-sequencing was performed in paired-end 150 base pair mode on the Illumina NovaSeq 6000 platform. Paired-end fragments were mapped to mm10, filtered, and assembled into transcripts using HISAT2 (Kim et al., 2019), SAMtools (Li et al., 2009), and StringTie (Pertea et al., 2015). Differential expression was evaluated using DESeq2 (Love et al., 2014), and further analyzed with in-house python scripts.

Mass spectrometry. Samples were precipitated by methanol/chloroform and redissolved in 8 M urea/100 mM TEAB, pH 8.5. Proteins were reduced with 5 mM tris(2-carboxyethyl)phosphine hydrochloride (TCEP, Sigma-Aldrich) and alkylated with 10 mM chloroacetamide (Sigma-Aldrich). Proteins were digested overnight at 37ºC in 2 M urea/100 mM TEAB, pH 8.5, with trypsin (Promega). The digested peptides were labeled with 10-plex TMT (Thermo product 90309, lot VI307195B), pooled samples were fractionated by basic reversed phase (Thermo 84868).

The TMT labeled lysate samples were analyzed on a Orbitrap Eclipse mass spectrometer (Thermo). Samples were injected directly onto a 25 cm, 100 μm ID column packed with BEH 1.7 μm C18 resin (Waters). Samples were separated at a flow rate of 300 nL/min on an EasynLC 1200 (Thermo). Buffer A and B were 0.1% formic acid in water and 90% acetonitrile, respectively. A gradient of 1–25% B over 100 min, an increase to 40% B over 20 min, an increase to 100% B over 10 min and held at 100% B for a 10 min was used for a 140 min total run time.

Peptides were eluted directly from the tip of the column and nanosprayed directly into the mass spectrometer by application of 2.5 kV voltage at the back of the column. The Eclipse was operated in a data dependent mode. Full MS1 scans were collected in the Orbitrap at 120k resolution. The cycle time was set to 3 sec, and within this 3 sec the most abundant ions per scan were selected for CID MS/MS in the ion trap. MS3 analysis with multinotch isolation (SPS3) was utilized for detection of TMT reporter ions at 60k resolution (McAlister et al., 2014). Monoisotopic precursor selection was enabled and dynamic exclusion was used with exclusion duration of 60 sec.

Protein and peptide identification were done with Integrated Proteomics Pipeline – IP2 (Integrated Proteomics Applications). Tandem mass spectra were extracted from raw files using RawConverter (He et al., 2015) and searched with ProLuCID (Xu et al., 2015) against Uniprot mouse database. The search space included all fully-tryptic and half-tryptic peptide candidates. Carbamidomethylation on cysteine and TMT on lysine and peptide N-term were considered as static modifications. Data was searched with 50 ppm precursor ion tolerance and 600 ppm fragment ion tolerance. Identified proteins were filtered to using DTASelect (Tabb et al., 2002) and utilizing a target-decoy database search strategy to control the false discovery rate to 1% at the protein level (Peng et al., 2003). Quantitative analysis of TMT was done with Census (Park et al., 2014) filtering reporter ions with 20 ppm mass tolerance and 0.6 isobaric purity filter.

Quantification and statistical analysis. Statistical analyses for **Figures 1E**, **2D**, **2G**, **S3G,** and **S8G** were performed using GraphPad Prism.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

ACKNOWLEDGMENTS

We thank the members of the Hetzer laboratory, Tony Hunter (Salk), Lorenzo Puri (Sanford Burnham Prebys), and Jongmin Kim (Massachusetts General Hospital) for the critical reading of the manuscript; Kenneth Diffenderfer and Aimee Pankonin (Stem Cell Core at the Salk Institute) for help with neurogenesis; Carol Marchetto and Fred Gage (Salk) for providing H9 embryonic stem cells; Lorenzo Puri, Alexandra Sacco, and Luca Caputo (Sanford Burnham Prebys) for helpful discussions and sharing mouse primary myoblasts. This work was supported by a Glenn Foundation for Medical Research Postdoctoral Fellowship in Aging Research (U.H.C.), the NOMIS foundation (M.W.H), and the National Institutes of Health (R01 NS096786 to M.W.H. and K01 AR080828 to U.H.C.). This work was also

supported by the Mass Spectrometry Core of the Salk Institute with funding from NIH-NCI CCSG: P30

014195 and the Helmsley Center for Genomic Medicine. We thank Jolene Diedrich and Antonio Pinto for

technical support.

AUTHOR CONTRIBUTIONS

REFERENCES

Aksenova, V., Smith, A., Lee, H., Bhat, P., Esnault, C., Chen, S., Iben, J., Kaufhold, R., Yau, K.C., Echeverria, C.*, et al.* (2020). Nucleoporin TPR is an integral component of the TREX-2 mRNA export pathway. Nat Commun *11*, 4577.

Al-Khalaf, M.H., Blake, L.E., Larsen, B.D., Bell, R.A., Brunette, S., Parks, R.J., Rudnicki, M.A., McKinnon, P.J., Jeffrey Dilworth, F., and Megeney, L.A. (2016). Temporal activation of XRCC1-mediated DNA repair is essential for muscle differentiation. Cell Discov *2*, 15041.

Alpha, K.M., Xu, W., and Turner, C.E. (2020). Paxillin family of focal adhesion adaptor proteins and regulation of cancer cell invasion. Int Rev Cell Mol Biol *355*, 1-52.

Araya, L.E., Soni, I.V., Hardy, J.A., and Julien, O. (2021). Deorphanizing Caspase-3 and Caspase-9 Substrates In and Out of Apoptosis with Deep Substrate Profiling. ACS Chem Biol *16*, 2280-2296.

Armand, A.S., Bourajjaj, M., Martinez-Martinez, S., el Azzouzi, H., da Costa Martins, P.A., Hatzis, P., Seidler, T., Redondo, J.M., and De Windt, L.J. (2008). Cooperative synergy between NFAT and MyoD regulates myogenin expression and myogenesis. J Biol Chem *283*, 29004-29010.

Bano, D., Dinsdale, D., Cabrera-Socorro, A., Maida, S., Lambacher, N., McColl, B., Ferrando-May, E., Hengartner, M.O., and Nicotera, P. (2010). Alteration of the nuclear pore complex in Ca(2+)-mediated cell death. Cell Death Differ *17*, 119-133.

Bloemberg, D., and Quadrilatero, J. (2014). Mitochondrial pro-apoptotic indices do not precede the transient caspase activation associated with myogenesis. Biochim Biophys Acta *1843*, 2926-2936.

Boonstra, K., Bloemberg, D., and Quadrilatero, J. (2018). Caspase-2 is required for skeletal muscle differentiation and myogenesis. Biochim Biophys Acta Mol Cell Res *1865*, 95-104.

Burgon, P.G., and Megeney, L.A. (2018). Caspase signaling, a conserved inductive cue for metazoan cell differentiation. Semin Cell Dev Biol *82*, 96-104.

Bustos, F., de la Vega, E., Cabezas, F., Thompson, J., Cornelison, D.D., Olwin, B.B., Yates, J.R., 3rd, and Olguin, H.C. (2015). NEDD4 regulates Pax7 levels promoting activation of the differentiation program in skeletal muscle precursors. Stem Cells *33*, 3138-3151.

Chang, L., Shen, L., Zhou, H., Gao, J., Pan, H., Zheng, L., Armstrong, B., Peng, Y., Peng, G., Zhou, B.P.*, et al.* (2019). ITCH nuclear translocation and H1.2 polyubiquitination negatively regulate the DNA damage response. Nucleic Acids Res *47*, 824-842.

Cho, U.H., and Hetzer, M.W. (2020). Nuclear periphery takes center stage: the role of nuclear pore complexes in cell identity and aging. Neuron *106*, 899-911.

Connolly, P.F., Jager, R., and Fearnhead, H.O. (2014). New roles for old enzymes: killer caspases as the engine of cell behavior changes. Front Physiol *5*, 149.

D'Angelo, M.A., Gomez-Cavazos, J.S., Mei, A., Lackner, D.H., and Hetzer, M.W. (2012). A change in nuclear pore complex composition regulates cell differentiation. Dev Cell *22*, 446-458.

Ding, A.X., Sun, G., Argaw, Y.G., Wong, J.O., Easwaran, S., and Montell, D.J. (2016). CasExpress reveals widespread and diverse patterns of cell survival of caspase-3 activation during development in vivo. Elife *5*.

Eigler, T., Zarfati, G., Amzallag, E., Sinha, S., Segev, N., Zabary, Y., Zaritsky, A., Shakked, A., Umansky, K.B., Schejter, E.D.*, et al.* (2021). ERK1/2 inhibition promotes robust myotube growth via CaMKII activation resulting in myoblast-to-myotube fusion. Dev Cell *56*, 3349-3363 e3346.

Fernando, P., Kelly, J.F., Balazsi, K., Slack, R.S., and Megeney, L.A. (2002). Caspase 3 activity is required for skeletal muscle differentiation. Proc Natl Acad Sci U S A *99*, 11025-11030.

Ferrando-May, E., Cordes, V., Biller-Ckovric, I., Mirkovic, J., Gorlich, D., and Nicotera, P. (2001). Caspases mediate nucleoporin cleavage, but not early redistribution of nuclear transport factors and modulation of nuclear permeability in apoptosis. Cell Death Differ *8*, 495-505.

Forler, D., Rabut, G., Ciccarelli, F.D., Herold, A., Kocher, T., Niggeweg, R., Bork, P., Ellenberg, J., and Izaurralde, E. (2004). RanBP2/Nup358 provides a major binding site for NXF1-p15 dimers at the nuclear pore complex and functions in nuclear mRNA export. Mol Cell Biol *24*, 1155-1167.

Gomez-Cavazos, J.S., and Hetzer, M.W. (2015). The nucleoporin gp210/Nup210 controls muscle differentiation by regulating nuclear envelope/ER homeostasis. J Cell Biol *208*, 671-681.

He, L., Diedrich, J., Chu, Y.Y., and Yates, J.R., 3rd (2015). Extracting accurate precursor information for tandem mass spectra by RawConverter. Anal Chem *87*, 11361-11367.

Hervy, M., Hoffman, L., and Beckerle, M.C. (2006). From the membrane to the nucleus and back again: bifunctional focal adhesion proteins. Curr Opin Cell Biol *18*, 524-532.

Hutten, S., and Kehlenbach, R.H. (2006). Nup214 is required for CRM1-dependent nuclear protein export in vivo. Mol Cell Biol *26*, 6772-6785.

Ibarra, A., Benner, C., Tyagi, S., Cool, J., and Hetzer, M.W. (2016). Nucleoporin-mediated regulation of cell identity genes. Genes Dev *30*, 2253-2258.

Jacinto, F.V., Benner, C., and Hetzer, M.W. (2015). The nucleoporin Nup153 regulates embryonic stem cell pluripotency through gene silencing. Genes Dev *29*, 1224-1238.

Jacquemont, C., and Taniguchi, T. (2007). Proteasome function is required for DNA damage response and fanconi anemia pathway activation. Cancer Res *67*, 7395-7405.

Kato, K., Tanaka, T., Sadik, G., Baba, M., Maruyama, D., Yanagida, K., Kodama, T., Morihara, T., Tagami, S., Okochi, M.*, et al.* (2011). Protein kinase C stabilizes X-linked inhibitor of apoptosis protein (XIAP) through phosphorylation at Ser(87) to suppress apoptotic cell death. Psychogeriatr *11*, 90-97.

Kihlmark, M., Imreh, G., and Hallberg, E. (2001). Sequential degradation of proteins from the nuclear envelope during apoptosis. J Cell Sci *114*, 3643-3653.

Kihlmark, M., Rustum, C., Eriksson, C., Beckman, M., Iverfeldt, K., and Hallberg, E. (2004). Correlation between nucleocytoplasmic transport and caspase-3-dependent dismantling of nuclear pores during apoptosis. Exp Cell Res *293*, 346-356.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915.

Labade, A.S., Salvi, A., Kar, S., Karmodiya, K., and Sengupta, K. (2021). Nup93 and CTCF modulate spatiotemporal dynamics and function of the HOXA gene locus during differentiation. J Cell Sci *134*.

Lee, J.E., Lee, J.Y., Wilusz, J., Tian, B., and Wilusz, C.J. (2010). Systematic analysis of cis-elements in unstable mRNAs demonstrates that CUGBP1 is a key regulator of mRNA decay in muscle cells. PLoS One *5*, e11201.

Li, F., Ambrosini, G., Chu, E.Y., Plescia, J., Tognin, S., Marchisio, P.C., and Altieri, D.C. (1998). Control of apoptosis and mitotic spindle checkpoint by survivin. Nature *396*, 580-584.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

Lian, Y.L., Chen, K.W., Chou, Y.T., Ke, T.L., Chen, B.C., Lin, Y.C., and Chen, L. (2020). PIP3 depletion rescues myoblast fusion defects in human rhabdomyosarcoma cells. J Cell Sci *133*.

Liang, Y., Franks, T.M., Marchetto, M.C., Gage, F.H., and Hetzer, M.W. (2013). Dynamic association of NUP98 with the human genome. PLoS Genet *9*, e1003308.

Lin, W.S., Lu, K.M., Chung, M.H., Liu, S.T., Chen, H.H., Chang, Y.L., Wang, W.M., and Huang, S.M. (2010). The subcellular localization and protein stability of mouse alpha-actinin 2 is controlled by its nuclear receptor binding motif in C2C12 cells. Int J Biochem Cell Biol *42*, 2082-2091.

Louis, M., Zanou, N., Van Schoor, M., and Gailly, P. (2008). TRPC1 regulates skeletal myoblast migration and differentiation. J Cell Sci *121*, 3951-3959.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Luo, S.W., Zhang, C., Zhang, B., Kim, C.H., Qiu, Y.Z., Du, Q.S., Mei, L., and Xiong, W.C. (2009). Regulation of heterochromatin remodelling and myogenin expression during muscle differentiation by FAK interaction with MBD2. EMBO J *28*, 2568-2582.

M, H.D., Tashakor, A., O'Connell, E., and Fearnhead, H.O. (2020). Apoptosome-dependent myotube formation involves activation of caspase-3 in differentiating myoblasts. Cell Death Dis *11*, 308.

Madureira, P.A., Hill, R., Lee, P.W., and Waisman, D.M. (2012). Genotoxic agents promote the nuclear accumulation of annexin A2: role of annexin A2 in mitigating DNA damage. PLoS One *7*, e50591.

Mahrus, S., Trinidad, J.C., Barkan, D.T., Sali, A., Burlingame, A.L., and Wells, J.A. (2008). Global sequencing of proteolytic cleavage sites in apoptosis by specific labeling of protein N termini. Cell *134*, 866-876.

Marchetto, M.C., Belinson, H., Tian, Y., Freitas, B.C., Fu, C., Vadodaria, K., Beltrao-Braga, P., Trujillo, C.A., Mendes, A.P.D., Padmanabhan, K.*, et al.* (2017). Altered proliferation and networks in neural cells derived from idiopathic autistic individuals. Mol Psychiatry *22*, 820-835.

McAlister, G.C., Nusinow, D.P., Jedrychowski, M.P., Wuhr, M., Huttlin, E.L., Erickson, B.K., Rad, R., Haas, W., and Gygi, S.P. (2014). MultiNotch MS3 enables accurate, sensitive, and multiplexed detection of differential expression across cancer cell line proteomes. Anal Chem *86*, 7150-7158.

Murray, T.V., McMahon, J.M., Howley, B.A., Stanley, A., Ritter, T., Mohr, A., Zwacka, R., and Fearnhead, H.O. (2008). A non-apoptotic role for caspase-9 in muscle differentiation. J Cell Sci *121*, 3786-3793.

Nagaraj, R., Sharpley, M.S., Chi, F., Braas, D., Zhou, Y., Kim, R., Clark, A.T., and Banerjee, U. (2017). Nuclear localization of mitochondrial TCA cycle enzymes as a critical step in mammalian zygotic genome activation. Cell *168*, 210-223 e211.

Nakanishi, K., Sudo, T., and Morishima, N. (2005). Endoplasmic reticulum stress signaling transmitted by ATF6 mediates apoptosis during muscle development. J Cell Biol *169*, 555-560.

Nix, D.A., Fradelizi, J., Bockholt, S., Menichi, B., Louvard, D., Friederich, E., and Beckerle, M.C. (2001). Targeting of zyxin to sites of actin membrane interaction and to the nucleus. J Biol Chem *276*, 34759-34767.

Ossovskaya, V., Lim, S.T., Ota, N., Schlaepfer, D.D., and Ilic, D. (2008). FAK nuclear export signal sequences. FEBS Lett *582*, 2402-2406.

Park, N., Schweers, N.J., and Gustin, K.E. (2015). Selective removal of FG repeat domains from the nuclear pore complex by Enterovirus 2A(pro). J Virol *89*, 11069-11079.

Park, S.K., Aslanian, A., McClatchy, D.B., Han, X., Shah, H., Singh, M., Rauniyar, N., Moresco, J.J., Pinto, A.F., Diedrich, J.K.*, et al.* (2014). Census 2: isobaric labeling data analysis. Bioinformatics *30*, 2208-2209.

Patre, M., Tabbert, A., Hermann, D., Walczak, H., Rackwitz, H.R., Cordes, V.C., and Ferrando-May, E. (2006). Caspases target only two architectural components within the core structure of the nuclear pore complex. J Biol Chem *281*, 1296-1304.

Peng, J., Elias, J.E., Thoreen, C.C., Licklider, L.J., and Gygi, S.P. (2003). Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. J Proteome Res *2*, 43-50.

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and Salzberg, S.L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol *33*, 290-295.

Qin, B., Yu, J., Nowsheen, S., Wang, M., Tu, X., Liu, T., Li, H., Wang, L., and Lou, Z. (2019). UFL1 promotes histone H4 ufmylation and ATM activation. Nat Commun *10*, 1242.

Schwarzerova, K., Bellinvia, E., Martinek, J., Sikorova, L., Dostal, V., Libusova, L., Bokvaj, P., Fischer, L., Schmit, A.C., and Nick, P. (2019). Tubulin is actively exported from the nucleus through the Exportin1/CRM1 pathway. Sci Rep *9*, 5725.

Serganov, A.A., Udi, Y., Stein, M.E., Patel, V., Fridy, P.C., Rice, C.M., Saeed, M., Jacobs, E.Y., Chait, B.T., and Rout, M.P. (2022). Proteomic elucidation of the targets and primary functions of the picornavirus 2A protease. J Biol Chem, 101882.

Shimbo, K., Hsu, G.W., Nguyen, H., Mahrus, S., Trinidad, J.C., Burlingame, A.L., and Wells, J.A. (2012). Quantitative profiling of caspase-cleaved substrates reveals different drug-induced and cell-type patterns in apoptosis. Proc Natl Acad Sci U S A *109*, 12432-12437.

Soledad, R.B., Charles, S., and Samarjit, D. (2019). The secret messages between mitochondria and nucleus in muscle cell biology. Arch Biochem Biophys *666*, 52-62.

Sugiyama, K., Aida, T., Nomura, M., Takayanagi, R., Zeilhofer, H.U., and Tanaka, K. (2017). Calpain-dependent degradation of nucleoporins contributes to motor neuron death in a mouse model of chronic excitotoxicity. J Neurosci *37*, 8830-8844.

Tabb, D.L., McDonald, W.H., and Yates, J.R., 3rd (2002). DTASelect and Contrast: tools for assembling and comparing protein identifications from shotgun proteomics. J Proteome Res *1*, 21-26.

Tang, H.L., Tang, H.M., Fung, M.C., and Hardwick, J.M. (2015). In vivo CaspaseTracker biosensor system for detecting anastasis and non-apoptotic caspase activity. Sci Rep *5*, 9015.

Toyama, B.H., Arrojo, E.D.R., Lev-Ram, V., Ramachandra, R., Deerinck, T.J., Lechene, C., Ellisman, M.H., and Hetzer, M.W. (2019). Visualization of long-lived proteins reveals age mosaicism within nuclei of postmitotic cells. J Cell Biol *218*, 433-444.

Ullman, K.S., Shah, S., Powers, M.A., and Forbes, D.J. (1999). The nucleoporin nup153 plays a critical role in multiple types of nuclear export. Mol Biol Cell *10*, 649-664.

Walker, E.J., Younessi, P., Fulcher, A.J., McCuaig, R., Thomas, B.J., Bardin, P.G., Jans, D.A., and Ghildyal, R. (2013). Rhinovirus 3C protease facilitates specific nucleoporin cleavage and mislocalisation of nuclear proteins in infected host cells. PLoS One *8*, e71316.

Xu, T., Park, S.K., Venable, J.D., Wohlschlegel, J.A., Diedrich, J.K., Cociorva, D., Lu, B., Liao, L., Hewel, J., Han, X.*, et al.* (2015). ProLuCID: An improved SEQUEST-like algorithm with enhanced sensitivity and specificity. J Proteomics *129*, 16-24.

Yamashita, T., Aizawa, H., Teramoto, S., Akamatsu, M., and Kwak, S. (2017). Calpain-dependent disruption of nucleo-cytoplasmic transport in ALS motor neurons. Sci Rep *7*, 39994.

Yi, X., Tesmer, V.M., Savre-Train, I., Shay, J.W., and Wright, W.E. (1999). Both transcriptional and posttranscriptional mechanisms regulate human telomerase template RNA levels. Mol Cell Biol *19*, 3989-3997.

Figure 1 | Caspases fully proteolyze 5 Nups during myogenesis (+ **Figures S1-5**)

Figure 1 | Caspases fully proteolyze 5 peripheral Nups during myogenesis. (A) Immunoblots showing the degradation of Nups, PARP, and αII-spectrin, the activation of caspase-3/12 and the upregulation of myogenin in differentiating C2C12 cells. **(B)** Caspase-dependent proteolysis of Nups in differentiating C2C12 cells assessed by immunoblotting. Q-VD-OPh used at 30 μM. **(C)** Nuclear-to-total mRNA ratio in C2C12 cells undergoing myogenesis. The average value of three replicates shown for each time point. **(D)** A stable C2C12 cell line that expresses NES_{Rev}-GFP. In dashed boxes are cells undergoing MB-to-MB or MB-to-MT fusion. **(E)** Cytoplasmic-to-nuclear ratio of NES_{Rev}-GFP was quantified for 22 donor myoblast / closest myonuclei pairs. Grey lines link each pair. **(F)** Localization of lamin B1 and α-tubulin in differentiating C2C12 cells assessed by immunoblotting. Nuclear α-tubulin marked with an arrow. 30 μg of protein loaded per lane. **(G)** Cytoplasmic and nuclear levels of focal adhesion proteins in differentiating C2C12 cells. 25 μg of protein loaded per lane. MT: myotubes, MB: myoblasts.

Figure 2 | Tpr and Pom121 C-terminal fragments dissociate from the NPC during myogenesis
(+ **Figures S6-7**)

Figure 2 | Tpr and Pom121 C-terminal fragments dissociate from the NPC during myogenesis. (A and B) C2C12 cells immunostained using two Nup153 and two Tpr antibodies. In dashed boxes are apoptotic nuclei. **(C)** C2C12 cells undergoing myogenesis immunostained for Crm1, Nup153, and Tpr. In dashed boxes are TprC⁻ cells. **(D)** Tpr C-term immunoreactivity (normalized against Nup153 internal domain immunoreactivity) determined for 50+ nuclei. **(E)** Nup153, Tpr, and caspase-3 fragments in subcellular fractions of differentiating C2C12 cells visualized by immunoblotting. Q-VD-OPh used at 30 μM. **(F)** C2C12 cells undergoing myogenesis immunostained for mAb414, Nup153, and Pom121. **(G)** Pom121 C-term immunoreactivity (normalized against Nup153 internal domain immunoreactivity) determined for 50+ nuclei.

Figure 3 I Quantitative proteomics reveals how NPC trimming impacts the nuclear proteome of differenti-ating myoblasts (+ **Supplemental Spreadsheet 1**)

Figure 3 I Quantitative proteomics reveals how NPC trimming impacts the nuclear proteome of differentiating myo-blasts. (A) Preparation of subcellular fractions for quantitative mass spectrometry. **(B)** Subcellular fractions were validated using multiple protein markers. **(C)** Volcano plot showing how protein levels in the "nuclear soluble" fraction change during the first 24 hours of myogenic differentiation. Each data point colored to represent their transcriptional change during the same time frame. **(D)** Zoomed-in view of the dotted square area in **(C)**.

Figure 4 | NPC trimming accompanies nuclear accumulation of cytoplasmic, plasma membrane, and mitochondrial proteins (+ **Supplemental Spreadsheet 1**)

Figure 4 | NPC trimming accompanies nuclear accumulation of cytoplasmic, plasma membrane, and mitochondrial proteins. (A) Protein groups detected in the "nuclear soluble" fraction. Red dotted line: y = 2. **(B)** Protein level change in the "nuclear soluble" fraction when C2C12 cells are differentiated with or without 30 μM Q-VD-OPh. Red dotted line: y = 1. Data points for proteins labeled in **(A)** are shown in the right box. **(C)** Visual summary of multiple events that take place during myoblast-to-myotube transition.

Figure 5 I Caspase-mediated NPC trimming occurs during neurogenesis and ER stress (+ **Figure S8**)

Figure 5 I Caspase-mediated NPC trimming occurs during neurogenesis and ER stress. (A) Schematic representation of caspase-mediated NPC trimming. **(B)** Down-/up-regulation of survivin, caspase-3, and myogenin in differentiating C2C12 cells. **(C)** Relative timing of key events in myogenesis. **(D)** Expression levels of Nup153, PARP, active caspase-3, survivin, and neuronal differentiation markers (Sox2 and βIII-tubulin) were determined in differentiating neurons. **(E)** Acute ER stress induced using 1 μg/mL tunicamycin in C2C12 myotubes and reserve cells. The levels of Nups, BiP, XIAP, myosin heavy chain (MHC), myogenin, and active caspase-3 levels were monitored by immunoblotting. **(F)** Chronic ER stress was induced in differentiated C2C12 cells for 7 days using low doses of tunicamycin. Nups, BiP, and active caspase-3 levels were assessed.

Figure 6 I Three different modes of NPC proteolysis

Figure 6 I Caspase activation and NPC proteolysis patterns in apoptosis, cell differentiation, and chronic ER stress.