RESEARCH ARTICLE

TITLE: H^+/K^+ ion pump enhances cytoskeletal polarity to drive gastrulation in sea

urchin embryo

Running title: H^+/K^+ ion pump and gastrulation

Kaichi Watanabe[1,*], Yuhei Yasui[1], Yuta Kurose[2], Masashi Fujii[1], Takashi Yamamoto[1],

Naoaki Sakamoto[1,*], Akinori Awazu[1,3,*]

[1] Graduate School of Integrated Sciences for Life, Hiroshima University, Higashi-

Hiroshima, 739-8526, Japan

[2] Department of Mathematical and Life Sciences, Graduate School of Science,

Hiroshima University, Higashi-Hiroshima, 739-8526, Japan

[3] Research Center for the Mathematics on Chromatin Live Dynamics, Hiroshima

University, Higashi-Hiroshima, Hiroshima, 739-8526, Japan

*Corresponding authors (kaichi1849@hiroshima-u.ac.jp (K.W.); naosaka@hiroshima-

u.ac.jp (N. S.); awa@hiroshima-u.ac.jp (A. A.))

KEY WORDS: Gastrulation, H^+/K^+ ion pump, sea urchin, cytoskeleton, pH

regulation

Summary statement: The H^+/K^+ ion pump is critical for normal gastrulation in

sea urchin through pH regulation of cell-position dependent intracellular actin

distributions.

ABSTRACT

Gastrulation is a universal process in the morphogenesis of many animal embryos. In sea urchin embryos, it involves the invagination of single-layered vegetal plate into blastocoel. Although morphological and molecular events have been well studied for gastrulation, the mechanical driving forces and their regulatory mechanism underlying the gastrulation is not fully understood. In this study, structural features and cytoskeletal distributions were studied in sea urchin embryo using an "exogastrulation" model induced by inhibiting the H^+/K^+ ion pump with omeprazole. The vegetal pole sides of the exogastrulating embryos had reduced roundness indices, intracellular pH polarization, and intracellular F-actin polarization at the pre-early gastrulation compared with the normal embryo. Gastrulation stopped when F-actin polymerization or degradation was inhibited by *RhoA* or *YAP1* knockout, although pH distributions were independent of such a knockout. A mathematical model of sea urchin embryos at the early gastrulation reproduced the shapes of both normal and exogastrulating embryos using cell-dependent cytoskeletal features based on F-actin and pH distributions. Thus, gastrulation required appropriate cell position-dependent intracellular F-actin distributions regulated by the H^+/K^+ ion pump through pH control.

INTRODUCTION

Gastrulation is a universal and essential morphogenic process in various animals,

wherein a ball of single layered cells (blastula) differentiates into a multilayered

gastrula or early embryo (Serrano Nájera and Weijer, 2020; Shindo, 2018; Stower and

Bertocchini, 2017). Species-specific variations in this process provides the basis for

particular animal morphologies and leads to the generation of internal organs. The basic

mechanism of whole embryonic transformations induced by organizational

rearrangements and cell movements are expected to be universal.

Sea urchin embryos are commonly used for studying various morphogenetic

processes universally observed in animals, such as the left–right symmetry breakage

(Aihara and Amemia, 2001; McCain and McClay, 1994; Takemoto et al., 2016; Warner

and McClay, 2014), neural network formations (Burke et al., 2014; McClay et al., 2018;

Yaguchi et al., 2010), and gastrulation (Dan and Okazaki, 1956; Gustafson and

Kinnander, 1960; Hardin and Cheng, 1986; Kominami and Takata, 2004; Martik and

McClay, 2017) due to its evolutionary position as an ancestral deuterostome.

Furthermore, the gene regulatory network controlling endomesoderm specification in

sea urchin embryo has been well studied (Davidson et al., 2002; Oliveri & Davidson,

2004). Sea urchin gastrulation progresses through the following four steps (Kominami

and Takata, 2004): step 1 (after hatching), the embryo becomes an epithelial monolayer

with a thickened vegetal pole; step 2 (pre-gastrulation), the vegetal plate bends to

invade the blastocoel; step 3 (early gastrulation), the tubular archenteron is formed, and

secondary mesenchyme cells (SMCs) appear at the tip; step 4 (mid-late gastrulation),

the archenteron elongates by pulling on the filopodia of the SMC. Steps 1 and 2 refer to

the primary invagination stages, and steps 3 and 4 are those of the secondary

invagination stages (Dan and Okazaki, 1956; Gustafson and Kinnander, 1956).

The embryos of some sea urchin species exhibit anomalous morphogenesis

(exogastrulation) under various perturbations where the vegetal plate of the embryo

prolapses outward instead of invaginating. For example, the archenteron of embryos

cultured with LiCl or sugar is completely evaginated due to the depolymerization of

SMC microtubules in the blastocoel space connecting the primary intestine and

ectoderm (Dan and Okazaki, 1956; Hardin and Cheng, 1986; Khurrum et al., 2004).

These studies suggest that lifting the primary intestine from the mid-gut by

microtubules plays an important role in step 4 of gastrulation. Exogastrulation was also

observed by Rab35 knockdown that was also suggested to disturb whole embryonic

cytoskeleton distribution (Remsburg et al., 2021).

Moreover, bending of the vegetal plate in step 2 of normally cultured embryos of

the sea urchin, *Lytechinus pictus* (LP), occurs even when it is surgically isolated

(Ettensohn, 1984). It is suggested that the tip of the archenteron should be bottle-shaped

(Nakajima and Burke, 1996), and the area around the vegetal plate should be wedge-

shaped (Burke et al., 1991) to initiate primary invagination (Kominami and Takata,

2004). Additionally, mathematical models of vegetal poles suggest that primary

invagination occurs under the appropriate force conditions for "apical constriction,"

"cell tractor," "apical contractile ring," "apicobasal contraction," and "gel swelling"

(Davidson et al., 1995; Odell et al., 1981). However, it is unclear which of these effects

are essential for development up to pre-early gastrulation (steps 2 and 3).

In this study, the roles of the dominant factors in steps 2 and 3 of gastrulation of sea

urchins were determined through experimental analyses and mathematical modeling.

First, specific inhibition of H^+/K^+ ion pump activity by omeprazole caused "partial"

exogastrulation without the loss of microtubules in the blastocoel space of embryos

from the sea urchin, *Hemicentrotus pulcherrimus* (HP). The H^+/K^+ ion pump regulates

the pH in cells and pH influences the structure and force of the actomyosin system

(Köhler et al., 2012). Therefore, the partial exogastrulation was likely due to anomalous

cytoskeletal behaviors. Second, knockout experiments on cytoskeleton-related

molecules were performed in HP embryos and analyzed by fluorescence imaging. Third,

simulations of morphogenic process of normal and H^+/K^+ ion pump inhibited

(exhibiting exogastrulation) HP embryo during the pre-early gastrulation were

performed using a mathematical model utilizing the cytoskeletal force parameters

determined by the above experiments. The modeled shapes of the H^+/K^+ ion pump-

inhibited embryos of the sea urchin, *Lytechinus variegatus* (LV,) were compared to

those of HP to reveal the contributions of the H^+/K^+ ion pump to the regulation of

cytoskeletal behaviors.

RESULTS

H^+/K^+ ion pump inhibition in HP embryos caused partial exogastrulation

The development of HP embryos exposed to the specific H^+/K^+ ion pump inhibitor,

omeprazole, immediately after fertilization exhibited anomalous morphogenesis with

partial exogastrulation observed at the primary-secondary invagination stages and 36

hours post fertilization (hpf) (Fig. 1). Detailed comparisons of the structural features

around the vegetal pole between the normal (control) and inhibited embryos at each

gastrulation step are as follows. In step 1, control and inhibited embryos exhibited

equivalent vegetal pole thickening (18-20 hpf). In step 2, primary invagination occurred

in both control and inhibited embryos at the same time. However, intestinal invagination

was slightly shallower in the inhibited embryos than in the control embryos (20-22 hpf).

In step 3, inhibited embryos did not show further elongation of the archenteron into the

blastocoel, and the shape of inhibited embryos elongated along the animal-vegetal axis

(22-24 hpf). In step 4 (secondary invagination stage), the outward protrusion of the

vegetal pole side was more pronounced in the inhibited embryo (24 hpf). The inhibited

embryo had comparatively little elongation of the archenteron into the blastocoel

compared to the control embryos. At 36 hpf, "partial" exogastrulation, in which the tip

of archenteron folded inward by the normal primary invagination but the remaining part

extended outward during the secondary invagination, was observed in the inhibited

embryo.

The migration of SMCs was observed in the control embryos and in the inhibited

embryos (Fig. 2) indicating that filopodia were formed. Previously reported

exogastrulation was induced by the loss of pseudopodia (Dan and Okazaki, 1956;

Hoshi, 1979; Khurrum et al., 2004), therefore, partial exogastrulation in this report was

expected to occur by a different mechanism.

H$^+$/K$^+$ ion pump inhibition perturbed whole embryonic pH and cytoskeletal

distributions

The intracellular distribution of F-actin and pH was analyzed at the primary

invagination stage (steps 1 and 2) to reveal the mechanism underlying the

morphological changes in the inhibited embryos at the secondary invagination stage

(steps 3 and 4), (Fig. 3A, B). F-actin was visualized by fluorescence of actinin-GFP, and

intracellular pH was estimated with pH indicator, whose fluorescent intensity increased

as pH decreased. The polarity of intracellular F-actin and pH distribution was evaluated

by estimating the fluorescence intensities ratio between the apical and basal sides of

cells.

Similar apical/basal ratio distributions of actinin-GFP intensities were obtained between

the control and inhibited embryos, while the ratios gradually increased closer to the

vegetal pole in both embryos at step 1 (Fig. 3C, D, Fig. S1A, B). The distribution of the

apical/basal ratios of actinin-GFP intensities in step 2 were similar between control and

inhibited embryos only in the animal half (Fig. 3E, Fig. S1C). The ratios of the control

embryos from the equator to the vegetal pole was larger than that of inhibited embryos,

while nearest the vegetal pole, control embryos showed smaller ratio than that of

inhibited embryos. Therefore, the formation of an anomalous embryonic shape of

inhibited embryos correlated with the deviation of intracellular F-actin polarity in the

vegetal half of the embryos in primary invagination.

The apical/basal ratio of the pH indicator was greater than one in both control and

inhibited embryos in step 2 (Fig. 3F, Fig. S1D, S2) indicating that the pH on the apical

side of each cell was always lower than that on the basal side. Additionally, this ratio

was higher in the vegetal half than that in the animal half in the control embryo, while

this pattern disappeared in the inhibited embryo (Fig. 3F, Fig. S2). Thus, the polarity of

the pH indicator fluorescent intensities was positively correlated with that of actinin-

GFP in whole embryos (Fig. 3E, F).

Meanwhile, inhibition of the H^+/K^+ ion pump did not influence the whole

embryonic distribution of fibropellin-1 (Fig. 3G, H), a F-actin scaffold in the apical

lamina of the apical pole (Burke et al., 1998; Nakajima and Burke, 1996).

Gastrulation starts but does not progress in embryos with knocked out

cytoskeleton regulators

CRISPR-Cas9-mediated knockout of the typical enhancer and repressor of F-actin

formation (*RhoA* and *YAP1*) (Beane et al., 2006; Dupont et al., 2011) was performed by

microinjection of sgRNAs designed with *Cas9* mRNA (Fig. S3) to elucidate the

contribution of F-actin to the gastrulation process. Although the mutation frequencies of

RhoA and *YAP1* knockout were 55.6% and 77.8%, respectively, and the frameshift rate

was 22.3% in either knockout (Fig. S3), the gastrulation stopped at step 2 at the primary

invagination stage in either *RhoA* or *YAP1* knockout embryos (Fig. 4A) and normal

gastrulation and exogastrulation were not observed. Interestingly, the phenotypes

between these gene knockout embryos were similar, although the gene functions are

different (Fig. S3). Pigmented cells were observed at 45 hpf indicating that development

had not ceased. This showed that primary invagination occurred even in embryos with

mild perturbation of F-actin function, but subsequent gastrulation processes required

appropriate F-actin-derived forces.

The whole embryonic distribution of average intensity and apical/basal ratios of the

pH indicator was unchanged in *RhoA*-knockout embryos compared with the control, and

therefore, perturbation of F-actin polymerization did not affect the pH gradient of the

cell (Fig. 4B, C).

Mathematical model of embryonic shape formations in early gastrulation

considering cell-dependent F-actin polarity

A mathematical model of cell motion at the cross section including animal and vegetal

poles during steps 1-3 of gastrulation was constructed to examine the influence of

intracellular F-actin distribution on the formation of embryonic shapes. The model was

constructed based on that of gastrulation in *Nematostella vectensis* (Tamulonis et al.,

2011) that consisted of cells constructed by springs and beads circularly connected to

form a 2-dimensional cross section of the embryo. The following assumptions were

made based on experimentally known facts.

I) Each embryo contained three cell types: pigment cells (Takata and Kominami, 2004),

wedge cells (Burke at al., 1991), and other cells (Fig. 5). The pigment cells near the

entrance to the cavity were assumed to be bottle-shaped (Takata and Kominami, 2004)

because of the site-specific force in these cells. This may explain the bending of the

vegetal plate occurring in embryos with knocked out F-actin polymerization regulators

(Fig. 4), or a surgically removed animal pole (Ettensohn, 1984). The apical length was

sufficiently shorter than the basal length and the apical sides of wedge cells were wider

than the basal side (Fig. 5A) to form a similar shape to previously observed embryos

(Kominami and Takata, 2004).

II) The width of the apical and basal sides changes in a cell-dependent manner in

control and inhibited embryos due to cytoskeletal forces generated by F-actin. The

apical side elongation was expected to correlate with F-actin concentration because

higher F-actin concentration was expected to generate stronger cell cortical forces.

Thus, the lengths of the apical and basal sides of cells, except pigment cells and wedge

cells, were determined based on their apical/basal ratios of actinin-GFP intensities at

step 2 (Fig. 5A).

III) The cell divisions and 3-dimensional mutual cell invasions mainly contributing to

late gastrulation were not included because these processes were rarely observed in

early gastrulation (Mizoguchi, 1999). Therefore, the present model described embryo

shape dynamics of the control and inhibited embryos as 2-dimensional motions and

deformations of the apical and basal sides of 64 cells (Fig. 5A).

The simulations showed different shapes (Fig. 5B), although each shape was like

that experimentally observed at step 3 (Fig. 1). Similar temporal changes in roundness

indices (Fig. 5C) of control embryos (0.85 ~ 0.9) and inhibited embryos (0.75 ~ 0.8)

were observed in the simulated data (Fig. 5D, Fig. S4A) and experimental data (Fig. 5E,

Fig. S4B).

The simulation where the elongation ratio of the apical side of each cell was a little

larger than that of control was also performed, with this model named: "over-polarized

embryo" (Fig. 5A, B). The whole embryonic shape of this model resulted in a wider and

larger roundness index than the control embryos (Fig. S4A). This suggested that the

apical/basal ratio polarity of F-actin distribution in each cell was sensitive to the whole

embryonic shape.

DISCUSSION

Inhibition of the H^+/K^+ ion pump resulted in abnormal embryo shape with partial

exogastrulation due to intracellular apical F-actin distribution at the vegetal pole (Fig. 3,

5). F-actin concentrations generating cell cortical forces were significantly larger at the

apical side of each cell than that of the basal side in the vegetal half of the control

embryos, except for cells at the vegetal pole. Therefore, the apical planes of wedge cells

and other cells in the vegetal pole side were expected to be significantly wider than the

basal planes. Conversely, the apical plane of cells in the vegetal half of the inhibited

embryo were not significantly wider compared to the basal sides due to lower apical F-

actin levels. The simulation results showed that such expected features in the vegetal

half may progress to normal early gastrulation in the control embryos and partial

exogastrulation in the inhibited embryos (Fig. 5). Therefore, appropriate embryonic

intracellular apical F-actin distribution in the vegetal half is required for gastrulation.

One recent study suggested an outer intestinal embryo is formed by Rab35

knockdown that might disturb the whole embryonic intracellular F-actin polarity

(Remsburg et al., 2021). The results of the present study support this suggestion, and

further indicate that the intracellular F-actin polarity on the vegetal pole side of the

embryo is particularly important for accurate gastrulation and the intracellular pH

controlled by H^+/K^+ ion pump is involved in this process.

Gastrulation does not proceed when apical lamina is inhibited (Burke et al., 1991).

The whole embryonic distributions of apical lamina along cells were nearly identical

between the control and inhibited embryos, although large differences were observed

between the cell shapes in the vegetal half and whole embryos. This indicated that the

generated force was independent of the relative amounts of apical lamina in each cell.

Embryos with partial knockout of F-actin regulating factors (*RhoA* or *YAP1*) stopped the

gastrulation at the primary invagination step (Fig. 4A, B). This indicated that the

seconday invagination is inhibited without appropriately regulated F-actin, even if the

apical lamina exists. Therefore, the apical lamina was suggested to be the scaffold for

apical F-actin to generate the force required for the deformation processes driving the

secondary invagination of gastrulation and whole embryonic deformation.

CRISPR-Cas9-mediated *RhoA* knockout showed intermediate level (55.6%) of

mutation frequency (Fig. S3) and the knockout embryos exhibited the primary

invagination. However, LV embryos expressing dominant negative RhoA failed to

initiate primary invagination (Beane et al., 2006). Therefore, contribution of F-actin

function is considerable for the primary invagination, but the extent of the dependency

on F-actin function is less and a mechanism other than the F-actin-related system might

be involved in this initial process. As mentioned above, F-actin function must be

essential for the secondary invagination.

The F-actin network formation generating cell cortical forces was drastically

enhanced with decreased pH *in vitro* (Köhler et al., 2012) (25). The lower pH at the

apical side compared with the basal side was maintained by H^+/K^+ ion pump activity.

Whole embryonic distributions of apical/basal ratios of pH indicator at step 2 correlated

with those of intracellular actinin-GFP (Fig. 3D, F) suggesting that the F-actin

concentration was high when the pH was low at the apical side of cells. Additionally,

the apical/basal F-actin polarity in each cell significantly decreased in the vegetal half of

the embryo by inhibiting H^+/K^+ ion pump activity (Fig. 3E). Conversely, the whole

embryonic distribution of intracellular pH and polarity was unchanged even when F-

actin polymerization was directly perturbed by knockout of F-actin regulating factors

(Fig. 4D). This suggested that the intracellular pH controlled by the H^+/K^+ ion pump is a

one-sided regulator of F-actin polymerization.

Inhibition of the H^+/K^+ ion pump did not influence on the whole embryonic

distribution of the apical lamina (Fig. 3G, H). This suggested that the regulation of F-

actin intracellular distribution by pH is independent of the apical lamina. The embryo

exhibited partial exogastrulation if the intracellular pH polarity at the primary

invagination stage was weaken by inhibiting H^+/K^+ ion pump activity. This indicated

that H^+/K^+ ion pump activity in the primary invagination stage provided a dominant

contribution to the progress of the normal secondary invagination to form appropriate

intracellular F-actin polarity in the whole embryo.

The influence of pH on F-actin network formation and force generation is a

universal biochemical feature confirmed by *in vitro* experiments (Köhler et al., 2012).

Inhibition of H^+/K^+ ion pump activity significantly decreases intracellular pH in the LV

embryo (Schatzberg et al., 2015), while an increase in the intracellular pH was observed

in the majority of cells in the HP embryo (Fig. S5). Decreases in the intracellular pH of

LV may induce increases in the apical/basal polarities of intracellular pH and F-actin

concentration like the "over-polarized embryo" model. The over-polarized embryo

model exhibited a wider embryonic shape with a larger roundness index than the control

embryos (Fig. 5B, S4B) which is consistent with LV imaging (Schatzberg et al., 2015).

This supported the suggestion that H^+/K^+ ion pump activity significantly contributed to

early gastrulation through the regulation of intracellular F-actin apical/basal

polarizations.

Omeprazole treatment of sea urchin embryos disrupts the left-right asymmetric

formations of the adult rudiment although gastrulation is completed (Bessodes et al.,

2012; Duboc et al., 2005; Hibino et al., 2006). The left-right asymmetric *nodal* gene

expression observed immediately after gastrulation is important for establishing this

asymmetry (Duboc et al., 2005). Frogs, chickens, zebrafish, and ascidians disrupt left-

right asymmetry by inhibiting H^+/K^+ ion pump activity (Kawakami et al., 2005; Levin

et al., 2002; Shimeld and Levin, 2006). Disrupted left-right asymmetry in HP is

observed at lower ($\sim$ 3/5 times) omeprazole concentrations (Hibino et al., 2006) than

those in this study. Inhibition of H^+/K^+ pump activity by dilute omeprazole solutions

induced weak perturbation of gut formation in this study, which may disrupt cell-cell

interactions and regulate whole embryonic *nodal* gene expression.

This report revealed the close relationship between pH, F-actin, and H^+/K^+ ion

pump activity inducing normal/abnormal gastrulation in sea urchins through

intracellular F-actin regulation. Embryonic F-actin distribution on the apical side of

each cell was important for normal gastrulation, and H^+/K^+ ion pump activity may be

involved in F-actin distribution. This finding provides key insights into the evolutionary

relationship between development and morphogenesis. Intracellular apical/basal

polarity of pH showed different cellular features between vegetal and animal poles in

step 2 of gastrulation. However, the formation of these position-dependent intracellular

states was not explored. Therefore, upstream regulators of H^+/K^+ ion pump activity

should be investigated to reveal the detailed morphogenesis molecular mechanism. The

universal effect of inhibiting H^+/K^+ ion pump activity on early embryo formation among

different animals will be tested in the future.

MATERIALS AND METHODS

Animals and embryos

Adult Japanese sea urchins (HP) were collected from Seto Inland Sea or Tateyama Bay.

Eggs and sperm were obtained by coelomic injection of 0.55 M KCl. Fertilized eggs

were cultured in filtered seawater at 16°C.

Inhibition of H^+/K^+ ion pump by omeprazole

A 100 mM stock solution of omeprazole (FUJIFILM Wako Pure Chemical Corporation,

Japan) dissolved in dimethyl sulfoxide (DMSO) was immediately added to fertilized

eggs to a final concentration of 100 μM and cultured at 16°C.

***In situ* nucleus and F-actin visualization in embryos**

Sea urchin embryos were fixed by immersion in fixative III (4% paraformaldehyde,

32.5% filtered seawater, 32.5 mM 3-(N-morpholino)propane sulfonic acid (MOPS), pH

7.0, 162.5 mM NaCl) for 16 h at 4°C. Embryos were washed three times using 0.5 mL 1

x phosphate buffered saline (PBS), stored in 100% ethanol at -20°C, and used within 1

week. Fixed embryos were washed three times with 1 x PBS, permeabilized with 0.5%

Triton X-100 for 20 min at room temperature, washed three times with 1 x PBS, and

stained with 100 nM phalloidin (in 1 x PBS) for 45 min at room temperature. Stained

embryos were washed three times with 1 x PBS, mounted in SlowFade Gold antifade

mountant with DAPI (Molecular Probes) and observed using a confocal laser scanning

microscopy (Carl Zeiss LSM 700, Germany) with laser illumination at 405/435 nm

excitation/emission for DAPI and 555/573 nm excitation/emission for phalloidin,

respectively.

Live imaging of pH, actinin, and apical lamina

Intracellular pH of sea urchin embryos was visualized using pHrodo Red AM

intracellular pH indicator or 5-(and-6)-carboxy SNARF-1 (C-1270) (ThermoFisher

Scientific, USA) at final concentrations of 10 μM and 5 μM, respectively, and stained at

16°C for 30 min. The embryos were washed in filtered seawater and observed using

confocal microscopy with laser illumination at 555/585 nm excitation/emission for

pHrodo and 555/573 m excitation/emission for SNARF-1, respectively.

Actinin and fibropellin-1 were visualized following fusion of the proteins with GFP.

RNA was extracted from mesenchyme blastula stage HP embryos using ISOGEN

(Nippongene, Japan) according to the manufacturer's instructions. Actinin and

fibropellin-1 coding sequences were amplified by RT-PCR using SuperScript™ III

Reverse Transcriptase (Thermo Fisher Scientific, USA) with the primers stated in Table

S1 and cloned into pGreenLantern2-derived plasmid at restriction sites EcoRI and

XhoI. mRNA (actinin-GFP, fibropellin-1-GFP, in that order) was synthesized *in vitro*

using an mMESSAGE mMACHINE T7 ultra transcription kit (ThermoFisher Scientific,

USA) and purified using an RNeasy mini kit (Qiagen, Nederland). mRNA was

microinjected into fertilized eggs as described by Liu et al., (2019).

Fluorescent images were acquired by confocal microscopy using an excitation of

488 nm, and emission of 515 nm. The images of the animal and vegetal poles of the

embryo were analyzed based on the *z*-stack image with the largest area by averaging

seven *z*-axis images with a total thickness of 6 µm.

Quantification of experimental images

The fluorescent images of embryos were acquired according to the angle θ (0°~180°)

between the vegetal and animal poles along the circumference (Fig. S6) and

transformed into a band-like image using the "Polar Transformer" function

(https://imagej.nih.gov/ij/plugins/polar-transformer.html) of ImageJ 2.1.0. Each band-

like image was filtered using a median filter (radius = 1.0). The filtered images were

binarized using "Mean" (pHrodo indicator), "IsoData" (actinin), and "Triangle"

(fibropellin-1 and SNARF-1) functions, respectively, to obtain the cellular regions of

the embryo. The apical/basal ratios of the pH indicator, actin-GFP, and fibropellin-1-

GFP for θ were determined using average fluorescent intensities over the regions with

the width of 3.126 μm from the apical and basal sides.

The fluorescent intensity of the pHrodo indicator was defined by the average

fluorescent intensity over the intracellular region with a width of 6.232 μm from the

apical side (Fig. S6). Fluorescent values were estimated by taking the ratio between the

observed fluorescence intensity and the average background fluorescence intensity

around the entire embryo.

F-actin and apical lamina were stained by mRNA microinjections of actinin-GFP

and fibropellin-1-GFP to fertilized eggs. Nonnegligible variations was inevitable in the

concentration of injected mRNA among fertilized egg samples, which would be

amplified during development. Therefore, only the apical/basal ratios were used to

evaluate the intracellular features of F-actin and apical lamina distributions.

Two data points were obtained for each θ from each fluorescence image of the

embryo since the left and right sides of the embryo were considered axis symmetric

against the animal-vegetal axis. Both data were used to estimate the sample average and

95% confidence interval of each value, where the number of samples (n) was given by 2

× [number of observed embryos].

Gene knockout by CRISPR-Cas9

Knock out of endogenous *RhoA* and *YAP1* in HP was performed using the method

described by Liu et al., (2019). The oligonucleotide sequences used in sgRNA

preparations were listed in Table S2.

Heteroduplex mobility assay (HMA) and DNA sequencing analysis

Analysis of knockout embryos by HMA and DNA sequencing was performed according

to Liu et al., (2019). The primer sequences used to amplify each target region were

listed in Table S3.

Mathematical model of sea urchin embryos

A 2-dimensional particles model describing cellular motion at the cross section of

animal poles and vegetal poles of sea urchin embryos during steps 1 to 3 of gastrulation

was constructed. The determined number of cells was based on that at the equatorial

plane of the embryo at the blastula stage (Mizoguchi, 1999). The following assumptions

were made: 1) each cell was represented by 16 particles with radii (r) of 1.125 μm, and

each embryo was represented by 64 cells connected in a ring (Fig. S7). 2) Each cell

perimeter was 36 μm and the height and width of the embryo at step 1 was 110 μm and

100 μm, respectively, which was consistent with sea urchin embryo observations

(results not shown). 3) The number of cells was constant because cell divisions and

cell invasions from other cross-sections were rarely observed during the primary

invagination stage (Mizoguchi, 1999). 4) The motion of each particle obeyed the

following overdamped limit of the equation of motion:

$\quad \frac{\partial X_{i,j}}{\partial t} = -\frac{1}{\gamma}\frac{\partial V}{\partial X_{i,j}}$ $\quad\quad$ (1),

$\quad$ where $X_{i,j} = (x_{i,j}(t), y_{i,j}(t))$ is the position of the j-th particle constructing the i-th

$\quad$ cell ($i = 0,1,2…63$ and $j = 0,1,2…15$) on the x–y plane at time t (Fig. 5, Fig. S7), γ is the

$\quad$ coefficient of the drag force acting on each particle, and V is the potential of the entire

$\quad$ system. 5) The y-axis was parallel to the animal-vegetal axis of the embryo model. We

$\quad$ rewrote $\frac{V}{\gamma} = U$ and U was calculated by:

$\quad$ $U = U_{bond} + U_{collision} + U_{actin} + U_{adhesion} + U_{ca} + U_{ea}$ $\quad\quad$ (2),

$\quad$ where U_{bond} is the elastic force potential between each neighboring pair of particles in

$\quad$ each cell to maintain each cell perimeter, denoted by:

$\quad\quad\quad\quad$ $U_{bond} = \sum_{i=0}^{63} \sum_{j=0}^{15} \frac{k_b}{2}(d_{(i,j),(i,j+1)} - 2r\,)^2$ $\quad\quad$ (3),

$\quad$ where k_b, $d_{(i,j),(i,j+1)}$, and r are the coefficient of elasticity, the distance between the j-

$\quad$ th and j +1-th particle ($j + 1 = 0$ for $j = 15$) in the i-th cell at time t, and the particle

$\quad$ radius, respectively.

$\quad$ $U_{collision}$ is the potential of excluded volume effects among all particles denoted as

$\quad$ follows:

$\quad$ $U_{collision} = \sum_{i<k} \sum_{j<l} \theta(2r - d_{(i,j),(k,l)})\frac{k_c}{2}(2r - d_{(i,j),(k,l)})^2$ $\quad\quad$ (4),

$\quad$ where k_c is the coefficient of repulsion between two particles, and $\theta(x)$ is the

$\quad$ Heaviside step function.

$\quad$ U_{actin} is the elastic force potential to form and sustain cell shape with the expansion

$\quad$ and contraction of the apical and basal sides by the cytoskeleton denoted as follows:

$U_{actin} = \sum_i \frac{k_a}{2}\left((d_{(i,10),(i,14)} - l_i^a)^2 + (d_{(i,2),(i,6)} - l_i^b)^2\right)$ (5),

where the apical and basal sides of the i-th cell consisted of particles with $j = 10 - 14$

and $j = 2 - 6$, respectively. The wideness of the apical and basal sides at time t were

given by $d_{(i,10),(i,14)}$ and $d_{(i,2),(i,6)}$, respectively. The basic wideness of the apical and

basal sides were given by l_i^a and l_i^b, respectively. k_a was assumed by the coefficient

of elasticity to sustain the wideness of the apical and basal sides of the cell.

$U_{adhesion}$ is the potential force for cell adhesion by proteins such as cadherin, and is

denoted as follows:

$U_{adhesion} = \sum_i \sum_j \frac{k_h}{2}(d_{(i,j),(i+1,k(j))} - 2r)^2$ (6),

where k_h is the coefficient for the adhesive forces between the particles of the i-th and

$i + 1$-th cells ($i + 1 = 0$ for $i = 63$), and $k(j) \equiv 6, 7, 8, 9, 10$ for $j \equiv 2, 1, 0, 15, 14,$

respectively (Fig. S7).

U_{ca} is the potential of the forces to conserve each cell area denoted as follows:

$U_{ca} = \sum_i \frac{k_{ca}}{2}(s_i - s_i^{ini})^2$ (7),

where k_{ca} is the coefficient of elasticity required to maintain each cell area. The area

of the i-th cell was estimated by $s_i = \sum_{j=0}^{15}|X_{i,j} - X_i^{cc}|$ with $X_i^{cc} = \frac{1}{16}\sum_{j=0}^{15} X_{i,j}$, and

s_i^{ini} referred to s_i given by $X_{i,j}$ at the initial state.

U_{ea} is the potential of the forces to maintain the area (volume) of the sea urchin

embryo and denoted as follows:

$U_{ea} = \frac{k_{ea}}{2}(S - S^{ini})^2$ (8),

where k_{ea} is the coefficient of elasticity required to maintain embryo area. The area

was estimated by $S = \sum_{i=0}^{63} |X_i^{cc} - X^{ec}|$ with $X^{ec} = \frac{1}{64} \sum_{i=0}^{63} X_i^{cc}$, and S^{ini} referred to

S given by $X_{i,j}$ at the initial state.

U_{ca} and U_{ea} were considered the alternatives of the volume-conserving forces in real

3-D cells and embryos as internal pressures in each cell and embryo are considered

isotropic.

Simulation method of mathematical model

The simulations of the present mathematical model were performed through the

integrals of equation of motion (1) using Euler method with time intervals of 0.000064

h with conserved $X^{ec} = (0, 0)$. In all models, the parameters

k_{bond}, $k_{collision}$, k_{actin}, $k_{adhesion}$, k_{ca}, and k_{ea} were given empirically as 9375, 6250,

8125, 8125, 625, and 0.00625 hour^{-1}, respectively, because there were no experiments to

measure or estimate them. The qualitative features of the results were independent of

the details of these values if the order was maintained. The model formed the embryo

shape at step 1 of gastrulation if the appropriate cell type-dependent values of l_i^a and

l_i^b, $= l_i^{a,ini}$ and $l_i^{b,ini}$ were given for the models of pigment cells, wedge cells, and

other cells (Fig. 5). This configuration gave the particle positions $X_{i,j}$ at time t = 0

(initial configuration) in all simulations.

The early gastrulation processes were simulated by the change in l_i^a and l_i^b

respectively, from $= l_i^{a,ini}$ and $l_i^{b,ini}$ to $= l_i^{a,fin}$ and $l_i^{b,fin}$ at time $t = 0$. Here, $l_i^{a,fin}$

and $l_i^{b,fin}$ were assumed to obey $l_i^{a,fin} + l_i^{b,fin} = l_i^{a,ini} + l_i^{b,ini} =$ constant among cells

except pigment cells based on the expectation that F-actin was constant. The model

showed similar structural behaviors to the early gastrulation as the relaxation process of

$d_{(i,10),(i,14)} \rightarrow l_i^{a,fin}$ and $d_{(i,2),(i,6)} \rightarrow l_i^{b,fin}$.

Statistical analysis

All experiments were performed independently 2 times or more with 4 replicates or

more per experiment. Statistical test was performed using SciPy library

(https://www.scipy.org/). The roundness index of the vegetal side of each embryo was

evaluated by: [embryo width half-way between the bottom and middle of the

embryo]/[embryo length] (Fig. 5C).

Acknowledgements

We thank Prof. Masato Kiyomoto (Tateyama Marine Laboratory, Ochanomizu

University) for supplying live sea urchins. We would like to thank Editage

(www.editage.com) for English language editing.

Competing interests

No competing interests declared.

Author Contributions

K.W., N.S., and A.A. conceived and designed the study; K.W., Y.K., and N.S. conducted

the experiments; K.W., Y.Y., M.F., and A.A. analyzed the data; K.W. and A.A.

conducted the mathematical model construction and simulations; K.W., N.S., and A.A.

wrote the manuscript with support from all authors; T.Y. supervised the work.

Funding information

This work was supported by JSPS KAKENHI [grant, award number: 17K05614,

21K06124 to A.A.]; and JSPS KAKENHI [grant, award number: 17K07241, 20K06602

to N.S.]; JSPS KAKENHI [grant, award number: 19K20382 to M.F]; SASAGAWA

SCIENTIFIC RESERCH GRANT, [award number: 2021-4023, to K.W.]; Hiroshima

University Graduate School Research Fellowship, 2021-2024 (InformationAI Field, to

K.W.);.

References

Aihara M. and Amemiya, S. (2001). Left-right positioning of the adult rudiment in sea

urchin larvae is directed by the right side. *Development* **128**(24), 4935-4948.

doi:10.1242/dev.128.24.4935

Beane, W. S., Gross, J. M. and McClay, D. R. (2006). RhoA regulates initiation of

invagination, but not convergent extension, during sea urchin gastrulation. *Dev.*

Biol. **292**(1), 213-225. doi:10.1016/j.ydbio.2005.12.031

Bessodes, N., Haillot, E., Duboc, V., Röttinger, E., Lahaye, F. and Lepage, T.

(2012). Reciprocal signaling between the ectoderm and a mesendodermal left-

right organizer directs left-right determination in the sea urchin embryo. *PLoS*

Genet. **8**(12), e1003121. doi:10.1371/journal.pgen.1003121

Burke, R. D., Lail, M. and Nakajima, Y. (1998). The apical lamina and its role in cell

adhesion in sea urchin embryos. *Cell Adhesion Comm*. **5**(2), 97-108.

doi:10.3109/15419069809040284

Burke, R. D., Moller, D. J., Krupke, O. A. and Taylor, V. J. (2014). Sea urchin

neural development and the metazoan paradigm of neurogenesis. *Genesis* **52**(3),

208-221. doi:10.1002/dvg.22750

Burke, R. D., Myers, R. L., Sexton, T. L. and Jackson, C. (1991). Cell movements

during the initial phase of gastrulation in the sea urchin embryo. *Dev. Biol*.

146(2), 542-557. doi:10.1016/0012-1606(91)90255-2

Dan, K. and Okazaki, K. (1956). Cytoembryological studies of sea urchins: III. Role

of the secondary mesenchyme cells in the formation of the primitive gut in sea

urchin larvae. *Biol. Bull.* **110**, 29-42.

Davidson, E. H., Rast, J. P., Oliveri, P., Ransick, A., Calestani, C., Yuh, C. H.,

Minokawa, T., Amore, G., Hinman, V., Arenas-Mena, C., Otim, O., Brown,

C. T., Livi, C. B., Lee, P. Y., Revilla, R., Schilstra, M. J., Clarke, P. J., Rust,

A. G., Pan, Z., Arnone, M. I., Rowen, L., Cameron, R. A., McClay, D. R.,

Hood, L. and Bolouri, H. (2002). A provisional regulatory gene network for

specification of endomesoderm in the sea urchin embryo. *Dev. Biol.*, **246**(1), 162-

190. doi: 10.1006/dbio.2002.0635

Davidson, L. A., Koehl, M. A., Keller, R. and Oster, G. F. (1995). How do sea

urchins invaginate? Using biomechanics to distinguish between mechanisms of

primary invagination. *Development* **121**(7), 2005-2018.

doi:10.1242/dev.121.7.2005

Duboc, V., Röttinger, E., Lapraz, F., Besnardeau, L. and Lepage, T. (2005). Left-

right asymmetry in the sea urchin embryo is regulated by nodal signaling on the

right side. *Dev. Cell*. **9**(1), 147-158. doi:10.1016/j.devcel.2005.05.008

Dupont, S., Morsut, L., Aragona, M., Enzo, E., Giulitti, S., Cordenonsi, M.,

Zanconato, F., Le Digabel, J., Forcato, M., Bicciato, S. et al. (2011). Role of

YAP/TAZ in mechanotransduction. *Nature* **474**(7350), 179-183.

doi:10.1038/nature10137

Ettensohn, C. A. (1984). Primary invagination of the vegetal plate during sea urchin

gastrulation. *Amer. Zool.* **24**(3), 571-588. doi:10.1093/icb/24.3.571

Gustafson, T. and Kinnander, H. (1956). Microaquaria for time-lapse

cinematographic studies of morphogenesis in swimming larvae and observations

on sea urchin gastrulation. *Exp. Cell Res*. **11**(1), 36-51. doi:10.1016/0014-

4827(56)90188-4

Gustafson, T. and Kinnander, H. (1960). Cellular mechanisms in morphogenesis of

the sea urchin gastrula: The oral contact. *Exp. Cell Res.* **21**(2), 361-373.

doi:10.1016/0014-4827(60)90268-8

Hardin, J. D. and Cheng, L. Y. (1986). The mechanisms and mechanics of

archenteron elongation during sea urchin gastrulation. *Dev. Biol*. **115**(2), 490-

501. doi:10.1016/0012-1606(86)90269-1

Hibino, T., Ishii, Y., Levin, M. and Nishino, A. (2006). Ion flow regulates left-right

asymmetry in sea urchin development. *Dev. Genes Evol*. **216**(5), 265-276.

doi:10.1007/s00427-005-0051-6

Hoshi, M. (1979). Exogastrulation induced by heavy water in sea urchin larvae. *Cell*

Differ. 8(6), 431-5. doi:10.1016/0045-6039(79)90039-3

Kawakami, Y., Raya, Á., Raya, R. M., Rodríguez-Esteban, C. and Izpisúa

Belmonte, J. C. (2005). Retinoic acid signalling links left-right asymmetric

patterning and bilaterally symmetric somitogenesis in the zebrafish embryo.

Nature **435**(7039), 165-171. doi:10.1038/nature03512

Khurrum, M., Hernandez, A., Eskalaei, M., Badali, O., Coyle-Thompson, C. and

Oppenheimer, S. B. (2004). Carbohydrate involvement in cellular interactions in

sea urchin gastrulation. *Acta Histochem*. **106**(2), 97-106. doi:

10.1016/j.acthis.2004.01.001

Kimberly, E. L. and Hardin, J. (1998). Bottle cells are required for the initiation of

primary invagination in the sea urchin embryo. *Dev. Biol*. **204**(1), 235-50.

doi:10.1006/dbio.1998.9075

Köhler, S., Schmoller, K. M., Crevenna, A. H., Bausch, A. R. (2012). Regulating

contractility of the actomyosin cytoskeleton by pH. *Cell Rep*. **2**(3), 433-439.

doi:10.1016/j.celrep.2012.08.014

Kominami, T. and Takata, H. (2004). Gastrulation in the sea urchin embryo: A model

system for analyzing the morphogenesis of a monolayered epithelium. *Dev.*

Growth Differ. **46**(4), 309-326. doi:10.1111/j.1440-169x.2004.00755.x

Levin, M., Thorlin, T., Robinson, K. R., Nogi, T. and Mercola, M. (2002).

Asymmetries in H^+/K^+-ATPase and cell membrane potentials comprise a very

early step in left-right patterning. *Cell* **111**(1), 77-89. doi:10.1016/S0092-

8674(02)00939-X

Liu, D., Awazu, A., Sakuma, T., Yamamoto, T. and Sakamoto, N. (2019).

Establishment of knockout adult sea urchins by using a CRISPR-Cas9 system.

Dev. Growth Differ. **61**(6), 378-388. doi:10.1111/dgd.12624

Martik, M. L. and McClay, D. R. (2017). New insights from a high-resolution look at

gastrulation in the sea urchin, *Lytechinus variegatus*. *Mech. Dev.* **148**, 3-10.

doi:10.1016/j.mod.2017.06.005

McCain, E.R. and McClay, D.R. (1994). The establishment of bilateral asymmetry in

sea urchin embryos. *Development* **120**(2), 395-404. doi:10.1242/dev.120.2.395

McClay, D. R., Miranda, E. and Feinberg, S. L. (2018). Neurogenesis in the sea

urchin embryo is initiated uniquely in three domains. *Development* **145**(21),

dev167742. doi: 10.1242/dev.167742

Mizoguchi, H. (1999). Cell numbers in the gut of the embryo of the sea urchin

Hemicentrotus pulcherrimus. *Zool. Sci.* **16**(2), 341-344. doi:10.2108/zsj.16.341

Nakajima, Y. and Burke, R. D. (1996). The initial phase of gastrulation in sea urchins

is accompanied by the formation of bottle cells. *Dev. Biol.* **179**(2), 436-446.

doi:10.1006/dbio.1996.0273

Odell, G. M., Oster, G., Alberch, P. and Burnside, B. (1981). The mechanical basis

of morphogenesis. I. Epithelial folding and invagination. *Dev. Biol*. **85**(2), 46-62.

doi:10.1016/0012-1606(81)90276-1

Oliveri, P. and Davidson, E. H. (2004). Gene regulatory network controlling

embryonic specification in the sea urchin. *Curr. Opin. Genet. Dev*., **14**(4), 351–

360. doi: 10.1016/j.gde.2004.06.004

Remsburg, C., Testa, M., Song, J. L. (2021). Rab35 regulates skeletogenesis and

gastrulation by facilitating actin remodeling and vesicular trafficking. *Cells &*

Dev., **165**(3), 203660. doi: 10.1016/j.cdev.2021.203660

Schatzberg, D., Lawton, M., Hadyniak, S. E., Ross, E. J., Carney, T., Beane, W. S.,

Levin, M. and Bradham, C. A. (2015). H^+/K^+ ATPase activity is required for

biomineralization in sea urchin embryos. *Dev. Biol.* **406**(2), 259-270.

doi:10.1016/j.ydbio.2015.08.014

Serrano Nájera, G. and Weijer, C. J. (2020). Cellular processes driving gastrulation

in the avian embryo. *Mech. Dev*. **163**, 103624. doi:10.1016/j.mod.2020.103624

Shimeld, S. M. and Levin, M. (2006). Evidence for the regulation of left-right

asymmetry in *Ciona intestinalis* by ion flux. *Dev. Dyn*. **235**(6), 1543-1553.

doi:10.1002/dvdy.20792

Shindo, A. (2018). Models of convergent extension during morphogenesis. (2018).

WIREs Dev. Biol. **7**(1), e293. doi:10.1002/wdev.293

Stower, M. J. and Bertocchini, F. (2017). The evolution of amniote gastrulation: the

blastopore-primitive streak transition. *Wiley Interdiscip. Rev. Dev. Biol*. **6**(2),

e262. doi: 10.1002/wdev.262

Takata, H. and Kominami, T. (2004). Pigment cells trigger the onset of gastrulation in

tropical sea urchin *Echinometra mathaei*. *Dev. Growth Differ*. **46**(1), 23-35.

doi:10.1111/j.1440-169X.2004.00726.x

Takemoto, A., Miyamoto, T., Simono, F., Kurogi, N., Shirae-Kurabayashi, M.,

Awazu, A., Suzuki, K.-i. T. and Sakamoto, N. (2016). Cilia play a role in

breaking left-right symmetry of the sea urchin embryo. *Genes Cells* **21**(6), 568-

578. doi:10.1111/gtc.12362

Tamulonis, C., Postma, M., Marlow, H. Q., Magie, C. R., de Jong, J. and

Kaandorp, J. (2011). A cell-based model of *Nematostella vectensis* gastrulation

including bottle cell formation, invagination and zippering. *Dev. Biol.* **351**(1),

217-228. doi:10.1016/j.ydbio.2010.10.017

Voiculescu, O., Bodenstein, L., Lau, I. J. and Stern, C. D. (2014). Local cell

interactions and self-amplifying individual cell ingression drive amniote

gastrulation. *eLife* **3**, e01817. doi: 10.7554/eLife.01817

Warner, J. F. and McClay, D. R. (2014). Left-right asymmetry in the sea urchin.

Genesis **52**(6), 481-487. doi:10.1002/dvg.22752

Yaguchi, S., Yaguchi, J., Angerer, R. C., Angerer, L. M. and Burke, R. D. (2010).

TGFβ signaling positions the ciliary band and patterns neurons in the sea urchin

embryo. *Dev. Biol.* **347**(1), 71-81. doi:10.1016/j.ydbio.2010.08.009

Figures

Fig. 1. Developmental stages of control and H^+/K^+ ion pump-inhibited HP

embryos. Typical bright-field images of the gastrulation process (steps 1-4) and 36 hpf

embryos (scale bars: 30 μm). Arrows indicate vegetal pole positions. Gastrulation did

not progress after step 2 in the inhibited embryos, and the outward protrusion of the

vegetal pole side was more pronounced at step 4. At 36 hpf, "partial" exogastrulation

was observed in the inhibited embryo.

Fig. 2. *In situ* SMC visualization of HP embryo development with inhibited H^+/K^+ ion pumps. Confocal microscopy images at 24 hpf (A), 27 hpf (B), and 29 hpf (C) (scale bars: 25 μm). The nucleus and F-actin in the embryos were stained with DAPI (magenta) and phalloidin (cyan), respectively. SMC migration was confirmed at all developmental stages as indicated by red triangles.

Fig. 3. Whole embryonic distributions of pH, actinin-GFP, and fibropellin-1-GFP.

(A-B) Confocal fluorescence microscopy images of co-stained intracellular pH (pH

indicator: red) and actinin (actinin-GFP: green) at gastrulation step 1 (A) and step 2 (B)

on the cross-section of animal poles and vegetal poles of control and inhibited embryos

(scale bars: 30 μm). (C) Definition of angle θ (0°~180°) from the vegetal pole (0°) to

the animal pole (180°) along the circumference of the embryo cross-section. (D-F)

Average apical/basal ratios and 95% confidence intervals (error bars) at angle θ of

actinin-GFP intensities at step 1 (D) and step 2 (E), pH indicator intensities at step 2 (F),

and fibropellin-1-GFP intensities at step 2 (G). Blue and orange curves and bars

represent the control and inhibited embryo values, respectively. Gray bars indicate

significantly different average values between the control and inhibited embryos

according to Welch's t-test ($p < 0.05$) (see also Fig. S1.). The correlation coefficients of

apical/basal ratios between actinin-GFP intensities (E) and pH indicator intensities (F)

in control and inhibited embryos were 0.56 and 0.64, respectively. (H) Fluorescence

images of fibropellin-1-GFP from control and inhibited embryos at gastrulation step 2

(scale bars: 30 μm).

Fig. 4. Effect of F-actin regulator knockout on gastrulation.

(A) Bright-field images of gastrulation (steps 2-3) in control, *RhoA* knockout, and *YAP1*

knockout embryos at selected time-points. PMCs and pigment cells were observed in all

embryos suggesting that development did not stop. The knockout embryos did not form

the structure like prism larva observed at 45 hpf (step 3) in the control embryo (scale

bars: 30 μm). (B-C) Average fluorescence intensities (arbitrary fluorescence units) and

95% confidence intervals (error bars) of intracellular pH indicator (B) and apical/basal

ratio of pH indicator (C) of *RhoA* knockout embryos and control embryos as a function

of angle θ. The meanings of the colors and θ are stated in Fig. 3 and Fig. S1.

(A) Embryo model

Pigment cells (vegetal plate)

Wedge cells

Other cells

$i = 32$

Cell index (i)

$i = 63$ $i = 0$

i : Cell index

Vegetal   Animal

Control (n = 34)
Inhibited (n = 34)

Control apical ($l_i^{a,fin}$)
Control basal ($l_i^{b,fin}$)
Inhibited apical ($l_i^{a,fin}$)
Inhibited basal ($l_i^{b,fin}$)
Over-polarized apical ($l_i^{a,fin}$)
Over-polarized basal ($l_i^{b,fin}$)

Apical/basal ratio of actinin-GFP

Step 2

Apical/basal length of model cells (μm)

(B) Simulated initial embryo shape at 18 hpf

Simulated embryo shapes at 26 hpf

Control embryo   Inhibited embryo   Over-polarized embryo

30 μm

(C)

Top

Distance

Width   Width

Bottom

$$\text{Roundness} = \frac{[\text{Width of embryo at 1/4 from vegetal pole}]}{[\text{Length of embryo}]}$$

$$= \frac{[\text{Width}]}{[\text{Distance}]}$$

(D)

Roundness

Control embryo model

Inhibited embryo model

Time (hpf)

(E)

Roundness

= 11
= 10
= 16
= 8   # = 11
Control embryos
= 10
Inhibited embryos
= 11
= 14   # = 14   # = 15

Time (hpf)

Fig. 5. Coarse-grained model and simulation of control, inhibited, and over-

polarized embryos. (A) Modeled cell lengths of apical and basal sides of control,

inhibited, and over-polarized embryos; $l_i^{a,fin}$ and $l_i^{b,fin}$ refer to the final length of the

apical and basal sides of the *i*-th cell, respectively (see Materials and methods).

Distributions of the final length of the former two models was determined based on the

distributions of apical/basal ratios of actinin-GFP intensities from control and inhibited

embryos in step 2. The top center panel was a modification of Fig. 3D. Red, green, and

blue circles in the top panel and panel (B) represent pigment cells, wedge cells, and

other cells, respectively. (B) Snapshots of the initial embryo shape at 18 hpf and at 26

hpf of the three models. (C) Definition of the roundness of the vegetal side of embryos

from simulation results and imaging. (D-E) Roundness indices of vegetal sides of

modeled control and inhibited embryos (D) and experimentally determined values (E).

Blue and orange # refer to the number of control and inhibited embryos at each time

point, error bars indicated 95% confidence intervals, and * indicated that the roundness

of control embryos was significantly larger than that of inhibited embryos according to

Welch's t-test (p < 0.05, see Fig. S4B).