Title: HCN1 channels mediate mu opioid receptor long-term depression at insular

cortex inputs to the dorsal striatum

Braulio Munoz[1], Brandon M. Fritz[1] and Brady K. Atwood[1,2].

[1]Department of Pharmacology & Toxicology, Indiana University School of Medicine,

Indianapolis, IN, 46202, USA.

[2]Stark Neurosciences Research Institute, Indiana University School of Medicine,

Indianapolis, IN, 46202, USA.

Author Contact information

Corresponding author: Brady K. Atwood, PhD.

Indiana University School of Medicine

Department of Pharmacology & Toxicology

320 W 15th St.

NB 400C

Indianapolis, IN, USA 46202

Tel: 317-274-8917

Fax: 317-274-7714

Email: bkatwood@iu.edu

Abstract

Mu opioid receptors (MORs) are expressed in the dorsal striatum, a brain region that mediates goal-directed (via the dorsomedial striatum), and habitual (via the dorsolateral striatum, DLS) behaviors. Our previous work indicates that glutamate transmission is depressed when MORs are activated in the dorsal striatum, inducing MOR-mediated long-term synaptic depression (MOR-LTD) or short-term depression (MOR-STD), depending on the input. In the DLS, MOR-LTD is produced by MORs on anterior insular cortex (AIC) inputs and MOR-STD occurs at thalamic inputs, suggesting input-specific MOR plasticity mechanisms. Here, we evaluated the mechanisms of induction of MOR-LTD and MOR-STD in the DLS using pharmacology and optogenetics combined with patch clamp electrophysiology. We found that cAMP/PKA signaling and protein synthesis are necessary for MOR-LTD expression, similar to previous studies of endocannabinoid-mediated LTD (eCB-LTD) in DLS. However, MOR-LTD does not require mTOR signaling as eCB-LTD does. MOR-STD does not utilize these same mechanisms. We characterized the role of presynaptic HCN1 channels in MOR-LTD induction as HCN1 channels expressed in AIC are necessary for MOR-LTD expression in the DLS. These results suggest a mechanism in which MOR activation needs HCN1 to induce MOR-LTD, suggesting a new target for pharmacological modulation of synaptic plasticity, providing new opportunities to develop novel drugs to treat alcohol and opioid use disorders.

Keywords: synaptic plasticity, long-term depression, dorsal striatum, corticostriatal, mu opioid receptor, insular cortex, HCN1 channel.

Introduction

The opioid system is expressed throughout the brain(Le Merrer, Becker, Befort, & Kieffer, 2009) and promotes synaptic plasticity in many brain regions(Bao et al., 2007; Dacher & Nugent, 2011a, 2011b; Drake, Chavkin, & Milner, 2007; Iremonger & Bains, 2009), including the dorsal striatum (DS)(Atwood, Kupferschmidt, & Lovinger, 2014; Hawes, Salinas, Lovinger, & Blackwell, 2017; Lovinger, 2010; Munoz, Fritz, Yin, & Atwood, 2018; Muñoz, Haggerty, & Atwood, 2020). The DS is the primary input nucleus of the basal ganglia and can be subdivided in two structures: the dorsomedial striatum (DMS), which controls goal-directed learning, and the dorsolateral striatum (DLS), which regulates habit formation(Burton, Nakamura, & Roesch, 2015; Corbit & Janak, 2016; Hilário & Costa, 2008; Lovinger, 2010; O'Tousa & Grahame, 2014). Drug and alcohol use shift from being outcome-driven to becoming more stimulus-driven and compulsive in nature during addiction development(Corbit & Janak, 2016; Everitt & Robbins, 2016; Hopf & Lesscher, 2014; Shen et al., 2018). This transition from flexible goal-directed drug use to more habitual or compulsive use, is paralleled by a transition in the neural activity from the DMS to the DLS(Corbit & Janak, 2016; Renteria, Cazares, & Gremel, 2020; Shen et al., 2018), thereby shifting the balance of action control from DMS to DLS. Previous reports have demonstrated the role of glutamatergic synaptic plasticity in behavior, showing that the disruption of glutamatergic long-term synaptic depression (LTD) at orbitofrontal cortex-DMS synapses prevents habit learning(Gremel et al., 2016), and the disruption of glutamatergic LTD in DLS promotes habit learning(DePoy et al., 2013; C. Nazzaro et al., 2012). Therefore, changes in the expression of excitatory synaptic plasticity affect specific DS-related behaviors.

Our previous work established that mu opioid receptor (MOR) activation induces long-term synaptic depression (MOR-LTD) or short-term depression (MOR-STD) of glutamate release in both the DLS and DMS(Atwood, Kupferschmidt, et al., 2014; Munoz et al., 2018; Muñoz et al., 2020). We demonstrated that MOR-mediated synaptic inhibition is expressed at multiple DS synapses (cortical, thalamic, amygdala and cholinergic), but only a subset of corticostriatal synapses expressed MOR-LTD(Atwood, Kupferschmidt, et al., 2014) that was disrupted by prior *in vivo* ethanol exposure(Munoz et al., 2018; Muñoz et al., 2020). We also found that mice that were exposed to alcohol and the opioid oxycodone in vivo had ablated corticostriatal MOR-LTD when measured in brain slices(Atwood, Kupferschmidt, et al., 2014; Munoz et al., 2018; Muñoz et al., 2020) MOR plasticity at other DS synapses was unaffected by alcohol exposure(Munoz et al., 2018; Muñoz et al., 2020). Of particular interest, in DLS, alcohol and opioid-sensitive MOR-LTD of glutamatergic input is completely restricted to anterior insular cortex (AIC) inputs(Atwood, Kupferschmidt, et al., 2014; Munoz et al., 2018). Altogether, these data led us to hypothesize that MOR plasticity utilizes different mechanisms at distinct glutamatergic synapses in DS. As we previously demonstrated that MOR-LTD and endocannabinoid LTD (eCB-LTD) are mutually occlusive in DS(Atwood, Kupferschmidt, et al., 2014), we predicted that these two forms of LTD would utilize similar mechanisms. cAMP/PKA signaling mediates eCB-LTD in the nucleus accumbens, amygdala, cerebellum, and hippocampus(Atwood, Lovinger, & Mathur, 2014). In addition, local protein translation in the presynaptic region is enhanced by the activation of CB1 receptor, which is important for the induction of inhibitory LTD in the hippocampus(Younts et al., 2016) and likely also in the DS(Yin, Davis, Ronesi, &

Lovinger, 2006). eCB-LTD at GABAergic synapses in the hippocampus is also mediated

by mTOR signaling(Younts et al., 2016). Therefore, we sought to test whether MOR-

LTD in DLS, and specifically at AIC-DLS synapses is mediated by cAMP/PKA, protein

translation and mammalian target of rampamycin (mTOR) signaling. As cAMP also

regulates hyperpolarization-activated cyclic nucleotide-gated (HCN) channels, these

channels can presynaptically regulate glutamate release, and the type 1 HCN (HCN1)

channel may be localized to corticostriatal neurons, we also explored a potential role for

HCN1 in mediating MOR-LTD in DLS(He, Chen, Li, & Hu, 2014; Z. Huang, Li, Aguado,

Lujan, & Shah, 2017; Moosmang, Biel, Hofmann, & Ludwig, 1999; Postea & Biel, 2011)

In the current study, we probed glutamatergic synapses in the DLS using ex vivo mouse

brain slice electrophysiology. We used broad electrical stimulation in C57BL/6J mice to

test all glutamatergic inputs. We used an optogenetics approach to probe specific

glutamate synapses in DLS, using Emx1-Ai32 mice to test all cortical inputs, VGluT2-

Ai32 mice to test all thalamic inputs, and C57BL/6J mice stereotaxically injected with an

AAV-ChR2 vector into the AIC to specifically probe AIC-DLS synapses. In combination

with these tools, we pharmacologically and genetically probed the mechanism involved

in the expression of MOR-LTD in the DLS. We demonstrate that cAMP/PKA signaling

and protein synthesis are necessary for MOR-LTD expression, but not for MOR-STD.

We describe a novel role of presynaptic AIC HCN1 channels in the induction of MOR-

LTD in DLS. We also report that mTOR signaling is required for eCB-LTD of glutamate

transmission in the DLS, but this is not required for MOR-LTD indicating a divergence in

the mechanisms of these two similar forms of synaptic plasticity.

Results

MOR activation express glutamatergic LTD in the DLS

We previously showed that the activation of MOR in DLS produces LTD of eEPSCs as
well as oEPSCs(Munoz et al., 2018). Using three approaches, we confirmed that 5 min
application of the MOR agonist DAMGO (0.3 µM) induces MOR-LTD. We found that
DAMGO produced a persistent decrease in eEPSC amplitude in DLS MSNs (72 ± 5%;
Figure 1A-D) in brain slices from C57Bl/6J mice. To generally probe corticostriatal
synapses, we used 470 nm light delivered through the microscope objective to evoke
oEPSCs in brain slices from Emx1-Ai32 mice (Figure 1E), where the activation of MOR
again induced MOR-LTD (71 ± 5%; Figure 1F-H). Finally, to specifically stimulate the
release of glutamate from AIC inputs to the DLS, we infused an AAV vector expressing
ChR2 in the AIC (Figure 1I), as we previously did(Munoz et al., 2018). As shown
previously, activation of MORs produced glutamatergic LTD of AIC-DLS transmission
(66 ± 7%; Figure 1J-L). We also confirmed that thalamostriatal inputs expressed MOR-
STD in the DLS using VGluT2-Ai32 transgenic mice (15-20 min: 70 ± 9%; 30-40 min: 93
± 9%; Figure supplement 1A-B) as we previously showed(Atwood, Kupferschmidt, et al.,
2014; Munoz et al., 2018). These data replicate our previous work that MORs mediated
LTD of AIC-DLS synapses and STD of thalamostriatal synapses in the DLS. We
previously showed that eCB-LTD and MOR-LTD were mutually occlusive in
DLS(Atwood, Kupferschmidt, et al., 2014). In order to determine whether we could infer
potential MOR-LTD mechanisms from what is known about eCB-LTD, we investigated
whether eCB-LTD also occurred at AIC-DLS synapses. We used the same AAV
approach described above (Figure supplement 2A) and tested the effects of stimulation-

induced LTD and responses to the CB1 receptor agonist WIN55,212-2 (1 μM). We

found that eCB-LTD (63 ± 7%; Figure 2B-D) and CB1 agonist-induced LTD (68 ± 8%;

Figure 2B, C, E) both occur at AIC-DLS synapses. Next, we explored the signaling

cascades that mediate MOR plasticity in DLS using what is known about eCB-LTD as a

guide for our experiments.

Inhibition of presynaptic PKA blocks MOR-mediated LTD

The activation of opioid receptors produces an inhibition of adenylyl cyclase (AC)

mediated by Gαi, decreasing cAMP levels, which leads to an attenuation of

cAMP/protein kinase A (PKA) signaling(Atwood, Lovinger, et al., 2014). However, little

is known about PKA's role in MOR-mediated LTD. Using KT5720 (1 μM), a specific and

cell permeable inhibitor of PKA, which brain slices were incubated in for 1h prior to

recordings, we demonstrated that the inhibition of PKA blocks the maintenance of MOR-

LTD of eEPSCs in DLS (101 ± 6%; Figure 3A-C). Corticostriatal MOR-LTD was also

blocked by PKA inhibition (90 ± 5%; Figure 3G-I), as was MOR-LTD at AIC-DLS

synapses (96 ± 4%; Figure 3M-O). We found that these effects were due to inhibiting

presynaptic PKA activity as at least 30 min treatment with the membrane-impermeable

PKA inhibitor PKI (1 μM), intracellularly infused through the recording micropipette into

the postsynaptic MSN, did not affect MOR-LTD in the DLS (eEPSCs: 84 ± 2%; Figure

3D-F) (corticostriatal oEPSCs: 54 ± 9%; Figure 3J-L) (AIC-DLS oEPSCs: 62 ± 6%;

Figure 3P-R). Furthermore, thalamostriatal inputs, that express MOR-mediated STD,

were not affected by broad inhibition of PKA (15-20 min: 84 ± 6%; 25-35 min: 107 ±

10%) or just inhibition of postsynaptic PKA (15-20 min: 87 ± 3%; 30-40 min: 111 ± 9%;

Figure supplement 1C-D). As we expected, our data demonstrate the role of presynaptic PKA uniquely in MOR-LTD.

The activation of adenylyl cyclase disrupts MOR-mediated LTD

Demonstrating a role for PKA signaling in MOR-LTD implicates adenylyl cyclase (AC) signaling as mediating MOR-LTD. To more specifically test this, we persistently activated AC using 20 μM forskolin throughout recording. In the presence of forskolin, DAMGO produced a transient increase of the EPSC amplitudes (Figure 4B, E and H), followed by a blockade MOR-LTD at all excitatory synapses (105 ± 4%; Figure 4A-C) and at corticostriatal (102 ± 3%; Figure 4D-F) and AIC inputs (93 ± 7%; Figure 4G-I). MOR-mediated STD at thalamic synapses was not affected by the application of forskolin (15-20 min: 83 ± 6%; 25-35 min: 96 ± 6%; Figure supplement 1E-F). Altogether, these findings indicate that cAMP/PKA signaling pathway is involved in the induction and maintenance of MOR-LTD in DLS.

Protein translation is required to produce MOR-mediated LTD

Next, we tested the role of protein translation in MOR-LTD. Using, bath application of 80 μM cycloheximide (selective inhibitor of protein synthesis) during the entire recording, we found that the inhibition of protein translation blocked MOR-LTD generally (104 ± 9%; Figure 5A-C) and specifically at cortical (105 ± 9%; Figure 5D-F) and AIC inputs (91 ± 2%; Figure 5G-I). Similar to our experiments targeting cAMP/PKA signaling, thalamic inputs were unaffected by cycloheximide treatment (15-20 min: 73 ± 6%; 30-40 min: 98 ± 9%; Figure supplement 1G-H).

It had been reported that eCB protein translation-mediated-LTD involves mTOR

signaling in the hippocampus. To test if mTOR is required for eCB-LTD also in the DLS,

we used bath application of 100 nM Torin 2 (a selective mTOR inhibitor). We

demonstrated that the inhibition of mTOR signaling disrupted eCB-LTD in the DLS

(Control: 77 ± 3% vs Torin 2: 103 ± 8%; Figure 6A-C). However, MOR-LTD does not

require mTOR signaling in the DLS (79 ± 2%; Figure 6D-F). These data suggest that

protein translation is necessary for MOR-mediated LTD, but mTOR signaling is not

involved and indicate that eCB- and MOR-LTD, while qualitatively similar, do not utilize

identical signaling mechanisms.

Calcium signaling is not required for MOR-mediated LTD

Our data to this point indicates that presynaptic cAMP/PKA and protein synthesis are

critical for the induction and maintenance of MOR-LTD. However, calcium is vital to

presynaptic neurotransmitter release and calcium-sensitive signaling mechanisms may

be required for LTD induction(Atwood, Lovinger, et al., 2014). Therefore, we next tested

whether Ca^{2+} is critical for MOR-LTD expression. First, we replaced Ca^{2+} for Sr^{2+}

throughout the duration of the recording, to induce asynchronous glutamate release. We

found that the replacement of Ca^{2+} did not impact MOR-LTD (74 ± 5%; Figure

supplement 2). Thus, we used another approach where we prepared a DAMGO solution

without Ca^{2+} in order to evaluate the role of Ca^{2+} in the induction of MOR-LTD. Our data

show that Ca^{2+} is not necessary to induce MOR-LTD (82 ± 7%; Figure supplement 2).

Presynaptic HCN1 channels are indispensable for MOR-mediated LTD

Given that we identified cAMP signaling as important for MOR-LTD expression (Figure

4), we also explored the possibility of HCN1 channels as key factors for MOR-LTD. To

do so, we applied the general HCN blocker ZD7288 (25 μM) to brain slices for at least 30 minutes before DAMGO application and during the entire recording. HCN inhibition blocked the expression of MOR-LTD (101 ± 7%; Figure 7A-C), suggesting a novel role for HCN channels in DLS glutamatergic synaptic plasticity. Next, we characterized the effects of HCN channels in glutamatergic DLS synaptic transmission, by bath applying 25 μM ZD7288 for at least 15 minutes before recording sEPSCs. We found that the inhibition of HCN channels caused an enhancement in the frequency and a decrease in the amplitude of sEPSCs (Figure 7D-F), without affecting rise and decay times (Figure 6C, G and H). These data suggest, that HCN channels are present on both pre- and postsynaptic sites and they mediate MOR-LTD in the DLS.

To further characterize this new role of HCN channels and establish that presynaptic HCN1 channels are specifically involved in the mechanism of expression of MOR-LTD, we used HCN1flox mice injected with AAV-cre vector into AIC to knock out the expression of HCN1 channel specifically in the AIC, and using AAV-GFP vector-infused mice as controls (Figure 7A, B). Using electrical stimulation and recording in the DLS, we found that the ablation of HCN1 in the AIC reduced the magnitude of MOR-LTD in AAV-cre-injected mice (Figure 7C-F), relative to AAV-GFP-injected mice (Figure 7C-G; cre: 90 ± 4% vs GFP: 76 ± 3%; Figure 7E). Next, we tested if the deletion of presynaptic HCN1 channels produced changes in basal striatal glutamatergic transmission. We found that the lack of AIC HCN1 channels did not affect any synaptic parameters (Figure 7H-L). These data suggest that HCN1 channels specifically on AIC inputs are necessary for MOR-LTD expression in the DLS.

Discussion

The current study demonstrates that the mechanism of MOR-mediated LTD is similar to the mechanism of induction of the inhibition of glutamate of CB1 or eCB-LTD in that it involves cAMP-PKA-protein translation signaling and both occur at AIC-DLS synapses (Atwood, Lovinger, et al., 2014; Lonart et al., 2003; Mato, Lafourcade, Robbe, Bakiri, & Manzoni, 2008; Yasuda, Huang, & Tsumoto, 2008; Yin et al., 2006; Younts et al., 2016) (Figures. 2-5). However, we found notable differences such as a lack of dependence on mTOR signaling and calcium (Figure 6 and supplement 2) and a novel mechanism involving HCN1 signaling for MOR-LTD (Figures 7 and 8). Figure Supplement 3 summarizes the findings of the paper. Furthermore, the MOR-LTD mechanism is different than that of MOR-STD (Figure supplement 1), suggesting that MOR-mediated plasticity does not utilize a common pathway at all glutamatergic synapses.

Our previous report demonstrates that MORs expressed in AIC are the sole source of MOR-mediated LTD at glutamate inputs to the DLS(Munoz et al., 2018). However in the DMS, MORs on medial prefrontal cortex, anterior cingulate cortex, and basolateral amygdala inputs are capable of inducing LTD(Muñoz et al., 2020). MOR-LTD also occurs locally at cholinergic interneuron synapses that co-release glutamate on to MSNs(Munoz et al., 2018; Muñoz et al., 2020). However, alcohol only disrupts MOR-LTD at AIC inputs to DLS and at medial prefrontal and anterior cingulate cortex inputs to DMS leaving MOR plasticity at thalamostriatal, amygdalostriatal and cholinergic interneurons unaffected(Munoz et al., 2018; Muñoz et al., 2020). We also found that the opioid oxycodone disrupts MOR-LTD in DLS(Atwood, Kupferschmidt, et al., 2014), and given that this is mediated by AIC-DLS MORs(Munoz et al., 2018), it follows that alcohol

and opioids have similar actions on MOR plasticity at this synapse. Given that alcohol

(and presumably oxycodone and other opioids) do not affect MOR plasticity at these

other non-corticostriatal synapses, differential mechanisms that underlie MOR plasticity

at each type of synapse may confer susceptibility to the deleterious effects of these

drugs and could account for their mechanisms of action in the development of habitual

or compulsive use of the drugs.

Previous work has shown that the inhibition of PKA disrupts eCB-LTD in the nucleus

accumbens(Mato et al., 2008) and that presynaptic PKA is required to induce

cannabinoid LTD in cerebellum(Lonart et al., 2003) and eCB synaptic plasticity in

hippocampus(Yasuda et al., 2008). Moreover, previous reports had shown that the

activation of AC by forskolin enhances corticostriatal EPSC amplitudes(Cho et al., 2008)

and blocked delta opioid receptor-mediated LTD of inhibitory postsynaptic

currents(Patton, Roberts, Lovinger, & Mathur, 2016). cAMP/PKA signaling has also

been implicated in type 2/3 metabotropic glutamate receptor- and type 1b serotonin

receptor-mediated LTD at various synapses in the brain(Atwood, Kupferschmidt, et al.,

2014). Until now, the role of cAMP/PKA signaling pathway in MOR-mediated plasticity

had not been addressed. Our results demonstrate that presynaptic PKA inhibition

blocked AIC-DLS MOR-LTD (Figure 3), but not thalamostriatal MOR-STD (Figure

supplement 1C-D). While the magnitudes and time courses differed slightly between

measures of eEPSCs and oEPSCs, it appears that PKA is not required for MOR-

mediated inhibition of glutamate release, but it is required for the maintenance of MOR-

LTD. On the other hand, we found that AC activation disrupted both acute MOR-

mediated inhibition (in fact turning it into a transient potentiation) and MOR-LTD (Figure

4), but had no effect on MOR-STD (Figure supplement 1E-F). Altogether these data

suggest that cAMP/PKA pathway may be a general mechanism whereby GPCRs that

signal through Gαi may induce LTD, but a receptor's ability to couple to Gαi does not

necessarily mean that it will induce cAMP/PKA-dependent LTD upon activation.

Protein translation is necessary for striatal eCB-LTD(Yin et al., 2006) and is involved in

GPCR-mediated LTD at some synapses, but not others(Atwood, Kupferschmidt, et al.,

2014; Younts et al., 2016) suggesting that protein translation can be an important

component of glutamatergic LTD, depending on the synapse. Indeed, local protein

translation can take place in the presynaptic region, where the activation of CB1

receptors enhances protein translation, being critical for the induction and not the

maintenance of inhibitory LTD(Younts et al., 2016). Here we demonstrated that protein

translation is necessary to produce MOR-LTD (Figure 5), but not MOR-STD (Figure

supplement 1G-H). Our data is consistent with the previous reports on protein

translation-dependent eCB-LTD, and suggest that it is presynaptic protein translation

that is involved in MOR-LTD expression, however more data is needed to determine

this definitively.

Although we have previously reported that MOR- and eCB-LTD are mutually occlusive

in the DLS(Atwood, Kupferschmidt, et al., 2014), and here we found that there were

similarities between MOR-LTD and eCB-LTD mechanisms, we found some differences

which make both types of LTD distinct. In the DLS, MOR-LTD of glutamate input is

restricted to those from the AIC(Munoz et al., 2018), which differs from the more broadly

expressed (and overlapping at AIC-DLS synapses, Figure supplement 2) eCB-

LTD(Gremel et al., 2016; Wu et al., 2015), suggesting that these two forms of LTD may

occur in distinct synaptic environments that could utilize differing mechanisms. It is likely

that even in synapses where both receptors are expressed (e.g. in AIC terminals) that

each receptor shares partially overlapping, but not identical signaling pathways.

A clear example of distinct signaling mechanisms identified here is in the role of the

mTOR pathway in eCB-LTD and MOR-LTD. The CB1 receptor can signal via mTOR in

the hippocampus, where the increase of protein translation induced by CB1 receptor

activation is mTOR-dependent producing inhibitory eCB-LTD(Younts et al., 2016). We

predicted that we would see the same effect in the DLS for both eCB-LTD and for MOR-

LTD. Partially correct in our prediction, we found that eCB-LTD is blocked after the

inhibition of mTOR in the DLS (Figure 6A-C), but the mechanism of MOR-LTD did not

utilize mTOR (Figure 6D-F). These novel data regarding eCB-mTOR signaling in DLS

provide rationale for future experiments addressing the role of eCB-mTOR signaling in

dorsal striatal- related behaviors.

Calcium signaling also plays an important role in the mechanism of synaptic

plasticity(Frank, Pielage, & Davis, 2009; Tien & Kerschensteiner, 2018). Postsynaptic

Ca^{2+} influx is needed to produce eCBs (Tanimura et al., 2010), which interacts with

presynaptic mechanism to produce LTD(Adermark, 2011; Adermark, Talani, & Lovinger,

2009; Cristiano Nazzaro et al., 2012; Plotkin et al., 2013; Wang et al., 2006). Some

forms of GPCR-mediated LTD, including eCB-LTD, also require presynaptic calcium

signaling as well(Atwood, Lovinger, et al., 2014). However, we demonstrated that Ca^{2+}

is not involved in MOR-LTD in DLS (Figure supplement 2), indicating another difference

between the mechanisms of MOR-LTD and eCB-LTD.

Presynaptic cations such as Na$^+$ and K$^+$ are important for the propagation of action

potentials and subsequent glutamate release(Chen & Lui, 2021). HCNs channels are

permeable to Na$^+$ and K$^+$ and modulated by cAMP(Sartiani, Mannaioni, Masi, Novella

Romanelli, & Cerbai, 2017), and they can decrease neurotransmission by restricting

presynaptic Ca^{2+} influx(Zhuo Huang et al., 2011). However, while we did not find that

Ca^{2+} plays a role in MOR-LTD, we did identify a critical role for the HCN1 channel in

MOR-LTD expression (Figures 7 and 8). It has been reported that HCN1 channels can

be found at presynaptic sites in the globus pallidus, where the inhibition by ZD7288

produced an increase of miniature inhibitory postsynaptic current frequency, with no

changes in amplitude(Boyes, Bolam, Shigemoto, & Stanford, 2007). The block or

deletion of HCN1 causes an increase in the frequency of miniature EPSCs in entorhinal

cortical pyramidal neurons(Zhuo Huang et al., 2011). Here we showed that broad

blockade of HCN channels enhanced glutamate release, but decreased postsynaptic

responses to glutamate (Figure 7D-H), presumably by acting non-specifically at both

presynaptic and postsynaptic sites. However, when we deleted just HCN1 from AIC

inputs to DLS, we did not observe any changes in basal glutamate transmission (Figure

8H-L). The differences in effects of broad blockade and synapse-specific deletion may

be attributed to the overall contribution of HCN1s at AIC inputs relative to the population

of glutamate synapses. Genetic deletion did not completely ablate MOR-LTD

expression. This could be due to an incomplete AAV vector transduction (most

recordings had a complete loss of LTD) or perhaps that HCN1 contributes to LTD

differently at AIC inputs to different subtypes of MSNs. Therefore, in the future, it will

also be important to investigate the role of HCN1 in MOR-LTD at AIC inputs to D1- or

D2-expressing MSNs (direct or indirect pathway respectively). It has recently been

reported that HCN1 contributes to alcohol preference(Salling & Harrison, 2020). Given

that alcohol disrupts MOR-LTD in DLS, this mechanism of MOR-HCN1 signaling may

be a target of alcohol's effects and contribute to alcohol-related, dorsal striatal-

dependent behaviors such as habitual and compulsive alcohol drinking. As the opioid

oxycodone also disrupts MOR-LTD it will be intriguing to determine the impact of both

oxycodone and alcohol on AIC-DLS HCN1 function. HCN1 may therefore have

therapeutic potential in alcohol and opioid use disorders.

In conclusion, these findings indicate that glutamatergic synapses in the DLS have

distinct mechanisms of plasticity induction mediated by the same types of receptors.

The specific signaling processes that result in these different forms of plasticity might be

the answer to why different types of plasticity are differentially sensitive to drugs of

abuse and alcohol(Atwood, Lovinger, et al., 2014; Munoz et al., 2018; Muñoz et al.,

2020). However, further work is required to resolve what components of these

mechanisms render some synapses susceptible to the deleterious effects of these

drugs while others are resistant. It will also be important to decipher the behavioral

relevance of AIC-DLS synapse HCN1 channels and eCB-mTOR signaling in the DLS.

Methods

Methods details

All experiments were performed similar to our previous dorsal striatal electrophysiological studies with some experiment-specific modifications(Atwood, Kupferschmidt, et al., 2014; Fritz, Munoz, & Atwood, 2019; Fritz, Munoz, Yin, Bauchle, & Atwood, 2018; Munoz et al., 2018). These methods are described in brief below.

Animals and materials

Animal care and experimental protocols for this study were approved by the Institutional Animal Care and Use Committee (IACUC) at the Indiana University School of Medicine and all guidelines for ethical protocols and care of experimental animals established by the NIH (National Institutes of Health, Maryland, USA) were followed. Male C57BL/6J mice were obtained from the Jackson Laboratory (JAX #000664, Bar Harbor, Maine, USA). Emx1Cre-Ai32, VGluT2Cre-Ai32, and HCN1fox mice were bred and genotyped in-house (Original stock strains: Ai32: JAX #012569; Emx1Cre: JAX #005628; VGluT2Cre: JAX #016963; HCN1f/f: JAX #028299). All mutant mice used in these studies were backcrossed to C57BL/6J mice for a minimum of 7 generations. The mice used in these studies were between PND 60-100 at the time of experimentation (with the exception of HCN1-flox AAV-cre-injected mice ~PND 98-154). Animals were group-housed in a standard 12-h light/dark cycle (lights on at 0800) at 50% humidity. Food and water were available *ad libitum*.

Reagents

We used the MOR agonist [D-Ala^2, NMe-Phe^4, Gly-ol^5]-enkephalin (DAMGO; H-2535,

Bachem), GABA$_A$ receptor antagonist picrotoxin (PTX, P1675, Sigma-Aldrich), PKI

(6221, Tocris), KT5720 (1288, Tocris), Forskolin (F3917, Sigma-Aldrich), Cycloheximide

(0970, Tocris), ZD7288 (1000, Tocris), WIN55,212-2 (1038, Tocris), and Torin 2 (4248,

Tocris). Other reagents used for making solutions were purchased from Sigma-Aldrich or

Fisher Scientific.

Brain slice preparation.

Mice were euthanized via decapitation under deep isoflurane anesthesia, and the brain

was quickly excised and placed in an ice-cold cutting solution containing (in mM): 194

sucrose, 30 NaCl, 4.5 KCl, 1 MgCl$_2$, 26 NaHCO$_3$, 1.2 NaH$_2$PO$_4$, 10 Glucose saturated

with a mixture of 95% O$_2$ and 5% CO$_2$, and sliced to a thickness of 280 μm on a

vibratome (Leica VT1200S, Germany). Slices were transferred to an artificial

cerebrospinal fluid (aCSF) solution containing (in mM): 124 NaCl, 4.5 KCl, 1 MgCl$_2$, 26

NaHCO$_3$, 1.2 NaH$_2$PO$_4$, 10 Glucose, 2 CaCl$_2$ (310-320 mOsm) saturated with 95%

O$_2$/5% CO$_2$ at 30°C for 1 hr before being moved to room temperature. When ready for

recording, slices were transferred to a recording chamber continuously perfused with

aCSF solution saturated with 95% O$_2$/5% CO$_2$.

Electrophysiology recordings

Whole-cell recordings of excitatory postsynaptic currents (EPSCs) in medium spiny

neurons (MSNs) were carried out at 29–32 C° and aCSF was continuously perfused at a

rate of 1-2 ml/min. Recordings were performed in the voltage clamp configuration using a

Multiclamp 700B amplifier and a Digidata 1550B (Molecular Devices, San Jose, CA).

Slices were visualized on an Olympus BX51WI microscope (Olympus Corporation of

America). MSNs were identified by their size, membrane resistance, and capacitance. Picrotoxin (50 µM) was added to the aCSF for recordings to isolate excitatory transmission. Patch pipettes were prepared from filament-containing borosilicate micropipettes (World Precision Instruments) using a P-1000 micropipette puller (Sutter Instruments, Novato, CA), having a 2.0-3.5 MΩ resistance. The internal solution contained (in mM): 120 CsMeSO$_3$, 5 NaCl, 10 TEA-Cl, 10 HEPES, 5 lidocaine bromide, 1.1 EGTA, 0.3 Na-GTP and 4 Mg-ATP (pH 7.2 and 290-310 mOsm). MSNs were voltage clamped at -60 mV for the duration of the recordings. To evaluate the effect of Ca^{2+}, aCSF Ca^{2+} was replaced with Sr^{2+} (2 mM) during the application of DAMGO (from 10 to 15 min after the start of recording) or through the entire recording (from 0 to 40 min). For electrically-evoked recordings, a twisted tungsten bipolar stimulating electrode (PlasticsONE, Roanoke, VA) was placed at the border of the white matter of the external capsule. Electrically-evoked excitatory postsynaptic currents (eEPSCs) were generated by a DS3 Isolated Current Stimulator (Digitimer, Ft. Lauderdale, FL) every 20 s and stimulus intensity was adjusted to produce stable eEPSCs of 200–600 pA in amplitude prior to the initiation of experimental recording. The spontaneous EPSC (sEPSC) recordings were made for 2 min and the analysis of frequency (Hz), decay constant (ms), rise constant (ms) and amplitude (pA) were measured to determine the effects of the deletion of HCN1 in AIC. The decay constant of sEPSCs was fitted as single exponential and both rise and decay-phase were fitted between 10 and 90% of the maximal amplitude. Data were acquired using Clampex 10.3 (Molecular Devices, San Jose, CA). Series resistance was monitored and only cells with a stable series resistance (less than 25 MΩ and that did not change more than 15% during recording) were included for data

analysis. Recordings were made 2–7 h after euthanasia.

Viral injections

C57BL/6J mice were anesthetized with isoflurane and stereotaxically-injected with the

adeno-associated viral (AAV) vector, AAV9.hSyn.ChR2(H134R)-eYFP (Addgene

#26973) to drive ChR2 expression in AIC neurons. Bilateral injections were made into

AIC: A/P: +2.4, M/L: ±2.3, D/V: −2.25 (50 nl/injection, 12.5 nl/min infusion rate).

To produce AIC projection neuron HCN1 knockout mice, HCN1-flox mice were

anesthetized with isoflurane and stereotaxically-injected with

AAV9.hSyn.HI.eGFP.Cre.WPRE.SV40 (Addgene #105540) or

AAV9.hSyn.eGFP.WPRE.bGH as control (Addgene #105539). Bilateral injections were

made into AIC at coordinates: A/P: +2.4, M/L: ±2.3, D/V: −2.25 (75 nl/injection, 12.5

nl/min infusion rate). HCN1-flox mice were allowed to recover for at least 4 weeks to

allow for adequate ablation of HCN1 channel expression before brain slices were made

for electrophysiological recordings. Prior to recording, brain slices were imaged via an

Olympus MVX10 microscope (Olympus Corporation of America) to verify GFP or GFP-

tagged cre-recombinase expression in injected HCN1flox mice.

Optogenetic Recordings

AAV-ChR2 injection in C57BL/6J mice was performed to target ChR2 expression to

inputs from AIC to DLS. Optically-evoked EPSCs (oEPSCs) in MSNs from Emx1-Ai32,

VGluT2-Ai32 and C57BL/6J mice injected with AAV-ChR2 were produced in brain slices

using 470-nm blue light (5-ms exposure time) delivered via field illumination through the

microscope objective. Light intensity was adjusted to produce stable oEPSCs of 200–

600-pA amplitude prior to experimental recording. oEPSCs were evoked every 30s. AIC-

DLS eCB-LTD was produced using optical high-frequency stimulation (oHFS) coupled

with depolarization (4 blue light pulses of 50 Hz, 10 s inter-pulse interval). Prior to

recording, brain slices were imaged via an Olympus MVX10 microscope (Olympus

Corporation of America) to verify YFP-tagged ChR2, GFP-tagged cre-recombinase, or

GFP expression in injected C57BL/6J mice.

Quantification and statistical analysis

Sample size

The target number of samples in each group for electrophysiological experiments was

determined from our previously published studies(Atwood, Kupferschmidt, et al., 2014;

Munoz et al., 2018; Muñoz et al., 2020).

Replication

All sample sizes indicated in figures for electrophysiological experiments represent

biological replicates. 1 neuron was recorded per brain slice and all experiments involved

recordings from at least 2 mice.

Data analyses

eEPSCs and oEPSCs were analyzed using Clampfit 10.1 (Molecular Devices, San

Jose, CA) and sEPSCs with MiniAnalysis 6.0 (Synaptosoft Inc). Unless otherwise

indicated, data are presented as the mean ± SEM. The analyses of normally distributed

data were performed using two-tailed unpaired or two-tailed paired Student's t tests

following an F test to confirm similar variances. Non-normally distributed data were

analyzed using two-tailed Welch's t test for unpaired data (Figure 7J-L). Data that were

analyzed using this test are indicated in the figure legends. Statistical analyses were

performed with Prism 9 (GraphPad, La Jolla, CA). The level of significance was set at P

< 0.05 for all analyses. Representative traces are the average baseline EPSC (1-10

min) and average post-treatment EPSC of final 10 min of recording. For VGluT2-Ai32

recordings, representative traces are the average baseline EPSC (1-10 min) average of

the peak of response after DAMGO (15-20 min) and average post-treatment EPSC of

final 10 min of recording Exclusion of individual data points was determined using a

ROUT outlier calculator (Q=1%) included in the Prism 9 software package.

Acknowledgments: The authors thank Fuqin Yin for expert technical assistance. All

reagents and other materials used for this work are commercially available. This work

was supported by NIH/NIAAA grant R01 AA027214.

Competing interests: The authors declare no conflict of interest.

Figures

482
Figure 1. MOR activation induces glutamatergic LTD in the DLS. A) Schematic
representation of coronal brain slice showing the recording of EPSCs evoked by focal
electric stimulation in the DLS of C57BL/6J mice. **B)** Representative electrically-evoked
EPSC traces before and after DAMGO (0.3 μM, 5 min) application. **C)** The activation
of MOR by DAMGO induced glutamatergic LTD in DLS MSNs of C57BL/6J (final
10 min of recording average: 72 ± 5%). **D)** eEPSC amplitudes in MSNs within DLS
were significantly reduced after DAMGO application (0–10 min baseline v. final
10 min of recording; paired t-test, P = 0.0091, t5 = 4.125, n = 6 neurons from 2
mice). **E)** Schematic representation of coronal brain slice showing the recording of
EPSCs evoked by focal optical stimulation (470-nm blue light for 5-ms exposure) in the
DLS of Emx1-Ai32 mice. **F)** Representative optically-evoked EPSC traces before and
after DAMGO (0.3 μM, 5 min) application. **G)** The activation of MOR by DAMGO
induced corticostriatal LTD in DLS MSNs of Emx1-Ai32 (final 10 min of recording
average: 71 ± 5%). **H)** oEPSC amplitudes were significantly reduced after DAMGO
application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.0017,
t7 = 4.9, n = 8 neurons from 3 mice). **I)** Schematic figure of the injection paradigm
showing an AAV vector encoding for ChR2 (AAV.hSyn.ChR2.YFP) in AIC in C57BL/6J
mice, this AAV was injected 2 weeks prior to recordings. Also, the next schematic
representation of coronal brain slice shows the recording of oEPSCs (470-nm blue light
for 5-ms exposure) in the DLS. **J)** Representative AIC-DLS oEPSC traces before and
after DAMGO (0.3 μM, 5 min) application. **K)** DAMGO induced AIC-DLS LTD (final
10 min of recording average: 66 ± 7%). **L)** DAMGO application significantly reduced

oEPSC amplitudes (0–10□min baseline v. final 10□min of recording; paired t-test,
P□=□0.0084, t4□=4.84, n□=□5 neurons from 2 mice). Data represent mean ± SEM.
**P□<□0.01.

Figure 2. Cannabinoid LTD occurs at AIC-DLS synapses. (A) Coronal brain slice showing the AAV-ChR2 infection of AIC and dorsal striatal terminal expression (bar scale□=□1000□µm). **(B)** Representative oEPSC traces showing the effects of pairing optical high frequency stimulation with postsynaptic depression (oHFS) or WIN55,212-2 (1□µM, 10□min) application in brain slices of AAV-ChR2 C57Bl/6J mice. **(C)** Both oHFS and WIN55,212-2 induced AIC-DLS LTD (final 10□min of recording average: oHFS: 63 ± 7%; WIN55,212-2: 68 ± 8%). **(D)** oHFS significantly reduced oEPSC amplitudes (0–10□min baseline v. final 10□min of recording; paired t-test, P□=□0.017, t3=4.78, n□=□4 neurons from 2 mice). E) WIN55,212-2 significantly reduced oEPSC amplitudes (0–10□min baseline v. final 10□min of recording; paired t-test, P□=□0.0038, t7=4.24, n□=□8 neurons from 4 mice). Data represent mean ± SEM. *P□<□0.05, **P□<□0.01.

Figure 3. Inhibition of Presynaptic PKA blocks MOR-mediated LTD. (A) Representative eEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the preincubation of PKA-selective inhibitor, KT5720 (1 μM, ≥1 hr). **(B-C)** The preincubation of KT5720 blocked glutamatergic MOR-LTD (101 ± 6%), the eEPSC amplitudes did not change after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.9487, t8 = 0.0664, n = 9 neurons from 3 mice). **(D)** Representative eEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the intracellular dialysis of PKI (1 μM, ≥30 min). **(E-F)** The inhibition of postsynaptic PKA did not alter MOR-LTD (84 ± 2%). eEPSC amplitudes were reduced after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P < 0.0001, t11 = 6.374, n = 12 neurons from 5 mice). **(G)** Representative oEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the preincubation of PKA-selective inhibitor, KT5720 (1 μM, ≥1 hr). **(H-I)** PKA inhibition blocked corticostriatal MOR-LTD (90 ± 5%), with no changes in oEPSC amplitudes after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.162, t8 = 1.541, n = 9 neurons from 4 mice). **(J)** Representative oEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the intracellular dialysis of PKI (1 μM, ≥30 min). **(K-L)** The inhibition of postsynaptic PKA did not alter MOR-LTD (54 ± 9%). oEPSC amplitudes were reduced after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.0014, t6 = 5.588, n = 7 neurons from 2 mice). **(M)** Representative AIC-DLS oEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the preincubation of PKA-selective inhibitor, KT5720 (1 μM, ≥1 hr). **(N-O)** KT5720 blocked corticostriatal MOR-LTD (96 ± 4%), with no changes in oEPSC amplitudes after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.2121, t9 = 1.343, n = 10 neurons from 3 mice). **(P)** Representative AIC-DLS oEPSC traces showing the effects of DAMGO (0.3 μM, 5 min) application after the intracellular dialysis of PKI (1 μM, ≥30 min). **(Q-R)** Postsynaptic PKA inhibition did not alter specific AIC MOR-mediated LTD (62 ± 6%). oEPSC amplitudes were reduced after DAMGO application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.0026, t5 = 5.576, n = 6 neurons from 3 mice). Data represent mean ± SEM. ns= no significant, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 4. The activation of adenylyl cyclase disrupts MOR-mediated LTD. (A) Representative eEPSC traces showing the effects of 20 μM forskolin before, during and after DAMGO (0.3 μM, 5 min) application. **(B-C)** AC activation disrupted glutamatergic MOR-LTD (105 ± 4%), showing no changes in eEPSC amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.2681, t8 =1.19, n = 9 neurons from 4 mice). **(D)** Representative oEPSC traces showing the effects of 20 μM forskolin before, during and after DAMGO (0.3 μM, 5 min) application in brain slices from Emx1-Ai32 mice. **(E-F)** AC activation disrupted corticostriatal MOR-LTD (102 ± 3%), with no effects in oEPSC amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.4031, t5 =0.913, n = 6 neurons from 3 mice). **(G)** Representative AIC-DLS oEPSC traces showing the effects of 20 μM forskolin before, during and after DAMGO (0.3 μM, 5 min) application. **(H-I)** AC activation blocked specifically AIC MOR-LTD (105 ± 4%), without changes in oEPSC amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.379, t5 =0.964, n = 6 neurons from 2 mice). Data represent mean ± SEM. ns= no significant.

Figure 5. Protein translation is required for MOR-mediated LTD expression. (A) Representative eEPSC traces showing the effects of 80 μM cycloheximide before, during and after DAMGO (0.3 μM, 5 min) application. **(B-C)** Protein translation inhibition disrupted glutamatergic MOR-LTD (104 ± 9%), showing no changes in eEPSC amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.7006, t8 =0.398, n = 9 neurons from 4 mice). **(D)** Representative oEPSC traces showing the effects of 80 μM cycloheximide before, during and after DAMGO (0.3 μM, 5 min) application in brain slices from Emx1-Ai32 mice. **(E-F)** Protein translation inhibition blocked corticostriatal MOR-LTD (105 ± 9%) with no effects in oEPSC amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.4992, t7 =0.713, n = 8 neurons from 3 mice). **(G)** Representative AIC-DLS oEPSC traces showing the effects of 80 μM cycloheximide before, during and after DAMGO (0.3 μM, 5 min) application. **(H-I)** Protein translation is needed to induce MOR-LTD (91 ± 2%) from AIC inputs. DAMGO did not change in oEPSC amplitude (0–10 min baseline v. final 10 min of recording; paired t-test, P = 0.1208, t5 =1.868, n = 6 neurons from 2 mice). Data represent mean ± SEM. ns= no significant.

593
Figure 6. mTOR pathway is involve in eCB-LTD but not in MOR-LTD in the DLS.
(A) Representative eEPSC traces before and after high-frequency stimulation (HFS)
coupled with depolarization (4 pulses of 100□Hz, 10□s inter-pulse interval), showing
the effects of torin 2 (100 nM) bath application in brain slices of C57BL/J. **(B-C)** mTOR
inhibition blocked eCB-LTD in DLS (0–10□min baseline v. final 10□min of recording;
Control: paired t-test, P□=□0.0002, t7□=7.08, n□= 8 neurons from 3 mice; torin 2:
paired t-test, P□=□0.9009, t7□=0.129, n□= 8 neurons from 3 mice). **(D)** Representative
eEPSC traces before and after DAMGO bath application (0.3□µM, 5□min), showing the
effects of torin 2 (100 nM) bath application in brain slices of C57BL/J. **(E-F)** mTOR
inhibition did not affect the expression of MOR-LTD in DLS (0–10□min baseline v. final
10□min of recording; paired t-test, P□<□0.0001, t7□=8.11, n□= 8 neurons from 3
mice). Data represent mean ± SEM. ***P□<□0.001, ****P□<□0.0001.
606

Figure 7. Inhibition of HCN channels disrupts MOR-mediated LTD. (A)
Representative eEPSC traces showing the effects of 25 μM ZD7288 before, during and
after DAMGO (0.3 μM, 5 min) application. **(B-C)** HCN channel inhibition disrupted
glutamatergic MOR-LTD (104 ± 9%) in the DLS, showing no changes in eEPSC
amplitude after DAMGO bath application (0–10 min baseline v. final 10 min of
recording; paired t-test, P = 0.8008, t7 =0.2621, n= 8 neurons from 5 mice). **(D)**
Representative synaptic current traces from C57BL/6J mice showing the spontaneous
excitatory postsynaptic currents (sEPSC) in absence (black trace) or presence of 25 μM
ZD7288 (gray trace, 15 min) application. **(E-H)** HCN channel inhibition increased
frequency (paired t-test, P = 0.043, t14 =2.223, n= 15 neurons from 2 mice) and
decreased amplitude (paired t-test, P = 0.0005, t14 =4.49, n= 15 neurons from 2
mice), without changes in rise (paired t-test, P = 0.542, t14 =0.625, n= 15 neurons
from 2 mice) and decay times (paired t-test, P = 0.324, t14 =1.023, n= 15 neurons
from 2 mice) of the sEPSC events in DLS. Data represent mean ± SEM. ns= no
significant, *P < 0.05, ***P < 0.001.

Figure 8. Genetic ablation of HCN1 channels reduces MOR-mediated LTD expression. (A) Schematic figure of coronal brain slice showing injection paradigm enabling knock out of HCN1 channel expression in AIC in HCN1-flox mice. An AAV vector encoding for either cre-recombinase (AAV.hSyn.cre) or eGFP (AAV.hSyn.eGFP) was injected at least 4 weeks prior to recordings. **(B)** Coronal brain slice showing the AAV-cre or AAV-GFP infection of AIC and dorsal striatal terminal expression (bar scale = 1000 µm). **(C)** Representative eEPSC traces showing the effects of DAMGO (0.3 µM, 5 min) application in brain slices of AAV-injected HCN1-flox mice. **(D-E)** The conditional KO of HCN1 channel from AIC, blunted MOR-LTD in DLS (cre: 90 ± 4% vs GFP: 76 ± 3%, Unpaired t-test, P=0.017, t17=2.64, n=10 neurons from 5 cre-injected mice and n=9 neurons from 3 GFP-injected mice). **(F)** HCN1 deletion did not fully disrupt MOR-LTD, because there was still a small, but significant reduction of eEPSC amplitude after DAMGO application (0–10 min baseline v. final 10 min of recording;

paired t-test, $P = 0.011$, $t9 = 3.18$, $n = 10$ neurons from 5 mice). **(G)** Control GFP-
injected mice showed normal decrease of eEPSC amplitude after DAMGO application
0–10 min baseline v. final 10 min of recording; paired t-test, $P < 0.0001$, $t8 = 7.51$,
$n = 9$ neurons from 3 mice). **(H)** Representative synaptic current traces from HCN1-flox
mice showing the spontaneous excitatory postsynaptic currents (sEPSCs) in AAV-cre
(orange trace) and AAV-GFP (green trace) injected mice. **(I-L)** HCN1 channel ablation
from AIC synapses did not affect frequency (cre: 2.6 ± 0.4 Hz vs GFP: 2.7 ± 0.3 Hz,
Unpaired t-test, $P=0.886$, $t19=0.146$, $n=10$ neurons from 3 cre-injected mice and $n=11$
neurons from 2 GFP-injected mice), amplitude (cre: 18.2 ± 1.2 pA vs GFP: 18.3 ± 0.6
pA, Unpaired t-test with Welch's correction, $P=0.957$, $t13=0.055$, $n=10$ neurons from 3
cre-injected mice and $n=11$ neurons from 2 GFP-injected mice), rise time (cre: $1.23 \pm$
0.11 ms vs GFP: 0.98 ± 0.04 ms, Unpaired t-test with Welch's correction, $P=0.0501$,
$t11=2.19$, $n=10$ neurons from 3 cre-injected mice and $n=11$ neurons from 2 GFP-
injected mice) and decay time (cre: 5.25 ± 0.43 ms vs GFP: 4.36 ± 0.14 ms, Unpaired t-
test with Welch's correction, $P=0.075$, $t11=1.97$, $n=10$ neurons from 3 cre-injected mice
and $n=11$ neurons from 2 GFP-injected mice) of the sEPSC events in DLS. Data
represent mean ± SEM. ns= no significant, $*P < 0.05$, $****P < 0.0001$.

References

Adermark, L. (2011). Modulation of endocannabinoid-mediated long-lasting disinhibition of striatal output by cholinergic interneurons. *Neuropharmacology, 61*(8), 1314-1320. doi:https://doi.org/10.1016/j.neuropharm.2011.07.039

Adermark, L., Talani, G., & Lovinger, D. M. (2009). Endocannabinoid-dependent plasticity at GABAergic and glutamatergic synapses in the striatum is regulated by synaptic activity. *The European journal of neuroscience, 29*(1), 32-41. doi:10.1111/j.1460-9568.2008.06551.x

Atwood, B. K., Kupferschmidt, D. A., & Lovinger, D. M. (2014). Opioids induce dissociable forms of long-term depression of excitatory inputs to the dorsal striatum. *Nat Neurosci, 17*(4), 540-548. doi:10.1038/nn.3652

Atwood, B. K., Lovinger, D. M., & Mathur, B. N. (2014). Presynaptic long-term depression mediated by Gi/o-coupled receptors. *Trends Neurosci, 37*(11), 663-673. doi:10.1016/j.tins.2014.07.010

Bao, G., Kang, L., Li, H., Li, Y., Pu, L., Xia, P., . . . Pei, G. (2007). Morphine and heroin differentially modulate in vivo hippocampal LTP in opiate-dependent rat. *Neuropsychopharmacology, 32*(8), 1738-1749. doi:10.1038/sj.npp.1301308

Boyes, J., Bolam, J. P., Shigemoto, R., & Stanford, I. M. (2007). Functional presynaptic HCN channels in the rat globus pallidus. *European Journal of Neuroscience, 25*(7), 2081-2092. doi:https://doi.org/10.1111/j.1460-9568.2007.05463.x

Burton, A. C., Nakamura, K., & Roesch, M. R. (2015). From ventral-medial to dorsal-lateral striatum: neural correlates of reward-guided decision-making. *Neurobiol Learn Mem, 117*, 51-59. doi:10.1016/j.nlm.2014.05.003

Chen, I., & Lui, F. (2021). Neuroanatomy, Neuron Action Potential. In *StatPearls*. Treasure Island (FL): StatPearls Publishing

Copyright © 2021, StatPearls Publishing LLC.

Cho, H. S., Lee, H. H., Choi, S. J., Kim, K. J., Jeun, S. H., Li, Q.-Z., & Sung, K.-W. (2008). Forskolin Enhances Synaptic Transmission in Rat Dorsal Striatum through NMDA Receptors and PKA in Different Phases. *kjpp, 12*(6), 293-297. doi:10.4196/kjpp.2008.12.6.293

Corbit, L. H., & Janak, P. H. (2016). Habitual Alcohol Seeking: Neural Bases and Possible Relations to Alcohol Use Disorders. *Alcoholism: Clinical and Experimental Research, 40*(7), 1380-1389. doi:10.1111/acer.13094

Dacher, M., & Nugent, F. S. (2011a). Morphine-induced modulation of LTD at GABAergic synapses in the ventral tegmental area. *Neuropharmacology, 61*(7), 1166-1171. doi:10.1016/j.neuropharm.2010.11.012

Dacher, M., & Nugent, F. S. (2011b). Opiates and plasticity. *Neuropharmacology, 61*(7), 1088-1096. doi:10.1016/j.neuropharm.2011.01.028

DePoy, L., Daut, R., Brigman, J. L., MacPherson, K., Crowley, N., Gunduz-Cinar, O., . . . Holmes, A. (2013). Chronic alcohol produces neuroadaptations to prime dorsal striatal learning. *Proc Natl Acad Sci U S A, 110*(36), 14783-14788. doi:10.1073/pnas.1308198110

Drake, C. T., Chavkin, C., & Milner, T. A. (2007). Opioid systems in the dentate gyrus. *Prog Brain Res, 163*, 245-263. doi:10.1016/s0079-6123(07)63015-5

Everitt, B. J., & Robbins, T. W. (2016). Drug Addiction: Updating Actions to Habits to Compulsions Ten Years On. *Annual Review of Psychology, 67*(1), 23-50. doi:10.1146/annurev-psych-122414-033457

Frank, C. A., Pielage, J., & Davis, G. W. (2009). A presynaptic homeostatic signaling system composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels. *Neuron, 61*(4), 556-569. doi:10.1016/j.neuron.2008.12.028

Fritz, B. M., Munoz, B., & Atwood, B. K. (2019). Genetic Selection for Alcohol Preference in Mice Alters
Dorsal Striatum Neurotransmission. *Alcohol Clin Exp Res, 43*(11), 2312-2321.
doi:10.1111/acer.14187
Fritz, B. M., Munoz, B., Yin, F., Bauchle, C., & Atwood, B. K. (2018). A High-fat, High-sugar 'Western' Diet
Alters Dorsal Striatal Glutamate, Opioid, and Dopamine Transmission in Mice. *Neuroscience,
372*, 1-15. doi:10.1016/j.neuroscience.2017.12.036
Gremel, C. M., Chancey, J. H., Atwood, B. K., Luo, G., Neve, R., Ramakrishnan, C., . . . Costa, R. M. (2016).
Endocannabinoid Modulation of Orbitostriatal Circuits Gates Habit Formation. *Neuron, 90*(6),
1312-1324. doi:10.1016/j.neuron.2016.04.043
Hawes, S. L., Salinas, A. G., Lovinger, D. M., & Blackwell, K. T. (2017). Long-term plasticity of
corticostriatal synapses is modulated by pathway-specific co-release of opioids through κ-opioid
receptors. *J Physiol, 595*(16), 5637-5652. doi:10.1113/jp274190
He, C., Chen, F., Li, B., & Hu, Z. (2014). Neurophysiology of HCN channels: from cellular functions to
multiple regulations. *Prog Neurobiol, 112*, 1-23. doi:10.1016/j.pneurobio.2013.10.001
Hilário, M. R. F., & Costa, R. M. (2008). High on habits. *Frontiers in neuroscience, 2*(2), 208-217.
doi:10.3389/neuro.01.030.2008
Hopf, F. W., & Lesscher, H. M. B. (2014). Rodent models for compulsive alcohol intake. *Alcohol, 48*(3),
253-264. doi:https://doi.org/10.1016/j.alcohol.2014.03.001
Huang, Z., Li, G., Aguado, C., Lujan, R., & Shah, M. M. (2017). HCN1 channels reduce the rate of
exocytosis from a subset of cortical synaptic terminals. *Sci Rep, 7*, 40257. doi:10.1038/srep40257
Huang, Z., Lujan, R., Kadurin, I., Uebele, V. N., Renger, J. J., Dolphin, A. C., & Shah, M. M. (2011).
Presynaptic HCN1 channels regulate CaV3.2 activity and neurotransmission at select cortical
synapses. *Nature Neuroscience, 14*(4), 478-486. doi:10.1038/nn.2757
Iremonger, K. J., & Bains, J. S. (2009). Retrograde opioid signaling regulates glutamatergic transmission
in the hypothalamus. *J Neurosci, 29*(22), 7349-7358. doi:10.1523/jneurosci.0381-09.2009
Le Merrer, J., Becker, J. A. J., Befort, K., & Kieffer, B. L. (2009). Reward processing by the opioid system in
the brain. *Physiological reviews, 89*(4), 1379-1412. doi:10.1152/physrev.00005.2009
Lonart, G., Schoch, S., Kaeser, P. S., Larkin, C. J., Südhof, T. C., & Linden, D. J. (2003). Phosphorylation of
RIM1alpha by PKA triggers presynaptic long-term potentiation at cerebellar parallel fiber
synapses. *Cell, 115*(1), 49-60. doi:10.1016/s0092-8674(03)00727-x
Lovinger, D. M. (2010). Neurotransmitter roles in synaptic modulation, plasticity and learning in the
dorsal striatum. *Neuropharmacology, 58*(7), 951-961. doi:10.1016/j.neuropharm.2010.01.008
Mato, S., Lafourcade, M., Robbe, D., Bakiri, Y., & Manzoni, O. J. (2008). Role of the cyclic-AMP/PKA
cascade and of P/Q-type Ca++ channels in endocannabinoid-mediated long-term depression in
the nucleus accumbens. *Neuropharmacology, 54*(1), 87-94.
doi:10.1016/j.neuropharm.2007.04.014
Moosmang, S., Biel, M., Hofmann, F., & Ludwig, A. (1999). Differential distribution of four
hyperpolarization-activated cation channels in mouse brain. *Biol Chem, 380*(7-8), 975-980.
doi:10.1515/BC.1999.121
Munoz, B., Fritz, B. M., Yin, F., & Atwood, B. K. (2018). Alcohol exposure disrupts mu opioid receptor-
mediated long-term depression at insular cortex inputs to dorsolateral striatum. *Nat Commun,
9*(1), 1318. doi:10.1038/s41467-018-03683-1
Muñoz, B., Haggerty, D. L., & Atwood, B. K. (2020). Synapse-specific expression of mu opioid receptor
long-term depression in the dorsomedial striatum. *Scientific Reports, 10*(1), 7234.
doi:10.1038/s41598-020-64203-0
Nazzaro, C., Greco, B., Cerovic, M., Baxter, P., Rubino, T., Trusel, M., . . . Tonini, R. (2012). SK channel
modulation rescues striatal plasticity and control over habit in cannabinoid tolerance. *Nat
Neurosci, 15*(2), 284-293. doi:10.1038/nn.3022

Nazzaro, C., Greco, B., Cerovic, M., Baxter, P., Rubino, T., Trusel, M., . . . Tonini, R. (2012). SK channel modulation rescues striatal plasticity and control over habit in cannabinoid tolerance. *Nature Neuroscience, 15*(2), 284-293. doi:10.1038/nn.3022

O'Tousa, D., & Grahame, N. (2014). Habit formation: Implications for alcoholism research. *Alcohol, 48*(4), 327-335. doi:https://doi.org/10.1016/j.alcohol.2014.02.004

Patton, M. H., Roberts, B. M., Lovinger, D. M., & Mathur, B. N. (2016). Ethanol Disinhibits Dorsolateral Striatal Medium Spiny Neurons Through Activation of A Presynaptic Delta Opioid Receptor. *Neuropsychopharmacology, 41*(7), 1831-1840. doi:10.1038/npp.2015.353

Plotkin, J. L., Shen, W., Rafalovich, I., Sebel, L. E., Day, M., Chan, C. S., & Surmeier, D. J. (2013). Regulation of dendritic calcium release in striatal spiny projection neurons. *J Neurophysiol, 110*(10), 2325-2336. doi:10.1152/jn.00422.2013

Postea, O., & Biel, M. (2011). Exploring HCN channels as novel drug targets. *Nat Rev Drug Discov, 10*(12), 903-914. doi:10.1038/nrd3576

Renteria, R., Cazares, C., & Gremel, C. M. (2020). Habitual Ethanol Seeking and Licking Microstructure of Enhanced Ethanol Self-Administration in Ethanol-Dependent Mice. *Alcoholism, clinical and experimental research, 44*(4), 880-891. doi:10.1111/acer.14302

Salling, M. C., & Harrison, N. L. (2020). Constitutive Genetic Deletion of Hcn1 Increases Alcohol Preference during Adolescence. *Brain Sci, 10*(11). doi:10.3390/brainsci10110763

Sartiani, L., Mannaioni, G., Masi, A., Novella Romanelli, M., & Cerbai, E. (2017). The Hyperpolarization-Activated Cyclic Nucleotide–Gated Channels: from Biophysics to Pharmacology of a Unique Family of Ion Channels. *Pharmacological Reviews, 69*(4), 354-395. doi:10.1124/pr.117.014035

Shen, F., Jin, S., Duan, Y., Liang, J., Zhang, M., Jiang, F., & Sui, N. (2018). Distinctive Changes of L-Type Calcium Channels and Dopamine Receptors in the Dorsomedial and Dorsolateral Striatum after the Expression of Habitual Cocaine-Seeking Behavior in Rats. *Neuroscience, 370*, 139-147. doi:10.1016/j.neuroscience.2017.07.049

Tanimura, A., Yamazaki, M., Hashimotodani, Y., Uchigashima, M., Kawata, S., Abe, M., . . . Kano, M. (2010). The Endocannabinoid 2-Arachidonoylglycerol Produced by Diacylglycerol Lipase α Mediates Retrograde Suppression of Synaptic Transmission. *Neuron, 65*(3), 320-327. doi:https://doi.org/10.1016/j.neuron.2010.01.021

Tien, N.-W., & Kerschensteiner, D. (2018). Homeostatic plasticity in neural development. *Neural Development, 13*(1), 9. doi:10.1186/s13064-018-0105-x

Wang, Z., Kai, L., Day, M., Ronesi, J., Yin, H. H., Ding, J., . . . Surmeier, D. J. (2006). Dopaminergic Control of Corticostriatal Long-Term Synaptic Depression in Medium Spiny Neurons Is Mediated by Cholinergic Interneurons. *Neuron, 50*(3), 443-452. doi:https://doi.org/10.1016/j.neuron.2006.04.010

Wu, Y. W., Kim, J. I., Tawfik, V. L., Lalchandani, R. R., Scherrer, G., & Ding, J. B. (2015). Input- and cell-type-specific endocannabinoid-dependent LTD in the striatum. *Cell reports, 10*(1), 75-87. doi:10.1016/j.celrep.2014.12.005

Yasuda, H., Huang, Y., & Tsumoto, T. (2008). Regulation of excitability and plasticity by endocannabinoids and PKA in developing hippocampus. *Proceedings of the National Academy of Sciences, 105*(8), 3106-3111. doi:10.1073/pnas.0708349105

Yin, H. H., Davis, M. I., Ronesi, J. A., & Lovinger, D. M. (2006). The Role of Protein Synthesis in Striatal Long-Term Depression. *The Journal of Neuroscience, 26*(46), 11811-11820. doi:10.1523/jneurosci.3196-06.2006

Younts, T. J., Monday, H. R., Dudok, B., Klein, M. E., Jordan, B. A., Katona, I., & Castillo, P. E. (2016). Presynaptic Protein Synthesis Is Required for Long-Term Plasticity of GABA Release. *Neuron, 92*(2), 479-492. doi:10.1016/j.neuron.2016.09.040