Title

Targeting current species ranges and carbon stocks fails to conserve biodiversity in a changing

climate: opportunities to support climate adaptation under 30x30

Authors

Lindsay M. Dreiss[1]*, L. Mae Lacey[1], Theodore C. Weber[2], Aimee Delach[2], Talia E. Niederman[1],

Jacob W. Malcom[1]

Affiliations

[1]Center for Conservation Innovation, Defenders of Wildlife, Washington, DC 20036, USA.

[2]Department of Landscape Conservation, Defenders of Wildlife, Washington, DC 20036, USA.

*Correspondence

Lindsay Dreiss, Defenders of Wildlife 1130 17th Street NW, Washington, DC 20036

lrosa@defenders.org, 202.772.3201

ABSTRACT

Protecting areas for climate adaptation will be essential to ensuring greater opportunity for

species conservation well into the future. However, many proposals for protected areas

expansion focus on our understanding of current spatial patterns, which may be ineffective

surrogates for future needs. A science-driven call to address the biodiversity and climate crises

by conserving at least 30% of lands and waters by 2030, 30x30, presents new opportunities to

inform the siting of new protections globally and in the U.S. Here we identify climate refugia

and corridors based on a weighted combination of currently available models; compare them to

current biodiversity hotspots and carbon-rich areas to understand how 30x30 protections siting

may be biased by data omission; and compare identified refugia and corridors to the Protected

Areas Database to assess current levels of protection. Available data indicate that 20.5% and

27.5% of identified climate adaptation areas (refugia and/or corridor) coincides with current

imperiled species hotspots and carbon-rich areas, respectively. With only 12.5% of climate

refugia and corridors protected, a continued focus on current spatial patterns in species and

carbon richness will not inherently conserve places critical for climate adaptation. However,

there is ample opportunity for establishing future-minded protections: 52% of the contiguous

U.S. falls into the top quartile of values for at least one class of climate refugia. Nearly 27% is

already part of the protected areas network but managed for multiple uses that may limit their

ability to contribute to the goals of 30x30. Additionally, nearly two-thirds of nationally identified

refugia coincide with ecoregion-specific refugia suggesting representation of nearly all

ecoregions in national efforts focused on conserving climate refugia. Based on these results, we

recommend that land planners and managers make more explicit policy priorities and strategic

decisions for future-minded protections and climate adaptation.

Keywords: Climate refugia, Climate corridors, Protected areas, Biodiversity conservation,

Carbon mitigation

INTRODUCTION

The spatial heterogeneity of shifting climatic conditions presents challenges and opportunities for large-scale biodiversity conservation, as impacts to habitat and species can vary significantly across the landscape (Baldwin et al. 2018). In North America, nearly half of species are already undergoing local extinctions (Wiens 2016), which are partially due to spatially variable changes in temperature and precipitation (Román-Palacios and Wiens 2020). In the contiguous U.S. (CONUS), the average annual temperature has risen 1.2-1.8 °C since the beginning of the 20th century (Vose et al. 2017) and precipitation patterns have shifted with large reductions in the Southeast and West (Fei et al. 2017, Wuebbles et al. 2017). As the effects of climate change accelerate, local biodiversity will either need to find locations that serve as refugia from extreme or rapid climatic changes or shift their ranges to better-suited habitat (Neilson et al. 2005, Keppel and Wardell-Johnson 2012, Franks and Hoffman 2012, Román-Palacios and Wiens 2020). Identifying and conserving important refugia habitats and dispersal routes will be one critical step in jointly addressing the biodiversity and climate crises (Pörtner et al. 2021). While expansion of the U.S. protected areas network has been identified as an important solution to lowering extinction risk and overall ecosystem degradation (Stolton et al. 2015, Gray et al. 2016, Dinerstein et al. 2017, 2019), efforts generally focus on current species distributions and may not effectively reflect future needs (Elsen et al. 2020, Maxwell et al. 2020).

Calls to address the joint biodiversity and climate crises by protecting at least 30% of Earth by 2030, known as "30x30" (Dinerstein et al. 2019), have been endorsed by government and conservation leaders at global (United Nations 2020), national (Biden 2021, U.S. DOI et al. 2021), and state levels (e.g., Newsom 2020). While the specifics have yet to be established (Büscher et al. 2016, Rights and Resources Initiative 2020, Simmons et al. 2021), efforts would hypothetically conserve areas needed to sustain essential ecological services and reverse extinction trends (Locke 2013, Dinerstein et al. 2017). Translating these commitments into national policy may prove challenging since the current distribution of protected areas excludes many areas important for biodiversity conservation (Scott et al. 2001, Jenkins et al. 2015, Venter et al. 2018) and carbon storage (Buotte et al. 2019, Melillo et al. 2015). While not all areas storing large amounts of carbon are also biodiversity hotspots, there are many instances of overlap; for instance, carbon-rich forests can provide important habitat and climate change

buffers (Berkessy and Wintle 2008, Strassburg et al. 2010). However, it is unclear how well the

current network and 30x30 goals can ensure the conservation of climate-resilient habitat in the

coming decades as climate change continues to accelerate.

Climate-resilient habitat can largely be delineated into refugia and corridors. Refugia are

areas where species, natural communities, or ecosystems can persist within a larger area that has

been rendered inhospitable. Such areas are relatively buffered from changes in regional

environmental conditions, and allow organisms to persist as long as local conditions remain

tolerable (Keppel and Wardell-Johnson, 2012; Morelli et al., 2016). Climate-change refugia may

only be temporary for a given species (Morelli et al., 2020), but can delay ecosystem transitions

for decades or longer (Krawchuk et al., 2020). They can serve as a "slow lane," protecting native

species and ecosystems from negative effects of climate change in the short term, providing

longer-term havens for overall biodiversity and ecosystem function, and reducing the risk of

extinction (Morelli et al., 2020). Although refugia can be examined over a continuum of spatial

scales, they are usually classified as either macrorefugia or microrefugia (Ashcroft, 2010).

Macrorefugia are identified at coarse scales, using global climate data or models, and are large

enough to maintain viable animal or plant populations. Microrefugia are identified at local

scales. For example, steep canyons and north-facing slopes are sheltered from solar radiation and

heat accumulation (Stralberg et al., 2020a) and wet areas can remain moist during droughts

(Morelli et al., 2016; Stralberg et al., 2020a).

At broad scales, refugia can be identified by various approaches, such as topodiversity,

climate exposure, and climate tracking (Michalak et al. 2020). Topodiversity models are based

on physical habitat data and highlight regions with varied land cover, climate, soil, and

topographic conditions (Ackerly et al. 2010, Groves et al. 2012, Carroll et al. 2018).

Topographically varying areas can contain features like deep valley bottoms or shaded slopes

that serve as microrefugia (Ashcroft 2010). Climatic exposure models are based on projected

climatic changes and represent the degree of climate change likely to be experienced by a species

or locale (Saxon 2011, Groves et al. 2012). Lastly, climate tracking models are based on

representative climate models and measure the proximity and accessibility of future suitable

climatic conditions (Hamann et al. 2015, Michalak et al. 2018).

However, to survive in the face of ongoing and worsening climate change impacts, some

species may need to relocate (Román-Palacios and Wiens 2020). Climate corridors can facilitate

long-distance movement to more suitable habitat if rising temperatures and other changing

conditions exceed an organism's tolerance (Stralberg et al., 2020b). Depending on model inputs,

corridors may emphasize movement toward cooler latitudes and topographies, along rivers and

streams, and/or through areas providing better habitat and less stress from disturbances (McGuire

et al. 2016, Stralberg 2020b, Carroll et al. 2018, Littlefield et al. 2017). For example, Parks et al.

(2020) found that human land modifications decrease climate connectivity.

Given the urgency of the biodiversity and climate crises, there is a pressing need to

include potential climate refugia and corridors in the conservation planning process. However,

some challenges exist. First, a growing body of available spatial data for identifying areas

important for climate adaptation means that planners must reconcile a diversity of data that

represent different mechanisms and priorities (Michalak et al. 2020, Carroll and Ray 2021).

Second, most conservation prioritization frameworks focus on the current state of species and

environments (Cushman et al. 2009, Lookingbill et al. 2010, Dickson et al. 2013, Belote et al.

2016, McClure et al. 2016) which may result in critical omissions in protected areas siting for

longer-term persistence of some target species (Monzón et al. 2011, Elsen et al. 2020). If this is

the case, consideration of future conditions may complement efforts to preserve current

biodiversity and ecosystem service hotspots, thereby reducing the threat of mass extinctions and

accompanying biosphere degradation. Last, other omissions of important local refugia may occur

if areas for climate resilience are identified and prioritized at a national scale (e.g. Kraus and

Hebb 2020). While this may identify the most important nationwide climate refugia, it could

favor some regions (e.g., mountain ranges) and omit others entirely. Evaluating at smaller scales

(e.g., stratifying by ecoregion) could include more locally important refugia. Taking additional

steps to identify refugia at multiple scales may help increase ecosystem representation and

protections for the unique species assemblages and services they harbor.

Proper identification, protection, and management of climate-informed refugia and

corridors are essential to ensuring greater opportunity for species conservation via migration and

adaptation. While previous research and policy discussion surrounding the extent and

distribution of protected land and water has identified areas important to conserving the current

state of biodiversity and natural carbon storage (Scott et al. 2001, Myers et al. 2000, Gray et al.

2016, Buotte et al. 2020), less attention has been paid to areas important for wildlife and plant
climate adaptation. To help close this knowledge gap, we:

1. identify areas in the contiguous U.S. critical to climate adaptation based on
   coincidence and complementarity among refugia (national and ecoregion-specific)
   and corridors models;
2. compare the spatial distribution of identified climate refugia and corridors with
   current biodiverse and carbon-rich areas; and
3. quantify the extent to which climate refugia and corridors are considered protected.

Given the opportunity to focus land conservation using a science-driven, ecologically-
meaningful (rather than opportunistic) approach, step #2 guides our understanding of how
protections siting under the 30x30 framework may be biased by data omission, and step #3 helps
to assess the current level of protection for identified climate refugia and distinguish where
stronger management might be needed. Our research contributes to a growing literature
demonstrating the importance of incorporating climate-informed data in place-based land
protection policy and practices and helping to identify specific areas for conservation. While
these analyses are not meant to serve as a map of priority lands for conservation, they help guide
discussion on operationalizing 30x30 for strategic, future-minded conservation decisions.

METHODS

We focus on spatial datasets based on climate models or topography to identify areas that
could serve as important refugia or migration routes for the contiguous U.S. (CONUS; Table 1).
All datasets using climate models are informed by an ensemble of three to seven General
Circulation Models (GCMs) for emission scenario Representative Concentration Pathway (RCP)
4.5 and projected for the time period 2071-2100 based on the Coupled Model Intercomparison
Project phase 5 (CMIP5) models. All datasets have been resampled and aligned at 1km
resolution. We combined datasets for refugia (n = 8) and corridors (n = 2) separately, accounting
for differences in underlying mechanisms in modeling method and landscape conservation
principles.

Climate refugia

We initially analyzed relationships between datasets through a principal components
analysis where each component helps define a refugia class. The principal components spatial
analyst tool in ArcPro transforms the data in the input raster bands to compress data and
eliminate redundancy. The result is a multiband raster and text output containing a covariance
matrix, correlation matrix, eigenvalues and eigenvectors. The number of principal components
selected was based on scree plot and broken stick model. All datasets were normalized to a scale
of 0 to 1 by calculating z-scores for all eight refugia datasets representing climate-, landscape-,
and species-based metrics (Table 1), which were then used as inputs for the principal
components analysis.

As with principal components, datasets were assigned to a class based on the sign and
size of the eigenvector. However, to avoid a tradeoff in refugia identification within a single
class, all datasets within the class were required to load together and in the same direction on a
principal component. In addition to presenting three separate classes that represented
topodiversity, climatic stability, and tree macrorefugia, we weighted all eight normalized datasets
such that each dataset contributed equally to its class and each class contributed equally to the
final pixel value. Unlike existing studies in this area (e.g., Michalak et al. 2020), this approach
groups datasets together by relationship and reduces the chances of over-emphasizing any one
variable in identifying refugia. Based on the relationships between refugia datasets, the weighted
combination was calculated as follows, where each variable in the equation represents the
normalized spatial dataset:

$Combined\ Refugia$

$= Z_{Bird\ Macrorefugia} + Z_{Current\ Climate\ Diversity} + Z_{Ecotypic\ Diversity}$

$+ Z_{Land\ Facet\ Diversity} + Z_{Landscape\ Diversity} + \left(Z_{Climatic\ Dissimilatiry} * 2.5\right)$

$+ \left(Z_{Climate\ Velocity} * 2.5\right) + \left(Z_{Tree\ Macrorefugia} * 5\right)$

We analyzed locations in the 80[th] percentile (i.e., the top 20% of values) of the distribution of
values for the combined data and for each refugia class separately. Additionally, we quantified
the degree of overlap in refugia classes.

In addition to CONUS-level analyses, we extracted refugia values for each ecoregion
separately (EPA level II; EPA 2006), classifying the locations that fell into the top 20% of the
distribution as areas of interest. The result was a map of ecoregion-specific refugia, ensuring

equal representation of all ecoregions relative to size. Results from the national- and ecosystem-

scale analyses were compared and contrasted using spatial overlays.

Climate corridors

We extracted raw data values on connectivity and climate flow (The Nature Conservancy

2020) for areas that were identified as 'climate-informed' corridors by using the categorical

connectivity and climate flow dataset as a mask (The Nature Conservancy 2020). The remaining

values were rescaled to fall between 0 and 1. A second climate corridor dataset on current flow

centrality (Carroll et al. 2018) was similarly rescaled. We then combined these two datasets and

analyzed locations in the 80th percentile of the distribution of combined values.

Combined Climate Corridors

$$= Z_{Climate\ Informed\ *\ Connectivity\ and\ Flow} + Z_{Current\ Flow\ Centrality}$$

Analyses

We used spatial overlay analysis to describe the extent to which the current protected

areas network covers identified climate refugia (based on national- and ecoregion-scales) and

corridors in CONUS. We quantified the extent to which identified refugia would be protected by

the 30x30 framework if it were to solely focus on current areas of high imperiled species

biodiversity and ecosystem carbon. Data on protected areas are from the PADUS 2.1 database

(USGS 2020). We use U.S. Geological Survey's Gap Analysis Program (GAP) codes, which are

specific to the management intent to conserve biodiversity. GAP 1 and 2 areas are managed in

ways typically consistent with conservation. Areas assigned a GAP 3 code are governed under

multiple-use mandates that may include biodiversity priorities but may also include incompatible

activities such as forestry and mining, and GAP 4 areas lack any conservation mandates or such

information is unknown as of 2020. Imperiled species richness was assessed from publicly

available range data (USGS GAP, International Union of Conservation of Nature - IUCN, and

U.S. Fish and Wildlife Service) for species defined as 'imperiled' (1,923 species). These include

species that are listed or under consideration for listing under the ESA, have a NatureServe G1-3

status and/or are in critically endangered, endangered or vulnerable IUCN categories. Modeled

current total ecosystem carbon is based on a high-resolution map of global above- and below-

ground carbon stored in biomass and soil (Soto-Navarro et al. 2020). We define hotspots for

biodiversity and carbon as the top 20% of values in the raw distribution of each dataset. We

selected the top 20% as a threshold to balance selectivity and breadth of coverage, but recognize

that other thresholds could be chosen. We used ArcPro v2.5 (ESRI, USA) to produce maps and

run analyses, with maps using the Albers Equal Area Conic projection.

RESULTS

Identifying refugia and corridors

Climate refugia datasets generally correlated well with others of similar methodology or

concept; three resulting classes generally represent topodiversity, climatic stability, and tree

macrorefugia (Tables 1 & S1). The main exception was for climate-based datasets with species

information, where bird macrorefugia correlated with datasets based on topodiversity, but tree

macrorefugia was the sole dataset in its class (Table S2). The three refugia classes exhibited very

little overlap with one another at the national scale: while 52% of CONUS falls into at least one

of the refugia classes, 7.5% falls into refugia identified by 2 or more classes (approx. 568,000

km^2, Fig. S1). Locations in the combined refugia layer that were within the top 20% of the

distribution of values represent these overlaps and are used for reporting the remainder of

statistics here.

34% of CONUS is identified as a climate refugia or corridor under one or more datasets

(approx. 2,652,000 km^2, Fig. 1). Climate refugia generally follow the Appalachian, Rocky, and

Cascade Mountain Ranges with additional refugia in the Ozarks and parts of California. Climate

corridors are somewhat complementary to national-scale refugia, with 28.9% of their area

(444,501 km^2) overlapping identified refugia locations. Overlaps occur in the central

Appalachians, Pacific Northwest, and portions of the Rockies, Sierra Nevadas and Ozarks.

Corridors that do not overlap with refugia are key in connecting parts of the Great Plains and

Mexico borderlands to refugia and in connecting refugia to northern locales.

Refugia identified in a stratified ecoregion approach were highly coincident with the

national scale analysis, with 63% of all national refugia overlapping with ecoregion refugia (Fig.

2). Overlaps between the two cover 12% of CONUS total land area (approx. 949,000 km^2). All

refugia combined (both from national and ecoregion-specific analyses) equal 26% of the total

CONUS land area (approx. 2.1 million km^2). Locations that were emphasized in the ecoregion-
specific approach include temperate and semi-arid prairies and places along the eastern coast.

Comparison to 30x30 objectives: biodiversity and carbon

Refugia and corridors are generally complementary on the landscape to areas of current
high biodiversity and carbon storage values (Fig. 3a&b). There is some overlap between current
biodiversity hotspots (i.e., top quartile of imperiled species richness values) and identified
national-scale refugia (36.8%) and corridors (9.3%; Table 2). Overlaps are generally
concentrated in western California and Appalachia/Ozarks regions. Overlap between carbon-rich
areas is greater in extent overall (refugia overlap = 32.5% and corridor overlap = 27.2%) and
similar in spatial pattern with greater overlap in northern areas: northern Appalachians, Crown of
the Continent and Pacific Northwest. When combining the two objectives (biodiversity and/or
carbon), 45.0% (approx. 1,000,000 km^2) of the land area representing at least one of these
objectives is also identified as part of a climate refuge or corridor.

Taking an ecoregion-specific approach to comparing refugia, corridors, biodiversity, and
carbon results in less coincidence: 22.0% and 21.7% of stratified refugia overlap with ecoregion-
specific biodiversity hotspots and carbon-rich areas, and 17.5% and 26.1% of corridors overlap
with ecoregion-specific biodiversity hotspots and carbon-rich areas, respectively (Fig. 3c&d;
Table 3).

Current protections for refugia and corridors

Overall, 12.5% of the combined network of refugia and corridors is managed consistently
with biodiversity conservation (i.e., GAP 1 or 2; 4.2% of CONUS or approx. 325,000 km^2; Fig.
4). The rest of this network falls on GAP 3 (26.5%) or GAP 4 (69.3%) lands, which represents
29.2% of CONUS (approx. 2,280,000 km^2). Proportions are similar when analyzing protection of
national-scale climate refugia and corridors separately (Table 2). Ecoregion-specific refugia fall
more heavily in GAP 4 categories with 12.2% of area on lands managed for biodiversity
conservation and 19.6% on those managed for multiple uses (Fig. 4, Table 3). Finally, the entire
set of CONUS lands representing either biodiversity conservation (GAP 1 or 2) or 30x30

objectives (biodiversity hotspots and/or carbon-rich areas) coincides with 44.5% of the national

climate refugia and corridor network.

DISCUSSION

Currently, the U.S. protected areas network and emerging conservation policy objectives

largely fail to represent valuable climate refugia and corridors. While there is some overlap,

solely using recent imperiled species ranges and carbon stores as conservation criteria will not

inherently protect climate-resilient lands. In the most protective situation - if all biodiversity

hotspots and carbon-rich areas were to be considered for strong conservation mandates (e.g.,

GAP 1 or 2 protections) - a majority (55.5%) of identified climate refugia or corridors would still

be left unprotected. The omission of landscapes for climate adaptation from planning initiatives

could inhibit the potential for longer-term conservation successes. While simply protecting

currently biodiverse or carbon-rich areas may not ensure the preservation of climate corridors

and refugia, conserving corridors and refugia will benefit imperiled species in biodiversity-rich

hotspots and promote carbon sequestration. This is particularly true in parts of the country (e.g.,

Appalachia and western California) where hotspots are not directly covered by climate corridors,

but adjacent to them, providing opportunities for migration to refugia or future climate analogs.

With over half of the contiguous U.S. identified as at least one type of climate refugia

(topodiversity, climatic stability, or tree macrorefugia), many opportunities exist for decision

makers interested in future-minded conservation. Like previous work, we demonstrate trade-offs

in using one refugia dataset over others: topodiversity data favor environmentally complex

regions like mountain ranges, whereas climatic exposure and tree macrorefugia highlight other

lands (Michalak et al. 2020). Through our ensemble approach to refugia identification we both

highlight the complementary information provided by these approaches (Belote et al. 2018) and

simplify varied complex datasets for greater interpretability. A weighted combination of the

datasets puts less pressure on the user to choose between mechanisms and on the decision maker

to have a deep understanding of the methodology when interpreting maps. However, clarification

of a specific refugia type may help states or local municipalities working to set priorities for

contributing to national refugia protections based on local environments and community needs.

In addition, taking a combined approach results in high overlap with an ecoregion-stratified

approach, suggesting representation of nearly all ecoregions in national efforts focused on

conserving climate refugia.

Currently unprotected climate refugia and corridors represent 29.2% of CONUS, of

which 38% is federally managed. Given the extent and distribution of land managers, protecting

valuable climate adaptation areas can help contribute to the 30% target numerically and

meaningfully. However, there will need to be a concerted effort by land managers in all

jurisdictions and leadership across jurisdictional boundaries.

Lands Administered by Government and Tribal Entities

Public lands can make significant contributions to achieving 30x30. The federal lands

estate is particularly expansive (20% of CONUS, 86% of PADUS; CRS 2020, Rosa and Malcom

2020) and federal land management agencies are required to varying degrees to prioritize

wildlife and habitat conservation. Currently, the majority (86%, representing 18.4% of CONUS)

of GAP 3 lands are managed by federal agencies, suggesting that substantial gains can be made

in focusing on existing statutory authorities to advance climate-smart conservation on these

lands. Of GAP 3 lands, over half are managed by the Bureau of Land Management (BLM) and

another third by the U.S. Forest Service (Rosa and Malcom 2020). Both agencies are guided by

multiple use management mandates that empower them to designate and manage lands to

enhance protection of areas recognized as having important conservation values (respectively,

the Federal Land Policy and Management Act of 1976, National Forest Management Act of

1976). The agencies can capitalize on existing land and water designation authorities - like

wilderness designation and BLM "areas of critical environmental concern" - to increase

protection for climate refugia and corridors.

Expansion of GAP 1 and 2 lands to cover more refugia and corridors can also ensure

greater conservation for climate adaptation. The U.S. Fish and Wildlife Service manages the

National Wildlife Refuge System (NWRS) expressly to conserve species and habitat, providing a

high level of federal land protection. Pursuing the acquisition of lands fundamental to species'

survival and sustainability, including climate refugia and climate corridors, to establish new

refuges would be consistent with the purview of NWRS. However, since federal land acquisition

and management decisions are often politically contentious, this may be a less feasible option for

conserving the additional 440 million acres of land needed to reach the 30% target.

State governments also manage significant acreage (approximately 4% of the U.S.),

including state forests, wildlife management areas, game lands, and natural area preserves. State

parks, or portions thereof, may also contribute to conservation refugia and corridors, but are

often categorized as GAP 4 (i.e., absent or unknown mandates for conservation). States can

contribute to 30x30 by upgrading GAP status and management of undeveloped state lands that

can further climate adaptation. The State Wildlife Action Planning (SWAP) process also requires

each state is to describe "locations and relative condition of key habitats and community types

essential to conservation of species" (USFWS & AFWA 2017). Results from this and other

studies can help inform this process and help states increasingly update their SWAPs to include

climate changes (NFWPCAN 2021).

Tribal nations hold over 56 million acres in trust by the Bureau of Indian Affairs and may

manage their lands in ways that afford more substantive protections for lands and species given

their lower rates of habitat modification (Lee-Ashley et al. 2019). A long history of managing

and observing their lands has provided many indigenous communities with valuable knowledge

and experience to inform land management and planning for climate adaptation and resilience

(BIA 2018). Respectful inclusion of indigenous systems of knowledges and perspectives "can

inform our understanding of how the climate is changing and strategies to adapt to climate

change impacts" (NFWPCAN 2021). As such, government-to-government relationships will be

important in addressing climate adaptation needs for species and peoples and may include cross-

landscape management, tribal involvement in federal and state planning, and more. The

Landscape Conservation Cooperative (LCC) program developed by Interior offers one such

mechanism to advance landscape-scale protections and coordinate climate-related land

conservation activities among Tribal Nations, federal agencies, state, local, and tribal

governments, and other stakeholders (NASEM 2016).

Private and Non-Governmental Organization Lands

As most land in the U.S. is privately owned, conservation efforts on private lands will be

critical to expanding protected areas. 62% of the refugia and 56% of corridors fall outside of the

protected areas network (GAP 4), but this only represents 20% of CONUS. This suggests that

well-targeted, voluntary acquisitions and easements could translate to large gains in private lands

conservation. Land trusts are uniquely positioned to scale-up conservation on private lands to

achieve the 30x30 target.

In addition to the role of land trusts, private working lands and associated conservation

programs can be important to achieving 30x30 (Garibaldi et al. 2020, American Farmland Trust

2021). For instance, Farm Bill programs administered by the U.S. Department of Agriculture

such as the Agriculture Conservation Easement Program (ACEP) could be targeted to lands

identified as climate refugia or connectivity areas and specify sensitive wetland habitats and

riparian areas as eligible lands for wetland easements, as these will be increasingly valuable for

supporting wildlife and ecosystem services as the climate changes (Theoharides 2014, Lewis et

al. 2019). Longer-term (30 year) contracts that offer a commitment to re-enrollment should be

encouraged to ensure enduring conservation measures. Additionally, Environmental Quality

Incentives Program (EQIP) and the Conservation Stewardship Program (CSP) can better reflect

climate adaptation needs by assigning higher ranking to practices designed to build resilient

landscapes (Theoharides 2014).

Limitations

To enhance species' resilience in the face of growing climate and biodiversity crises,

corridors and refugia must be preserved across both lands and waters. Due to some limitations of

data and our analyses, we recommend against siting protections based on the coincidence of

current biodiversity/carbon hotspots and climate refugia/corridors alone. For one,

complementarity of species assemblages is not accounted for and there may be biases toward

conserving certain taxa. Additionally, while we include aquatic species in our biodiversity

metric, and wetland/riparian areas in some topographic measures of refugia, we did not explicitly

include aquatic refugia. Currently, there is no complete national dataset to represent aquatic

refugia. Because cold-water aquatic organisms are among the most vulnerable taxa to climate

change, future analyses should focus on identifying freshwater refugia and corridors where

sufficient data exists. Given the international scope of 30x30 and the benefits of larger-scale

connectivity, future work on climate adaptation in 30x30 implementation should look beyond
terrestrial habitats and political boundaries to cover all ecosystems of North America.

Our analysis demonstrates the need to make climate adaptation a more explicit objective
in conservation planning for addressing the biodiversity crisis. Without direct consideration for
climate refugia and corridors, a 30x30 implementation focused on current species ranges and
carbon stocks may be ineffective for the longer-term persistence of species. The key to
operationalizing 30x30 and subsequent efforts will be growing a protected areas network that
ensures a long-term commitment to biodiversity and climate. By incorporating climate refugia
and corridors, the U.S. can work to protect places that will continue to serve wildlife and human
populations now and in the future.

ACKNOWLEDGEMENTS
We thank M. Anderson, J. Grand, J. Lawler, R. List, J. Michalak, S. Saunders, and R. Wynn-
Grant for their thoughtful review of the project and engaging discussion over key concepts.
Additional gratitude goes to those organizations that make these datasets publicly available to
enable this and other research.
Recent work in the field of conservation science and STEM at large has identified a bias in
citation practices such that papers from women and minorities are relatively under-cited (see
Rudd et al. 2021, Larivière et al. 2013, and others). While we did not proactively choose
references that reflect the diversity of the field in thought, form or contribution, gender, and other
factors for this work, we recognize the biases that may have been unintentionally introduced. We
look forward to future work that can help us to better understand how to support equitable
practices in conservation science.

DATA AVAILABILITY
The data that support the findings of this study are all publicly available. Those directly
generated by this work can be downloaded from the following OSF repository:
https://osf.io/jksyx/

LITERATURE CITED

Ackerly, D. D., Loarie, S. R., Cornwell, W. K., Weiss, S. B., Hamilton, H., Branciforte, R., &

Kraft, N. J. B. (2010). The geography of climate change: Implications for conservation

biogeography: Geography of climate change. *Diversity and Distributions*, *16*(3), 476–

487. https://doi.org/10.1111/j.1472-4642.2010.00654.x

AdaptWest Project. (2015). Gridded climatic velocity data for North America at 1km resolution.

https://adaptwest.databasin.org/pages/adaptwest-velocitywna/

American Farmland Trust. (2021). Agriculture's Role in 30x30: Partnering with Farmers and

Ranchers to Protect Land, Biodiversity, and the Climate. American Farmland Trust.

https://s30428.pcdn.co/wp-content/uploads/2021/04/AFT_-

_Agricultures_Role_in_30x30_-_4-5-2021.pdf

Ashcroft, M. B. (2010). Identifying refugia from climate change. *Journal of Biogeography 37*,

1407–1413. https://doi.org/10.1111/j.1365-2699.2010.02300.x

Baldwin, R. F., Trombulak, S. C., Leonard, P. B., Noss, R. F., Hilty, J. A., Possingham, H. P.,

Scarlett, L., & Anderson, M. G. (2018). The Future of Landscape Conservation.

BioScience, 68(2), 60–63. https://doi.org/10.1093/biosci/bix142

Berkessy, S. A., & Wintle, B. A. (2008). Using Carbon Investment to Grow the Biodiversity

Bank. *Conservation Biology*, 22(3), 510–513. https://doi.org/10.1111/j.1523-

1739.2008.00943.x

Belote, R. T., Dietz, M. S., McRae, B. H., Theobald, D. M., McClure, M. L., Irwin, G. H.,

McKinley, P. S., Gage, J. A., & Aplet, G. H. (2016). Identifying Corridors among Large

Protected Areas in the United States. *PLOS ONE, 11(4)*, e0154223.

https://doi.org/10.1371/journal.pone.0154223

Belote, R. T., Carroll, C., Martinuzzi, S., Michalak, J., Williams, J. W., Williamson, M. A., &

Aplet, G. H. (2018). Assessing agreement among alternative climate change projections

to inform conservation recommendations in the contiguous United States. *Scientific*

Reports, *8*(1), 9441. https://doi.org/10.1038/s41598-018-27721-6

Biden, J.R. (2021). Executive Order on Tackling the Climate Crisis at Home and Abroad.

https://www.whitehouse.gov/briefing-room/presidential-actions/2021/01/27/executive-

order-on-tackling-the-climate-crisis-at-home-and-abroad/

Buotte, P. C., Law, B. E., Ripple, W. J., & Berner, L. T. (2020). Carbon sequestration and

biodiversity co-benefits of preserving forests in the western United States. *Ecological*

Applications, 30(2), https://doi.org/10.1002/eap.2039

Bureau of Indian Affairs (BIA). (2018). National Climate Assessment: Indigenous Peoples'

Resilience Actions. https://biamaps.doi.gov/nca/

Büscher, B., Fletcher, R., Brockington, D., Sandbrook, C., Adams, W. M., Campbell, L., Corson,

C., Dressler, W., Duffy, R., Gray, N., Holmes, G., Kelly, A., Lunstrum, E., Ramutsindela,

M., & Shanker, K. (2017). Half-Earth or Whole Earth? Radical ideas for conservation,

and their implications. *Oryx, 51(3),* 407-410.https://doi.org/10.1017/S0030605316001228

Carroll, C., Roberts, D.R., Michalak, J.L. et al. (2017). Scale-dependent complementarity of

climatic velocity and environmental diversity for identifying priority areas for

conservation under climate change. Global Change Biology Early View.

https://doi.org/10.1111/gcb.13679

Carroll, C., Parks, S. A., Dobrowski, S. Z., & Roberts, D. R. (2018). Climatic, topographic, and

anthropogenic factors determine connectivity between current and future climate analogs

in North America. *Global Change Biology,* gcb.14373. https://doi.org/10.1111/gcb.14373

Carroll, C., & Ray, J. C. (2021). Maximizing the effectiveness of national commitments to
protected area expansion for conserving biodiversity and ecosystem carbon under climate
change. *Global Change Biology,* gcb.15645. https://doi.org/10.1111/gcb.15645
504

505 Cushman, S. A., McKelvey, K. S., & Schwartz, M. K. (2009). Use of Empirically Derived
506   Source-Destination Models to Map Regional Conservation Corridors. *Conservation*
507   *Biology, 23(2),* 368–376. https://doi.org/10.1111/j.1523-1739.2008.01111.x
508

509 Dickson, B. G., Roemer, G. W., McRae, B. H., & Rundall, J. M. (2013). Models of Regional
510   Habitat Quality and Connectivity for Pumas (Puma concolor) in the Southwestern United
511   States. *PLoS ONE, 8(12),* e81898. https://doi.org/10.1371/journal.pone.0081898
512

513 Dinerstein, E., Vynne, C., Sala, E., Joshi, A. R., Fernando, S., Lovejoy, T. E., Mayorga, J.,
514   Olson, D., Asner, G. P., Baillie, J. E. M., Burgess, N. D., Burkart, K., Noss, R. F., Zhang,
515   Y. P., Baccini, A., Birch, T., Hahn, N., Joppa, L. N., & Wikramanayake, E. (2019). A
516   Global Deal For Nature: Guiding principles, milestones, and targets. *Science Advances,*
517   *5(4),* eaaw2869. https://doi.org/10.1126/sciadv.aaw2869
518

Dinerstein, Eric, Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., Hahn, N.,
Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E. C., Jones, B., Barber,
C. V., Hayes, R., Kormos, C., Martin, V., Crist, E., … Saleem, M. (2017). An Ecoregion-
Based Approach to Protecting Half the Terrestrial Realm. *BioScience, 67(6),* 534–545.
https://doi.org/10.1093/biosci/bix014

Doyle, Martin W. & F. Douglas Shields. (2012). Compensatory Mitigation for Streams Under the
Clean Water Act: Reassessing Science and Redirecting Policy. *Journal of the American*
Water Resources Association (JAWRA) 48(3), 494-509.

Elsen, P. R., Monahan, W. B., Dougherty, E. R., & Merenlender, A. M. (2020). Keeping pace
with climate change in global terrestrial protected areas. *Science Advances, 6(25),*
eaay0814. https://doi.org/10.1126/sciadv.aay0814
530

Environmental Protection Agency (EPA). (2006). Ecoregions of North America.

https://www.epa.gov/eco-research/ecoregions-north-america

Favata, C. A., A. Maia, M. Pant, V. Nepal, & R. E. Colombo. (2018). Fish assemblage change

following the structural restoration of a degraded stream. *River Research and*

Applications 34(8), 927-936. https://doi.org/10.1002/rra.3336

Fei, S., J. M. Desprez, K. M. Potter, I. Jo, J. A. Knott & C. M. Oswalt. Divergence of species

responses to climate change. (2017). *Science Advances 3(5),* e1603055.

https://doi.org/10.1126/sciadv.1603055

Franks, S. J., & Hoffmann, A. A. (2012). Genetics of Climate Change Adaptation. *Annual*

Review of Genetics, 46(1), 185–208. https://doi.org/10.1146/annurev-genet-110711-

155511

Garibaldi, L. A., Oddi, F. J., Miguez, F. E., Bartomeus, I., Orr, M. C., Jobbágy, E. G., Kremen,

C., Schulte, L. A., Hughes, A. C., Bagnato, C., Abramson, G., Bridgewater, P., Carella,

D. G., Díaz, S., Dicks, L. V., Ellis, E. C., Goldenberg, M., Huaylla, C. A., Kuperman, M.,

… Zhu, C. (2021). Working landscapes need at least 20% native habitat. Conservation

Letters, 14(2). https://doi.org/10.1111/conl.12773

Gray, C. L., Hill, S. L. L., Newbold, T., Hudson, L. N., Börger, L., Contu, S., Hoskins, A. J.,

Ferrier, S., Purvis, A., & Scharlemann, J. P. W. (2016). Local biodiversity is higher

inside than outside terrestrial protected areas worldwide. *Nature Communications, 7(1),*

12306. https://doi.org/10.1038/ncomms12306

Groves, C. R., Game, E. T., Anderson, M. G., Cross, M., Enquist, C., Ferdaña, Z., Girvetz, E.,

Gondor, A., Hall, K. R., Higgins, J., Marshall, R., Popper, K., Schill, S., & Shafer, S. L.

(2012). Incorporating climate change into systematic conservation planning. *Biodiversity*

and Conservation, 21(7), 1651–1671. https://doi.org/10.1007/s10531-012-0269-3

Hamann, A., Roberts, D. R., Barber, Q. E., Carroll, C., & Nielsen, S. E. (2015). Velocity of

climate change algorithms for guiding conservation and management. *Global Change*

Biology, *21*(2), 997–1004. https://doi.org/10.1111/gcb.12736

Hausfather, Z., & Peters, G. P. (2020). RCP8.5 is a problematic scenario for near-term

emissions. *Proceedings of the National Academy of Sciences, 117(45),* 27791–27792.

https://doi.org/10.1073/pnas.2017124117

International Energy Agency (IEA). (2019). World Energy Outlook 2019.

https://www.iea.org/reports/world-energy-outlook-2019

Intergovernmental Pannel of Climate Change (IPCC). (2014). Climate Change 2014: Synthesis

Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of

the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and

L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland. https://www.ipcc.ch/report/ar5/syr/

Isaak, D.J.; Wenger, S.J.; Peterson, E.E.; Ver Hoef, J.M.; Hostetler, S.W.; Luce, C.H.; Dunham,

J.B.; Kershner, J.L.; Roper, B.B.; Nagel, D.E.; Chandler, G.L.; Wollrab, S.P.; Parkes,

S.L.; Horan, D.L. (2016). NorWeST modeled summer stream temperature scenarios for

the western U.S. Fort Collins, CO: Forest Service Research Data Archive.

https://doi.org/10.2737/RDS-2016-0033.

Jantarasami, L.C., Novak, R., Delgado, R., Marino, E., McNeeley, S., Narducci, C., Raymond-

Yakoubian, J., Singletary, L., and K. Powys Whyte. (2018). Tribes and Indigenous

Peoples. In Impacts, Risks, and Adaptation in the United States: Fourth National Climate

Assessment, Volume II [Reidmiller, D.R., Avery, C.W., Easterling, D.R., Kunkel, K.E.,

Lewis, K.L.M., Maycock, T.K., & Stewart, B.C. (eds.)]. U.S. Global Change Research

Program, Washington, DC, USA, pp. 572–603.

https://doi.org/10.7930/NCA4.2018.CH15

Jayakarana, A.D., Z.T. Smoot, D.M. Park, and D.R. Hitchcock. (2016). Relating stream function

593   and land cover in the Middle Pee Dee River Basin, SC. *Journal of Hydrology Regional*

594   *Studies, 5,* 261–275.

596   Jenkins, C. N., Van Houtan, K. S., Pimm, S. L., & Sexton, J. O. (2015). US protected lands

597   mismatch biodiversity priorities. *Proceedings of the National Academy of Sciences,*

598   *112(16),* 5081–5086. https://doi.org/10.1073/pnas.1418034112

600   Keppel, G., & Wardell-Johnson, G. W. (2012). Refugia: Keys to climate change management.

601   *Global Change Biology, 18(8),* 2389–2391. https://doi.org/10.1111/j.1365-

602   2486.2012.02729.x

604   Kraus, D., & Hebb, A. (2020). Southern Canada's crisis ecoregions: Identifying the most

605   significant and threatened places for biodiversity conservation. *Biodiversity and*

606   *Conservation, 29(13),* 3573–3590. https://doi.org/10.1007/s10531-020-02038-x

608   Krawchuk, M. A., G. W. Meigs, J. M. Cartwright, J. D. Coop, R. Davis, A. Holz, C. Kolden,

609   and AJH Meddens. (2020). Disturbance refugia within mosaics of forest fire, drought,

610   and insect outbreaks. *Frontiers in Ecology and the Environment, 18(5)*, 235–244.

611   https://doi.org/10.1002/fee.2190

613   Larivière, V., Ni, C., Gingras, Y., Cronin, B., & Sugimoto, C. R. (2013). Bibliometrics: Global

614   gender disparities in science. *Nature*, 504(7479), 211–213.

615   https://doi.org/10.1038/504211a

617   Lee-Ashley, M., Rowland-Shea, J. & Richards, R. (2019). The Green Squeeze. Washington,

618   D.C., Center for American Progress.

619   https://www.americanprogress.org/issues/green/reports/2019/10/22/476220/the-green-

620   squeeze/

Letcher, B., et al. (2017). Brook Trout Probability of Occurrence. Northeast U.S. North Atlantic

Landscape Conservation Cooperative.

Lewis, K. E., Rota, C. T., Lituma, C. M., & Anderson, J. T. (2019). Influence of the Agricultural

Conservation Easement Program wetland practices on winter occupancy of Passerellidae

sparrows and avian species richness. PLOS ONE, 14(1), e0210878.

https://doi.org/10.1371/journal.pone.0210878

Littlefield, C. E., McRae, B. H., Michalak, J. L., Lawler, J. J., & Carroll, C. (2017). Connecting

today's climates to future climate analogs to facilitate movement of species under climate

change: Climate Change and Species' Movement. *Conservation Biology*, *31*(6), 1397–

1408. https://doi.org/10.1111/cobi.12938

Locke, H. (2013). Nature needs half: A necessary and hopeful new agenda for protected areas.

PARKS, 19(2), 13–22. https://doi.org/10.2305/IUCN.CH.2013.PARKS-19-2.HL.en

Lookingbill, T. R., Gardner, R. H., Ferrari, J. R., & Keller, C. E. (2010). Combining a dispersal

model with network theory to assess habitat connectivity. *Ecological Applications, 20(2),*

427–441. https://doi.org/10.1890/09-0073.1

Maxwell, S. L., Cazalis, V., Dudley, N., Hoffmann, M., Rodrigues, A. S. L., Stolton, S.,

Visconti, P., Woodley, S., Kingston, N., Lewis, E., Maron, M., Strassburg, B. B. N.,

Wenger, A., Jonas, H. D., Venter, O., & Watson, J. E. M. (2020). Area-based

conservation in the twenty-first century. *Nature*, *586*(7828), 217–227.

https://doi.org/10.1038/s41586-020-2773-z

McClure, M. L., Hansen, A. J., & Inman, R. M. (2016). Connecting models to movements:

Testing connectivity model predictions against empirical migration and dispersal data.

Landscape Ecology, 31(7), 1419–1432. https://doi.org/10.1007/s10980-016-0347-0

McGuire, J. L., Lawler, J. J., McRae, B. H., Nuñez, T. A., & Theobald, D. M. (2016). Achieving

climate connectivity in a fragmented landscape. *Proceedings of the National Academy of*

Sciences, *113*(26), 7195–7200. https://doi.org/10.1073/pnas.1602817113

Melillo, J. M., Lu, X., Kicklighter, D. W., Reilly, J. M., Cai, Y., & Sokolov, A. P. (2016).

Protected areas' role in climate-change mitigation. *Ambio, 45(2),* 133–145.

https://doi.org/10.1007/s13280-015-0693-1

Merriam, E. R., J. T. Petty, & J. Clingerman. (2019). Conservation planning at the intersection of

landscape and climate change: brook trout in the Chesapeake Bay watershed. *Ecosphere,*

10(2), e02585. https://doi.org/10.1002/ecs2.2585

Michalak, J. L., Lawler, J. J., Roberts, D. R., & Carroll, C. (2018). Distribution and protection of

climatic refugia in North America: Climatic Refugia. *Conservation Biology*, *32*(6), 1414–

1425. https://doi.org/10.1111/cobi.13130

Michalak, J. L., D. Stralberg, J. M. Cartwright, & J. J. Lawler. (2020). Combining physical and

species-based approaches improves refugia identification, *Front Ecol Environ 18(5),* 254–

260.

Monzón, J., Moyer-Horner, L., & Palamar, M. B. (2011). Climate Change and Species Range

Dynamics in Protected Areas. *BioScience, 61(10),* 752–761.

https://doi.org/10.1525/bio.2011.61.10.5

Morelli, T. L., C. Daly, S. Z. Dobrowski, D.M. Dulen, J. L. Ebersole, & S. T. Jackson, et al. (2016).

Managing climate change refugia for climate adaptation. *PLoS ONE, 11(8),* e0159909.

https://doi.org/10.1371/journal.pone.0159909

Morelli, T. L., C. W Barrows, A. R. Ramirez, J. M. Cartwright, D. D. Ackerly, T. D. Eaves, J. L.

Ebersole, et al. (2020). Climate-change refugia: biodiversity in the slow lane. *Front Ecol*

Environ, 18(5), 228–234. https://doi.org/10.1002/fee.2189

Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J. (2000).

Biodiversity hotspots for conservation priorities. *Nature, 403(6772),* 853–858.

https://doi.org/10.1038/35002501

National Fish, Wildlife, and Plants Climate Adaptation Network (NFWPCAN). (2021).

Advancing the national fish, wildlife, and plants climate adaptation strategy into a new

decade. Association of Fish and Wildlife Agencies, Washington, DC.

https://www.fishwildlife.org/application/files/4216/1161/3356/Advancing_Strategy_Rep

ort_FINAL.pdf

National Academy of Sciences, Engineering, and Medicine (NASEM). (2016). A review of the

Landscape Conservation Cooperative. Washington, DC: The National Academies Press.

https://doi.org/10.17226/21829

Nature Conservancy, The. (2020). Resilient Land Mapping tool.

https://maps.tnc.org/resilientland/

Neilson, R. P., Pitelka, L. F., Solomon, A. M., Nathan, R., Midgley, G. F., Fragoso, J. M. V.,

Lischke, H., & Thompson, K. (2005). Forecasting Regional to Global Plant Migration in

Response to Climate Change. *BioScience, 55(9),* 749. https://doi.org/10.1641/0006-

3568(2005)055[0749:FRTGPM]2.0.CO;2

Newsom, G. (2020). Executive Order N-82-20. https://www.gov.ca.gov/wp-

content/uploads/2020/10/10.07.2020-EO-N-82-20-signed.pdf

Parks, S. A., Carroll, C., Dobrowski, S. Z., Allred, B. W. (2020). Human land uses reduce

climate connectivity across North America. *Global Change Biology 26(5)*, 2944-2955.

https://doi.org/10.1111/gcb.15009

Pörtner, H.O., Scholes, R.J., Agard, J., Archer, E., Arneth, A., Bai, X., Barnes, D., Burrows, M.,

Chan, L., Cheung, W.L., Diamond, S., Donatti, C., Duarte, C., Eisenhauer, N., Foden,

W., Gasalla, M. A., Handa, C., Hickler, T., Hoegh-Guldberg, O., Ichii, K., Jacob, U.,

Insarov, G., Kiessling, W., Leadley, P., Leemans, R., Levin, L., Lim, M., Maharaj, S.,

Managi, S., Marquet, P. A., McElwee, P., Midgley, G., Oberdorff, T., Obura, D.,

Osman, E., Pandit, R., Pascual, U., Pires, A. P. F., Popp, A., ReyesGarcía, V.,

Sankaran, M., Settele, J., Shin, Y. J., Sintayehu, D. W., Smith, P., Steiner, N.,

Strassburg, B., Sukumar, R., Trisos, C., Val, A.L., Wu, J., Aldrian, E., Parmesan, C.,

Pichs-Madruga, R., Roberts, D.C., Rogers, A.D., Díaz, S., Fischer, M., Hashimoto, S.,

Lavorel, S., Wu, N., Ngo, H.T. (2021). IPBES-IPCC co-sponsored workshop report on

biodiversity and climate change; IPBES and IPCC.

https://doi.org/10.5281/zenodo.4782538

Rights and Resources Initiative. (2020). Rights-Based Conservation: The path to preserving

Earth's biological and cultural diversity? Washington, DC: Rights and Resources

Initiative. https://rightsandresources.org/publication/rights-based-conservation/

Román-Palacios, C., & Wiens, J. J. (2020). Recent responses to climate change reveal the drivers

of species extinction and survival. *Proceedings of the National Academy of Sciences,*

117(8), 4211–4217. https://doi.org/10.1073/pnas.1913007117

Rosa, L. & J. Malcom. (2020). Getting to 30x30: Guidelines for decision makers. Defenders of

Wildlife. https://defenders.org/sites/default/files/2020-07/getting-to-30x30-guidelines-

for-decision-makers.pdf

Rudd, L. F., Allred, S., Bright Ross, J. G., Hare, D., Nkomo, M. N., Shanker, K., Allen, T.,

Biggs, D., Dickman, A., Dunaway, M., Ghosh, R., González, N. T., Kepe, T., Mbizah, M.

M., Middleton, S. L., Oommen, M. A., Paudel, K., Sillero-Zubiri, C., & Dávalos, A.

(2021). Overcoming racism in the twin spheres of conservation science and practice.

Proceedings of the Royal Society B: *Biological Sciences*, 288(1962), 20211871.

https://doi.org/10.1098/rspb.2021.1871

Saxon, E. (2008). Noah's Parks: A partial antidote to the Anthropocene extinction event.

Biodiversity, *9*(3–4), 5–10. https://doi.org/10.1080/14888386.2008.9712901

Scott, J. M., Davis, F. W., McGhie, R. G., Wright, R. G., Groves, C., & Estes, J. (2001). Nature

reserves: Do they capture the full range of America's biological diversity? *Ecological*

Applications, 11(4), 999–1007. https://doi.org/10.1890/1051-

0761(2001)011[0999:NRDTCT]2.0.CO;2

Soto-Navarro, C., Ravilious, C., Arnell, A., de Lamo, X., Harfoot, M., Hill, S. L. L., Wearn, O.

R., Santoro, M., Bouvet, A., Mermoz, S., Le Toan, T., Xia, J., Liu, S., Yuan, W., Spawn,

S. A., Gibbs, H. K., Ferrier, S., Harwood, T., Alkemade, R., … Kapos, V. (2020).

Mapping co-benefits for carbon storage and biodiversity to inform conservation policy

and action. Philosophical Transactions of the Royal Society B: Biological Sciences,

375(1794), 20190128. https://doi.org/10.1098/rstb.2019.0128

Simmons, B. A., Nolte, C., & McGowan, J. (2021). Delivering on Biden's 2030 conservation

commitment [Preprint]. *Ecology*. https://doi.org/10.1101/2021.02.28.433244

Stolton, S., Dudley, N., Çokçalışkan, B. A., Hunter, D., Ivanić, N., Kanga, E., Kettunen, M.,

Kumagai, Y., Maxted, N., Senior, J., Wong, M., Keenleyside, K., Mulrooney, D., &

Withaka, J. (2015). Values and benefits of protected areas. In *Protected Area Governance*

and Management. ANU Press.

Stralberg, D., Carroll, C., Pedlar, J. H., Wilsey, C. B., McKenney, D. W., & Nielsen, S. E.

(2018). Macrorefugia for North American trees and songbirds: Climatic limiting factors

and multi-scale topographic influences. *Global Ecology and Biogeography*, *27*(6), 690–

703. https://doi.org/10.1111/geb.12731

Stralberg, D., D. Arseneault, J. L. Baltzer, Q. E. Barber, E. M. Bayne, Y. Boulanger, et al.

(2020a). Climate-change refugia in boreal North America: what, where, and for how

long? *Front Ecol Environ 18(5),* 261–270. https://doi.org/10.1002/fee.2188

Stralberg, D., C. Carroll, & S. E. Nielsen. (2020b). Toward a climate-informed North American

protected areas network: Incorporating climate-change refugia and corridors in

conservation planning. *Conservation Letters,* e12712. https://doi.org/10.1111/conl.12712

Strassburg, B. B. N., Kelly, A., Balmford, A., Davies, R. G., Gibbs, H. K., Lovett, A., Miles, L.,

Orme, C. D. L., Price, J., Turner, R. K., & Rodrigues, A. S. L. (2010). Global congruence

of carbon storage and biodiversity in terrestrial ecosystems. *Conservation Letters*, 3(2),

98–105. https://doi.org/10.1111/j.1755-263X.2009.00092.x

Stranko, S. A., R. H. Hilderbrand, R. P. Morgan II, M. W. Staley, A. J. Becker, A. Roseberry-

Lincoln, E. S. Perry, & P. T. Jacobson. (2008). Brook Trout Declines with Land Cover

and Temperature Changes in Maryland. *North American Journal of Fisheries*

Management, 28, 1223-1232. https://doi.org/10.1577/M07-032.1

Sullivan, S. M. P. & M. C. Watzin. (2009). Stream–floodplain connectivity and fish assemblage

diversity in the Champlain Valley, Vermont. *Journal of Fish Biology, 74,* 1394–1418.

https://doi.org/10.1111/j.1095-8649.2009.02205.x

Sweeney, B. W. & J. D. Newbold. (2014). Streamside Forest Buffer Width Needed to Protect

Stream Water Quality, Habitat, and Organisms: A Literature Review. *Journal of the*

American Water Resources Association (JAWRA), 50(3), 560-584.

https://doi.org/10.1111/jawr.12203

Theoharides, K. (2014). Seeds of Resilience: Safeguarding Wildlife and Habitat from Climate

Change through the Farm Bill Conservation Programs. Defenders of Wildlife.

https://defenders.org/sites/default/files/publications/seeds-of-resilience-safeguarding-

wildlife-and-habitat-from-climate-change-through-the-farm-bill-conservation-

programs.pdf

United Nations Environment Program Convention on Biological Diversity (2020). Zero draft of

the post-2020 global biodiversity framework.

https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-en.pdf

U.S. Fish and Wildlife Service (USFWS) & Association of Fish and Wildlife Agencies (AFWA).

(2017). Guidance for Wildlife Action Plan Review and Revision.

https://fawiki.fws.gov/download/attachments/5931146/2017%20Guidance%20on%20Wil

dlife%20Action%20Plan%20Review%20and%20Revision.pdf?version=1&modification

Date=1512664258000&api=v2

U.S. Geological Survey (USGS) Gap Analysis Project (GAP). (2020). Protected Areas Database

of the United States (PAD-US) 2.1. U.S. Geological Survey data release,

https://doi.org/10.5066/P92QM3NT.

U.S. Departments of Interior (DOI), Agriculture, Commerce, and the Council on Environmental

Quality. (2021). Conserving and Restoring American the Beautiful.

https://www.doi.gov/sites/doi.gov/files/report-conserving-and-restoring-america-the-

beautiful-2021.pdf

Venter, O., Magrach, A., Outram, N., Klein, C. J., Possingham, H. P., Di Marco, M., & Watson,

J. E. M. (2018). Bias in protected-area location and its effects on long-term aspiration of

biodiversity conventions. Conservation Biology, 32(1), 127-134.

https://doi.org/10.1111/cobi.12970

Vose, R. S., Easterling, D. R., Kunkel, K. E., LeGrande, A. N., & Wehner, M. F. (2017). Ch. 6:

Temperature Changes in the United States. Climate Science Special Report: Fourth

National Climate Assessment, Volume I. U.S. Global Change Research Program.

https://doi.org/10.7930/J0N29V45

Wiens, J. J. (2016). Climate-Related Local Extinctions Are Already Widespread among Plant

and Animal Species. *PLOS Biology, 14(12),* e2001104.

https://doi.org/10.1371/journal.pbio.2001104

Wuebbles, D. J., Fahey, D. W., Hibbard, K. A., Dokken, D. J., Stewart, B. C., & Maycock, T. K.

(2017). Climate Science Special Report: Fourth National Climate Assessment, Volume I.

U.S. Global Change Research Program. https://doi.org/10.7930/J0J964J6

 TABLES AND FIGURES

863 **Table 1.** Description of refugia datasets. All datasets are currently based on the Coupled Model

864 Intercomparison Project phase 5 (CMIP5) models. Classes are based on results from a principal

865 components analysis where component 1 (topodiversity) explained 33.8%, component 2 (climate

866 stability) explained 15.9% and component 3 (tree macrorefugia) explained near 13.8% of

867 variation. See SI for additional details.

Refugia Class	Dataset	Description	Source
Topodiversity	Current Climate Diversity	*Climate-Based.* Based on 11 bioclimatic variables using climate data for a 30-year climate normal period (1981-2010).	Carroll et al., 2017
	Ecotypic Diversity	*Landscape-based and climate-based.* Derived from edaphic, climatic, and landcover data.	Carroll et al., 2017
	Land Facet Diversity	*Landscape-based.* Incorporated elevation, latitude-adjusted elevation, topographic position index, slope, modified heat load index, and soil.	Carroll et al., 2017
	Landscape Diversity	*Landscape-based.* Described the diversity of microhabitats and climatic gradients. Microclimates were measured by quantifying elevation range, the variety of small-scale landforms, and the density and configuration of wetlands in a 100-acre neighborhood.	The Nature Conservancy, 2020
	Bird Macrorefugia	*Climate-based and species-based.* Focused on regions where the current and projected future species ranges overlap. Input based on current species niches for 268 songbird species; climate velocity based on 4 representative GCMs, RCP 4.5, 2080s.	Stralberg et al., 2018
Climatic Stability	Climatic Dissimilarity	*Climate-based.* Described how different the future climate at a location will be from its current climate. Measured in terms of multivariate climate characteristics, via a principal components analysis (PCA) of 11 biologically-relevant temperature and precipitation variables, RCP 4.5, 2080s.	Belote et al., 2018

		Climate-based. Velocity was calculated by dividing the rate of climate change by the rate of spatial climate variability to focus on regions where climatic conditions move more slowly across the landscape. Input based on A2 emissions scenarios implemented by seven GCMs of the CMIP3 multimodel dataset, RCP 4.5, 2080s.	AdaptWest Project, 2015
	Climate Velocity		
Tree Macrorefugia	Tree Macrorefugia	Climate-based and species-based. Focused on regions where the current and projected future species ranges overlapped. Input based on current species niches for 324 tree species; climate velocity based on 4 representative GCMs, RCP 4.5, 2080s.	Stralberg et al., 2018

889 **Table 2.** Overlays of national-level datasets representing protected areas, carbon stores,

890 biodiversity, climate refugia, and climate corridors. Values represent the percent of each top line

891 item (column) that falls within each row. Values in parentheses are the percent of total CONUS

892 area represented by the overlay.

% of top line items that fall into each of the following:	GAP 1 & 2	GAP 3	Top 20% Carbon	Top 20% Biodiversity	Top 20% Refugia	Top 20% Climate-Informed Corridors
GAP 1 & 2	100 (7.5)	0.0 (0.0)	12.7 (2.4)	3.7 (0.7)	13.3 (2.6)	13.8 (2.7)
GAP 3	0.0 (0.0)	100 (16.6)	20.2 (3.9)	5.1 (1.0)	25.0 (4.8)	30.4 (6.0)
Top 20% Carbon	32.8 (2.4)	23.3 (3.9)	100 (20.0)	28.8 (5.6)	32.5 (6.2)	27.2 (5.4)
Top 20% Biodiversity	12.3 (0.7)	8.7 (1.0)	32.0 (5.6)	100 (20.0)	30.8 (5.9)	11.2 (1.8)
Top 20% Refugia	34.2 (2.6)	29.0 (4.8)	32.8 (6.2)	36.8 (5.9)	100 (20.0)	28.7 (5.7)
Top 20% Climate-Informed Corridors	36.5 (2.7)	36.3 (6.0)	25.8 (5.4)	9.3 (1.8)	29.6 (5.7)	100 (20.0)

Table 3. Overlays of ecoregion-specific datasets representing protected areas, carbon stores, biodiversity, climate refugia, and climate corridors. Values represent the percent of each top line item (column) that falls within each row. Values in parentheses are the percent of total CONUS area represented by the overlay.

% of top line items that fall into each of the following:	GAP 1 & 2	GAP 3	Top 20% Carbon	Top 20% Biodiversity	Top 20% Refugia	Top 20% Climate-Informed Corridors
GAP 1 & 2	100 (7.5)	0.0 (0.0)	1.1 (2.2)	8.6 (1.5)	12.2 (2.4)	13.8 (2.7)
GAP 3	0.0 (0.0)	100 (16.6)	17.9 (3.5)	17.7 (3.1)	19.6 (3.8)	30.4 (6.0)
Top 20% Carbon	29.8 (2.2)	21.2 (3.5)	100 (20.0)	28.0 (4.9)	21.7 (4.2)	26.1 (5.2)
Top 20% Biodiversity	18.4 (1.5)	17.5 (3.1)	26.4 (4.9)	100 (20.0)	22.0 (4.2)	17.5 (3.3)
Top 20% Refugia	31.4 (2.4)	22.7 (3.8)	21.3 (4.2)	24.2 (4.2)	100 (20.0)	25.9 (5.1)
Top 20% Climate-Informed Corridors	36.5 (2.7)	36.3 (6.0)	26.3 (5.2)	18.8 (3.3)	26.7 (5.1)	100 (20.0)

Figure 1. A) National-scale and B) ecoregion-specific refugia (top 20% of all three refugia classes combined) with climate-informed corridors (ecoregions are outlined in black). The full raster datasets were used to identify refugia in national analyses. Ecoregion-specific analyses employ a stratified approach, where refugia are identified for each ecoregion separately before combining them together. Ecoregions are outlined in black in map B.

934 **Figure 2.** Coincidence between national-scale and ecoregion-specific refugia. The full raster

935 datasets were used to identify refugia in national analyses. Ecoregion-specific analyses employ a

936 stratified approach, where refugia are identified for each ecoregion separately before combining

937 them together. Ecoregions are outlined in black.

Figure 3. Overlap between national-scale (A,B) and ecoregion-scale (C,D) refugia and corridors with carbon stocks (B,D) and biodiversity hotspots (A,C). The full raster datasets were used to identify refugia in national analyses. Ecoregion-specific analyses employ a stratified approach, where refugia are identified for each ecoregion separately before combining them together. Ecoregions outlined in black in maps C and D.

Figure 4. Overlap between national-scale refugia (A), climate corridors (B), and either refugia or corridors (C) with the protected areas database of the US (PADUS). GAP codes are specific to the management intent to conserve biodiversity; GAP 1 and 2 areas are managed in ways typically consistent with conservation and GAP 3 areas are governed under multiple-use mandates that may include biodiversity priorities but may also include incompatible activities.