**The unfolded protein response triggers the ancestral immune deficiency pathway**

Lindsay C. Sidak-Loftis[1], Kristin L. Rosche[1], Natasha Pence[2#], Jessica K. Ujczo[3], Joanna

Hurtado[1,4], Elizabeth A. Fisk[1], Susan M. Noh[1,3], John W. Peters[2], Dana K. Shaw[1,4]*

[1]Program in Vector-borne Disease, Department of Veterinary Microbiology and Pathology,

Washington State University, Pullman, WA, USA.

[2]Institute of Biological Chemistry, Washington State University, Pullman, WA USA.

[3]United States, Department of Agriculture, Agricultural Research Service, Animal Disease

Research Unit, Pullman, WA, USA.

[4]School of Molecular Biosciences, Washington State University, Pullman, Washington, USA.

[#]Present address: Department of Chemistry, Dartmouth College, Hanover, NH, USA.

**Corresponding Author**: *Dana K. Shaw; Dana.Shaw@wsu.edu

**ABSTRACT**

The insect immune deficiency (IMD) pathway is a defense mechanism that senses and

responds to Gram negative bacteria. Ticks lack genes encoding upstream components that

initiate the IMD pathway. Despite this deficiency, core signaling molecules are present and

functionally restrict tick-borne pathogens. The molecular events preceding activation are

unclear. Here, we show that the Unfolded Protein Response (UPR) initiates the IMD network in

*Ixodes scapularis* ticks. The endoplasmic reticulum (ER) stress receptor, IRE1α, is

phosphorylated in response to tick-borne bacteria, but does not splice the mRNA encoding

XBP1. Instead, through protein modeling and reciprocal pulldowns, we show that *Ixodes* IRE1α

complexes with TRAF2. Disrupting IRE1α-TRAF2 signaling blocks IMD pathway activation and

diminishes the production of antimicrobial effectors. Through *in vitro*, *in vivo* and *ex vivo*

techniques we demonstrate that the UPR-IMD pathway circuitry limits the Lyme disease-

causing spirochete *Borrelia burgdorferi* and the rickettsial agents *Anaplasma phagocytophilum*

and *A. marginale* (anaplasmosis). Altogether, our study uncovers the upstream signaling

requirements of the IMD pathway in ticks. We propose that this mode of IMD network activation

is evolutionarily ancient, predating the classically defined pathway in insects.

## INTRODUCTION

Arthropod-borne diseases continue to be a substantial source of morbidity and mortality worldwide[1]. Factors influencing the ability of arthropods to harbor and transmit pathogens are incompletely understood, although progress on this front has been made in recent years. It is now acknowledged that arthropod immunity is an important force in shaping vector competency[2–8]. For example, humoral defense networks such as the Immune Deficiency (IMD) pathway recognize and restrict invading microbes. As classically defined in *Drosophila melanogaster*, IMD pathway signaling events are similar to the tumor necrosis factor receptor (TNFR) pathway in mammals, but instead respond to Gram negative bacteria by sensing DAP (diaminopimelic acid)-type peptidoglycan[9,10]. Pathway initiating receptors PGRP-LC and PGRP-LE (peptidoglycan recognition proteins LC and LE) recruit adapter molecules IMD and FADD (fas-associated protein with death domain)[11,12], the latter pairing with DREDD (death-related ced-3/Nedd2-like protein)[13] which cleaves IMD. The E3 ubiquitin ligase IAP2 (inhibitor of apoptosis 2) and E2 conjugating enzymes Bendless, Uev1a and Effette then promote (K)63 polyubiquitylation of IMD[9,10,14]. The resulting signaling scaffold leads to cleavage of the NF-κB signaling molecule Relish, which translocates to the nucleus and promotes antimicrobial peptide (AMP) expression[10,14].

Significant advances in elucidating arthropod immunity have been possible owing the insect model organism, *Drosophila*. Less is known about immunity in more ancient arthropods, although deviations from classically defined fly immunity have been reported. For example, arachnids, including mites and spiders, and several hemimetabolous insects (those with incomplete metamorphosis) such as lice, bed bugs, psyllids, squash bugs, and whiteflies do not have upstream components of the IMD pathway[15–27]. Triatomine bugs recently had many IMD pathway components identified, but are missing the gene encoding IMD itself[32–34]. The *Ixodes scapularis* genome lacks genes encoding upstream regulators of the IMD pathway including

transmembrane *PGRPs*, *imd* and *fadd*[15,27,28,30]. Despite the absence of upstream regulators,

core IMD signaling molecules are active against infection[28,30,33–35]. For example, activity of the

*Ixodes* IMD pathway hinges on Bendless, Uev1a, XIAP (X-linked inhibitor of apoptosis), p47,

Relish, and the negative regulator Caspar, which functionally restricts tick-borne pathogens

*Borrelia burgdorferi* (Lyme disease) and *Anaplasma phagocytophilum* (granulocytic

anaplasmosis)[5,28,30,31]. In the absence of classically defined pathway initiators, functionality of

the core IMD cascade suggests that an alternative mode of activation exists.

66            Cellular stress responses are well-conserved across eukaryotes and respond to adverse

environmental conditions, such as infection[36–45]. Herein, we demonstrate that a stress-response

network, the Unfolded Protein Response (UPR), initiates the IMD pathway in arthropods. *B.*

*burgdorferi* and *A. phagocytophilum* activate the endoplasmic reticulum (ER) stress receptor

IRE1α (inositol-requiring enzyme 1α), which pairs with a TRAF2 (TNF receptor associated factor

2) signaling molecule. Through molecular modeling, biochemical interactions, pharmacological

manipulations and RNAi, we show that the tick IRE1α-TRAF2 axis functionally restricts *B.*

*burgdorferi* and *A. phagocytophilum* in *I. scapularis* ticks and initiates production of antimicrobial

effectors. IRE1α-TRAF2 signaling also restricts the cattle pathogen *Anaplasma marginale* in

*Dermacentor andersoni* ticks and potentiates IMD pathway-specific AMPs in *Drosophila*.

Collectively, we show a fundamentally distinct mode of IMD pathway activation that is

conserved across arthropods. We propose that the IRE1α-TRAF2 mediated form of IMD

network activation is ancestral to the classically defined pathway in *Drosophila* and functions as

an ancient form of immunity against infection.

**RESULTS**

*The Ixodes UPR responds to tick-borne pathogens and restricts bacterial colonization*

The absence of IMD pathway initiating molecules led us to hypothesize that the core

signaling components may be induced through crosstalk with other molecular circuits. The UPR

is a response network that is activated by ER stress through the transmembrane receptors

IRE1α, PERK (PKR-like ER kinase) and ATF6 (Activating transcription factor 6). Downstream

signaling cascades ensue with the goal of restoring cellular homeostasis. In a non-stressed

state, the sensor molecule BiP (binding immunoglobulin protein) keeps all receptors in an

inactive state by binding to them[36–38] (Fig 1A). We first evaluated whether tick-borne pathogens

induce the UPR in *I. scapularis* by quantifying gene expression in *A. phagocytophilum*-infected

nymphs. Relative to uninfected ticks (dotted baseline), significant increases were observed with

*BiP, IRE1α,* and *traf2,* suggesting that the tick UPR responds to infection (Fig 1B).

To determine how the UPR impacts pathogen survival in ticks, we used pharmacological

inducers or RNAi with the ISE6 *I. scapularis* cell line. Tick cells were treated with low doses of

either thapsigargin or tunicamycin to induce ER stress prior to *A. phagocytophilum* infection.

Thapsigargin inhibits the sarco/endoplasmic reticulum $Ca^{2+}$ ATPase (SERCA), which decreases

calcium levels in the ER[46]. Tunicamycin blocks N-linked glycosylation, leading to an increase of

misfolded proteins[47]. Both treatments resulted in significantly less *A. phagocytophilum* (Fig 1C-

D). We also used an RNAi-based approach to overactivate the UPR by decreasing expression

of the negative regulator BiP. In agreement with pharmacological induction, transcriptional

silencing of *BiP* caused a decrease in *A. phagocytophilum* colonization (Fig 1E). Altogether, this

demonstrates that *A. phagocytophilum* induces the UPR in ticks, which functionally restricts

bacterial colonization and survival.

*Infection induces IRE1α activation, but not XBP1*

Transcripts induced by *A. phagocytophilum* are associated with the IRE1α signaling axis

(Fig 1A-B), which is the most conserved branch of the UPR among eukaryotes[48]. When

activated, IRE1α autophosphorylates and either splices the mRNA *xbp1* (X-box binding protein

1) or signals through TRAF2[36,37,49,50] (Fig 1A). Unspliced *xbp1* mRNA (*xbp1$^U$*) is held in an

inactive state in the cytoplasm by forming a hairpin structure that inhibits translation. The RNase

domain of IRE1α splices an internal intron from *xpb1$^U$* allowing it to be translated into a protein

that functions as a transcription factor[49,51–54] (Fig 1A). Alternatively, IRE1α can recruit the

signaling molecule TRAF2 to produce proinflammatory responses through NF-κB signaling[36–

38,50]. We aligned mammalian sequences from the IRE1α pathway with tick homologs and

observed sequence similarity with BiP, IRE1α, XBP1 and TRAF2 (Supplementary Figure 1A-D).

Notably, the IRE1α kinase domain, RNase domain, and the activity-inducing phospho-serine

(Supplementary Figure 1B) were well-conserved with human sequences. Given this sequence

conservation, we used an antibody against human phosphorylated IRE1α to examine the

posttranslational activation status of IRE1α in ticks. ISE6 tick cells were either treated with

tunicamycin or infected with *A. phagocytophilum*. Increased amounts of phosphorylated IRE1α

were observed by immunoblot under both conditions indicating receptor activation (Fig 2A).

When IRE1α phosphorylation was blocked in tick cells with the pharmacological inhibitor

KIRA6[55], significant increases in *A. phagocytophilum* were observed (Fig 2B). Similarly,

reducing the expression of IRE1α through RNAi also increased *A. phagocytophilum* bacterial

burden (Fig 2C). These data show that IRE1α signaling in ticks is activated by infection and

restricts bacterial colonization *in vitro*.

To delineate the signaling events downstream from IRE1α, *xbp1$^U$* was next examined in

infected ISE6 cells. Primers flanking the *xbp1* intron (Supplementary Fig 1E) were used to

delineate spliced and unspliced transcripts by PCR. Unspliced *xbp1$^U$* migrates as a single 459

bp band. In contrast, spliced *xbp1$^s$* presents as a trimer on an agarose gel, consisting of spliced

transcripts (*xbp1$^S$*, 434 bp), unspliced transcripts (*xbp1$^U$*) and an *xbp1$^U$-xbp1$^S$* heterodimer that

is an artifact of PCR and migrates slightly higher. Spliced *xbp1$^s$* was observed in thapsigargin-

treated tick cells under all conditions. In contrast, *A. phagocytophilum* infection did not induce

*xbp1$^U$* splicing at any time points *in vitro* (Fig 2D). We next probed *in vivo* samples from replete

*I. scapularis* nymphs that were either fed on uninfected mice or those infected with *A.*

*phagocytophilum* or *B. burgdorferi*. Across all samples, *xbp1$^U$* remained unspliced (Fig 2E).

These results indicate that although tick IRE1α is activated by infection and restricts bacterial

burden, this phenotype is not carried out through XBP1 activity.

Since XBP1 is not responsive to infection, we sought to determine whether IRE1α is

signaling through the *Ixodes* TRAF2 protein. Reducing the expression of *traf2* through RNAi in

*Ixodes* ISE6 cells caused a significant increase in *A. phagocytophilum* (Fig 2F), correlating with

the phenotype observed when silencing *ire1α* transcripts (Fig 2C). These data, together with

upregulated *traf2* expression in *A. phagocytophilum*-infected *I. scapularis* nymphs (Fig 1B),

suggests that IRE1α is signaling through TRAF2 to restrict pathogen colonization.

*IRE1α interfaces with TRAF2 in I. scapularis ticks*

To determine whether *I. scapularis* IRE1α can interface with TRAF2, we performed

structural modeling and *in silico* docking analysis. BLAST was used to identify the human IRE1α

crystal structure[56] (PDB code 6URC) as a homology template for modeling the cytosolic

RNase/kinase domain of *I. scapularis* IRE1α. The structure was modeled in the active state

quaternary structure proposed to be necessary for autophosphorylation and RNase activity[57]

(Supplementary Figure 2A-B). Next, a human TRAF2 crystal structure[58] (PDB code 1CA9) was

identified by BLAST as a modeling template for *I. scapularis* TRAF2. The modeled form of *I.*

*scapularis* TRAF2 C-terminal region features part of a coiled-coil domain and the highly

conserved TRAF-C domain (Supplementary Fig 2C). In addition, the homology model is a trimer

where the coiled-coil domain is a single alpha helix and the TRAF-C domain forms an eight-

stranded antiparallel β-sandwich.

How TRAF2 interacts with the RNase/kinase domain of IRE1α remains poorly defined.

To gain structural insight into the *I. scapularis* IRE1α-TRAF2 complex*,* we used a prediction-

driven docking approach[59]. This tactic combines the utility of interface prediction with *ab initio*

docking and is a useful alternative to *ab initio* docking alone when examining protein-protein

complex formation. CPORT (Consensus Prediction of interface Residues in Transient

complexes)[59] was used to assign active and passive residues at the interface of the trimeric

TRAF-C domains and the RNase/kinase domain of IRE1α (Fig 3A). Residues were then used to

filter the docking process by HADDOCK 2.2[60], which optimizes residue conformations at the

interface before proceeding to refinement. The docking model places the trimeric TRAF2

interface at the kinase domain of IRE1α with a buried surface area of 3262.16 Å$^2$ (Fig 3B).

Importantly, trimeric TRAF2 is positioned in a manner that does not interfere with the IRE1α

dimer interface and is away from the C-terminal transmembrane domain (circled) that anchors

IRE1α to the ER (Fig 3B). Five salt bridge interactions were identified that define how the

TRAF2 trimer is positioned onto the kinase domain of IRE1α (Fig 3C). Each chain of TRAF2

participates in salt bridge interactions with the kinase domain of IRE1α. Therefore, the

oligomeric state of TRAF2 seems to play an important role in docking specificity with the

RNase/kinase domain of IRE1α. Altogether, *in silico* docking analyses with *Ixodes* IRE1α and

TRAF2 suggest that these two molecules can directly interface with one another.

To confirm that IRE1α and TRAF2 specifically interact, we used a Human Embryonic

Kidney (HEK) 293T cell transfection system with plasmids expressing *Ixodes* IRE1α and TRAF2

fused to affinity tags (Fig 3D). Recombinant protein expression was confirmed by

immunoblotting transfected cells with antibodies for FLAG and HA tags (IRE1α-FLAG and

TRAF2-HA). When IRE1α and TRAF2 are co-expressed, immunoprecipitating with antibodies

against the FLAG tag demonstrates that IRE1α specifically pulls down TRAF2 and vice versa

(Fig 3D). Altogether, these data demonstrate that *I. scapularis* IRE1α and TRAF2 directly and

specifically interact.

*IRE1α and TRAF2 restrict in vivo bacterial colonization of Ixodes ticks*

We next determined whether the pathogen-restricting activity of IRE1α and TRAF2

observed *in vitro* had similar impacts *in vivo*. To knock down gene expression, unfed *I.*

*scapularis* nymphs were microinjected with siRNA targeting *ire1α* and *traf2* or with a scrambled

control (scRNA). Nymphs were rested overnight and then fed to repletion on *A.*

*phagocytophilum*-infected mice. Gene silencing and bacterial burden were both quantified by

qRT-PCR. Similar to *in vitro* experiments, reducing the expression of *ire1α* and *traf2* lead to an

increase in *A. phagocytophilum* burdens in *I. scapularis* nymphs (Fig 4A-B).

*I. scapularis* take a blood meal once per life stage, with ticks initially becoming infected

during the larval phase[61]. Since gene expression can vary depending on arthropod life stage[62–

64], we examined the impact of IRE1α and TRAF2 on pathogen colonization in larvae. We

silenced *traf2* and *ire1α* in *I. scapularis* larvae using a modified immersion protocol where ticks

were submerged in siRNA or scrambled controls overnight[65]. Following immersion, larvae were

rested for 24 hours before feeding to repletion on *A. phagocytophilum*-infected mice. Significant

knockdown of *ire1α* and *traf2* was observed in siRNA-treated larvae with this method, which

caused an increase in *A. phagocytophilum* numbers (Fig 4C-D).

Soon after *A. phagocytophilum* is acquired, the bacteria migrate to the salivary glands

where they persist throughout the tick life cycle[61,66,67]. To understand how IRE1α influences

bacterial colonization in tick tissue subsets, we employed an *ex vivo* tick organ culture

system[68,69]. Midguts and salivary glands from adult *I. scapularis* ticks were dissected and

treated with the IRE1α inhibitor KIRA6 prior to infection with *A. phagocytophilum* (Fig 5E).

Similar to *in vitro* and *in vivo* findings, inhibiting the activity of IRE1α lead to significantly higher

*A. phagocytophilum* burden in *ex vivo* salivary gland and midgut cultures (Fig 4F-G),

demonstrating that this signaling axis functionally restricts bacterial colonization in disparate tick

tissues.

We next asked whether the activity of IRE1α-TRAF2 signaling was restrictive to different

tick-borne microbes, such as the Lyme disease-causing spirochete *B. burgdorferi*. Expression of

*ire1α* and *traf2* was knocked down through RNAi in both *I. scapularis* nymphs and larvae using

the same methods described above and ticks were fed to repletion on *B. burgdorferi*-infected

mice. In agreement with the phenotype observed with *A. phagocytophilum*, significantly higher

*B. burgdorferi* levels were observed in siRNA-treated ticks at both the nymph and larval life

stages (Fig 5A-D). These data show that IRE1α-TRAF2 signaling is broadly responsive to

multiple *I. scapularis*-transmitted pathogens and is functionally restrictive to microbial

colonization during different tick life stages.

*The IMD pathway is triggered by IRE1α in arthropods.*

TRAF2 is a component of the mammalian TNFR network, which is functionally

analogous to the arthropod IMD pathway. This parallel led us to ask whether the antimicrobial

activity of the IRE1α-TRAF2 axis operates through arthropod immunity. AMPs specific to the

IMD pathway have not yet been identified in ticks. Instead, the *Drosophila* S2* cell line can be

used as a surrogate model[28]. To examine whether ER stress induces an immune response in

the absence of microbes, we treated S2* cells with the UPR inducer thapsigargin. AMPs

corresponding to the IMD pathway (*diptericin*, *attacin A* and *cecropin A2*) were significantly

induced in a dose-dependent manner over unstimulated controls (Fig 6A). In contrast, the Toll

pathway AMP *IM1* was not significantly different, demonstrating that ER stress leads to IMD

pathway activation independent of microbial agonists.

It is known that the IMD pathway is responsive to tick-transmitted pathogens *A.*

*phagocytophilum* and *B. burgdorferi*[28,30]. Since tick-borne microbes also activate the UPR (Figs

1B and 2A) and ER stress induces the IMD network (Fig 6A), we asked whether blocking IRE1α

during infection would inhibit the IMD pathway. S2* cells that were treated with the IRE1α

inhibitor KIRA6 prior to *A. phagocytophilum* or *B. burgdorferi* infection (MOI 50) showed

significantly reduced IMD pathway AMPs (Fig 6B-C). Collectively, our results provide strong

evidence that the IRE1α-TRAF2 axis functions as an IMD pathway-activating mechanism.

*IRE1α-TRAF2 signaling potentiates antimicrobial products*

A complementary immune mechanism to the IMD pathway is the production of reactive

oxygen species (ROS), which cause bactericidal damage to nucleic acids, proteins and

membrane lipids[15,70,71]. Because *B. burgdorferi* and *A. phagocytophilum* are both sensitive to

killing by ROS[72–75] and the mammalian UPR can lead to ROS production[76,77], we investigated

whether ROS can be induced by the *Ixodes* IRE1α-TRAF2 pathway. ISE6 cells were stimulated

with either thapsigargin, tunicamycin or a vehicle control and monitored for ROS with the

fluorescent indicator *2′,7′*-dichlorofluorescin diacetate. Pharmacological inducers caused

significantly higher fluorescence, indicating that the tick UPR potentiates ROS (Fig 6D).

Infection with *A. phagocytophilum* and *B. burgdorferi* also elicited ROS production in tick cells

(Fig 6E-F). However, blocking IRE1α activity with KIRA6 either reduced or completely mitigated

ROS (Fig 6E-F), demonstrating that infection-induced ROS is controlled by IRE1α.

*IRE1α-TRAF2 signaling restricts pathogens across tick vectors*

Since the UPR is conserved across eukaryotes, we explored the possibility that the

microbe-restricting activity of IRE1α-TRAF2 signaling could functionally impact other arthropod

vectors. *D. andersoni* ticks are important disease vectors that transmit several pathogens

including the obligate intracellular rickettsia, *A. marginale*[78]. When inducing the UPR in the *D.*

*andersoni* tick cell line DAE100 with tunicamycin and thapsigargin (Fig 7A-B) or blocking IRE1α

with KIRA6 (Fig 7C), we observed significant changes in *A. marginale* invasion and replication,

comparable to what was observed with *I. scapularis* and *A. phagocytophilum* (Figs 1C-D, 2B).

Moreover, higher bacterial loads were also observed in *D. andersoni ex vivo* midgut and salivary

gland cultures when IRE1α activity was blocked with KIRA6 (Fig 7D-F). Altogether, this

demonstrates that the microbe-restricting activity of IRE1α-TRAF2 signaling is conserved across

tick species and is active against disparate pathogens, including intracellular bacteria (*A.*

*phagocytophilum* and *A. marginale*) and extracellular spirochetes (*B. burgdorferi*).

**DISCUSSION**

How arthropod immunity responds to infection is a fundamental factor influencing the

ability of vectors to harbor and transmit pathogens[2–8]. The IMD pathway is increasingly

recognized as being divergent across species, with classically defined upstream regulators

missing in many arthropod genomes[15–27,32,34,35]. This suggests that an alternative activation

mechanism exists. In this article we demonstrate that the *I. scapularis* IMD pathway is initiated

through the IRE1α-TRAF2 axis of the UPR. Colonization and replication of *A. phagocytophilum*

and *B. burgdorferi* are restricted in ticks by IRE1α and TRAF2 both *in vitro* and *in vivo*.

Moreover, we show that production of IMD pathway-specific AMPs and ROS in response to *A.*

*phagocytophilum* and *B. burgdorferi* are dependent on IRE1α activity and that this mode of

antibacterial restriction is conserved across arthropods. Collectively, our findings provide an

explanation for how the core IMD pathway is activated in the absence of canonical upstream

regulators.

To our knowledge, this is the first time that cellular stress responses have been

implicated in influencing vector competency. Why host cell stress responses are triggered by *A.*

*phagocytophilum* and *B. burgdorferi* remains unclear. Ticks do not appear to suffer pathological

consequences from the microbes they transmit. The connection between host cell stress and

immune outcomes supports a model where transmissible pathogens would benefit most by

decreasing infection-induced stress. This model is reenforced by the absence of common

inflammatory pathogen-associated molecular patterns (PAMPs) in many tick-transmitted

pathogens. For example, all *Ixodes*-transmitted bacteria lack lipopolysaccharide (LPS) and

DAP-PGN[79–82]. *B. burgdorferi* flagella are housed in the periplasm, effectively shielded from

recognition by host cells[83]. During coevolution with ticks, *Ixodes*-transmitted pathogens may

have lost inflammatory PAMPs with the benefit of reducing cellular stress and host responses,

thereby promoting persistence and transmission. Nevertheless, our data shows that *A.*

*phagocytophilum* and *B. burgdorferi* impart at least some stress on ticks. Whether this is

through resource competition, damage to host cells or secreted microbial molecules is not

presently known. Since immune responses are energetically costly to the host[84,85], we speculate

that the tick response is tuned to match the level of threat imposed by infection, ultimately

striking a balance that conserves resources and preserves tick fitness.

The absence of upstream IMD pathway regulators appears to be a shared trait among

chelicerates and hemimetabolous insects[15–24,26–30,32–34,84]. Considering this observation,

classically defined IMD pathway initiating molecules may have evolved with holometabolous

insects (those with complete metamorphosis), such as *Drosophila* and mosquitos, which

undergo dramatic tissue remodeling between life stages[86,87]. The classically described IMD

pathway is speculated to have arisen as a defense mechanism against microbiome-resident

bacteria that are liberated with midgut breakdown during metamorphosis[34]. In contrast,

arthropods with incomplete metamorphosis, such as ticks and hemimetabolous insects, undergo

some tissue remodeling during molting but immature stages generally resemble adult

morphologies[86,88,89]. Since these arthropods are phylogenetically more ancient than

holometabolous insects[86,87,90–92], we propose that UPR activation of core IMD signaling

molecules is ancestral to the classically defined pathway in *Drosophila*. Furthermore, our data

supports that this network is conserved across arthropods. Specifically, IRE1α-mediated IMD

pathway activation was observed both in *Drosophila* cells (Fig 6B-C) and in two species of ticks

(Fig 4,5 and 7). Collectively, this suggests that the UPR-IMD signaling network evolved in a

common ancestor of hexapods and chelicerates.

In summary, we have discovered a linkage between cellular stress responses and

arthropod immunity where the IRE1α-TRAF2 signaling axis initiates an ancestral IMD pathway

(Supplementary Figure 3). The previous "orphaned" status of the IMD pathway in ticks was a

perception that arose from comparative studies with the holometabolous insect, *Drosophila*.

This revelation underscores the importance of studying fundamental processes outside of model

organisms, which may be valuable for determining concepts that are basally applicable across

species. Our findings are conceptually important given that the IMD pathway widely impacts

vector competence in many arthropods. With this commonality, one can envision a scenario

where a conserved network across species may be an attractive target for future transmission

intervention strategies.

**METHODS**

*Bacteria and animal models*

*E. coli* cultures were grown in lysogeny broth (LB) supplemented with ampicillin at 100

µg µl$^{-1}$. Cultures were grown overnight at 37°C with shaking between 230-250 RPM.

*A. phagocytophilum* (HZ) was cultured in HL60 cells with Roswell Park Memorial

Institute (RPMI) 1640 medium supplemented with 10% heat-inactivated fetal bovine serum

(Atlanta Biologicals, S11550) and 1X Glutamax (Gibco, 35050061). Cells were maintained

between $1\times10^5$ - $1\times10^6$ ml$^{-1}$ at 37°C, 5% $CO_2$. *A. phagocytophilum* was enumerated as

previously described[93]. Briefly, the percentage of infected cells is multiplied the average number

of microcolonies per cell, termed 'morulae' (5), the average bacteria per morulae (19) and the

average amount of bacteria typically recovered from the isolation procedure (50%). Host cell-

free *A. phagocytophilum* was isolated by syringe lysis with a 27 gauge needle as previously

described[3].

*B. burgdorferi* B31 (MSK5[94]) was grown in modified Barbour-Stoenner-Kelly (BSK) II

medium supplemented with 6% normal rabbit serum (NRS, Pel-Freez, 31126-5) at 37°C, 5%

$CO_2$[94,95]. Spirochete density and growth phase were monitored by dark field microscopy. Prior to

infection, plasmid profiles of all *B. burgdorferi* cultures were screened by PCR, as described

previously[94].

Uninfected *I. scapularis* ticks where provided by the Biodefense and Emerging Infectious

Diseases (BEI) Research Resources Repository from the National Institute of Allergy and

Infectious Diseases (NIAID) ([www.beiresources.org](www.beiresources.org)) at the National Institutes of Health (NIH) or

from Oklahoma State University (Stillwater, OK, USA). Ticks were maintained in a 23°C

incubator with 16/8 hours light/dark photoperiods and 95-100% relative humidity. C3H/HeJ mice

were purchased from Jackson Laboratories and C57BL/6 mice were obtained from colonies

maintained at Washington State University. 6-10 week old male mice were used for all

experiments. C57BL/6 mice were infected intraperitoneally with $1\times10^7$ host cell-free *A.*

*phagocytophilum* HZ. C3H/HeJ mice were inoculated intradermally with $1\times10^5$ low passage *B.*

*burgdorferi*. All mice were confirmed for infection status prior to tick placement by collecting 25-

50 µl of blood from the lateral saphenous vein of each mouse 7 days post infection. *A.*

*phagocytophilum* burdens were enumerated by quantitative PCR (*16s* relative to mouse *β-*

*actin*[96,97]). *B. burgdorferi*-infected blood was subcultured in BSK-II media and examined for the

presence of spirochetes by dark field microscopy[98,99]. Experiments involving mice were carried

out according to guidelines and protocols approved by the American Association for

Accreditation of Laboratory Animal Care (AAALAC) and by the Office of Campus Veterinarian at

Washington State University (Animal Welfare Assurance A3485-01). The animals were housed

and maintained in an AAALAC-accredited facility at Washington State University in Pullman,

WA. All procedures were approved by the Washington State University Biosafety and Animal

Care and Use Committees.

*D. melanogaster and tick cell cultures*

*D. melanogaster* S2* cells were cultured with Schneider's *Drosophila* Medium (Gibco,

21720024) supplemented with 10% heat inactivated FBS (Sigma, SH30070) and 1X Glutamax.

Cell were maintained in T75 culture flasks (Corning, 353136) at 28°C with 0% $CO_2$.

The *I. scapularis* tick cell line, ISE6, was cultured at 32°C, 1% $CO_2$ in L15C-300 medium

supplemented with 10% heat inactive FBS (Sigma, F0926), 10% Tryptose Phosphate Broth

(TPB, BD, B260300) and 0.1% Lipoprotein Bovine Cholesterol (LPBC, MP Biomedicals,

219147680)[100]. The *D. andersoni* tick cell line, DAE100, was maintained at 34°C and cultured in

L15B medium supplemented with 5% FBS, 10% TBP, and 1% LPBC as previously

described[101,102].

*Polyacrylamide gel electrophoresis and Western blotting*

Protein concentrations were quantified using BCA assays per manufacture protocol

(Pierce, 23225). 10-15 µg of protein per sample were separated on a 4-15% MP TGX precast

cassette (Bio-Rad, 4561083) at 100V for 1 hour 25 minutes before being transferred to a PVDF

membrane. Membranes were blocked with 5% milk in PBS-T (1X phosphate-buffered saline

containing 0.1% Tween-20) for 1-2 hours at room temperature before being incubated at 4°C

overnight with a primary antibody in PBS-T and 5% BSA (Bovine Serum Albumin) or 0.5%-5%

milk. Primary antibodies used for immunoblotting are as follows: α-phospho-IRE1α (Abcam,

ab124945, 1:1000), α-Actin (Sigma, A2103, 1:1000), α-HA (Pierce, 26183, 1:1000), and α-

FLAG-HRP (Sigma, A8592, 1:500). Secondary antibodies were applied for 1-2 hours at room

temperature and are as follows: Goat α-Rabbit-HRP (Abcam, ab97051, 1:5000), Donkey α-

Rabbit-HRP (Thermo Fisher Scientific, A16023, 1:2000), Rabbit α-Mouse-HRP (Bio-Rad,

STAR13B, 1:2000), and Rec-G-Protein-HRP (Thermo Fisher Scientific, 101223, 1:2000). Blots

were visualized with Enhanced Chemiluminescence (ECL) Western blotting substrate (Thermo

Fisher Scientific, 32106). If necessary, blots were stripped with Western Blot Stripping Buffer

(Thermo Fisher Scientific, 21059) for 15-20 minutes at room temperature with shaking.

*Plasmid construction*

Both IRE1α and TRAF2 were codon optimized for expression in human cell lines

(GenScript). Primers listed in Supplementary Table 1 were used to amplify full length *I.*

*scapularis ire1α* for cloning into pCMV/hygro-Negative Control Vector (SinoBiological, CV005)

with *HindIII* sites. Full length *I. scapularis traf2* was amplified and cloned into pCMV-HA (New

MCS) vector (Received as a gift from Christopher A. Walsh; Addgene plasmid #32530) using

*XhoI* and *EcoRV*. All constructs were confirmed by sequencing (Eurofins Genomics).

*Maintenance and Transfection of HEK 293T cells*

HEK 293T cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Sigma,

D6429) supplemented with 10% heat inactivated FBS (Atlanta Biologicals, S11550) and 1X

Glutamax. Cells were maintained in T75 culture flasks (Corning, 353136) at 37°C, 5% $CO_2$. For

transfection, $1\times10^6$ HEK 293T cells were seeded into 6-well plates and allowed to attach

overnight. The following day cells were transfected with 2.5 µg of pCMV-TRAF2-HA and/or

pCMV-IRE1α-FLAG plasmid DNA using 10 µl of Lipofectamine 2,000 (Invitrogen, 11668027) in

Opti-MEM I Reduced Serum Medium (Gibco, 31985062). After 5 hours, media containing the

plasmid-Lipofectamine 2,000 complex was removed and replaced with complete DMEM for 48

hours at 33°C, 5% $CO_2$. The transfected cells were lysed with 500 µl of 25 mM Tris-HCL pH 7.4,

150 mM NaCl, 1% NP-40, 1 mM EDTA, 5% glycerol with 1X protease and phosphatase inhibitor

cocktail (Thermo Scientific, 78440) for 15 minutes on ice.

*Co-immunoprecipitation assay*

IRE1α-FLAG and TRAF2-HA expression was validated by immunoblotting whole cell

lysates with α-FLAG-HRP (Sigma, A8592, 1:500) and α-HA (Pierce, 26183, 1:1000). After

protein expression was confirmed, cross-linked agarose beads (α-FLAG M2: Sigma, A2220; α-

HA: Pierce, 26181) were washed 2X with TBS (50mM Tris, 150mM NaCl, pH 7.5) and incubated

with lysis buffer at 4°C for 1 hour. Approximately 1-2 mg of cell lysate was combined with 80 µl

(packed volume) of cross-linked agarose beads and incubated overnight at 4°C. Beads were

washed 3 times with TBS and protein was eluted by boiling in 50 µl of 4X Laemmli buffer for 5

minutes. Protein interactions were evaluated by immunoblot as described above.

*Template-based homology modeling of tick TRAF2 and the RNase/kinase domain of IRE1α*

A BLAST search in the Protein Data Bank (PDB) using the *I. scapularis* TRAF2

sequence returned the candidate template crystal structure of the TRAF-C domain from human

TRAF2 (39.64% sequence identity). The human TRAF2 crystal structure (PDB code 1CA9) was

used as reference for building the homology model of the TRAF-C domain and part of the

coiled-coil domain for *I. scapularis* TRAF2 (residues 176-357) in SWISS-MODEL[58,103].

QMEANdisCo was used to obtain a quality score, which defines how well the homology model

aligns to reference structures in the PDB. Scores closer to 1 indicate that the homology model

matches well to other reference structures[104]. Quality assessment of the TRAF2 homology

model in QMEANDisCo gave a score of 0.69. The GalaxyRefine server was used to then further

refine the *I. scapularis* TRAF2 homology model, which increased the quality score in

QMEANDisCo to 0.71[105].

A PDB BLAST search for *I. scapularis* IRE1α returned the candidate template crystal

structure of the RNase/kinase domain from human IRE1α (62.20% sequence identity). A

homology model for the cytosolic RNase/kinase domain of tick IRE1α (residues 525-944) was

built using the crystal structure of the RNase/kinase domain from humans (PDB code 6URC)

with SWISS-MODEL[56,103]. Quality assessment of the tick IRE1α homology model in

QMEANDisCo gave a score of 0.78.

*Prediction-driven docking of tick TRAF2 and the RNase/kinase domain of IRE1α*

A consensus interface predictor, CPORT (Consensus Prediction of interface Residues in

Transient complexes), was used to assign residues at the interface of TRAF2 and the IRE1α

RNase/kinase domain[59]. Predicted residues were used to define the docking interface between

TRAF2 and IRE1α for docking in HADDOCK2.2[60]. The docked model was immersed in a

solvent shell using the TIP3P water model and a short 300K MD simulation was ran to optimize

side chains and improve interaction energetics[60]. The cluster with the lowest Z-score was

chosen for further analysis. Docking models were then screened based on salt bridge

interactions at the docking interface and the model with the best chemical complementarity was

used in the final analysis. PyMOL version 2.2.3 was used for all distance measurements of salt-

bridge interactions (<4 Å cutoff) (The PyMOL Molecular Graphics System, Schrodinger, LLC).

*ROS assay*

ISE6 cells were seeded at a density of $1.68 \times 10^5$ cells per well in a black-walled, clear-

bottom 96-well plate (Thermo Scientific, 165305) with L15C-300 media. The cells were

maintained in growth conditions described above for the length of experiments. All wells were

treated for 1 hour with 10 µM 2', 7'-dichlorofluorescin diacetate (DCF-DA, Sigma, D6883) in

Ringer buffer (155 mM NaCl, 5 mM KCl, 1 mM $MgCl_2 \cdot 6H_2O$, 2 mM $NaH_2PO_4 \cdot H_2O$, 10 mM

HEPES, and 10 mM glucose)[106] alone or with 5 µM diphenyleneidonium chloride (DPI, Sigma,

D2926), 1 µM KIRA6 (Cayman Chemical, 19151) or 0.1% DMSO. Buffer was removed; cells

were washed with room temperature 1X PBS and incubated for 72 hours in L15C-300 alone or

with *A. phagocytophilum* (MOI 200), *B. burgdorferi* (MOI 200), 10 nM thapsigargin (TG, Sigma,

T9033), or 50 nM tunicamycin (TU, Sigma T7765). Fluorescence was measured at 504 nm

(excitation), 529 nm (emission). Data is graphed as fold change of relative fluorescence units

(RFU) normalized to the negative control ± standard errors of the means (SEM).

*Pharmacological treatments, RNAi silencing, quantitative reverse transcriptase-PCR*

ISE6 cells were seeded at $1\times10^6$ cells per well and DAE100 cells were seeded at $5\times10^5$

cells per well in a 24-well plate and pre-treated with KIRA6, thapsigargin or tunicamycin for

indicated times and concentrations prior to infection. Cells were infected with *A.*

*phagocytophilum* (ISE6) or *A. marginale* (DAE100) at an MOI 50 for 18 hours before collection

in Trizol (Invitrogen, 15596026). For DAE100 experiments, all incubations occurred at 34°C in a

BD campy bag with no gaspak. RNA was extracted using the Direct-zol RNA microprep Kit

(Zymo, R2062). cDNA was synthesized from 300-500 ng total RNA with the Verso cDNA

Synthesis Kit (Thermo Fisher Scientific, AB1453B). Bacterial burden and gene silencing were

assessed by quantitative reverse transcription-PCR (qRT-PCR) with the iTaq Universal SYBR

Green Supermix (Bio-Rad, 1725125) using primers listed in Supplementary Table 1. Cycle

conditions are as recommended by the manufacturer.

For transfection experiments, siRNAs and scrambled controls (scRNAs) were

synthesized following directions from the Silencer siRNA construction Kit (Invitrogen, AM1620)

using the primers listed in Supplementary Table 1. siRNA or scRNA (3 µg) was used to transfect

$1\times10^6$ ISE6 cells overnight with 2.5 µl of Lipofectamine 2,000. Cells were infected with *A.*

*phagocytophilum* (MOI 50) for 18 hours before being collected in Trizol. RNA was isolated and

transcripts were quantified by qRT-PCR as described above. All data are expressed as means ±

SEM.

*Ex vivo I. scapularis and D. andersoni*

Ten male and female unfed adult *I. scapularis* ticks were surface sterilized with

continuous agitation in 10% benzalkonium chloride (Sigma, 12060) for 10 minutes, washed

twice with sterile water, dried on sterile filter paper under aseptic conditions and transferred to a

sterile tube. Midgut and salivary glands were excised on a microscope slide in a pool of sterile

1X PBS + 100 I.U./ml penicillin and 100 µg/ml streptomycin (Gibco, 15140122). Tissues were

placed in individual wells of a 96-well plate (Costar, 3595) with 100 µl of L15C-300 and

incubated at 32°C with 1% $CO_2$. Tissues were treated with 1 µM of KIRA6 or 1% DMSO for 1

hour before the addition of $1x10^6$ *A. phagocytophilum*. 24 hours post-infection, samples were

collected following the addition of 100 µl of Trizol. Tissues were homogenized using

QIAshredder columns (Qiagen, 79654) according to the manufacturer's instructions prior to

RNA extraction and qRT-PCR analysis, performed as previously described.

Twenty male unfed adult *D. andersoni* ticks were surface sterilized and dissected as

above. Tissues were placed in individual wells of a 96-well plate with 100 µl of L15B. Tissues

were pretreated with KIRA6 or vehicle control (DMSO) as previously stated prior to the addition

of $1\times10^6$ *A. marginale* for 22 hours. Samples were collected and processed as above with qRT-

PCR standard curves using primers listed in Supplementary Table 1. All data are expressed as

means ± SEM

*I. scapularis microinjections*

*I. scapularis* nymphs were microinjected as described previously[93,100]. 10 µl Drummond

microdispensers (DrummondSci, 3000203G/X) were drawn to fine point needles using a

Narishige PC-100 micropipette puller. *I. scapularis* nymphs were microinjected with 25 nl of

siRNA or scRNA (~1000 ng/µl) into the anal pore using a Dummond Nanoject III Nanoliter

Injector (DrummondSci, 3000207). Ticks were allowed to rest overnight before being placed

between the ears and on the back of an infected mouse. Each group was placed on a single

mouse and fed to repletion (5-7 days). Nymphs were flash frozen with liquid nitrogen,

individually crushed with a plastic pestle and suspended in Trizol for RNA extraction.

*RNAi silencing in larvae*

*I. scapularis* larvae were pre-chilled at 4°C for 5 minutes. Approximately 150 larvae were

transferred to a 1.5 ml tube with 40-50 µl of either siRNA or scRNA (~1000 ng/µl). The tubes

were centrifuged at 3000 x g for 5 minutes to encourage submersion of the larvae in the dsRNA

and were then incubated overnight at 15°C. The following day, ticks were dried and rested

overnight before being placed on mice to feed until repletion (3-7 days). Larvae were flash

frozen in liquid nitrogen and individually crushed with a plastic pestle. Trizol was added before

proceeding to RNA isolation and qRT-PCR analysis, performed as previously stated. All data

are expressed as means ± SEM.

*xbp1 PCR and agarose gel electrophoresis*

RNA was isolated from both ISE6 cells or replete *I. scapularis* nymphs (uninfected, *A.*

*phagocytophilum*-infected or *B. burgdorferi*-infected). ISE6 cells were treated with either 0.5 µM

thapsigargin or *A. phagocytophilum* at an MOI of 50. Cells were collected 1, 3, 8, and 24 hours

post-treatment in Trizol. RNA was isolated and cDNA synthesized as previously described. The

cleavage status of *xbp1* was assessed via PCR using DreamTaq Green PCR Mastermix

(Thermo Scientific, K1082), using the *xbp1* primers listed in Supplementary Table 1 and the

cycling protocol according to the manufacturer's instructions. Samples were analyzed using a

3% agarose (Thermo Fisher, BP160) gel in 1X Tris-Borate EDTA (TBE, Thermo Fisher,

BP1333) with 0.5 µg/ml of ethidium bromide (Thermo Fisher, BP102) and imaged with a Protein

Simple AlphaImager HP system.

*UPR and IMD Gene Expression Profiling*

Untreated *I. scapularis* nymphs were fed to repletion on *A. phagocytophilum*-infected

mice or uninfected mice and frozen. The expression levels of UPR genes were assessed in

individual ticks by qRT-PCR as previously described. Primers specific for *bip*, *ire1α*, *xbp1*, and

*traf2* are listed in Supplementary Table 1. Data are expressed as means ± SEM.

519           $1\times10^6$ *D. melanogaster* S2* cells were seeded in Schneider's media with 1 µM of 20-

hydroxyecdysone to prime the IMD pathway, as previous reported[107]. Cells were treated with

indicated concentrations of thapsigargin for 6 hours or with 10 µM of KIRA6 for 1 hour prior to

infection with *A. phagocytophilum* (MOI 50) or *B. burgdorferi* (MOI 50) for 6 hours. Samples

were collected in Trizol and RNA was isolated. IMD pathway and Toll pathway-specific AMPs

were quantified by qRT-PCR as previously described with data expressed with primers listed in

Supplementary Table 1.

*Gene alignment*

527           UPR gene sequences were identified by querying the *I. scapularis* genome with *Homo*

*sapiens* protein sequences using NCBI (National Center for Biotechnology Information) protein

BLAST. Human sequences include BiP (NP_005338.1), IRE1α (NP_001424.3), TRAF2

(NP_066961.2) and XBP1 (NP_005071.2). Human and tick sequences were aligned using Jal

view[108]. Shaded regions indicate amino acid physiochemical property conservation. EMBL-EBI

(European Bioinformatics Institute) Pfam 34.0 was used to identify and annotate protein

domains[109].

*Statistical analysis*

535           *In vitro* experiments were performed with 3-5 replicates. *In vivo* experiments involved the

use of 10-20 ticks. Data were expressed as means ± SEM and analyzed with either unpaired

Student's t-test or Welch's t-test. Calculations and graphs were created with GraphPad Prism

version 9.0. $P < 0.05$ was considered statistically significant.

**REFERENCES**

1.    WHO | Vector-borne diseases. *WHO* http://www.who.int/mediacentre/factsheets/fs387/en/.
2.    Hillyer, J. F., Schmidt, S. L. & Christensen, B. M. Hemocyte-mediated phagocytosis and
543           melanization in the mosquito *Armigeres subalbatus* following immune challenge by
544           bacteria. *Cell Tissue Res.* **313**, 117–127 (2003).
545     3.    Garver, L. S., Dong, Y. & Dimopoulos, G. Caspar controls resistance to *Plasmodium*
546           *falciparum* in diverse *Anopheline* species. *PLOS Pathog* **5**, e1000335 (2009).
547     4.    Blumberg, B. J., Trop, S., Das, S. & Dimopoulos, G. Bacteria- and IMD pathway-
548           independent immune defenses against *Plasmodium falciparum* in *Anopheles gambiae*.
549           *PloS One* **8**, e72130 (2013).
550     5.    Oliva Chávez, A. S., Shaw, D. K., Munderloh, U. G. & Pedra, J. H. F. Tick humoral
551           responses: marching to the beat of a different drummer. *Front. Microbiol.* **8**, (2017).
552     6.    De la Fuente, J. *et al.* Tick-pathogen interactions and vector competence: identification of
553           molecular drivers for tick-borne diseases. *Front. Cell. Infect. Microbiol.* **7**, (2017).
554     7.    Lane, R. S. Competence of ticks as vectors of microbial agents with an emphasis on
555           *Borrelia burgdorferi*. In Sonenshine DE, Mather TN, editors. in *Ecological dynamics of tick-*
556           *borne zoonoses.* 45–67 (Oxford University Press, 1994).
557     8.    Goddard, J. *Infectious Diseases and Arthropods.* (Humana Press, 2000).
558     9.    Buchon, N., Silverman, N. & Cherry, S. Immunity in *Drosophila melanogaster*--from
559           microbial recognition to whole-organism physiology. *Nat. Rev. Immunol.* **14**, 796–810
560           (2014).
561     10.   Kleino, A. & Silverman, N. The *Drosophila* IMD pathway in the activation of the humoral
562           immune response. *Dev. Comp. Immunol.* **42**, 25–35 (2014).
563     11.   Kaneko, T. *et al.* PGRP-LC and PGRP-LE have essential yet distinct functions in the
564           *Drosophila* immune response to monomeric DAP-type peptidoglycan. *Nat. Immunol.* **7**,
565           715–723 (2006).
566     12.   Naitza, S. *et al.* The *Drosophila* immune defense against gram-negative infection requires
567           the death protein dFADD. *Immunity* **17**, 575–581 (2002).
568     13.   Meinander, A. *et al.* Ubiquitylation of the initiator caspase DREDD is required for innate
569           immune signaling. *EMBO J.* **31**, 2770–2783 (2012).
570     14.   Paquette, N. *et al.* Caspase-mediated cleavage, IAP binding, and ubiquitination: linking
571           three mechanisms crucial for *Drosophila* NF-kappaB signaling. *Mol. Cell* **37**, 172–182
572           (2010).
573     15.   Palmer, W. J. & Jiggins, F. M. Comparative genomics reveals the origins and diversity of
574           arthropod immune systems. *Mol. Biol. Evol.* **32**, 2111–2129 (2015).
575     16.   Arp, A. P., Hunter, W. B. & Pelz-Stelinski, K. S. Annotation of the Asian citrus psyllid
576           genome reveals a reduced innate immune system. *Front. Physiol.* **0**, (2016).
577     17.   Saha, S. *et al.* Improved annotation of the insect vector of citrus greening disease:
578           biocuration by a diverse genomics community. *Database J. Biol. Databases Curation* **2017**,
579           bax032 (2017).
580     18.   Shelby, K. S. Functional immunomics of the squash bug, *Anasa tristis* (De Geer)
581           (Heteroptera: *Coreidae*). *Insects* **4**, 712–730 (2013).
582     19.   Zhang, C.-R. *et al.* The immune strategy and stress response of the Mediterranean
583           species of the *Bemisia tabaci* complex to an orally delivered bacterial pathogen. *PLOS*
584           *ONE* **9**, e94477 (2014).
585     20.   Chen, W. *et al.* The draft genome of whitefly *Bemisia tabaci* MEAM1, a global crop pest,
586           provides novel insights into virus transmission, host adaptation, and insecticide resistance.
587           *BMC Biol.* **14**, 110 (2016).
588     21.   Kim, J. H. *et al.* Comparison of the humoral and cellular immune responses between body
589           and head lice following bacterial challenge. *Insect Biochem. Mol. Biol.* **41**, 332–339 (2011).

22.  Elsik, C. G. The pea aphid genome sequence brings theories of insect defense into
question. *Genome Biol.* **11**, 106 (2010).

23.  Benoit, J. B. *et al.* Unique features of a global human ectoparasite identified through
sequencing of the bed bug genome. *Nat. Commun.* **7**, 10165 (2016).

24.  Kirkness, E. F. *et al.* Genome sequences of the human body louse and its primary
endosymbiont provide insights into the permanent parasitic lifestyle. *Proc. Natl. Acad. Sci.*
**107**, 12168–12173 (2010).

25.  Bechsgaard, J. *et al.* Comparative genomic study of arachnid immune systems indicates
loss of beta-1,3-glucanase-related proteins and the immune deficiency pathway. *J. Evol.*
*Biol.* **29**, 277–291 (2016).

26.  Capelli-Peixoto, J. *et al.* The transcription factor Relish controls *Anaplasma marginale*
infection in the bovine tick *Rhipicephalus microplus*. *Dev. Comp. Immunol.* **74**, 32–39
(2017).

27.  Gulia-Nuss, M. *et al.* Genomic insights into the *Ixodes scapularis* tick vector of Lyme
disease. *Nat. Commun.* **7**, 10507 (2016).

28.  Shaw, D. K. *et al.* Infection-derived lipids elicit an immune deficiency circuit in arthropods.
*Nat. Commun.* **8**, ncomms14401 (2017).

29.  Rosa, R. D. *et al.* Exploring the immune signalling pathway-related genes of the cattle tick
*Rhipicephalus microplus*: from molecular characterization to transcriptional profile upon
microbial challenge. *Dev. Comp. Immunol.* **59**, 1–14 (2016).

30.  Carroll, E. E. M. *et al.* p47 licenses activation of the immune deficiency pathway in the tick
*Ixodes scapularis*. *Proc. Natl. Acad. Sci.* **116**, 205–210 (2019).

31.  Severo, M. S. *et al.* The E3 ubiquitin ligase XIAP restricts *Anaplasma phagocytophilum*
colonization of *Ixodes scapularis* ticks. *J. Infect. Dis.* **208**, 1830–1840 (2013).

32.  Mesquita, R. D. *et al.* Genome of *Rhodnius prolixus*, an insect vector of Chagas disease,
reveals unique adaptations to hematophagy and parasite infection. *Proc. Natl. Acad. Sci.*
*U. S. A.* **112**, 14936–14941 (2015).

33.  Salcedo-Porras, N., Guarneri, A., Oliveira, P. L. & Lowenberger, C. *Rhodnius prolixus*:
identification of missing components of the IMD immune signaling pathway and functional
characterization of its role in eliminating bacteria. *PLoS ONE* **14**, e0214794 (2019).

34.  Zumaya-Estrada, F. A., Martínez-Barnetche, J., Lavore, A., Rivera-Pomar, R. & Rodríguez,
M. H. Comparative genomics analysis of triatomines reveals common first line and
inducible immunity-related genes and the absence of IMD canonical components among
hemimetabolous arthropods. *Parasit. Vectors* **11**, 48 (2018).

35.  Rolandelli, A., Nascimento, A. E. C., Silva, L. S., Rivera-Pomar, R. & Guarneri, A. A.
modulation of IMD, Toll, and Jak/STAT immune pathways genes in the fat body of
*Rhodnius prolixus* during *Trypanosoma rangeli* infection. *Front. Cell. Infect. Microbiol.* **0**,
(2021).

36.  Grootjans, J., Kaser, A., Kaufman, R. J. & Blumberg, R. S. The unfolded protein response
in immunity and inflammation. *Nat. Rev. Immunol.* **16**, 469–484 (2016).

37.  Hetz, C. The unfolded protein response: controlling cell fate decisions under ER stress and
beyond. *Nat. Rev. Mol. Cell Biol.* **13**, 89–102 (2012).

38.  Schröder, M. & Kaufman, R. J. The mammalian unfolded protein response. *Annu. Rev.*
*Biochem.* **74**, 739–789 (2005).

39.  Zhou, Y. *et al.* Porcine reproductive and respiratory syndrome virus infection induces
stress granule formation depending on protein kinase R-like endoplasmic reticulum kinase
(PERK) in MARC-145 cells. *Front. Cell. Infect. Microbiol.* **7**, (2017).

40.  Kültz, D. Evolution of the cellular stress proteome: from monophyletic origin to ubiquitous
function. *J. Exp. Biol.* **206**, 3119–3124 (2003).

41.  Kültz, D. Molecular and evolutionary basis of the cellular stress response. *Annu. Rev.*
*Physiol.* **67**, 225–257 (2004).

42.  Celli, J. & Tsolis, R. M. Bacteria, the endoplasmic reticulum and the unfolded protein
response: friends or foes? *Nat. Rev. Microbiol.* **13**, 71–82 (2015).

43.  Keestra-Gounder, A. M. *et al.* NOD1 and NOD2 signalling links ER stress with
inflammation. *Nature* **532**, 394–397 (2016).

44.  Byndloss, M. X., Keestra-Gounder, A. M., Bäumler, A. J. & Tsolis, R. M. NOD1 and NOD2:
new functions linking endoplasmic reticulum stress and inflammation. *DNA Cell Biol.* **35**,
311–313 (2016).

45.  Rosche, K. L., Sidak-Loftis, L. C., Hurtado, J., Fisk, E. A. & Shaw, D. K. Arthropods under
pressure: stress responses and immunity at the pathogen-vector interface. *Front. Immunol.*
**11**, (2021).

46.  Thastrup, O., Cullen, P. J., Drobak, B. K., Hanley, M. R. & Dawson, A. P. Thapsigargin, a
tumor promoter, discharges intracellular Ca2+ stores by specific inhibition of the
endoplasmic reticulum Ca2(+)-ATPase. *Proc. Natl. Acad. Sci.* **87**, 2466–2470 (1990).

47.  Wu, J. *et al.* Tunicamycin specifically aggravates ER stress and overcomes
chemoresistance in multidrug-resistant gastric cancer cells by inhibiting N-glycosylation. *J.*
*Exp. Clin. Cancer Res.* **37**, 272 (2018).

48.  Zhang, L., Zhang, C. & Wang, A. divergence and conservation of the major UPR branch
IRE1-bZIP signaling pathway across eukaryotes *Sci. Rep.* **6**, (2016).

49.  Adams, C. J., Kopp, M. C., Larburu, N., Nowak, P. R. & Ali, M. M. U. Structure and
molecular mechanism of ER stress signaling by the unfolded protein response signal
activator IRE1. *Front. Mol. Biosci.* **6**, (2019).

50.  Hu, P., Han, Z., Couvillon, A. D., Kaufman, R. J. & Exton, J. H. Autocrine tumor necrosis
factor alpha links endoplasmic reticulum stress to the membrane death receptor pathway
through IRE1alpha-mediated NF-kappaB activation and down-regulation of TRAF2
expression. *Mol. Cell. Biol.* **26**, 3071–3084 (2006).

51.  Back, S. H., Lee, K., Vink, E. & Kaufman, R. J. Cytoplasmic IRE1α-mediated XBP1 mRNA
splicing in the absence of nuclear processing and endoplasmic reticulum stress*. *J. Biol.*
*Chem.* **281**, 18691–18706 (2006).

52.  Calfon, M. *et al.* IRE1 couples endoplasmic reticulum load to secretory capacity by
processing the XBP-1 mRNA. *Nature* **415**, 92–96 (2002).

53.  Shen, X. *et al.* Complementary signaling pathways regulate the unfolded protein response
and are required for *C. elegans* development. *Cell* **107**, 893–903 (2001).

54.  Yoshida, H., Matsui, T., Yamamoto, A., Okada, T. & Mori, K. XBP1 mRNA is induced by
ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription
factor. *Cell* **107**, 881–891 (2001).

55.  Ghosh, R. *et al.* Allosteric inhibition of the IRE1α RNase preserves cell viability and
function during endoplasmic reticulum stress. *Cell* **158**, 534–548 (2014).

56.  Harnoss, J. M. *et al.* Disruption of IRE1α through its kinase domain attenuates multiple
myeloma. *Proc. Natl. Acad. Sci.* **116**, 16420–16429 (2019).

57.  Mm, A. *et al.* Structure of the IRE1 autophosphorylation complex and implications for the
unfolded protein response. *EMBO J.* **30**, 894–905 (2011).

58.  Park, Y. C., Burkitt, V., Villa, A. R., Tong, L. & Wu, H. Structural basis for self-association
and receptor recognition of human TRAF2. *Nature* **398**, 533–538 (1999).

59.  Vries, S. J. de & Bonvin, A. M. J. J. CPORT: A consensus interface predictor and its
performance in prediction-driven docking with HADDOCK. *PLOS ONE* **6**, e17695 (2011).

60.  van Zundert, G. C. P. *et al.* The HADDOCK2.2 web server: user-friendly integrative
modeling of biomolecular complexes. *J. Mol. Biol.* **428**, 720–725 (2016).

61.  Hodzic, E. *et al.* Acquisition and transmission of the agent of human granulocytic
ehrlichiosis by *Ixodes scapularis* ticks. *J. Clin. Microbiol.* **36**, 3574–3578 (1998).

62.  Fellous, S. & Lazzaro, B. P. Potential for evolutionary coupling and decoupling of larval
and adult immune gene expression. *Mol. Ecol.* **20**, 1558–1567 (2011).

63. League, G. P., Estévez-Lao, T. Y., Yan, Y., Garcia-Lopez, V. A. & Hillyer, J. F. *Anopheles gambiae* larvae mount stronger immune responses against bacterial infection than adults: evidence of adaptive decoupling in mosquitoes. *Parasit. Vectors* **10**, 367 (2017).

64. Critchlow, J. T., Norris, A. & Tate, A. T. The legacy of larval infection on immunological dynamics over metamorphosis. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, (2019).

65. Galay, R. L. *et al.* Induction of gene silencing in *Haemaphysalis longicornis* ticks through immersion in double-stranded RNA. *Ticks Tick-Borne Dis.* **7**, 813–816 (2016).

66. Liu, L. *et al. Ixodes scapularis* salivary gland protein P11 facilitates migration of *Anaplasma phagocytophilum* from the tick gut to salivary glands. *EMBO Rep.* **12**, 1196–1203 (2011).

67. Hodzic, E. *et al.* Granulocytic ehrlichiosis in the laboratory mouse. *J. Infect. Dis.* **177**, 737–745 (1998).

68. Grabowski, J. M. *et al.* Dissecting flavivirus biology in salivary gland cultures from fed and unfed *Ixodes scapularis* (black-legged tick). *mBio* **10**, (2019).

69. Grabowski, J. M., Offerdahl, D. K. & Bloom, M. E. The use of *ex vivo* organ cultures in tick-borne virus research. *ACS Infect. Dis.* **4**, 247–256 (2018).

70. Ha, E.-M. *et al.* Coordination of multiple dual oxidase–regulatory pathways in responses to commensal and infectious microbes in *Drosophila* gut. *Nat. Immunol.* **10**, 949–957 (2009).

71. Chakrabarti, S., Poidevin, M. & Lemaitre, B. The *Drosophila* MAPK p38c regulates oxidative stress and lipid homeostasis in the intestine. *PLOS Genet.* **10**, e1004659 (2014).

72. Lin, M. & Rikihisa, Y. Degradation of p22phox and inhibition of superoxide generation by *Ehrlichia chaffeensis* in human monocytes. *Cell. Microbiol.* **9**, 861–874 (2007).

73. Boylan, J. A. & Gherardini, F. C. Determining the cellular targets of reactive oxygen species in *Borrelia burgdorferi*. *Methods Mol. Biol. Clifton NJ* **431**, 213–221 (2008).

74. Hyde, J. A., Shaw, D. K., Smith Iii, R., Trzeciakowski, J. P. & Skare, J. T. The BosR regulatory protein of *Borrelia burgdorferi* interfaces with the RpoS regulatory pathway and modulates both the oxidative stress response and pathogenic properties of the lyme disease spirochete. *Mol. Microbiol.* **74**, 1344–1355 (2009).

75. Hyde, J. A., Shaw, D. K., Smith, R., Trzeciakowski, J. P. & Skare, J. T. Characterization of a conditional bosR mutant in *Borrelia burgdorferi*. *Infect Immun* **78**, 265–274 (2010).

76. Abuaita, B. H., Burkholder, K. M., Boles, B. R. & O'Riordan, M. X. The endoplasmic reticulum stress sensor inositol-requiring enzyme 1α augments bacterial killing through sustained oxidant production. *mBio* **6**, (2015).

77. Abuaita, B. H., Schultz, T. L. & O'Riordan, M. X. Mitochondria-derived vesicles deliver antimicrobial reactive oxygen species to control phagosome-localized *Staphylococcus aureus*. *Cell Host Microbe* **24**, 625-636.e5 (2018).

78. Okafor, C. C. *et al.* Factors associated with seroprevalence of *Anaplasma marginale* in Kentucky cattle. *Vet. Parasitol. Reg. Stud. Rep.* **13**, 212–219 (2018).

79. Dunning Hotopp, J. C. *et al.* Comparative genomics of emerging human ehrlichiosis agents. *PLoS Genet.* **2**, e21 (2006).

80. Beck, G., Benach, J. L. & Habicht, G. S. Isolation, preliminary chemical characterization, and biological activity of *Borrelia burgdorferi* peptidoglycan. *Biochem. Biophys. Res. Commun.* **167**, 89–95 (1990).

81. Rikihisa, Y. *Anaplasma phagocytophilum* and *Ehrlichia chaffeensis*: subversive manipulators of host cells. *Nat. Rev. Microbiol.* **8**, 328–339 (2010).

82. Takayama, K., Rothenberg, R. J. & Barbour, A. G. Absence of lipopolysaccharide in the lyme disease spirochete, *Borrelia burgdorferi*. *Infect. Immun.* **55**, 2311–2313 (1987).

83. Holt, S. C. Anatomy and chemistry of spirochetes. *Microbiol. Rev.* **42**, 114–160 (1978).

84. Shaw, D. K. *et al.* Vector immunity and evolutionary ecology: the harmonious dissonance. *Trends Immunol.* **39**, 862–873 (2018).

85. Moret, Y. Survival for immunity: the price of immune system activation for bumblebee workers. *Science* **290**, 1166–1168 (2000).

86.  Truman, J. W. & Riddiford, L. M. The origins of insect metamorphosis. *Nature* **401**, 447–
452 (1999).
87.  Misof, B. *et al.* Phylogenomics resolves the timing and pattern of insect evolution. *Science*
**346**, 763–767 (2014).
88.  Tettamanti, G. & Casartelli, M. Cell death during complete metamorphosis. *Philos. Trans.*
*R. Soc. B Biol. Sci.* **374**, 20190065 (2019).
89.  Grigoryeva, L. A. Morpho-functional changes in the midgut of ixodid ticks (Acari: *Ixodidae*)
during the life cycle. *Entomol. Rev.* **90**, 405–409 (2010).
90.  Giribet, G. & Edgecombe, G. D. The phylogeny and evolutionary history of arthropods.
*Curr. Biol.* **29**, R592–R602 (2019).
91.  Truman, J. W. & Riddiford, L. M. The evolution of insect metamorphosis: a developmental
and endocrine view. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20190070 (2019).
92.  Thomas, G. W. C. *et al.* Gene content evolution in the arthropods. *Genome Biol.* **21**, 15
(2020).
93.  Shaw, D. K. *et al.* Infection-derived lipids elicit an immune deficiency circuit in arthropods.
*Nat. Commun.* **8**, 14401 (2017).
94.  Labandeira-Rey, M. & Skare, J. T. Decreased infectivity in *Borrelia burgdorferi* strain B31
is associated with loss of linear plasmid 25 or 28-1. *Infect. Immun.* **69**, 446–455 (2001).
95.  Zückert, W. R. Laboratory maintenance of *Borrelia burgdorferi*. *Curr. Protoc. Microbiol.* **4**,
12C.1.1-12C.1.10 (2007).
96.  Sukumaran, B. *et al.* Receptor interacting protein-2 contributes to host defense against
*Anaplasma phagocytophilum* infection. *FEMS Immunol. Med. Microbiol.* **66**, 211–219
(2012).
97.  Pedra, J. H. F. *et al.* ASC/PYCARD and Caspase-1 regulate the IL-18/IFN-γ axis during
*Anaplasma phagocytophilum* infection. *J. Immunol.* **179**, 4783–4791 (2007).
98.  James, A. E., Rogovskyy, A. S., Crowley, M. A. & Bankhead, T. Characterization of a DNA
adenine methyltransferase gene of *Borrelia hermsii* and its dispensability for murine
infection and persistence. *PLOS ONE* **11**, e0155798 (2016).
99.  Bankhead, T. & Chaconas, G. The role of VlsE antigenic variation in the Lyme disease
spirochete: persistence through a mechanism that differs from other pathogens. *Mol.*
*Microbiol.* **65**, 1547–1558 (2007).
100. Severo, M. S. *et al.* The E3 ubiquitin ligase XIAP restricts *Anaplasma phagocytophilum*
colonization of *Ixodes scapularis* ticks. *J. Infect. Dis.* **208**, 1830–1840 (2013).
101. Munderloh, U. G. & Kurtti, T. J. Formulation of medium for tick cell culture. *Exp. Appl.*
*Acarol.* **7**, 219–229 (1989).
102. Simser, J. A., Palmer, A. T., Munderloh, U. G. & Kurtti, T. J. Isolation of a spotted fever
group *Rickettsia, Rickettsia peacockii*, in a rocky mountain wood tick, *Dermacentor*
*andersoni*, cell line. *Appl. Environ. Microbiol.* **67**, 546–552 (2001).
103. Waterhouse, A. *et al.* SWISS-MODEL: Homology modelling of protein structures and
complexes. *Nucleic Acids Res.* **46**, W296–W303 (2018).
104. Benkert, P., Biasini, M. & Schwede, T. Toward the estimation of the absolute quality of
individual protein structure models. *Bioinformatics* **27**, 343–350 (2011).
105. Ko, J., Park, H., Heo, L. & Seok, C. GalaxyWEB server for protein structure prediction and
refinement. *Nucleic Acids Res.* **40**, W294–W297 (2012).
106. Abuaita, B. H., Schultz, T. L. & O'Riordan, M. X. Mitochondria-derived vesicles deliver
antimicrobial reactive oxygen species to control phagosome-localized *Staphylococcus*
*aureus*. *Cell Host Microbe* **24**, 625-636.e5 (2018).
107. Rus, F. *et al.* Ecdysone triggered PGRP-LC expression controls *Drosophila* innate
immunity. *EMBO J.* **32**, 1626–1638 (2013).

108. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics*
**25**, 1189–1191 (2009).
109. Mistry, J. *et al.* Pfam: the protein families database in 2021. *Nucleic Acids Res.* **49**, D412–
D419 (2021).
797

 **FIGURE LEGENDS**

**Figure 1 / The tick UPR responds to and restricts bacterial colonization.** (**a**) Graphic

representation of the unfolded protein response (UPR) in mammals. (**b**) UPR gene expression

in *A. phagocytophilum*-infected *I. scapularis* nymphs relative to uninfected controls (dotted line).

Each point is representative of 1 nymph. Gene expression was quantified by qRT-PCR. (**c-e**)

ISE6 cells ($1 \times 10^6$) were infected with *A. phagocytophilum* MOI 50 for 18 hours following a 24

hour treatment with either (**c**) thapsigargin, (**d**) tunicamycin or (**e**) siRNA targeting the negative

regulator *bip*. Gene silencing and *A. phagocytophilum* load (16s rDNA) was measured by qRT-

PCR. All data shown are representative of 5 biological replicates with least two technical

replicates ± SEM.. Student's t-test. *P < 0.05. scRNA, scrambled RNA; siRNA, small interfering

RNA; NS, not significant. See also Supplementary Figure 1 and Supplementary Table 1.

**Figure 2 | The IRE1α branch of the UPR is induced by *A. phagocytophilum* through**

**TRAF2. (a)** Phosphorylated IRE1α immunoblot against ISE6 ($1 \times 10^6$) cells treated with either the

ER stress inducer tunicamycin (24 hours) or infected with *A. phagocytophilum* (*A.p.*). Protein

expression differences were quantified by ImageJ and are expressed as a ratio of

phosphorylated IRE1α (~110 kDa) to the internal loading control, β-actin (45 kDa). ISE6 cells

were treated with (**b**) the IRE1α inhibitor KIRA6 (1 hour) or (**c** and **f**) siRNAs to silence gene

expression prior to *A. phagocytophilum* (*A.p.*; MOI 50) infection for 18 hours. Gene silencing

and *A. phagocytophilum* burden were measured by qRT-PCR. Experiments shown are

representative of at least two technical replicates ± SEM. Student's t-test. *P < 0.05. (**d**) ISE6

cells ($1 \times 10^6$) were either untreated (-), stimulated with 0.5 μM of thapsigargin (TG) or infected

with *A. phagocytophilum* (*A.p.*; MOI 50) for indicated time points. (**e**) Replete *I. scapularis*

nymphs fed either on uninfected mice (-), *A. phagocytophilum* (*A.p.*)-infected or *B. burgdorferi*

(*B.b.*)-infected mice. (**d-e**) cDNA was synthesized from RNA and used to evaluate *xbp1* splicing

by PCR. Samples were analyzed on a 3% agarose gel. scRNA, scrambled RNA; siRNA, small

interfering RNA. See also Supplementary Figure 1 and Supplementary Table 1.

**Figure 3 | IRE1α - TRAF2 molecular interactions in *I. scapularis*.** (**a**) The interfaces assigned

by CPORT for the TRAF2 trimer and IRE1α homology models. Active central (cyan) and

passive peripheral (navy blue) residues shown as spheres were used to filter the docking

solutions in HADDOCK 2.2. (**b**) Final docking model between TRAF2 and IRE1α places TRAF2

away from the dimer interface and the C-terminus of IRE1α (black circle), which anchors IRE1α

to the ER. (**c**) Salt bridges were determined between all three chains of TRAF2 and IRE1α with

a measured distance between 2.7 - 2.8 Å. Negatively charged Asp and Glu residues (red

spheres) pair with positively charged Lys, Arg, and His residues (blue spheres). (**d**)

Immunoprecipitation (IP) analysis followed by Western blotting (WB) showing interaction

between FLAG-tagged IRE1α and HA-tagged TRAF2 expressed in HEK 293T cells. WB is

representative of two biological replicates. See also Supplementary Figure 2 and

Supplementary Table 1.

**Figure 4 | Vector competence for *A. phagocytophilum* is influenced by IRE1α and TRAF2**

**at multiple tick life stages *in vivo.*** *I. scapularis* (**a-b**) nymphs or (**c-d**) larvae had *ire1α* and

*traf2* expression silenced through RNAi prior to feeding on *A. phagocytophilum*-infected mice.

Silencing levels and bacterial load were measured in whole *I. scapularis* nymphs or larvae. (**e**)

Schematic of *ex vivo I. scapularis* midgut and salivary gland cultures. (**f-g**) Midguts and salivary

glands from *I. scapularis* adults were dissected, cultured and treated with 1 µM of KIRA6 (1

hour) followed by *A. phagocytophilum* infection for 24 hours. Silencing levels and *A.*

*phagocytophilum* load (16s rDNA) were measured by qRT-PCR. Each point represents 1 tick,

midgut, or pair of salivary glands (two technical replicates each), ± SEM. Welch's t-test. *P <

0.05. scRNA, scrambled RNA; siRNA, small interfering RNA. See also Supplementary Table 1.

**Figure 5 | IRE1α and TRAF2 restrict *B. burgdorferi* colonization *in vivo* at multiple tick life stages.** RNAi silencing of *ire1α* and *traf2* in *I. scapularis* (**a-b**) nymphs or (**c-d**) larvae prior to feeding on *B. burgdorferi*-infected mice. Silencing levels and *B. burgdorferi* (*flaB*) were measured in whole *I. scapularis* nymphs or larvae. Each point represents 1 tick (two technical replicates each) ± SEM. Welch's t-test. *P < 0.05. scRNA, scrambled RNA; siRNA, small interfering RNA. Also see Supplementary Table 1.

**Figure 6 | The UPR stimulates antimicrobial products in arthropod cells.** (**a**) Indicated concentrations of thapsigargin (TG) were used to treat S2* cells ($1 \times 10^6$) for 6 hours prior to examining gene expression differences. (**b-c**) S2* cells ($1 \times 10^6$) were pretreated with KIRA6 (1 hour) before (**b**) *A. phagocytophilum* (*A.p.*; MOI 50) or (**c**) *B. burgdorferi* (*B.b.*; MOI 50) infection (6 hours). Gene expression is relative to *rp49*. Dotted line denotes unstimulated controls. Data shown are representative of 4-5 biological replicates and two technical replicates. (**d**) Thapsigargin (TG; 10 nM) or tunicamycin (Tu; 50 nM) were used to stimulate ISE6 cells ($1.68 \times 10^5$). (**e-f**) ISE6 cells were pretreated with DPI (5 µM) or KIRA6 (1 µM) for 1 hour prior to (**e**) *A. phagocytophilum* (MOI 50) or (**f**) *B. burgdorferi* (MOI 50) infection. (**d-f**) ROS was determined by relative fluorescence units (RFU) after 72 hours. Data shown is representative of 3 biological replicates and 2 technical replicates, ± SEM. Student's t-test. *P < 0.05. (-), non-stimulated. DPI, diphenyleneidonium. See also Supplementary Figure 3 and Supplementary Table 1.

**Figure 7 | IRE1α and TRAF2-mediated pathogen restriction is conserved across arthropod vectors.** DAE100 cells ($5 \times 10^5$) were treated with indicated concentrations of (**a**) thapsigargin, (**b**) tunicamycin or (**c**) KIRA6 followed by infection with *A. marginale* (MOI 50) for 18 hours. Student's t-test. *P < 0.05. (**d**) Schematic of *ex vivo D. andersoni* midgut and salivary gland cultures. (**e-f**) Midguts and salivary glands from *D. andersoni* adults were dissected, cultured and treated with 1 µM of KIRA6 (1 hour) followed by *A. marginale* infection for 22

871    hours. *A. marginale* (*rpoH*) was quantified by qRT-PCR and relative to *β-actin.* Welch's t-test. *P

872    < 0.05. Each point representative of 1 tick, midgut, or pair of salivary glands (two technical

873    replicates), ± SEM. See also Supplementary Table 1.

# Figure 1 | The tick UPR responds to and restricts bacterial colonization

# Figure 2 | The IRE1α branch of the UPR is induced by *A. phagocytophilum* through TRAF2

**Figure 3 | IRE1α - TRAF2 molecular interactions in *I. scapularis***

# Figure 4 | Vector competence for *A. phagocytophilum* is influenced by IRE1α and TRAF2 at multiple tick life stages *in vivo*

# Figure 5 | IRE1α and TRAF2 restrict *B. burgdorferi* colonization *in vivo* at multiple tick life stages

# Figure 6 | The UPR stimulates antimicrobial products in arthropod cells

# Figure 7 | IRE1α and TRAF2-mediated pathogen restriction is conserved across arthropod vectors

**Acknowledgements**

We grateful to Ulrike Munderloh (University of Minnesota) for providing ISE6 and DAE100 tick cell lines; Jon Skare (Texas A&M Health Science Center) for providing *B. burgdorferi* B31 (MSK5); Alan Goodman (Washington State University) for providing S2* cells; BEI Resources and Oklahoma State University for *Ixodes scapularis* ticks, and Addgene plasmid #32530 received as a gift from Christopher A. Walsh.

**Funding**

This work is supported by the National Institutes of Health (R21AI139772 to D.K.S.), the WSU Intramural CVM grants program, funded in part by National Institute of Food and Agriculture and the, Joseph and Barbara Mendelson Endowment Research Fund (to D.K.S.) and Washington State University, College of Veterinary Medicine. Additional support to L.C.S-L. came from The Fowler Emerging Diseases Graduate Fellowship funded by Ralph and Maree Fowler. J.H. is a trainee under the Institutional Training Grant T32 from the National Institute of Allergy and Infection Diseases (T32GM008336). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Allergy and Infection Diseases or the National Institutes of Health.

**Author contributions**

L.C.S-L., K.L.R. and D.K.S. designed the study. L.C.S-L., K.L.R., N.P., J.K.U, J.H., and D.K.S. performed experiments. N.P. and J.W.P. performed homology modeling and IRE1α-TRAF2 docking. S.M.H. and J.K.U. contributed reagents and performed *D. andersoni* experiments. E.A.F. designed and constructed Supplementary Figure 3. L.C.S-L, K.L.R., N.P., J.W.P. and D.K.S. analyzed data. All authors provided intellectual input into the study. L.C.S-L., K.L.R., N.P. and D.K.S. wrote the manuscript; all authors contributed to editing.