A genetically linked pair of NLR immune receptors show contrasting patterns of evolution

Motoki Shimizu[1], Akiko Hirabuchi[1], Yu Sugihara[2], Akira Abe[1], Takumi Takeda[1], Michie Kobayashi[1,3], Yukie Hiraka[1], Eiko Kanzaki[1], Kaori Oikawa[1], Hiromasa Saitoh[1,4], Thorsten Langner[5], Mark J. Banfield[6], Sophien Kamoun[5] and Ryohei Terauchi[1, 2]

[1] Iwate Biotechnology Research Center, Kitakami, Iwate 024-0003, Japan

[2] Laboratory of Crop Evolution, Graduate School of Agriculture, Kyoto University, Mozume, Muko, Kyoto 617-0001, Japan

[3] Institute of Agrobiological Sciences, National Agriculture and Food Research Organization (NARO), Kannondai 2-1-2, Tsukuba, Ibaraki 305-8602, Japan

[4] Department of Molecular Microbiology, Tokyo University of Agriculture, Setagaya-ku, Tokyo 156-8502, Japan

[5] The Sainsbury Laboratory, University of East Anglia, Norwich, NR4 7UH, UK

[6] Department of Biochemistry and Metabolism, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK

Correspondence to:
Motoki Shimizu (m-shimizu@ibrc.or.jp) and
Ryohei Terauchi (terauchi@ibrc.or.jp)

Abstract

Throughout their evolution, plant nucleotide-binding leucine-rich-repeat receptors (NLRs) have acquired widely divergent unconventional integrated domains that enhance their ability to detect pathogen effectors. However, the functional dynamics that drive the evolution of NLRs with integrated domains (NLR-IDs) remain poorly understood. Here, we reconstructed the evolutionary history of an NLR locus prone to unconventional domain integration and experimentally tested hypotheses about the evolution of NLR-IDs. We show that the rice (*Oryza sativa*) NLR Pias recognizes the effector AVR-Pias of the blast fungal pathogen *Magnaporthe oryzae*. Pias consists of a functionally specialized NLR pair, the helper Pias-1 and the sensor Pias-2, and is allelic to the

previously characterized Pia pair of NLRs: the helper RGA4 and the sensor RGA5. Remarkably, Pias-2 carries a C-terminal DUF761 domain at a similar position to the heavy metal–associated (HMA) domain of RGA5. Phylogenomic analysis showed that Pias-2/RGA5 sensor NLRs have undergone recurrent genomic recombination within the genus *Oryza*, resulting in up to six sequence-divergent domain integrations. Allelic NLRs with divergent functions have been maintained trans-species in different *Oryza* lineages to detect sequence-divergent pathogen effectors. By contrast, Pias-1 has retained its NLR helper activity throughout evolution and is capable of functioning together with the divergent sensor-NLR RGA5 to recognize AVR-Pia. These results suggest that opposite selective forces have driven the evolution of paired NLRs: highly dynamic domain integration events maintained by balancing selection for sensor NLRs, in sharp contrast to purifying selection and functional conservation of immune signaling for helper NLRs.

Keywords:
nucleotide-binding leucine-rich-repeat receptors (NLRs), paired NLR, integrated domains, evolution, rice, *Magnaporthe oryzae*

Significance statement

Plants have evolved sophisticated defense mechanisms to fend off pathogens. Plant nucleotide-binding leucine-rich repeat receptor (NLR) proteins play crucial roles in detecting pathogen molecules inside plant cells and mounting defense responses. Here, we identified the *Pias* gene from rice, which encodes the NLR pair Pias-1 "helper" and Pias-2 "sensor." These proteins function together to detect the pathogen molecule AVR-Pias of *Magnaporthe oryzae* and defend against rice blast disease. *Pias* is allelic to the previously reported *Pia* gene. A comparison of Pias/Pia alleles among *Oryza* species showed that Pias/Pia helper is evolutionarily and functionally conserved, whereas Pias/Pia sensor shows highly dynamic evolution, with various host domains integrated into similar positions, allowing it to detect a wide variety of pathogen molecules.

Introduction

Plants are continually attacked by a multitude of microbial pathogens. Pathogens secrete effector molecules to enable the invasion of their hosts (Hogenhout et al. 2009). To counter this, plants have evolved a surveillance system detecting pathogen effectors in the cytosol, leading to effector-triggered immunity (ETI; Jones and Dangl, 2006). Nucleotide-binding leucine-rich-repeat receptors (NLRs) play pivotal roles in ETI, which frequently leads to hypersensitive response (HR)-mediated cell death (Kourelis and van der Hoorn, 2018; Adachi et al. 2019a). NLR genes are highly expanded in most plant genomes (~150 in *Arabidopsis thaliana* and ~500 in rice) and are among the most variable genes in plants, pointing

to strong selection pressure from pathogens (Clark et al. 2007; Van de Weyer et al. 2019). NLR proteins are characterized by a conserved nucleotide-binding (NB) domain and a leucine-rich-repeat (LRR) domain. NLRs are divided into two major groups depending on the type of N-terminal domain: NLRs with the N-terminal coiled-coil (CC) domain are called CNLs (CC-NLRs), and those with the N-terminal Toll-like domain are called TNLs (TIR-NLRs). CNLs are widespread in the plant kingdom, whereas TNLs have been detected only in dicot plants (Pan et al. 2000). ADP/ATP exchange at the NB domain (Bernoux et al. 2016; Wang et al. 2019a) and oligomerization of NLRs (Ma et al. 2020; Martin et al. 2020; Sharif et al. 2019; Tenthorey et al. 2017; Wang et al. 2019b; Zhang et al. 2015) trigger ETI signaling. The Arabidopsis CNL ZAR1 forms a pentamer "resistosome" complex after binding to a host protein and a pathogen effector. This leads to the protrusion of the N-terminal alpha helix, which perturbs the plasma membrane to trigger the HR (Wang et al. 2019ab), a process potentially mediated by Ca^{2+} influx (Bi et al. 2021). The Arabidopsis TNL RPP1 forms a tetramer "resistosome" upon binding to its cognate effector ATR1, followed by an increase in its NAD^{+}ase activity (Ma et al. 2020), which triggers cell death (Wan et al. 2019; Horsefield et al. 2019). These recent breakthroughs have started to reveal the biochemical links between NLR molecular structure and HR induction (Adachi et al. 2019b).

NLRs can function as singletons, in pairs or in networks (Adachi et al. 2019a). Arabidopsis ZAR1 (Wang et al. 2019a), RPP1 (Krasileva et al. 2010), and many other NLRs function as singleton NLRs and recognize AVRs directly or indirectly. Two NLR proteins encoded by genetically linked genes function together as paired NLRs. One of the paired NLRs frequently has a non-canonical domain called the integrated domain (ID), which shares amino acid sequence similarity with domains in the host protein (Kroj et al. 2016; Sarris et al. 2016). Examples of paired NLRs include RPS4/RRS1 in Arabidopsis (Williams et al. 2014; Le Roux et al. 2015) and RGA4/RGA5 (Okuyama et al. 2011; Cesari et al. 2013), Pik1/Pik2 (Ashikawa et al. 2008; Maqbool et al. 2015), and Pii1/Pii2 in rice (Takagi et al. 2013; Fujisaki et al. 2017). Multiple NLRs encoded by unlinked genes may function together; these network NLRs (Wu et al. 2017) include NRC2, NRC3, and NRC4 in Solanaceae plants. When NLRs function as a pair or network, one NLR is involved in the recognition of the AVR effector (the "sensor NLR"), whereas the other plays a role in signaling (the "helper NLR"). Recent studies have expanded our understanding of the genetic architecture of NLR pairs and networks; however, how they function together and how they evolved remain elusive.

A phylogenetic study of 4,184 NLRs in the genomes of seven Poaceae species including rice, wheat, and other grass species grouped the NLRs into 24 major clades, including three clades (MIC1, MIC2, and MIC3) containing the majority of NLRs with IDs (Bailey et al. 2018). NLRs in clade MIC1 (including RGA5) are characterized by a wide variety of IDs integrated into similar positions in the NLR after the LRR domain. Bailey et al. (2018) reported a conserved 43-amino acid motif between LRR and ID named the CID (conservation and association with IDs) motif. The authors proposed a possible evolutionary mechanism of ID generation and ID shuffling mediated by the CID motif. However, this study examined only a few genomes from each species, with a focus on inter-species diversity within the Poaceae, and provided limited information about the recent diversification of NLRs within a single genus.

In addition, experimental validation of the hypotheses underpinning the functional diversity of NLRs with IDs remains limited.

Rice blast disease caused by blast fungus (*Magnaporthe oryzae* [syn. *Pyricularia oryzae*]) is a major disease of rice that threatens world food security (Pennisi 2010). The best way to control this disease is to deploy cultivars with resistance (*R*) genes. To date, 40 *R*-genes against blast have been reported and 25 have been cloned, most of which encode NLRs (Liu et al. 2010; Wu et al. 2015). However, the molecular interactions between rice NLRs and blast AVRs are understood for only a small number of cases, including Pita and AVR-Pita (Orbach et al. 2000; Jia et al. 2000), Pia and AVR-Pia/AVR1-CO39 (Cesari et al. 2013), Pik and AVR-Pik (Kanzaki et al. 2012; Maqbool et al. 2015), Pii and AVR-Pii (Takagi et al. 2013; Fujisaki et al. 2015; 2017), and Pizt and AVR-Pizt (Park et al. 2016, Wang et al. 2016).

The rice Pia pair, consisting of the NLR helper RGA4 and the NLR sensor RGA5, is one of the most well-characterized NLR pairs (Okuyama et al. 2011; Cesari et al. 2013). Overexpression of *RGA4* in *Nicotiana benthamiana* and rice protoplasts caused HR-like cell death in the respective plant species, suggesting that the helper RGA4 is responsible for HR signaling. RGA4-mediated cell death was suppressed by co-expression of RGA5, indicating that RGA5 negatively regulates RGA4-mediated defense signaling. Finally, co-expression of RGA5, RGA4, and AVR-Pia triggered cell death via the direct binding of AVR-Pia to HMA ID of the sensor RGA5 (Cesari et al. 2013). This study, together with a study of the Arabidopsis RPS4/RRS1 pair (Williams et al. 2014; Le Roux et al. 2015), supported a negative regulation model for paired NLRs in which helper NLRs function in HR signaling and sensor NLRs in the suppression of helper NLR activity. Upon binding or modification of the sensor-NLR IDs by pathogen AVRs, this suppression is released, allowing HR signaling to proceed.

Here, we describe the rice NLR pair Pias, which recognize the *M. oryzae* effector AVR-Pias. *Pias* encodes the helper Pias-1 and the sensor Pias-2 and is allelic to the previously characterized *Pia* gene, encoding the NLR pair helper RGA4 and sensor RGA5. Pias-2 carries a C-terminal DUF761 domain at a similar position to the heavy metal–associated (HMA) domain of RGA5. We show that the Pias/Pia helper NLR lineage is evolutionarily and functionally conserved, while its sensor NLR lineage shows highly dynamic evolution with various host domains integrated into similar positions, possibly allowing it to detect a wide variety of pathogen molecules.

Results

Isolation of rice *Pias* NLR genes and the matching gene *AVR-Pias* from *Magnaporthe oryzae*

As part of a large genetic screen to identify novel rice blast resistance genes, we crossed *japonica*-type rice cultivar Hitomebore to 20 different rice cultivars representing the worldwide genetic diversity of *Oryza sativa*, resulting in the generation of recombinant inbred lines (RILs) of the F_7-F_9 generations. We inoculated the parental rice lines with a panel of *M. oryzae* isolates and recorded their resistance or susceptibility to each isolate. Since Hitomebore was susceptible to *M. oryzae* isolate 2012-1 but *indica-*

type accession WRC17 (cultivar Keiboba; Kojima et al. (2005)) was resistant to this isolate, we set out to isolate the resistance genes in WRC17 rice against the 2012-1 pathogen. The 58 RILs derived from a cross between Hitomebore and WRC17 segregated into 52 resistant and 6 susceptible lines (**Fig. 1A**), indicating that WRC17 likely contains more than one locus conferring resistance against 2012-1.

To identify the resistance genes, we performed whole-genome sequencing and RNA-seq of WRC17 and conducted an association study using a bioinformatics pipeline we named "RaIDeN" (https://github.com/YuSugihara/RaIDeN) using the RILs segregating for these phenotypes (**Fig. 1; Suppl. Fig. 1**). We sequenced the genomes WRC17, Hitomebore, and six RILs showing susceptibility to the isolate 2012-1 on an Illumina DNA sequencer (**Suppl. Table 1**) and subjected the short reads of WRC17 to *de novo* assembly with *DISCOVAR* (https://www.broadinstitute.org/software/discovar/blog/), resulting in the WRC17 reference genome sequence (**Suppl. Table 2**). We also performed RNA-seq of WRC17 leaves that had been inoculated with *M. oryzae* (2012-1). The RNA-seq reads were mapped to the WRC17 reference genome, revealing 22,561 genes expressed from the WRC17 genome. The short reads obtained from Hitomebore and six susceptible (S-) RILs were aligned to the genome sequences of the 22,561 expressed genes. We reasoned that Hitomebore and the six S-RILs share the identical DNA sequences in the candidate genes responsible for their resistance that are different from the sequences in resistant WRC17 (**Suppl. Fig. 1**). Two types of DNA polymorphisms were considered: (1) presence/absence of the genes and (2) SNPs in the genes. We identified 14 genes that were present in WRC17 but absent from Hitomebore and the six S-RILs, in addition to 839 genes with shared SNPs among Hitomebore and the S-RILs that were different from the sequences of WRC17.

From this group of 853 genes (**Suppl. Dataset 1**), we selected Resistance Gene Analogs (RGAs) using "RGAugury" (Li et al. 2016), which predicts genes encoding putative NLRs, Receptor Like Kinases (RLKs), and Receptor Like Proteins (RLPs). This analysis identified 38 RGAs as the candidate genes (**Suppl. Table 3**). Since most rice resistance genes against *M. oryzae* reported to date are *NLR*s, we focused on 18 *NLR* (11 *CNL* and 7 *NL*) genes as the candidate resistance genes of WRC17 against *M. oryzae* isolate 2012-1. Among the 18 NLR genes, only one (*NL-04*) showed presence/absence polymorphisms and the rest contained SNPs associated with the phenotypes. We developed DNA markers in the candidate *NLR* genes and studied their association with phenotypes using 58 RILs segregating for resistance and susceptibility to the isolate 2012-1 (**Suppl. Fig. 2A**). This analysis showed that 16 *NLR*s were tightly linked to each other and that the two other *NLR*s were linked to each other, suggesting that the two loci (designated *Pi-W17-1* and *Pi-W17-2*) are involved in the resistance of WRC17 against 2012-1 (**Fig. 1B**). Genotyping of the 52 RILs using the markers located in the two loci, *Pi-W17-1* (*CNL-04*) and *Pi-W17-2* (*NL-07* and *W17.11718*), suggested that the lines became resistant when either of the two loci contained the WRC17-type allele (**Fig. 1A; Suppl. Fig. 2A**).

The genomic position of *Pi-W17-1* corresponds to that of the previously reported *Pia* locus (Okuyama et al. 2011), and the position of *Pi-W17-2* corresponds to that of the *Pita* (=*NL-07*) and *Ptr* (=*W17.11718*) loci (Zhao et al. 2018; Orbach et al. 2000; Meng et al. 2020). Among the progeny showing resistance against 2012-1, we selected 19 RILs containing *PiW17-1* but not *PiW17-2*. These lines shared

10 candidate NLR genes within the *PiW17-1* region. (**Fig. 1B; Suppl. Fig. 2B**). We performed RNAi-mediated gene silencing of eight of these genes (encoding proteins over 900 amino acids long) (**Suppl. Fig. 2C**) using the RIL HW-RIL7, which contains *Pi-W17-1* but lacks *Pi-W17-2*. When *CNL-04* or *CNL-05* was silenced, its resistance against 2012-1 became compromised (**Suppl. Fig. 2CD**). We confirmed this result by generating *CNL-04* and *CNL-05* knockout mutant lines by CRISPR/Cas9-mediated genome editing (**Fig. 1C; Suppl. Fig. 3**). These data suggest that *Pi-W17-1*-mediated resistance requires both the neighboring NLRs *CNL-04* and *CNL-05*. Indeed, the position of *CNL-04* corresponds to that of *RGA4*, whereas the position of *CNL-05* corresponds to that of *RGA5* of *Pia* (**Fig. 1D**; Okuyama et al. 2011). Interestingly, *CNL-04* is similar to *RGA4* in terms of both structure and DNA sequence (96.6% DNA sequence identity), whereas *CNL-05* has a distinct structure and a DNA sequence that diverged from *RGA5* (59.8% DNA sequence identity) (**Suppl. Fig. 4**). *Pia RGA5* encodes a protein with a heavy metal–associated (HMA) domain in its C terminus (Okuyama *et al.* 2011), whereas *CNL-05* encodes a protein with a 19-amino acid motif corresponding to domain of unknown function 761 (DUF761) near its C terminus. The physical distance between *CNL-04* and *CNL-05* is 8.7 kb, which is longer than that between *RGA4* and *RGA5* (3.7 kb) (**Fig. 1D**). In view of the substantial differences between *CNL-05* and *RGA5*, we decided to name this WRC17 allele *Pias*, *CNL-04* as *Pias-1*, and *CNL-05* as *Pias-2* (**Suppl. Fig. 5**).

To isolate the *AVR-Pias* avirulence gene cognate of rice NLR *Pias*, we performed an association study of expressed genes encoding candidate effector proteins (See **Suppl. Fig. 6** and **Suppl. Table 4, 5** for details). This analysis identified three genes (*G9141*, *G9435*, and *G9532*) as candidates of *AVR-Pias*. We selected *M. oryzae* isolate Ao92-06-2, which is compatible with HW-RIL7, transformed it with each of the candidate genes, and tested their interactions with HW-RIL7 (**Suppl. Fig. 7**). Transformation with one of the candidate genes, *G9532*, rendered Ao92-06-2 incompatible with HW-RIL7 (**Suppl. Fig. 7A**), suggesting that *G9532* is *AVR-Pias*. To validate this result, we generated a knockout mutant of *G9532* in the 2012-1 background, which became compatible with HW-RIL7 (**Fig. 1E; Suppl. Fig. 8**). These results indicate that *G9532* is *AVR-Pias*, which is recognized by *Pias*. Both *Pias-1* and *Pias-2* are required for the recognition of *AVR-Pias*, as knockout of either *NLR* gene abrogated resistance (**Suppl. Fig. 7B**). AVR-Pias is a 91-amino acid protein with a secretion signal peptide (**Fig. 1F**). This protein contains a 12-amino acid Toxin18-like motif in its C terminus.

Various host domains are integrated into Pias/Pia sensor NLRs

Pias and *Pia* are allelic to each other, and both are composed of a pair of NLRs. The helper NLRs (*Pias-1* and *RGA4*) are conserved, whereas the sensor NLRs (*Pias-2* and *RGA5*) are divergent, with different integrated domains. To explore the diversity and evolution of *Pias/Pia* NLRs in the entire *Oryza* genus, we obtained the genomic DNA sequences of the *Pias/Pia* locus from 171 accessions representing 11 *Oryza* species as well as four non-*Oryza* species of Poaceae, including *Setaria italica* (foxtail millet), *Panicum hallii* (Hall's panicgrass), *Hordeum vulgare* (barley), and *Aegilops tauschii* (Tausch's goatgrass) (**Suppl. Table 6**). Remarkably, Pias-2/RGA5 sensor proteins contain a wide variety of IDs, with up to nine different domains including HMA, DUF761, DUF677, Zinc_ribbon_12, Pkc_MAPKK (PKc_M), Pkc_like,

and WRKY inserted into the identical position after the LRR domain (**Fig. 2A**). Between the ID and LRR domains, we identified a conserved 145- to 146-amino acid motif named the LRR-ID intervening (LII) region (**Fig. 2B**), which partially overlaps but is different from the previously reported CID motif (Bailey et al. 2018; **Suppl. Fig. 9**).

We studied the frequency of different IDs in the 11 *Oryza* species used to reconstruct a phylogenetic tree based on whole-genome sequences (**Fig. 2C; Suppl. Table 6, 7**). The IDs of the cultivated rice *O. sativa* (44 samples) are shared by the HMA and DUF761 domains with similar frequencies. *O. rufipogon* (13 samples), the wild progenitor of *O. sativa*, contains DUF677 and PKc_M in addition to HMA and DUF761. *Oryza* species belonging to the A-genome group (*O. sativa*, *O. rufipogon*, *O. barthii*, *O. glumaepatula*, and *O. meridionalis*) contain a higher proportion of the DUF761 ID, suggesting its importance in their defense. However, outside the A-genome species, the DUF761 ID is absent. Instead, PKc_M (*O. brachyantha* and *O. punctata*), DUF677 (*O. officinalis*), and HMA (*O. australiensis*) are dominant, indicating that the integration of the DUF761 domain into Pias-2/RGA5 likely occurred in the immediate ancestor of the A-genome species that diverged from other *Oryza* lineages over 2.3 Mya (TimeTree; http://www.timetree.org/).

A comparison of the genome sequences around *Pias/Pia* sensor-NLR genes between *O. punctata* and *O. sativa* cultivar Nipponbare revealed an interesting conserved region (**Fig. 2DE**). Both of their coding regions contain conserved LII sequences, whereas the IDs downstream of the LII are different: PKc_M for *O. punctata* W1582 and DUF761 for *O. sativa* Nipponbare. However, in the 30.8-kb region downstream of the *O. sativa* Nipponbare *Pias-2* gene, two regions (block **a** and **b**) share high DNA sequence similarity with the LII and ID sequences of *O. punctata* W1582 Pias-2 homolog (**Fig. 2DE; Suppl. Fig. 10**): block **a** corresponds to region LII to exon-6, and block **b** corresponds to exon-7 to -10 of *O. punctata* W1582 *Pias-2* homolog. Perhaps this conserved sequence downstream of the *O. sativa* Nipponbare *Pias* sensor is a footprint of the replacement of the ID from PKc_M to DUF761 via homologous recombination at the LII region (**Fig. 2F**). A survey of 51 A-genome *Oryza* accessions with the DUF761 ID revealed that the *O. punctata* W1582 LII-ID-like sequence is widely conserved in *O. sativa*, *O. rufipogon*, *O. meridionalis*, and an accession of *O. barthii* (Accession W1702), but not in the majority of *O. barthii* and *O. glumaepatula* accessions (**Suppl. Table 8; Suppl. Fig. 11**). These results indicate that this possible recombination occurred only once and that the footprint was probably lost in the latter two species. We did not detect similar footprints in other sensor *Pias/Pia* NLRs with non-DUF761 IDs.

Contrasting patterns of evolution of *Pias/Pia* sensors and helpers

To explore the evolutionary patterns of the *Pias/Pia* NLR locus, we reconstructed phylogenetic trees of *Pias*/*Pia* NLR pairs separately for the helper NLR (Pias-1 and RGA4) and the sensor NLR (Pias-2 and RGA5) using the amino acid sequences of the helper NLRs (full-length protein) and sensor NLRs (full-length protein except the ID domain) of 22 *Oryza* accessions and four accessions from other Poaceae genera. Note that Pias/Pia helper and sensor NLRs belong to distantly related NLR clades (Bailey et al.

2018). Overall, the helper NLR tree (**Fig. 3A**) is consistent with the species tree (**Fig. 2C**) with a few exceptions. Species in the AA genome group (*O. sativa* subsp. *japonica*, *O. sativa* subsp. *indica*, *O. rufipogon*, *O. barthii*, *O. glumaepatula*, and *O. meridionalis*) clustered together. Only the placement of *O. australiensis* (accession W0008) is not congruent between the Pias-1/RGA4 helper NLR tree and the species tree. On the other hand, the sensor-NLR tree is quite distinct from the helper NLR tree and is not congruent to the species tree. Remarkably, the sensor NLRs show a higher level of divergence than the helper NLRs (**Fig. 3A**). The sensor phylogeny includes two major clades (C1and C2) separated by deep branches. RGA5 of *O. sativa* cv. Sasanishiki (Okuyama et al, 2011) belongs to the C1 clade, whereas Pias-2 of *O. sativa* WRC17 belongs to the C2 clade. Four species of the AA genome group (*O. sativa* subsp. *japonica*, *O. sativa* subsp. *indica*, *O. rufipogon*, *O. barthii*, and *O. meridionalis*) and *O. punctata* (BB genome) have sensor-NLR alleles both from the C1 and C2 clades, whereas *O. australiensis* (EE genome), *O. granulata* (GG genome), and *O. brachyantha* (FF genome) have C1 alleles and *O. glumaepatula* (AA genome), and *O. officinalis* (CC genome) have C2 alleles. Therefore, phylogenetic analysis of the sensor NLR of the *Pias/Pia* locus pointed to trans-species polymorphism, which is reminiscent of the major histocompatibility (MHC) gene polymorphism in vertebrates (Edwards and Hedrick, 1998).

We then tested the signatures of positive selection in the *Pias-1/RGA4* helper and *Pias-2/RGA5* sensor-NLR genes. We calculated the ratio of nonsynonymous to synonymous base substitutions (ω: Yang and Nielsen, 2000) for the pair of NLR gene sequences obtained from different *Oryza* species, separately for the helper and sensor NLRs (**Fig 3B**). Note that the ID sequences were removed from the sensor-NLR sequences before analysis. Remarkably, the ω values were < 0.4 between the pair of helper NLRs, which is suggestive of purifying selection, whereas larger ω values were obtained between the pair of sensor NLRs, indicating positive selection ($\omega > 1.0$) in some combinations. These results suggest that the tightly linked helper and sensor NLRs of the *Pias/Pia* locus in the genus *Oryza* are undergoing contrasting modes of evolution.

Pias-1 helper functions with RGA5 sensor to recognize AVR-Pia

Pia comprises the helper NLR RGA4 and the sensor-NLR RGA5. The expression of RGA4 triggers HR cell death in rice as well as *N. benthamiana*, and RGA5 suppresses RGA4-mediated cell death (Césari et al. 2014). Upon the binding of AVR-Pia to HMA ID of RGA5, this suppression is released, and HR-like cell death is triggered (Césari et al. 2014). To address the functional conservation of helper/sensor NLRs of the *Pias/Pia* locus, we examined the functions of helper NLRs by transiently overexpressing five Pias/Pia helper NLRs in *N. benthamiana* using agroinfiltration (**Suppl. Fig. 12**). Only RGA4 supported strong cell death, and other alleles including Pias-1 caused weaker cell death, indicating that the role of RGA4 as a strong HR inducer is not typical among the tested helper NLRs (**Suppl. Fig. 12**). Next, we tested the effects of co-expressing Pias-1 and Pias-2, which surprisingly resulted in stronger cell death than that caused by Pias-1 expression alone (**Suppl. Fig. 13**). These results suggest that the function of Pias-1/Pias-2 is different from that of the NLR pair RGA4/RGA5 NLR.

We then examined the effects of exchanging various helper and sensor Pias/Pia NLRs by performing a transient expression assay in *N. benthamiana* leaves (**Fig. 4A, Suppl. Fig. 14**). Co-expression of the sensor RGA5 with the two heterologous helpers (Pias-1 and RGA4-Ogr from *O. granulata*) suppressed HR-like cell death, indicating that RGA5 can suppress the cell death induced by helper NLRs other than RGA4 (**Fig. 4A, Suppl. Fig. 14**). Furthermore, this RGA5-mediated suppression of the Pias-1-triggered HR was released via co-expression of AVR-Pia, which resulted in cell death (**Fig. 4A; Suppl. Fig. 14**). A similar result was obtained using RGA4-Ogr helper (**Fig. 4A; Suppl. Fig. 14**). These results indicate that the sensor RGA5 properly functions with helpers other than RGA4 to recognize AVR-Pia and mount the HR in *N. benthamiana*.

Finally, we generated transgenic rice plants (HW-RIL7:35S-RGA5) expressing the *RGA5* transgene driven by the cauliflower mosaic virus 35S (CaMV35S) promoter in the *Pias* (*Pias-1* and *Pias-2*) background. We challenged these lines with two *M. oryzae* isolates: Ao92-62-2 harboring the *AVR-Pia* transgene and Ao92-62-2 harboring the *AVR-Pias* transgene (**Fig. 4B; Suppl. Fig. 15**). Remarkably, HW-RIL7:35S-RGA5 exhibited resistance against *M. oryzae* containing *AVR-Pia* (**Fig. 4B**). Although the HW-RIL7:35S-RGA5 lines consistently showed resistance against *M. oryzae* containing *AVR-Pia*, their resistance against *M. oryzae* containing *AVR-Pias* varied. We crossed an *rga4* mutant in the Sasanishiki background (Sas1493; Okuyama et al. 2011) with HW-RIL7 and obtained F1-B plants. These plants, harboring intact *RGA5* as well as *Pias-1* and *Pias-2*, now recognized and triggered resistance against *M. oryzae* containing *AVR-Pia*, but not *AVR-Pias* (**Fig. 4B**). Similar results were obtained for F1-A plants generated by crossing Sasanishiki (WT) with HW-RIL7 (**Fig. 4B**). These results suggest that *Pia* function is dominant over *Pias* function in terms of the recognition of *AVR-Pia* and *AVR-Pias*.

These results suggest that Pias-1 helper functions together with RGA5 in rice to recognize and mount resistance against *M. oryzae* containing *AVR-Pia*, suggesting that the helper function has been conserved over the long history of Pias/Pia evolution.

Discussion

Our study investigated the evolution of a pair of genetically linked NLRs in the genus *Oryza* and provided experimental evidence that the two paired NLRs have evolved in dramatically contrasting fashions. This study points to the evolution of a modular architecture of paired NLRs. Division of roles between a conserved helper NLR for signaling and a divergent sensor NLR with a cassette-like receptor domain for pathogen sensing may have conferred plants with the ability to efficiently fend off rapidly evolving microbe pathogens.

We identified and functionally characterized the rice R-gene *Pias*. This gene encodes the paired NLRs Pias-1 helper and Pias-2 sensor, which recognizes the *M. oryzae* effector AVR-Pias. *Pias* is allelic to the well-studied *R*-gene *Pia*, encoding the NLRs RGA4 helper and RGA5 sensor (Okuyama et al. 2011), which recognizes the effectors AVR-Pia (Yoshida et al. 2009) and AVR1-CO39 (Cesari et al. 2013). The allelic sensor-NLRs Pias-2 and RGA5 carry different domains at their C termini. In RGA5, the integrated

domain HMA directly binds to and recognizes two *M. oryzae* effectors, AVR-Pia and AVR1-CO39 (Cesari et al. 2013). We have not yet detected direct binding between DUF761 ID of Pias-2 and AVR-Pias despite several attempts. Perhaps the recognition of AVR-Pias by Pias-2 requires other host components; indeed, the recognition of AVR-Pii by Pii-2 requires the rice protein OsExo70-F3, which binds to both AVR-Pii and Pii-2 (Fujisaki et al. 2015, 2017).

What is the origin of the integrated DUF761 domain of Pias-2? The rice sensor-NLRs RGA5 and Pik-1 contain HMA domains as IDs. The ID HMA shares high amino acid sequence similarity with rice small heavy metal–associated domain proteins (sHMAs: Oikawa et al. 2020; Białas et al. 2021). We revealed that the *M. oryzae* effector AVR-Pik binds to and stabilizes sHMA proteins, likely to promote pathogen infection (Oikawa et al. 2020; Maidment et al. 2021). To identify the proteins that provide the DUF761 domain to Pias-2 ID, we performed BLAST searches against the rice protein database using the short Pias-2 DUF761 sequence (19 amino acids) as a query (**Suppl. Fig.16**). This identified 15 proteins with a similarity threshold E < 10. Most of the 15 proteins contained DUF761 at their C termini. Only a few functional studies of DUF761-containing proteins have been performed. A study in cotton (*Gossypium hirsuta*) showed that GhCFE1A, containing DUF761 and DUF4408 domains, binds to actin proteins and localized to the endoplasmic reticulum (ER) upon overexpression in *N. benthamiana* (Lv et al. 2015). Knockdown of *GhCFE1A* did not cause any phenotypic changes, while its overexpression led to delayed cotton fiber cell elongation. These results suggest that GhCFE1A is a linker protein that mediates the formation of the ER network and actin cytoskeleton. Arabidopsis *A70*, encoding a DUF761-containing protein, is specifically induced in the incompatible interaction with *Pseudomonas syringae*, but not in the compatible interaction (Truman et al. 2007). Knockout of the Arabidopsis DUF761-containing protein gene *DUF761-1* did not have any phenotypic effects, whereas overexpressing *DUF761-1* altered plant morphology and resulted in a constitutive defense response, leading to enhanced resistance against *P. syringae* (Zhang et al. 2019). These findings suggest that DUF761-containing proteins function in defense, presumably mediated by the actin-ER network. Future studies should investigate how AVR-Pias is recognized by Pias and how AVR-Pias interferes with host cellular processes.

Divergent sensor NLRs in the Pias/Pia lineage

The Pias-2/RGA5 sensor-NLR lineage is extremely divergent among *Oryza* species, with up to six different ID motifs integrated at their C termini (**Fig. 2**). These hugely divergent IDs may mediate the detection of a diversity of effectors from the blast fungus and possibly other pathogens. We hypothesize that the diversity of the Pias-2/RGA5 lineage has been maintained by natural selection to maintain various IDs that recognize the invasion of pathogens by directly binding to effectors or guarding host factors that are modified by effectors.

Within *Oryza* species, there are two major clades, C1 and C2. Notably, alleles from both clades are maintained within the species *O. sativa*, *O. rufipogon*, *O. barthii*, *O. meridionalis*, and *O. punctata*. The observed trans-species allelic divergence and their roles in detecting pathogen molecules are similar to those of the major host incompatibility (MHC) locus of vertebrates (Edwards and Hedrick, 1998).

Individuals with higher heterozygosity at the MHC locus might have higher fitness (overdominance) due to their ability to bind to a larger number of pathogen peptides (Takahata and Nei, 1990; Piertney and Oliver, 2006). Perhaps in the ancestral outcrossing *Oryza* species, heterozygous plants with a larger repertoire of NLRs with different IDs had a selective advantage against a multitude of pathogens. It is also possible that frequency-dependent selection helped maintain this polymorphism. When the frequency of a pathogen effector in a population increases, the frequency of an allele for a cognate sensor NLR will increase, resulting in a reduction in the frequency of pathogen alleles in the population. In turn, the frequency of another effector gene will increase in the pathogen, and the frequency of the cognate sensor NLR will increase. According to this Red Queen model, the allele frequencies of both effector and sensor NLRs oscillate and may be maintained for a long time by balancing selection (Takahata and Nei, 1990; Woolhouse et al. 2002; Terauchi and Yoshida, 2010). In summary, the highly divergent evolution of Pias/Pia sensor NLRs with variable IDs seems driven by the fitness gain obtained by an enhanced recognition capability of pathogen effectors.

Genetic mechanism of ID switching in Pias/Pia NLRs

The Pias/Pia sensor NLRs contain various ID sequences at the identical position downstream of the LRR domain. This suggests the presence of a mechanism for "cassette"-like exchange of IDs between sensor NLRs. Between the LRR and ID, we identified a highly conserved stretch of 145 amino acid sequences named LII. Bailey et al. (2018) reported a 43-amino acid CID motif conserved in the region between the LRR and ID of the MIC1 NLR clade by studying the NLRs of seven Poaceae species. The LII motif identified encompasses the CID motif (**Suppl. Fig. 9**) but extends to the N-terminal direction by approximately 100 amino acids. Bailey hypothesized that the CID could serve as a recombination point of integration of genomic sequences. Despite the difference in the conserved motif, our data basically support Bailey (2018)'s hypothesis that this region serves as the recombination point for the integration of endogenous sequences matching various protein domains. In support of this idea, the downstream sequence of a Pias sensor NLR with DUF761 contains a DNA sequence similar to the LII and ID regions of the *O. punctata* Pias sensor homolog. Perhaps PKc_M ID of the original *O. punctata* Pias was replaced by the ID DUF761 and this switch caused the translocation of PKc_M to the region downstream of the sensor NLR gene (**Fig. 2F**). In view of the high sequence conservation between *O. punctata* PKc_M ID and the downstream sequence, it is also possible that the downstream sequence was functional until the recent past, the Pias sensor contained the dual IDs DUF761 and PKc_M in the same molecule, or Pias switched between these two IDs, possibly via alternative splicing. Future study should address the mechanism of LII-mediated recombination.

Function of *Pias/Pia* paired NLRs

Pia has been extensively studied (Césari et al. 2014) and serves as a paradigm for paired NLRs together with the paired NLRs RPS4/RRS1 (Williams et al. 2014). In these two cases, the helper NLR is regarded as a cell death inducer and the sensor NLR as a suppressor that maintains the complex in an inactive state when the pathogen is absent. Once the sensor has been modified by direct binding (RGA5) or modification (RRS1) of the ID by pathogen effectors, the suppression of the helper is released and HR-like cell death occurs. However, Pias NLRs do not function according to this model. In the *N. benthamiana* assay, Pias-1 functioned as a weak cell death inducer and Pias-2 did not suppress Pias-1-mediated cell death. Indeed, Pias-1 and Pias-2 together triggered stronger cell death than that caused by Pias-1 alone. A recent functional study of rice Pikp, another paired NLR, also showed that the helper NLR alone does not cause cell death in the *N. benthamiana* system and that the helper and sensor cooperate to trigger HR (Zdrzałek et al. 2020). Therefore, the hypothesized functional roles of the helper as a cell death inducer and the sensor as a cell death suppressor as well as a detector of effector molecules may not be as prevalent as assumed. The system of negative regulation of a cell death inducer by a suppressor encoded by genetically separate factors carries tremendous risks given that a loss-of-function mutation in the suppressor gene kills the carrier cells and incurs a genetic load. Therefore, such an extreme negative regulation system is unlikely to be maintained over a long period of evolution. We predict that cooperative NLRs in pairs or networks are more prevalent (Wu et al. 2017). It is also possible that the Pias/Pia paired NLR system is regulated by additional components in rice cells that are absent from *N. benthamiana*. Further studies are needed to understand the full regulatory network of Pias/Pia paired NLR-mediated resistance.

Helper NLRs in the Pias/Pia lineage are functionally conserved

Phylogenetic reconstruction of the Pias/Pia NLR locus revealed that the helper NLR Pias-1/RGA4 is conserved, whereas the sensor NLR Pias-2/RGA5 is highly divergent (**Fig. 3AB**). We validated these findings from molecular evolutionary analyses with *in planta* experiments. A functional study in *N. benthamiana* showed that Pias-1-mediated cell death was suppressed by RGA5 and that Pias-1 together with RGA5 function in the recognition of AVR-Pia leading to cell death (**Fig. 4A**). Similarly, an RGA4 homolog of *O. granulata* that is phylogenetically distant from *O. sativ*a also functions with RGA5 to recognize AVR-Pia (**Fig. 4A**). Moreover, a HW-RIL7 rice line harboring *Pias* as well as an *RGA5* transgene recognized AVR-Pia (**Fig. 4B**). These results suggest that the function of the Pias-1/RGA4 helper lineage is conserved, which is in line with its conserved amino acid sequences. It is possible that the separation of the roles of NLRs between the conserved helper and divergent sensor allowed for higher flexibility of pathogen recognition compared to singleton NLRs. This flexibility would allow the plant to cope with the rapid evolution of pathogens, which exhibit larger population sizes and shorter generation times than the host plants. The functional understanding of the modular structure of paired NLRs revealed here provides a basis for engineering NLRs to detect various effectors and to confer resistance to crops against pathogens.

Materials and Methods

Rice pathogenicity assays

Rice leaf blade punch inoculation was performed using the *M. oryzae* isolates. A conidial suspension (3 × 10^5 conidia mL^{-1}) was punch inoculated onto a rice leaf 1 month after seed sowing. The inoculated plants were placed in a dew chamber at 27°C for 24 h in the dark and transferred to a growth chamber with a 16-h-light/8-h-dark photoperiod. Disease lesions were scanned 10 days post-inoculation (dpi), and lesion size was measured using Image J software (Schneider et al. 2012).

RNA-seq of rice and barley leaves infected with *M. oryzae* 2012-1 isolate

Total RNA was extracted from rice and barley infected leaves using an SV Total RNA Isolation System (Promega, WI, USA). One microgram of total RNA was used to prepare each sequencing library with an RNA Sample Prep Kit v. 2 (Illumina, CA, USA). The two types of libraries, created from infected barley and rice leaves, were sequenced by Paired-End (PE) and Single-End (SE) sequencing using the NextSeq 500 platform.

DNA-seq for the RaIDeN pipeline

Genomic DNA was extracted from WRC17, Hitomebore, and S-RILs leaves using a NucleoSpin Plant II Kit (Macherey Nagel Co, Du□ren, Germany). Libraries for PE short reads were constructed using an Illumina TruSeq DNA LT Sample Prep Kit (Illumina, CA, USA). The PE library of WRC17 and other libraries were sequenced on the Illumina MiSeq and HiSeq 4000 platforms.

Generation of *PiW17-1* with knocked-down expression of candidate NLR genes in rice and quantitative RT-PCR

The eight types of gene knockdown (RNAi) constructs (pANDA-*PiW17-1* candidate NLRs) were generated by PCR amplification of a specific fragment of each *PiW17-1* candidate NLR gene from WRC17 WT cDNA. The sequences were cloned into the Gateway vector pENTR/D-TOPO (Invitrogen, CA, USA) and transferred into recombination sites of the pANDA vector (Miki and Shimamoto, 2004) using LR Clonase (Invitrogen, CA, USA). The resulting vectors with eight types of pANDA-PiW17-1 candidate NLR genes were introduced into *Agrobacterium tumefaciens* (strain EHA105) and used for *Agrobacterium*-mediated transformation of HW-RIL7 following the method of Okuyama et al. (2011). Total RNA was extracted from leaves using an SV Total RNA Isolation System (Promega, WI, USA) and used for quantitative RT-PCR (qRT-PCR). cDNA was synthesized from 500 ng total RNA using a Prime Script RT Reagent Kit (Takara Bio, Otsu, Japan). qRT-PCR was performed using a StepOne Real-time PCR Instrument (Applied Biosystems, CA, USA) with KAPA SYBR FAST PCR Master Mix (Kapa Biosystems, MA, USA). Melting curve analysis (from 60 to 95°C) was included at the end of the cycles to

ensure the consistency of the amplified products. The comparative Ct (ΔΔCt) method was used to calculate the expression of *CNL-04* (*CNL-05*) relative to the rice *ACTIN* gene (*LOC_Os03g50885*) as an internal control. The data presented are the average and standard deviations from three experimental replications. The primers used to generate the RNAi construct and for qRT-PCR are listed in **Suppl. Table 9**.

Generation of rice mutants of *CNL-04* and *CNL-05* by CRISPR/Cas9-mediated genome editing

Rice knockout mutants of *CNL-04* and *CNL-05* were generated using the CRISPR/Cas9 system developed by Mikami et al. (2015). Sense and antisense target sequences were designed using the web-based service CRISPR direct (http://crispr.dbcls.jp), annealed, and cloned into the pU6::ccdB::gRNA cloning vector following digestion with *Bbs*I as the target sequence. The target sequence with the *OsU6* promoter was cloned into the pZH::gYSA::MMCas9 vector following digestion with *Asc*I and *Pac*I. The resulting vectors (pZH::gYSA::MMCas9-*CNL-04* and -*CNL-05*) were introduced into *Agrobacterium tumefaciens* (strain EHA105) and used for *Agrobacterium*-mediated transformation of HW-RIL7 following the method of Okuyama et al. (2011). The resulting regenerated T0 plants were sequenced, and the mutation type was confirmed using primers listed in **Suppl. Table 9**.

Genetic complementation of the candidate AVR-Pias

Three candidate gene constructs (pCB1531-*AVR-Pias* candidate) were generated by PCR amplification of the coding sequences of the *AVR-Pias* candidate genes from cDNA prepared from *M. oryzae* 2012-1-infected barley leaves. The sequences were digested with *Xba*I and *BamH*I and cloned into pCB1531-pex22p-EGFP (Yoshida et al. 2009) that had been linearized by digestion with *Xba*I and *Bam*HI. The resulting vectors were used to transform Ao92-06-2 (lacking *AVR-Pias*) following the method of Sweigard et al. (1997). *M. oryzae* isolate 2012-1 mutated in *G9532* was generated using the CRISPR/Cas9 system developed by Arazoe et al. (2015). Sense and antisense target sequences were designed using the web-based service CRISPR direct (http://crispr.dbcls.jp), annealed, and cloned into the pCRISPR/Cas-U6-1 cloning vector following the method of Arazoe et al. (2015). To generate the targeting vector TV-*G9532*, the 5□ flanking region of *G9532* was amplified and cloned into pCB1636 (Sweigard et al. 1997) containing a hygromycin resistance gene that had been linearized by inverse PCR using primers pCB1636iv2fwd and pCB1636iv2rev as described by Shimizu et al. (2019) by In-fusion cloning (Clontech, Madison, WI, USA). Subsequently, the 3′ flanking region was amplified and cloned into a plasmid containing the 5′ flanking region that had been linearized by inverse PCR using primers pCB1636iv1fwd and pCB1636iv1rev as described by Shimizu et al. (2019) by In-fusion cloning (Clontech, Madison, WI, USA). The resulting vectors were used to transform 2012-1 isolate (containing *AVR-Pias*) following the method of Sweigard et al. (1997). The primers used for construct generation are listed in **Suppl. Table 9**.

Genome sequences used for the study

Genome sequences of 171 accessions of Poaceae plants were used, including 167 *Oryza* accessions as well as one accession each from *Aegilops tauschii*, *Hordeum vulgare*, *Panicum hallii*, and *Setaria italica*. These

also included sequences of *O. sativa* 'WRC17' (this study) and *O. sativa* 'Sasanishiki' (Okuyama et al. 2011). Genome sequences of 66 accessions were obtained from public databases: 52 accessions of *O. sativa* and *O. rufipogon* (Zhao et al. 2018), *O. sativa* subsp. *indica* 'ShuHui498' (Du et al. 2017), *O. sativa* subsp. *japonica* 'Nipponbare' (IRGSP. 2004), *O. rufipogon* 'W1943' (National Center for Gene Research), 5 accessions from *O. barthii*, *O. glumaepatula*, *O. meridionalis*, *O. punctata*, and *O. brachyantha* (The Oryza Map Alignment Project (OMAP)), *O. officinalis* 'W0002' (National Institute of Genetics, Japan), *O. granulata* 'W0067B' (Wu et al. 2018), *Ae. tauschii* 'AL8/78' (Zimin et al. 2017), *H. vulgare* 'Morex' (Mascher et al. 2017), *P. hallii* 'FIL2' (DOE Joint Genome Institute), and *S. italica* 'Yugu18' (Bennetzen et al. 2012) (see **Suppl. Table 6** for details). For 101 accessions of the wild *Oryza* species (*O. barthii*, *O. glumaepatula*, *O. meridionalis*, *O. punctata*, *O. officinalis, O. brachyantha*, and *O. granulate*), NGS reads (fastq format) of whole-genome sequences were retrieved from the NGS NCBI database (**Suppl. Table 6**) and used for *de novo* assembly by MaSuRCA (Zimin et al. 2013). For the two wild *Oryza* accessions, *O. glumaepatula* W2184 and *O. punctata* W1582, DNA sequencing was performed using Oxford Nanopore Technology (ONT) using genomic DNA extracted from their leaves. Sequencing was performed using the PromethION system with a FLO-PRO002 flow cell (ONT). Base calling of ONT reads was performed on FAST5 files using Guppy (ONT). Subsequently, low-quality reads were filtered out and *de novo* assembly was performed using NECAT (https://github.com/xiaochuanle/NECAT/) software. To further improve the accuracy of the assembly, Racon software (https://github.com/lbcb-sci/racon) was applied twice, and Medaka (https://github.com/nanoporetech/medaka) was used to correct mis-assembly. One round of consensus correction was performed using BWA (Li and Durbin, 2010) and HyPo (https://github.com/kensung-lab/hypo) on Illumina short reads for the accessions.

Gene annotation and detection of Pias/Pia orthologs

We used both publicly available RNA-seq and newly generated RNA-seq data in addition to the genome sequences to annotate the Pias/Pia homologs in *O. rufipogon* 'W1943', *O. barthii* 'IRGC105608', *O. glumaepatula* 'GEN1233_2', *O. glumaepatula* 'W2184', *O. meridionalis* 'W2112', *O. punctata* 'IRGC105690', *O. punctata* 'W1582', *O. officinalis* 'W0002', *O. australiensis* 'W0008', *O. brachyantha* 'IRGC101232', *O. brachyantha* 'W0655', and *O. granulata* 'W0067B'. The genome-guided transcriptome assembly was performed with StringTie (Pertea et al, 2015) by parsing the alignments produced by Hisat2 (Kim et al, 2015). The generated transcripts were processed by TransDecoder (https://github.com/TransDecoder/) to infer potential open reading frames (ORFs). Using the *de novo* assembly and the predicted ORFs, Pias/Pia homologs were predicted by Exonerate (http://www.ebi.ac.uk/~guy/exonerate) with Pias-1/RGA4 and Pias-2/RGA5 amino acid sequences as queries (**Suppl. Dataset 2**).

Prediction of ID sequences in the Pias/Pia sensor NLRs of the *Oryza* genus

To explore the diversity of ID in *Pias/Pia* sensor NLRs of the entire *Oryza* genus, the ID sequences of *Oryza* accessions were determined by Exonerate (http://www.ebi.ac.uk/~guy/exonerate) with 14 annotated sequences as queries in **Suppl. Dataset 2**.

Amino acid sequences and accession numbers of Pias/Pia orthologs

Amino acid sequences of Pias/Pia homologs used in this study were retrieved from the Eukaryotic Genome Annotation of the NCBI database (Du et al. 2017; Mascher et al. 2017); accession numbers for Pias-1/RGA4 homologs and Pias-2/RGA5 homologs are as follows: XP_015617251.1 and XP_015617810.1 for *O. sativa* subsp. *japonica* 'Nipponbare', OsR498G1119642600.01 and OsR498G1119642700.01 for *O. sativa* subsp. *indica* 'ShuHui498', XP_020148260.1 and XP_020148256.1 for *Aegilops tauschii* 'AL8/78', HORVU.MOREX.r2.4HG0288000.1, HORVU.MOREX.r2.4HG0288010.1 for *Hordeum vulgare* 'Morex', XP_025826635.1, XP_025827327.1 for *Panicum hallii* 'FIL2', and XP_004979045.1, XP_004979046.2 for *Setaria italica* 'Yugu18'.

Phylogenetic analysis of Pias-1/RGA4 and Pias-2/RGA5 orthologs

The protein sequences of the Pias-1/RGA4 and Pias-2/RGA5 orthologs were aligned using webPRANK (https://www.ebi.ac.uk/goldman-srv/webprank/) (Löytynoja et al. 2010). We used the whole amino acid sequences of the Pias-1/RGA4 orthologs but only the partial amino acid sequences (CC-NB-LRR-LII domains) of the Pias-2/RGA5 orthologs due to the very low sequence similarity after the LII domains. A maximum likelihood tree was constructed with IQ-TREE v2.0.3 (Minh et al. 2020) using 1,000 ultrafast bootstrap replicates (Hoang et al., 2018). The models were automatically selected by ModelFinder (Kalyaanamoorthy et al. 2017) in IQ-TREE (Minh et al. 2020). ModelFinder (Kalyaanamoorthy et al. 2017) selected "JTT + G4" for Pias-1/RGA4 orthologs and "JTT + R2" for Pias-2/RGA5 orthologs. Finally, the phylogenetic trees were drawn with FigTree v1.2.3 (http://tree.bio.ed.ac.uk/software/figtree/).

Analysis of selection pressure

First, we conducted codon-based alignment of the coding sequences of *Pias-1/RGA4* and *Pias-2/RGA5* orthologs with webPRANK (https://www.ebi.ac.uk/goldman-srv/webprank/) (Löytynoja et al., 2010). In this analysis, we used the whole coding sequences of the *Pias-1/RGA4* orthologs and the partial coding sequences (CC-NB-LRR-LII domains) of the *Pias-2/RGA5* orthologs. Second, we used the YN00 program (Yang and Nielsen, 2000) in PAML v4.8 (Yang 1997) to estimate the positive selection pressure on the *Pias-1/RGA4* and *Pias-2/RGA5* orthologs. The YN00 program can estimate the pairwise ω (also known as *dN/dS*) in the input sequence alignment; each resulting ω was graphically displayed in a heatmap (**Fig. 3B**).

Expression constructs used in the cell death assay

Expression constructs for five types of helper-NLRs, Pias-1 (*O. sativa* subsp. *indica*, WRC17), RGA4 (*O. sativa* subsp. *japonica*, Sasanishiki), RGA4-Oru (*O. rufipogon* accession W1943), RGA4-Oau (*O. australiensis* accession W0008), and RGA4-Ogr (*O. granulata* accession W0067B) (pCambia1300S-

"helper-NLR":HA), were generated by PCR amplification of the coding sequences from cDNA generated from leaf material and cloned into the binary vector pCambia1300S (http://www.cambia.org) that had been linearized by digestion with *Pst*I and *Spe*I. Expression constructs for two types of sensor NLRs, Pias-2 (*O. sativa* subsp. *indica*, WRC17) and RGA5 (*O. sativa* subsp. *japonica*, Sasanishiki) (pCambia1300S-FLAG: "sensor NLR"), were generated by PCR amplification of the coding sequences from cDNA generated from leaf material and cloned into the binary vector pCambia1300S (http://www.cambia.org) that had been linearized by digestion with *Sal*I and *Pst*I. The resulting vectors were introduced into *Agrobacterium tumefaciens* (strain GV3101). The primers used to generate the expression constructs are listed in **Suppl. Table 9**.

Immunoblot analysis

Proteins were expressed in *Nicotiana benthamiana* leaves and extracted from leaf tissue (approximately 100 mg) in 200 µL of extraction buffer (50 mM Tris-HCl [pH7.5], 150 mM NaCl and 0.1 % NP-40 [v/v]). The extracts were subjected to SDS-PAGE, followed by immunoblotting using anti-FLAG-HRP antibody, anti-HA-HRP antibody, or anti-AVR-Pia antibody. Anti-mouse IgG-HRP was used as a secondary antibody following incubation with anti-AVR-Pia antibody. Antibody-antigen complexes were detected using a luminescent image analyzer (ImageQuant LAS-4000) (Cytiva, Tokyo, Japan). Equal loading of proteins in the PAGE gel was confirmed by Coomassie blue staining.

Cell death assay in *N. benthamiana*

Transient expression of Pias/Pia allelic NLR and AVR-Pia was performed by infiltrating 4- to 5-week-old *N. benthamiana* plants with *A. tumefaciens* carrying the expression vector. *A. tumefaciens* suspensions in infiltration buffer (10 mM MES, 10 mM MgCl$_2$, and 150 µM acetosyringone, pH 5.6) were adjusted to the densities shown in **Suppl. Table 10**. The autofluorescence value under UV light was scored using a luminescent image analyzer (ImageQuant LAS-4000) (Cytiva, Tokyo, Japan).

Acknowledgements

This study was supported by JSPS KAKENHI 15H05779, 20H00421 and 16H06279 (PAGS) to RT and 21K14834 to MS, the Royal Society UK-Japan International exchange grants JPJSBP120215702 and IEC\R3\203081 to RT, SK and MB, the Biotechnology and Biological Sciences Research Council (UKRI-BBSRC, UK, grant BB/P012574), the European Research Council (ERC BLASTOFF project, 743165), the John Innes Foundation and the Gatsby Charitable Foundation. Computations were partially performed on the NIG supercomputer at ROIS National Institute of Genetics. We also thank the National Institute of Genetics (National Bioresource Project), Japan, and the Genebank at the National Agriculture and Food Research Organization, Japan, for providing the seeds of wild rice and WRC17, respectively.

Competing interests

S.K. receives funding from industry on NLR biology.

References in the main text

Adachi H, Derevnina L, Kamoun S. NLR singletons, pairs, and networks: evolution, assembly, and regulation of the intracellular immunoreceptor circuitry of plants. Curr Opin Plant Biol. 2019:121-131. doi: 10.1016/j.pbi.2019.04.007.

Adachi H, Kamoun S, Maqbool A. A resistosome-activated 'death switch'. Nat Plants. 2019b 5:457-458. doi: 10.1038/s41477-019-0425-9.

Ashikawa I, Hayashi N, Yamane H, Kanamori H, Wu J, Matsumoto T, Ono K, Yano M. Two adjacent nucleotide-binding site-leucine-rich repeat class genes are required to confer *Pikm*-specific rice blast resistance. Genetics. 2008 180:2267-76. doi: 10.1534/genetics.108.095034

Bailey PC, Schudoma C, Jackson W, Baggs E, Dagdas G, Haerty W, Moscou M, Krasileva KV. Dominant integration locus drives continuous diversification of plant immune receptors with exogenous domain fusions. Genome Biol. 2018 19:23. doi: 10.1186/s13059-018-1392-6.

Bernoux M, Burdett H, Williams SJ, Zhang X, Chen C, Newell K, Lawrence GJ, Kobe B, Ellis JG, Anderson PA, Dodds PN. Comparative Analysis of the Flax Immune Receptors L6 and L7 Suggests an Equilibrium-Based Switch Activation Model. Plant Cell. 2016 28:146-59. doi: 10.1105/tpc.15.00303.

Bi G, Su M, Li N, Liang Y, Dang S, Xu J, Hu M, Wang J, Zou M, Deng Y, Li Q, Huang S, Li J, Chai J, He K, Chen YH, Zhou JM. The ZAR1 resistosome is a calcium-permeable channel triggering plant immune signaling. Cell. 2021 S0092-867400600-0. doi: 10.1016/j.cell.2021.05.003.

Białas A, Langner T, Harant A, et al. Two NLR immune receptors acquired high-affinity binding to a fungal effector through convergent evolution of their integrated domain. Elife. 2021;10:e66961. Published 2021 Jul 21. doi:10.7554/eLife.66961

Cesari S, Thilliez G, Ribot C, Chalvon V, Michel C, Jauneau A, Rivas S, Alaux L, Kanzaki H, Okuyama Y, Morel JB, Fournier E, Tharreau D, Terauchi R, Kroj T. The rice resistance protein pair RGA4/RGA5 recognizes the *Magnaporthe oryzae* effectors AVR-Pia and AVR1-CO39 by direct binding. Plant Cell. 2013 25:1463-81. doi: 10.1105/tpc.112.107201.

Césari S, Kanzaki H, Fujiwara T, Bernoux M, Chalvon V, Kawano Y, Shimamoto K, Dodds P, Terauchi R, Kroj T. The NB-LRR proteins RGA4 and RGA5 interact functionally and physically to confer disease resistance. EMBO J. 2014 33:1941-59. doi: 10.15252/embj.201487923.

Clark RM, Schweikert G, Toomajian C, Ossowski S, Zeller G, Shinn P, Warthmann N, Hu TT, Fu G, Hinds DA, Chen H, Frazer KA, Huson DH, Schölkopf B, Nordborg M, Rätsch G, Ecker JR, Weigel D. Common sequence polymorphisms shaping genetic diversity in *Arabidopsis thaliana*. Science. 2007 317:338-42. doi: 10.1126/science.1138632.

Edwards SV, Hedrick PW. Evolution and ecology of MHC molecules: from genomics to sexual selection. Trends Ecol Evol. 1998 13:305-11. doi: 10.1016/s0169-5347(98)01416-5.

Fujisaki K, Abe Y, Ito A, Saitoh H, Yoshida K, Kanzaki H, Kanzaki E, Utsushi H, Yamashita T, Kamoun S, Terauchi R. Rice Exo70 interacts with a fungal effector, AVR-Pii, and is required for AVR-Pii-triggered immunity. Plant J. 2015 83:875-87. doi: 10.1111/tpj.12934.

Fujisaki K, Abe Y, KanzakiE, Ito K, Utsushi H, Saitoh H, Białas A, Banfield MJ, Kamoun S, Terauchi R. An unconventional NOI/RIN4 domain of a rice NLR protein binds host EXO70 protein to confer fungal immunity. bioRxiv. 2017 doi:10.1101/239400.

Hogenhout SA, Van der Hoorn RA, Terauchi R, Kamoun S. Emerging concepts in effector biology of plant-associated organisms. Mol Plant Microbe Interact. 2009 22:115-22. doi: 10.1094/MPMI-22-2-0115.

Horsefield S, Burdett H, Zhang X, Manik MK, Shi Y, Chen J, Qi T, Gilley J, Lai JS, Rank MX, Casey LW, Gu W, Ericsson DJ, Foley G, Hughes RO, Bosanac T, von Itzstein M, Rathjen JP, Nanson JD, Boden M, Dry IB, Williams SJ, Staskawicz BJ, Coleman MP, Ve T, Dodds PN, Kobe B. NAD$^+$ cleavage activity by animal and plant TIR domains in cell death pathways. Science. 2019 365:793-799. doi: 10.1126/science.aax1911.

Jia Y, McAdams SA, Bryan GT, Hershey HP, Valent B. Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. EMBO J. 2000 19:4004-14. doi: 10.1093/emboj/19.15.4004

Jones JD, Dangl JL. The plant immune system. Nature. 2006 444:323-9. doi: 10.1038/nature05286

Jupe F, Witek K, Verweij W, Sliwka J, Pritchard L, Etherington GJ, Maclean D, Cock PJ, Leggett RM, Bryan GJ, Cardle L, Hein I, Jones JD. Resistance gene enrichment sequencing (RenSeq) enables

reannotation of the NB-LRR gene family from sequenced plant genomes and rapid mapping of resistance loci in segregating populations. Plant J. 2013 76:530-44. doi: 10.1111/tpj.12307.

Kanzaki H, Yoshida K, Saitoh H, Fujisaki K, Hirabuchi A, Alaux L, Fournier E, Tharreau D, Terauchi R. Arms race co-evolution of *Magnaporthe oryzae AVR-Pik* and rice *Pik* genes driven by their physical interactions. Plant J. 2012 72:894-907. doi: 10.1111/j.1365-313X.2012.05110.x.

Kojima Y, Ebana K, Fukuoka S, Nagamine T, Kawase M. Development of an RFLP-based rice diversity research set of germplasm. Breeding Sci. 2005 55:431-440. doi: 10.1270/jsbbs.55.431

Kourelis J, van der Hoorn RAL. Defended to the Nines: 25 Years of Resistance Gene Cloning Identifies Nine Mechanisms for R Protein Function. Plant Cell. 2018 30:285-299. doi: 10.1105/tpc.17.00579.

Krasileva KV, Dahlbeck D, Staskawicz BJ. Activation of an *Arabidopsis* resistance protein is specified by the in planta association of its leucine-rich repeat domain with the cognate oomycete effector. Plant Cell. 2010 22:2444-58. doi: 10.1105/tpc.110.075358.

Kroj T, Chanclud E, Michel-Romiti C, Grand X, Morel JB. Integration of decoy domains derived from protein targets of pathogen effectors into plant immune receptors is widespread. New Phytol. 2016 210:618-26. doi: 10.1111/nph.13869.

Le Roux C, Huet G, Jauneau A, Camborde L, Trémousaygue D, Kraut A, Zhou B, Levaillant M, Adachi H, Yoshioka H, Raffaele S, Berthomé R, Couté Y, Parker JE, Deslandes L. A receptor pair with an integrated decoy converts pathogen disabling of transcription factors to immunity. Cell. 2015 161:1074-1088. doi: 10.1016/j.cell.2015.04.025.

Li P, Quan X, Jia G, Xiao J, Cloutier S, You FM. RGAugury: a pipeline for genome-wide prediction of resistance gene analogs (RGAs) in plants. BMC Genomics. 2016 17:852. doi: 10.1186/s12864-016-3197-x.

Liu J, Wang X, Mitchell T, Hu Y, Liu X, Dai L, Wang GL. Recent progress and understanding of the molecular mechanisms of the rice-*Magnaporthe oryzae* interaction. Mol Plant Pathol. 2010 11:419-27. doi: 10.1111/j.1364-3703.2009.00607.x.

Lv F, Wang H, Wang X, et al. GhCFE1A, a dynamic linker between the ER network and actin cytoskeleton, plays an important role in cotton fibre cell initiation and elongation. J Exp Bot. 2015;66(7):1877-1889. doi:10.1093/jxb/eru530

Ma S, Lapin D, Liu L, Sun Y, Song W, Zhang X, Logemann E, Yu D, Wang J, Jirschitzka J, Han Z, Schulze-Lefert P, Parker JE, Chai J. Direct pathogen-induced assembly of an NLR immune receptor complex to form a holoenzyme. Science. 2020 370:eabe3069. doi: 10.1126/science.abe3069.

Maidment JHR, Franceschetti M, Maqbool A, Saitoh H, Jantasuriyarat C, Kamoun S, Terauchi R, Banfield MJ. Multiple variants of the fungal effector AVR-Pik bind the HMA domain of the rice protein OsHIPP19, providing foundation to engineer plant defense. J Biol Chem 2021 196:100371. doi: 10.1016/j.jbc.2021. 100371.

Maqbool A, Saitoh H, Franceschetti M, Stevenson CE, Uemura A, Kanzaki H, Kamoun S, Terauchi R, Banfield MJ. Structural basis of pathogen recognition by an integrated HMA domain in a plant NLR immune receptor. Elife. 2015 4:e08709. doi: 10.7554/eLife.08709.

Martin R, Qi T, Zhang H, Liu F, King M, Toth C, Nogales E, Staskawicz BJ. Structure of the activated Roq1 resistosome directly recognizing the pathogen effector XopQ. bioRxiv. 2020 doi: https://doi.org/10.1101/2020.08.13.246413.

Meng X, Xiao G, Telebanco-Yanoria MJ, Siazon PM, Padilla J, Opulencia R, Bigirimana J, Habarugira G, Wu J, Li M, Wang B, Lu GD, Zhou B. The broad-spectrum rice blast resistance (*R*) gene *Pita2* encodes a novel R protein unique from *Pita*. Rice (N Y). 2020 13:19. doi: 10.1186/s12284-020-00377-5.

Oikawa K, Fujisaki K, Shimizu M, Takeda T, Saitoh H, Hirabuchi A, Hiraka Y, Białas A, Langner T, Kellner R, Bozkurt T, Cesari S, Kroj T, Maidment J, Banfield M, Kamoun S, Terauchi R. The blast pathogen effector AVR-Pik binds and stabilizes rice heavy metal-associated (HMA) proteins to co-opt their function in immunity. bioRxiv. 2020 doi:10.1101/2020.12.01.406389.

Okuyama Y, Kanzaki H, Abe A, Yoshida K, Tamiru M, Saitoh H, Fujibe T, Matsumura H, Shenton M, Galam DC, Undan J, Ito A, Sone T, Terauchi R. A multifaceted genomics approach allows the isolation of the rice *Pia*-blast resistance gene consisting of two adjacent NBS-LRR protein genes. Plant J. 2011 66:467-79. doi: 10.1111/j.1365-313X.2011.04502.x.

Orbach MJ, Farrall L, Sweigard JA, Chumley FG, Valent B. A telomeric avirulence gene determines efficacy for the rice blast resistance gene Pi-ta. Plant Cell. 2000 12:2019-32. doi: 10.1105/tpc.12.11.2019.

Pan Q, Wendel J, Fluhr R. Divergent evolution of plant NBS-LRR resistance gene homologues in dicot and cereal genomes. J Mol Evol. 2000 50:203-13. doi: 10.1007/s002399910023.

Park CH, Shirsekar G, Bellizzi M, Chen S, Songkumarn P, Xie X, Shi X, Ning Y, Zhou B, Suttiviriya P, Wang M, Umemura K, Wang GL. The E3 Ligase APIP10 Connects the Effector AvrPiz-t to the NLR Receptor Piz-t in Rice. PLoS Pathog. 2016 12:e1005529. doi: 10.1371/journal.ppat.1005529.

Pennisi E. Armed and dangerous. Science. 2010 327:804-5. doi: 10.1126/science.327.5967.804.

Piertney SB, Oliver MK. The evolutionary ecology of the major histocompatibility complex. Heredity (Edinb). 2006 96:7-21. doi: 10.1038/sj.hdy.6800724.

Sarris PF, Cevik V, Dagdas G, Jones JD, Krasileva KV. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. BMC Biol. 2016 14:8. doi: 10.1186/s12915-016-0228-7.

Sharif H, Wang L, Wang WL, Magupalli VG, Andreeva L, Qiao Q, Hauenstein AV, Wu Z, Núñez G, Mao Y, Wu H. Structural mechanism for NEK7-licensed activation of NLRP3 inflammasome. Nature. 2019 570:338-343. doi: 10.1038/s41586-019-1295-z.

Takagi H, Uemura A, Yaegashi H, Tamiru M, Abe A, Mitsuoka C, Utsushi H, Natsume S, Kanzaki H, Matsumura H, Saitoh H, Yoshida K, Cano LM, Kamoun S, Terauchi R. MutMap-Gap: whole-genome resequencing of mutant F2 progeny bulk combined with de novo assembly of gap regions identifies the rice blast resistance gene *Pii*. New Phytol. 2013 200:276-283. doi: 10.1111/nph.12369.

Takahata N, Nei M. Allelic genealogy under overdominant and frequency-dependent selection and polymorphism of major histocompatibility complex loci. Genetics. 1990 124:967-78.

Tenthorey JL, Haloupek N, López-Blanco JR, Grob P, Adamson E, Hartenian E, Lind NA, Bourgeois NM, Chacón P, Nogales E, Vance RE. The structural basis of flagellin detection by NAIP5: A strategy to limit pathogen immune evasion. Science. 2017 358:888-893. doi: 10.1126/science.aao1140.

Terauchi R, Yoshida K. Towards population genomics of effector-effector target interactions. New Phytol. 2010 187:929-39. doi: 10.1111/j.1469-8137.2010.03408.x.

Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M. Arabidopsis systemic immunity uses conserved defense signaling pathways and is mediated by jasmonates. Proc Natl Acad Sci U S A. 2007 104:1075-80. doi: 10.1073/pnas.0605423104.

Van de Weyer AL, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones JDG, Dangl JL, Weigel D, Bemm F. A Species-Wide Inventory of NLR Genes and Alleles in *Arabidopsis thaliana*. Cell. 2019 178:1260-1272.e14. doi: 10.1016/j.cell.2019.07.038.

Wan L, Essuman K, Anderson RG, Sasaki Y, Monteiro F, Chung EH, Osborne Nishimura E, DiAntonio A, Milbrandt J, Dangl JL, Nishimura MT. TIR domains of plant immune receptors are NAD^+-cleaving enzymes that promote cell death. Science. 2019 365:799-803. doi: 10.1126/science.aax1771.

Wang J, Wang J, Hu M, Wu S, Qi J, Wang G, Han Z, Qi Y, Gao N, Wang HW, Zhou JM, Chai J. Ligand-triggered allosteric ADP release primes a plant NLR complex. Science. 2019a 364:eaav5868. doi: 10.1126/science.aav5868.

Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang HW, Zhou JM, Chai J. Reconstitution and structure of a plant NLR resistosome conferring immunity. Science. 2019b 364:eaav5870. doi: 10.1126/science.aav5870.

Wang R, Ning Y, Shi X, He F, Zhang C, Fan J, Jiang N, Zhang Y, Zhang T, Hu Y, Bellizzi M, Wang GL. Immunity to Rice Blast Disease by Suppression of Effector-Triggered Necrosis. Curr Biol. 2016 26:2399-2411. doi: 10.1016/j.cub.2016.06.072.

Williams SJ, Sohn KH, Wan L, Bernoux M, Sarris PF, Segonzac C, Ve T, Ma Y, Saucet SB, Ericsson DJ, Casey LW, Lonhienne T, Winzor DJ, Zhang X, Coerdt A, Parker JE, Dodds PN, Kobe B, Jones JD. Structural basis for assembly and function of a heterodimeric plant immune receptor. Science. 2014 344:299-303. doi: 10.1126/science.1247357.

Woolhouse ME, Webster JP, Domingo E, Charlesworth B, Levin BR. Biological and biomedical implications of the co-evolution of pathogens and their hosts. Nat Genet. 2002 32:569-77. doi: 10.1038/ng1202-569.

Wu CH, Abd-El-Haliem A, Bozkurt TO, Belhaj K, Terauchi R, Vossen JH, Kamoun S. NLR network mediates immunity to diverse plant pathogens. Proc Natl Acad Sci U S A. 2017 114:8113-8118. doi: 10.1073/pnas.1702041114.

Wu Y, Xiao N, Yu L, Pan C, Li Y, Zhang X, Liu G, Dai Z, Pan X, Li A. Combination Patterns of Major R Genes Determine the Level of Resistance to the *M. oryzae* in Rice (*Oryza* sativa L.). PLoS One. 2015 10:e0126130. doi: 10.1371/journal.pone.0126130.

Yang Z, Nielsen R. Estimating synonymous and nonsynonymous substitution rates under realistic evolutionary models. Mol Biol Evol. 2000 17:32-43. doi: 10.1093/oxfordjournals.molbev.a026236.

Yoshida K, Saitoh H, Fujisawa S, Kanzaki H, Matsumura H, Yoshida K, Tosa Y, Chuma I, Takano Y, Win J, Kamoun S, Terauchi R. Association genetics reveals three novel avirulence genes from the rice blast fungal pathogen *Magnaporthe oryzae*. Plant Cell. 2009 21:1573-91. doi: 10.1105/tpc.109.066324.

Zdrzałek R, Kamoun S, Terauchi R, Saitoh H, Banfield MJ. The rice NLR pair Pikp-1/Pikp-2 initiates cell death through receptor cooperation rather than negative regulation. PLoS One. 2020 15:e0238616. doi: 10.1371/journal.pone.0238616.

Zhao H, Wang X, Jia Y, Minkenberg B, Wheatley M, Fan J, Jia MH, Famoso A, Edwards JD, Wamishe Y, Valent B, Wang GL, Yang Y. The rice blast resistance gene *Ptr* encodes an atypical protein required for broad-spectrum disease resistance. Nat Commun. 2018 9:2039. doi: 10.1038/s41467-018-04369-4.

Zhang L, Chen S, Ruan J, Wu J, Tong AB, Yin Q, Li Y, David L, Lu A, Wang WL, Marks C, Ouyang Q, Zhang X, Mao Y, Wu H. Cryo-EM structure of the activated NAIP2-NLRC4 inflammasome reveals nucleated polymerization. Science. 2015 350:404-9. doi: 10.1126/science.aac5789.

Zhang Y, Zhang F, Huang X. Characterization of an *Arabidopsis thaliana* DUF761-containingprotein with a potential role in development and defense responses. Theor. Exp. Plant Physiol. 2019 31:303-316.

References in the MM and the supplemental data (Figure and Table)

Arazoe T, Miyoshi K, Yamato T, Ogawa T, Ohsato S, Arie T, Kuwata S. Tailor-made CRISPR/Cas system for highly efficient targeted gene replacement in the rice blast fungus. Biotechnol Bioeng. 2015 112:2543-9. doi: 10.1002/bit.25662. Epub 2015 Jul 14. PMID: 26039904.

Bennetzen JL, Schmutz J, Wang H, et al. Reference genome sequence of the model plant Setaria. Nat Biotechnol. 2012;30:555-561. Published 2012 May 13. doi:10.1038/nbt.2196

Du H, Yu Y, Ma Y, et al. Sequencing and *de novo* assembly of a near complete indica rice genome. Nat Commun. 2017;8:15324. Published 2017 May 4. doi:10.1038/ncomms15324

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol. 2018 35:518-522. doi: 10.1093/molbev/msx281. PMID: 29077904; PMCID: PMC5850222.

Hyon GS, Naga NTT, Chuma I, Inoue Y, Asano H, Murata N, Tosa Y. Characterization of interactions between barley and various host-specific subgroups of *Magnaporthe oryzae* and *M. grisea*. J. Gen. Plant Pathol. 2012 78: 237-246.

International Rice Genome Sequencing Project. The map-based sequence of the rice genome. Nature. 2005;436:793-800. doi:10.1038/nature03895

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods. 2017 14:587-589. doi: 10.1038/nmeth.4285. Epub 2017 May 8. PMID: 28481363; PMCID: PMC5453245.

Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. Nat Methods. 2015;12:357-360. doi:10.1038/nmeth.3317

Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics. 2010;26:589-595. doi:10.1093/bioinformatics/btp698

Löytynoja A, Goldman N. webPRANK: a phylogeny-aware multiple sequence aligner with interactive alignment browser. BMC Bioinformatics. 2010 11:579. doi: 10.1186/1471-2105-11-579. PMID: 21110866; PMCID: PMC3009689.

Mascher M, Gundlach H, Himmelbach A, et al. A chromosome conformation capture ordered sequence of the barley genome. Nature. 2017;544:427-433. doi:10.1038/nature22043

Mikami M, Toki S, Endo M. Comparison of CRISPR/Cas9 expression constructs for efficient targeted mutagenesis in rice. Plant Mol Biol. 2015 88:561-72. doi: 10.1007/s11103-015-0342-x. Epub 2015 Jul 19. PMID: 26188471; PMCID: PMC4523696.

Miki D, Shimamoto K. Simple RNAi vectors for stable and transient suppression of gene function in rice. Plant Cell Physiol. 2004 45:490-5. doi: 10.1093/pcp/pch048. PMID: 15111724.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol

Evol. 2020 May 1;37(5):1530-1534. doi: 10.1093/molbev/msaa015. Erratum in: Mol Biol Evol. 2020 37:2461. PMID: 32011700; PMCID: PMC7182206.

Okuyama Y, Kanzaki H, Abe A, Yoshida K, Tamiru M, Saitoh H, Fujibe T, Matsumura H, Shenton M, Galam DC, Undan J, Ito A, Sone T, Terauchi R. A multifaceted genomics approach allows the isolation of the rice *Pia*-blast resistance gene consisting of two adjacent NBS-LRR protein genes. Plant J. 2011 66:467-79. doi: 10.1111/j.1365-313X.2011.04502.x.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol. 2015;33:290-295. doi:10.1038/nbt.3122

Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Methods. 2012 9:671-5. doi: 10.1038/nmeth.2089. PMID: 22930834; PMCID: PMC5554542.

Shimizu M, Nakano Y, Hirabuchi A, Yoshino K, Kobayashi M, Yamamoto K, Terauchi R, Saitoh H. RNA-Seq of in planta-expressed *Magnaporthe oryzae* genes identifies MoSVP as a highly expressed gene required for pathogenicity at the initial stage of infection. Mol Plant Pathol. 2019 20:1682-1695. doi: 10.1111/mpp.12869. Epub 2019 Sep 27. PMID: 31560822; PMCID: PMC6859710.

Sweigard J, Chumley F, Carroll A, Farrall L, Valent B. A series of vectors for fungal transformation. Fungal Genet. Newsl. 1997 44, 52–55.

Wu Z, Fang D, Yang R, et al. *De novo* genome assembly of *Oryza granulata* reveals rapid genome expansion and adaptive evolution. Commun Biol. 2018;1:84. Published 2018 Jun 29. doi:10.1038/s42003-018-0089-4

Yang Z. PAML: a program package for phylogenetic analysis by maximum likelihood. Comput Appl Biosci. 1997 13:555-6. doi: 10.1093/bioinformatics/13.5.555. PMID: 9367129.

Yang Z, Nielsen R. 2000. estimating synonymous and nonsynonymous substitution rates under realistic evolutionary models. Mol Biol Evol17:32–43. doi:10.1093/oxfordjournals.molbev.a026236

Yoshida K, Saitoh H, Fujisawa S, Kanzaki H, Matsumura H, Yoshida K, Tosa Y, Chuma I, Takano Y, Win J, Kamoun S, Terauchi R. Association genetics reveals three novel avirulence genes from the rice blast fungal pathogen *Magnaporthe oryzae*. Plant Cell. 2009 21:1573-91. doi: 10.1105/tpc.109.066324.

Zhao Q, Feng Q, Lu H, et al. Pan-genome analysis highlights the extent of genomic variation in cultivated and wild rice [published correction appears in Nat Genet. 2018 Aug;50:1196]. Nat Genet. 2018;50:278-284. doi:10.1038/s41588-018-0041-z

Zimin AV, Puiu D, Luo MC, et al. Hybrid assembly of the large and highly repetitive genome of Aegilops tauschii, a progenitor of bread wheat, with the MaSuRCA mega-reads algorithm. Genome Res. 2017;27(5):787-792. doi:10.1101/gr.213405.116

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA genome assembler. Bioinformatics. 2013;29:2669-2677. doi:10.1093/bioinformatics/btt476

Figure legends

Figure 1. The *Pias* gene of rice line WRC17 (cultivar Keiboba) encodes a CC-NLR protein and is allelic to *Pia*.

(A) Segregation of the resistance and susceptibility traits among the 58 RILs derived from a cross between WRC17 (cultivar Keiboba) and Hitomebore. Disease symptoms of WRC17, Hitomebore, and six RILs showing a susceptible phenotype after punch inoculation of *Magnaporthe oryzae* isolate 2012-1 (leaf photographs) and the frequency distribution of disease lesion area of the 58 RILs (bar graphs). (B) Linkage maps of candidate NLR genes at the *PiW17-1* and *PiW17-2* loci. (C) Both *CNL-04* and *CNL-05* are required for *PiW17-1*-mediated resistance against *M. oryzae* 2012-1. HW-RIL7 contains only *PiW17-1* and is resistant to *M. oryzae* 2012-1. Knockout of *CNL-04* (*cnl-04*) and *CNL-05* (*cnl-05*) in HW-RIL7 rendered plants susceptible to 2012-1. (D) Gene structures of *Pia* consisting of *RGA4* and *RGA5* and *Pias* consisting of *Pias-1* and *Pias-2*. The positions of protein domains (CC, NB-ARC, LRR, HMA, and DUF761) encoded by the NLRs are indicated. (E) The *M. oryzae* 2012-1 *AVR-Pias* knockout mutant became virulent to HW-RIL7. (F) Amino acid sequence of G9532 protein (AVR-Pias). The secretion signal is indicated by red letters and the Toxin18-like motif by blue letters.

Figure 2. Recurrent integration of extraneous domains in Pias/Pia sensor NLRs.

(A) A simplified scheme of the structures of the Pias/Pia NLR pairs. Pias-1/RGA4 helper NLRs are shown in green, and Pias-2/RGA5 sensor NLRs are shown in white. The conserved LRR-ID intervening motif (LII) is indicated by black ellipses. Integrated domains are shown by different-colored hexagons. (B) A sequence logo showing conserved amino acids of the LRR-ID Intervening (LII) motif. (C) Distribution of ID motifs among *Oryza* species. The pie charts show the frequencies of different ID motifs in a given species. The colors correspond to the ID colors in (A). The numbers below the pie charts indicate the sample numbers. A cladogram showing the phylogenetic relationships of 11 *Oryza* species and four other Poaceae species (*Setaria italica*, *Panicum hallii*, *Hordeum vulgare*, and *Aegilops tauschii*) based on Time

Tree, the time scale of life web database (http://www.timetree.org/). The numbers on the branches indicate the estimated time of the splitting of lineages (MYA: million years ago). (D) DNA sequence similarity between *O. punctata Pias-2/RGA5 sensor NLR* and the downstream sequence of *O. sativa* (Nipponbare) *Pias-2/RGA5*. LII: LRR-ID intervening motif. (E) Dot-plot analysis of the *O. punctata* (W1582) *Pias-2/RGA5 sensor NLR* and *O. sativa* (Nipponbare) *Pias-2/RGA5 NLR* downstream sequences using the Dotmatcher tool (http://emboss.bioinformatics.nl/cgi-bin/emboss/dotmatcher/). (F) Possible evolutionary process of ID replacement that might have occurred between the *O. punctata* and the *O. sativa Pias-2/RGA5* lineages. We still do not know the mode of interaction between the AVR-Pias effector and DUF761-containing protein, so it is indicated by "?".

Figure 3. Contrasting evolutionary patterns of the helper and sensor NLRs of the *Pias/Pia* locus.
(A) Phylogenetic tree of the *Pias-1/RGA4 helper NLR* gene (left) and *Pias-2/RGA5 sensor NLR* (right) gene based on the full-length amino acid sequence of Pias-1/RGA4 and the sequence in the region CC to LII for Pias-2/RGA5. *Pias-2/RGA5* sensor NLRs form two major clades (C1 and C2). The numbers indicate bootstrap values. (B) Pairwise comparison of ω, the ratio of nonsynonymous to synonymous nucleotide substitutions for *Pias-1/RGA4* (above the diagonal: the full-length NLR DNA sequence was used) and *Pias-2/RGA5* (below the diagonal: the sequence of the CC to LII region was used).

Figure 4. The NLR helper Pias-1 is functionally conserved.
(A) Representative images of *N. benthamiana* leaves after agroinfiltration with Pias-1:HA, Pias-1:HA/FLAG:RGA5, FLAG:RGA5/AVR-Pia, and Pias-1:HA/FLAG:RGA5/AVR-Pia (left) and RGA4-Ogr:HA derived from *O. granulata*, RGA4-Ogr:HA/FLAG:RGA5, FLAG:RGA5/AVR-Pia, and RGA4-Ogr:HA/FLAG:RGA5/AVR-Pia (right). Autofluorescence under UV light is shown. (B) Pias-1 cooperates with RGA5 to recognize AVR-Pia and induces resistance in rice. The rice line HW-RIL7 with *Pias* (*Pias-1* and *Pias-2*) recognizes the Ao-92-06-2 strain with *AVR-Pias* (Ao92-06-2+*pex22p:AVR-Pias*) and induces resistance. However, HW-RIL7 cannot recognize the Ao02-06-2 strain with *AVR-Pia* (Ao92-06-2+*pex22p:AVR-Pia*). Two lines (Line #1 and #2) contain the *35S-RGA5* transgene in the HW-RIL7 background. F1-A is a progeny derived from a cross between Sasanishiki with *Pia* (*RGA4* and *RGA5*) and HW-RIL7. F1-B is a progeny derived from a cross between a Sasanishiki mutant (Sas1493) with *pia* (*rga4* and *RGA5*) and HW-RIL7.

Legends to Supplementary Figures

Suppl. Fig. 1. The RaIDeN method used to identify candidate genes responsible for the resistance of rice line WRC17 to *M. oryzae* 2012-1.
(A) Segregation of the resistance and susceptibility traits among the 58 RILs derived from a cross between WRC17 (cultivar Keiboba) and Hitomebore. (B) The whole-genome sequence of the resistant parent WRC17 was obtained by Illumina sequencing. RNA-seq of WRC17 leaves inoculated with *M. oryzae* was

also performed. Next, short reads of whole-genome sequences of the susceptible parent Hitomebore and the six susceptible RILs were obtained by Illumina sequencing. (C) *De novo* assembly of R-parental line WRC17 was performed using *DISCOVAR* software. (D) Expressed genes of WRC17 were identified by mapping RNA-seq reads to the WRC17 reference genome. (E) Mapping of short reads of the susceptible parent (Hitomebore) and susceptible RILs to the predicted genes identified DNA polymorphisms shared by the susceptible lines using the RaIDeN pipeline (the script of the RaIDeN pipeline and details about the pipeline are available at https://github.com/YuSugihara/RaIDeN).

Suppl. Fig. 2. Mapping of *NLR*s in the *PiW17-1* and *PiW17-2* loci that confer resistance against *M. oryzae* 2012-1 to WRC17. (A) Association between resistance/susceptible phenotypes and genotypes among the RILs (the primers used for genotyping are shown in Suppl. Table 9). *CNL-04* and *CNL-05*, *CNL-06* and *CNL-07*, and *CNL-08*, *NL-2*, and *CNL-09* are located on the same contigs. (B) Fine maps of candidate NLR genes in *PiW17-1* and *PiW17-2*. Blue and yellow indicate WRC17- and Hitomebore-type genotypes, respectively. The 10 NLRs shown in the red rectangle were considered to be the candidates of *Pi-W17-1*.

Suppl. Fig. 2 (continued). Results of RNAi-mediated silencing of candidate *NLR* genes.
(C) RNAi-mediated silencing of eight candidate genes encoding proteins over 900 amino acids long was performed. Gene silencing of *CNL-04* and *CNL-05* made plants susceptible to *M. oryzae* 2012-1. (D) Gene expression levels of *CNL-04* and *CNL-05* in plants transformed with the RNAi constructs. Control (Con) is WRC17 transformed with empty vector.

Suppl. Fig. 3. **CRISPR/Cas9-mediated knockout of *CNL-04* abolishes *PiW17-1*-mediated resistance against *M. oryzae* isolate 2012-1.**
(A) The position of guide DNA targeted to *CNL-04*. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. (B) Sequences of the sgRNA positions in the *CNL-04* knockout lines. (C) Resistance/susceptible phenotypes of the selfed progeny of the *cnl-04_1.14.2* line heterozygous for wild-type (*a*) and mutated (*b*: 4-bp deletion) alleles. The *b/b* homozygous plants became susceptible.

Suppl. Fig. 3 (**continued**). **CRISPR/Cas9-mediated knockout of *CNL-05* abolishes *PiW17-1* mediated resistance against *M. oryzae* isolate 2012-1.**
(D) The positions of guide DNAs targeted to *CNL-05*. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. (E) Sequences of the sgRNA positions in the *CNL-05* knockout lines. (F) Resistance/susceptible phenotypes of the selfed progeny of *cnl-05_3.7.2* line heterozygous for wild-type (*a*) and mutated (*b*: 1-bp insertion) alleles. The *b/b* homozygous plants became susceptible.

Suppl. Fig. 4. Comparison of the gene structures of *Pia* (*RGA4* and *RGA5*) and *Pias* (*Pias-1* and *Pias-2*).
DNA sequence similarities obtained by ClustalW are shown.

Suppl. Fig. 5. Gene structures and predicted amino acid sequences of *Pias-1* and *Pias-2*.

(A) Gene structures of *Pias-1* and *Pias-2*. (B) Amino acid sequences of Pias-1 and Pias-2 with the putative domains.

Suppl. Fig. 6. Isolation of *AVR-Pias* from *M. oryzae* isolate 2012-1.

A simplified scheme showing the procedure used to isolate *AVR-Pias*. First, the genome of isolate 2012-1 was sequenced on the Illumina platform, and the short reads were used for *de novo* assembly with *DISCOVAR*, resulting in the 2012-1 genome reference (**Suppl. Table 4, 5**). Next, the isolate 2012-1 was used to inoculate barley (*Hordeum vulgare*) cultivar 'Nigrate', which is highly susceptible to *M. oryzae* (Hyon et al. 2012), and the infected leaves were subjected to RNA-seq (Shimizu et al. 2019), resulting in the identification of 10,991 genes in 2012-1 that were expressed during host infection. Next, eight *M. oryzae* isolates (r1–r8) were selected and used for inoculation of rice line HW-RIL7 with *Pias* but without *Pi-W17-2*. The eight isolates were compatible with HW-RIL7 (result shown in the bottom), suggesting that they lack *AVR-Pias*. The eight isolates were then subjected to genome resequencing on the Illumina platform, and the RaIDeN method was used to identify only presence/absence polymorphisms (**Suppl. Table 4**). The short reads of eight isolates were aligned to the expressed genes of the 2012-1 isolate, resulting in the identification of 87 expressed genes that were specific to isolate 2012-1. These transcripts were further filtered based on four criteria: (1) genes showing a higher level of expression 24 h after inoculation (FPKM > 200) and genes encoding (2) a putative secreted protein, (3) a non-transmembrane protein, and (4) a small protein (<150 amino acids). This analysis identified three transcripts (*G9141*, *G9435*, and *G9532*) as the candidates of *AVR-Pias*.

Suppl. Fig. 7. The expressed gene *G9532* of *M. oryzae* isolate 2012-1 is *AVR-Pias*.

(A) Results of inoculation of rice line HW-RIL7 with *Pias* or Hitomebore without *Pias* with *M. oryzae* isolate Ao92-06-2 wild type (WT) or Ao92-06-2 with the *G9141*, *G9435*, or *G9532* transgene, all driven by the *pex22* promoter. When Ao92-06-2 contained *G9532*, the interaction became incompatible, indicating that *G9532* is *AVR-Pias*. (B) Introduction of *pex22p*:*G9532* into *M. oryzae* isolate Ao92-06-2 confers avirulence to the pathogen against HW-RIL7 with *Pias-1* and *Pias-2*. Knockout of the host gene *Pias-1* or *Pias-2* compromises the avirulence caused by *G9532*.

Suppl. Fig. 8. Deletion of *AVR-Pias* from *M. oryzae* isolate 2012-1 causes a loss of avirulence against rice line HW-RIL7 with *Pias*.

(A) Schematic overview of the target gene replacement (TGR) strategy at the *AVR-Pias* locus using RNA-guided nuclease. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. The primers used for PCR analysis are indicated by horizontal arrows. (B) Inoculation assay of rice line HW-RIL7 with *Pias* wild type (WT) and TGR transformants. (C) PCR analysis of TGR events at the *AVR-Pias* locus. Upper and lower images show PCR results using *AVR-Pias*- and *Hygromycin* (*Hyg*)-

specific primers, respectively. pCB1636: Replacement vector containing the hygromycin resistance gene and genomic regions neighboring *AVR-Pias*. The isolates corresponding to lanes 2–5 show compatibility with rice line HW-RIL7 with *Pias*, whereas the wild-type isolate (2012-1) in lane 6 is incompatible with HW-RIL7.

Suppl. Fig. 9. Positions of the LII motif (this study) and the CID motif (Bailey et al. 2018).

Amino acid sequence alignment of RGA5, Pias-2, and LOC_Os12g18360 and the positions of the LII motif (this study) and the CID motif as described by Bailey et al. (2018). The aqua and orange boxes indicate the LII and CID motif regions, respectively.

Suppl. Fig. 10. The *O. punctata* W1582 LII-ID sequence is conserved in the downstream sequence of *Pias* in *O. sativa* cv. Nipponbare.

Alignment of the DNA sequences of the *O. punctata* W1582 LII-ID and the W1582 LII-ID like sequence in *O. sativa* Nipponbare. The blue boxes and green lines indicate the exon sequence of *O. punctata* W1582 Pias-2 homolog and the LII region, respectively.

Suppl. Fig. 11. The LII + ID sequence of the *O. punctata* (W1582) *Pias-2* homolog is conserved in the downstream sequence of *O. meridionalis* (W2112) Pias-2.

Dot-plot analysis of *O. punctata* (W1582) *Pias-2* homolog and the genome sequence of *O. meridionalis* (W2112). The purple line corresponds to block (a) in Figure 2. The block (b) sequence in Figure 2 is deleted in *O. meridionalis* (W2112).

Suppl. Fig. 12 HR-like cell death after overexpression of helper NLRs of the RGA4/Pias-1 lineage in *N. benthamiana*.

(A) Box plots of autofluorescence values after transient expression of RGA4/Pias-1 homologs. GUS vector was included as a control. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Representative image of Pias-1/RGA4-meditated HR in *N. benthamiana*. (C) Immunoblot analysis of Pias-1:HA, RGA4:HA, RGA4-Oau:HA (cloned from *O. australiensis* accession W0008), RGA4-Oru:HA (cloned from *O. rufipogon* accession W1943), and RGA4-Ogr:HA (cloned from *O. granulata* accession W0067B) proteins detected by anti-HA (α-HA) antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size marker are indicated on the left (kDa)

Suppl. Fig. 13 HR-like cell death after overexpression of Pias-1:HA helper and Pias-1/RGA5 sensors in *N. benthamiana*.

(A) Box plots of autofluorescence values after transient expression of Pias-1:HA, FLAG:Pias-2, and FLAG:RGA5 separately or in combination in *N. benthamiana*. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Representative image of HR after transient

expression of Pias-1:HA, FLAG:Pias-2, and FLAG:RGA5 separately or in combination in *N. benthamiana*. (C) Immunoblot analysis of Pias-1:HA protein detected by anti-HA (*α*-HA) antibody and FLAG:Pias-2 and FLAG:RGA5 proteins detected by anti-FLAG antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size marker are indicated on the left (kDa).

Suppl. Fig. 14. HR-like cell death caused by Pias1 and RGA4-Ogr expression is suppressed by RGA5 expression, and additional AVR-Pia expression induces cell death.

(A) Box plots of autofluorescence values after transient expression of Pias-1:HA, FLAG:RGA5, and AVR-Pia separately or in combination in *N. benthamiana* leaves. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Immunoblot analysis of Pias-1:HA protein detected by anti-HA (*α*-HA) antibody, FLAG:RGA5 protein detected by anti-FLAG antibody, and AVR-Pia detected by anti-AVR-Pia antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size markers are indicated on the left (kDa). (C) Box plots of autofluorescence values after transient expression of RGA4-Ogr:HA, FLAG:RGA5, and AVR-Pia separately or in combination in *N. benthamiana* leaves.The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (D) Immunoblot analysis of RGA4-Ogr:HA protein detected by anti-HA (*α*-HA) antibody, FLAG:RGA5 protein detected by anti-FLAG (*α*-FLAG) antibody, and AVR-Pia detected by anti-AVR-Pia (*α*-AVR-Pia) antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size markers are indicated on the left (kDa).

Suppl. Fig. 15. Results of inoculation of HW-RIL7, Sasanishiki, and Sas1493 rice plants with *M. oryzae* isolates Ao92-06-2, Ao92-06-2 +pex22p:AVR-Pia, and Ao92-06-2 +pex22p:AVR-Pias.

Suppl. Fig. 16. The DUF761-containing gene family in rice.
A BLASTP search was performed using the amino acid sequence of the DUF761 domain of Pias-2 as a query. Fifteen DUF761 domain-containing genes were retrieved using a cutoff e-value < 10. The middle panel shows the amino acid sequence alignment, and the right panel shows the domain structure of each protein.

Figure 1. The *Pias* gene of rice line WRC17 (cultivar Keiboba) encodes a CC-NLR protein and is allelic to *Pia*.

(A) Segregation of the resistance and susceptibility traits among the 58 RILs derived from a cross between WRC17 (cultivar Keiboba) and Hitomebore. Disease symptoms of WRC17, Hitomebore and six RILs showing a susceptible phenotype after punch inoculation of *Magnaporthe oryzae* isolate 2012-1 (leaf photographs) and the frequency distribution of disease lesion area of the 58 RILs (bar graphs). (B) Linkage maps of candidate NLR genes at the *PiW17-1* and *PiW17-2* loci. (C) Both *CNL-04* and *CNL-05* are required for *PiW17-1*-mediated resistance against *M. oryzae* 2012-1. HW-RIL7 contains only *PiW17-1* and is resistant to *M. oryzae* 2012-1. Knockout of *CNL-04* (*cnl-04*) and *CNL-05* (*cnl-05*) in HW-RIL7 rendered plants susceptible to 2012-1. (D) Gene structures of *Pia* consisting of *RGA4* and *RGA5* and *Pias* consisting of *Pias-1* and *Pias-2*. The positions of protein domains (CC, NB-ARC, LRR, HMA, and DUF761) encoded by the NLRs are indicated. (E) The *M. oryzae* 2012-1 *AVR-Pias* knockout mutant became virulent to HW-RIL7. (F) Amino acid sequence of G9532 protein (AVR-Pias). The secretion signal is indicated by red letters and the Toxin18-like motif by blue letters.

(A) STEP1: Phenotyping of RILs

(B) STEP2: Whole genome sequencing of parental lines and susceptible RILs (S-RILs).

(C) STEP3: *De novo* assembly of R-parental line "WRC17" using "*DISCOVAR*" software.

(D) STEP4: Gene prediction of WRC17, the resistant parent, using RNA-seq data.

(E) STEP5: Mapping short reads of susceptible parent and RILs to the predicted genes of WRC17.

Suppl. Fig. 1. The RaIDeN method used to identify candidate genes responsible for the resistance of rice line WRC17 to *M. oryzae* 2012-1.
(A) Segregation of the resistance and susceptibility traits among the 58 RILs derived from a cross between WRC17 (cultivar Keiboba) and Hitomebore. (B) The whole-genome sequence of the resistant parent WRC17 was obtained by Illumina sequencing. RNA-seq of WRC17 leaves inoculated with *M. oryzae* was also performed. Next, short reads of whole-genome sequences of the susceptible parent Hitomebore and the six susceptible RILs were obtained by Illumina sequencing. (C) *De novo* assembly of R-parental line WRC17 was performed using *DISCOVAR* software. (D) Expressed genes of WRC17 were identified by mapping RNA-seq reads to the WRC17 reference genome. (E) Mapping of short reads of the susceptible parent (Hitomebore) and susceptible RILs to the predicted genes identified DNA polymorphisms shared by the susceptible lines using the RaIDeN pipeline (the script of the RaIDeN pipeline and details about the pipeline are available at https://github.com/YuSugihara/RaIDeN).

Suppl. Table 1. Summary of Illumina sequencing results used for RaIDeN methods

Line	Sequencing platform	Trim adapters and quality filter	Total no. of paired-end (PE) reads after filtering	Max length of short reads	Purpose
<u>Whole genome sequencing</u>					
R-parental line					
WRC17	Miseq	No	36,047,742	250	for *de novo* assemble
S-parental line					
Hitomebore	Hiseq4000	Yes	32,016,561	126	for RaIDeN
S-RILs					for RaIDeN
#22	Hiseq4000	Yes	30,830,739	126	for RaIDeN
#25	Hiseq4000	Yes	39,330,775	126	for RaIDeN
#34	Hiseq4000	Yes	36,109,189	126	for RaIDeN
#37	Hiseq4000	Yes	34,054,341	126	for RaIDeN
#39	Hiseq4000	Yes	42,282,097	126	for RaIDeN
#53	Hiseq4000	Yes	34,013,397	126	for RaIDeN
<u>RNAseq</u>					
M. oryzae infected leaf of WRC17	Nextseq500	Yes	39,744,107	75	for RaIDeN

Suppl. Table 2. Summary metrics of genome assembly of the resistance rice line WRC17. Genome assembly was performed by *DISCOVAR* software.

No. of contigs (>1,000bp)	25,278
Contig N50 (kb)	55.1
Largest contig (kb)	459.5
Total size of the assembled genome (Mb)	356.2

Suppl. Table 3. Summary of 38 candidate resistance gene analogs (RGAs) expressed in WRC17 leaves and possibly responsible for WRC17 resistance against 2012-1 as predicted by RGAugury (Li et al. 2016). These were selected from the 853 candidate genes as identified by the RaIDeN method. From the 38 RGAs, we selected 18 genes belonging to the categories of CNL (Coiled-Coil, NBS and Leucine rich repeat protein genes: 11 genes) and NL (NBS and Leucine rich repeat protein genes: 7 genes) as the candidate NLRs conferring resistance to WRC17 against *M. oryzae* 2012-1.

	NBS encoding					TX	Other	RLP	RLK	TM-CC
NBS	CNL	TNL	CN	TN	NL					
3	11	0	1	0	7	0	0	1	12	3

CC, C: coiled-coil       TX: TIR-unknown domain

T: Toll/Interleukin-1 receptor    RLK: receptor like kinase

NBS, N: nucleotide-binding site    RLP: receptor like protein

L: leucine rich repeat       TM: transmembrane

Suppl. Fig. 2. Mapping of *NLR*s in the *PiW17-1* and *PiW17-2* loci that confer resistance against *M. oryzae* 2012-1 to WRC17. (A) Association between resistance/susceptible phenotypes and genotypes among the RILs (the primers used for genotyping are shown in Suppl. Table 9). *CNL-04* and *CNL-05*, *CNL-06* and *CNL-07*, and *CNL-08*, *NL-2* and *CNL-09* are located on the same contigs. (B) Fine maps of candidate NLR genes in *PiW17-1* and *PiW17-2*. Blue and yellow indicate WRC17- and Hitomebore-type genotypes, respectively. The 10 NLRs shown in the red rectangle were considered to be the candidates of *Pi-W17-1*.

C

#4 #6 #8   #1 #3 #5   CNL-06 CNL-07 CNL-08 CNL-09 NL-02 NL-04

CNL-04.RNAi **(*Pias-1*)**   ***CNL-05.RNAi*** **(*Pias-2*)**

D

Suppl. Fig. 2 (continued). Results of RNAi-mediated silencing of candidate *NLR* genes.
(C) RNAi-mediated silencing of eight candidate genes encoding proteins over 900 amino acids long was performed. Gene silencing of *CNL-04* and *CNL-05* made plants susceptible to *M. oryzae* 2012-1. (D) Gene expression levels of *CNL-04* and *CNL-05* in plants transformed with the RNAi constructs. Control (Con) is WRC17 transformed with empty vector.

A

B

	Sequence (sgRNA4.1)	Modification
WT	GCCGCCGTACAGACGCTATGCA–TGGAGGACCTC	n/a
cnl-04_1.18.3	GCCGCCGTACAGACGCTATGC‑‑TGGAGGACCTC	1bp deletion, premature stop 127aa
cnl-04_1.14.2	GCCGCCGTACAGACGCTATGCA–TGGAGGACCTC	n/a
	GCCGCCGTACAGACGCTA‑‑‑‑‑TGGAGGACCTC	4bp deletion, premature stop 126aa
cnl-04_1.21.2	GCCGCCGTACAGACGCTATGCACTGGAGGACCTC	1bp insertion, premature stop 162aa

C

Suppl. Fig. 3. **CRISPR/Cas9-mediated knockout of *CNL-04* abolishes *PiW17-1*-mediated resistance against *M. oryzae* isolate 2012-1.**
(A) The position of guide DNA targeted to *CNL-04*. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. (B) Sequences of the sgRNA positions in the *CNL-04* knockout lines. (C) Resistance/susceptible phenotypes of the selfed progeny of the *cnl-04_1.14.2* line heterozygous for wild-type (*a*) and mutated (*b*: 4-bp deletion) alleles. The *b/b* homozygous plants became susceptible.

D

CNL-05

ATG
127bp
216bp
404bp

CC = CC
NB-ARC
LRR
DUF761

CCAAGCCACAGGGCAAGATACTGGATGGATGAG
sgRNA5.3

CCAAGCCACAGGGCAAGATACTGGATGGATGAG
sgRNA5.2

CTGAGCCACTGAAGGACGGAATCGAGCTCCTCA
sgRNA5.1

E

	Sequence (sgRNA5.1)	Modification
WT	CTGAGCCACTGAAGGACGGAATCGAGCTCCTCA	n/a
cnl-05_1.6.2	CTGAGCCACTG-AGGACGGAATCGAGCTCCTCA	1bp deletion, premature stop 45aa

	Sequence (sgRNA5.2)	Modification
WT	CCAAGCCACAGGGCAAGATACTGGATGGATGAG	n/a
cnl-05_2.4.2	CCAAGCCACA-GGCAAGATACTGGATGGATGAG	1bp deletion, premature stop 109aa

	Sequence (sgRNA5.3)	Modification
WT	GCTCGCCAGGA-TCGCGGAGCTCAGGGCTCTGGT	n/a
cnl-05_3.7.2	GCTCGCCAGGA-TCGCGGAGCTCAGGGCTCTGGT	n/a
	GCTCGCCAGGACTCGCGGAGCTCAGGGCTCTGGT	1bp insertion, premature stop 148aa

F

#1 #2 #3 #4 #5 #6 #7 #8

Phenotype S R R R S S S R

Genotype *b/b* *a/b* *a/b* *a/b* *b/b* *b/b* *b/b* *a/a*

a: Wild type allele
b: Mutated type (+1bp) allele

Suppl. Fig. 3 (**continued**). **CRISPR/Cas9-mediated knockout of** *CNL-05* **abolishes** *PiW17-1* **mediated resistance against** *M. oryzae* **isolate 2012-1.**
(D) The positions of guide DNAs targeted to *CNL-05*. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. (E) Sequences of the sgRNA positions in the *CNL-05* knockout lines. (F) Resistance/susceptible phenotypes of the selfed progeny of *cnl-05_3.7.2* line heterozygous for wild-type (*a*) and mutated (*b*: 1-bp insertion) alleles. The *b/b* homozygous plants became susceptible.

Suppl. Fig. 4. Comparison of the gene structures of *Pia* (*RGA4* and *RGA5*) and *Pias* (*Pias-1* and *Pias-2*). DNA sequence similarities obtained by ClustalW are shown.

A

B

Pias-1

RX-CC
```
MEAALLSGFIKAILPRLFSLVNDKLNLHKGVKGDIDFLIKELRMIVGAIDDDLSVEHGAAAAAAV
QTLCMEDLRELAHGIEDCIDGVLYRAAREQRRSSSLLPRTVRATKKLLQTNQHLAQELQRLKRMVE
E
```
```
ANQRKQRYTAAAPGQHGQVYSSAAAQVDEPWPSCSSASDPRIHEADLVG
```

NB-ARC
```
VDADRAELLEQLAERQPEQLKVIAIVGFCGLGKTALAAEAYNRETRGGRFERHAWVCAAHRSAREV
LGELLRRIDAACHGDSDAGQLCVDIRQQLEKKRYFIVIDDIQTEDQWKSIKSAFPTDKDIGSRIVV
TTTIQSVANACCSANGYLHKMSRLDKNCSKQLLSKKACPERYSHYKQPDSAAILKKCDGQPLALVT
IGEFLQANGWPTGPNCEDLCNRLHYHLENDKTLERMRRVLVRNYTSLPGHALKACLLYFGMFPSDH
PIRRKSLLRRWLAEGFVEPV
```
```
SSSSNLDSTAAFDVLMDRNIIEPINVSNNDKVKTCQTYGMMREFISHMSISQNFVTFFCDDKFLPK
YVRRLSLHGDTVVNGDNFNGIDLSLVRSLAVFGEAGTTVLDFSKYQLLRVLDLEKCDDLNDDHLKE
ICNLVL
```

LRR
```
LKYLSLGGNISKLPKDIAKLKD
LEALDVRRSKVKIMPVEVFGLPC
LIHLLGKFKLSDKVKQKTEVQEFLLKGKSN
LQTLAGFASNGSEGFLHLMRYMNKLRKLKIWCTSSAGSTDWTDLREA
IQQFILDEKEANIGTRSLSLHFTGCSEDAINSLKEPCY
LSSLKLHGNFPQLPQFVTSLRG
LKELCLSSTKFTTGLLEALSNLSYLQYLKLVADELEKFIIKVQGFPRLLR
LCIVLQCPTFPVIEEGALPFLVT
LQLLCKDLHGLSDIQIECFKH
```
```
LQEVTLHSGVTPATRQEWVKAAKEHPNRPKVLLLKSVDTAESEHTDVDSVMEAVKSETTEYYIAPE
GPEQVIDMNNKMQLDHGLESSSVLNKQNNFADQSSSKDQLHYSFNNMGLSDVSPAVSELPNGMVPS
CT
```

Pias-2

RX-CC
```
MDAPVSFSLGAMGPLLRKLDSLPVAPEIRLPEPLKDGIELLKEDLEEIGAAVVEQSTVDSPSHRAR
YWMDEVRDLSYHIEDCIDTMFSMRCGGDDGKPRSVRR
```
```
HKVGRVKVDGFSKTQKPCTRLARIAELRALVREASERHERYQLGDGRASSSSSSSSHRVFTAHGQVP
APCRNLVGM
```

NB-ARC
```
DEPKTKLTNMLTDEAELHMKVVCILGSAGIGKTTLAEQVYRKLRWQFDCHAFVRASRKPDMRRLLG
AILSQVQPRIRISDTSTVQSLIDNLWEYLQKKRYFIVIDELYETATWDIITSAFPEDNNCSRIMTT
AGIEGVALECCSYHSVNIFKMKPLGLDDSAKLFFNRVFGSEQQCPYELSEVSYRITAKCGGLPLAV
IIIAGLLASLPCKTELWYNIDGCLCSSVTTDIDLDEILKEIISLGYDNLPHYLKTCLLYLSLYSEG
FIIWTADLLKQWISEGFIAVIDG
```
```
EDIEEVAESYFYNLVNRGMIQSVKTKYNNQVLCTVHHTVFDLIIHKSKEEKFISAIDYSQTMPGNS
LEARRLSFHFSNTRYATEVAGITLSQVRSFAFLGLLKCMPSIMEFKL
```

LRR
```
LRVLILEFWGDNHGCMSFNVARICRLFQ
LRYLKISSQIIIELPAQIRGLKY
LETLEIDARVTAVPSDIIHLRS
LLHLYFQDGIVLPDGIGCIRSLRT
LKYFDLGSNSEENIRSLGQLTNLRDLHLTCSAPKSNQQAKRNLVILASYTGKLGN
LKSVKFSPGDSGMDISFLFYGIGISVDRSRTASS
LPVSVRTLELPSICIFARLPDWIGQ
LRKLHTLNLAVRELIENDIDS
LAGLPDLIVLSMHIMKAPMERIVFNRKAFPV
LKYFKFICGTLRMAFQAGAMANLHR
LKLGFNAH
```
```
KGEKYDNILVGIEHLLNLKKIAVRIGGAAEEAKESDRMAAEAAALKEAIRKHLMFLDDLDIARVECVK
EEYKCIKKKHKIKIEDSISEKNGDSKKQHSVEKKAVWGKTMKNIADSGVFPEDYTMSREQRVAEGF
VVGIEKCRAEDAAERIIRNVPVDYDGLGQVSTSKIQDHLPELAPRAVQNEKFGSSNDLSIMIQINK
YARLPSYEWRDTDISKLNFRLLRAPMLLEAVTARCHLLDLILIGSNN
```

LRR
```
ITVLDLGRPTITKLPASIECLPN
LRYLRLQGTQLKSLSEVIVKMPT
IRGLDIKNTKTEELPQGILRMKK
```
```
LSHLSMGEKQKNIQVFMEKMQTLAETVQDSDDLSDETEGIADDDEGEFSTRANASTPKVD
```

DUF761
```
EDEVDRRANNFIARFRKQI
```
```
TIRNSGFPKRRAASTNDCGHEISMSANSPREVDDFKKKFDEIIARNRHTWKPIESPRSVKHGKYFV
RCPPSLITSAVFSFFFLSIFLVWYSLRFRILSGKP
```

Suppl. Fig. 5. Gene structures and predicted amino acid sequences of *Pias-1* and *Pias-2*.
(A) Gene structures of *Pias-1* and *Pias-2*. (B) Amino acid sequences of Pias-1 and Pias-2 with the putative domains.

Suppl. Fig. 6. Isolation of *AVR-Pias* from *M. oryzae* isolate 2012-1.
A simplified scheme showing the procedure used to isolate *AVR-Pias*. First, the genome of isolate 2012-1 was sequenced on the Illumina platform, and the short reads were used for *de novo* assembly with *DISCOVAR*, resulting in the 2012-1 genome reference (**Suppl. Table 4, 5**). Next, the isolate 2012-1 was used to inoculate barley (*Hordeum vulgare*) cultivar 'Nigrate', which is highly susceptible to *M. oryzae* (Hyon et al. 2012), and the infected leaves were subjected to RNA-seq (Shimizu et al. 2019), resulting in the identification of 10,991 genes in 2012-1 that were expressed during host infection. Next, eight *M. oryzae* isolates (r1–r8) were selected and used for inoculation of rice line HW-RIL7 with *Pias* but without *Pi-W17-2*. The eight isolates were compatible with HW-RIL7 (result shown in the bottom), suggesting that they lack *AVR-Pias*. The eight isolates were then subjected to genome resequencing on the Illumina platform, and the RaIDeN method was used to identify only presence/absence polymorphisms (**Suppl. Table 4**). The short reads of eight isolates were aligned to the expressed genes of the 2012-1 isolate, resulting in the identification of 87 expressed genes that were specific to isolate 2012-1. These transcripts were further filtered based on four criteria: (1) genes showing a higher level of expression 24 h after inoculation (FPKM > 200) and genes encoding (2) a putative secreted protein, (3) a non-transmembrane protein and (4) a small protein (<150 amino acids). This analysis identified three transcripts (*G9141*, *G9435*, and *G9532*) as the candidates of *AVR-Pias*.

Suppl. Table 4. Summary of whole genome sequencing of the nine *M. oryzae* isolates and RNA sequencing of *M. oryzae* isolate 2012 for identification of *AVR-Pias*

Line	Race code	Sequencing platform	Trim adapters and quality filter	Total no. of paired-end (PE) reads after filtering	Max length of short reads	Purpose
Whole genome sequencing						
2012-1		Miseq	No	3,332,000	250	for *de novo* assemble
S-pathogen for HW-RIL7						
Ina87T-56A	**r1**	Miseq	Yes	1,776,902	125	for RaIDeN
Ao92-06-2	**r2**	Miseq	Yes	2,691,934	125	for RaIDeN
85-141	**r3**	Miseq	Yes	3,034,450	125	for RaIDeN
SL91-48D	**r4**	Miseq	Yes	3,108,538	125	for RaIDeN
2403-1	**r5**	Miseq	Yes	3,388,670	125	for RaIDeN
H98-315-1	**r6**	Miseq	Yes	2,933,673	125	for RaIDeN
0423-1	**r7**	Miseq	Yes	2,996,542	125	for RaIDeN
Ina85-182	**r8**	Miseq	Yes	3,148,657	125	for RaIDeN
RNAseq						
M. oryzae isolate 2012-1 inoculated to the leaf of *Hordeum vulgare* var. Nigrate				**Total no. of reads (SE) after filtering**		
24H after inoculation		Nextseq500	Yes	15,658,123	75	for RaIDeN and expression analysis
48H after inoculation		Nextseq500	Yes	15,770,613	75	for RaIDeN and expression analysis

Suppl. Table 5. Summary metrics of genome assembly of *M. oryzae* 2012-1 isolate. Genome assembly was performed by *DISCOVAR* software.

No. of contigs (>500bp)	2,414
Contig N50 (kb)	154.7
Largest contig (Mb)	1.0
Total size of the assembled genome (Mb)	41.0

A

M. oryzae: Ao92-06-2

+*pex22p*: -   *G9141*   *G9435*   ***G9532***

WT   #2   #4   #9   #11   #2   #3   #4   #2   #3   #4

Rice   HW-RIL7 (*Pias⁺*)   Hitomebore (*Pias⁻*)

B

Ao92-06-2

WT   +*pex22p:G9532*

	S	R	S	S
Pias-1		+	-	+
Pias-2		+	+	-

Suppl. Fig. 7. The expressed gene *G9532* of *M. oryzae* isolate 2012-1 is *AVR-Pias*.
(A) Results of inoculation of rice line HW-RIL7 with *Pias* or Hitomebore without *Pias* with *M. oryzae* isolate Ao92-06-2 wild type (WT) or Ao92-06-2 with the *G9141*, *G9435* or *G9532* transgene, all driven by the *pex22* promoter. When Ao92-06-2 contained *G9532*, the interaction became incompatible, indicating that *G9532* is *AVR-Pias*. (B) Introduction of *pex22p:G9532* into *M. oryzae* isolate Ao92-06-2 confers avirulence to the pathogen against HW-RIL7 with *Pias-1* and *Pias-2*. Knockout of the host gene *Pias-1* or *Pias-2* compromises the avirulence caused by *G9532*.

A

B

Suppl. Fig. 8. Deletion of *AVR-Pias* from *M. oryzae* isolate 2012-1 causes a loss of avirulence against rice line HW-RIL7 with *Pias*.
(A) Schematic overview of the target gene replacement (TGR) strategy at the *AVR-Pias* locus using RNA-guided nuclease. The PAM is marked with blue letters, and the sgRNA sequence is marked with red letters. The primers used for PCR analysis are indicated by horizontal arrows. (B) Inoculation assay of rice line HW-RIL7 with *M. oryzae* 2012-1 wild type (WT) and TGR transformants. (C) PCR analysis of TGR events at the *AVR-Pias* locus. Upper and lower images show PCR results using *AVR-Pias*- and *Hygromycin* (*Hyg*)-specific primers, respectively. pCB1636: Replacement vector containing the hygromycin resistance gene and genomic regions neighboring *AVR-Pias*. The isolates corresponding to lanes 2–5 show compatibility with rice line HW-RIL7 with *Pias*, whereas the wild-type isolate (2012-1) in lane 6 is incompatible with HW-RIL7.

Figure 2. Recurrent integration of extraneous domains in Pias/Pia sensor NLRs.
(A) A simplified scheme of the structures of the Pias/Pia NLR pairs. Pias-1/RGA4 helper NLRs are shown in green, and Pias-2/RGA5 sensor NLRs are shown in white. The conserved LRR-ID intervening motif (LII) is indicated by black ellipses. Integrated domains are shown by different-colored hexagons. (B) A sequence logo showing conserved amino acids of the LRR-ID Intervening (LII) motif. (C) Distribution of ID motifs among *Oryza* species. The pie charts show the frequencies of different ID motifs in a given species. The colors correspond to the ID colors in (A). The numbers below the pie charts indicate the sample numbers. A cladogram showing the phylogenetic relationships of 11 *Oryza* species and four other Poaceae species (*Setaria italica*, *Panicum hallii*, *Hordeum vulgare*, and *Aegilops tauschii*) based on Time Tree, the time scale of life web database (http://www.timetree.org/). The numbers on the branches indicate the estimated time of the splitting of lineages (MYA: million years ago). (D) DNA sequence similarity between *O. punctata Pias-2/RGA5 sensor NLR* and the downstream sequence of *O. sativa* (Nipponbare) *Pias-2/RGA5*. LII: LRR-ID intervening motif. (E) Dot-plot analysis of the *O. punctata* (W1582) *Pias-2/RGA5 sensor NLR* and *O. sativa* (Nipponbare) *Pias-2/RGA5 NLR* downstream sequences using the Dotmatcher tool (http://emboss.bioinformatics.nl/cgi-bin/emboss/dotmatcher/). (F) Possible evolutionary process of ID replacement that might have occurred between the *O. punctata* and the *O. sativa* of *Pias-2/RGA5* lineages. We still do not know the mode of interaction between the AVR-Pias effector and DUF761-containing protein, so it is indicated by "?".

The Sequence Manipulation Suite: Multiple Align Show

```
RGA5 (1116 aa)         A          -                          - - - -          K      E       - - - -          V S - - -      - - H  V          T  K      66
Pias-2 (1368 aa)                   -                          P  - - - -       E      D       - - - -          A A - - - V      - - Q          S  R      66
LOC_Os12g18360 (1033)  A P A   A Q  V I  R S  T S      L Q P P E P  P P A Q  S S  R K  E R K K  L    R G    R H L L D D Y Y L      P P S D  A P P  D S T  A    80

RGA5 (1116 aa)         F                          - - -  S            E          - -  I          K P          M                          L      141
Pias-2 (1368 aa)       Y                          - - -  C            V          - - V  V        T Q          L                          H      141
LOC_Os12g18360 (1033)  C  A K      E      D V D  F L  E L T T Q L L H H R    G    S S T A G A K K M I S S M I   R L R    E L N R R R W I A - - D E V T L F S  R K      I R  Q    158

RGA5 (1116 aa)              - - - - - - - - - - - - - - -      - - - - V C        P V        D    R A R P L H H G  S A          V  F          N      S  E G            A      203
Pias-2 (1368 aa)           - - - - - - - - - - - - - - -      G  A S  S S      H          H  Q V P - - - -    P C R          M  P          T N M    T      A L  M      V C  L    203
LOC_Os12g18360 (1033)  S  H    R R T S S S R P R E E V D D   D  E D  A  N E R R  F L S L T F G M D - - D A    H G Q        R  I S M Q    V      A  - G    K          S      235

RGA5 (1116 aa)         P          A      L D      - - - -  E                          Q    G          R - - Q          A Y A D S                          277
Pias-2 (1368 aa)                   E Q V      K L      - - - -      H                    R      A          - -  I  I      T S - - -                          W  Y      274
LOC_Os12g18360 (1033)      G  V          F      L  G R R L D A P          T P        T K I  T D M      L R  Q H Q H Q      V W - - - E  D R  L E T I    T      312

RGA5 (1116 aa)         R      L      G          V  N A N          V  F          D  Q            K Y D Y  M R  E        S L      K  V      K          D      357
Pias-2 (1368 aa)       K          V    E Y      T      I T      E D  N      M T    G  G          S H  V N  F  K      L      A          R      Q      354
LOC_Os12g18360 (1033)          E D      A S S M      V S R G L    N              T E    P      A      N  E H  I  I D      D  V  S Q        S G  V  Q G N E F    392

RGA5 (1116 aa)         P    K      N T  L E              S        G  - Q  E N P V    D Y V T  Y          -  G          K D V V      T      S  S      P F      435
Pias-2 (1368 aa)       Y    S      Y R  T A              V          A  -  C K T E    Y N  D G C          - V T  D I D  D E I L      I I S  G  D        Y      432
LOC_Os12g18360 (1033)  G H  T      H D M I K          T  T  R H F K  Q  L D G M Q Q    N H  Q  S  T T  N  K K          Q G - M R Q V      I          C      A    471

RGA5 (1116 aa)         L L Y L G M Y P D G H I M L K A D L M K Q W S A E G F V S A N E A K D T E E I V D K Y F D E L V N R G I L E P V E I N K N G K V L S C T L H H A V H D L V M P K    515
Pias-2 (1368 aa)       L L Y L S L Y S E G F I I W T A D L L K Q W I S E G F I A V I D G E D I E E V A E S Y F N L V N R G M I Q S V K T K Y N N Q V L - C T V H H T V F D L I I H K    511
LOC_Os12g18360 (1033)  L L Y L S I Y K E D Y I I R K A N L V R Q W M A E G F I N S I E N K V M E E V A G N Y F D E L V G R G L V Q P V D V N C K N E V L S C V V H H M V L N F I R C K    551

RGA5 (1116 aa)         F N D D K F T M S V D Y S Q T I T G P S T M V R R L S L H F S S T R Y A T K P A G I I L S R V R S L A F F G L L N C M P C I G E F K L L R V L I L E F W G S H G    595
Pias-2 (1368 aa)       S K E E K F I S A I D Y S Q T M P G N S L E A R R L S F H F S N T R Y A T E V A G I T L S Q V R S F A F L G L L K C M P S I M E F K L L R V L I L E F W G D N H    591
LOC_Os12g18360 (1033)  S I E E N F S I T L D H S Q T T V R H A D K V R R L S L H F S N A H D T T P L A G L R L S Q V R S M A F F G Q V K C M P S I A D Y R L L R V L I L C F W A D Q -    630

RGA5 (1116 aa)         E Q R S L N L I P V C R L F Q L R Y L K T S G D V V V Q L P A Q I S G L Q Y L E T L E I D A R V S A V P F D L V H L P N L L H L Q L Q D E T K L P D G I G C M R    675
Pias-2 (1368 aa)       G C M S F N V A R I C R L F Q L R Y L K I S S Q I I E L P A Q I R G L K Y L E T L E I D A R V T A V P S D I I H L R S L L H L Y F Q D G I V L P D G I G C I R    671
LOC_Os12g18360 (1033)  E K T S Y D L T S I S E L L Q L R Y L K I T G N I T V K L P E K I Q G L Q H L Q T L E A D A R A T A V L L D I V H T Q C L L H L R L V L L D L L P H - - - C H R    707

RGA5 (1116 aa)         S L R T L Q Y F D L G N N S V D N L R G L G E L T N L Q D L H L S Y S A P S S N E G L M I N L N A I T S S L S R L S N L K S L I L S P G A I S M V I F F - - - -    751
Pias-2 (1368 aa)       S L R T L K Y F D L G S N S E E N I R S L G Q L T N L R D L H L T C S A P K S N Q Q A K R N L V I L A S Y T G K L G N L K S V K F S P G D S G M D I S F L F Y G    751
LOC_Os12g18360 (1033)  - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -    707

RGA5 (1116 aa)         - - - - - D I S S I I S V V P V F L Q R L E L L P P I C I F C R L P K S I G Q L H K L C I L K V S V R E L L T T D I D N L T G L P S L T V L S L Y A Q T A P E G    826
Pias-2 (1368 aa)       I G I S V D R S R T A S S L P V S V R T L E - L P S I C I F A R L P D W I G Q L R K L H T L N L A V R E L I E N D I D S L A G L P D L I V L S M H I M K A P M E    830
LOC_Os12g18360 (1033)  - - - - - - - - - - - - - - - - - - - - - - Y I F T S I P K W T G K L N N L R I L N I A V M Q I S Q D D L D T L K G L G S L T A L S L L V R T A P A Q    760

RGA5 (1116 aa)         R F I F K D G T L P V L K Y F K F G C G E L C L A F M A G A M P N L Q R L K L V F N I R K S E K Y R H T L F G I E H L V S L Q D I A T R I G V D T S T G E S D R    906
Pias-2 (1368 aa)       R I V F N R K A F P V L K Y F K F I C G T L R M A F Q A G A M A N L H R L K L G F N A H K G E K Y D N I L V G I E H L L N L K K I A V R I G G A A E A K E S D R    910
LOC_Os12g18360 (1033)  R I V A A N E G F G S L K Y F M F V C T A P C M T F V E G A M P S V Q R L N L R F N A N E F K Q Y D S K E T G L E H L V A L A E I S A R I G G - T D D D E S N K    839

RGA5 (1116 aa)         R A A E S A F K E T V N K H P -                          R C L R S S L Q W V V - - S T E E E S H P L E K Q H H - - - - - - - -    946
Pias-2 (1368 aa)       M A A E A A L K E A I R K H L M F L D D D L D I A R V E C V K E E Y K C I K K K H K I K I E D S I S E K N G D S K K Q H S V E K K A V W G K T M K N I A D S G V F    990
LOC_Os12g18360 (1033)  T E V E S A L R T A I R K H P T P - - - - - - - - - - - - - - - S T L M V D I Q W - V D W I F G A E G R D L D E D L A - - - - - - - - - - - - - - - -    882

RGA5 (1116 aa)         - - - - - K R E K G S S A G H G V - L E K E S V E D S E K - - - - - - - - - - - - - - - N T D R V Q T L L S - - - - - - - - - -    979
Pias-2 (1368 aa)       P E D Y T M S R E Q R V A E G F V V G I E K C R A E D A A E R I I R N V P V D Y D G L G Q V S T S K I Q D H L P E L A P R A V Q N E K F G S S N D L S I M I Q I    1070
LOC_Os12g18360 (1033)  - - - - - - - - - Q Q D D H G F C M L P E S S R L Q S R E - - - - - - - - - - - - - - - I S I E E D M P - - - - - - - - - - - - -    911

RGA5 (1116 aa)         - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - P Q L S N M E S V V E S - - - - - - - A L T G    995
Pias-2 (1368 aa)       N K Y A R L P S Y E W R D T D I S K L N F R L L R A P M L L E A V T A R C H L L D L I L I G S N N I T V L D L G R P T I T K L P A S I E C L P N L R Y L R L Q G    1230
LOC_Os12g18360 (1033)  - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - T G A E G T V I V P H T M E Q F A L H M    931

RGA5 (1116 aa)         Q R - - - - T K I V V K V - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - H M P C G K S R - - - - - - - A K A M A L A A S V N G V D S V E - - - - - I T G    1032
Pias-2 (1368 aa)       T Q L K S L S E V I V K M P T I R G L D I K N T K T E E L P Q G I L R M K K L S H L S M G E K Q K N I Q V F M E K M Q T L A E T V Q D S D D L S D E T E G I A D    1230
LOC_Os12g18360 (1033)  S Q A K Q S H K L V V I Q - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - F T T S R C P A S R Y I A P A F T E Y A K E F A G A V F I K V N V - - - - - - - - -    977

RGA5 (1116 aa)         E D K D R L V V V G R G I D P V - - - - - - - - - R L V A L L R E K - - - - - - - - - - - C G L A E L L M - - - - - - - V E L V E K E K T Q L    1076
Pias-2 (1368 aa)       D D E G E F S T R A N A S T P K V D E D E V D R R A N N F I A R F R K Q I T I R N S G F P K R R A A S T N D C G - H E I S M S A N S P R E V D D F K K K F D E I    1309
LOC_Os12g18360 (1033)  - D S D E L E S V T D W Y D I E - - - - - - - - - G I V P T F - - - - - - - - - - - - - - - - - - - F F - - - - - - - V K D G E K - I D K I    1010

RGA5 (1116 aa)         A G G K K G A Y K - - - - - - - - - K H P T Y N L S P F D Y V E Y P P S A P I M Q D I N P C S T M - - - - - - - - - -    1116
Pias-2 (1368 aa)       I A R N R H T W K P I E S P R S V K H G K Y - - - - - - F V R C P P S L - I T S A V F S F F F L S I F L V W Y S L R F R I L S G K P    1368
LOC_Os12g18360 (1033)  P G A N K E L L R - - - - - - - - - - - - - - - - - - A K I R R H T A S P Y F L R - - - - - - - - - - - - - - - -    1033
```

☐ LII motif (aqua box)
☐ CID motif (orange box)

Suppl. Fig. 9. Positions of the LII motif (this study) and the CID motif (Bailey et al. 2018).
Amino acid sequence alignment of RGA5, Pias-2 and LOC_Os12g18360 and the positions of the LII motif (this study) and the CID motif as described by Bailey et al. (2018). The aqua and orange boxes indicate the LII and CID motif regions, respectively.

Suppl. Fig. 10. The *O. punctata* W1582 LII-ID sequence is conserved in the downstream sequence of *Pias-2* in *O. sativa* cv. Nipponbare (NP).
Alignment of the DNA sequences of the *O. punctata* W1582 LII-ID and the W1582 LII-ID like sequence in *O. sativa* cv. Nipponbare. The blue boxes and green lines indicate the exon sequence of *O. punctata* W1582 Pias-2 homolog and the LII region, respectively.

Suppl. Fig. 10. Continued.

Suppl. Fig. 11. The LII + ID sequence of the *O. punctata* (W1582) *Pias-2* homolog is conserved in the downstream sequence of *O. meridionalis* (W2112) Pias-2.
Dot-plot analysis of *O. punctata* (W1582) *Pias-2* homolog and the genome sequence of *O. meridionalis* (W2112). The purple line corresponds to block (a) in Figure 2. The block (b) sequence in Figure 2 is deleted in *O. meridionalis* (W2112).

Suppl. Table 8. A survey of 51 A-genome *Oryza* accessions with DUF761 ID revealed that the *O. punctata* LII-ID-like downstream sequence is widely conserved.
BLASTN search was performed on each genome sequence using *O. punctata* W1582 LII-ID-like sequence of Nipponbare cv. as a query. Block (a) and block (b) region correspond to those in Figure 2.DE.

Species	with DUF761 as ID	*O. punctata* W1582 LII-ID-like sequence conserved	
		block (a) and (b)	block (a) only
O. sativa	19	19	0
subsp. *indica*	4	4	0
subsp. *japonica*	15	15	0
O. rufipogon	4	4	0
O. barthii	8	1	0
O. glumipatula	11	0	0
O. meridionalis	9	1	7

Figure 3. Contrasting evolutionary patterns of the helper and sensor NLRs of the *Pias/Pia* locus.
(A) Phylogenetic tree of the *Pias-1/RGA4 helper NLR* gene (left) and *Pias-2/RGA5 sensor NLR* (right) gene based on the full-length amino acid sequence of Pias-1/RGA4 and the sequence in the region CC to LII for Pias-2/RGA5. *Pias-2/RGA5* sensor NLRs form two major clades (C1 and C2). The numbers indicate bootstrap values. (B) Pairwise comparison of ω, the ratio of nonsynonymous to synonymous nucleotide substitutions for *Pias-1/RGA4* (above the diagonal: the full-length NLR DNA sequence was used) and *Pias-2/RGA5* (below the diagonal: the sequence of the CC to LII region was used).

Suppl. Fig. 12 HR-like cell death after overexpression of helper NLRs of the Pias-1/RGA4 lineage in *N. benthamiana.*

(A) Box plots of autofluorescence values after transient expression of Pias-1/RGA4 homologs. GUS vector was included as a control. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Representative image of Pias-1/RGA4-meditated HR *in N. benthamiana*. (C) Immunoblot analysis of Pias-1:HA, RGA4:HA, RGA4-Oau:HA (cloned from *O. australiensis* accession W0008), RGA4-Oru:HA (cloned from *O. rufipogon* accession W1943), and RGA4-Ogr:HA (cloned from *O. granulata* accession W0067B) proteins detected by anti-HA (*a*-HA) antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size marker are indicated on the left (kDa).

Suppl. Fig. 13 HR-like cell death after overexpression of Pias-1:HA helper and Pias-2/RGA5 sensors in *N. benthamiana*.
(A) Box plots of autofluorescence values after transient expression of Pias-1:HA, FLAG:Pias-2 and FLAG:RGA5 separately or in combination in *N. benthamiana*. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Representative image of HR after transient expression of Pias-1:HA, FLAG:Pias-2, and FLAG:RGA5 separately or in combination in *N. benthamiana*. (C) Immunoblot analysis of Pias-1:HA protein detected by anti-HA (α-HA) antibody and FLAG:Pias-2 and FLAG:RGA5 proteins detected by anti-FLAG antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size marker are indicated on the left (kDa).

A

B

Figure 4. The NLR helper Pias-1 is functionally conserved.

(A) Representative images of *N. benthamiana* leaves after agroinfiltration with Pias-1:HA, Pias-1:HA/FLAG:RGA5, FLAG:RGA5/AVR-Pia, and Pias-1:HA/FLAG:RGA5/AVR-Pia (left) and RGA4-Ogr:HA derived from *O. granulata*, RGA4-Ogr:HA/FLAG:RGA5, FLAG:RGA5/AVR-Pia, and RGA4-Ogr:HA/FLAG:RGA5/AVR-Pia (right). Autofluorescence under UV light is shown. (B) Pias-1 cooperates with RGA5 to recognize AVR-Pia and induces resistance in rice. The rice line HW-RIL7 with *Pias* (*Pias-1* and *Pias-2*) recognizes the Ao-92-06-2 strain with *AVR-Pias* (Ao92-06-2+*pex22p:AVR-Pias*) and induces resistance. However, HW-RIL7 cannot recognize the Ao02-06-2 strain with *AVR-Pia* (Ao92-06-2+*pex22p:AVR-Pia*). Two lines (Line #1 and #2) contain the *35S-RGA5* transgene in the HW-RIL7 background. F1-A is a progeny derived from a cross between Sasanishiki with *Pia* (*RGA4* and *RGA5*) and HW-RIL7. F1-B is a progeny derived from a cross between a Sasanishiki mutant (Sas1493) with *pia* (*rga4* and *RGA5*) and HW-RIL7.

Suppl. Fig. 14. HR-like cell death caused by Pias1 and RGA4-Ogr expression is suppressed by RGA5 expression, and additional AVR-Pia expression induces cell death.
(A) Box plots of autofluorescence values after transient expression of Pias-1:HA, FLAG:RGA5 and AVR-Pia separately or in combination in *N. benthamiana* leaves. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (B) Immunoblot analysis of Pias-1:HA protein detected by anti-HA (α-HA) antibody, FLAG:RGA5 protein detected by anti-FLAG antibody and AVR-Pia detected by anti-AVR-Pia antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size markers are indicated on the left (kDa). (C) Box plots of autofluorescence values after transient expression of RGA4-Ogr:HA, FLAG:RGA5, and AVR-Pia separately or in combination in *N. benthamiana* leaves. The number of spots inoculated with *Agrobacterium* are indicated at the top of the boxplot. (D) Immunoblot analysis of RGA4-Ogr:HA protein detected by anti-HA (α-HA) antibody, FLAG:RGA5 protein detected by anti-FLAG (α-FLAG) antibody and AVR-Pia detected by anti-AVR-Pia (α-AVR-Pia) antibody. Coomassie blue staining of Rubisco small subunit shows equal protein loading. The protein bands expressed from the constructs are marked by red asterisks. The positions of molecular size markers are indicated on the left (kDa).

Host: Rice

M. oryzae	HW-RIL7 (Pias-1, Pias-2)	P	Sasanishiki (RGA4, RGA5)	P	Sas1493 (rga4, RGA5)	P
Ao92-06-2 (WT)		S		S	-	
+ pex22p::AVR-Pia		S		R		S
+ pex22p:: AVR-Pias		R		S	-	

Suppl. Fig. 15. Results of inoculation of HW-RIL7, Sasanishiki and Sas1493 rice plants with *M. oryzae* isolates Ao92-06-2, Ao92-06-2 + pex22p:AVR-Pia and Ao92-06-2 + pex22p:AVR-Pias.

Suppl. Fig. 16. The DUF761-containing gene family in rice.
A BLASTP search was performed using the amino acid sequence of the DUF761 domain of Pias-2 as a query.
Fifteen DUF761 domain-containing genes were retrieved using a cutoff e-value < 10. The middle panel shows
the amino acid sequence alignment and the right panel shows the domain structure of each protein.

Suppl. Table 9. List of primers used in this study.

List of primers used for genotyping of RILs

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Genotyping method
CNL-01	GCAGGTTCCCGCACAAACTATGG	GGTTGCCACGGGTATTTGTAATG	CAPS (MspI)
CNL-02	GGTTGATGTTGTGGAGTGAGAC	TAGCCTGGGCCCTGAGGAGATG	CAPS (MspI)
CNL-03	CAAAACATCAACAATTATGGAACC	TTGAAGCTCTCATATAGAAGTGTCC	CAPS (EcoRI)
CNL-04	TATTGCACCAGAGGGACCAGAAC	GAATCCACCTGATAGGGAATTCG	CAPS (MspI)
CNL-06	CGTGAAAGTTGCCAAATCTATAG	GTTCGAAGACGTAGAAGTTTGAG	CAPS (SacI)
CNL-09	CTACAGAAACTGCAAGTGTCCAC	CATAATTATTTGCAGAGCCGAAC	P/A
CNL-10	TTGCTAGTTGTACGGCTTCGATG	AACCCCATGGTTATGAACAGGAAC	CAPS (MspI)
CNL-11	GCTAAGACCTCTCATCATCATGG	GCGGATGAACTCTGTAACCATAA	CAPS (XhoI)
NL-01	GGGGATGGAGAATAACAGACAAC	GCCTTTTTACTGAACAGGAGGAG	CAPS (SacI)
NL-02	GTGAAACAAATCTTCACCAAAGC	ATCTTTCCTTTATGGCACTTTTC	CAPS (XhoI)
NL-03	GATGTCATCCCTCTGCAAACTTG	TCCTTCGGTGGATCATTTTCATC	CAPS (NsiI)
NL-04	CAAGATTCGCAGGCTCTCAATAC	CCTGCAGAGCTGCCATGTTAGC	P/A
NL-05	TATAAGACAGCTCCAATCCATGAC	TCGGAATTCATGAAAGAATCTC	P/A
NL-06	AGACGGTGGAGATGACAAGGTG	CTGTGATGGACCAAAGCCTGTTG	P/A
NL-07	ACTTCCTCGACGAGCTAACGAC	AAGACTGATGCTGATGTTGTGG	CAPS (SalI)
W17.11718	TAATGTTGCAGGTGCTAAGAGAAGC	GATACAACACGCGTTGGGATCTTC	CAPS (PvuII)

List of primers used for RNAi construct

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
RNAi.CNL-04	CTTCCCCCAATTGCCTCAGT	TGGCCTATTCGGATGCTCCTT	Construction of the plasmid pANDA-CNL04
RNAi.CNL-05	CTGCCATCTTATGAGTGGAGGG	CGCGTTGGCCCTTGTAGAGAAC	Construction of the plasmid pANDA-CNL05
RNAi.CNL-06	GCACTCAAGCCCAGGTCTAC	AAGTCTTTGTGCAGGCTCGT	Construction of the plasmid pANDA-CNL06
RNAi.CNL-07	CAAGGGCACATTGGACTACC	TACGGACCTCTGCAACATGC	Construction of the plasmid pANDA-CNL07
RNAi.CNL-08	TAATTGTCCGAGGGCAGTGG	ACATGCTGCATTTTCCGGTG	Construction of the plasmid pANDA-CNL08
RNAi.CNL-09	ACTACGGAGCGTGTGGATTG	GGGCACCGCCTATAATCTCC	Construction of the plasmid pANDA-CNL09
RNAi.NL-02	AACTACGGAGCGTGTGGATTG	ACCGCCTATGATCTCCAAGTG	Construction of the plasmid pANDA-NL02
RNAi.NL-04	CAAGATTCGCAGGCTCTCAATAC	CCTGCAGAGCTGCCATGTTAGC	Construction of the plasmid pANDA-NL04

List of primers used for q-PCR of CNL-04 and CNL-05

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
q-PCR_CNL-04	CTTAGGGAGGCCATTCAACA	AACAAACTGAGGCAATTGGG	qPCR for CNL-04
q-PCR_CNL-05	AAAGCAGCATTCTGTCGAAAAG	CTTTTCAATGCCCACTACGAAG	qPCR for CNL-05
OsACTIN1	ACCATTGGTGCTGAGCGTTT	CGCAGCTTCCATTCCTATGAA	qPCR for OsACTIN1

List of primers used for CRISPR/Cas9 construct of CNL-04 and CNL-05

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
sgRNA4.1	GTTGCGTACAGACGCTATGCATGG	AAAACCCATGCATAGCGTCTGTACG	Construction of the plasmid pZH::gYSA::MMCas9-CNL-04.1
sgRNA5.1	GTTGAGCTCGATTCCGTCCTTCAG	AAACCTGAAGGACGGAATCGAGCT	Construction of the plasmid pZH::gYSA::MMCas9-CNL-05.1
sgRNA5.2	GTTGCCATCCAGTATCTTGCCCTG	AAACCAGGGCAAGATACTGGATGG	Construction of the plasmid pZH::gYSA::MMCas9-CNL-05.2
sgRNA5.3	GTTGCCATCCAGTATCTTGCCCTG	AAACCAGGGCAAGATACTGGATGG	Construction of the plasmid pZH::gYSA::MMCas9-CNL-05.3
CNL-04 sel	TCTCCCTCGTCAACGACAAGCTC	TCTTCTTCTCCAACTGTTGTCTG	Selection of CNL-04 knockout rice line
CNL-05 sel	ACTGCATCGACACCATGTTTTC	CCAAGAAGCCTCCTCATGTCAG	Selection of CNL-05 knockout rice line

List of primers used for AVR-Pias candidate transgene construct

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
G9141	GCAtctagaGGCAGACACAATGCAGTTCAACATCAAGAC	CGAggatccTTAAGTGCAGTTGCTTTTGCTG	Construction of the plasmid pCB1531-pex22p-G9141
G9435	GCAtctagaGGCAGACACAATGGTTTCTTTTACTTGGACCTATTTG	CGAggatccTTATGCAGGGGGCGAAGCGCCTG	Construction of the plasmid pCB1531-pex22p-G9435
G9532	GCAtctagaGGCAGACACAATGCGTTTTTTCATCTATTCC	CGAggatccTTATTCAGGATAACCAAAAACAG	Construction of the plasmid pCB1531-pex22p-G9532

List of primers used for CRISPR/Cas9 construct of AVR-Pias

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
AVR-Pias-target1	ttcgTGCTATTGCCGCCCAAATTC	aaacGAATTTGGGCGGCAATAGCA	Construction of the plasmid pCRISPR-Cas-U6-1-AVR-Pias.1
AVR-Pias-target2	ttcgAACGCCGAAATGAAAATTAT	aaacATAATTTTCATTTCGGCGTT	Construction of the plasmid pCRISPR-Cas-U6-1-AVR-Pias.2
AVR-Pias-U1	TGGGTACCAACGCAACGGGAGTACAAATATC	CCCTCGACGAAGTAGAGGGTTGTGGAGATGGAC	Construction of the plasmid for TGR of AVR-Pias
AVR-Pias-L1	TTGATATCAAAGCTGTTGGAATAGAGCAGAAC	GCTCTAGACGGCGGTTTGAGACCTCTGGTGAG	Construction of the plasmid for TGR of AVR-Pias
AVR-Pias-sel	CACCGCCCAAATTCAGGATTCTTC	TTATTCAGGATAACCAAAAAC	Selection of AVR-Pias knockout *M. oryzae* line
Hyg-sel	CGACAGCGTCTCCGACCTGA	CGACCTCGTATTGGGAATCC	Selection of AVR-Pias knockout *M. oryzae* line

List of primers used for Pias/RGA5 allelic genes cloning

Primer name	Foward: sequence (5'-3')	Reverse: sequence (5'-3')	Usage in this study
Pias-1-HA	cctctagagtcgaccATGGAGGCCGCGCTTCTG	cgtacggataactagtTGTACAAGAAGGCACCATG	Construction of the plasmid pCambia1300S-Pias-1:HA
RGA4-HA	cctctagagtcgaccATGGAGGCCGCGCTTTTGAG	cgtacggataactagtCTCACAGCAGGAGACATCTG	Construction of the plasmid pCambia1300S-RGA4:HA
RGA4-Oau	cctctagagtcgaccATGGAGGCCGCGCTTTTGAG	cgtacggataactagtTGTACAAGAAGGTACCATGCC	Construction of the plasmid pCambia1300S-RGA4-Oau:HA
RGA4-Oru	cctctagagtcgaccATGGAGGCCGCCCTTCTGAG	cgtacggataactagtTGTACAAGAAGGCACCATG	Construction of the plasmid pCambia1300S-RGA4-Oru:HA
RGA4-Ogr	cctctagagtcgaccATGGAGGCCGTCCTTCTG	cgtacggataactagtTGTACAAGAAGGCACCATG	Construction of the plasmid pCambia1300S-RGA4-Ogr:HA
FLAG-Pias-2	atcctctagagtcgacATGGATGCTCCGGTGAG	tcgagcttgcatgccTTAGGGTTTTCCTGACAA	Construction of the plasmid pCambia1300S-FLAG:Pias-2
FLAG-RGA5	atcctctagagtcgacATGGATGCTCCGGCGAG	tcgagcttgcatgccTCACATGGTTGAGCAAGG	Construction of the plasmid pCambia1300S-FLAG:RGA5

Suppl. Table 10. List of NLRs and AVR-Pia used in the cell death assays.
OD$_{600}$ of *Agrobacterium tumefaciens* used for agroinfiltration is shown.

Protein name	Epitope tag	Density (OD$_{600}$)
Pias-1	C-terminal HA	0.2
RGA4	C-terminal HA	0.2
RGA4-Oru	C-terminal HA	0.2
RGA4-Oau	C-terminal HA	0.2
RGA4-Ogr	C-terminal HA	0.2
Pias-2	N-terminal FLAG	0.4
RGA5	N-terminal FLAG	0.4
AVR-Pia	Untagged	0.3