1   **Major article**

3   **Redistribution of the novel *C. difficile* spore-adherence receptor, E-cadherin, by TcdA and**

4   **TcdB increases spore-binding to adherens junctions.**

6   Pablo Castro-Córdova[1,#,$], Macarena Otto-Medina[2,#], Nicolás Montes-Bravo[1,#], Christian Brito-

7   Silva[1], D. Borden Lacy[3], Daniel Paredes-Sabja [1,4]*

9   [#] Authors contributed equally to this work.

10   [1] ANID – Millennium Science Initiative Program – Millennium Nucleus in the Biology of the

11   Intestinal Microbiota, Santiago, Chile.

12   [2].Departamento de Ciencias Biológicas, Facultad de Ciencias de la Vida, Universidad Andrés

13   Bello, Santiago, Chile.

14   [3] Department of Pathology, Microbiology, and Immunology, Vanderbilt University School of

15   Medicine, Nashville, Tennessee, USA.

16   [4] Department of Biology, Texas A&M University, College Station, Texas, USA.

17   [$]Present address: IMPACT, Center of Interventional Medicine for Precision and Advanced

18   Cellular Therapy, Santiago, Chile.

Short Title: Association of *C. difficile* spores to adherens junctions.

Key words*: C. difficile* spores, E-cadherin, adherens junctions

*Corresponding Author. Dr. Daniel Paredes-Sabja, Department of Biology, Texas A&M

University, College Station, TX, 77843, USA. E-mail: dparedes-sabja@bio.tamu.edu

Word Count

Abstract: 167 words

Text (introduction, methods, results, and discussion): 3493 words

References: 50

ABSTRACT

Clostridioides difficile causes antibiotic-associated diseases in humans ranging from mild

diarrhea to severe pseudomembranous colitis and death. A major clinical challenge is the

prevention of disease recurrence, which affects nearly ~20 to 30 % of the patients with a primary

C. difficile infection (CDI). During CDI, *C. difficile* forms metabolically dormant spores that are

essential for recurrence of CDI (R-CDI). In prior studies, we have shown that *C. difficile* spores

interact with intestinal epithelial cells (IECs), which contributes to R-CDI. However, this

interaction remains poorly understood. Here, we provide evidence that *C. difficile* spores interact

with E-cadherin, contributing to spore-adherence and internalization into IECs. *C. difficile* toxins

TcdA/TcdB lead to adherens junctions opening and increase spore-adherence to IECs. Confocal

micrographs demonstrate that *C. difficile* spores associate with accessible E-cadherin; spore-E-

cadherin association increases upon TcdA/TcdB intoxication. The presence of anti-E-cadherin

antibodies decreased spore adherence and entry into IECs. By ELISA, immunofluorescence, and

immunogold labelling, we observed that E-cadherin binds to *C. difficile* spores, specifically to the

hair-like projections of the spore, reducing spore-adherence to IECs. Overall, these results expand

our knowledge of how *C. difficile* spores bind to IECs by providing evidence that E-cadherin acts

as a spore-adherence receptor to IECs and by revealing how toxin-mediated damage affects spore

interactions with IECs.

Key Words:

C. difficile spores, E-cadherin, spore-host interaction, *C. difficile* toxins TcdA/TcdB.

INTRODUCTION

C. difficile is a Gram-positive, anaerobic, spore-forming bacterium and the main causative agent of antibiotic-associated diarrhea, causing death in ~5% of cases (1, 2). R-CDI is a major clinical challenge affecting ~30% of CDI patients (2, 3). CDI is primarily driven by the *C. difficile* toxins TcdA and TcdB, with contributions from the binary transferase toxin in some cases (4-6). Both TcdA and TcdB target their intracellular glucosyltransferase activity towards RhoA, Rac1, and Cdc42 (7, 8). This inhibitory modification of the Rho GTPase family leads to actin depolymerization and disruption of tight and adherens junctions (9-13), causing disruption of epithelial paracellular barrier function (14). Clinical studies showed that administration of anti-TcdB antibody (bezlotoxumab) in combination with antibiotics reduces R-CDI rates (15). The antibody may prevent toxin-mediated stem cell dysfunction and therefore prevent the delay in recovery and repair of the intestinal epithelium (10).

During infection, *C. difficile* produces spores (16). Strains unable to form spores do not cause R-CDI in animal models (17), indicating that the spores are critical for R-CDI. EDTA-disruption of cell-cell junctions in Caco-2 cells leads to increased adherence of *C. difficile* spores at the cell-cell junction, suggesting a tropism for cell-cell junctions (18). *C. difficile* spores bind to fibronectin and vitronectin, located in the basolateral membrane (19). These molecules contribute to *C. difficile* spore internalization into IECs in an integrin-dependent manner, and spore-entry *in vivo* contributes to the R-CDI (20).

The intestinal barrier is an impermeable layer that separates the apical and basolateral cell sides (21), and is formed by junctions between adjacent IECs. The cell-cell junctions between IECs

are constituted of E-cadherin containing adherens junctions and tight junctions formed by claudins,

occludins, and zonula occludens. These junctions provide the gut barrier with impermeability

properties and restrict the translocation of pathogens and commensal bacteria. However, some

pathogens can alter or disrupt the cell junctions, to translocate through the host (22).

Several enteric pathogens such as *Streptococcus pneumoniae*, *Fusobacterium nucleatum*,

Campylobacter jejuni, and *Listeria monocytogenes* take advantage of disrupted intestinal adherens

junctions to internalize in host cells (23-27). How *C. difficile* spores interact with adherens

junctions remains poorly described. Herein, we show that accessible E-cadherin is reduced in

differentiated Caco-2 cells. Also, TcdA/TcdB intoxication increases the spore adherence that is

correlated with an increase in E-cadherin accessibility in IECs. Next, we observed that *C. difficile*

spores interact with E-cadherin in healthy IECs and the binding is increased in TcdA/TcdB-

intoxicated cells. Also, the blocking of the cellular E-cadherin with monoclonal antibodies reduces

the spore adherence and entry to IECs. Finally, we observed that E-cadherin binds to the hair-like

extensions of the exosporium.

METHODS

Bacterial growth conditions.

C. difficile was grown under anaerobic conditions at 37 °C in a Bactron III–2 chamber

(Shellab, OR, USA) in BHIS medium; 3.7% brain–heart infusion broth (BD, USA) supplemented

with 0.5% of yeast extract (BD, USA), and 0.1% L–cysteine (Merk, USA) liquid or on plates with

1.5% agar (BD, USA). *C. difficile* strain R20291 wild-type B1/NAP1/PCR-ribotype 027 strain

obtained from CRG0825 stock at Nottingham Clostridia Research Group (28). *C. difficile*

630Δ*erm* is a toxigenic (*tcdA*⁺ *tcdB*⁺) strain of *C. difficile* isolates from a patient with

pseudomembranous colitis during an outbreak of CDI (29). *C. difficile* 630Δ*erm bclA1::CT*,

630Δ*erm bclA2::CT* and 630Δ*erm bclA3::CT* are insertional inactivation clostron mutant strains

in the exosporium collagen-like proteins BclA1, BclA2 and BclA3 (30), and were a kind gift from

Dr. Simon Cutting (School of Biological Sciences, Royal Holloway, University of London).

Bacillus megaterium transformed with the plasmid pHIS 1622 that contains the cloned genes *tcdA*

and *tcdB* of *C. difficile* VPI 10463 (31), were aerobically grown at 37 °C in Luria-Bertani (BD)

with 10 µg/mL tetracycline.

Cell lines and culture conditions

Caco-2 cells (ATCC, USA) were grown at 37 °C with 5% CO_2 with Dulbecco's modified

Eagle's minimal essential medium High Glucose (DMEM; Hyclone, USA) supplemented with

10% of fetal bovine serum (FBS; Hyclone, USA) and 50 U mL⁻¹ penicillin, and 50 µg mL⁻¹

streptomycin (Corning Cellgro, USA) as previously has been described For experiments, cells

were plated over glass coverslips in 24–well plates and were cultured for 2–days post–confluence

(undifferentiated) or 8–days post–confluence (differentiated), changing the media every other day.

To polarize Caco-2 cells were cultured onto 12mm Transwell (Corning, USA) with pores size of

0.4 µm until transepithelial electric resistance >250 Ω × cm².

Recombinant *C. difficile* toxin Purification

C. difficile toxins, TcdA and TcdB, were expressed in *B. megaterium* transformed with the

plasmid pHIS1622, containing the cloned genes *tcdA* and *tcdB* of *C. difficile* VPI 10463 as

previously described (31). Briefly, transformed *B. megaterium* were cultured in Luria–Bertani

medium (BD, USA) supplemented with 10 µg mL^{-1} tetracycline, and 35 mL of an overnight

culture was used to inoculate 1L of media. Bacteria were grown at 37 °C with 220 rpm. Toxin

expression was induced with 0.5% D–xylose once the culture reached OD$_{600}$ = 0.5. Cells were

harvested after 4 h by centrifugation at 4,750×g and resuspended in 200 mL of binding buffer (20

mM Tris [pH 8.0], 100 mM NaCl for TcdA and 20 mM Tris [pH 8.0], 500 mM NaCl for TcdB)

supplemented with 0.16 µg mL^{-1} DNase, 10 mg mL^{-1} lysozyme and protease inhibitors (P8849;

Sigma). Cell lysis was performed using an Emulsiflex homogenizer, and lysates were centrifuged

at 48,000×g for 30 min. The toxins were purified from the supernatant by Nickel–affinity, anion

exchange, and size–exclusion chromatography. Toxins were eluted and stored in 20 mM HEPES

(pH 7.0), 50 mM NaCl.

Spore preparation and purification

C. difficile spores were purified as we previously described (32). Briefly, *C. difficile* spore

was cultured overnight in BHI medium in a Bactron III–2 anaerobic chamber (Shellab, USA).

Then a 1:500 dilution was plated in TY agar plates; 3% trypticase soy and 0.5% yeast extract and

1.5% Bacto agar (BD, USA) and incubated for 7 days at 37 °C in anaerobiosis. Colonies were

harvested with sterile, cold Milli–Q water, washed 3 times and purified with 50% Nicodenz. The

spore suspension was 99% free of vegetative cells, sporulating cells, and debris. Spores were

quantified in a Neubauer chamber and stored at –80 °C until use.

Immunofluorescence of cell monolayers.

To evaluate E-cadherin distribution in undifferentiated and differentiated Caco-2 cells,

cells were washed with PBS and were fixed with PBS–4% paraformaldehyde for 20 min at 4 °C,

blocked with PBS–3% BSA 1 h at room temperature (RT). To stain the accessible E-cadherin,

cells were incubated with 1: 50 of primary antibody mouse polyclonal anti–human E-cadherin (sc-

8426, Santa Cruz Biotechnology, USA) in PBS–3% BSA, for 2 h at RT followed by two washed

with PBS and 1:200 secondary antibody donkey anti–mouse-Alexa Fluor 568 (ab175700, Abcam,

USA). To detect the total E-cadherin, fixed samples were permeabilized with PBS-0.2% Triton

100-X for 10 min at RT and blocked with PBS–3% BSA for 1 h at RT. The total protein was

detected with the same primary antibody used accessible E-cadherin; primary antibody mouse

polyclonal anti–human E-cadherin (sc-8426, Santa Cruz Biotechnology, USA) in PBS–3% BSA

that was incubated for 2 h at RT, then were washed and was incubated with 1:200 secondary

antibody donkey anti–mouse-Alexa Fluor 568 (ab175700, Abcam, USA). The non–permeabilized

samples stained for accessible E-cadherin were permeabilized with PBS-0.2% Triton 100-X for

10 min at RT, washed, and to stain the F–actin, cells were incubated with 1:1,000 Phalloidin i–

Fluor 488 (ab176753, Abcam, USA) in PBS for 30 min at RT. Cells were washed, and nuclei were

stained with 8 μg mL^{-1} Hoechst 33342 (ThermoFisher, USA) in PBS for 10 min at RT. Then

samples were washed and mounted using Dako Fluorescent Mounting Medium (Dako, Denmark).

Samples were visualized in the confocal microscope Leica TCS LSI (Leica, Germany). Please see

below.

162    To study the redistribution of E-cadherin in TcdA/TcdB intoxicated cells, differentiated

Caco-2 cells cultured into 24–well plates were washed twice with DPBS (Hyclone, USA) and were

intoxicated with 600 pM TcdA and 600 pM TcdB for 3, 6, and 8 h in DMEM, without serum. As

a control, cells were incubated in the same solution without TcdA/TcdB. Then cells were rinsed

with PBS, fixed with PBS–4% paraformaldehyde, and washed with PBS and blocked with PBS–

1% BSA at 4 °C overnight. To stain the surface, accessible E-cadherin cells were incubated with

1:200 mouse mAb anti–E-cadherin (SC8426, Santa Cruz Biotechnologies, USA), 1 h at RT,

washed, and followed incubation with 1:400 donkey anti–mouse IgG Alexa Fluor 488 (ab150109,

Abcam, USA) for 1 h at RT. To detect the total protein, samples were permeabilized with PBS–

0.2% Triton 100-X for 10 min at RT and blocked with PBS–1% BSA for 1 h at RT. The total

protein was detected with the same primary antibody used before, washed, and samples were

incubated with donkey anti–mouse IgG Alexa Fluor 568 (ab175700, Abcam, USA). Samples were

mounted using Dako Fluorescent Mounting Medium (Dako, Denmark) and were visualized in the

confocal microscope Leica Sp8 (Leica, Germany; please see below.)

To evaluate *C. difficile* spore association with accessible E-cadherin in TcdA/TcdB

intoxicated IECs, differentiated Caco-2 cells were intoxicated with 600 pM TcdA and 600 pM

TcdB for 8 h at RT, as described above. Then cells were infected with *C. difficile* spores in an MOI

of 10 for 1 h at 37 °C. Then cells were washed, fixed with PBS–4% paraformaldehyde for 10 min

at RT, blocked with PBS–1% BSA overnight at 4 °C, and *C. difficile* spores were stained with

1:1,000 primary polyclonal anti–*C. difficile* spore IgY batch 7246 antibodies (33) (Aveslab, USA)

in PBS–1% BSA for 1 h at RT. Following two PBS washed, samples were incubated with 1:400

goat anti–chicken IgY secondary antibodies Alexa–Fluor 488 (#ab150173 Abcam, USA) in PBS–

3% BSA for 1 h at RT, washed two times with PBS, and accessible E–cadherin was stained as it

is described above. Cellular nuclei were stained with 8 µg mL^{-1} Hoechst 33342 (ThermoFisher,

USA) for 15 min at RT. Finally, samples were mounted and analyzed by confocal microscopy

Leica SP8.

The Leica microscopes TCS LSI and SP8 at the Confocal Microscopy Core Facility of the

Universidad Andrés Bello were used. To evaluate E-cadherin distribution in undifferentiated and

differentiated Caco-2 cells, Leica TCS LSI was used with 63× ACS APO oil objective numerical

aperture 1.3, with an optical zoom of 2×. Micrographs were acquired using excitation wavelengths

of 405, 488, and 532 nm, and signals were detected with an ultra–high dynamic photomultiplier

(PMT) spectral detector (430–750nm). Emitted fluorescence was split with four dichroic mirrors

(QD 405, 488, 561, and 635 nm). Images 1,024 × 1,024 pixels were acquired each 0.3 μm. To

observe the redistribution of accessible E-cadherin and total E-cadherin in TcdA and TcdB

intoxicated cells, Leica SP8 was used with HPL APO CS2 40× oil, numerical aperture 1.30. For

signal detection, three PMT spectral detectors were used; PMT1 (410–483) DAPI, PMT2 (505–

550), Alexa–Fluor 488, and PMT3 (587–726) Alexa–Fluor 555. Emitted fluorescence was split

with dichroic mirrors DD488/552. Pictures of 1,024 × 1,024 pixels were captured each 0.3 μm.

Images of 3–dimensional reconstructions of the Caco-2 monolayers were performed using ImageJ

software (NIH, USA). In undifferentiated and differentiated cells, accessible E-cadherin cells were

quantified as the number of cells with positives staining. To quantify the redistribution of E-

cadherin, Fl. Int. of each slide was measured using the plug–in "Measure stack" of Fiji (ImageJ,

NHI, USA), and the sum of the Fl. Int. of each slide is considered as the Fl. Int. of the z–stack of

a field for accessible and total E-cadherin. For each field, the same number of slides was used.

Quantification of *C. difficile* spore adherence and internalization.

To evaluate spore adherence and internalization in TcdA/TcdB intoxicated cells,

differentiated Caco-2 cells were incubated with 600 pM TcdA and 600 pM TcdB for 3, 6, and 8 h

in DMEM, without serum. As a control, cells were incubated in the same solution without

TcdA/TcdB or cells were incubated with TcdA/TcdB previously heat inactivated at 70°C for 5

min. Then cells were infected with *C. difficile* spores an MOI 10 for 3 h at 37 °C. Unbound spores

were rinsed off with PBS, and cells were fixed with PBS–4% paraformaldehyde for 10 min, and

adhered spores were stained as described below.

To evaluate spore adherence and entry in presence of anti–E-cadherin, undifferentiated,

differentiated, and polarized Caco-2 cells were incubated with 0,1; 1,0; 10; 100 µg mL^{-1} of mouse

monoclonal anti–human E-cadherin (SC8426, Santa Cruz Biotechnology, USA) or as control non–

immune IgG antibody of rabbit serum (I5006, Sigma-Aldrich, USA) for 2 h at 37 °C. Then cells

were infected for 3 h at 37 °C with *C. difficile* spores, previously incubated with 20 µL of FBS per

well for 1 h at 37 °C. Cells were washed two times with PBS to remove unbound spores and were

fixed with PBS–4% paraformaldehyde for 10 min at RT, and adhered spores were stained as

described below.

To evaluate spore adherence and entry in presence of purified E-cadherin, undifferentiated

and differentiated Caco-2 cells were infected with *C. difficile* spores previously incubated with 1,

2.5, 5, 10 and 25µg mL^{-1} of human purified E-cadherin (5085, Sigma-Aldrich, USA) in DMEM

or DMEM without FBS as control for 1 h at 37 °C. Then cells were infected and were incubated

for 3 h at 37 °C. Cells were washed two times with PBS to remove unbound spores, fixed with

PBS–4% paraformaldehyde for 10 min at RT, and adhered spores were stained as described below.

To evaluate whether the presence of E-cadherin decreases the spore adherence in

TcdA/TcdB intoxicated cells, undifferentiated and differentiated Caco-2 cells were 8 h-

TcdA/TcdB intoxicated and were infected for 1 h at 37 °C with *C. difficile* spores previously

incubated with 25 µg mL^{-1} of E-cadherin for 1 h at 37 °C. Cells were washed two times with PBS

to remove unbound spores, fixed with PBS–4% paraformaldehyde for 10 min at RT, and adhered

spores were stained as described below.

241   In undifferentiated, differentiated, and polarized *C. difficile* spore infected Caco-2 cells,

adhered *C. difficile* spores were immunostained using the intracellular spore exclusion

immunostaining assay as previously described (20). For this *C. difficile* spore–infected cells were

washed three times with PBS at RT, blocked with PBS–3% BSA overnight at 4 °C, and samples

were incubated with anti–*C. difficile* spore goat serum 1:50 (PAC 5573, Pacific Immunology,

USA) in PBS–1% BSA for 1 h at RT. The cells were washed 3 times with PBS and incubated with

1:400 secondary antibody donkey IgG anti–goat CFL 488 (SC362255, Santa Cruz Biotechnology,

USA) for 1 h at RT. Cells were washed, and DNA was stained with 8 µg mL^{-1} Hoechst 33342

(ThermoFisher, USA) for 15 min at RT. In the case of polarized cells, to stain the F–actin, samples

were incubated with 1:100 Phalloidin Alexa Fluor 568 (A12380, Invitrogen, USA) for 30 min at

RT. Samples were washed three times with PBS and one time with distilled water and mounted

with Dako Fluorescent Mounting Medium (Dako, Denmark). Samples were visualized in the

epifluorescence microscope Olympus BX53 fluorescence microscopy with UPLFLN 100× oil

objective (numerical aperture 1.30). Images were captured with the camera for fluorescence

imaging Qimaging R6 Retiga, and pictures were analyzed with ImageJ (NIH, USA). Adhered

spores were considered as spores in phase–contrast that were marked in fluorescence. Internalized

spores were considered spores in phase–contrast, but that do not have fluorescence. In the case of

polarized cells, it was only possible to quantify adhered spores. The number or adhered spores

were quantified, and data were normalized to the control.

ELISA of E-cadherin bound to *C. difficile* spores.

Ninety six-well plate were coated overnight at 4ºC with 100 µL/well containing 1.6×10^8

spores mL^{-1} or with 100 µL of PBS containing 1 µg of BSA in PBS (as a control). overnight at 4

°C. Wells were washed 5 times with PBS–0.05% Tween-20 (PBS–Tween) to remove unbound

spores and blocked with 200 µL of PBS–Tween–2% BSA (Sigma Aldrich, USA) for 1 h at 37 °C.

The spores were washed and incubated with 50 µL of 0.2, 2.0, 20, and 200 nM of recombinant

human purified E-cadherin (5085, Sigma-Aldrich, USA) for 1 h at 37 °C. Unbonded E-cadherin

was washed 5 times with 150 µL PBS–Tween and was incubated with 50 µL of 1:50 IgG of mouse

anti–E-cadherin (SC8426-Santa Cruz Biotechnology, USA) in PBS–Tween–1% BSA for 1 h at 37

°C. Wells were washed five times with PBS–Tween and then incubated with 50 µL of 1: 5,000

secondary antibody anti–mouse-HRP (610-1302, Rockland, USA) 1 h at 37 °C. Then, wells were

washed five times with PBS–Tween and once with carbonate buffer (pH 9.6). Finally, wells were

incubated with 50 µL of buffer substrate (17 mM citric acid, 65 mM potassium phosphate, 0.15%

hydrogen peroxide, and 0.4% θ-phenylenediamine) for 20 min at RT. The reaction was stopped

with 25 µL of 4.5 N H_2SO_4. The color intensity was quantified at 450 nm using the plate reader

Infinite F50, (Tecan, Switzerland).

Immunofluorescence of E-cadherin bound to *C. difficile* spores.

The immunofluorescence was performed as previously described with modification (19).

Briefly, 4×10^7 *C. difficile* spores were incubated with 1, 2.5, 5.0, 10, and 25 µg mL^{-1} of

recombinant human purified E-cadherin (5085, Sigma-Aldrich, USA) in PBS for 1 h at 37 °C.

Spores were washed 5 times with PBS and fixed with PBS–4% paraformaldehyde in poly–L-

lysine-treated 12 mm glass–coverslips (Marienfeld GmbH, Germany). The spores were washed 3

times with PBS and were blocked with PBS–3% BSA for 1 h at RT. Then, samples were incubated

with 1:50 mouse IgG anti–E-cadherin (SC8426, Santa Cruz Biotechnology, USA) for 2 h at RT

and washed 3 times with PBS and once with distilled H_2O. Finally, samples were mounted with

Dako Fluorescent Mounting Medium (Dako, Denmark), and images were captured in an

epifluorescence microscope Olympus BX53. The fluorescence intensity was quantified using Fiji

(ImageJ, NHI, USA). Data were normalized as (Fl. Int/area)$_{sample}$ − (Fl. Int/area)$_{background.}$ The

Surface Plot was generated with the plug–in 3D surface Plot of Fiji (ImageJ, NHI, USA).

SDS-PAGE and western blot of intoxicated Caco-2 cells.

Undifferentiated and differentiated Caco-2 cells were intoxicated for 3, 6 and 8 h with TcdA/TcdB

in 24-well plates. The cells were washed and frozen at −80 °C. The frozen cells were lysed with

80 µL/well of cold RIPA buffer (50 mM Tris-HCl; 150 mM NaCl; 0.5% Deoxycholate; 1% NPO_4;

1mM EGTA; 1mM EDTA; 0.5% SDS; protease inhibitor) in ice and centrifuged at 18,400×g for

15 min at 4 °C. The supernatant was collected and 20 µg of protein was suspended in 2× SDS-

PAGE sample loading buffer, boiled and electrophoresed on 12% and 4% acrylamide SDS-PAGE

gels. Proteins were transferred to nitrocellulose membrane and blocked with 5% non-fat milk in

TBS for 1 h at RT. Membranes were probed for non-glucosylated Rac1 in Tris -buffered saline

containing 0.1% Tween (TTBS) with 5% milk and 1:1,000 mouse anti-Rac1 (Clone 102/Rac1

610650; BD Bioscience, USA) and then incubated overnight at RT and 6 h at RT. The membranes

were washed with TTBS and incubated with 1:5,000 goat anti-mouse HRP for 1 h at RT (Rockland

610-1302) in 5% milk-TTBS. Samples were washed 3 times with TTBS and HRP activity was detected with a chemiluminescence detection system (Li-core) using PicoMax sensitive chemiluminescence HRP substrate (Biorad, USA). To detect total Rac-1 in undifferentiated cells, membranes were incubated twice with mild stripping solution (1.5% glycine, 0.1% SDS, 1% Tween 20; pH 2.2) for 7 min at RT, washed twice with PBS and twice with TTBS 0.05%, and blocked with 5% non-fat milk-TBS 1 h at RT. For differentiated cells, total Rac1 was detected in the other membrane loaded simultaneously with the same concentration of protein. Membranes were incubated with 1:1,000 mouse anti-Rac1 (clone 23A8 05-389; Merck Millipore, USA) overnight at 4 °C and 3 h at RT and then washed 3 times with TTBS. Membranes were then incubated with 1:5,000 goat anti-mouse HRP (Rockland 610-1302) 1 h at RT and washed 3 times with TTBS. HRP activity was detected using Li-core as described for non-glucosylated Rac1.

Transmission electron microscopy of E-cadherin bound to *C. difficile* spores.

2.5×10^8 *C. difficile* spores were incubated with $10 \, \mu g \, mL^{-1}$ of recombinant human purified E-cadherin (5085, Sigma-Aldrich, USA) with PBS for 1 h at 37 °C. Spores were washed 5 times with PBS and centrifugation at $18{,}400 \times g$ for 5 min and were blocked with PBS–1% BSA for 20 min at RT. Then samples were incubated with 1:50 IgG of mouse anti–E-cadherin (SC8426, Santa Cruz Biotechnologies, USA) in PBS–1% BSA for 1 h at RT, washed 3 times by centrifugation at $18{,}400 \times g$ for 5 min with PBS. Spores were incubated with IgG anti–mouse conjugated to 10nm gold nanoparticles (G7777, Sigma-Aldrich, USA) for 1 h at RT. Then were washed 3 times with PBS–0.1% BSA by centrifugation at $18{,}400 \times g$ for 5 min. Subsequently, spores were processes for TEM, as previously have been described (19). Briefly, spores were fixed in 0.1 M cacodylate buffer–2.5% glutaraldehyde with 1% paraformaldehyde and then fixed with 0.1 M cacodylate

buffer–1% osmium tetroxide and stained with 1% tannic acid for 30 min. Samples were dehydrated

with increasing acetone concentrations; 30% acetone (with 2% uranyl acetate only in this step) for

30 min, 50% acetone for 30 min, 70% acetone overnight, and 90% acetone for 30 min, and 2 times

with 100% acetone. Dehydrated samples were embedded in resin: acetone in a ratio of 3:1, 1:1,

and 1: 3 for 40 min each. Finally, samples were resuspended in resin for 4 h and incubated for 12

h at 65 °C. With a microtome, 90 nm sections cut and placed in a carbon–coated grid for negative

staining and double staining with 2% uranyl acetate with lead citrate. Sections were observed in a

transmission electron microscope Phillip Tecnai 12 bioTwin of the Pontificia Universidad Católica

de Chile. The distribution of thin or thick exosporium was quantified according to previously

described morphotypes (34).

RESULTS

Characterization of accessible E-cadherin in monolayers of Caco-2 cells. Caco-2 cells have

been employed to assess *C. difficile* interaction with IECs (18-20, 32, 33, 35-37). Early work

demonstrated that *C. difficile* spores had a tropism towards the cell-cell junction of EDTA-treated

Caco-2 cells (18). Therefore, to understand the dynamic of adherens junction closure during Caco-

2 cell differentiation, we assessed the accessibility of E-cadherin in undifferentiated and

differentiated monolayers of Caco-2 cells. First, Caco-2 cell differentiation was assessed for the

appearance of microvilli by TEM and sucrase-isomaltose as a differentiation marker in monolayers

cultured for 2- or 8-days post-confluency (19). Next, differentiated and undifferentiated Caco-2

cells were immunostained for accessible E-cadherin or permeabilized and stained for total E-

cadherin (Figure 1*A, B*). Results demonstrate that in undifferentiated Caco-2 cells, ~84% of the

cells were immunolabeled for accessible E-cadherin (Figure 1*C*), whereas, in differentiated Caco-

2 cells, only ~7% of the cells were stained for accessible E-cadherin (Figure 1*C*). As a control,

cells were permeabilized and stained for total E-cadherin, and we observed that ~100% of both

undifferentiated and differentiated Caco-2 cells were immunolabeled for E-cadherin (Figure 1*D*-

F). These results demonstrate that after 8-days of differentiation, most of the cells lack accessible

E-cadherin.

Intoxication with *C. difficile* TcdA and TcdB increases accessible E-cadherin. *C. difficile*

toxins TcdA and TcdB cause tight and adherens junction dissociation (9-11). TcdA, but not TcdB,

causes a displacement of E-cadherin from the cell-cell junctions to be located around the cells in

Caco-2 cells (11) while in colonic epithelium, TcdB causes adherens junctions disruption (10).

Since most of the *C. difficile* strains causing disease produces TcdA and TcdB (38), we asked

whether TcdA and TcdB affect the accessibility of E-cadherin in differentiated Caco-2 cells.

TcdA/TcdB-intoxicated differentiated Caco-2 cells for 3, 6, and 8 h were immunostained for

accessible E-cadherin (non-permeabilized) using an antibody that detects an extracellular domain

of E-cadherin, followed by fluorescent-labeled secondary antibody. For total E-cadherin staining,

cells were permeabilized and immunostained with the same primary antibody and a fluorescent-

labeled secondary antibody with another Alexa Fluor dye (See Methods). Then, we acquired

confocal micrographs, and the fluorescence intensity (Fl. Int.) was quantified for every slice of the

z-stack for accessible and total E-cadherin. The sum of the Fl. Int. of accessible E-cadherin was

significantly higher (175% relative to unintoxicated monolayers) after 8h-intoxication (Figure 2*A*,

B, Supplementary Figure 1). To evaluate if this increase of Fl. Int. occurs in the apical or basal half

of the monolayer, we split the number of slices of the z-stack into 2-halves (apical and basal) with

the same number of slices. We summed the Fl. Int. for accessible E-cadherin in the apical (Figure

2C) or basal (Figure 2D) halves of the monolayer. We observed a significant increase in accessible

E-cadherin Fl.Int. in both the apical and basal halves, 177% and 176% respectively, after 8h-

intoxication (Figure 2C, D). Using the same strategy, we quantified the total E-cadherin Fl. Int. in

the complete *z*-stack, and we observed a significant decrease of 19% after 6h- and an increase of

12.2% after 8h-intoxication compared to unintoxicated cells (Figure 2E). Then, we observed that

these variations were in the apical and not in the basal half of the monolayer. We observed a

decrease of 35% after 6h- and an increase of 44% after 8h-intoxication compared to unintoxicated

cells (Figure 2F, G). To confirm the degree of TcdA/TcdB intoxication, we evaluated the

glucosyltransferase activity by measuring the levels of glucosylated Rac1 in TcdA/TcdB

intoxicated differentiated and undifferentiated monolayers using strategies previously described

(39). After 3 h of intoxication no modification of Rac1 was observed. However, after 6 and 8 h of

intoxication, the unglucosylated Rac1 signal disappeared, indicating a complete glucosylation of

Rac1 (Figure 2H). These results indicate that the cells were fully intoxicated by 6 hours and

accessible E-cadherin is increased in differentiated Caco-2 cells after 8h-intoxication.

***C. difficile* toxins TcdA and TcdB increase the adherence to IECs.** Several studies have shown

that vegetative *C. difficile*-adherence to IECs is enhanced following TcdA or CDT intoxication

(11, 40, 41). To expand the understanding of how intoxication of IECs impacts spore-adherence,

differentiated and undifferentiated Caco-2 cells were TcdA/TcdB intoxicated for 8 h, or incubated

8 h with heat inactivated TcdA/TcdB as a control. Then cells were washed and infected for 1 h

with *C. difficile* spores in the absence of serum. Cells were fixed and immunostained without

permeabilization. In TcdA/TcdB intoxicated Caco-2 cells, we observed more adhered spores

compared to unintoxicated cells (Figure 3*A*). After quantitative analysis, we observed in undifferentiated Caco-2 cells that spore adherence in 8h-TcdA/TcdB-intoxicated cells was ~105% higher than in unintoxicated monolayers, and no differences in spore adherence were observed in cells 8 h-incubated with heat-inactivated TcdA/TcdB (Figure 3*B*). Similarly, in differentiated cells we observed an increase in ~321% in 8 h-TcdA/TcdB-intoxicated cells over that in unintoxicated monolayers, and no differences were observed in cells 8 h-incubated with heat-inactivated TcdA/TcdB (Figure 3*C*). Collectively these data indicate that TcdA/TcdB intoxication of undifferentiated and differentiated Caco-2 cells increases *C. difficile* spore adherence.

Intoxication of IECs increases *C. difficile* spore association with E-cadherin. Since we observed that TcdA/TcdB-intoxicated differentiated Caco-2 cells lead to an increase of accessible E-cadherin and adherence of *C. difficile* spores, we tested whether this could be attributed to *C. difficile* spores associating with accessible E-cadherin. For this, unintoxicated or TcdA/TcdB-intoxicated differentiated Caco-2 cells were infected with *C. difficile* spores for 1 h. Then, accessible E-cadherin and spores were immunostained and visualized by confocal microscopy (Figure 4*A, D*). Plot profile analysis of unintoxicated Caco-2 cells (Figure 4*A*) reveals that ~21% of adhered *C. difficile* spores were associated with accessible E-cadherin (Figure 4*B, C,* and *G*). Notably, upon TcdA/TcdB-intoxication (Figure 4*D*), ~72% of adhered spores were associated with accessible E-cadherin (Figure 4*E, F, G*). These results indicate that adhered *C. difficile* spores can associate with accessible E-cadherin in IECs, and this association is enhanced upon TcdA/TcdB-intoxication.

Antibody blocking of accessible E-cadherin reduces adherence and internalization of *C. difficile* spores to IECs. Next, we assessed whether E-cadherin is involved in *C. difficile* spore adherence and internalization. We have shown that *C. difficile* spore internalization occurs in the presence of serum (20). Therefore, undifferentiated, differentiated, and polarized Caco-2 cells were incubated with anti-E-cadherin antibody for 2h and subsequently infected for 3h with FBS-incubated *C. difficile* spores. To identify intracellular spores, we used the intracellular spore exclusion immunostaining assay (20), where only extracellular spores are immunostained in non-permeabilized cells, and total spores are visualized by phase-contrast microscopy. Using this strategy in undifferentiated Caco-2 cells, we observed that *C. difficile* spore adherence to IECs was significantly decreased upon increasing concentrations of anti-E-cadherin antibody; spore adherence decreased by 38, 53, and 73% compared to the control when cells were incubated with 1, 10, and 100 µg mL^{-1} of anti-E-cadherin antibody, respectively (Figure 5*A*). Upon quantifying internalization in undifferentiated Caco-2 cells, we observed a significant decrease of 38, 43, and 69% compared to the control in cells incubated with 1, 10, 100 µg mL^{-1} of anti-E-cadherin, respectively (Figure 5*B*). Similarly, in differentiated Caco-2 cells, a significant decrease in spore adherence in 18, 32, 60, and 70% compared to the control was observed in cells incubated with 0.1, 1, 10, and 100 µg mL^{-1} of anti-E-cadherin, respectively (Figure 5*C*). For internalization, we observed a significant decrease in 15, 35, 42, and 54% compared to the control in cells incubated with 0.1, 1, 10, and 100 µg mL^{-1} of anti-E-cadherin, respectively (Figure 5*D*). Finally, in polarized Caco-2 cells, we observed a decrease in 32, 56, 63, and 79% in spore adherence compared to the control in presence with 0.1, 1, 10, and 100 µg mL^{-1} of anti-E-cadherin, respectively (Figure 5*E*). These data indicate that the antibody-mediated blockage of accessible E-cadherin reduces *C. difficile* spore adherence and internalization to IECs.

Purified recombinant human E-cadherin competitively inhibits spore adherence in undifferentiated, differentiated and TcdA/TcdB intoxicated IECs cells. Subsequently, to provide further evidence that E-cadherin acts as a spore adherence receptors, we evaluated whether infected healthy undifferentiated Caco-2 monolayers with *C. difficile* spores preincubated with 1, 2.5, 5, 10, and 25 µg mL^{-1} of E-cadherin, and we observed a decrease in spore adherence of 12%, 26%, 38%, 47% and 64% respectively compared to spores not incubated with E-cadherin (Fig. 6*E*). In differentiated Caco-2 cells, we observed a decrease in spore adherence of 27%, 30%, 30%, 27% and 53% for spores incubated with 1, 2.5, 5, 10, and 25 µg mL^{-1} of E-cadherin, respectively (Figure. 6*F*). Finally, we also assessed whether the increased spore adherence that we previously observe in TcdA/TcdB intoxicated Caco-2 cells was E-cadherin specific. As expected, significant *C. difficile* spore adherence to TcdA/TcdB intoxicated Caco-2 cells was observed in the absence of E-cadherin; however, upon of TcdA/TcdB intoxicated differentiated and undifferentiated Caco-2 cells with spores previously incubated with 25µg mL^{-1} of E-cadherin for 1 h at 37°C, and we observed a significant decrease of 41% in undifferentiated (Figure 6*G*) and 55% in differentiated Caco-2 cells (Figure 6*H*) compared to TcdA/TcdB intoxicated cells. Overall, these results show that E-cadherin interacts with *C. difficile* spores in a dose-dependent manner and that E-cadherin-ligand binding sites are non-homogeneously distributed on the *C. difficile* spore surface. There results indicate that competition experiments with purified E-cadherin reduces the *C. difficile* spore adherence in healthy and intoxicated cells.

E-cadherin binds in a concentration-dependent manner to *C. difficile* spores. To provide evidence that *C. difficile* spores interact with E-cadherin, we used ELISA with *C. difficile* spore-

coated wells incubated with increasing concentrations of E-cadherin, or BSA as a control. E-

cadherin binding activity was evident in *C. difficile* spore-coated wells (Figure. 7A). The

specificity of this interaction was demonstrated by the dose-dependent binding. Additionally, we

evaluated the distribution of binding of E-cadherin to the spore surface by immunofluorescence.

Upon micrograph analysis, we observed that anti-E-cadherin immunostained spores exhibited a

non-homogeneous Fl. Int surrounding the spore (Figure. 7B). In spores, the Fl. Int. for E-cadherin

was 156, 189, 258, 304, and 369% relative to the control upon incubation with 1, 2.5, 5, 10, and

25 µg mL^{-1} of E-cadherin (Figure. 7C). Next, we established a Fl. Int cut-off limit for those *C.*

difficile spores that were visually associated with 2.5 µg mL^{-1} E-cadherin. Based on this threshold,

59, 65, and 66% of the spores were positive for E-cadherin in the presence of 5, 10, and 25 µg

mL^{-1} E-cadherin, respectively. Finally, we evaluated E-cadherin Fl. Int distribution for positive

immunodetected spores, and we observed that they were mainly distributed between 1.4-2.1

normalized Fl. Int. arbitrary units (Figure. 7D). Taken together, these results indicate that E-

cadherin binds in a concentration dependent manner to *C. difficile* spores, in a heterogeneous and

skewed manner to *C. difficile* spores.

E-cadherin interacts with hair-like projections of *C. difficile* spores. The outermost structure

of *C. difficile* spores are the hair-like projections, formed in part by BclA3, that are common in

epidemically relevant strains (16, 20, 42). Isogenic sporulating cultures of *C. difficile* lead to the

formation of spores with two exosporium morphotypes that exhibit hair-like projections, spores

with a thick exosporium layer display distinctive electron-dense bumps, and spores with a thin

exosporium lack these bumps (34, 42-44), suggesting that this exosporium morphotype diversity

might be contributing to E-cadherin spore-binding variability. Consequently, to gain ultrastructural

insight of the spore-E-cadherin interaction, *C. difficile* spores incubated with E-cadherin were

imaged by TEM and immunogold-labeling of E-cadherin. We observed E-cadherin-gold particles

in close contact with the exosporium of *C. difficile* spores, particularly with the hair-like

projections (Figure 8*A-E*). No gold particles were observed in spores without the E-cadherin

incubation. (Figure 8*A*). Next, within the E-cadherin positive spores, we observed gold particles

in 52/236 (22.1%) of the spores (Figure 8*F*). Of these, 19/52 (36.5%) and 33/52 (63.5%) of the E-

cadherin-bound spores corresponded to spores with thin and thick exosporiums, respectively

(Figure 8*G*). These results are representative of a 90 nm-slice of the sample, and therefore the

number of gold particles corresponds to those only present in the slice and not of a complete spore.

These results suggest that E-cadherin binds preferentially to thick-exosporium layer spores, which

seem to have enriched in hair-like projections.

and specifically to the hair-like projections.

Most *C. difficile* genomes encode three *bclA* orthologues encodes three collagen-like

proteins of the BclA family of proteins (45). However, *C. difficile* strain R20291 only encodes two

bclA orthologues (i.e., *bclA2* and *bclA3*) (45). Therefore, we reasoned to use previously

characterized *bclA1*, *bclA2* and *bclA3* mutants in strain 630 (30) to address by ELISA whether

these collagen-like proteins could be implicated in E-cadherin binding to *C. difficile* spores. As

with R20291 spores, E-cadherin binding to 630 spore coated wells was evident (Figure 8E).

Notably, upon coating wells with spores of *bclA1*, *bclA2* or *bclA3* spores, a dose dependent binding

of E-cadherin to spore-coated wells that as similar to that of wild-type spores, with the exception

of *bclA3* spores which exhibited higher binding of E-cadherin (Figure 8E). Overall, these results

suggest the presence of an unidentified E-cadherin-specific spore surface ligand.

DISCUSSION

A major clinical challenge that remains unresolved is how to address the high rates of CDI recurrence (46-48). *C. difficile* spores are regarded as the key virulence factors required for recurrence and transmission of the disease (16, 17). Although significant work has advanced our understanding of the biology of *C. difficile* spores (16, 42), the mechanisms through which *C. difficile* spores interact with the host within the environment of the colon remain poorly described. The integrity of the intestinal epithelial barrier is key in preventing pathogen translocation (22). However, in the healthy intestinal mucosa, the apical- and basolateral-cell sides are not always separated. The adherens junctions, where E-cadherin is located, can be accessible in certain situations, i.e., along lateral membranes of goblet cells (20, 49, 50), extruding apoptotic sites, produced transiently when senescent cells are expelled and detached from the epithelium (49, 51, 52) and in epithelial folds of the intestinal villus produced by tension and constriction forces (49). In this context, we have demonstrated that E-cadherin is a novel spore adherence-receptor that promotes binding of *C. difficile* spores at adherens junctions, and this binding is enhanced in intoxicated IECs.

Work by others have demonstrated that E-cadherin is hijacked by several pathogens, including *Listeria monocytogenes*, *Fusobacterium nucleatum*, and *Streptococcus pneumoniae* (23- 26, 49). This report expands the list of pathogens that hijack E-cadherin for adherence and entry to include *C. difficile* spores. Our prior work had demonstrated that *C. difficile* spores adhere to IECs *in vitro* (18) and gain entry into host cells via fibronectin and vitronectin in an integrin- dependent pathway (20). The ~70-80% reduction in adherence in undifferentiated, differentiated and polarized monolayers of Caco-2 cells upon blockade with anti-E-cadherin antibodies or competition with purified E-cadherin, suggests that E-cadherin is a major cellular adherence

receptor for *C. difficile* spores. Fibronectin, vitronectin, and integrin receptors are typically located

in the basolateral membrane (41, 53-56), where E-cadherin is localized. An outstanding question

is whether E-cadherin acts as an anchoring protein allowing *C. difficile* spores to bind

fibronectin/vitronectin and their cognate integrins to trigger internalization, or whether spore-

binding to E-cadherin alone is capable of triggering a signal for internalization of *C. difficile*

spores. This question will be addressed in future studies.

540   Another conclusion of this work is that *C. difficile* spores bind to E-cadherin in a dose-

dependent manner. *L. monocytogenes* interacts with human and not with mouse E-cadherin

through InlA (26). While the exosporium of *C. difficile* spores has no orthologs of InlA (57), we

show E-cadherin binding to the hair-like projections of the exosporium. *C. difficile* spores form

two exosporium morphotypes during sporulation (i.e., a thick and a thin exosporium layer) (42);

among the main differences is the presence of electron-dense bumps underlying the hair-like

projections. Whether there are differences in the hair-like projections and in their numbers between

both morphotypes, is unclear, and matter of study in our laboratory. The relevance of the

exosporium thickness in E-cadherin binding. Notably, spores of *bclA1* and *bclA2* orthologue in

strain 630 were able to bind to E-cadherin to a similar extent as wild-type spores, suggesting either

redundance in function. It was surprising to observe that *bclA3* mutant spores bind higher levels

of E-cadherin, suggesting that absence of BclA3 might uncover an alternative spore surface

ligand(s) implicated in binding of E-cadherin to the spore surface. Current ongoing work in our

laboratory aims to identify the spore surface ligand(s) that bind to E-cadherin and its implication

in disease.

A final and major conclusion of this work is that TcdA/TcdB intoxication increases spore

adherence to IECs in a E-cadherin-specific manner. This is supported by increased association

between basolateral E-cadherin and *C. difficile* spores, and by the observation that soluble E-

cadherin can reduce the adherence of *C. difficile* spores to TcdA/TcdB intoxicated IECs. *C. difficile*

toxins TcdA and TcdB cause a disruption of the cell-cell junctions (10, 12, 13), causing a loss of

cellular polarity (9-11). This disruption is caused by a dissociation of tight and adherens junction

proteins. In this context, our work contributes to understanding how *C. difficile* spores interact

with TcdA/TcdB-intoxicated IECs. Our results demonstrate that differentiated TcdA/TcdB-

intoxicated monolayers show an increased accessibility of E-cadherin that correlates with an

increase in spore adherence to TcdA/TcdB-intoxicated cells. These observations suggest that

during CDI, TcdA and TcdB increase the E-cadherin accessibility, contributing to the spore

adherence and entry into the IECs *in vivo*. Indeed, since intracellular *C. difficile* spores contribute

to disease recurrence (20), TcdA/TcdB -intoxication may significantly contribute to *C. difficile*

spore adherence and internalization during CDI, contributing overall to disease recurrence. These

hypotheses are in agreement with clinical observations in CDI patients where high titers of

endogenous antibodies against the *C. difficile* toxins correlate with reduced R-CDI rates (58), and

with the reduced R-CDI observed in the clinical trials of bezlotoxumab (15). The development of

multifactorial treatment with intervention in TcdA/TcdB, spore adherence, and entry to IECs might

importantly reduce the R-CDI rates.

NOTES

CONFLICT OF INTEREST. The authors declare that they have no conflict of interest.

FUNDING. This work was funded by FONDECYT Regular 1191601, and by Millennium Science

Initiative Program – NCN17_093, and Startup funds from the Department of Biology at Texas

A&M University to D.P-S. Additional funding support to P.C-C from ANID-PCHA/Doctorado

Nacional/ 2016-21161395.

REFERENCES

1.   Redelings MD, Sorvillo F, Mascola L. 2007. Increase in *Clostridium difficile*-related
mortality rates, United States, 1999-2004. Emerg Infect Dis 13:1417-9.
2.   Stevens VW, Nelson RE, Schwab-Daugherty EM, Khader K, Jones MM, Brown KA, Greene
T, Croft LD, Neuhauser M, Glassman P, Goetz MB, Samore MH, Rubin MA. 2017.
Comparative Effectiveness of Vancomycin and Metronidazole for the Prevention of
Recurrence and Death in Patients With *Clostridium difficile* Infection. JAMA Intern Med
177:546-553.
3.   Di X, Bai N, Zhang X, Liu B, Ni W, Wang J, Wang K, Liang B, Liu Y, Wang R. 2015. A meta-
analysis of metronidazole and vancomycin for the treatment of *Clostridium difficile*
infection, stratified by disease severity. Braz J Infect Dis 19:339-49.
4.   Lyerly DM, Lockwood DE, Richardson SH, Wilkins TD. 1982. Biological activities of toxins
A and B of *Clostridium difficile*. Infect Immun 35:1147-50.
5.   Lyras D, O'Connor JR, Howarth PM, Sambol SP, Carter GP, Phumoonna T, Poon R, Adams
V, Vedantam G, Johnson S, Gerding DN, Rood JI. 2009. Toxin B is essential for virulence of
Clostridium difficile. Nature 458:1176-9.
6.   Kuehne SA, Cartman ST, Heap JT, Kelly ML, Cockayne A, Minton NP. 2010. The role of toxin
A and toxin B in *Clostridium difficile* infection. Nature 467:711-3.
7.   Just I, Selzer J, Wilm M, von Eichel-Streiber C, Mann M, Aktories K. 1995. Glucosylation of
Rho proteins by *Clostridium difficile* toxin B. Nature 375:500-3.
8.   Just I, Wilm M, Selzer J, Rex G, von Eichel-Streiber C, Mann M, Aktories K. 1995. The
enterotoxin from *Clostridium difficile* (ToxA) monoglucosylates the Rho proteins. J Biol
Chem 270:13932-6.
9.   Nusrat A, von Eichel-Streiber C, Turner JR, Verkade P, Madara JL, Parkos CA. 2001.
Clostridium difficile toxins disrupt epithelial barrier function by altering membrane
microdomain localization of tight junction proteins. Infect Immun 69:1329-36.
10.   Mileto SJ, Jardé T, Childress KO, Jensen JL, Rogers AP, Kerr G, Hutton ML, Sheedlo MJ,
Bloch SC, Shupe JA, Horvay K, Flores T, Engel R, Wilkins S, McMurrick PJ, Lacy DB, Abud
HE, Lyras D. 2020. *Clostridioides difficile* infection damages colonic stem cells via TcdB,
impairing epithelial repair and recovery from disease. Proc Natl Acad Sci U S A 117:8064-
8073.
11.   Kasendra M, Barrile R, Leuzzi R, Soriani M. 2014. *Clostridium difficile* toxins facilitate
bacterial colonization by modulating the fence and gate function of colonic epithelium. J
Infect Dis 209:1095-104.
12.   Hecht G, Pothoulakis C, LaMont JT, Madara JL. 1988. *Clostridium difficile* toxin A perturbs
cytoskeletal structure and tight junction permeability of cultured human intestinal
epithelial monolayers. J Clin Invest 82:1516-24.
13.   Hecht G, Koutsouris A, Pothoulakis C, LaMont JT, Madara JL. 1992. *Clostridium difficile*
toxin B disrupts the barrier function of T84 monolayers. Gastroenterology 102:416-23.

14.   Leslie JL, Huang S, Opp JS, Nagy MS, Kobayashi M, Young VB, Spence JR. 2015. Persistence
and toxin production by *Clostridium difficile* within human intestinal organoids result in
disruption of epithelial paracellular barrier function. Infect Immun 83:138-45.
15.   Wilcox MH, Gerding DN, Poxton IR, Kelly C, Nathan R, Birch T, Cornely OA, Rahav G, Bouza
E, Lee C, Jenkin G, Jensen W, Kim YS, Yoshida J, Gabryelski L, Pedley A, Eves K, Tipping R,
Guris D, Kartsonis N, Dorr MB, Investigators MIaMI. 2017. Bezlotoxumab for Prevention
of Recurrent *Clostridium difficile* Infection. N Engl J Med 376:305-317.
16.   Paredes-Sabja D, Shen A, Sorg JA. 2014. *Clostridium difficile* spore biology: sporulation,
germination, and spore structural proteins. Trends Microbiol 22:406-16.
17.   Deakin LJ, Clare S, Fagan RP, Dawson LF, Pickard DJ, West MR, Wren BW, Fairweather NF,
Dougan G, Lawley TD. 2012. The *Clostridium difficile spo0A* gene is a persistence and
transmission factor. Infect Immun 80:2704-11.
18.   Paredes-Sabja D, Sarker MR. 2012. Adherence of *Clostridium difficile* spores to Caco-2
cells in culture. J Med Microbiol 61:1208-1218.
19.   Mora-Uribe P, Miranda-Cardenas C, Castro-Cordova P, Gil F, Calderon I, Fuentes JA, Rodas
PI, Banawas S, Sarker MR, Paredes-Sabja D. 2016. Characterization of the Adherence of
Clostridium difficile Spores: The Integrity of the Outermost Layer Affects Adherence
Properties of Spores of the Epidemic Strain R20291 to Components of the Intestinal
Mucosa. Front Cell Infect Microbiol 6:99.
20.   Castro-Córdova P, Mora-Uribe P, Reyes-Ramírez R, Cofré-Araneda G, Orozco-Aguilar J,
Brito-Silva C, Mendoza-León MJ, Kuehne SA, Minton NP, Pizarro-Guajardo M, Paredes-
Sabja D. 2021. Entry of spores into intestinal epithelial cells contributes to recurrence of
Clostridioides difficile infection. Nat Commun 12:1140.
21.   Buckley A, Turner JR. 2018. Cell Biology of Tight Junction Barrier Regulation and Mucosal
Disease. Cold Spring Harb Perspect Biol 10.
22.   Paradis T, Bègue H, Basmaciyan L, Dalle F, Bon F. 2021. Tight Junctions as a Key for
Pathogens Invasion in Intestinal Epithelial Cells. Int J Mol Sci 22.
23.   Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW. 2013. *Fusobacterium nucleatum*
promotes colorectal carcinogenesis by modulating E-cadherin/β-catenin signaling via its
FadA adhesin. Cell Host Microbe 14:195-206.
24.   Anderton JM, Rajam G, Romero-Steiner S, Summer S, Kowalczyk AP, Carlone GM,
Sampson JS, Ades EW. 2007. E-cadherin is a receptor for the common protein
pneumococcal surface adhesin A (PsaA) of *Streptococcus pneumoniae.* Microb Pathog
42:225-36.
25.   Elmi A, Nasher F, Jagatia H, Gundogdu O, Bajaj-Elliott M, Wren B, Dorrell N. 2016.
Campylobacter jejuni outer membrane vesicle-associated proteolytic activity promotes
bacterial invasion by mediating cleavage of intestinal epithelial cell E-cadherin and
occludin. Cell Microbiol 18:561-72.
26.   Mengaud J, Ohayon H, Gounon P, Mege R-M, Cossart P. 1996. E-cadherin is the receptor
for internalin, a surface protein required for entry of *L. monocytogenes* into epithelial
cells. Cell 84:923-32.
27.   Ribet D, Cossart P. 2015. How bacterial pathogens colonize their hosts and invade deeper
tissues. Microbes Infect 17:173-83.

28.   Ng YK, Ehsaan M, Philip S, Collery MM, Janoir C, Collignon A, Cartman ST, Minton NP.
668   2013. Expanding the repertoire of gene tools for precise manipulation of the *Clostridium*
669   *difficile* genome: allelic exchange using *pyrE* alleles. PLoS One 8:e56051.
670   29.   Wüst J, Sullivan NM, Hardegger U, Wilkins TD. 1982. Investigation of an outbreak of
671   antibiotic-associated colitis by various typing methods. J Clin Microbiol 16:1096-101.
672   30.   Phetcharaburanin J, Hong HA, Colenutt C, Bianconi I, Sempere L, Permpoonpattana P,
673   Smith K, Dembek M, Tan S, Brisson MC, Brisson AR, Fairweather NF, Cutting SM. 2014.
674   The spore-associated protein BclA1 affects the susceptibility of animals to colonization
675   and infection by *Clostridium difficile*. Mol Microbiol 92:1025-38.
676   31.   Chandrasekaran R, Kenworthy AK, Lacy DB. 2016. *Clostridium difficile* Toxin A Undergoes
677   Clathrin-Independent, PACSIN2-Dependent Endocytosis. PLoS Pathog 12:e1006070.
678   32.   Calderon-Romero P, Castro-Cordova P, Reyes-Ramirez R, Milano-Cespedes M, Guerrero-
679   Araya E, Pizarro-Guajardo M, Olguin-Araneda V, Gil F, Paredes-Sabja D. 2018. *Clostridium*
680   *difficile* exosporium cysteine-rich proteins are essential for the morphogenesis of the
681   exosporium layer, spore resistance, and affect *C. difficile* pathogenesis. PLoS Pathog
682   14:e1007199.
683   33.   Pizarro-Guajardo M, Diaz-Gonzalez F, Alvarez-Lobos M, Paredes-Sabja D. 2017.
684   Characterization of Chicken IgY Specific to *Clostridium difficile* R20291 Spores and the
685   Effect of Oral Administration in Mouse Models of Initiation and Recurrent Disease. Front
686   Cell Infect Microbiol 7:365.
687   34.   Pizarro-Guajardo M, Calderon-Romero P, Castro-Cordova P, Mora-Uribe P, Paredes-Sabja
688   D. 2016. Ultrastructural Variability of the Exosporium Layer of *Clostridium difficile* Spores.
689   Appl Environ Microbiol 82:2202-2209.
690   35.   Schöttelndreier D, Langejürgen A, Lindner R, Genth H. 2020. Low Density Lipoprotein
691   Receptor-Related Protein-1 (LRP1) Is Involved in the Uptake of *Clostridioides difficile* Toxin
692   A and Serves as an Internalizing Receptor. Front Cell Infect Microbiol 10:565465.
693   36.   Tam J, Icho S, Utama E, Orrell KE, Gómez-Biagi RF, Theriot CM, Kroh HK, Rutherford SA,
694   Lacy DB, Melnyk RA. 2020. Intestinal bile acids directly modulate the structure and
695   function of *C.difficile* TcdB toxin. Proc Natl Acad Sci U S A 117:6792-6800.
696   37.   Landenberger M, Nieland J, Roeder M, Nørgaard K, Papatheodorou P, Ernst K, Barth H.
697   2021. The cytotoxic effect of *Clostridioides difficile* pore-forming toxin CDTb. Biochim
698   Biophys Acta Biomembr 1863:183603.
699   38.   Aktories K, Schwan C, Jank T. 2017. *Clostridium difficile* toxin biology. Annu Rev Microbiol
700   71:281-307.
701   39.   Kroh HK, Chandrasekaran R, Zhang Z, Rosenthal K, Woods R, Jin X, Nyborg AC, Rainey GJ,
702   Warrener P, Melnyk RA, Spiller BW, Lacy DB. 2018. A neutralizing antibody that blocks
703   delivery of the enzymatic cargo of *Clostridium difficile* toxin TcdB into host cells. J Biol
704   Chem 293:941-952.
705   40.   Schwan C, Stecher B, Tzivelekidis T, van Ham M, Rohde M, Hardt WD, Wehland J, Aktories
706   K. 2009. *Clostridium difficile* toxin CDT induces formation of microtubule-based
707   protrusions and increases adherence of bacteria. PLoS Pathog 5:e1000626.
708   41.   Schwan C, Kruppke AS, Nolke T, Schumacher L, Koch-Nolte F, Kudryashev M, Stahlberg H,
709   Aktories K. 2014. *Clostridium difficile* toxin CDT hijacks microtubule organization and

710 reroutes vesicle traffic to increase pathogen adherence. Proc Natl Acad Sci U S A
711 111:2313-8.

42. Paredes-Sabja D, Cid-Rojas F, Pizarro-Guajardo M. 2022. Assembly of the exosporium layer in Clostridioides difficile spores. Curr Opin Microbiol 67:102137.

43. Pizarro-Guajardo M, Calderon-Romero P, Paredes-Sabja D. 2016. Ultrastructure Variability of the Exosporium Layer of *Clostridium difficile* Spores from Sporulating Cultures and Biofilms. Appl Environ Microbiol 82:5892-8.

44. Pizarro-Guajardo M, Calderón-Romero P, Romero-Rodríguez A, Paredes-Sabja D. 2020. Characterization of Exosporium Layer Variability of C*lostridioides difficile* Spores in the Epidemically Relevant Strain R20291. Front Microbiol 11:1345.

45. Pizarro-Guajardo M, Olguin-Araneda V, Barra-Carrasco J, Brito-Silva C, Sarker MR, Paredes-Sabja D. 2014. Characterization of the collagen-like exosporium protein, BclA1, of *Clostridium difficile* spores. Anaerobe 25:18-30.

46. Lessa FC, Mu Y, Bamberg WM, Beldavs ZG, Dumyati GK, Dunn JR, Farley MM, Holzbauer SM, Meek JI, Phipps EC, Wilson LE, Winston LG, Cohen JA, Limbago BM, Fridkin SK, Gerding DN, McDonald LC. 2015. Burden of *Clostridium difficile* infection in the United States. N Engl J Med 372:825-34.

47. Kelly CP. 2012. Can we identify patients at high risk of recurrent *Clostridium difficile* infection? Clin Microbiol Infect 18:21-7.

48. Zhang S, Palazuelos-Munoz S, Balsells EM, Nair H, Chit A, Kyaw MH. 2016. Cost of hospital management of Clostridium difficile infection in United States-a meta-analysis and modelling study. BMC Infect Dis 16:447.

49. Nikitas G, Deschamps C, Disson O, Niault T, Cossart P, Lecuit M. 2011. Transcytosis of *Listeria monocytogenes* across the intestinal barrier upon specific targeting of goblet cell accessible E-cadherin. J Exp Med 208:2263-77.

50. Gessain G, Tsai YH, Travier L, Bonazzi M, Grayo S, Cossart P, Charlier C, Disson O, Lecuit M. 2015. PI3-kinase activation is critical for host barrier permissiveness to *Listeria monocytogenes*. J Exp Med 212:165-83.

51. Pentecost M, Otto G, Theriot JA, Amieva MR. 2006. *Listeria monocytogenes* invades the epithelial junctions at sites of cell extrusion. PLoS Pathog 2:e3.

52. Pentecost M, Kumaran J, Ghosh P, Amieva MR. 2010. *Listeria monocytogenes* internalin B activates junctional endocytosis to accelerate intestinal invasion. PLoS Pathog 6:e1000900.

53. Singh B, Su YC, Riesbeck K. 2010. Vitronectin in bacterial pathogenesis: a host protein used in complement escape and cellular invasion. Mol Microbiol 78:545-60.

54. Henderson B, Nair S, Pallas J, Williams MA. 2011. Fibronectin: a multidomain host adhesin targeted by bacterial fibronectin-binding proteins. FEMS Microbiol Rev 35:147-200.

55. Hamzaoui N, Kernéis S, Caliot E, Pringault E. 2004. Expression and distribution of beta1 integrins in in vitro-induced M cells: implications for *Yersinia* adhesion to Peyer's patch epithelium. Cell Microbiol 6:817-28.

56. Clark MA, Hirst BH, Jepson MA. 1998. M-cell surface beta1 integrin expression and invasin-mediated targeting of *Yersinia pseudotuberculosis* to mouse Peyer's patch M cells. Infect Immun 66:1237-43.

753   57.   Diaz-Gonzalez F, Milano M, Olguin-Araneda V, Pizarro-Cerda J, Castro-Cordova P, Tzeng
754   SC, Maier CS, Sarker MR, Paredes-Sabja D. 2015. Protein composition of the outermost
755   exosporium-like layer of *Clostridium difficile* 630 spores. J Proteomics 123:1-13.
756   58.   Kelly CP, Poxton IR, Shen J, Wilcox MH, Gerding DN, Zhao X, Laterza OF, Railkar R, Guris
757   D, Dorr MB. 2020. Effect of Endogenous *Clostridioides difficile* Toxin Antibodies on
758   Recurrence of *C. difficile* Infection. Clin Infect Dis 71:81-86.

Figure Legends

Figure 1. Distribution of accessible and total E-cadherin in undifferentiated and differentiated Caco-2 cells. 3D projection of a representative confocal micrograph of healthy *A*, undifferentiated, and *B*, differentiated Caco-2 cells immunostaining for accessible E-cadherin. Accessible E-cadherin (shown as acc Ecad) is in green, F-actin in red, and nuclei in blue. *C*, Repartition of cells that were positive (Ecad+) or negative (Ecad−) stained for accessible E-cadherin. 3D projection of a representative confocal micrograph of a healthy *D*, undifferentiated and *E*, differentiated Caco-2 cells immunostaining for total E-cadherin and *F*, Repartition of cells that were positive (Ecad+) or negative (Ecad−) stained for total E-cadherin. Fluorophores colors were digitally reassigned for a better representation. Micrographs are representative of 6 fields of 2 independent experiments. Scale bar, 20μm. Error bars indicate mean ± SEM. Statistical analysis was performed by two-tailed unpaired Student's *t*-test, ns, $p > 0.05$; ****, $p \leq 0.0001$.

Figure 2. The accessibility of E-cadherin increases in TcdA and TcdB intoxicated IECs. Differentiated TcdA/TcdB intoxicated Caco-2 cells for 3, 6, or 8 h in DMEM without serum. As a control, cells were incubated with DMEM without serum. Non-permeabilized cells were stained for accessible E-cadherin (shown as acc Ecad; green), permeabilized, and total E-cadherin was stained (shown as total Ecad; red) and nuclei (blue). *A*, Representative confocal microscopy images 3D (XY) projection of control cells (left) and TcdA/TcdB intoxicated cells for 8h (right); below the orthogonal view (XZ; accessible E-cadherin, total E-cadherin and merged) of the dotted line of the XY. Representative confocal images for 3h and 8h not shown. Relative Fl. Int. measured as the sum of each z-step of *B*, accessible E-cadherin, and its abundance in the *C*, apical side, or *D* in the basal side of the cell. Relative Fl.Int. of *E*, total E-cadherin, its abundance in the *F*, apical side,

and the *G*, basal side of the cell. Controls were set 100%. Error bars indicate the mean ± SEM

from at least 9 fields (*n* = 3). *H*, immunoblotting anti non-glucosylated Rac1 and total Rac1 of cell

lysates of undifferentiated and differentiated Caco-2 cells intoxicated with TcdA/TcdB for 3, 6 or

8h. For undifferentiated in non-glucosylated Rac1 was evaluated, then stripped and membranes

were tested for total Rac1. Meanwhile for differentiated cells, the same sample was solved in 2

independent membranes simultaneously and were independently tested for non-glucosylated and

total Rac1. Western blot are representative of 3 independent experiment. Statistical analysis was

performed by two-tailed unpaired Student's *t*-test, ns, *p* > 0.05; ****, *p* < 0.0001. Bars, 5μm.

Figure 3. TcdA and TcdB increase *C. difficile* spore interaction with IECs.

Differentiated TcdA/TcdB intoxicated Caco-2 cells for 8h or incubated with DMEM without

serum as control and then, were washed with PBS and infected for 3h at an MOI 10 with *C. difficile*

spores. Unbound spores were rinsed off. Non-permeabilized cells were stained for adhered spores.

A, Representative epifluorescence microscopy images of TcdA and TcdB intoxicated cells (or

DMEM as control) for 8h then washed and infected with *C. difficile* spore. *B*, *C. difficile* spore

adherence relative in *B*, undifferentiated and *C*, differentiated cells. In bars, each dot represents

one independent well from 2 independent experiment, Per well, a total of 5 fields were analyzed.

Error bars indicate the mean ± SEM.. Statistical analysis was performed by unpaired *t*-test, ns, *p* >

0.05. Bars, 10 μm.

Figure 4. E-cadherin of IECs interacts with *C. difficile* spores increasing in TcdA and TcdB

intoxicated cells.

Differentiated Caco-2 cells were intoxicated with TcdA and TcdB for 8 h at 37 °C in DMEM

without serum. As a control, cells were incubated with DMEM without serum. Then cells were

infected with *C. difficile* spores for 1 h at 37 °C. Non-permeabilized cells were stained for

accessible E-cadherin (shown as acc Ecad; green), spores (red), and nuclei (blue). *A*,

Representative 3D projection confocal micrograph of healthy cells infected with *C. difficile* spores

(Ctrl). On the right, magnifications of representative *C. difficile* spores associated or not with E-

cadherin. *B*, *C,* Plot profile Fl.Int. of *C. difficile* spores (red line) and accessible E-cadherin (shown

as Ecad, green line) performed in the white line of *A*. *D*, Representative 3D projection confocal

micrograph of TcdA and TcdB intoxicated cells infected with *C. difficile* spores. On the right

magnifications of representative *C. difficile* spores associated or not with E-cadherin. *E, F*, Plot

profile Fl.Int. of *C. difficile* spores (red) and accessible E-cadherin (shown as Ecad, in green)

performed in the white line of *D*. Repartition of spores that were positive (Ecad+) or negative

(Ecad−) associated with fluorescence signal for accessible E-cadherin was shown as the average

associated/non-associated spores with E-cadherin for each field, a total of ~500 spores were

analyzed. Fluorophores were digitally reassigned for a better representation. Micrographs are

representative of 6 independent fields of 3 different experiments. Error bars indicate mean ± SEM.

Statistical analysis was performed by two-tailed unpaired Student's *t*-test, ns, $p > 0.05$; **, $p \leq$

0.01; ***, $p \leq 0.001$. Bars, 10μm.

Figure 5. E-cadherin blocking antibodies decrease the spore interaction with IECs.

Undifferentiated, differentiated, and polarized Caco-2 cells were infected with *C. difficile* spores

preincubated with FBS in the presence of anti-E-cadherin or non-immune IgG as control for 2 h at

37 °C. *A*, adherence, and *B*, internalization of *C. difficile* spores in undifferentiated Caco-2 cells

or *C*, adherence and *D*, internalization in differentiated Caco-2. *E, C. difficile* spore adherence in

polarized Caco-2 cells. Data showed in each panel are normalized to 0 µg mL^{-1} of antibody.

The data are the mean of 6 independent wells collected in 2 independent experiments. In bars, each

dot represents one independent well collected from 2 independent experiments. For each bar, a

total of ~1,300 spores were analyzed. Error bars indicate mean ± SEM. Statistical analysis was

performed by two-tailed unpaired Student's *t*-test, ns, *p* > 0.05; ****, *p* < 0.0001.

Figure 6. E-cadherin competitively reduces spore adherence to healthy and intoxicated

IECs.

A Undifferentiated and differentiated Caco-2 cells where infected for 3 h at a MOI of 10 with *C.*

difficile spores pretreated with 1, 2.5, 5, 10 and 25µg/mL of human purified E-cadherin, and

adhered cells where identified and quantified as described in the Method section. *B*. TcdA/TcdB

intoxicated undifferentiated and differentiated Caco-2 cells where infected for 3 h at a MOI of 10

with *C. difficile* spores pretreated with 25µg/mL of human purified E-cadherin, and adhered cells

where identified and quantified as described in the Method section. The data are the mean of 6

independent wells collected in 2 independent experiments. Adherence of *C. difficile* spores

preincubated with E-cadherin (or DMEM as control) in 8 h-TcdA/TcdB intoxicated cells. The

data are the mean of 6 independent wells collected in 2 independent experiments.

Figure 7. E-cadherin interacts with *C. difficile* spore.

A, Solid-phase binding assay of *C. difficile* R20291 spores or BSA as control with increasing

concentrations (i.e., 0.2, 2.0, 20 and 200 nM) of human purified E-cadherin. The data represented

are the mean of 9 independent wells collected in 3 independent experiments. *B*, Representative

phase-contrast micrograph, immunofluorescence against E-cadherin, and immunofluorescence

plot profile of 4×10^7 *C. difficile* spores that were incubated with increasing concentrations (i.e.,

1, 2.5, 5, 10, 25 µg mL^{-1}) of human purified E-cadherin for 1 h at 37 °C. *C*, Quantitative analysis

of Fl.Int. *D*, Distribution of E-cadherin fluorescence intensity of *C. difficile* incubated with

increasing concentrations (i.e., 1, 2.5, 5, 10, 25 µg mL^{-1}) of human purified E-cadherin. A total of

~450 spores were analyzed for each condition. Error bars indicate mean ± SEM. Statistical analysis

was performed by two-tailed unpaired Student's *t*-test, ns, $p > 0.05$; *, $p < 0.05$; ****, $p < 0.0001$.

Figure 8. Immunoelectron microscopy of E-cadherin-gold complex binds to *C. difficile*

spores.

A,B TEM of *C. difficile* spores incubated with 10 µg mL^{-1} E-cadherin or without E-cadherin

(negative control). Then incubated with anti-E-cadherin and anti-mouse-gold 12 nm complex.

White arrows denote the mouse anti-E-cadherin and anti-mouse-gold 12 nm complex. Spores

where stratified in thick and thin exosporium spores. Lower panels are magnification of the black

squares highlighting immunogold spots, Quantification of the *C. difficile* spore associated with E-

cadherin ("−" and "+" denote negative or positive association, respectively). *C,D* Classification of

E-cadherin positively associated *C. difficile* spores according to the morphology of the spore, thin

or thick exosporium, represented as the percentage of the total E-cadherin positive spores. Figures

show representative spores with a thin or thick exosporium layer, and are representative of 2

independent experiments. A total of 100 spores were quantified. *E* solid-phase binding assay of

human E-cadherin to *C. difficile* spores of 630 wild-type and isogenic mutant strains of *bclA1*,

bclA2 and *bclA3* or BSA as a control with increasing concentrations (i.e., 0.2, 2.0, 20 and 200 nM)

of human purified E-cadherin. The data represented are the mean of 9 independent wells collected

in 3 independent experiments. Statistical analysis was performed by ONE-WAY ANOVA

followed by Bonferroni for multicomparisons, ns, $p > 0.05$; *, $p < 0.05$, ****, $p < 0.0001$. Colors

correspond to the mutant strain being compared to wild-type.

Supplementary figures review only.

Supplementary confocal micrographs of Fig. 1.

These are supplementary figures from the experiment detailed in Fig. 4. Differentiated Caco-2

cells intoxicated with 600 pM of TcdA and TcdB for 3, 6, or 8 h in DMEM without serum. As a

control, cells were treated with DMEM without serum. Non-permeabilized cells were stained for

accessible E-cadherin (shown as acc Ecad) in green, then permeabilized and stained total E-

cadherin (total Ecad) shown in red nuclei shown in blue. Representative confocal microscopy

images 3D (XY) projection of intoxicated cells for 3 h (left) or 6 h (right) with its control; below

a magnified slide (XY), and the orthogonal view (XZ). Representative images from at 9 fields (*n*

= 3). Bars, 5μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplementary Fig 1