1
10 **Small proline-rich proteins (SPRRs) are epidermally produced antimicrobial proteins**
11 **that defend the cutaneous barrier by direct bacterial membrane disruption**
12
13

14 Chenlu Zhang[1*], Zehan Hu[2,3], Abdul G. Lone[1], Methinee Artami[1], Marshall Edwards[1], Christos
15 C. Zouboulis[4], Maggie Stein[1], Tamia A. Harris-Tryon[1,2*#]
16
17
18
[1]Department of Dermatology, University of Texas Southwestern Medical Center, Dallas, TX,
75390
[2]Department of Immunology, University of Texas Southwestern Medical Center, Dallas, TX,
75390
[3]State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of
Metabolic & Developmental Sciences, School of Life Sciences and Biotechnology, Shanghai
Jiao Tong University, Shanghai, China, 200240
[4]Department of Dermatology, Venereology, Allergology and Immunology, Dessau Medical
Center, Brandenburg Medical School Theodore Fontane and Faculty of Health Sciences
Brandenburg, 06847 Dessau, Germany.
29
30
31
32 *Correspondence to:
33 tamia.harris-tryon@utsouthwestern.edu
34 chenlu.zhang@utsouthwestern.edu

Abstract

Human skin functions as a physical barrier, preventing the entry of foreign pathogens while also accommodating a myriad of commensal microorganisms. A key contributor to the skin landscape is the sebaceous gland. Mice devoid of sebocytes are prone to skin infection, yet our understanding of how sebocytes function in host defense is incomplete. Here we show that the small proline-rich proteins, SPRR1 and SPRR2 are bactericidal in skin. SPRR1B and SPPR2A were induced in human sebocytes by exposure to the bacterial cell wall component lipopolysaccharide (LPS). Further, LPS injected into mouse skin triggered the expression of the mouse SPRR orthologous genes, *Sprr1a* and *Sprr2a*, through stimulation of MYD88. Both mouse and human SPRR proteins displayed potent bactericidal activity against MRSA (methicillin-resistant *Staphylococcus aureus*), *Pseudomonas aeruginosa* and skin commensals. Thus, *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice are more susceptible to MRSA and *Pseudomonas aeruginosa* skin infection. Lastly, mechanistic studies demonstrate that SPRR proteins exert their bactericidal activity through binding and disruption of the bacterial membrane. Taken together, these findings provide insight into the regulation and antimicrobial function of SPRR proteins in skin and how the skin defends the host against systemic infection.

Introduction

The skin is the human body's largest organ, with direct contact with the external environment [1]. As a result, the skin surface continuously encounters a diverse microbial community including bacteria, fungi, viruses and parasites [2–4]. When host defense is impaired, skin infection results. Thus, skin and soft tissue infections pose a considerable public health threat [5]. The majority of infections of the skin are caused by *Staphylococcus aureus* (*S. aureus*) [6]. Additionally, *Pseudomonas aeruginosa* (*P. aeruginosa*) infections in burn patients are one of the most common causes of mortality [7]. Adding to the challenges of treating infections posed by these pathogens has been the development of antibiotic resistant strains of bacteria such as methicillin resistant *Staphylococcus aureus* (MRSA) [8].

Skin antimicrobial proteins (AMPs) play an essential role in defending the host from the invasion of pathogens [9,10]. Mammalian antimicrobial proteins are evolutionarily ancient immune effectors that rapidly kill bacteria by targeting bacterial cell wall or cell membrane structures [11,12]. Several distinct antimicrobial protein families, such as β-defensins, cathelicidins, resistin and S100 proteins, have been identified and characterized in skin [13,14]. However, we still have a limited understanding of the arsenal of antimicrobial proteins expressed by the skin, the regulatory networks that control the expression of AMPs, and how AMPs function to protect mammalian skin surfaces. Even less is known about the contribution that skin appendages make to host defense.

Sebaceous glands (SGs) are specialized epithelial cells that cover the entire skin surface except the palms and soles. SGs excrete a lipid-rich and waxy substance called sebum to the skin surface [15–17]. SGs are believed to contribute to the antimicrobial functions of the skin [13], yet few

current studies have examined the role of SGs in skin host defense. Here we show the impact

of the bacterial cell wall component lipopolysaccharide (LPS) on human sebocytes and

demonstrate that bacterial lipoproteins stimulate the expression of members of the small

proline-rich protein (SPRR) family. SPRR proteins were originally identified as markers of

terminal differentiation and function as substrates of transglutaminase in the crosslinked

cornified envelope present at the skin surface [18,19]. In this study, we demonstrate that SPRR

proteins function as antimicrobial proteins. *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice are more susceptible to

MRSA and *P. aeruginosa* skin infection, revealing that SPRR proteins protects against

pathogenic bacterial infections of the skin. We also show mechanistically that the bactericidal

activity of SPRR proteins is mediated through the binding and disruption of bacterial

membranes. Taken together our findings support a novel antimicrobial function of SPRR

proteins in skin.

102
103 **Results**
104

SPRR proteins are induced by LPS in human sebocytes

As a first step towards understanding the role of the sebaceous gland in skin host defense, we

performed whole transcriptome RNA-sequencing to compare transcript abundances in human

immortalized sebaceous gland cells (SZ95) treated with lipopolysaccharide (LPS) to untreated

sebocytes (Fig 1A). LPS, a lipoprotein which coats the surface of Gram-negative bacteria, had

broad impacts on gene expression in human sebocytes, including the increased expression of

inflammatory cytokines, chemokines and antimicrobial proteins (Fig 1A). Notably, three

members of the small proline rich family of proteins (SPRR) were markedly upregulated in

sebocytes after LPS treatment (Fig 1A). To confirm that the expression of SPRR genes were

induced by LPS, we used quantitative reverse transcription PCR (qRT-PCR) to analyze the

change of the human *SPRR1B, SPRR2A, SPRR2D* transcript abundance in SZ95 cells.

Consistent with our RNA-seq results, LPS-treated SZ95 cells displayed significant increase in

the relative expression of *SPRR* family genes compared to vehicle treated cells (Fig 1B).

Additionally, dose response and time course experiments revealed that 1μg LPS treatment for

16 hours is optimal for induction of *SPRR* family genes expression (fig S1). Consistent with

the qRT-PCR results, SPRR proteins expression also increased after LPS treatment in SZ95

cells (Fig 1C, 1D). LPS is a pattern associated molecular pattern (PAMP) known to trigger gene

transcription through the pattern recognition receptor, Toll-like receptor (TLR) 4 [20]. We

therefore tested a panel of PAMPs to see if other foreign stimuli would trigger the expression

of the *SPRR* genes in human sebocytes. Interestingly, two other lipoproteins, Pam3CSK4 and

FSL-1, stimulate the expression of *SPRR1B*, *SPRR2A,* and *SPRR2D* genes in human sebocytes

(fig S2), indicating that sebocytes respond to bacterial lipoproteins by expressing SPRR1 and

SPRR2 proteins.

Next, we decided to further explore whether heat-inactivated bacteria could induce the

expression of SPRR genes in sebocytes. We used qRT-PCR to analyze the change of the human

SPRR2A transcript abundance in SZ95 cells after various heat-inactivated bacteria treatment.

Interestingly, only the Gram-negative bacteria tested could trigger SPRR2A expression (fig

S3). Taken together, these data establish that bacterial lipoproteins triggers the expression of

SPRR genes in human sebocytes.

SPRR proteins are upregulated by the injection of LPS in mouse skin

Next, we sought to examine whether commensal skin microbiota colonization could induce the

expression of *Sprr* family genes *in vivo*. *Sprr1b* is not expressed in mouse skin, so we tested

the ability of the microbiota to stimulate the expression of the mouse orthologs, *Sprr1a* and

*Sprr2*a. In contrast to what we observed in human sebocytes in culture, *Sprr* gene expression

was similar between germ free and conventional mice (Fig 2A), indicating that bacterial

colonization alone is insufficient to trigger increased expression of *Sprr* genes in mouse skin.

As sebaceous glands cells are located deeper in the dermis, we next chose to test whether

intradermal injection of LPS – mimicking wounding and infection conditions — could

stimulate *Sprr1a* and *Sprr2a* expression in mouse skin. Indeed, qRT-PCR analysis revealed that

intradermal injection of LPS stimulates the expression of *Sprr1a* and *Sprr2a* in mouse skin (Fig

2B). Canonically, LPS stimulates gene expression through activation of TLR4 and the TLR

signaling adaptor MYD88. Thus, LPS injected into the skin of *Myd88*$^{-/-}$ mice did not trigger

increased expression of the *SPRR* genes (Fig 2B), indicating that LPS induces SPRR protein

expression through the TLR-MYD88 signaling pathway. Consistent with these data, both live

and heat-inactivated *Escherichia coli* (*E. coli*) bacteria were able to induce SPRR expression,

suggesting that stimulation of *Sprr1a* and *Sprr2a* in mouse skin does not depend on the presence

of bacterial metabolites (Fig 2C). As has been shown by others [21], we also found that *Sprr1a*

and *Sprr2a* transcript abundance is locally induced by crosshatch skin wounding in mice (Fig

2D).

Next, to investigate the expression pattern of SPRR proteins in mouse skin, we performed

immunofluorescence experiment using a commercial antibody specific to SPRR1A. SPRR1A

is expressed in the sebaceous gland and non-specifically at the junction of the sebaceous gland

and hair follicle (Fig 2E). This suggests SPRR proteins expressed in the sebocyte may be

secreted to the hair canal and are present at the skin surface. Since our immunofluorescence

images and prior studies in keratinocytes indicated that SPRR1A and SPRR2A are expressed

by keratinocytes [22,23], we sought to study whether LPS could stimulate SPRRs expression in

keratinocyte cells. We first made use of hTert cells originally derived from human foreskin

keratinocytes and immortalized by expression of human telomerase and mouse Cdk4 and

treated the hTert cells with LPS for 24 hours. The qRT-PCR results showed that LPS cannot

induce the expression of *SPRR* family genes in human keratinocytes (fig S4). To further

confirm this result, we isolated primary mouse keratinocytes from neonatal mice, and cultured

them under both low and high calcium condition. Under low calcium conditions the

keratinocyte cell can proliferate but won't differentiate into a stratified layer. High calcium did

initiate the differentiation process and triggered increased expression of both mouse *Sprr1a* and

Sprr2a (fig S5). However, LPS did not stimulate the expression of *Sprr* genes in mouse

keratinocyte under either condition (fig S5). Taken together, these data show that SPRR

proteins are unregulated by LPS in human sebocytes in culture and by LPS dermal injection in

mouse skin.

SPRR proteins are bactericidal against skin pathogens and commensals

The SPRR family proteins have highly conserved amino acid sequences with numerous

cysteine and proline repeats. The antimicrobial properties of proline rich proteins have not been

described in humans. However, in insects and lower vertebrates, proteins rich in cysteines and

prolines have been shown to kill microbes [24,25]. Based on these findings, we hypothesized that

the SPRR proteins might function in cutaneous host defense as antimicrobial proteins. To test

the antimicrobial ability of SPRR proteins, we produced recombinant human SPRR1B,

SPRR2A and mouse SPRR1A protein in the baculovirus insect cell expression system and

further purified to homogeneity using size exclusion chromatography (fig S6). We then tested

the antimicrobial function of SPRR proteins against skin commensals and pathogens *in vitro.*

When bacteria were exposed to SPRR proteins, we observed a marked dose-dependent

reduction in the viability of *Staphylococcus epidermidis* (*S. epidermidis*), Methicillin Resistant

Staphylococcus aureus (MRSA), and *P. aeruginosa* (Fig 3A). with is a more than 90% decline

in bacterial viability after 2-hour exposure to 2.5 µM of SPRR proteins (Fig 3B). In contrast,

SPRR proteins had no impact on the viability of the prototypical Gram-negative bacteria

Escherichia coli (fig S7). Lastly, to visualize morphological changes of *P. aeruginosa* after

exposure to SPRR proteins, we used transmission electron microscopy and observed distinct

bacterial cell membrane damage and cytoplasmic leakage (Fig 3C).

Since SPRR proteins exhibit similar spectrum of bactericidal activity, we chose to use human

SPRR1B and mouse SPRR1A protein to further delineate the mechanism underlying its

bactericidal activity. We first used the propidium iodide (PI) uptake assay to assess the capacity

of SPRR proteins to permeabilize bacterial membranes. Human SPRR1B and mouse SPRR1A

proteins promoted the dose-dependent uptake of the membrane-impermeant small molecule

dye, PI, by *P. aeruginosa* (Fig 3D). Next, we examined whether SPRR proteins can directly

bind lipid molecules by incubating proteins with lipid strips dotted with different lipids. Both

human SPRR1B and mouse SPRR1A bind to lipids bearing negatively charged lipid head

groups, but not to neutral lipids (Fig 3E). These data are consistent with the idea that most

bacterial cell membranes are negatively charged and thus susceptible to SPRR1 binding. Lastly,

to determine whether SPRR proteins disrupt bacterial membrane, we incubated human

SPRR1B and mouse SPRR1A protein with liposomes loaded with the fluorescent dye

carboxyfluorescein (CF, ~10 Å Stokes diameter) and quantified dye release after exposure.

Phosphatidylcholine (PC)/ phosphatidylserine (PS) liposomes are composed of 85% of the

neutral lipid PC and 15% of the negatively charged lipid PS —similar to that of bacterial

membranes. Both human SPRR1B and mouse SPRR1A protein induced rapid dye efflux from

negatively charged PC/PS liposomes in a dose-dependent manner (Fig. 3F). These findings

confirm that SPRR proteins exert bactericidal activity through membrane disruption. Taken

together, these results suggest that SPRR proteins are bactericidal and compromise the

membrane of specific skin bacteria including the pathogen *P. aeruginosa*.

SPRR proteins limit MRSA and *P. aeruginosa* infection

Given that SPRR1A and SPRR2A protein exhibited bactericidal activity against a panel of skin

pathogens *in vitro* (Figure 3), we predicted that the removal of these proteins might limit MRSA

and *P. aeruginosa* skin infection *in vivo*. To test this hypothesis, we created mice lacking

SPRR1A and SPRR2A. We used a CRISPR/Cas9-mediated in vitro fertilization method to

delete the entire mouse *Sprr1a* locus of the *Sprr2a*$^{-/-}$ mice [26] and verified that SPRR1A and

SPRR2A are absent in *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$mice skin at both the transcript and protein level (fig

S8). *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$ mice were healthy and showed normal skin barrier with no signs of

immune infiltration and no significant change in transepidermal water loss (TEWL) (fig S9).

To test the *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$ mice for susceptibility to infection, we inoculated the skin of

wild-type and *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$ mice with 1 x 10^6 CFU of a bioluminescent strain of MRSA

by topical application. The luminescence was monitored daily with serial photography and

quantified for 3 consecutive days (Fig 4A). Mice devoid of SPRR1A and SPRR2A expression

had greater bacterial burdens in the first 3 days of MRSA infection (Fig 4B, 4C). Next, we

aimed to assess the susceptibility of *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$ mice to *P. aeruginosa* skin infection.

Since damaged or burned skin has increased risk of *P. aeruginosa* infection [7], we first

introduced skin wounding in mice by superficial crosshatching method, and then inoculated 1

x 10^6 CFU of a bioluminescent strain of *P. aeruginosa* at the wounding site. Intravital imaging

and quantification revealed that *Sprr1a*$^{-/-}$;*Sprr2a*$^{-/-}$mice are more susceptible to *P. aeruginosa*

skin infection (Fig 4D-4F). Collectively, we confirm that SPRR proteins limit MRSA and *P.*

aeruginosa infection in mice skin.

233
234   **Discussion**
235

Skin is in direct contact with the microbe-filled outer world. Thus, defending the host from

invasion by pathogens like *S. aureus* and *P. aeruginosa* is one of the chief functions of the skin

[6,27]. In this study we have identified SPRR1 and SPRR2 proteins as a previously uncharacterized

group of antibacterial proteins expressed by keratinocytes and sebaceous gland cells, which can

rapidly kill pathogens through membrane disruption and limit bacterial skin infection.

Several earlier studies have shown that SPRR proteins are upregulated in the GI tract, urinary

tract and the airway after exposure to stress and other inflammatory stimuli [28]. Further, Hu et

al. recently revealed that SPRR2A has antimicrobial actions in the gastrointestinal tract during

helminth infection [26]. Our data highlight a previously unappreciated role of SPRR family

proteins in skin immune defense, demonstrating that both SPRR1 and SPRR2 are bactericidal

proteins induced in sebocytes by the bacterial cell wall component LPS (Fig 1). The major role

of sebaceous glands in mammals is to produce sebum, a mixture of non-polar lipids and proteins

required for normal skin ecology [29]. Sebum secretion can also act as a delivery system for AMPs

[30]. Though beneficial to the host, AMPs can damage mammalian membranes, so confining

expression of AMPs to the non-viable parts of skin through sebum delivery is optimal. Sebum

also excretes AMPs to the surface within an acidic milieu, which is often required for AMP

activity [31].

Interestingly, our data also show that bacterial colonization of germ-free mice did not

induce expression of SPRR proteins *in vivo* (Fig 2). Further, LPS was not able to stimulate the

production of SPRR proteins in mouse keratinocytes (fig S4-5). Taken together, our findings

suggest that deeper penetration of bacterial stimuli may be required for MYD88 mediated

SPRR protein production in skin. These finding are distinct from what has been observed in the

gastrointestinal tract, where colonization of germ-free mice is sufficient to stimulate SPRR2A

expression [32]. As SPRR family proteins also function as cross-linking proteins in terminal

differentiation of keratinocytes and formation of the cornified cell envelope (Candi et al., 2005),

there are likely other regulatory networks that control SPRR proteins expression in skin in

response to wounding and other stimuli. Additional studies with sebaceous gland-specific

deletion of TLRs and MYD88 will be required to confirm that stimulation of SPRR proteins by

LPS requires interaction with TLRs on sebocytes.

In this study, we also reveal that SPRR family proteins have potent bactericidal activity

against *Staphylococcal* species and the gram-negative bacteria *P. aeruginosa* (Fig 3A-C), with

effective inhibition at low micromolar concentrations. Further, our biochemical data show that

mouse and human SPRR1 proteins interact with negatively charged lipid membranes,

suggesting broad spectrum antimicrobial capability of these proteins (Fig 3D-F). Additionally,

skin devoid of SPRR1 and SPRR2 is more susceptible to skin infection by MRSA and

Pseudomonas aeruginosa, both pathogens that cause skin infections requiring hospitalization

(Fig 4). Treatments for these infections are currently limited. SPRR proteins might be

promising alternatives to traditional antibiotics used to cure MRSA or *P. aeruginosa* infection.

Human SPRR proteins are 6–18 kDa in size and comprise four subclasses (SPRR1,

SPRR2, SPRR3, and SPRR4) with a similar structural organization. Among them, two SPRR1

and seven SPRR2 proteins are characterized with a much higher homogeneity, as they contain

a similar consensus repeat sequence [18]. Our findings that SPRR family proteins have

antimicrobial function may have implications for other proline-rich proteins expressed at other

mucosal sites. Altogether, these findings expand our current understanding of the molecules

involved in cutaneous host defense and provide insight into how the sebaceous gland

contributes to the fight against skin infection.

282
283

Materials and Methods

Mice

All animal protocols were approved by the Institutional Animal Care and Use Committees of

the University of Texas (UT) Southwestern Medical Center. Age- and sex-matched mice 8-12

weeks old were used for all experiments. Conventionally raised C57BL/6 wild type (WT),

Myd88$^{-/-}$ and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* on C57BL/6 background were bred and maintained in the

specific pathogen free (SPF) barrier facility at the UT Southwestern Medical Center on standard

chow diet. The generation of *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice is described below. Germ-free (GF)

C57BL/6 mice were bred and maintained in flexible film vinyl isolators in the gnotobiotic

mouse facility at UT Southwestern with autoclaved diet (Hooper Lab Diet 6F5KAI, Lab Diet,

St. Louis, MO) and autoclaved nanopore water. All mice were housed under a 12 hour light-

dark cycle, and mice were randomly assigned to treatment groups.

Sprr1a$^{-/-}$;Sprr2a$^{-/-}$ mice were generated using CRISPR/Cas9-mediated *in vitro* fertilization

with a guide RNA targeting regions upstream and downstream of the *Sprr1a* locus on the

basis of *Sprr2a$^{-/-}$* mice (Figure S8). *Sprr2a$^{-/-}$* mice were obtained with the permission of Dr.

Lora Hooper [26]. Guide RNAs were injected into fertilized female *Sprr2a$^{-/-}$* mice embryos

by the UT Southwestern Transgenic Core facility. The resulting litters were screened by

genomic sequencing to detect the deletion of *Sprr1a* and *Sprr2a* locus. Mice harboring the

deleted allele were backcrossed with WT to homozygosity.

Bacterial strains

Bacteria were grown in species-specific growth media: *Escherichia coli* (ATCC; ATC-PTA-

7555) and *Citrobacter rodentium* (ATCC51459) were grown in Luria Broth. *Pseudomonas*

aeruginosa (ATCC27853), bioluminescent *Pseudomonas aeruginosa* strain Xen05 (Perkin

Elmer 119228) and *Enterococcus faecalis* (ATCC29212) were grown in Brain Heart Infusion

Broth (BD Biosciences). Methicillin resistant *Staphylococcus aureus* (ATCC25923),

bioluminescent MRSA strain SAP430 *lux* [33], *Staphylococcus epidermidis* (clinical isolate),

Staphylococcus xylosus (clinical isolate) were grown in Tryptic Soy Broth.

Sebocyte cell culture and treatments

SZ95 cells are an immortalized human sebocyte cell line generated from the face of an 87-year-

old female and transformed with Simian virus 40. These cells were previously obtained from

Cristos Zouboulis [14]. SZ95 sebocytes were maintained in Sebomed Basal Medium (Fischer

Scientific NC9711618) supplemented with 10% fetal bovine serum (GeminiBio 100-106), 5

ng/ml human epidermal growth factor (ThermoFisher PHG0313) and 1% Antibiotic-

Antimycotic (Gibco 15240062). Cells were cultured in 5% CO_2 incubator at 37°C. Cells were

stimulated with 1μg/ml LPS (Sigma L4524). 16 hours post-stimulation cells were harvested

and analyzed as described below. For TLR agonists treatment, reagents in human TLR1-9

Agonist kit (InvivoGen, tlrl-kit1hw) were diluted to working concentration in PBS. 100ng/ml

Pam3CSK4, 1 x 10^8 cells/ml HKLM, 500ng/ml Poly (I:C), 500ng/ml Flagellin, 100ng/ml FSL-

1 and 1μg/ml Imiquimod were used for stimulation of SZ95 cells. For heat-inactivated bacteria

treatment, bacteria were grown to mid-logarithmic phase, spun down, washed and resuspended

in PBS. Bacteria were heat-inactivated by incubation at 95 °C for 20 minutes. Then 1 x10^8

cells/ml were used for each treatment.

Immunofluorescence microscopy

Mouse skin samples were fixed in formalin and embedded in paraffin by the UT Southwestern

histology core. Samples were deparaffined with xylene followed by rehydration with

decreasing concentrations of ethanol. Boiling in 10mM sodium citrate buffer with 0.2% tween

for 15 minutes for antigen retrieval. Slides were blocked with 10% FBS, 1% BSA and 1%

Triton X-100 in PBS, and then incubated with primary antibodies against mouse SPRR1A

(Thermo Fisher PA5-26062), mouse CD36 (R&D AF2519), rabbit IgG isotype control (Thermo

Fisher 02-6102) or goat IgG isotype control (Thermo Fisher 02-6202) using 1:100 dilutions at

4°C overnight. Secondary antibodies Alex Fluor 594 or Alexa Fluor 647 (Thermo Fisher) were

diluted 1:350 in blocking buffer and applied to slides for 1 hour at room temperature in the

dark. Slides were then washed with PBST (PBS with 0.2% tween) and mounted with DAPI

Fluoromount-G (Southern Biotechnology 0100-20). Images were captured using an Echo

Revolve 4 microscope.

Quantitative real-time PCR

RNA was extracted from cells or mouse skin using the RNAeasy Plus universal kit (Qiagen

73404). RNA was quantified by absorbance at 260 nm, and its purity was evaluated by the

ratios of absorbance at 260/280 nm. 2μg of RNA was used for cDNA synthesis (Thermofischer

4368814, High Capacity cDNA reverse transcription kit). Quantitative real-time PCR was

performed using PowerUp SYBR Green Gene Expression Assays (Thermofischer A25741) and

a QuantStudio 7 Flex Real-Time PCR System (Applied Biosystems). Relative expression

values were calculated using the comparative Ct ($\Delta\Delta$Ct) method, and transcript abundances

were normalized to GAPDH transcript abundance. The primer sequences are shown in table

S1.

Whole transcriptome sequencing and data analysis

RNA was extracted from SZ95 sebocyte cells using the RNAeasy Plus universal kit (Qiagen

73404). RNA quality was assessed using Agilent 2100 Bioanalyzer. Truseq RNA sample

preparation kit v2 (Illumina) was used for the preparation of sequencing libraries. Sequencing

was performed on an Illumina HiSeq 2500 for signal end 50 bp length reads. Sequence reads

were mapped against the hg19 genome using TopHat. For each gene, read counts were

computed using HTSeq-count and analyzed for differential expression using DESeq2.

Western blot

SZ95 cells were harvested by applying 200 μL of diluted 1x sample buffer (Thermo Fisher

39000) directly to a 6-well plate, scraping down the cell sample to disrupt the membranes, then

boiling for 15 minutes before loading. Equal amounts of protein were loaded onto a 4-20%

gradient SDS-PAGE and transferred to a PVDF membrane. After blocking with 5% milk in

TBST, the membranes were incubated with anti-SPRR1B antibody (Thermo Fisher PA5-

26062), anti-SPRR2A (Abcam ab125385) or anti-GAPDH (Abcam ab181602) at 4°C

overnight. Membranes were then incubated with anti-rabbit secondary antibodies conjugated

with HRP (Abcam). Membranes were visualized using a Bio-Rad ChemiDoc Touch system

and bands were quantified by Image Lab software.

Keratinocyte cell line culture and treatments

hTERT (ATCC CRL-4048) and primary mouse keratinocyte cells were cultured in KSFM

medium (Invitrogen, 37010022) supplemented with 0.05 mM $CaCl_2$ (Sigma, C7902), 0.05

μg/mL hydrocortisone (Sigma, H0888), 5 ng/mL EGF (Invitrogen, 10450-013), 7.5 μg/mL

bovine pituitary extract (Invitrogen, 13028-014), 0.5 μg/mL insulin (Sigma, I9278), 100 U/mL

penicillin, 100 μg/mL streptomycin, and 25 μg/mL of amphotericin B (Invitrogen, 10450-013).

Mouse primary keratinocytes were isolated through dispase digestion. Before digestion, the

subcutaneous fat was first removed from the mouse skin. Skin tissue from 3-5 days neonatal

mice was floated on 1 U/ml dispase (Corning, 354235) in HBSS (Gibco, 14170) for 16 h at 4°C

with the dermis side down. The next day, the skin was placed in a new dish with the epidermis

side down, and the epidermis was peeled and placed into a new dish with HBSS. After being

washed, the cells were collected into a new 15 mL tube. The epidermis was cut into small

pieces, resuspended in HBSS, gently pipetted up and down several times, and then combined

with the cells in the 15 mL tube. The cell solution was filtered with a 70 μm cell strainer,

centrifuged at 1,000 rpm for 4 minutes, and resuspended in complete KSFM medium. The cells

were gently washed once and seeded in culture dishes with complete KSFM medium. The

culture dishes were precoated with collagen (Advanced Biomatrix, 5005-B). Keratinocytes

from each mouse were seeded in one 10 cm dish, and fresh complete KSFM medium was

supplied after 24 hours. After 3-4 days, the primary keratinocytes reached approximately 80%

confluence under normal culture conditions in a 5% CO_2, 37°C incubator. For keratinocyte

differentiation, cells were treated with 1 mM $CaCl_2$ for 2 days. For treatment of keratinocyte

cells, 1 μg/mL LPS (Sigma L4524) were supplemented in the KSFM complete medium. 16

hours post-stimulation cells were harvested for qRT-PCR or Western blot analysis.

Intradermal injection of mice

The mouse dorsal hair was removed by shaving (Andis ProClip), followed by depilatory cream

(Nair™) one day before injection. LPS (Sigma L4524) was dissolved in PBS and further diluted

to a concentration of 1 mg/mL. For intradermal injection of LPS, each mouse was injected with

50 μL of LPS solution to the back skin. For intradermal injection of bacteria, various bacteria

were grown in species-specific media to mid-logarithmic phase, spun down, washed and

resuspended in PBS to a concentration of 1 x10^8 cells/mL. Bacteria were heat-inactivated by

incubation at 95 °C for 20 minutes. 100 μL of live or heat-inactivated bacteria was injected to

the mouse back skin. The site of injection was circled with permanent marker. After 8 hours,

mice were sacrificed and the injection site skin was analyzed.

Protein expression and purification

SPRR genes containing a C-terminal 6xHis tag were cloned into a pFastBac1 vector and

heterologously expressed in Sf9 cells (Thermo Fisher). One liter of cells (2.5×10^6 cells/mL)

was infected with 10 mL baculovirus at 28°C. Cells were cultured in suspension and harvested

48 hours after infection. Harvested cells were resuspended in the buffer containing 25 mM Tris-

HCl pH 8.0, 150 mM NaCl and 1 mM Phenylmethylsulfonyl fluoride (PMSF), and lysed by

sonication. The mixture was pelleted by centrifugation at 10,000g for 30 minute and the

supernatant was loaded onto a Ni^{2+} metal affinity column (Qiagen). The column was washed

with buffer containing 30 mM imidazole and the protein was eluted in buffer containing 300

mM imidazole. The eluate was concentrated in a 3K cutoff Amicon Ultra centrifugal device

(Millipore) and further purified by size exclusion chromatography on a Superdex 75™ 10/300

GL column (GE Healthcare Life Sciences) in standard assay buffer (10 mM MES, pH 6.0 and

25 mM NaCl).

Bacterial killing assays

Bacteria were grown in species-specific growth media as described above. 10 mL bacterial

cultures were grown to mid-logarithmic phase and then pelleted and washed twice in assay

buffer (10 mM MES, pH 6.0, 25 mM NaCl). Approximately 5×10^6 cells/mL bacteria were then

incubated at 37°C for 2 hours in assay buffer with varying concentrations of recombinant SPRR

protein or BSA (Gemini 700-106P). Surviving colony-forming units (CFU) were quantified by

dilution plating onto agar plates and calculated as a percentage of the remaining colonies in the

assay buffer only control sample.

Lipid strip assay

Membrane lipid strips (Echelon, P-6002) were used following the manufacturer's protocol.

Briefly, the lipid strips were blocked with blocking buffer (10 mM MES, pH 6.0, 25 mM NaCl,

2% BSA, and 0.05% Tween-20) for 1 hour at room temperature. Purified recombinant SPRR

proteins were diluted to 1 μg/mL in blocking buffer and incubated with the lipid strip overnight

at 4°C. After three washes with washing buffer (10 mM MES, pH 6.0, 25 mM NaCl and 0.05%

Tween-20), the lipid strip was sequentially incubated with anti-SPRR1B antibody and HRP-

conjugated secondary antibody. Dots were detected with ECL reagent (Bio-Rad, 1705060)

using a Bio-Rad ChemiDoc system.

Electron microscopy

For electron microscopy of bacteria, 10 mL *P. aeruginosa* cultures were grown to mid-

logarithmic phase and then pelleted and washed in 10 mL standard assay buffer (10 mM MES

pH 6.0 and 25 mM NaCl). Bacteria were then resuspended in 1 mL standard assay buffer.

Purified recombinant SPRR proteins was added at a final concentration of 5 μM to 300 μL of

resuspended bacteria and incubated at 37°C for 2 hours. Bacteria were centrifuged for 10

minutes at 16,000 g, resuspended in cross-linking reagent (4% paraformaldehyde and 4%

glutaraldehyde in 0.1 M sodium phosphate buffer, pH 7.4), and incubated overnight at 4°C.

After three washes, bacterial pellets were embedded in 3% agarose, sliced into small blocks (1

mm^3), and fixed with 1% osmium tetroxide and 0.8% potassium ferricyanide in 0.1 M sodium

phosphate buffer for 1.5 hours at room temperature. Cells were rinsed with water then stained

with 1% uranyl acetate in water for 1 hour. Cells were dehydrated with increasing

concentrations of ethanol, transitioned into propylene oxide, infiltrated with Embed-812 resin,

and polymerized overnight in a 60°C oven. Blocks were sectioned with a diamond knife

(Diatome) on Leica Ultracut 7 ultramicrotome (Leica Microsystems), collected onto copper

grids, and post-stained with 2% aqueous uranyl acetate and lead citrate. Images were captured

on a Tecnai G2 spirit transmission electron microscope (Thermo Fisher) equipped with a LaB6

source using a voltage of 120 kV.

Liposome disruption assay

Unilamellar liposomes were prepared using lipids from Avanti. 85% 1-palmitoyl-2-oleoyl-

sn-glycero-3-phosphocholine or POPC (AVANTI PC #850457C) and 15% 1,2-dioleoyl-

sn-glycero-3-phospho-L-serine or DOPS (AVANTI PS #840035C) dissolved in chloroform

were mixed in specific molar ratios in glass tubes and then dried under a stream of N_2, followed

by drying under vacuum overnight to ensure complete removal of organic solvents. Dried lipids

were then combined with 5(6)-carboxyfluorescein (CF) (Sigma #21877) and vortexed for 5

minutes. Lipids were transferred to 2 mL cryotubes and subjected to five freeze thaw cycles in

liquid N_2 and stored at −80 °C freezer. Lipids were then thawed and passed through a 100

nm pore membrane using a mini-extruder kit (Avanti Polar Lipids #610000) and purified

on a PD-10 column to remove excess dye. Prepared liposomes were diluted in standard

assay buffer (10 mM MES, pH 6.0, 25 mM NaCl) to a working concentration of 100 μM.

QuantaMaster 300 fluorometer (Photon Technology International) was used to monitor

fluorescence. SPRR protein was added to the system at increasing concentrations. At the

end time point, 1% v/v n-octylglucoside detergent (OG, Anatrace #O311) was added to

completely disrupt the liposomes. Fluorescence was measured over time in seconds and as

a percentage of total CF dye released by the detergent OG.

Dye uptake assay

Bacterial cultures were grown to mid-logarithmic phase and then pelleted and washed in

standard assay buffer (10 mM MES, pH 6.0 and 25 mM NaCl). Bacteria were then diluted to

approximately 5×10^8 cells/mL in standard assay buffer containing 5.5 μg/mL Propidium iodide

(PI) (Thermo Fisher, P3566). Then bacterial samples (90 μL each well) were added to black

96-well Costar plates (Fisher, 07-200-567) and placed into a Spectramax plate reader

(Molecular Devices) that was pre-equilibrated to 37°C for 10 minutes. After an initial reading,

10 μL of recombinant purified SPRR proteins at varying concentrations or BSA were added

and fluorescence outputs (excitation, 535 nm; emission, 617 nm) were measured every 5

minutes for 1 hour. PI uptake activity was measured against the maximum fluorescence output

from the positive control added with 0.05% SDS.

Skin infections

The dorsal hair was removed from C57BL6 $Sprr1a^{-/-};Sprr2a^{-/-}$ or WT male mice by shaving

(Andis ProClip), followed by depilatory cream (Nair). For MRSA infection, one day after

shaving, bioluminescent MRSA strain SAP430-*lux* were grown to mid-logarithmic phase in

Tryptic Soy Broth and then pelleted and washed twice in PBS. Approximately 1 x 10^6 CFU

bacteria in 100 μL PBS were placed on rectangular gauze. For *P. aeruginosa*- infection, after

24 hours, the dorsal skin was superficially abraded in a crosshatch pattern by a 15-blade scalpel

(Fine Scientific Tools 10115-10) to introduce skin wounding. One day post-wounding, the

bioluminescent *P. aeruginosa* strain Xen05 (Perkin Elmer 119228) were grown in Brain Heart

Infusion Broth (BD Biosciences) and around 1 x 10^6 CFU bacteria in 100 μL PBS were placed

on a gauze rectangle. The gauze was applied to the dorsal skin of $Sprr1a^{-/-};Sprr2a^{-/-}$ or WT

mice and held in place with 2 tegaderm (3M 9505W) and a Band-Aid Sheer Strips (BAND-

AID) for 3 days. The luminescent was monitored daily with serial photography and

quantification using a IVIS Lumina3 imager machine.

TEWL measurement

Transepidermal water loss (TEWL) of mice dorsal skin was measured using Vapometer (Delfin

Technologies) according to manufacturer instructions.

Data availability

RNA-seq data (Figures 1B,1C) has been submitted to the Gene Expression Omnibus with an

accession number: GSE182756

Statistical analysis

Statistical details of experiments can be found in the figure legends, including how significance

was defined and the statistical methods used. Data represent mean ± standard error of the mean.

Formal randomization techniques were not used; however, mice were allocated to experiments

randomly and samples were processed in an arbitrary order. Mouse skin samples that were

determined to be in the anagen hair cycle were excluded. All statistical analyses were performed

with GraphPad Prism software. To assess the statistical significance of the difference between

two treatments, we used two-tailed Student's t-tests. To assess the statistical significance of

differences between more than two treatments, we used one-way ANOVAs.

Acknowledgements:

We would like to thank Dr. Lora Hooper for helpful discussion and sharing $Sprr2a^{-/-}$ mice

strain. We thank Dr. Richard Wang for sharing the hTert immortalized human keratinocyte cell

line. We thank Dr. Zheng Kuang for his help with analysis of RNA-sequencing data. This work

was supported by a Harold Amos Award through the Robert Wood Johnson Foundation,

National Institutes of Health K08 award, a UT Southwestern Disease Oriented Clinical Scholars

Program award, and a Burroughs Wellcome Fund Career Award for Medical Scientists.

Author Contributions:

C.Z. designed and performed most experiments. Z.H. performed protein expression and

purification and liposome assays in fig S6 and Fig 5. A.L. performed skin infection experiments

in Fig 4. M.A. and M.E. provided technical assistance with the mouse experiments and assisted

with bacterial killing assays in Fig 3A, 3B. M.S. completed the dose response and time course

experiments in fig S1. C.C.Z revised the manuscript and is the original source of the SZ95 cells.

C.Z. and T.A.H. designed experiments, interpreted results, and wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Fig. 1. The expression of *SPRR* family genes are up-regulated by lipopolysaccharide (LPS)

in human sebaceous gland cells. (A) Schematic of facial skin derived immortalized human

SZ95 sebaceous gland cells treated with LPS for RNA sequencing. (B) Volcano plot with

significantly up-regulated genes (fold change>1.4 and adjusted *p* value<0.01) highlighted in

red (up regulated in LPS treated). (C) Heat map of significantly up-regulated genes, represented

as Z-scored RPKM (reads per kilo base per million reads). *SPRR1B, SPRR2A* and *SPRR2D* are

highlighted in red. (D) Reverse transcription-quantitative polymerase chain reaction (RT-

qPCR) analysis of *SPRR1B, SPRR2A* and *SPRR2D* transcript in the vehicle and LPS treated

human SZ95 sebocytes. (E) Western blot analysis of SPRR1B and SPRR2A was performed on

vehicle or LPS treated SZ95 cells. GAPDH was used as the loading control. (F) Quantification

of Western blot in (E). Means ± SEM are plotted, *$P<0.05$, **$P<0.01$, ***$P<0.001$ was

determined by unpaired t-test.

Fig. 2. The expression of *SPRR* family genes are up regulated by LPS in mice. (A-D) qRT-
PCR analysis of *Sprr1a* and *Sprr2a* gene expression in mouse dorsal skin tissue. (A) Germ-free
mice (GF) compared to conventionally raised mice (CV). (B) PBS treated mouse skin compared
to LPS intradermal injection mouse skin in WT or MYD88$^{-/-}$ mouse. (C) Mouse intradermally-
injected by vehicle (PBS), *E.coli* or heat-inactivated (HI) E.coli. (D) Untreated mouse skin
compared to wounded skin abraded in a crosshatch pattern by a 15-blade scalpel. Means ± SEM
are plotted. *P < 0.05; **P < 0.01; ****P < 0.0001ns, not significant by unpaired t test. (E)
Immunofluorescence staining of SPRR1A expression in mice skin. CD36 was used as marker
of sebocyte cells. Nuclei are stained with DAPI (Blue). Epidermis and sebaceous gland
indicated with an arrow. White dashed line separates the epidermis and dermis. Yellow dashed
line indicates the outline of SG. Scale bar, 50 μm.

Fig. 3. SPRR family proteins exert bactericidal activity against various skin commensal
and pathogenic bacteria by membrane disruption. (A) Increasing concentrations of purified
recombinant SPRR proteins were added to mid-logarithmic phase *MRSA*, *S. epidermidis*,
P.aeruginosa for 2 hours and surviving bacteria were quantified by dilution plating. (B) 2.5 μM
of SPRR proteins was added to midlogarithmic phase bacteria for 2 hours and surviving bacteria
were quantified by dilution plating. Remaining Colony forming units (CFUs) are expressed as
a percentage of untreated bacteria control. (C) Transmission electron microscopy of *P.*
aeruginosa after incubation with 5 μM purified recombinant SPRR proteins. BSA was used as
negative control. Examples of cell surface damage and cytoplasmic leakage are indicated with
red rectangular box. Upper scale bars, 500 nm. Lower scale bars, 100 nm. (D) Propidium iodide
(PI) uptake by *P. aeruginosa* in the presence of increasing concentrations of mouse SPRR1A
and human SPRR1B protein. (E) Membranes displaying various lipids were incubated with
1μg/mL SPRR proteins, and detected with specific antibody. (F) Carboxyfluorescein (CF)-
loaded liposomes were treated with increasing concentrations of mouse SPRR1A and human
SPRR1B protein, and dye efflux was monitored over time and was expressed as a percentage

of total efflux in the presence of the detergent octyl glucoside (OG). All assays were performed

in triplicate. Means ± SEM are plotted. *P < 0.05; ***P < 0.001; ****P < 0.0001; ns, not

significant by two-tailed t-test.

Fig. 4. SPRR family proteins protect against skin MRSA and *P. aerugonisa* infection. (A-

C) WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice were epicutaneously challenged with MRSA (1 x 10^6

CFU) on the shaved dorsal skin for 3 consecutive days. (A) Representative *in vivo*

bioluminescent imaging (BLI) photographs from Day 0 to Day 3. Quantification of MRSA total

flux (photons/s) for each mouse (B) or plotted as Means ± SEM (C). (D-F) WT and

Sprr1a$^{-/-}$;Sprr2a$^{-/-}$ mice were superficially abraded in a crosshatch pattern by a 15-blade

scalpel to introduce skin wounding. One day post crosshatch wounding, WT and

*Sprr1a$^{-/-}$;Sprr2a$^{-/-}$*mice were topically applied with 1 x 10^6 CFU *P. aerugonisa* on the back

skin for 3 days. (D) Representative in vivo bioluminescent imaging (BLI) signals from Day 0

to Day 3. Quantification of *P. aerugonisa* total flux (photons/s) for each mouse (E) or plotted

as Means ± SEM (F). *p < 0.05, ns, not significant by unpaired t-test.

fig. S1. Dose response and time course analysis of LPS treatment on human sebocyte cells.

qRT-PCR analysis of *SPRR1B and SPRR2A* transcript in the vehicle and various LPS treated

human SZ95 sebocytes. (A) SZ95 sebocyte cells were treated with different concentration of

LPS for 16 hours. (B) SZ95 cell were treated with 1 μg/mL LPS for different length of time.

Means ± SEM are plotted. **P < 0.01; ***P < 0.001;ns, not significant by unpaired t-test.

fig. S2. The expression of SPRR family genes are up regulated by TLR2 and TLR4

agonists in human sebaceous gland cells. (A) A list of different TLR agonists. (B) qRT-PCR

analysis of *SPRR* family genes expression in the vehicle and various TLR agonists treated SZ95

cells. Means ± SEM are plotted. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; ns, not

significant by one-way ANOVAs.

fig. S3. Gram-negative bacteria can trigger *Sprr2a* gene expression in human sebaceous

gland cells. qRT-PCR analysis of *Sprr2a* transcript in the vehicle and various heat inactivated

bacteria treated human SZ95 sebocyte cells. Means ± SEM are plotted. ****P < 0.0001; ns, not

significant by one-way ANOVAs.

fig. S4. LPS cannot induce the expression of *SPRR* family genes in human keratinocyte
cells. hTERT immortalized human keratinocyte cells were treated with LPS for 16 hours. qRT-
PCR analysis of *SPRR1B, SPRR2A* and *SPRR2D* transcript in the vehicle and LPS treated
human hTERT keratinocytes. Means ± SEM are plotted. ns, not significant by unpaired t-test.

fig. S5. LPS cannot trigger the expression of *Sprr* family genes in primary mouse
keratinocytes. Primary mouse keratinocytes were isolated from 3-5 days old neonatal mice
through dispase digestion. With 0.05 mM $CaCl_2$ (Low Ca^{2+}), keratinocytes can proliferate but
won't differentiate into a stratified layer. Under 1 mM $CaCl_2$ (High Ca^{2+}) condition,
keratinocytes will initiate differentiation process. (A) qRT-PCR analysis of *Sprr1a* and *Sprr2a*
transcript in the vehicle and LPS treated primary mouse keratinocyte cells cultured under Low
Ca^{2+} or High Ca^{2+} condition. (B) Western blot analysis of SPRR1A protein was performed on
vehicle or LPS treated primary mouse keratinocytes under both Low Ca^{2+} and High Ca^{2+}
condition. (C) Quantification of the Western blot in (B). Means ± SEM are plotted. *P < 0.05;
***P < 0.001 by unpaired t-test.

fig. S6. Recombinant expression and purification of SPRR proteins. (A and C)
Recombinant mouse SPRR1A protein (A) and human SPRR1B protein (C) was expressed using
a baculovirus expression system and purified by size-exclusion chromatography (Superdex 75
10/300 GL) [26]. (B and D) Fractions from (A) and (C) were visualized by Coomassie blue
staining following SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

fig. S7. SPRR1A protein was resistant to Gram-negative bacteria *Escherichia coli*.
Increasing concentrations of purified recombinant mouse SPRR1A protein was added to mid-
logarithmic phase *Escherichia coli* for 2 hours and surviving bacteria were quantified by
dilution plating. Remaining Colony forming units (CFUs) are expressed as a percentage of
untreated bacteria control. BSA was used as negative control. Means ± SEM are plotted.

fig. S8. Generation and validation of *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$*mice by CRISPR/Cas9 genomic
targeting. (A) Schematic diagram of two-step strategy using CRISPR/Cas9-mediated gene
targeting to delete the entire *Sprr2a* and *Sprr1a* locus. (B) qRT-PCR analysis of *Sprr1a* and
Sprr2a expression in the skin of WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice. Values were normalized to
Gapdh expression. Means ± SEM are plotted. *P < 0.05; ***P < 0.001 by unpaired t-test. (C)
Western blot analysis of SPRR1A protein in the skin of WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$*mice.
GAPDH was used as the loading control.

fig. S9. *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice does not show signs of inflammation or impaired skin
barrier. (A) Transepidermal water loss (TEWL) of WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$*mice skin was
analyzed. Means ± SEM is plotted, ns, not significant by unpaired t test. (B) Representative
H&E staining of mouse skin from WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$*. Scale bar, 200 μm.

References:

1. Belkaid, Y. & Segre, J. A. Dialogue between skin microbiota and immunity. *Science* (2014) doi:10.1126/science.1260144.

2. Duerkop, B. A. & Hooper, L. V. Resident viruses and their interactions with the immune system. *Nature Immunology* (2013) doi:10.1038/ni.2614.

3. Oh, J., Byrd, A. L., Park, M., Kong, H. H. & Segre, J. A. Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).

4. Findley, K. *et al.* Topographic diversity of fungal and bacterial communities in human skin. *Nature* (2013) doi:10.1038/nature12171.

5. Dryden, M. S. Skin and soft tissue infection: microbiology and epidemiology. *Int. J. Antimicrob. Agents* (2009) doi:10.1016/S0924-8579(09)70541-2.

6. Stulberg, D. L., Penrod, M. A. & Blatny, R. A. Common bacterial skin infections. *Am. Fam. Physician* (2002) doi:10.1080/00325481.1993.11701692.

7. Huebinger, R. M. *et al.* Targeting bacterial adherence inhibits multidrug-resistant Pseudomonas aeruginosa infection following burn injury. *Sci. Rep.* **6**, 1–8 (2016).

8. Klevens, R. M. *et al.* Invasive methicillin-resistant Staphylococcus aureus infections in the United States. *J. Am. Med. Assoc.* (2007) doi:10.1001/jama.298.15.1763.

9. Zhang, L. J. & Gallo, R. L. Antimicrobial peptides. *Current Biology* (2016) doi:10.1016/j.cub.2015.11.017.

10. Nakatsuji, T. & Gallo, R. L. Antimicrobial peptides: Old molecules with new ideas. *Journal of Investigative Dermatology* (2012) doi:10.1038/jid.2011.387.

11. Mukherjee, S. *et al.* Antibacterial membrane attack by a pore-forming intestinal C-type lectin. *Nature* (2014) doi:10.1038/nature12729.

12. Mishra, A. K., Choi, J., Moon, E. & Baek, K. H. Tryptophan-rich and proline-rich antimicrobial peptides. *Molecules* (2018) doi:10.3390/molecules23040815.

13. Gallo, R. L. & Hooper, L. V. Epithelial antimicrobial defence of the skin and intestine. *Nature Reviews Immunology* vol. 12 503–516 (2012).

14. Harris, T. A. *et al.* Resistin-like Molecule a Provides Vitamin-A- Dependent
Antimicrobial Protection in the Skin Article Resistin-like Molecule a Provides
Vitamin-A-Dependent Antimicrobial. *Cell Host Microbe* **25**, 1–12 (2019).

15. Fischer, H. *et al.* Holocrine Secretion of Sebum Is a Unique DNase2-
Dependent Mode of Programmed Cell Death. *J. Invest. Dermatol.* **137**, 587–
594 (2017).

16. Zouboulis, C. C., Yoshida, G. J., Wu, Y., Xia, L. & Schneider, M. R.
Sebaceous gland: Milestones of 30-year modelling research dedicated to the
"brain of the skin". *Exp. Dermatol.* (2020) doi:10.1111/exd.14184.

17. Zouboulis, C. C. *et al.* Beyond acne: Current aspects of sebaceous gland
biology and function. *Reviews in Endocrine and Metabolic Disorders* (2016)
doi:10.1007/s11154-016-9389-5.

18. Cabral, A. *et al.* Structural Organization and Regulation of the Small Proline-
rich Family of Cornified Envelope Precursors Suggest a Role in Adaptive
Barrier Function. *J. Biol. Chem.* (2001) doi:10.1074/jbc.M100336200.

19. Candi, E., Schmidt, R. & Melino, G. The cornified envelope: A model of cell
death in the skin. *Nature Reviews Molecular Cell Biology* vol. 6 328–340
(2005).

20. Vaishnava, S. *et al.* The antibacterial lectin RegIIIγ promotes the spatial
segregation of microbiota and host in the intestine. *Science (80-.).* (2011)
doi:10.1126/science.1209791.

21. Vermeij, W. P. & Backendorf, C. Skin cornification proteins provide global
link between ROS detoxification and cell migration during wound healing.
PLoS One **5**, e11957–e11957 (2010).

22. Fischer, D. F., Gibbs, S., van De Putte, P. & Backendorf, C. Interdependent
transcription control elements regulate the expression of the SPRR2A gene
during keratinocyte terminal differentiation. *Mol. Cell. Biol.* **16**, 5365–5374
(1996).

23. MW, S., DF, F., E, de M., P, van de P. & C, B. AP-1 and ets transcription
factors regulate the expression of the human SPRR1A keratinocyte terminal
differentiation marker. *J. Biol. Chem.* **273**, 24683–24692 (1998).

24. Scocchi, M., Tossi, A. & Gennaro, R. Proline-rich antimicrobial peptides:
Converging to a non-lytic mechanism of action. *Cellular and Molecular Life*
Sciences (2011) doi:10.1007/s00018-011-0721-7.

25. Stotz, H. U., Thomson, J. G. & Wang, Y. Plant defensins: defense,
development and application. *Plant Signal. Behav.* **4**, 1010–1012 (2009).

26. Hu Zehan, Zhang Chenlu….H. L. V. Small proline rich protein 2A is a
bactericidal protein that defends the intestinal barrier during helminth infection.
(Accepted) *Science (80-.).* (2021).

27. Grice, E. A. *et al.* A diversity profile of the human skin microbiota. *Genome*
Res. (2008) doi:10.1101/gr.075549.107.

28. Demetris, A. J. *et al.* Small proline-rich proteins (SPRR) function as SH3
domain ligands, increase resistance to injury and are associated with epithelial-
mesenchymal transition (EMT) in cholangiocytes. *J. Hepatol.* (2008)
doi:10.1016/j.jhep.2007.09.019.

29. Chen, Y. E., Fischbach, M. A. & Belkaid, Y. Skin microbiota-host interactions.
Nat. Publ. Gr. (2018) doi:10.1038/nature25177.

30. Lovászi, M., Szegedi, A., Zouboulis, C. C. & Törőcsik, D. Sebaceous-
immunobiology is orchestrated by sebum lipids. *Dermatoendocrinol.* **9**,
e1375636 (2017).

31. Malik, E., Dennison, S. R., Harris, F. & Phoenix, D. A. pH Dependent
Antimicrobial Peptides and Proteins, Their Mechanisms of Action and
Potential as Therapeutic Agents. *Pharmaceuticals* **9**, (2016).

32. Hooper, L. V. *et al.* Molecular Analysis of Commensal Host-Microbial
Relationships in the Intestine. *Science (80-.).* **291**, 881–884 (2001).

33. Liu, H. *et al.* Staphylococcus aureus Epicutaneous Exposure Drives Skin
Inflammation via IL-36-Mediated T Cell Responses. *Cell Host Microbe* **22**,
653-666.e5 (2017).

Figure 1

Fig. 1. The expression of *SPRR* family genes are up-regulated by lipopolysaccharide (LPS) in human sebaceous gland cells. (A) Schematic of facial skin derived immortalized human SZ95 sebaceous gland cells treated with LPS for RNA sequencing. (B) Volcano plot with significantly up-regulated genes (fold change>1.4 and adjusted *p* value<0.01) highlighted in red (up regulated in LPS treated). (C) Heat map of significantly up-regulated genes, represented as Z-scored RPKM (reads per kilo base per million reads). *SPRR1B, SPRR2A* and *SPRR2D* are highlighted in red. (D) Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) analysis of *SPRR1B, SPRR2A* and *SPRR2D* transcript in the vehicle and LPS treated human SZ95 sebocytes. (E) Western blot analysis of SPRR1B and SPRR2A was performed on vehicle or LPS treated SZ95 cells. GAPDH was used as the loading control. (F) Quantification of Western blot in (E). Means ± SEM are plotted, *$P<0.05$, **$P<0.01$, ***$P<0.001$ was determined by unpaired t-test.

Figure 2

Fig. 2. The expression of *SPRR* family genes are up regulated by LPS in mice. (A-D) qRT-PCR analysis of *Sprr1a* and *Sprr2a* gene expression in mouse dorsal skin tissue. (A) Germ-free mice (GF) compared to conventionally raised mice (CV). (B) PBS treated mouse skin compared to LPS intradermal injection mouse skin in WT or MYD88$^{-/-}$ mouse. (C) Mouse intradermally-injected by vehicle (PBS), *E.coli* or heat-inactivated (HI) E.coli. (D) Untreated mouse skin compared to wounded skin abraded in a crosshatch pattern by a 15-blade scalpel. Means ± SEM are plotted. *P < 0.05; **P < 0.01; ****P < 0.0001ns, not significant by unpaired t test. (E) Immunofluorescence staining of SPRR1A expression in mice skin. CD36 was used as marker of sebocyte cells. Nuclei are stained with DAPI (Blue). Epidermis and sebaceous gland indicated with an arrow. White dashed line separates the epidermis and dermis. Yellow dashed line indicates the outline of SG. Scale bar, 50 μm.

Figure 3

Fig. 3. SPRR family proteins exert bactericidal activity against various skin commensal and pathogenic bacteria by membrane disruption. (A) Increasing concentrations of purified recombinant SPRR proteins were added to mid-logarithmic phase *MRSA*, *S. epidermidis*, *P.aeruginosa* for 2 hours and surviving bacteria were quantified by dilution plating. (B) 2.5 μM of SPRR proteins was added to midlogarithmic phase bacteria for 2 hours and surviving bacteria were quantified by dilution plating. Remaining Colony forming units (CFUs) are expressed as a percentage of untreated bacteria control. (C) Transmission electron microscopy of *P. aeruginosa* after incubation with 5 μM purified recombinant SPRR proteins. BSA was used as negative control. Examples of cell surface damage and cytoplasmic leakage are indicated with red rectangular box. Upper scale bars, 500 nm. Lower scale bars, 100 nm. (D) Propidium iodide (PI) uptake by *P. aeruginosa* in the presence of increasing concentrations of mouse SPRR1A and human SPRR1B protein. (E) Membranes displaying various lipids were incubated with 1μg/mL SPRR proteins, and detected with specific antibody. (F) Carboxyfluorescein (CF)-loaded liposomes were treated with increasing concentrations of mouse SPRR1A and human SPRR1B protein, and dye efflux was monitored over time and was expressed as a percentage of total efflux in the presence of the detergent octyl glucoside (OG). All assays were performed in triplicate. Means ± SEM are plotted. *P < 0.05; ***P < 0.001; ****P < 0.0001; ns, not significant by two-tailed t-test.

Figure 4

Fig. 4. SPRR family proteins protect against skin MRSA and *P. aerugonisa* infection. (A-C) WT and *Sprr1a⁻/⁻;Sprr2a⁻/⁻* mice were epicutaneously challenged with MRSA (1 x 10⁶ CFU) on the shaved dorsal skin for 3 consecutive days. (A) Representative *in vivo* bioluminescent imaging (BLI) photographs from Day 0 to Day 3. Quantification of MRSA total flux (photons/s) for each mouse (B) or plotted as Means ± SEM (C). (D-F) WT and *Sprr1a⁻/⁻;Sprr2a⁻/⁻* mice were superficially abraded in a crosshatch pattern by a 15-blade scalpel to introduce skin wounding. One day post crosshatch wounding, WT and *Sprr1a⁻/⁻;Sprr2a⁻/⁻* mice were topically applied with 1 x 10⁶ CFU *P. aerugonisa* on the back skin for 3 days. (D) Representative in vivo bioluminescent imaging (BLI) signals from Day 0 to Day 3. Quantification of *P. aerugonisa* total flux (photons/s) for each mouse (E) or plotted as Means ± SEM (F). *p < 0.05, ns, not significant by unpaired t-test.

figure 1

fig. S1. Dose response and time course analysis of LPS treatment on human sebocyte cells. qRT-PCR analysis of *SPRR1B and SPRR2A* transcript in the vehicle and various LPS treated human SZ95 sebocytes. (A) SZ95 sebocyte cells were treated with different concentration of LPS for 16 hours. (B) SZ95 cell were treated with 1 μg/mL LPS for different length of time. Means ± SEM are plotted. **P < 0.01; ***P < 0.001;ns, not significant by unpaired t-test.

A

TLR Ligands	Targets	Description
Pam3CSK4	TLR1/2	Tripalmitoylated lipopeptide that mimics the bacterial lipoprotein
HKLM	TLR2	Heat-killed *Listeria monocytogenes*
Poly(I:C) HMW	TLR3	Double-strand RNA (1.5-8kb)
Poly(I:C) LMW	TLR3	Double-strand RNA (0.2-1kb)
LPS	TLR4	Lipopolysaccharide
Flagellin	TLR5	Component of the bacterial flagellar filament
FSL-1	TLR6/2	Lipopeptide that mimics the lipoprotein LP44 of *Mycoplasma salivarium*
Imiquimod	TLR7	An imidazoquinoline amine analog to guanosine

B

fig. S2. The expression of SPRR family genes are up regulated by TLR2 and TLR4 agonists in human sebaceous gland cells. (A) A list of different TLR agonists. (B) qRT-PCR analysis of *SPRR* family genes expression in the vehicle and various TLR agonists treated SZ95 cells. Means ± SEM are plotted. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; ns, not significant by one-way ANOVAs.

figure 3

fig. S3. Gram-negative bacteria can trigger *Sprr2a* gene expression in human sebaceous gland cells. qRT-PCR analysis of *Sprr2a* transcript in the vehicle and various heat inactivated bacteria treated human SZ95 sebocyte cells. Means ± SEM are plotted. ****P < 0.0001; ns, not significant by one-way ANOVAs.

figure 4

fig. S4. LPS cannot induce the expression of *SPRR* family genes in human keratinocyte cells. hTERT immortalized human keratinocyte cells were treated with LPS for 16 hours. qRT-PCR analysis of *SPRR1B, SPRR2A* and *SPRR2D* transcript in the vehicle and LPS treated human hTERT keratinocytes. Means ± SEM are plotted. ns, not significant by unpaired t-test.

figure 5

fig. S5. LPS cannot trigger the expression of *Sprr* family genes in primary mouse keratinocytes. Primary mouse keratinocytes were isolated from 3-5 days old neonatal mice through dispase digestion. With 0.05 mM CaCl$_2$ (Low Ca^{2+}), keratinocytes can proliferate but won't differentiate into a stratified layer. Under 1 mM CaCl$_2$ (High Ca^{2+}) condition, keratinocytes will initiate differentiation process. (A) qRT-PCR analysis of *Sprr1a* and *Sprr2a* transcript in the vehicle and LPS treated primary mouse keratinocyte cells cultured under Low Ca^{2+} or High Ca^{2+} condition. (B) Western blot analysis of SPRR1A protein was performed on vehicle or LPS treated primary mouse keratinocytes under both Low Ca^{2+} and High Ca^{2+} condition. (C) Quantification of the Western blot in (B). Means ± SEM are plotted. *P < 0.05; ***P < 0.001 by unpaired t-test.

figure 6

fig. S6. Recombinant expression and purification of SPRR proteins. (A and C) Recombinant mouse SPRR1A protein (A) and human SPRR1B protein (C) was expressed using a baculovirus expression system and purified by size-exclusion chromatography (Superdex 75 10/300 GL) [26]. (B and D) Fractions from (A) and (C) were visualized by Coomassie blue staining following SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

figure 7

fig. S7. SPRR1A protein was resistant to Gram-negative bacteria *Escherichia coli*. Increasing concentrations of purified recombinant mouse SPRR1A protein was added to mid-logarithmic phase *Escherichia coli* for 2 hours and surviving bacteria were quantified by dilution plating. Remaining Colony forming units (CFUs) are expressed as a percentage of untreated bacteria control. BSA was used as negative control. Means ± SEM are plotted.

figure 8

A   **Step 1: Deletion of *Sprr2a* locus**

B

fig. S8. Generation and validation of *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice by CRISPR/Cas9 genomic targeting. (A) Schematic diagram of two-step strategy using CRISPR/Cas9-mediated gene targeting to delete the entire *Sprr2a* and *Sprr1a* locus. (B) qRT-PCR analysis of *Sprr1a* and *Sprr2a* expression in the skin of WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice. Values were normalized to *Gapdh* expression. Means ± SEM are plotted. *P < 0.05; ***P < 0.001 by unpaired t-test. (C) Western blot analysis of SPRR1A protein in the skin of WT and *Sprr1a$^{-/-}$;Sprr2a$^{-/-}$* mice. GAPDH was used as the loading control.

figure 9

fig. S9. *Sprr1a⁻/⁻;Sprr2a⁻/⁻* mice does not show signs of inflammation or impaired skin barrier. (A) Transepidermal water loss (TEWL) of WT and *Sprr1a⁻/⁻;Sprr2a⁻/⁻* mice skin was analyzed. Means ± SEM is plotted, ns, not significant by unpaired t test. (B) Representative H&E staining of mouse skin from WT and *Sprr1a⁻/⁻;Sprr2a⁻/⁻*. Scale bar, 200 µm.