Redox-dependent Structure and Dynamics of Macrophage Migration Inhibitory Factor Reveal Sites of Latent Allostery

Erin Skeens,[1] Meagan Gadzuk-Shea,[2] Dilip Shah,[3] Vineet Bhandari,[3] Devin K. Schweppe,[2] Rebecca B. Berlow,[4]* and George P. Lisi[1]*

[1]Department of Molecular Biology, Cell Biology, & Biochemistry, Brown University, Providence, RI 02903
[2]Department of Genome Sciences, University of Washington, Seattle, WA 98195
[3]Section of Neonatology, Department of Pediatrics, Cooper University Hospital, Camden, NJ 08103
[4]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037

*To whom correspondence should be addressed: George P. Lisi (george_lisi@brown.edu)
Rebecca B. Berlow (rberlow@scripps.edu)

ORCID: Erin Skeens (0000-0002-0580-1291)
Rebecca B. Berlow (0000-0003-1934-0139)
George P. Lisi (0000-0001-8878-5655)
Meagan Gadzuk-Shea (0000-0001-9496-529X)

Abstract

Macrophage migration inhibitory factor (MIF) is a multifunctional immunoregulatory protein that is a key player in the innate immune response. Given its overexpression at sites of inflammation in a wide range of diseases marked by increasingly oxidative cellular environment, a comprehensive structural understanding of how cellular redox conditions may impact the structure and function of MIF is necessary. We used solution NMR spectroscopy and mass spectrometry to investigate structural and dynamic signatures of MIF under varied solution redox conditions. Our results indicate that the MIF structure is modified and becomes increasingly dynamic in an oxidative environment, which may be a means to alter the MIF functional response in a redox-dependent manner. We identified latent allosteric sites within MIF that are redox-sensitive and mutational analysis reveals that loss of redox-responsive residues attenuates activation of the coreceptor CD74. Leveraging sites of redox-sensitivity therefore reveals an avenue to modulate MIF function in its "disease state" via structure-based drug design.

Introduction

Macrophage migration inhibitory factor (MIF) is a ubiquitous human protein that plays a crucial role in the pathophysiology of inflammation.[1] MIF is expressed by nearly every cell in the body[1] and many of its effects are elicited by its binding to the chemokine receptors CXCR2 and CXCR4[2,3] or the coreceptor CD74.[4-6] Circulating levels are reported in the low μM range in healthy humans,[7-9] but MIF is overexpressed under inflammatory conditions and has been linked to glucocorticoid overriding activity,[10] asthma,[7,11] rheumatoid arthritis,[12] colitis,[13] pancreatitis,[14] acute respiratory distress syndrome (ARDS),[15] and cancer[16-20] as well as complications from COVID-19.[21] Efforts to elucidate the pathogenic mechanism of MIF have attempted to connect

its disease state to cellular redox properties,[11] most recently by investigating its cysteine

residues as conformational switches.[22] Mutation of either C56 or C59 in a canonical [56]CXXC[59]

(XX = AL in MIF) motif modulates the enzymatic and macrophage activation functions of MIF.[23,24]

An oxidative cellular environment is one of the hallmarks of inflammatory disease,

characterized by excess reactive oxygen species (ROS) at sites of inflammation.[25] Despite the

acknowledged contribution of redox imbalance to asthma,[26,27] ARDS,[28] pulmonary fibrosis,[29] and

cancer,[30] as well as the link between these pathologies and MIF, very little structural work has

directly addressed the redox behavior of MIF. We wondered whether MIF could utilize redox

imbalances to act as a pro-inflammatory sensor to toggle its structure based on the chemical

environment of the cell. Altered conformations of MIF at local sites of inflammation could then

modulate its function or create different modes of interaction with accessory proteins, thereby

controlling downstream biological responses. In a first step toward exploring such a mechanism,

we investigated the impact of solution redox conditions on the MIF structure and identified

regions of the protein that are sensitive to oxidative environments. We used solution nuclear

magnetic resonance (NMR) spectroscopy to assess the redox-dependent dynamics of MIF with

spin relaxation experiments and used mass spectrometry to identify redox-modified cysteine

residues. We then leveraged these findings, using further NMR structural studies and an *in vivo*

assay, to confirm redox-sensitive amino acids as latent points of functional control in MIF.

Results

Oxidizing solution alters MIF structural stability

To assess the effect of solution redox potentials on the MIF structure, we performed far-

UV circular dichroism experiments on redox-neutral MIF samples as well as MIF samples that

had been oxidized or reduced. CD spectra of redox-altered MIF show characteristic α-β

structure, with only minor differences (**Figure S1A**). These data are consistent with numerous

MIF structures in the Protein Data Bank (PDB) that have no obvious alterations in the presence

of substrates, inhibitors, or mutations. Despite similarly folded structures, temperature-

dependent CD experiments (**Figure S1B**) show that an oxidizing environment destabilizes MIF

to unfolding relative to reduced and redox-neutral MIF (ΔT_m = - 4.56 °C, $\Delta\Delta G$ = - 2.65 kcal/mol,

Figure S1C). This energetic difference (expressed per trimer) is modest, but suggests a

loosening of the MIF structure that may affect its conformational ensemble.

Redox-dependent NMR spectra of MIF highlight multiple conformations and altered dynamics

We used solution NMR to pinpoint regions of the MIF structure that are sensitive to the

redox environment. $^1H^{15}N$ TROSY-HSQC spectral overlays of reduced, oxidized, and redox-

neutral MIF (**Figure S2**) show that while reduced and redox-neutral samples are spectrally

similar, more significant spectral changes are observed for MIF under oxidizing conditions.

Redox-sensitive chemical shift perturbations (relative to redox-neutral MIF) are observed

throughout the protein, including for residues at the N-terminal enzymatic active site, solvent

channel, and monomer-monomer interface (**Figures 1A**). Additionally, for oxidized MIF samples,

duplicate resonances are observed for 22 residues (**Figures 1B, S3**), suggesting that an

oxidizing environment modulates an equilibrium between two conformations of MIF on the NMR

timescale. Sites of slow exchange localize to the monomer-monomer interface (**Figure 1C**),

indicating that the structure of this region of the protein is strongly affected by oxidizing solution,

consistent with the lower thermal stability observed by CD. Perturbations to MIF are not exclusive

to redox-active Cys residues, at times occurring 10 - 20 Å from the [56]CALC[59] motif or C80.

However, Cys and Met residues serve as nucleation points for our observations, with a majority

of redox-dependent changes to the MIF structure occurring proximal to these traditionally

sensitive sites (**Figure S4**). These data are consistent with an alternate oxidized MIF

conformation, "oxMIF," most recently probed in a synthetic peptide comprising the MIF [56]CALC[59]

sequence.[22]

Figure 1. Redox-dependent changes in MIF detected by NMR. **(A)** $^1H^{15}N$ combined chemical shift perturbations ($\Delta\delta$) for oxidized and reduced MIF, relative to a redox-neutral sample. **(B)** Selected resonances from $^1H^{15}N$ HSQC spectral overlays of reduced (blue) and oxidized (red) MIF. Sites of slow exchange observed under oxidizing conditions are indicated by arrows. Populations of MIF conformers determined from the volumes of slow exchange resonances are mapped onto the MIF trimer (PDB: 1MIF) in **(C)**, where the percentage indicated in the legend denotes the minor species (*i.e.* satellite peak). Volumes and populations of slow exchange resonances can be found in Figure S3.

85   Analysis of longitudinal (R_1) and transverse (R_2) NMR relaxation rates[31] highlights local

86   differences caused by redox conditions. Consistent with the altered thermodynamic parameters

87   of the oxidized MIF samples, individual relaxation rates, as well as the R_1R_2 product, qualitatively

imply that MIF exhibits heightened flexibility under oxidizing conditions (**Figure S5A**). Analysis of R_1R_2 values[31] highlights 21 sites suggestive of µs – ms flexibility in oxMIF (R_1R_2 >1.5σ above the 10% trimmed mean of the data), 7 of which do not appear flexible in both redox-neutral and reduced MIF. Despite local patches of altered flexibility, the dynamic profiles of reduced and redox-neutral MIF are similar to each other. ^{1}H-[^{15}N] NOE values and order parameters also hint at patches of local flexibility in oxMIF, though fast timescale dynamics do not differ substantially between samples. Residues with relaxation parameters suggestive of µs – ms flexibility are mapped onto the MIF trimer and monomer structures in **Figure S5**. Correlation plots of R_1R_2 and S^2 parameters for reduced and oxidized MIF reveal a weaker correlation for R_1R_2, suggesting that MIF experiences greater dynamic changes on the µs – ms timescale over a range of redox conditions (**Figure 2A**). Contributions to the variations in R_1R_2 parameters are most strongly affected by oxidizing solution conditions (**Figure 2B, left panel**), while reducing conditions contribute more significantly to the ps – ns dynamics of MIF via S^2 (**Figure 2B, right panel**). Redox-sensitive residues with apparent µs – ms flexibility include T30, G50, L58, S63, I64*, K66, R73, L83, A84, R86, R93, I96, N97*, N109*, N110, T112, and A114, and those with apparent ps – ns fluctuations are F3, T7, L26, T30, D44, I64*, K66, K77, L83, L87, S90, Y95*, D100, A103, and S111* (**Figure 3C**). Asterisks indicate residues previously identified as controlling MIF enzymatic activity and CD74 activation via direct interaction or through allosteric signaling,[6,32,33] indicating redox-dependent perturbations in MIF may have functional consequences.

Figure 2. (A) Correlation plots of the R_1R_2 (left) and S^2 (right) parameters for oxidized and reduced MIF. Gray shaded areas represent $\pm 1.5\sigma$ from the linear least-squares fits of the data (black lines). **(B)** Per-residue differences in R_1R_2 (left) and S^2 (right) plotted as oxidized – reduced. Bars in the shaded red area denote a greater value in oxMIF, while those in the blue area denote a greater value in reduced MIF. **(C)** Residues outside of the 1.5σ correlation boundaries in **(A)** are mapped onto the MIF monomer (PDB: 1MIF), highlighting regions of the protein with redox-dependent differences in μs – ms (left) and ps – ns (right) dynamics.

To confirm the influence of solution redox potential on μs – ms dynamics in MIF, we quantified the exchange contribution to transverse relaxation, R_{ex}, via CPMG relaxation dispersion experiments. The average global R_{ex} value ($<R_{ex}>$) for MIF is fairly similar across the redox regimes tested, with local differences arising primarily under oxidizing conditions (**Figure 3A**). Numerous sites of μs – ms dynamics are observed in all redox-modulated MIF samples (**Figure S6**), however, oxidizing solution stimulates the most widespread relaxation dispersion, affecting 39 residues (~34 %) in the MIF structure. These data are consistent with R_1R_2 values that estimated a stronger contribution from oxidizing solution to μs – ms motions (**Figure 2**).

Regions of MIF with known functional importance become flexible, including the monomer interface (L26, I37, G50) and solvent channel allosteric site (I96, Y99). Despite similar dynamics in large portions of MIF, **Figure 3B** highlights selected sites that only appear flexible under oxidizing conditions. With the exception of the solvent channel, these flexible sites have not been well characterized for functional impact, suggesting that studies of MIF redox dependence can reveal new target residues for structure-function correlations.

Figure 3. (A) Per-residue exchange contribution to relaxation (R_{ex}) for reduced (blue), redox-neutral (black), and oxidized MIF (red). **(B)** CPMG relaxation dispersion profiles for selected MIF residues collected in reducing (left), redox-neutral (center), or oxidizing (right) solution. These sites display evidence of $\mu s - ms$ flexibility under oxidizing conditions.

Cysteine 80 is a selectively modified redox switch

Prior work with a MIF epitope[22] suggested that C80 functions as a redox molecular switch. However, redox sensitivity of C80 in the full-length trimeric protein is unknown. We performed quantitative mass spectrometry to explore the possibility that C80 could be selectively modified by redox reagents. Stepwise alkylation of MIF under each redox condition with iodoacetamide (IAA) and N-ethylmaleimide (NEM) yields critical insight into the oxidation state of C80 in the intact MIF trimer (**Figures 4, S7, S8**). Using NEM-bound C80 as a proxy for the abundance of cystine residues reveals an increase from 75.1% to 91.3% cystine abundance between redox-neutral and oxidizing conditions (**Figure 4B**). Under redox-neutral conditions, C80 exhibits the

greatest variance in cystine abundance, further highlighting the reactivity and sensitivity of this

residue to environmental conditions. The relative abundance of cystine is also sensitive to the

amount of oxidative stress (**Figure S8A**). Increasing the concentration of oxidant in solution from

1:1 to 2:1 shifts the C80 cystine relative abundance from 83.1% to 91.3%. For comparison,

Figure S7B shows the expected contributions to the total ion intensity of C80 modified with IAA

or NEM if C80 is in a fully reduced, 50% reduced, or fully oxidized state. In addition to C80, MIF

also contains two cysteine residues in its ^{56}CALC^{59} motif; however, the peptide fragment

containing C56 and C59 is a large 34-mer, precluding quantitation by the MS methods utilized

here. Although we focus on C80, we cannot rule out a significant role for other cysteines in MIF

redox dependence.

Figure 4. (A) Stepwise alkylation of MIF cysteine residues to determine the relative abundance of oxidized residues via mass spectrometry. The alkylation workflow involved three reaction steps (red, top panel). MIF equilibrated under reducing conditions was treated with only one alkylating agent and served as a quantitative control (blue, bottom panels). **(B)** Relative quantitation of C80 alkylation revealed conformational dependence and sensitivity to the surrounding redox environment.

Mutation of redox-sensitive residues alters MIF structure and biological function

To test the possibility that the redox sensitivity of MIF could reveal sites of functional impact that are not otherwise obvious, we created single point mutations at two residues displaying redox-dependent structural and/or dynamic changes, lysine 66 (K66) and cysteine 80 (C80). K66 was found to have a large variation in the R_1R_2 parameter in oxMIF, as well as a low ^{1}H-[^{15}N] NOE in reduced MIF (Figures 2, S5), indicating that it undergoes significant changes in flexibility between the oxidized and reduced states. C80 also exhibits dynamic changes suggestive of redox-altered μs – ms chemical exchange and was shown by mass spectrometry to be selectively modified under oxidizing conditions (Figures 4, S5-8). ^{1}H^{15}N TROSY-HSQC NMR spectra of the K66A and C80A variants were collected in redox-neutral, reducing, and oxidizing solutions. Baseline structural changes caused by the K66A and C80A mutations are significant under redox-neutral conditions, particularly for the C80A variant, though CD spectroscopy indicates the variants are fully folded at NMR sample temperatures (Figures S9, S10). Interestingly, chemical shift perturbations observed for K66A MIF under oxidizing and reducing conditions are modest when compared to a redox-neutral K66A sample (Figures 5A, 5B). Most notably, oxidized K66A exhibits a chemical shift profile much smaller in magnitude than that observed in the corresponding wt-MIF experiments. In addition, reduced K66A exhibits larger chemical shift perturbations than oxidized in both number and magnitude, the inverse of what was found for wt-MIF. These data alone suggest that K66 may play a role in modulating the redox-dependent structural changes of wt-MIF, which are substantially diminished upon mutation. Likewise, the C80A variant strongly attenuates redox-sensitive NMR chemical shift changes in both reduced and oxMIF. These data, along with mass spectrometry, indicate that C80 modulates the conformational landscape of MIF under oxidative conditions.

Figure 5. Mutation of the redox-sensitive K66 and C80 residues alters the redox-dependent MIF structure and hinders the ability of MIF to activate CD74 *in vivo*. K66A and C80A mutations alter the combined ^{1}H-^{15}N chemical shift profiles of reduced **(A)** and oxidized **(B)** MIF, relative to redox-neutral K66A and C80A samples used as references. Blue lines denote 1.5σ above the 10% trimmed mean of all shifts. Combined ^{1}H-^{15}N chemical shift plots for reduced and oxidized wt-MIF are repeated here from Figure 1 for comparison. **(C)** K66A and C80A mutations significantly decreased CD74-dependent neutrophil recruitment in murine lungs *in vivo*. **(D)** Attenuated CD74-dependent neutrophil recruitment by K66A and C80A also led to significantly decreased pulmonary edema *in vivo*. (n = 4 in each group, *p<0.05, ***p≤0.001 and ****p≤0.0001). Data are expressed as mean ± SEM.

To understand whether this altered redox behavior foreshadowed changes in MIF function, we evaluated the K66A and C80A variants in neutrophil recruitment assays in murine lungs, where we measured total immune cells, percentage of neutrophils in BAL fluid, and protein content (as a marker of alveolar capillary leak/pulmonary edema). This *in vivo* assay quantifies neutrophil recruitment on the surface of alveolar macrophages, an established metric for the

MIF-induced activation of the pro-inflammatory CD74 receptor.[34] The assay is indirect, using

chemokines released by activated macrophages to stimulate CXCR2 receptors on neutrophils

(that do not directly express CD74) to induce their migration to murine lungs.[35] We found that

wt-MIF significantly increased the total BAL fluid cell counts over saline controls, while the K66A

and C80A variants showed a reduced number of total BAL fluid immune cells (**Figure S11**). The

neutrophil influx and pulmonary edema protein levels were significantly increased in the wt-MIF

group relative to vehicle controls (**Figures 5C, 5D**). In contrast, we observed significant

decreases in the groups to which the MIF variants were delivered, as compared to wt-MIF.

Together, our data indicate that the K66A and C80A variants have a functional effect in reducing

the inflammatory response of MIF in the murine lung. These findings also suggest that structural

and/or dynamic sensitivity of residues within MIF can provide a novel route for achieving

functional control beyond the known MIF active sites.

Discussion

We used solution redox potentials to modulate the MIF structure and dynamics in order

to reveal changes in conformational equilibria and redox-sensitive residues that may be latent

functional sites within its structure. This work builds upon prior studies implying redox sensitivity

of MIF[11,22,36,37] as well as our recent paper highlighting a series of allosteric residues linking the

enzymatic (tautomerase) and CD74 binding sites of MIF.[38] Here, we demonstrate that

examination of MIF redox sensitivity is another useful avenue for probing function beyond its

established active sites. Since MIF participates in both intracellular and extracellular

interactions,[39] where the redox environments differ, understanding the implications of its redox

dependence may also inform its pro-inflammatory mechanisms and functional promiscuity,

which has been investigated at the biochemical, but not the structural, level. A previous

characterization of the association of MIF with p53 tumor suppressor suggested the interaction

to be redox-dependent, having shown a significant decrease in coprecipitation in the presence

of a reductant.[40] Further, interaction of MIF with ribosomal protein S-19 (RPS-19) was shown to

occur more strongly under reducing conditions.[41] In addition, MIF is known to be post-

translationally modified by glycation in the brain tissue of Alzheimer's patients. Most notably, an

altered electrophoretic migratory behavior of glycated MIF was observed and demonstrated *in*

vitro by incubating MIF with glucose under oxidizing conditions.[42]

Redox signatures of inflammatory diseases, most of which implicate MIF, can also be

markedly different from those of healthy humans. The redox potential of epithelial lining fluid is -

110 to -40 mV in the lungs of asthmatics, which is significantly more oxidizing than that of healthy

humans (-150 to -200 mV).[27] Further, the synovial fluid of rheumatoid arthritis patients has been

shown to be highly oxidizing relative to that of a non-arthritic control group.[43,44] A significant body

of literature suggests that when overexpressed at sites of inflammation,[16] MIF can endure large

swings in environmental redox potential, consistent with our determination of the MIF redox

potential *in vitro* using solution NMR[45,46] (**Figure S12**, midpoint E_0' = -215 $\pm$ 5 mV, near normal

physiological range). NMR spin relaxation studies show redox-dependent modulation of local

flexibility in MIF, with motion observed on multiple timescales. oxMIF, in particular, exhibits the

most significant structural and dynamic changes, with widespread NMR spectral perturbations

and CPMG relaxation dispersion that show oxidizing potentials stimulate flexibility in ~34% of

the MIF structure, including its redox-active residues. However, the majority of these flexible

regions have not been confirmed to harbor any functional impact in prior work.

It remains unclear how this relatively compact (12.5 kDa monomer) and symmetrical

homotrimer accommodates a wide range of non-overlapping activities. A 2018 study showed

that nanosecond dynamics were important for organization of the MIF-CD74 binding interface,[6]

and studies of a MIF epitope containing its $^{56}CALC^{59}$ motif showed selective binding to the Fab

fragment of the BaxB01 antibody under oxidizing conditions, after rearrangement of the Cys

sulfhydryls.[22] Oxidizing conditions appear to promote a more flexible solvent channel, which we

recently showed to be central to an allosteric mechanism regulating tautomerase and CD74

activation involving motion on multiple timescales.[38] Given the number of MIF functions that have

yet to be explored at the molecular level and the fact that ~100 X-ray crystal structures of MIF

mutants, substrate-, and inhibitor-bound complexes have negligible variation in static structure

but vastly different NMR spectra, it is plausible, if not likely, that additional regions of functional

importance exist in MIF.

Here, we asked if changes in solution redox potential could reveal some of these latent

sites by modulating the structural and/or dynamic signatures of MIF. We experimentally altered

the redox potential of wt-MIF NMR samples, enabling us to identify redox-dependent structural

changes in residues that may be important for function. We mutated two such residues, K66 and

C80, that showed redox-sensitivity in wt-MIF and observed structural effects on MIF redox

behavior as well as altered functional responses. Having further confirmed the propensity of C80

for redox-driven modification by mass spectrometry, we speculate that MIF could act as a sensor

that toggles its conformation depending on its cellular environment to engage with varied binding

partners, as was previously suggested in a conformational control scheme for the $^{56}CALC^{59}$

motif. The more extensive but partially-overlapping set of residues affected by the redox

environment suggests the regulatory network controlling non-overlapping MIF functions may be

larger than previously reported, where additional "allosteric nodes" may be revealed only when

MIF is exposed to stimuli, such as environmental factors associated with inflammation, serving

as a natural means of activating latent residues to expand MIF functions under altered cellular

conditions. Speculating further, targeting small molecules to redox sensitive residues may

prevent MIF from engaging in certain downstream cascades. Future studies of protein-protein

interactions with supposed binding partners of MIF, such as thioredoxin[47,48] or ribosomal protein

S19,[41] as well as with small molecules, can evaluate this mechanism. Another interesting avenue

warranting further exploration is the redox-dependent interaction of MIF with its known inhibitors

or with novel compounds, as it presents an opportunity to preferentially target oxMIF, a form of

the protein that may appear predominately within the inflammatory state.

Materials and Methods

Protein expression, purification, and redox sample preparation

Wild-type (wt) and/or human MIF variants cloned into a pET11b vector were grown in

lysogeny broth (LB) for biochemical studies and in isotopically enriched M9 minimal medium

supplemented with $CaCl_2$, $MgSO_4$, and MEM vitamins for NMR, after adapting BL21(DE3) cells

to D_2O. Small cultures of MIF were grown overnight in LB at 37 °C and used to inoculate cultures

containing 50% D_2O the following morning, which were grown 8 – 10 hours at 37 °C and then

used to inoculate cultures containing 95% D_2O, which were incubated at 37 °C for 12 hours. The

cells were collected by centrifugation and resuspended in M9 medium (1 L, 100% D_2O)

supplemented with $^{15}NH_4Cl$ (Cambridge Isotope Laboratories) and $^{12}C_6H_{12}O_6$ as the sole

nitrogen and carbon sources, respectively. Cultures were grown to an OD_{600} of 0.8 – 1.0 and

induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG). Following 16-18 hours of post-

induction growth, the cells were harvested by centrifugation.

The cells were resuspended in ~30 mL of an ice-cold buffer of 20 mM Tris-HCl, 20 mM

NaCl, and 1 mM EDTA at pH 7.4 supplemented with protease inhibitors, lysed by ultrasonication,

clarified by centrifugation, filtered, and purified with a 120 mL Macro-Prep High Q

column(BioRad) pre-equilibrated with the lysis buffer. MIF does not bind the Q resin and is found

in the flow-through. MIF was purified to its final form on a HiLoad 26/600 Superdex 200 size

exclusion column (GE Healthcare). For NMR samples, MIF was dialyzed into a buffer containing

20 mM sodium phosphate, 1 mM EDTA, and 7.5% D_2O at pH 7.0. Final concentrations of MIF

were 0.5 – 1 mM determined using ε_{280} = 12,950 M^{-1} cm^{-1}.[49]

Oxidation of MIF was achieved by addition of H_2O_2 to a final concentration of 6 mM in an

aliquot of 0.5 – 1 mM MIF. Reduced MIF was prepared with a 6 mM final concentration of DTT.

Redox-neutral samples were prepared with equimolar amounts of DTT/H_2O_2. Stocks of the

oxidizing and reducing agents were freshly prepared in a degassed buffer of 20 mM sodium

phosphate, 1 mM EDTA, and 7.5% D_2O at pH 7.0. Following addition of oxidant or reductant to

MIF samples, the NMR tubes were briefly degassed with nitrogen and sealed with Parafilm

(Bemis). The samples were equilibrated in the presence of redox reagents for at least six hours.

Circular Dichroism (CD) Spectroscopy

CD spectra and thermal unfolding experiments were recorded on a JASCO J-815

spectropolarimeter equipped with a variable temperature Peltier device. Denaturation curves of

10 μM MIF were recorded at 218 nm in a 2 mm quartz cuvette. The temperature range for each

scan was 20 – 100 °C (293 K – 373 K). Thermodynamic parameters were extracted via nonlinear

curve fitting of CD data to **Equation 1** in GraphPad Prism:

$Ellipticity\ (T) = \dfrac{[(m_f T + b_f) + (m_u T + b_u)] exp\left[\left(-\frac{\Delta H_{D,vH}}{R}\right)\left(\frac{1}{T} - \frac{1}{T_m}\right)\right]}{1 + exp\left[\left(-\frac{\Delta H_{D,vH}}{R}\right)\left(\frac{1}{T} - \frac{1}{T_m}\right)\right]}$ (1)

where m_f, b_f, and m_u, b_u are the slopes and y-intercepts of the folded (low temperature) and

unfolded (high temperature) regions of the melting curve, R is the gas constant, T_m and $\Delta H_{D,vH}$

are the unfolding midpoint and van't Hoff enthalpy of denaturation at T_m, respectively. Free

energy analysis was performed as described elsewhere,[50] using values of ΔC_p, the unfolding

heat capacity, estimated from the report of Privalov.[51,52] The differences in apparent unfolding

enthalpies, entropies, and free energies at a reference temperature (T_{REF}) of 298 K were

calculated as;

$$\Delta H_{D,app} = \Delta H_{D,vH} + \Delta C_p(T_{REF} - T_m) \tag{2}$$

$$\Delta S_{D,app} = \frac{\Delta H_{D,vH}}{T_m} + \Delta C_p ln\left(\frac{T_{REF}}{T_m}\right) \tag{3}$$

$$\Delta G(T) = \Delta H_{D,vH}\left(1 - \frac{T_{REF}}{T_m}\right) + \Delta C_p\left[(T_{REF} - T_m) - T_{REF}ln\left(\frac{T_{REF}}{T_m}\right)\right] \tag{4}$$

These data are reported as "apparent" values due to the ~90% reversibility of the MIF unfolding

transition.

NMR Spectroscopy

NMR experiments were performed on a Bruker Avance NEO 600 MHz or Bruker

Avance III HD 850 MHz spectrometer at 30 °C. NMR data were processed using NMRPipe[53]

and analyzed in Sparky[54] along with in-house scripts. The pH of NMR samples was monitored

after addition of redox reagents to ensure that NMR spectral changes were due to

reduction/oxidation. NMR assignments for wt-MIF and the K66A and C80A variants were

confirmed from standard triple resonance experiments. Combined chemical shift perturbations

($\Delta\delta$) were determined by $\Delta\delta = \sqrt{(\delta_{HN}^2) + (\delta_N^2/25)}$. The HSQC of oxidized MIF was processed

with an exponential window function in both the direct and indirect dimensions in

NMRPipe. The peak intensity and volume for each pair of slow exchanging peaks were

calculated in SPARKY, where peak volume was calculated by integrating each peak as a

Lorentzian. The minor state populations were estimated by taking the volume of the minor

peak and dividing by the total volume of the slow exchanging pairs.

NMR spin relaxation experiments were performed using TROSY-based pulse

sequences adapted from Palmer and coworkers.[55] Relaxation data were acquired with the ^{1}H

and ^{15}N carriers set to the water resonance and 120 ppm, respectively. Longitudinal relaxation

rates (R_1) were measured with T_1 delays of 0, 20, 60, 100, 200, 600, 800, and 1200 ms.

Transverse relaxation rates (R_2) were collected with 1.0 ms spacing between 180° CPMG

pulses at total relaxation delays of 0, 1, 2, 4, 8, and 10, and 12 ms. The recycle delay in these

experiments was 2.5 s. Longitudinal and transverse relaxation rates were extracted by non-

linear least squares fitting of the peak heights (major peaks in cases of slow exchange) to a

single exponential decay using in-house software. Uncertainties in these rates were

determined from replicate spectra. The heteronuclear cross-relaxation rate (^{1}H-[^{15}N] NOE) was

obtained by interleaving pulse sequences with and without proton saturation and calculated

from the ratio of peak heights from these experiments. Carr-Purcell-Meiboom-Gill (CPMG)

NMR experiments were performed at 30 °C with a constant relaxation period of 40 ms, a 2.0 s

recycle delay, and τ_{cp} points of 0, 25, 50, 75, 100, 150, 250, 500, 750, 800, 900, and 1000 ms.

Relaxation dispersion profiles were generated by plotting R_2 vs. $1/\tau_{cp}$ and exchange

parameters were obtained from fits of these data carried out with in-house scripts.

Uncertainties were obtained from replicate spectra.

NMR spin-relaxation rates were fit to one of five semi-empirical forms of the spectral

density function using model-free analysis.[56,57] Fitting of motional parameters was performed in

RELAX.[58,59] The criteria for inclusion of residues in the diffusion tensor estimate relied on the

method of Tjandra and coworkers.[47] N-H bond lengths were assumed to be 1.02 Å and the ^{15}N

chemical shift anisotropy was set to -160 ppm. During the model selection process, the diffusion

tensor parameters were optimized simultaneously, and model selection was repeated until the

optimized tensor parameters and order parameter (S^2) did not differ from those of the previous

iteration.

Mass Spectrometry

Cysteine and cystine residues of purified MIF equilibrated under each of the redox

conditions were alkylated in a stepwise process with iodoacetamide (IAA) followed by *N*-

ethylmaleimide (NEM). Parallel reaction monitoring-mass spectrometry captured the oxidation

states of C80 under different redox conditions. Skyline software[60] was used to calculate the

relative abundance of cysteine and cystine at the C80 position, indicated by binding to IAA or

NEM. Additional methodological details can be found in the **Supporting Information**.

In Vivo Neutrophil Recruitment Assay

All animal studies were approved by the Institutional Animal Care and Use Committee of

Cooper University Healthcare, Camden, NJ. The wt mice of genetic background strain

(C57BL6/J) were purchased from the Jackson Laboratory (Bar Harbor, ME). Mice were housed

at the pathogen-free animal facility at Cooper University Healthcare. All experiments were done

in 9-10 week old male mice. Mice were administered a one-time intratracheal instillation of 100

μl of normal saline alone (vehicle) or normal saline solution containing 1 μg of wt-MIF, K66A MIF

or C80A MIF. These mice were sacrificed after 6 h of vehicle only or vehicle + experimental

agent administration. Bronchoalveolar lavage (BAL) was performed by cannulating the trachea

with a blunt 22-gauge needle and lavaging both lungs with 800 μl of sterile PBS solution.

Bronchoalveolar lavage fluid (BALF) was harvested, and total cell counts in BALF were

determined using the TC20 automated cell counter (Bio-Rad Laboratories, Inc., Hercules, CA).

The differential cell counts were performed on cells cytocentrifuged onto glass slides (Fisher

Scientific) stained with the Hema 3 Staining System (Fisher Diagnostics, Middletown, VA) and

cell differential was tabulated using light microscopy. Total protein concentration in the BAL fluid

was measured using the Pierce^{TM} BCA assay kit (Thermo Scientific, Rockford, IL), as previously

described.[6]

Acknowledgments

The authors thank J. Patrick Loria for helpful discussions. This work was supported by Rhode

Island Foundation Grant GR5290658 and funds from the Office of the Vice President for

Research at Brown University (to GPL).

Competing Interests

The authors declare no competing interests.

References

(1)   Bernhagen, J. r.; Bucala, R.: MIF Family Cytokines in Innate Immunity and Homeostasis.
In *Progress in Inflammation Research,*; 1st ed.; Springer International Publishing : Imprint: Springer,:
Cham, 2017.
(2)   Leng, L.; Metz, C. N.; Fang, Y.; Xu, J.; Donnelly, S.; Baugh, J.; Delohery, T.; Chen, Y.;
Mitchell, R. A.; Bucala, R. MIF signal transduction initiated by binding to CD74. *J Exp Med* **2003**, *197*,
1467-1476.
(3)   Bernhagen, J.; Krohn, R.; Lue, H.; Gregory, J. L.; Zernecke, A.; Koenen, R. R.; Dewor,
M.; Georgiev, I.; Schober, A.; Leng, L.; Kooistra, T.; Fingerle-Rowson, G.; Ghezzi, P.; Kleemann, R.;
McColl, S. R.; Bucala, R.; Hickey, M. J.; Weber, C. MIF is a noncognate ligand of CXC chemokine
receptors in inflammatory and atherogenic cell recruitment. *Nat Med* **2007**, *13*, 587-596.
(4)   Meyer-Siegler, K. L.; Iczkowski, K. A.; Leng, L.; Bucala, R.; Vera, P. L. Inhibiton of
Macrophage Migration Inhibitory Factor or its Receptor (CD74) Attenuates Growth and Invasion of DU-
145 Prostate Cancer Cells. *J. Immunol.* **2006**, *177*, 8730-8739.

415   (5) Cournia, Z.; Leng, L.; Gandavadi, S.; Du, X.; Bucala, R.; Jorgensen, W. L. Discovery of
416 Human Macrophage Migration Inhibitory Factor (MIF)-CD74 Amtagonists via Virtual Screening. *J. Med.*
417 *Chem.* **2009**, *52*, 416-424.

418   (6) Pantouris, G.; Ho, J.; Shah, D.; Syed, M. A.; Leng, L.; Bhandari, V.; Bucala, R.; Batista,
419 V. S.; Loria, J. P.; Lolis, E. J. Nanosecond Dynamics Regulate the MIF-induced Activity of CD74. *Angew.*
420 *Chem. Int. Ed.* **2018**, *57*, 7116-7119.

421   (7) Mizue, Y.; Ghani, S.; Leng, L.; McDonald, C.; Kong, P.; Baugh, J.; Lane, S. J.; Craft, J.;
422 Nishihira, J.; Donnelly, S. C.; Zhu, Z.; Bucala, R. Role for Macrophage Migration Inhibitory Factor in
423 Asthma. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 14410-14415.

424   (8) Joshi, P. C.; Poole, G. V.; Sachdev, V.; Zhou, X.; Jones, Q. Trauma Patients with Positive
425 Cultures have Higher Levels of Circulating Macrophage Migration Inhibitory Factor. *Res. Commun. Mol.*
426 *Pathol. Pharmacol.* **2000**, *107*, 13-20.

427   (9) Kleeman, R.; Hausser, A.; Geiger, G.; Mischke, R.; Burger-Kentischer, A.; Flieger, O.;
428 Johannes, F. J.; Roger, T.; Calandra, T.; Kapurniotu, A.; Grell, M.; Finkelmeier, D.; Brunner, H.;
429 Bernhagen, J. Intracellular Action of the Cytokine MIF to Modulate AP-1 Activity and the Cell Cycle
430 Through JAB-1. *Nature* **2000**, *408*, 211-216.

431   (10) Calandra, T.; Bernhagen, J.; Metz, C. N.; Spiegel, L. A.; Bacher, M.; Donnelly, T.; Cerami,
432 A.; Bucala, R. MIF as a Glucocorticoid-Induced Modulator of Cytokine Production. *Nature* **1995**, *377*,
433 68-71.

434   (11) Thiele, M.; Bernhagen, J. Link between Macrophage Migration Inhibitory Factor and
435 Cellular Redox Regulation. *Antioxid. Redox Signal.* **2005**, *7*, 1234-1248.

436   (12) Morand, E. F.; Leech, M.; Bernhagen, J. MIF: A New Cytokine Link between Rheumatoid
437 Arthritis and Atherosclerosis. *Nat. Rev. Drug Discov.* **2006**, *5*, 399-410.

438   (13) de Jong, Y. P.; Abadia-Molina, A. C.; Satoskar, A. R.; Clarke, K.; Rietdijk, S. T.; Faubion,
439 W. A.; Mizoguchi, E.; Metz, C. N.; Alsahli, M.; ten Hove, T.; Keates, A. C.; Lubetsky, J. B.; Farrell, R.
440 J.; Michetti, P.; van Deventer, S. J.; Lolis, E.; David, J. R.; Bhan, A. K.; Terhorst, C. Development of
441 Chronic Colitis is Dependent on the Cytokine MIF. *Nat. Immunol.* **2001**, *2*, 1061-1066.

442   (14) Makhija, R.; Kingsnorth, A.; Demaine, A. Gene Polymorphisms of the Macrophage
443 Migration Inhibitory Factor and Acute Pancreatitis. *J.Pancreas* **2007**, *8*, 289-295.

444   (15) Donnelly, S. C.; Haslett, C.; Reid, P. T.; Grant, I. S.; Wallace, W. A.; Metz, C. N.; Bruce,
445 L. J.; Bucala, R. Regulatory Role for Macrophage Migration Inhibitory Factor in Acute Respiratory
446 Distress Syndrome. *Nat. Med.* **1997**, *3*, 320-323.

447   (16) Bucala, R.; Donnelly, S. C. Macrophage Migration Inhibitory Factor: A Probable Link
448 between Inflammation and Cancer. *Immunity* **2007**, *26*, 281-285.

449   (17) Balogh, K. N.; Templeton, D. J.; Cross, J. V. Macrophage Migration Inhibitory Factor
450 Protects Cancer Cells from Immunogenic Cell Death and Impairs Anti-tumor Responses. *PLOS One* **2018**,
451 *13*, e0197702.

452   (18) Richard, V.; Kindt, N.; Saussez, S. Macrophage Migration Inhibitory Factor Involvement
453 in Breast Cancer. *Int. J. Oncol.* **2015**, *47*, 1627-1633.

454   (19) Agarwal, R.; Whang, D. H.; Alvero, A. B.; Visintin, I.; Lai, Y. L.; Segal, E. A.; Schwartz,
455 P.; Ward, D.; Rutherford, T.; Mor, G. Macrophage Migration Inhibitory Factor Expression in Ovarian
456 Cancer. *Am. J. Obstet. Gynecol.* **2007**, *196*, 348.e341-348.e345.

457   (20) Nobre, C. C. G.; de Arajuo, J. M. G.; Fernandes, T. A. A. M.; Cobucci, R. N. O.; Lanza,
458 D. C. F.; Andrade, V. S.; Fernandes, J. V. Macrophage Migration Inhibitory Factor (MIF): Biological
459 Activities and Relation with Cancer. *Pathol. Oncol. Res.* **2017**, *23*, 235-244.

(21)   Bleilevens, C.; Soppert, J.; Hoffmann, A.; Breuer, T.; Bernhagen, J.; Martin, L.; Stiehler, L.; Marx, G.; Dreher, M.; Stoppe, C.; Simon, T. P. Macrophage Migration Inhibitory Factor (MIF) Plasma Concentration in Critically Ill COVID-19 Patients: A Prospective Observational Study. *Diagnostics (Basel)* **2021**, *11*.

(22)   Schinagl, A.; Kerschbaumer, R. J.; Sabarth, N.; Douillard, P.; Scholz, P.; Voelkel, D.; Hollerweger, J. C.; Goettig, P.; Brandstetter, H.; Scheiflinger, F.; Thiele, M. Role of the Cysteine 81 Residue of Macrophage Migration Inhibitory Factor as a Molecular Redox Switch. *Biochemistry* **2018**, *57*, 1523-1532.

(23)   Kleemann, R.; Kapurniotu, A.; Frank, R. W.; Gessner, A.; Mischke, R.; Flieger, O.; Juttner, S.; Brunner, H.; Bernhagen, J. Disulfide analysis reveals a role for macrophage migration inhibitory factor (MIF) as thiol-protein oxidoreductase. *J Mol Biol* **1998**, *280*, 85-102.

(24)   Kleemann, R.; Mischke, R.; Kapurniotu, A.; Brunner, H.; Bernhagen, J. Specific reduction of insulin disulfides by macrophage migration inhibitory factor (MIF) with glutathione and dihydrolipoamide: potential role in cellular redox processes. *FEBS Lett* **1998**, *430*, 191-196.

(25)   Mittal, M.; Siddiqui, M. R.; Tran, K.; Reddy, S. P.; Malik, A. B. Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal* **2014**, *20*, 1126-1167.

(26)   Fahy, J. V.; Corry, D. B.; Boushey, H. A. Airway Inflammation and Remodeling in Asthma. *Curr. Opin. Pulm. Med.* **2000**, *6*, 15-20.

(27)   Fitzpatrick, A. M.; Teague, W. G.; Holguin, F.; Yeh, M.; Brown, L. A. Airway Glutathione Homeostasis is Altered in Children with Severe Asthma: Evidence for Oxidant Stress. *J. Allergy Clin. Immunol.* **2009**, *123*, 146-152.

(28)   Watson, W.; Ritzenthaler, J. D.; Roman, J. Lung Extracellular Matrix and Redox Regulation. *Redox Biol.* **2016**, *8*, 303-315.

(29)   Kinnula, V. L.; Fattman, C. L.; Tan, R. J.; Oury, T. D. Oxidative Stress in Pulmonary Fibrosis: A Possible Role for Redox Modulatory Therapy. *Am. J. Respir. Crit. Care Med.* **2005**, *172*, 417-422.

(30)   Bakalova, R.; Zhelev, Z.; Aoki, I.; Saga, T. Tissue Redox Activity as a Hallmark of Carcinogenesis: From Early to Terminal Stages of Cancer. *Clin. Cancer Res.* **2013**, *19*, 2503-2517.

(31)   Kneller, J. M.; Lu, M.; Bracken, C. An Effective Method for the Discrimination of Motional Anisotropy and Chemical Exchange. *J. Am. Chem. Soc.* **2002**, *124*, 1852-1853.

(32)   Pantouris, G.; Syed, M. A.; Fan, C.; Rajasekaran, D.; Cho, T. Y.; Rosenberg, E. M., Jr.; Bucala, R.; Bhandari, V.; Lolis, E. J. An Analysis of MIF Structural Features that Control Functional Activation of CD74. *Chem Biol* **2015**, *22*, 1197-1205.

(33)   Lubetsky, J. B.; Dios, A.; Han, J.; Aljabari, B.; Ruzsicska, B.; Mitchell, R.; Lolis, E.; Al-Abed, Y. The tautomerase active site of macrophage migration inhibitory factor is a potential target for discovery of novel anti-inflammatory agents. *J Biol Chem* **2002**, *277*, 24976-24982.

(34)   Takahashi, K.; Koga, K.; Linge, H. M.; Zhang, Y.; Lin, X.; Metz, C. N.; Al-Abed, Y.; Ojamaa, K.; Miller, E. J. Macrophage CD74 contributes to MIF-induced pulmonary inflammation. *Respir Res* **2009**, *10*, 33.

(35)   Mehrad, B.; Strieter, R. M.; Moore, T. A.; Tsai, W. C.; Lira, S. A.; Standiford, T. J. CXC chemokine receptor-2 ligands are necessary components of neutrophil-mediated host defense in invasive pulmonary aspergillosis. *J Immunol* **1999**, *163*, 6086-6094.

(36)   Nguyen, M. T.; Beck, J.; Lue, H.; Funfzig, H.; Kleemann, R.; Koolwijk, P.; Kapurniotu, A.; Bernhagen, J. A 16-residue peptide fragment of macrophage migration inhibitory factor, MIF-(50-65), exhibits redox activity and has MIF-like biological functions. *J Biol Chem* **2003**, *278*, 33654-33671.

(37)   Takahashi, M.; Nishihira, J.; Shimpo, M.; Mizue, Y.; Ueno, S.; Mano, H.; Kobayashi, E.;
Ikeda, U.; Shimada, K. Macrophage migration inhibitory factor as a redox-sensitive cytokine in cardiac
myocytes. *Cardiovasc Res* **2001**, *52*, 438-445.

(38)   Pantouris, G.; Khurana, L.; Ma, A.; Skeens, E.; Reiss, K.; Batista, V. S.; Lisi, G. P.; Lolis,
E. J. Regulation of MIF Enzymatic Activity by an Allosteric Site at the Central Solvent Channel. *Cell*
Chem Biol **2020**, *27*, 740-750 e745.

(39)   Calandra, T.; Roger, T. Macrophage migration inhibitory factor: a regulator of innate
immunity. *Nat Rev Immunol* **2003**, *3*, 791-800.

(40)   Jung, H.; Seong, H. A.; Ha, H. Critical role of cysteine residue 81 of macrophage migration
inhibitory factor (MIF) in MIF-induced inhibition of p53 activity. *J Biol Chem* **2008**, *283*, 20383-20396.

(41)   Filip, A. M.; Klug, J.; Cayli, S.; Frohlich, S.; Henke, T.; Lacher, P.; Eickhoff, R.; Bulau,
P.; Linder, M.; Carlsson-Skwirut, C.; Leng, L.; Bucala, R.; Kraemer, S.; Bernhagen, J.; Meinhardt, A.
Ribosomal protein S19 interacts with macrophage migration inhibitory factor and attenuates its pro-
inflammatory function. *J Biol Chem* **2009**, *284*, 7977-7985.

(42)   Kassaar, O.; Pereira Morais, M.; Xu, S.; Adam, E. L.; Chamberlain, R. C.; Jenkins, B.;
James, T. D.; Francis, P. T.; Ward, S.; Williams, R. J.; van den Elsen, J. Macrophage Migration Inhibitory
Factor is subjected to glucose modification and oxidation in Alzheimer's Disease. *Sci Rep* **2017**, *7*, 42874.

(43)   Maurice, M. M.; Nakamura, H.; van der Voort, E. A.; van Vliet, A. I.; Staal, F. J.; Tak, P.
P.; Breedveld, F. C.; Verweij, C. L. Evidence for the role of an altered redox state in hyporesponsiveness
of synovial T cells in rheumatoid arthritis. *J Immunol* **1997**, *158*, 1458-1465.

(44)   Wu, S. M.; Pizzo, S. V. alpha(2)-Macroglobulin from rheumatoid arthritis synovial fluid:
functional analysis defines a role for oxidation in inflammation. *Arch Biochem Biophys* **2001**, *391*, 119-
126.

(45)   Taylor, S. L.; Crawley-Snowdon, H.; Wagstaff, J. L.; Rowe, M. L.; Shepherd, M.;
Williamson, R. A.; Howard, M. J. Measuring protein reduction potentials using 15N HSQC NMR
spectroscopy. *Chem Commun (Camb)* **2013**, *49*, 1847-1849.

(46)   Zimmermann, J.; Kuhne, R.; Sylvester, M.; Freund, C. Redox-regulated conformational
changes in an SH3 domain. *Biochemistry* **2007**, *46*, 6971-6977.

(47)   Tjandra, N.; Feller, S. E.; Pastor, R. W.; Bax, A. Rotational diffusion anisotropy of human
ubiquitin from N-15 NMR relaxation. *J Am Chem Soc* **1995**, *117*, 12562-12566.

(48)   Son, A.; Kato, N.; Horibe, T.; Matsuo, Y.; Mochizuki, M.; Mitsui, A.; Kawakami, K.;
Nakamura, H.; Yodoi, J. Direct Association of Thioredoxin-1 (TRX) with Macrophage Migration
Inhibitory Factor (MIF): Regulatory Role of TRX on MIF Internalization and Signaling *Antioxid. Redox*
Signal. **2009**, *11*, 2595-2605.

(49)   Lubetsky, J. B.; Swope, M.; Dealwis, C.; Blake, P.; Lolis, E. Pro-1 of macrophage
migration inhibitory factor functions as a catalytic base in the phenylpyruvate tautomerase activity.
Biochemistry **1999**, *38*, 7346-7354.

(50)   Zhong, F.; Lisi, G. P.; Collins, D. P.; Dawson, J. H.; Pletneva, E. V. Redox-dependent
stability, protonation, and reactivity of cysteine-bound heme proteins. *Proc Natl Acad Sci U S A* **2014**,
111, E306-315.

(51)   Makhatadze, G. I.; Privalov, P. L. Heat capacity of proteins. I. Partial molar heat capacity
of individual amino acid residues in aqueous solution: hydration effect. *J Mol Biol* **1990**, *213*, 375-384.

(52)   Privalov, P. L.; Makhatadze, G. I. Contribution of hydration and non-covalent interactions
to the heat capacity effect on protein unfolding. *J Mol Biol* **1992**, *224*, 715-723.

(53)   Delaglio, F.; Grzesiek, S.; Vuister, G. W.; Zhu, G.; Pfeifer, J.; Bax, A. NMRPipe: a
multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* **1995**, *6*, 277-293.

(54)   Lee, W.; Tonelli, M.; Markley, J. L. NMRFAM-SPARKY: enhanced software for
biomolecular NMR spectroscopy. *Bioinformatics* **2015**, *31*, 1325-1327.

(55)   Loria, J. P.; Rance, M.; Palmer, A. G. A relaxation-compensated Carr-Purcell-Meiboom-
Gill sequence for characterizing chemical exchange by NMR spectroscopy. *J Am Chem Soc* **1999**, *121*,
2331-2332.

(56)   Clore, G. M.; Szabo, A.; Bax, A.; Kay, L. E.; Driscoll, P. C.; Gronenborn, A. M. Deviations
from the Simple 2-Parameter Model-Free Approach to the Interpretation of N-15 Nuclear Magnetic-
Relaxation of Proteins. *J Am Chem Soc* **1990**, *112*, 4989-4991.

(57)   Lipari, G.; Szabo, A. Model-Free Approach to the Interpretation of Nuclear Magnetic-
Resonance Relaxation in Macromolecules .2. Analysis of Experimental Results. *J Am Chem Soc* **1982**,
104, 4559-4570.

(58)   d'Auvergne, E. J.; Gooley, P. R. Optimisation of NMR dynamic models I. Minimisation
algorithms and their performance within the model-free and Brownian rotational diffusion spaces. *J
Biomol NMR* **2008**, *40*, 107-119.

(59)   d'Auvergne, E. J.; Gooley, P. R. Optimisation of NMR dynamic models II. A new
methodology for the dual optimisation of the model-free parameters and the Brownian rotational diffusion
tensor. *J Biomol NMR* **2008**, *40*, 121-133.

(60)   MacLean, B.; Tomazela, D. M.; Shulman, N.; Chambers, M.; Finney, G. L.; Frewen, B.;
Kern, R.; Tabb, D. L.; Liebler, D. C.; MacCoss, M. J. Skyline: an open source document editor for creating
and analyzing targeted proteomics experiments. *Bioinformatics* **2010**, *26*, 966-968.