1   **Title: Systems analysis shows a role of cytophilic antibodies in shaping innate**

2   **tolerance to malaria**

Authors: Maximilian Julius Lautenbach[1,2], Victor Yman[1,3], Nadir Kadri[1,2,4], David Fernando

Plaza[1,2], Sina Angenendt[1,2], Klara Sondén[1,2], Anna Färnert[1,2], Christopher Sundling[1,2]*

Affiliations:

[1]Division of Infectious Diseases, Department of Medicine Solna, and Center for Molecular

Medicine, Karolinska Institutet, Stockholm, Sweden.

[2]Department of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden.

[3]Department of Infectious Diseases, South Stockholm Hospital, Stockholm, Sweden

[4]Science for Life Laboratory, Department of Medicine Solna, Karolinska Institute.

*Corresponding author. Email: christopher.sundling@ki.se (C.S.)

One Sentence Summary: A systems immunology analysis on natural malaria sheds light on

disease tolerance mechanism associated with gamma delta T cell expansion (134/150 with

spaces)

Abstract:

The mechanism of acquisition and maintenance of natural immunity against *Plasmodium falciparum* malaria remains unclear. Although, clinical immunity develops over time with repeated malaria episodes, disease tolerance is more rapidly acquired compared to protective immunity. It remains unclear, how pre-existing immune responses impacts the mechanism responsible for disease tolerance. Here, we investigated a cohort of returning travelers treated for acute symptomatic *P. falciparum* malaria, either infected for the first time, or with a previous history of malaria. Through repeated sampling over one year in a malaria free setting, we were able to study the acute and longitudinal effects of the infection. We combined comprehensive immune cell and plasma protein profiling with integrated and data driven analysis, describing the immune landscape from acute disease to one year after infection. We identified a strong association between pro-inflammatory signatures and γδ T cell expansion. The association was significantly impacted by previous exposure to malaria, resulting in a dampened pro-inflammatory response, which translated to reduced Vδ2$^+$ γδ T cell expansion compared to primary infected individuals. The dampened inflammatory signal was associated with early expansion of FcγRIII+ monocytes and parasite-specific antibodies of IgG1 and IgG3 isotypes.

Our data suggest that the interplay of FcγRIII+ monocytes and a cytophilic parasite-specific IgG during the early blood stage infection lead to lower parasitemia and a dampened pro-inflammatory response with reduced γδ T cell expansion. This enhanced control and reduced inflammation points to a potential mechanism on how tolerance is established following repeated malaria exposure. (244/250 words)

Main Text:

INTRODUCTION

Malaria remains a global burden with an estimated 229 million malaria cases leading to
approximately 409 000 deaths in 2019 *(1)*. Compared to other pathogens, immunity to malaria is
slow to develop *(2)*. In malaria endemic areas, partial immunity is acquired over time after repeated
infections. Clinical symptoms are reduced while low grade subclinical infections remain and allow
for continued parasite transmission *(2, 3)*. However, protection from severe forms of malaria,
despite high levels of parasitemia, seem to develop already after a few infections *(4)*. A recent
study by Nahrendorf and colleagues suggests a tolerance mechanism where better control of host-
damaging factors that are part of the natural immune response to infection are an underlying cause
for the reduced severity *(5)*. Currently no study has investigated how long tolerance persists and
how it is maintained in response to malaria.

Clinical symptoms are due to the blood stage of the infection, where merozoites invade
erythrocytes that then sequester, followed by parasite multiplication before bursting out to infect
new erythrocytes *(6)*. The clinical manifestations are partly due to the strong innate pro-
inflammatory response, during the parasite blood stage/to the *P. falciparum* blood stage *(7)*.

Much of our current understanding of the immune responses to *P. falciparum* infection is based
on hypothesis-driven research, where selected cell subsets, inflammatory markers, or clinical
features are investigated at a time *(8, 9)*. Improved technology and bioinformatic tools enable the
analysis of high-dimensional parameters in limited sample material, giving us the opportunity to
study complex immune responses at a systems level to provide new insights into complex
immunological networks that represent the immune response during infection *(10)*. Systems level
approaches to study febrile/non-febrile children in malaria endemic countries *(11)* as well as the
comparison of malaria naïve and Africans in controlled human malaria infection (CHMI) studies
(12) have generated valuable findings and new hypotheses. While studies in endemic populations
allow investigation of acute naturally acquired malaria and persistent (largely asymptomatic)
infections in the context of previous exposure. Despite CHMI studies enable more control over
prior parasite exposure, infectious dose, and follow-up after infection, although only in the context
of very low level parasitemia and often before symptoms having appeared. Although compensating
for some limitations, CHMI cannot fully mirror a natural infection, due to early treatment not
allowing observation of potential effects derived from a strong natural symptomatic infection. This

leaves a knowledge gap on how natural infection affects the immune response in the absence of

potential parasite re-exposure. This gap can be filled by investigating the immune response after

naturally acquired malaria in individuals leaving the endemic area, seeking healthcare in a setting

without risk of re-exposure.

Here, we study a prospective cohort of returning travelers treated for acute *P. falciparum* malaria

at the Karolinska University Hospital in Sweden, followed over one year after infection. Using a

systems immunology approach and data-driven analysis, we combined plasma protein and cell

profiling to study dynamic changes of the immune response over time in these patients. Based on

our results, we propose a model where antibody-derived memory modulates the pro-

inflammatory cytokine response which in turn impact $\gamma\delta$ T cell expansion and improve disease

tolerance.

RESULTS

Subhead 1: Longitudinal profiling of peripheral blood – Immune dynamics after natural

infection in a cohort of returning travelers

We aimed to comprehensively profile the immune response dynamics longitudinally after natural

infection with *P. falciparum* malaria, to appreciate the immunological changes occurring during

the acute disease and up to one year after diagnosis. We included 53 returning travelers, repeatedly

sampled in a non-endemic setting over one year after hospital admission *(13–16)* (Fig. 1A).

To profile the immune response in the cohort over time, we performed broad FACS-based

immunophenotyping using a 17-marker panel and gated for subsets of monocytes ($CD14^+$), T cells

($CD3^+CD56^-$), and NK cell ($CD3^-CD56^+$) (Fig. S1A). We also included data from B cell subsets

previously phenotyped from the same donors *(16)* (Fig. S1B). Plasma protein analysis was

performed using the Target 96-plex Inflammation panel from Olink Proteomics *(17)* targeting 92

immune response-related proteins. In total, we profiled 182 samples from 53 subjects, creating

more than 23,000 data points (Fig. 1B).

The cohort consisted of individuals infected with *P. falciparum* malaria for the first time (primary

infected, n = 17) and those infected before, having grown up in malaria endemic areas and reported

previous malaria (previously exposed, n = 36) (Fig. 1C). Comparing individuals with primary

infection versus previous exposure enabled us to investigate potential memory effect on the

immune response. The individuals with previous exposure moved from malaria endemic areas to

non-endemic Sweden on average 11.5 years before now experiencing acute malaria after visiting

an endemic area (median = 11.5, range 0-46 years, Tab. S1).

Although all individuals sought healthcare, primary infected patients had significantly higher

levels of parasitemia at hospital admission (median 1.10 vs 0.25 % infected red blood cells, p =

0.038, Fig1D), more signs of severe malaria (Tab. S1) and were on average admitted to hospital

care significantly longer than previously exposed patients (median 3 vs 2 days, p = 0.002, Fig1E,

Tab. S1).

Fig. 1. *Systems level profiling of peripheral blood from individuals of a prospective cohort of returning travelers with P. falciparum malaria. (A) Overview of prospective longitudinal malaria cohort of returning travelers, (B) temporal sample information symptom onset (black), admission to Karolinska University Hospital (blue), six sampling time points (x) for immune profiling (acute, after 10 days, 1 month, 3 months, 6 months, 1 year) and convalescence average (green lines). Patients with primary infection (n = 17) are colored in purple, and patients with previous malaria exposure (n = 36) are colored in orange. (C) A total of 53 returning travelers were longitudinally profiled for plasma proteins using Olink Proteomics PEA platform (n = 171 samples, i = 92 proteins) and their PBMC immune cells (n = 137 samples, i = 54 subsets), generating 23,130 unique data points. Comparison of (D) parasitemia at diagnosis (percentage of infected red blood cells) and (E) length of hospital admission (days) for primary infected and previously exposed individuals. Statistical differences between groups were assessed using the non-parametric Wilcoxon-test. *p < 0.05, **p < 0.01.*

Subhead 2: Integrated analysis of immune response on time axis after symptom onset

To explore the immune response dynamics after natural *P. falciparum* malaria on a systems level
and in a data driven manner, we used multi-omics factor analysis (MOFA). The unsupervised
nature of MOFA allows the model to capture both biological and technical variability in the low-
dimensional factors space *(18, 19)*. Here we used MEFISTO, a recent extension of MOFA which
allows for multi-omics integration of data while controlling for time-dependent variance in
repeated samples *(20)*. MOFA can disentangle the sources of heterogeneity in diverse data types
and accept missing data points using matrix factorization (Fig. S2A). Using MEFISTO, the dataset
was described by two latent factors. These factors explained 37% to 60% and 58% to 72 % of the
immune cell subset and plasma protein dynamics, respectively (Fig. 2A). Factor1 was primarily
associated with time-dependent changes in the immune response to the infection (Fig. 2B and
S2B). Positive factor values were associated with the acute phase and negative factor values were
associated with the time after treatment and transition towards convalescence (Fig. 2B and S2C).
The positive factor values were driven by increased levels of IL10, pro-inflammatory cytokines
such as IFN-gamma, IL-6, TNF, CCL3, IL8, CDCP1, and chemokines such as CXCL9, CXCL10,
CXCL11. For immune cells, $CD4^+$ and $CD8^+$ T cells expressing activation markers CD38 and
HLA-DR, intermediate monocytes ($CD14^+CD16^+$) and plasmablasts
($CD19^+CD20^{lo}CD38^{hi}CD27^{hi}$) were associated with Factor1 (Fig. 2C). The negative factor values
were associated with the transition phase towards convalescence. Here, several subsets of NK cells
expressing CD57, CD8 T effector memory cells, and γδ T cells were variables driving Factor1 for
the cellular immune response after treatment (Fig. 2C). We confirmed the accuracy of the model
parameters by comparing measured data between the acute and convalescent samples (internal
control; Fig. S2C-F) and to healthy control samples (external control; Fig. S3A-D).

We then used the integrated MEFISTO model to impute missing data points (plasma proteins data
modality, n = 11; immune cell modality, n = 64; Fig. S2A) to generate a heatmap of the integrated
immune landscape, visualizing the longitudinal immune system dynamics after disease. The
landscape dynamics, show a rapidly contracting pro-inflammatory response after treatment with a
temporary increase in primarily activated T cell subsets followed by a transition into a more long-
term post-infection response (Fig. 2D and Fig. S3).

In summary, our integrated analysis of immune cell subsets and plasma protein data after symptom onset allows us to draw a data driven descriptive immune landscape, describing the transition from the clinical acute phase towards convalescence over one year.

Fig. 2. ***Integrated analysis of immune response dynamics for one year after malaria.*** *Integrated multi omics factor analysis method for the Functional Integration of Spatial and Temporal Omics data (MEFISTO) model was utilized to integrate data modalities, plasma proteins and manually gated cell subsets, respectively, with their temporal covariate of time after symptom onset. (**A**) Variance explained by the MEFISTO model. Differences due to previous parasite exposure were assigned to groups in order to model group-specific time-dependent immune response dynamics. (**B**) Plotting Factor1 value against the time after symptom onset (weeks) highlights the captured variance in Factor1 with positive factor values associated with acute phase samples and negative factor values associated with non-acute phase samples. (**C**) Top10 features of Factor1 for each of the immune system views based on negative and positive scaled feature weights - immune cells (purple) and plasma proteins (green). (**D**) The heatmaps visualize immune response dynamics for*

Factor1 driving features. The characteristic immune variables for each data modality, aligned to time after symptom onset, reveals contrasting patterns characterizing the transition from acute phase towards convalescence. Missing data points have been imputed based on model Factor1. Cell counts and relative protein levels are shown as rows, and each column represents an individual patient. Rows were clustered using Euclidean distance and column were arranged according to MEFISTO model group-aligned time after onset. Cell counts and protein level values were converted to z-scores.

Subhead 3: Relationship between acute cytokine milieu and cellular response

Cytokines and chemokines are known to orchestrate an immunological response to drive recruitment, activation, and subsequently proliferation of immune cell populations. Hence, we sought to analyze possible associations between levels of the acute phase responses and immune cells subsets, characterizing the transition from acute towards convalescence.

Focusing on the top immune variables determined by the integrated model, we correlated acute samples with all remaining sample time points using spearman rank correlation (Fig. 3A). We discovered strong and significant correlations (|rho| > 0.7, after adjusted FDR < 0.05) between acute phase cytokines and immune cell subsets at all sample time points after treatment (D10, M3, M6 and Y1) (Fig. 3B).

In particular, the numbers of γδ T cell subsets at the 10-day sample time-point were strongly associated with several pro-inflammatory cytokines at the acute time-point. The total level of γδ T cells positively correlated with TNF (rho = 0.704, p = 0.049), IL6 (rho = 0.781, p = 0.009), CCL3 (rho = 0.712, p = 0.041), and CXCL10 (rho = 0.707, p = 0.045) (Fig. 3C). Especially the Vδ2$^+$ subset of γδ T cells has been shown to be important in the immune response to malaria parasites *(21–23)*. When investigating these subsets, we observed that the number of Vδ2$^+$ γδ T cells was positively correlated with levels of IL6 (rho = 0.767, p = 0.013), IFN-gamma (rho = 0.753, p = 0.018), CXCL11 (rho = 0.74, p = 0.024), IL10 (rho = 0.723, p = 0.034), TNF (rho = 0.724, p = 0.034), and CXCL9 (rho = 0.707, p = 0.045) (Fig. 3D). The number of Vδ2$^-$ γδ T cells in contrast was only positively correlated with levels of IL8 (rho = 0.77, p = 0.013), and IL6 (rho = 0.72, p = 0.033) (Fig. 3E).

Acute phase levels of IL8 were also positively correlated with the number of activated
(CD38+HLA-DR+) CD4+ T cells (rho = 0.728, p = 0.031) at the 10-day sample time-point. In
contrast, neurotrophin-3 (NT-3) levels were negatively correlated with the number of activated
CD8+ T cells in several different subsets (CD57+CD38+; rho = –0.77, p = 0.012 and
CD57+CD38+HLA-DR+; rho = –0.726, p = 0.042) (Fig. 3F, G).

These results point towards a strong association of the levels of pro-inflammatory cytokines during
the early acute phase and the size of immune cell subsets after the acute phase. The highest degree
of association was identified for γδ T cells, and especially the Vδ2+ subset, which was strongly
positively correlated with the levels of pro-inflammatory cytokines.

Fig. 3. Associations of the acute phase plasma protein response with immune cell subsets. (A)
Heatmap based on overall Spearman correlation (rho) acute phase plasma proteins and post-

*acute phase cell immune cell subsets. Blue and red colors symbolize positive and negative correlations, respectively. **(B)** Chord diagram based on strong (rho > |0.7|) and significant correlations values (after adjusted FDR < 0.05). **(C-G)** Scatter plots to show acute phase plasma protein levels and immune cell subset counts at day 10 post disease for significant correlations of **(C)** Total γδ T cells, **(D)** Vδ2⁻ γδ T cells, **(E)** Vδ2⁺ γδ T cells, **(F)** activated CD4+ T cells and **(G)** activated CD8+ T cells. All stated p-values are FDR corrected for multiple testing.*

Subhead 4: Impact of previous *P. falciparum* exposure on γδ T cell responses

To further characterize the association between acute phase cytokine levels and the γδ T cell response, we sought to examine if previous exposure to malaria parasites impacts this association. Overall, both groups show similar trajectories of their immune dynamics when plotting Factor1 values against time (Fig. 4A). However, primary infected individuals were associated with both higher and lower factor values during the acute phase and convalescent phase, respectively (Fig. S2C). This indicates differences in protein and immune cell subset magnitudes attributed to previous malaria.

Due to the strong correlation of acute phase proteins and γδ T cell subset numbers at day 10 (Fig. 4B, excerpt of Fig. 3B), we aimed to investigate the association further in the context of previous exposure.

When plotting γδ T cells against the time after symptom onset, we observed that Vδ2⁺ γδ T cells expanded to a greater extent in primary infected individuals while Vδ2⁻ γδ T cells expanded similarly for both groups (Fig. 4C). We confirmed that especially the Vδ2⁺ subset was significantly more expanded after the acute time-point in primary infected individuals (Fig. 4D). All acute phase proteins that correlated with Vδ2⁺ γδ T cells (Fig. 4B) followed a similar trend, although only IFN-gamma, CXCL9 and CXCL11 were significantly higher in primary infected compared to previously exposed individuals (Fig. 4E).

Here, we could show that primary infected individuals have significantly higher levels of pro-inflammatory cytokines at the acute phase, as well as a significantly larger expansion of Vδ2⁺ γδ T cells after the acute phase compared with individuals previously exposed to malaria.

Fig. 4. ***Impact of previous P. falciparum exposure on immune dynamics after clinical malaria.***
(A) Smooth fit of MEFISTO Factor1 values against time after symptom onset in weeks visualizing
transition from acute to convalescence for primary infected (purple) and previously exposed
(orange) individuals. (B) Associated inflammatory signature with γδ T cells based on spearman
rank correlation. (C) γδ T cell dynamics over time after symptom onset. (D) Longitudinal γδ T cell
levels. Statistical differences between groups for γδ T cell subtypes were assessed using linear
mixed model fit with restricted maximum likelihood and t-tests with FDR correction for multiple
testing. Error bars denote standard error of the mean. (E) Comparison between primary infected
(purple) and previously exposed (orange) individuals for acute phase proteins significantly
associated with γδ T cells at the acute time-point. Statistical differences between groups were
assessed using the non-parametric Wilcoxon-test, FDR adjusted p-values to correct for multiple
testing. *p < 0.05, **p < 0.01, ***p < 0.001.

Subhead 5: Effect of previous malaria exposure on Vδ2⁺ γδ T cell characteristics

The Vδ2$^+$ subset of γδ T cells have been shown to play an important role in combating blood stage

malaria, via TCR-induced cytotoxicity and CD16 mediated phagocytosis *(24, 25)*. We therefore

focused on characterizing the CD16$^+$Vδ2$^+$ γδ T cell response further using linear-mixed effects

models to assess the impact of previous exposure to parasites on γδ T cell functional markers.

First, we assessed CD16 expression among γδ T cells. The expression was similar between primary

infected and previously exposed individuals for total and Vδ2 positive and negative subsets (Fig.

5A). The frequency of CD16$^+$ cells increased somewhat in both groups between the acute time-

point and 10 days, for Vδ2$^-$ cells. A similar expansion was also observed for the Vδ2$^+$ cells, but

only in primary infected individuals (Fig. 5A), potentially reflecting an upregulation in response

to infection. To further assess if the Vδ2$^+$ γδ T cells responded to the infection, we next investigated

the activation status of the CD16$^+$Vδ2$^+$ γδ T cells. Approximately 60 % of all Vδ2$^+$ γδ T cells had

upregulated the activation marker CD38 at the acute infection. The primary infected individuals

then retained a significantly higher frequency of CD38$^+$ cells at the day 10 time-point after which

the levels reduced over 3-6 months until reaching baseline levels (Fig. 5B). The increased

activation-status in primary infected individuals was also reflected by a significantly higher co-

expression of CD38 and HLA-DR at the acute time-point (Fig. 5B).

We also assessed the expression of CD56 and CD57, associated with NK cell cytotoxicity *(26)*

and replicative senescence *(27)*, respectively. Although there were no differences between the

groups, there were some changes in cells expressing the markers over time, such as a temporary

reduction in the frequency of CD56$^+$ cells after the acute infection, while CD57$^+$ cells increased

over time (Fig. 5C). These effects were especially observed in primary infected individuals,

potentially reflecting the stronger γδ T cell response in these individuals.

To further determine which subsets of Vδ2$^+$ γδ T cells that responded during infection, we assessed

the levels of CCR7 and CD45RA to identify different naïve or effector populations *(28)*. Almost

all CD16$^+$ Vδ2$^+$ γδ T cells were negative for CCR7 (median 98 % over all time-points), indicating

that these cells display an effector-phenotype. During the acute phase of the response, both groups

displayed similar distribution of effector memory (T$_{EM}$, CD45RA$^-$) and T$_{EMRA}$ (CD45RA$^+$)

γδ T cells (Fig. 5D). However, as the number of Vδ2$^+$ γδ T cells expanded in primary infected

individuals the frequency of effector memory cells significantly increased, while the frequency in

previously exposed individuals was relatively stable over time. This suggests that it was primarily effector memory Vδ2$^+$ γδ T cells that expanded after the acute infection.

Collectively these results show that despite Vδ2$^+$ γδ T cells only expanding in primary infected individuals, the cells became activated in both groups after infection. The level of activation and subsequent changes in host effector and differentiation markers were however different with a more robust effect in primary infected individuals.

Fig. 5. Previous P. falciparum exposure impacts γδ T cell expansion. *Linear mixed effect modelling by maximum likelihood for γδ T cell frequencies for primary infected (purple) and previously exposed (orange) individuals, error bars denote the standard error of the mean. **(A)** Percentage of CD16$^+$ cells of total γδ T cells, Vδ2$^-$ or Vδ2$^+$ γδ T cells. **(B)** Percent of Vδ2$^+$ CD16$^+$γδ T cells expressing activation markers CD38 alone or together with HLA-DR. **(C)** Percent of Vδ2$^+$ CD16$^+$γδ T cells expressing CD56$^+$ or CD57$^+$ **(D)** Vδ2$^+$ γδ T cells effector memory (CD45RA$^-$) or T$_{EMRA}$ (CD45RA$^+$). Difference in cell frequency at each time point was evaluated by repeated t-tests with FDR correction. *$p < 0.05$, ****$p < 0.0001$.*

Subhead 6: γδ T cell function is retained while expansion is affected due to previous malaria exposure

It has been described that especially Vδ2$^+$ γδ T cells respond strongly during malaria but that this effect is reduced upon subsequent infections *(29)*. This pattern is supported by our data, as the Vδ2$^+$ γδ T cell activation and especially expansion were reduced in previously exposed compared

to primary infected individuals (Fig. 4D). Previous descriptions indicate that the reduced

responsiveness can be due to changes in the imprinting through changes in the methylation patterns

(30, 31), although it remains unclear if such imprinting can last for this long. To assess if the

function of the γδ T cells in our cohorts were inherently affected in their capacity to respond to

stimulation, we performed blinded cultures of PBMCs from primary infected and previously

exposed individuals. We stimulated the cells with PMA and ionomycin that together bypass the T

cell receptor complex and measured production of TNFα, IL17, and IL10. PMA and ionomycin

stimulation led to increased production of TNFα, but not IL17 or IL10 (Fig. 6A). Comparing the

response of Vδ2$^+$ and Vδ2$^-$ γδ T cells, the Vδ2$^+$ cells responded with more TNFα production (Fig.

6B), consistent with a proinflammatory effector function. For Vδ2$^-$ γδ T cells, the production of

TNFα was similar between the acute and convalescent time-point and healthy controls. For Vδ2$^+$

γδ T cells, however, the frequency of TNFα producing cells was reduced during the acute response,

but then recovered at the convalescent phase, for which frequencies were similar as in healthy

controls (Fig. 6C). Further separating the Vδ2$^+$ and Vδ2$^-$ subsets into primary infected and

previously exposed individuals showed no differences between the groups for Vδ2$^-$ cells, while

primary infected individuals responded with lower numbers of TNFα-producing Vδ2$^+$ cells at the

acute time-point (Fig. 6D), potentially reflecting a larger proportion of already activated cells being

restimulated.

In summary, Vδ2$^+$ γδ T cells from previously exposed individuals do not display apparent intrinsic

dysfunctionality upon *ex vivo* reactivation. This further supports a role for extrinsic factors

regulating Vδ2$^+$ γδ T cell activation and expansion in malaria.

Fig. 6. γδ T cell function in response to restimulation. *(A) PBMCs from donors with primary infection (n = 4) and previous parasite exposure (n = 4) at the acute and convalescent time-point (6-12 months after infection) and healthy controls (n = 6) were stimulated with PMA and ionomycin (open boxes) or left unstimulated (Mock, open circles) for 5 hours. Frequencies of TNFα, IL17A, and IL10-producing cells were then measured. (B) TNF⁺ cells were compared between Vδ2⁺ (open boxes) and Vδ2⁻ (open circles) cell subsets. (C) Comparison of TNF⁺ frequencies in Vδ2⁻ and Vδ2⁺ cell subsets at the acute time-point (pink circles), convalescent time-point (green boxes) and healthy controls (grey triangles). (D) Comparison of TNF⁺ Vδ2⁻ and Vδ2⁺ cells between primary infected (purple circles) and previously exposed (orange boxes) individuals. Statistical comparisons for A-C were done using a matched pair two-way ANOVA followed by Tukey's post-hoc test, while statistics in D was evaluated by two-way ANOVA followed by Sidak's post-hoc test. *p < 0.05, **p < 0.01, ****p < 0.0001, ns = not significant.*

Subhead 7: Acute malaria specific IgG3 levels are associated with the level of inflammation and γδ T cell expansion

A long-lived memory imprint to pathogenic encounter is in general mediated by the adaptive immune system. Here, we report that previously exposed individuals without re-exposure to parasites for many years (average 11.5 years since leaving an endemic region), at the time of a new acute infection respond with a reduced pro-inflammatory cytokine response upon re-infection as well as reduced Vδ2⁺ γδ T cell expansion. Hence, we hypothesized that the reduced pro-

inflammatory response and γδ T cell expansion could be mediated by existing long-lived adaptive
humoral immune memory responses.

To investigate this, we repurposed a previously published dataset, investigating the
immunoglobulin G subclass response (IgG1-4) to five malaria *P. falciparum* blood-stage antigens
(AMA1, MSP1, MSP2, MSP3, and RH5) in 51 matching individuals from the same cohort *(15)*.
We established an IgG subclass and time point specific score, the cumulative response score (CRS)
that summarizes the responses to the separately measured malaria specific antigens from the
previous study (Fig. S5). The CRS dynamics correspond to the average breadth of the antibody
response for total IgG and IgG subclasses (Fig. 7A).

Spearman rank correlation showed that levels of several acute pro-inflammatory cytokines were
strongly inversely correlated with IgG subclass CRS (Fig.7B), in particularly IgG3 and IFN-
gamma levels (Fig. 7C). The IgG2 and IgG3 subclass CRSs at the acute infection were also
negatively correlated with day 10 total and $V\delta2^+$ γδ T cell numbers, respectively (Fig. 7D), with
an especially strong negative correlation between IgG3 levels and $V\delta2^+$ γδ T cell numbers (Fig.
7E).
Cytophilic immunoglobulins such as IgG3 can interact with CD16 expressing immune cells,
including γδ T cells, and intermediate ($CD16^+CD14^{high}$) and non-classical ($CD16^+CD14^{lo}$)
monocytes. Interestingly, we observed higher numbers of intermediate and non-classical
monocytes at the acute time point for previously exposed compared to primary infected individuals
(Fig. 7F). These observations, together with an increased IgG3 response, points towards an
antibody-phagocytosis mediated mechanism that could partly explain the lower parasitemia
observed in individuals with prior exposure to malaria parasites (Fig. 1E). To assess if this could
be the case, we used linear regression to determine which associated combinations of immune
factors that best explain the difference in expansion of $V\delta2^+$ γδ T cells for the two exposure groups.
Using Directed Acyclic Graphs (DAG) *(32)*, we determined the minimal sufficient adjustment sets
of covariates for estimating the total effect of a given associated component on acute IFN-gamma
levels and day 10 $V\delta2^+$ γδ T cell numbers, respectively (described in Supplementary Methods and
Fig. S7). Comparing the reference model with exposure group as explanatory variable to the DAG-
based adjusted immune response component model, we concluded that the IFN-gamma response
was not directly explained by the parasitemia (Fig. S7A). In contrast, differences in $V\delta2^+$ γδ T cell

expansion was partly explained by parasitemia, but more directly by cytophilic antibody levels and CD16[+] intermediate monocytes (Fig. S7B-C). This suggests a potentially more direct antibody-mediated effect by regulating cytokine levels. Based on these results we suggest a working model where early production of parasite-specific antibodies interact with innate immune cells to regulate parasitemia and cytokine levels, which in turn control subsequent γδ T cell expansion (Figure 7G).

Fig. 7*. *Adaptive response due to prior malaria exposure linked to dampened immune inflammation. (A) Cumulative IgG subclass response score (CRS) to 5 blood-stage antigens from Yman et al., BMC Medicine, 2019, over sample time points for primary infected (purple) and

previously exposed (orange) individuals (n = 52), shaded area denotes the 95% confidence

*interval. **(B)** Correlation network of plasma proteins and IgG subclass CRS values at the acute*

*time-point, FDR corrected p < 0.05 **(C)** Scatter plots for significant correlations, FDR adjusted p-*

*values are stated. **(D)** Correlation network of acute IgG subclass CRS values and immune cell*

*subsets at the 10-day sample time point, FDR corrected p < 0.05, rho < –0.7. **(E)** Scatter plots for*

*significant correlations, FDR corrected p < 0.05. **(F)** Cell counts of monocyte subsets at the acute*

time point. Statistical differences between groups were assessed using the non-parametric

*Wilcoxon-test, FDR adjusted p-values to correct for multiple testing. *p < 0.05, **p < 0.01 **(G)***

Schematic model to explain improved disease tolerance in previously exposed individuals.

DISCUSSION

Immunity to malaria is slow to develop. It has been proposed that this is partly due to immune perturbations during infection *(41)*. Understanding how the immune system is affected during natural infection with malaria parasites is important to identify the potential role of individual immune mediators in this process. Here, we used systems immunology to comprehensively investigate the immune landscape after natural malaria and over one year after infection in the absence of re-exposure to parasites. By sampling each individual over time, we could reduce potential noise associated with inter-individual variability in their immune response to the infection. Overall, the integrated immune landscape identified the acute phase followed by a transitional phase leading into convalescence up to one year after disease. The acute phase was characterized by activated CD4$^+$ and CD8$^+$ T cells and high levels of inflammatory cytokines and chemokines, consistent with previous literature *(2, 33)*. Here, a comprehensive analysis of immune components over time revealed that acute inflammatory response signature associated with long-term changes to cellular immune response.

The composition of our cohort, consisting of both primary infected and previously exposed individuals, allowed us to study the impact of memory on the acute response and its associations with the transition towards immunological convalescence. We could show that previously exposed individuals produce lower levels of pro-inflammatory cytokines during natural *P. falciparum* infection. Similar findings have been observed at the blood transcriptome level by Tran et al., where Malian adults exhibited a dampened inflammatory response compared to first time CHMI-infected Dutch adults *(34)*. Further, a study from a cohort of children in Uganda showed that older children have lower levels of serum cytokines during acute malaria compared to younger children *(35)*. These previous studies, together with our findings of reduced pro-inflammatory response confirms that previous malaria exposure, possibly cumulative, induces a form of tolerance response. Of note, all individuals in our cohort experienced clinical malaria, suggesting that the level of tolerance could be dependent on the amount of prior exposure to the parasite. However, the individuals with previous exposure had left malaria endemic areas on average 11.5 years earlier before being reinfected/experiencing a new episode of acute *P. falciparum* malaria after visiting endemic area, indicating that the mechanism is significantly long-lived.

Antibodies are highly associated with protection from malaria and a key building block of naturally acquired immunity. We have recently reported that previously exposed individuals from our cohort responded to infection by producing high levels of *P. falciparum*-specific IgG, particularly of the cytophilic subclasses IgG1 and IgG3 *(15)*. Memory B cells and long-lived plasma cells constitute a likely source of the increased parasite specific antibodies *(15, 36)*, as memory can be maintained for up to 16 years without re-exposure in a different cohort of travelers *(37)*. Re-purposing the previously published antibody data *(15)* enabled us to link the effect of adaptive responses, in the form of an increased cytophilic antibodies, to enhanced functionality of the innate response. As reported here, previously exposed individuals responded to malaria with an overall dampened inflammatory response but significantly higher numbers of intermediate monocytes *(38)*. Their expression of CD16 (FcγRIII), a receptor for cytophilic antibodies of the subclasses IgG1 and IgG3, can mediate parasite clearance through antibody-dependent phagocytosis (ADP) and/or antibody-dependent cellular cytotoxicity (ADCC) *(39)*. Indeed, higher numbers of CD16$^+$ monocytes and a stronger IgG3 response against blood stage antigens indicate an interplay which could promote lower levels of parasitemia, compared to individuals without previous exposure. Supporting such a mechanisms, vaccination with RTS,S/AS01 leads to CD16-mediated phagocytosis associated with protection *(40)*.

We propose that antibody-mediated parasite control is an important component in shaping the pro-inflammatory cytokine milieu towards a reduced pro-inflammatory response which could be important for the development of immune tolerance *(41)*. Complementary to the described adaptive-innate interplay, several recent studies suggest that monocytes are modulated towards a tolerance phenotype *(5, 41)*. The mechanism of this imprint remains unclear, but a rodent model point towards transcriptomic reprogramming of monocytes in the spleen *(5)*, while data from *in-vitro* stimulated monocytes from a cohort in Mali point towards epigenetic reprogramming of myeloid progenitor cells in the bone marrow *(41)*. However, to our knowledge, no study has shown that epigenetic remodeling can remain and affect innate cells at re-stimulation for that long. An alternative hypothesis, that overcomes the fact that monocytes have a short lifespan *(42)*, is that they are remodeled in their function due to constant stimuli by remaining hemozoin in the spleen *(5, 43, 44)*.

Both hypotheses of epigenetic imprinting and through splenic remodeling need further

investigation. In our cohort, where study participants moved away from malaria endemic areas 10

years prior, we propose an additional mechanism responsible for the tolerogenic response via the

interplay of innate and adaptive immunity. We suggest that tolerance develops over several

exposures to parasites and further that it is sustained long-term through adaptive memory

responses. However, these above hypotheses are not mutually exclusive and could potentially

synergize.

In agreement with previous reports, we observed expanding γδ T cells after the initial inflammatory

response in response to the infection *(45, 46)*. γδ T cells are suggested to have an important role in

the control of malaria *(29)* and direct anti-parasite functions during the blood stage have been

reported *(24)*. In this study, we observed that the acute phase inflammatory response was positively

associated with expansion of Vδ2$^+$ γδ T cells, and further that this was strongly impacted by

previous malaria exposure, resulting in a dampened inflammatory response and less γδ T cell

expansion. Interestingly, this effect was primarily observed for the subset of Vδ2$^+$ and not Vδ2$^-$

γδ T cells. The strong association could indicate that the pro-inflammatory response directly or

indirectly shapes the subsequent γδ T cell expansion and capacity to adapt to prolonged and high

levels of blood stage parasitemia.

Vδ2$^+$ γδ T cells are known to activate and expand during a primary *P. falciparum* infection in

response to malaria phosphoantigens and that their activity is modulated upon subsequent

infections *(47)*. Given the recently described role of Vδ2$^+$ γδ T cells in antiparasitic activities via

antibody-CD16 dependent phagocytosis of infected erythrocytes and cytotoxicity *(24, 25)*,

associations to possible activity modulating factors are of interest. The benefit of reduced

expansion of γδ T cells due to a dampened inflammatory response could be to reduce overall

inflammatory responses, which are otherwise potentially detrimental to the host *(48)*. This is a

common hypothesis as the γδ T cells are known to acquire a dysfunctional or tolerance phenotype

over time with repeated episodes of malaria *(23)*. This reduction in γδ T cell effector function was

associated with continuous malaria exposure *(23)*, although it remains unclear what the

underlaying tolerance mechanism is, and how long-lived such an effect could be. To assess if the

reduced activation and expansion of Vδ2$^+$ γδ T cells could be due to an inherent inability of the γδ

T cells to respond, we re-stimulated cells from primary infected and previously exposed

individuals. However, we did not observe any reduced response that could have been associated

with intrinsic down-regulation of effector functions. Based on this, we instead hypothesized that

the regulation of Vδ2$^+$ γδ T cell activation and expansion was likely due to extrinsic mechanisms.

Among the cytokines and chemokines associated with Vδ2$^+$ γδ T cell expansion, we found IFN-

gamma and CXCL11 significantly reduced in previously exposed individuals. CXCL11 is

chemotactic for activated T cells, and it was reported that individuals with asymptomatic

P. falciparum malaria had lower levels in an endemic setting *(49)*. IFN-gamma and the IFN-

gamma inducible chemokine CXCL9 are associated in a different context with regulatory crosstalk

of pro-inflammatory γδ T cell effects *(50)*, which could explain the association of these cytokines

with subsequent expansion. Consistent with these observations, both IFN-gamma and CXCL11

were reduced due to previous parasite exposure in a mouse model, suggesting some type of

memory response *(5)*.

In summary, our results, together with previous research in the field, supports a model where early

control of host self-damage through increased tolerance is needed until a more broad, diverse, and

protective antibody repertoire is achieved (see Figure 7G). The tolerance response (dampened pro-

inflammatory response and reduced Vδ2$^+$ γδ T cell expansion) could be mediated in two ways: 1)

an expanding antibody repertoire that enhances CD16$^+$ mediated phagocytosis and effector

functions, direct parasite neutralization and blocking further RBC invasion, rosetting and

sequestration by merozoites, and 2) training/priming during previous malaria episodes via

epigenetic or transcriptional remodeling of the monocyte population, potentially leading to rapid

generation of CD16-expressing monocytes with enhanced antibody effector functions.

From an evolutionary perspective, this type of dampened inflammatory response could be

important to generate an efficient B cell response, as recently shown in an influenza vaccine study

(51) and further supported by mouse models of malaria *(52)*. Although further studies are needed

to evaluate these mechanisms on a cellular and molecular level, the comprehensive systems levels

analysis performed here provides a dynamic description for how proteins, cells, and antibodies,

interact in the host response to malaria during and after primary and repeated infection.

MATERIALS AND METHODS

Study design/cohort

A prospective study enrolling adult patients diagnosed with *P. falciparum* malaria at Karolinska University Hospital in Stockholm, Sweden. Fifty-three patients that were admitted with acute *P. falciparum* infection between 2011 and 2017, and where both cells and plasma was frozen were included in the current study. We stratified the patients based on previous exposure to compare the immune responses in malaria-naive individuals of primarily Swedish origin, who contracted malaria for the first time (n = 17, denoted as primary infected), with individuals originating from malaria-endemic areas in Sub-Saharan Africa and reporting previous malaria (n = 36, denoted as previously exposed). Patients were invited for sampling at the time of malaria diagnosis (Acute) and then at approximately 10 days (D10), 1 month (M1), 3 months (M3), 6 months (M6), and 12 months (Y1) (see Fig.1A). On each sampling, venous blood was collected for plasma and serum isolation, PBMC preparation and blood chemistry. Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll-Paque density gradient separation, resuspended in 90 % fetal calf serum supplemented with 10% DMSO, and stored at - 150°C. Clinical data was extracted from medical records, and a questionnaire relating to the participant's health status, previous traveling, and malaria exposure was filled in by all participants (Table S1).

Part of the cohort has been described previously for clinical *(53)*, parasitological *(13)* and immunological aspects *(15, 16, 54)*.

In addition, we profiled peripheral blood samples of eight healthy controls to compare relative measures of protein expression and cell populations with healthy/normal values (Figure S2).

Clinical diagnostics

P. falciparum parasites were detected and enumerated by light microscopy of Fields stained thick and thin blood smears at the Department of Clinical Microbiology at Karolinska University Hospital in addition to PCR as described previously *(13)*. Leukocyte, neutrophil, monocyte, and platelet cell differential counts were performed at the Department of Clinical Chemistry at Karolinska University Hospital.

Data acquisition

Immune cell phenotyping

Immune cells were phenotyped by Flow cytometry (LSRFortessa, BD) and a panel of 17 antibodies covering major immune cell populations and subpopulations (Fig. S1) for 137 samples. Frozen PBMCs were thawed in a 37°C water bath and mixed with 1 equal volume cold Iscove's Modified Dulbecco's Medium supplemented with L-glutamine (2 mM), penicillin (100 U/ml), streptomycin (100 µg/ml), and 10 % heat-inactivated fetal bovine serum (all from Thermo Fischer Scientific). Cells were then rested 20 minutes on ice before being washed twice in DPBS lacking magnesium or calcium. After washing, the cells were incubated with Aqua Live/Dead stain (Thermo Fisher Scientific) for 20 minutes followed by further washing in DPBS supplemented with 2% FBS. The cells were then incubated for 20 minutes on ice, in 2 steps with 2 washes in between, using an antibody mix targeting surface antigens (Table S2). After staining, the cells were washed twice in DPBS with 2% FBS before acquisition on a 5-laser BD LSRFortessa flow cytometer. Gating was done with FlowJo X software version 10.4.2, with the gating strategy shown in Figure S1.

Plasma protein profiling

Plasma proteins relevant for the immune response were profiled using the Proximity Extension Assay (PEA) (Olink Proteomics, Sweden) to analyze 171 samples of 53 individuals using Olink Target 96 Inflammation panel (92-plex). The method has been described previously *(17)*. Briefly, paired oligonucleotide-coupled antibodies bind to target proteins, leading to hybridization of the oligonucleotides when the antibody-pair is in close proximity, forming a PCR template for real-time PCR detection. Resulting data is normalized using assay internal controls and transformed into Normalized Protein eXpression (NPX) values, representing an arbitrary relative quantification unit on log2 scale. After quality control and removal of markers with a missing frequency greater than 50%, 74 proteins remained for downstream analysis.

***In vitro* stimulation for detection of γδ T cells cytokine production**

For intracellular staining, PBMCs were thawed, washed, resuspended in complete RMPI media supplemented with 10% FCS and then incubated at 37°C over-night. The cells were then stimulated with PMA (0.4 µM) and ionomycin (13 µM) per 10^6 cells, (Nordic Biosite AB) for 5 hours. Golgi plug (BD Biosciences) was added after the first hour of stimulation. Activated cells were subsequently stained with antibodies targeting the surface markers CD3, γδ TCR, and Vδ2,

according to manufacturers' instruction. Cells were then fixed and permeabilized using the BD

Cytofix/Cytoperm reagents (BD Biosciences) and stained with intracellular antibodies for TNF,

IL17A, and IL10 (Table S2). For the exclusion of dead cells, LIVE/DEAD aqua fixable viability

staining kit (Invitrogen) was used. Cells were acquired using a BD LSR Fortessa BD (BD

Biosciences) and the data analyzed using FlowJo Version 7.6 (FlowJo, Ashland, OH).

Bioinformatics

Immune cell data normalization

Prior to bioinformatic analysis, cell proportions from manually gates (Fig. S1) were normalized to

the number of cells per 1000 live cells. Subsequently, we adjusted the experimentally obtained live

cell counts with lymphocyte and monocyte counts from clinical blood chemistry counts for each

individual and timepoint to obtain cells per microliter blood values (Fig. S5A). Monocyte gates

containing values (Fig. S1A) were normalized according to:

$$\frac{\text{cells}}{1000 \text{ live cells}} * (\text{lymphocyte counts} + \text{monocyte counts}) = \frac{\text{cells}}{\text{microliter blood}}$$

Values of pure lymphocyte containing gates (Fig. S1B) were normalized according to:

$$\frac{\text{cells}}{1000 \text{ live cells}} * \text{lymphocyte counts} = \frac{\text{cells}}{\text{microliter blood}}$$

To standardize the cells-per-microliter values and to normalize the data set for extreme values, we

performed log2 transformation. Standardized data was comparable to not-standardized data (Fig.

S5B).

Antibody cumulative response score (CRS) calculation

A previous study on individuals of the traveler cohort investigated the immunoglobulin G subclass

response (IgG1-4) s to five malaria *P. falciparum*-blood-stage antigens (AMA1, MSP1, MSP2,

MSP3, RH5) in 52 this current study matching cohort individuals *(15)*. For the current study, we

though to look at the overall IgG subclass response to merozoite antigens instead of specific

antigens. To summarize the adjusted signals, measured for each antigen and each subclass, we

rank normalized the signals to achieve normal distribution for each antigen (Fig. S5A). Normally

distributed antigen response values were then summarized for each IgG subclass and time point,

creating the cumulative response score (CRS), presenting an average breadth value of the antibody

response for IgG subclasses (Fig. S5B).

Statistical analysis and visualization

Multi-Omics-Factor Analysis

Multi-Omics Factor analysis (MOFA) is a powerful tool for omics integration. It reduces high-dimensional data into a few latent factors by capturing multi-dataset spanning variance in these factors. The unsupervised nature of MOFA+ allows the model to capture both biological and technical variability in the low-dimensional factors space *(18, 19)*. A recent development (MEFISTO) in addition allows for multi-omics integration of data with continues structures given by temporal relationships*(20)*. Here, we utilized MEFISTO to analyze the underlying key drivers for the time course of one year after infection on our cohort of returning travelers with *P. falciparum* malaria.

For the data-driven-integration approach, we used 15,396 immune feature data points out of 23,130 possible. We included manually gated immune cell subset (features = 44) data and highly multiplexed plasma protein data (features = 60) and including the time after symptom onset (0-70 weeks) for each sample to disentangle time dependent variation on a systems-level.

To standardize the actual time covariate of each sample, we used the time after reported symptom onset for each sample as a temporal covariate for the model. Moreover, we assigned the time-associated samples to groups of primary infected (n = 57) and previously exposed (n = 125) to account for previous *P. falciparum* exposure. Immune parameters without temporal variation were excluded prior to model training.

We trained the MEFISTO model using default model options, but adjusted training options (drop-factor-threshold = 0.05; maxiter = 10,000; convergence_mode = slow). To align the covariates across groups, warping was set "TRUE" and "primary infected" was set as warping reference group. The optimal model was determined by the MOFA+ function run_mofa() with the setting "use_basilisk = T".

All data wrangling, analysis and visualization was done using R (www.r-project.org) using the tidyverse package *(55)*. Spearman correlation analysis with FDR/BH correction for multiple testing *(56)* was done using the correlation package *(57)*. If not stated otherwise, non-parametric data distribution was assumed, and statistical difference was assessed using unpaired Wilcox Test from rstatix (https://github.com/kassambara/rstatix).

Results were visualized using r packages circlize *(58)*, complexHeatmap *(59)* and ggpubr

(https://github.com/kassambara/ggpubr9). Linear-mixed-effect models, with subjectID as random

effect, were fitted in GraphPad prism version 9.1.2 using restricted maximum likelihood followed

by t-tests using predicted least squares means with Benjamini and Hochberg FDR correction for

multiple testing *(56)*.

Supplementary Materials

Table S1. Descriptive statistics prospective cohort of returning travelers.

Table S2. Staining panel for flow cytometry

Fig. S1. Flow cytometry manual gating strategy – T cell and B cell panel.

Fig. S2. MEFISTO model and cohort internal feature evaluation.

Fig. S3. Cohort external evaluation - comparison of plasma protein and PBMC profiles at Acute

and Y1 compared to healthy controls.

Fig. S4. Comparison of MEFISTO Factor1 values for exposure groups on time points.

Fig. S5. Cumulative Response Score for malaria antigen specific IgG subclass response of Yman

et al 2019 data set.

Fig. S6. Leukocyte count adjusted cell counts.

Fig. S7. Directed Acyclic Graphs – DAGs and linear regression

References and Notes

1. World Health Organisation, *World malaria report 2020* (Geneva, 2020;

https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2020).

2. J. Langhorne, F. M. Ndungu, A. M. Sponaas, K. Marsh, Immunity to malaria: More questions

than answers*Nature Immunology* **9**, 725–732 (2008).

3. F. H. A. Osier, G. Fegan, S. D. Polley, L. Murungi, F. Verra, K. K. A. Tetteh, B. Lowe, T.

Mwangi, P. C. Bull, A. W. Thomas, D. R. Cavanagh, J. S. McBride, D. E. Lanar, M. J.

Mackinnon, D. J. Conway, K. Marsh, Breadth and magnitude of antibody responses to multiple

Plasmodium falciparum merozoite antigens are associated with protection from clinical malaria, *Infection and Immunity* (2008), doi:10.1128/IAI.01585-07.

4. B. P. Gonçalves, C.-Y. Huang, R. Morrison, S. Holte, E. Kabyemela, D. R. Prevots, M. Fried, P. E. Duffy, Parasite Burden and Severity of Malaria in Tanzanian Children, *New England Journal of Medicine* **370**, 1799–1808 (2014).

5. W. Nahrendorf, A. Ivens, P. J. Spence, Inducible mechanisms of disease tolerance provide an alternative strategy of acquired immunity to malaria, *eLife* **10** (2021), doi:10.7554/eLife.63838.

6. A. F. Cowman, J. Healer, D. Marapana, K. Marsh, Malaria: Biology and Disease*Cell* **167**, 610–624 (2016).

7. P. D. Crompton, J. Moebius, S. Portugal, M. Waisberg, G. Hart, L. S. Garver, L. H. Miller, C. Barillas-Mury, S. K. Pierce, Malaria Immunity in Man and Mosquito: Insights into Unsolved Mysteries of a Deadly Infectious Disease, *Annual Review of Immunology* **32**, 157–187 (2014).

8. C. Loiseau, M. M. Cooper, D. L. Doolan, Deciphering host immunity to malaria using systems immunology, *Immunological Reviews* **293**, 115–143 (2020).

9. T. M. Tran, P. D. Crompton, Decoding the complexities of human malaria through systems immunology*Immunological Reviews* **293**, 144–162 (2020).

10. M. M. Davis, C. M. Tato, D. Furman, Systems immunology: Just getting started*Nature Immunology* **18**, 725–732 (2017).

11. T. M. Tran, R. Guha, S. Portugal, J. Skinner, A. Ongoiba, J. Bhardwaj, M. Jones, J. Moebius, P. Venepally, S. Doumbo, E. A. DeRiso, S. Li, K. Vijayan, S. L. Anzick, G. T. Hart, E. M. O'Connell, O. K. Doumbo, A. Kaushansky, G. Alter, P. L. Felgner, H. Lorenzi, K. Kayentao, B. Traore, E. F. Kirkness, P. D. Crompton, A Molecular Signature in Blood Reveals a Role for p53 in Regulating Malaria-Induced Inflammation, *Immunity* (2019), doi:10.1016/J.IMMUNI.2019.08.009.

12. S. E. Jong, V. Unen, M. D. Manurung, K. A. Stam, J. J. Goeman, S. P. Jochems, T. Hamp, N. Pezzotti, Y. D. Mouwenda, M. Eunice Betouke Ongwe, F.-R. Lorenz, Y. C. M Kruize, S. Azimi, M. H. Kamp, A. Vilanova, E. Eisemann, B. P. F Lelieveldt, M. Roestenberg, B. Kim Lee Sim, M. J. T Reinders, R. Fendel, S. L. Hoffman, P. G. Kremsner, F. Koning, B. Mordmamp, B. Lell, M. Yazdanbakhsh, Systems analysis and controlled malaria infection in Europeans and Africans elucidate naturally acquired immunity, *Nature Immunology* (2021), doi:10.1038/s41590-021-00911-7.

13. M. Vafa Homann, S. N. Emami, V. Yman, C. Stenström, K. Sondén, H. Ramström, M.

Karlsson, M. Asghar, A. Färnert, Detection of Malaria Parasites After Treatment in Travelers: A

12-months Longitudinal Study and Statistical Modelling Analysis, *EBioMedicine* **25**, 66–72

(2017).

14. M. Asghar, V. Yman, M. V. Homann, K. Sondén, U. Hammar, D. Hasselquist, A. Färnert,

Cellular aging dynamics after acute malaria infection: A 12-month longitudinal study, *Aging Cell*

(2018), doi:10.1111/acel.12702.

15. V. Yman, M. T. White, M. Asghar, C. Sundling, K. Sondén, S. J. Draper, F. H. A. A. Osier,

A. Färnert, Antibody responses to merozoite antigens after natural Plasmodium falciparum

infection: Kinetics and longevity in absence of re-exposure, *BMC Medicine* (2019),

doi:10.1186/s12916-019-1255-3.

16. C. Sundling, C. Rönnberg, V. Yman, M. Asghar, P. Jahnmatz, T. Lakshmikanth, Y. Chen, J.

Mikes, M. N. Forsell, K. Sondén, A. Achour, P. Brodin, K. E. M. Persson, A. Färnert, B cell

profiling in malaria reveals expansion and remodeling of CD11c+ B cell subsets, *JCI Insight*

(2019), doi:10.1172/jci.insight.126492.

17. E. Assarsson, M. Lundberg, G. Holmquist, J. Björkesten, S. Bucht Thorsen, D. Ekman, A.

Eriksson, E. Rennel Dickens, S. Ohlsson, G. Edfeldt, A.-C. Andersson, P. Lindstedt, J. Stenvang,

M. Gullberg, S. Fredriksson, J. D. Hoheisel, Ed. Homogenous 96-Plex PEA Immunoassay

Exhibiting High Sensitivity, Specificity, and Excellent Scalability, *PLoS ONE* **9**, e95192 (2014).

18. R. Argelaguet, B. Velten, D. Arnol, S. Dietrich, T. Zenz, J. C. Marioni, F. Buettner, W.

Huber, O. Stegle, Multi-Omics Factor Analysis—a framework for unsupervised integration of

multi-omics data sets, *Molecular Systems Biology* (2018), doi:10.15252/msb.20178124.

19. R. Argelaguet, D. Arnol, D. Bredikhin, Y. Deloro, B. Velten, J. C. Marioni, O. Stegle,

MOFA+: A statistical framework for comprehensive integration of multi-modal single-cell data,

Genome Biology **21**, 111 (2020).

20. B. Velten, J. M. Braunger, D. Arnol, R. Argelaguet, O. Stegle, Identifying temporal and

spatial patterns of variation from multi-modal data using MEFISTO, ,

doi:10.1101/2020.11.03.366674.

21. P. Jagannathan, F. Lutwama, M. J. Boyle, F. Nankya, L. A. Farrington, T. I. McIntyre, K.

Bowen, K. Naluwu, M. Nalubega, K. Musinguzi, E. Sikyomu, R. Budker, A. Katureebe, J. Rek,

B. Greenhouse, G. Dorsey, M. R. Kamya, M. E. Feeney, Vδ2+ T cell response to malaria

correlates with protection from infection but is attenuated with repeated exposure, *Scientific*

Reports **7**, 1–12 (2017).

22. L. A. Farrington, P. C. Callaway, H. M. Vance, K. Baskevitch, E. Lutz, L. Warrier, T. I.

McIntyre, R. Budker, P. Jagannathan, F. Nankya, K. Musinguzi, M. Nalubega, E. Sikyomu, K.

Naluwu, E. Arinaitwe, G. Dorsey, M. R. Kamya, M. E. Feeney, G. Hart, Ed. Opsonized antigen

activates Vδ2+ T cells via CD16/FCγRIIIa in individuals with chronic malaria exposure, *PLOS*

Pathogens **16**, e1008997 (2020).

23. P. Jagannathan, C. C. Kim, B. Greenhouse, F. Nankya, K. Bowen, I. Eccles-James, M. K.

Muhindo, E. Arinaitwe, J. W. Tappero, M. R. Kamya, G. Dorsey, M. E. Feeney, Loss and

dysfunction of Vδ2+γδ T cells are associated with clinical tolerance to malaria, *Science*

Translational Medicine **6**, 251ra117-251ra117 (2014).

24. C. Junqueira, R. B. Polidoro, G. Castro, S. Absalon, Z. Liang, S. sen Santara, Â. Crespo, D.

B. Pereira, R. T. Gazzinelli, J. D. Dvorin, J. Lieberman, γδ T cells suppress Plasmodium

falciparum blood-stage infection by direct killing and phagocytosis, *Nature Immunology* , 1–11

(2021).

25. L. A. Farrington, P. C. Callaway, H. M. Vance, K. Baskevitch, E. Lutz, L. Warrier, T. I.

McIntyre, R. Budker, P. Jagannathan, F. Nankya, K. Musinguzi, M. Nalubega, E. Sikyomu, K.

Naluwu, E. Arinaitwe, G. Dorsey, M. R. Kamya, M. E. Feeney, G. Hart, Ed. Opsonized antigen

activates Vδ2+ T cells via CD16/FCγRIIIa in individuals with chronic malaria exposure, *PLOS*

Pathogens **16**, e1008997 (2020).

26. H. H. van Acker, A. Capsomidis, E. L. Smits, V. F. van Tendeloo, CD56 in the immune

system: More than a marker for cytotoxicity?*Frontiers in Immunology* **8**, 1 (2017).

27. W. Xu, A. Larbi, Heterogeneity in Vδ2pos T cell homeostasis in response to

stress*EBioMedicine* **43**, 31 (2019).

28. V. Appay, R. A. W. van Lier, F. Sallusto, M. Roederer, Phenotype and function of human T

lymphocyte subsets: Consensus and issues*Cytometry Part A* **73**, 975–983 (2008).

29. K. Deroost, J. Langhorne, Gamma/Delta T Cells and Their Role in Protection Against

Malaria, *Frontiers in Immunology* **9**, 2973 (2018).

30. H. Hsu, S. Boudova, G. Mvula, T. H. Divala, R. G. Mungwira, C. Harman, M. K. Laufer, C.

D. Pauza, C. Cairo, Prolonged PD1 Expression on Neonatal Vδ2 Lymphocytes Dampens

Proinflammatory Responses: Role of Epigenetic Regulation, *The Journal of Immunology* **197**, 1884–1892 (2016).

31. K. W. Dantzler, P. Jagannathan, γδ T cells in antimalarial immunity: New insights into their diverse functions in protection and tolerance*Frontiers in Immunology* (2018), doi:10.3389/fimmu.2018.02445.

32. A. Ankan, I. M. N. Wortel, J. Textor, Testing Graphical Causal Models Using the R Package "dagitty," *Current Protocols* **1** (2021), doi:10.1002/cpz1.45.

33. T. W. Ademolue, Y. Aniweh, K. A. Kusi, G. A. Awandare, Patterns of inflammatory responses and parasite tolerance vary with malaria transmission intensity, *Malaria Journal* **16**, 145 (2017).

34. T. M. Tran, M. B. Jones, A. Ongoiba, E. M. Bijker, R. Schats, P. Venepally, J. Skinner, S. Doumbo, E. Quinten, L. G. Visser, E. Whalen, S. Presnell, E. M. O'connell, K. Kayentao, O. K. Doumbo, D. Chaussabel, H. Lorenzi, T. B. Nutman, T. H. M. Ottenhoff, M. C. Haks, B. Traore, E. F. Kirkness, R. W. Sauerwein, P. D. Crompton, Transcriptomic evidence for modulation of host inflammatory responses during febrile Plasmodium falciparum malaria OPEN, (2016), doi:10.1038/srep31291.

35. L. Farrington, H. Vance, J. Rek, M. Prahl, P. Jagannathan, A. Katureebe, E. Arinaitwe, M. R. Kamya, G. Dorsey, M. E. Feeney, Both inflammatory and regulatory cytokine responses to malaria are blunted with increasing age in highly exposed children, *Malaria Journal* **16**, 499 (2017).

36. P. Jahnmatz, C. Sundling, V. Yman, L. Widman, M. Asghar, K. Sondén, C. Stenström, C. Smedman, F. Ndungu, N. Ahlborg, A. Färnert, Memory B-Cell Responses Against Merozoite Antigens After Acute Plasmodium falciparum Malaria, Assessed Over One Year Using a Novel Multiplexed FluoroSpot Assay, *Frontiers in Immunology* **11**, 619398 (2021).

37. F. M. Ndungu, K. Lundblom, J. Rono, J. Illingworth, S. Eriksson, A. Färnert, Long-lived Plasmodium falciparum specific memory B cells in naturally exposed Swedish travelers, *European Journal of Immunology* (2013), doi:10.1002/eji.201343630.

38. A. Ortega-Pajares, S. J. Rogerson, The Rough Guide to Monocytes in Malaria Infection*Frontiers in immunology* **9**, 2888 (2018).

39. S. P. Kurup, N. S. Butler, J. T. Harty, T cell-mediated immunity to malaria*Nature Reviews Immunology* **19**, 457–471 (2019).

40. T. J. Suscovich, J. K. Fallon, J. Das, A. R. Demas, J. Crain, C. H. Linde, A. Michell, H.

Natarajan, C. Arevalo, T. Broge, T. Linnekin, V. Kulkarni, R. Lu, M. D. Slein, C. Luedemann,

M. Marquette, S. March, J. Weiner, S. Gregory, M. Coccia, Y. Flores-Garcia, F. Zavala, M. E.

Ackerman, E. Bergmann-Leitner, J. Hendriks, J. Sadoff, S. Dutta, D. A. Lauffenburger, E.

Jongert, U. Wille-Reece, G. Alter, *Mapping functional humoral correlates of protection against*

malaria challenge following RTS,S/AS01 vaccination (2020).

41. R. Guha, A. Mathioudaki, S. Doumbo, D. Doumtabe, J. Skinner, G. Arora, S. Siddiqui, S. Li,

K. Kayentao, A. Ongoiba, J. Zaugg, B. Traore, P. D. Crompton, J. W. Kazura, Ed. Plasmodium

falciparum malaria drives epigenetic reprogramming of human monocytes toward a regulatory

phenotype, *PLOS Pathogens* **17**, e1009430 (2021).

42. A. A. Patel, Y. Zhang, J. N. Fullerton, L. Boelen, A. Rongvaux, A. A. Maini, V. Bigley, R.

A. Flavell, D. W. Gilroy, B. Asquith, D. Macallan, S. Yona, The fate and lifespan of human

monocyte subsets in steady state and systemic inflammation, *Journal of Experimental Medicine*

214, 1913–1923 (2017).

43. P. A. Buffet, I. Safeukui, G. Deplaine, V. Brousse, V. Prendki, M. Thellier, G. D. Turner, O.

Mercereau-Puijalon, The pathogenesis of Plasmodium falciparum malaria in humans: Insights

from splenic physiology*Blood* **117**, 381–392 (2011).

44. S. Kho, L. Qotrunnada, L. Leonardo, B. Andries, P. A. I. Wardani, A. Fricot, B. Henry, D.

Hardy, N. I. Margyaningsih, D. Apriyanti, A. M. Puspitasari, P. Prayoga, L. Trianty, E.

Kenangalem, F. Chretien, I. Safeukui, H. A. del Portillo, C. Fernandez-Becerra, E. Meibalan, M.

Marti, R. N. Price, T. Woodberry, P. A. Ndour, B. M. Russell, T. W. Yeo, G. Minigo, R.

Noviyanti, J. R. Poespoprodjo, N. C. Siregar, P. A. Buffet, N. M. Anstey, Hidden Biomass of

Intact Malaria Parasites in the Human Spleen, *New England Journal of Medicine* **384**, 2067–

2069 (2021).

45. M. R. Mamedov, A. Scholzen, R. v. Nair, K. Cumnock, J. A. Kenkel, J. H. M. Oliveira, D. L.

Trujillo, N. Saligrama, Y. Zhang, F. Rubelt, D. S. Schneider, Y. hsiu Chien, R. W. Sauerwein,

M. M. Davis, A Macrophage Colony-Stimulating-Factor-Producing γδ T Cell Subset Prevents

Malarial Parasitemic Recurrence, *Immunity* **48**, 350-363.e7 (2018).

46. A. Trampuz, M. Jereb, I. Muzlovic, R. M. Prabhu, Clinical review: Severe malaria*Critical*

Care **7**, 315–323 (2003).

47. J. Howard, I. Zaidi, S. Loizon, O. Mercereau-Puijalon, J. Déchanet-Merville, M. Mamani-

Matsuda, Human Vγ9Vδ2 T Lymphocytes in the Immune Response to P. falciparum Infection,

Frontiers in Immunology **9**, 2760 (2018).

48. D. I. Stanisic, J. Cutts, E. Eriksson, F. J. I. Fowkes, A. Rosanas-Urgell, P. Siba, M. Laman,

T. M. E. Davis, L. Manning, I. Mueller, L. Schofield, γδ T cells and CD14+ Monocytes Are

Predominant Cellular Sources of Cytokines and Chemokines Associated With Severe Malaria,

The Journal of Infectious Diseases **210**, 295–305 (2014).

49. J. N. Che, O. P. G. Nmorsi, B. P. Nkot, C. Isaac, B. C. Okonkwo, Chemokines responses to

Plasmodium falciparum malaria and co-infections among rural Cameroonians, *Parasitology*

International **64**, 139–144 (2015).

50. M. N. Ajuebor, Y. Jin, G. L. Gremillion, R. M. Strieter, Q. Chen, P. A. Adegboyega, γδT

Cells Initiate Acute Inflammation and Injury in Adenovirus-Infected Liver via Cytokine-

Chemokine Cross Talk, *Journal of Virology* **82**, 9564–9576 (2008).

51. D. L. Hill, S. Innocentin, J. Wang, E. A. James, J. C. Lee, W. Kwok, M. Zand, E. J. Carr, M.

A. Linterman, Impaired HA-specific T follicular helper cell and antibody responses to, ,

doi:10.1101/2021.04.07.21255038.

52. V. Ryg-Cornejo, L. J. Ioannidis, A. Ly, C. Y. Chiu, J. Tellier, D. L. Hill, S. P. Preston, M.

Pellegrini, D. Yu, S. L. Nutt, A. Kallies, D. S. Hansen, Severe Malaria Infections Impair

Germinal Center Responses by Inhibiting T Follicular Helper Cell Differentiation, *Cell Reports*

14, 68–81 (2016).

53. K. Sondén, K. Wyss, I. Jovel, A. Vieira Da Silva, A. Pohanka, M. Asghar, M. V. Homann, L.

L. Gustafsson, U. Hellgren, A. Färnert, High Rate of Treatment Failures in Nonimmune

Travelers Treated With Artemether-Lumefantrine for Uncomplicated Plasmodium falciparum

Malaria in Sweden: Retrospective Comparative Analysis of Effectiveness and Case Series,

Clinical Infectious Diseases Malaria Treatment Failures in Travelers • CID **2017**, 199–206.

54. M. Asghar, V. Yman, M. V. Homann, K. Sondén, U. Hammar, D. Hasselquist, A. Färnert,

Cellular aging dynamics after acute malaria infection: A 12-month longitudinal study, *Aging Cell*

17, e12702 (2018).

55. H. Wickham, M. Averick, J. Bryan, W. Chang, L. D', A. Mcgowan, R. François, G.

Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Lin Pedersen, E. Miller, S. M. Bache, K.

Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K.

Woo, H. Yutani, RStudio 2 cynkra 3 Redbubble 4 Erasmus University Rotterdam 5 Flatiron

Health 6 Department of Integrative Biology, *Journal of Open Source Software* **4**, 1686 (2019).

56. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing, *Journal of the Royal Statistical Society: Series B*

(Methodological) **57**, 289–300 (1995).

57. D. Makowski, M. Ben-Shachar, I. Patil, D. Lüdecke, Methods and Algorithms for

Correlation Analysis in R, *Journal of Open Source Software* **5**, 2306 (2020).

58. Z. Gu, L. Gu, R. Eils, M. Schlesner, B. Brors, Genome analysis circlize implements and

enhances circular visualization in R, **30**, 2811–2812 (2014).

59. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in

multidimensional genomic data, *Bioinformatics* (2016), doi:10.1093/bioinformatics/btw313.

60. FOR THE TREATMENT OF MALARIA GUIDELINES, (2015) (available at

www.who.int).

Acknowledgments:

We would firstly like to acknowledge the contribution of all the study participants. We would

also like to thank all involved clinicians, nurses especially Irene Nordling and Debbie Ribjer for

sampling and Fariba Foroogh for sample processing and organization. Further, we would like to

thank the team from the Translational Plasma Profile Facility at SciLifeLab for support and the

generation of data for this project. Figure 1A and 7G were created with BioRender.com.

Funding:

The Swedish Research Council grant 2019-01940 (CS)

Magnus Bergvall foundation grant 2017-02043 and 2018-02656 (CS)

Åke Wiberg foundation grant M18-0076 (CS)

Swedish Research Council 2015-02977, 2018-02688 and 2018-04468 (AF)

Region Stockholm 20150135 and 20180409 (AF)

Marianne and Marcus Wallenberg Foundation (AF)

Author contributions:

Conceptualization: CS, MJL, AF

Methodology: CS, MJL, NK, SA

Investigation: CS, MJL, KS, VY, NK, SA, DFP, AF

Visualization: CS, MJL

Funding acquisition: CS, AF

Project administration: CS

Supervision: CS, AF

Writing – original draft: CS, MJL

Writing – review & editing: CS, MJL, AF, KS, VY, NK, SA, DFP

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

The flow cytometry data and plasma protein data (Olink Inflammation panel) generated during

this study are available upon request. Code used in the analyzes is available in repository:

https://github.com/LautenbachMJ/MalariaTraveller.