ISL1 is necessary for auditory neuron development and contributes towards tonotopic organization

Authors: Iva Filova[1,7], Kateryna Pysanenko[2,7], Mitra Tavakoli[1], Simona Vochyanova[1], Martina Dvorakova[1], Romana Bohuslavova[1], Ondrej Smolik[1], Valeria Fabriciova[1], Petra Hrabalova[1], Sarka Benesova[3], Lukas Valihrach[3], Jiri Cerny[4], Ebenezer N. Yamoah[5], Josef Syka[2], Bernd Fritzsch[6], and Gabriela Pavlinkova[1*]

Affiliations:
[1]Laboratory of Molecular Pathogenetics, Institute of Biotechnology CAS, 25250 Vestec, Czechia
[2]Department of Auditory Neuroscience, Institute of Experimental Medicine CAS, 14220 Prague, Czechia
[3]Laboratory of Gene Expression, Institute of Biotechnology CAS, 25250 Vestec, Czechia
[4]Laboratory of Light Microscopy, Institute of Molecular Genetics CAS, 14220 Prague, Czechia
[5]Department of Physiology, School of Medicine, University of Nevada Reno, NV 89557, USA
[6]Department of Biology, Department of Otolaryngology, University of Iowa, Iowa City, IA 52242-1324, USA
[7]These authors contributed equally: Iva Filova, Kateryna Pysanenko.
[*] Email: Gabriela.Pavlinkova@ibt.cas.cz

Abstract

A cardinal feature of the auditory pathway is frequency selectivity, represented in a tonotopic map from the cochlea to the cortex. The molecular determinants of the auditory frequency map are unknown. Here, we discovered that the transcription factor ISL1 regulates the molecular and cellular features of auditory neurons, including the formation of the spiral ganglion and peripheral and central processes that shape the tonotopic representation of the auditory map. We selectively knocked out *Isl1* in auditory neurons using *Neurod1^{Cre}* strategies. In the absence of *Isl1*, spiral ganglion neurons migrate into the central cochlea and beyond, and the cochlear wiring is profoundly reduced and disrupted. The central axons of *Isl1* mutants lose their topographic projections and segregation at the cochlear nucleus. Transcriptome analysis of spiral ganglion neurons shows that *Isl1* regulates neurogenesis, axonogenesis, migration, neurotransmission-related machinery, and synaptic communication patterns. We show that peripheral disorganization in the cochlea affects the physiological properties of hearing in the midbrain and auditory behavior. Surprisingly, auditory processing features are preserved despite the significant hearing impairment, revealing central auditory pathway resilience and plasticity in *Isl1* mutant mice. Mutant mice have a reduced acoustic startle reflex, altered prepulse inhibition, and characteristics of compensatory neural hyperactivity centrally. Our findings show that ISL1 is one of the obligatory factors required to sculpt auditory structural and functional tonotopic maps. Still, upon *Isl1* deletion, the ensuing central compensatory plasticity of the auditory pathway does not suffice to overcome developmentally induced peripheral dysfunction of the cochlea.

Keywords

spiral ganglion neurons, inferior colliculus, auditory nuclei, inner ear, brain plasticity, *Isl1* mutation, auditory maps, auditory behavior

Introduction

Spiral ganglion neurons (SGNs) are bipolar, extending peripheral processes to the hair cells within the sensory epithelium (the organ of Corti) and central axons towards the cochlear nucleus (CN) complex, the first auditory nuclei in the brain. Sound-induced vibrations that reach the cochlea are amplified by the outer hair cells (OHCs) organized in three rows and innervated by type II SGNs. The inner hair cells (IHCs) receive, transduce, and transmit the auditory signal to type I SGNs that convey the signal via the vestibulocochlear cranial nerve to the CN of the brainstem. The auditory neurons are organized within the cochlea in an orderly fashion according to frequency, with high frequencies at the base and low frequencies at the apex [1, 2]. The tonotopic organization of type I SGNs corresponds to multiple diversities in their molecular profiles, connectivity patterns, and physiological features along the tonotopic axis [3, 4, 5]. The cochleotopic or tonotopic pattern is maintained throughout the auditory pathways in the brain [6]. The central auditory pathway transmits ascending acoustic information from the CN through the lateral lemniscus complex, the inferior colliculus in the midbrain, the medial geniculate nucleus of the thalamus to the auditory cortex [7]. The efferent motor neurons consist of the medial olivocochlear motor neurons, which modulate cochlear sound amplification by OHCs. In contrast, lateral olivocochlear motor neurons innervate afferent type I sensory neurons and regulate cochlear nerve excitability [7, 8, 9].

The cellular and molecular regulation of neuronal migration and the establishment of tonotopic connections to the hair cells or neurons of the hindbrain's first auditory nuclei, the CN, are not fully understood. Several transcription factors govern the development of inner ear neurons, including NEUROG1 [10], NEUROD1 [11], GATA3 [12, 13], and POU4F1 [14]. The transcription factor NEUROD1 is vital for the differentiation and survival of inner ear neurons [15, 16, 17]. We previously demonstrated that *Isl1^Cre*-mediated *Neurod1* deletion (*Neurod1CKO*) results in a disorganized cochleotopic projection from SGNs [18], affecting acoustic information processing in the central auditory system of adult mice at the physiological and behavioral levels [19]. During inner ear development, ISL1 is expressed in the differentiating neurons and sensory precursors [18, 20, 21, 22, 23]. *Isl1* is expressed in all four populations of SGNs (type Ia, Ib, Ic, and type II) identified by single-cell RNA sequencing [4]. Studies suggest that ISL1 plays a role in developing neurons and sensory cells, but there has been no direct evaluation of ISL1 function in the inner ear. Using *Neurod1^Cre* [24], we conditionally deleted *Isl1* (*Isl1CKO*) in neurons without directly affecting the development of the inner ear sensory epithelium.

This work provides the first genetic and functional evidence for ISL1's role in establishing the spiral ganglion peripheral projection map and proper central auditory circuitry. Most *Isl1CKO* neurons migrated into the center of the cochlear modiolus. They extended outside the cartilaginous/bony otic capsule, in contrast, to control animals with an arrangement of SGNs in parallel to the spiraling cochlear duct. Additionally, we analyzed the transcriptome of neurons, hearing function, sound information processing in the inferior colliculus, and auditory behavior of *Isl1CKO* to demonstrate how abnormal SGN development affects the formation, wiring, and operation of the auditory sensory maps.

Results

ISL1 controls neuronal phenotype in the cochlea

To investigate the role of ISL1 in inner ear neuron development, we eliminated *Isl1* by crossing *Isl1^loxP/loxP* mice [25] with *Neurod1^Cre* mice [24]. *Neurod1^Cre* is expressed in sensory neurons but not in the sensory epithelium, as visualized by *tdTomato* reporter expression (Supplementary Fig. 1).

Figure 1. *Isl1* **deletion results in abnormalities in cochlear innervation and the formation of the spiral ganglion.** (**a**) Diagram of the organization of the cochlea. Spiral ganglion neurons (SGNs) are organized tonotopically from the apex to the base of the cochlea. High-frequency sounds maximally stimulate the base of the cochlea, whereas the largest response to low-frequency sounds occurs in the cochlear apex. (**b**) The top view diagram onto the sensory epithelium shows outer hair cells (OHCs), inner hair cells (IHCs), and SGNs. Type I neurons extend radial fibers (RF) toward IHCs (5-30 neurons innervate one IHC), and type II neurons (representing 5% of SGNs) receive input from OHCs. The central processes of SGNs relay acoustic information to the brain. Efferent axons from the superior olivary complex, forming the intraganglionic spiral bundle (IGSB), innervate OHCs. (**c-f**) Representative images of whole-mount immunolabeling of the base and apex with anti-Myo7a (hair cell marker) and anti-α-tubulin (nerve fibers) show reduced and disorganized RF and missing or
(Continued)

Figure 1 *continued*

altered efferent fibers forming IGSB (arrowhead) in *Isl1CKO* compared to control. Abnormalities in innervation, including larger gaps between radial fiber bundles, and fiber bundles bypassing OC (arrowhead in f), are noticeable in *Isl1CKO*. **(g-i)** The shape of the spiral ganglion (SG) is shown in the basal and apical half of the whole-mounted cochlea of control-*Ai14* reporter mice at P0. TdTomato$^+$ neurons form a spiral located in the Rosenthal's canal and parallel to the OC (anti-Myo7a labeled hair cells), nerve fibers are labeled with anti-tubulin. **(j-l)** In *Isl1CKO*, only some tdTomato$^+$ neurons are in the Rosenthal's canal (delineated by white dotted lines), but many neurons are in the centre of the cochlea, a conical-shaped structure, the modiolus (yellow arrowhead). **(m, n)** The vibratome sections of the cochlea labeled by anti-NeuN (a nuclear marker of differentiated neurons) and anti-tubulin (nerve fibers) show the unusual position of cochlear neurons in the modiolus, and neurons entwined in central axons in *Isl1CKO* compared to control (white lines delineate a normal position of the SG). **(o)** The length of radial fibers was measured in the confocal images from cochlear whole-mount preparations in the apex, mid-apex, mid-base, and base. Data are expressed as mean ± SEM (n = 3 radial fiber bundles/each region/3 cochlea/genotype); unpaired *t*-test. **(p)** Fiber density was quantified in the base, mid-base, and mid-apex (n = 3 mice per genotype). Error bars represent mean ± SD; unpaired *t*-test. Scale bars: 100 μm.

Analyses of ISL1 expression in *Isl1CKO* confirmed efficient recombination by *Neurod1^{Cre}* with
virtually no expression of ISL1 during the differentiation of neurons in the inner ear ganglion as
early as embryonic day E10.5, and later in the cochlear neurons, shown at E14.5 and postnatal
day 0 (P0), in contrast to high levels of ISL1 in neurons in the control inner ear (Supplementary
Fig. 2). It is worth mentioning that no difference was observed in the density of Schwann cells
(Supplementary Fig. 2e, f).
The organization of the sensory epithelium in the organ of Corti of *Isl1CKO* was
comparable to controls with three rows of OHCs and one row of IHCs (Fig. 1a-f), as shown by
immunolabeling with the hair cell differentiation marker, myosin VIIA (Myo7a). Whole-mount
anti-tubulin staining of innervation showed reduced and disorganized radial fibers. Significant
gaps were found between radial fiber bundles, crisscrossing fibers, and an unusual dense
innervation in the apex, with some large fiber bundles bypassing the organ of Corti and extending
to the lateral wall in the *Isl1CKO* cochlea (arrowhead, Fig. 1f).
Next, we evaluated the effects of *Isl1* deletion on the formation of the spiral ganglion. The
SGN somas were well restricted to the Rosenthal's canal in control animals, arranged in a spiral
parallel to the cochlear duct (Fig. 1g-i). In contrast, unevenly distributed neurons were found in
Isl1CKO with many SGNs accumulated beyond the Rosenthal's canal, in the conical central part
of the cochlea, the modiolus (Fig. 1j-l), which usually carries only afferent and efferent fibers. The
unusual accumulation of cochlear neurons intertwined with the central axons in the modiolus of
Isl1CKO compared to the control mice is depicted in the immunolabeled sections of the cochlea
(Fig. 1m, n). To determine whether neuron number was affected by apoptosis during inner ear
development, we performed immunostaining for cleaved Caspase-3 (Supplementary Fig. 3). We
found moderately increased apoptosis of neurons in *Isl1CKO* at E12.5 but not at E10.5, although
the size of inner ear ganglia was unaffected at E12.5. These results imply that missing neurons in
the Rosenthal's canal are due to the altered migration of SGNs rather than neuronal death in the
developing inner ear. Correspondingly, most apoptotic neurons were found outside the cochlea in
the vestibular ganglion area in E14.5 *Isl1CKO*. In line with the unusual position of SGNs in the
Isl1CKO cochlea, the radial fibers were significantly lengthened in all regions of the *Isl1CKO*
cochlea (Fig. 1o), but the overall radial fiber density was reduced (Fig. 1p).
Interestingly, *Isl1* deficiency caused a shortening of the cochlea, which was on average
~20% and 25% shorter at P0 and in adults, respectively (Fig. 2a, b). Since *Neurod1^{Cre}* is expressed

Figure 2. The length of the organ of Corti is shortened, the apical end sensory epithelium is disorganized, and presynaptic ribbons are reduced in *Isl1CKO*. (**a**) The organ of Corti of *Isl1CKO* is shorter than control, as shown by anti-Myo7a labeling of hair cells in the cochlea at P0. (**b**) The length of the organ of Corti was measured at postnatal day P0 and in 2-month-old mice. Error bars represent mean ± SD; unpaired *t*-test. (**c**) Immunohistochemistry for prestin (a marker for OHCs) and calretinin (a marker for IHCs) shows a comparable cytoarchitecture of the organ of Corti in the cochlear mid-base of controls and *Isl1CKO* but disorganized rows of OHCs in the apical end with multiple OHC rows and ectopic calretinin[+] IHCs among OHCs (arrowheads) in *Isl1CKO* at P35. (**d**) Higher-magnification images show a detail of anti-tubulin labeled innervation with fibers forming three parallel outer spiral bundles (arrows) that innervate multiple OHCs and turn toward the base in the control cochlea. Guiding defects in the extension of these fibers to OHCs are obvious in *Isl1CKO* with some fibers randomly turned toward the apex (arrowheads); note the disorganization of radial fiber bundles in *Isl1CKO*. (**e**) Representative images of whole-mount immunolabeling of the cochlear mid-base with calretinin (IHCs) and anti-CtBP2 (presynaptic ribbons) of 2-month-old mice. (**f**) CtBP2[+] synapses were counted in 10 IHCs per each cochlear region (n = 6 mice per genotype). Box plots indicate median (middle line), 25th, 75th percentile (box) and min to max (whiskers); multiple *t*-tests, ****P < 0.000001. HS, Hoechst nuclear staining; IHC, inner hair cell; OHC, outer hair cell. Scale bars: 100 μm (a), 50 μm (c), 20 μm (d), 10 μm (e).

only in neurons, the shortening of the cochlea appears to be a secondary effect of the abnormalities

in neuronal development. A similar phenotype of truncated growth of the cochlea was reported for
Neurod1 and *Neurog1* mutants because of alternations in spatiotemporal gene expression [10, 18, 19,]
[26]. Correspondingly to the truncated cochlear phenotype [10, 18, 19, 26], detailed analyses showed
abnormalities in the epithelium at the apical end with disorganized rows of OHCs and ectopic
IHCs among OHCs in the *Isl1CKO* cochlea (Fig. 2c). The disorganized apical epithelium
represented ~11.8% of the length of the organ of Corti in adults (n = 6). Anti-tubulin labeling
revealed a reduced and disorganized innervation in the region of OHCs in *Isl1CKO* (Fig. 2d). In
the controls, fibers extending through the tunnel of Corti were oriented towards the base to form
three parallel outer spiral bundles. In contrast, guiding defects in the extension of these fibers in
Isl1CKO were apparent, with neurites extending with random turns towards the base and the apex
(arrowheads in Fig. 2d). To further investigate cochlear histopathology in *Isl1CKO*, we evaluated
synaptic contacts from type I SGNs onto IHCs in 2-month-old mice (Fig. 2e). CtBP2$^+$ ribbon
synapse number was reduced in *Isl1CKO*, representing ~46%, 51%, 56%, and 69% ribbons of the
control base, mid-base, mid-apex, and apex, respectively (Fig. 2f, Supplementary Table 1).
Altogether, these results demonstrate the prominent effects of *Isl1* deletion on the
formation of the spiral ganglion and innervation patterns in the cochlea. The truncated cochlea in
Isl1CKO indicates an altered spatial base-to-apex organization of structural and physiological
characteristics of auditory neurons.
145
ISL1 regulates neuronal identity and differentiation programs of SGNs
To gain insight at the molecular level on how the *Isl1* elimination causes the neuronal phenotype
in the cochlea, we sought to identify potential ISL1 targets through global transcriptome analysis.
We opted to use Bulk-RNA sequencing to obtain sequencing depth and high-quality data [27]. Six
biological replicates were used per genotype, and each replicate contained a total of 100 tdTomato$^+$
SGNs isolated from the E14.5 cochlea. Spiral ganglia were dissected, dissociated into single cells,
and fluorescent tdTomato$^+$ cells were FACS-sorted (Fig. 3a). Compared to controls, 650 protein-
coding genes were differentially expressed in *Isl1CKO* neurons (adjusted p-value < 0.05, and fold
change > 2 cut-off values, see Methods), 332 genes down- and 318 genes up-regulated (Fig. 3b,
Supplementary Data 1). Gene ontology (GO) term enrichment analysis for the GO term category
biological process revealed highly enriched GO terms associated with neuron development,
including "neurogenesis", "neuron differentiation", and "nervous system development", in both
up- and down-differentially expressed genes (Fig. 3c, Supplementary Data 2). The most enriched
and specific GO categories for downregulated genes were associated with neurotransmission-
related machineries, such as "transmembrane transporter", "voltage-gated channel", "cation and
ion transport", and "membrane potential regulation", indicating changes in neuronal cell functions.
The analysis identified enrichment of downregulated genes involved in axon development,
guidance, axonogenesis, and neuronal migration. These genes included members of all four classes
of axon guidance molecules and their cognate receptors, the ephrins-Eph, semaphorins-plexin,
netrin-unc5, and slit-roundabout [28], for instance, *Epha5, Epha4, Pdzrn3, Sema3e, Sema6d, Plxna4,*
Ntn3, Ntng1, Ntng2, Kirrel3, Unc5b, Unc5c, Slitrk3, and *Slitrk1*. The neurotrophic tyrosine kinase
receptors (*Ntrk2* and *Ntrk3*) and a G-protein-coupled chemokine receptor (*Cxcr4*), important
regulators of neuronal migration, were also downregulated [29].
In contrast, upregulated genes particularly enriched in *Isl1CKO* neurons were associated
with "regulation of synapse organization, synapse structure or activity" and "regulation of synapse
assembly", indicating compensatory changes relating to the ability of neurons to form neuronal
projections towards their targets. Upregulated genes encoding molecules critical for synapse

Figure 3. ISL1-mediated transcription signature in cochlear neurons. (a) Image of the whole-mount cochlea with genetically labeled tdTomato neurons in the spiral ganglion at E14.5 [Hoechst staining (blue), Scale bar: 100 μm]; and workflow depicts microdissection, dissociation, FACS sorting of single tdTomato⁺ spiral ganglion neurons for a bulk of 100 cells RNAseq analysis. **(b)** The volcano plot shows the change in protein-coding gene expression levels in the *Isl1CKO* compared to control spiral ganglion neurons (adjusted p-value < 0.05, and fold change > 2 cutoff values). The complete list of identified 332 down- and 318 up-differentially expressed genes is in Supplementary Data 1. **(c)** Enrichment map of down- and upregulated gene ontology (GO) sets visualized by the network. Each node represents a GO term; edges depict shared genes between nodes. Node size represents a number of genes of the mouse genome per the GO term, and node fill color represents a GO term significance. Each GO set cluster was assigned with representative keywords; a list of GO sets is available in Supplementary Data 2).

formation and adhesion included several members of the cadherin superfamily (*Cdh7, Cdh10,*
Pcdh8, Pcdh9, Pcdh10, Pcdh18, and *Pcdh19*), adhesion-related genes (*Ptprd, Ptprs, Cntnap2,*
Itga4, Itga5), and the ephrin ligands (*Efna3, Efna5*). Interestingly, "synaptic signaling" was among
the GO terms with significant representation in both up- and down-regulated genes, suggesting
changes in synaptic circuits and neuronal activity. We also found many genes encoding
transcription factors and signaling molecules, indicating that ISL1 may act through transcription
networks instead of defined target genes. The expression of neural-specific bHLH factors
important for differentiation and maturation of neurons was increased in *Isl1CKO*, including
members of NeuroD (*Neurod6, Neurod2*), Nscl (*Nhlh2*), and Olig (*Bhlhe23*) families [30]. Genes
encoding LIM domain proteins, the transcription factors *Lhx1, Lhx2*, *Lmo2*, *Lmo3*, a core member

of the PCP signaling paradigm, *Prickle1*, and neuronal homeobox genes, *Emx2*, *Uncx*, and *Pknox2* were upregulated in *Isl1CKO* neurons. *Prickle1* is an essential regulator of neuron migration and neurons' distal and central projections in the cochlea [31]. Some of the identified genes encoding regulatory molecules shown to be essential for neuronal development, the formation of SGNs, and their projections were downregulated in *ISL1CKO* neurons, such as the signaling molecules *Shh* [32], *Wnt3* [3], and *Fgfs* (*Fgf10, Fgf11,* and *Fgf13*) [33, 34], and the transcription factors *Gata3* [12, 13], *Irx1*, *Irx2*, *Pou3f2*, *Pou4f2* [22], and *Runx1* [3, 4].

Consistent with the idea that ISL1 may control a gene expression program mediating different aspects of neuronal development in the cochlea, several identified genes are known to affect the formation of SGNs and their innervation patterns. For instance, a mouse model with EphA4 deficient signaling in SGNs results in aberrant mistargeting of type I projections to OHCs [35] and SGN fasciculation defects [36]. Targeted disruptions of the neurotrophin receptors, *Ntrk2* and *Ntrk3*, are associated with misplaced SGNs in the modiolus, disorganized projections, and reduced neuronal survival [37]. The neurotrophin receptor p75NTR (*Ngfr*)-deficient SGNs demonstrate enhanced neurite growth behavior [38]. Mutation of *Prickle1* results in disorganized apical afferents [31]. Reduced and aberrant cochlear innervation was shown in mice with *Gata3*-specific SGN deletion [13]. *Nrp2* expressed in SGNs is required for axon path finding and a normal pattern of cochlear innervation [39]. The axon guidance molecule SLIT2 and its receptor ROBO1/2 control precise innervation patterns, and their deletions induce misplacement of SGNs [40]. Mice lacking *Dcc* have the more severe phenotype of mislocated SGNs and disorganized central projections, although most SGNs of *Dcc*$^{-/-}$ remained within Rosenthal's canal [41].

Other significantly deregulated genes mediate several aspects of neural-circuit formation in the brain, including neurite elongation, axon guidance, and interactions between neurons and microglia. For instance, deletion of *Epha5* alters retinal axon guidance within the midbrain [42]; inhibition of *Ntng1* reduces microglial accumulation and loss of cortical neurons [43]; the absence of either *Bhlhb22* or *Prdm8* alters neural circuit assembly [44]; LHX1 controls neuronal migration and axonal projections of motor neurons [45]; and LHX2 influences axon guidance and the topographical sorting of axons by regulating ROBO1/2 [46]. These molecular differences dovetail well with abnormalities in the innervation pattern and SGN migration defects *Isl1CKO*. Selected differentially expressed genes from the RNAseq analysis, namely *Dcc, Epha5, Ntng1, Prdm8, Robo2, Slit2, Tbx3, Uncx, Gata3, Lhx1, Lhx2, Cdh7, Nhlh2, Ntrk2,* and *Ntrk3,* were further validated by RT-qPCR of RNA isolated from whole inner ears of E14.5 embryos (Supplementary Fig. 4). These changes indicate that ISL1 regulates transcriptional networks that underlie neuronal identity and function during differentiation of SGNs and that *Isl1* elimination results in a major impairment in the development, axonogenesis, migration, and molecular characteristics of these neurons.

Isl1CKO mice have altered hearing function

Considering the substantial abnormalities in the formation of the spiral ganglion, innervation, and molecular neuron features, we assessed the hearing function of *Isl1CKO* mice. We evaluated distortion product otoacoustic emissions (DPOAE) to determine the robustness of OHC function and cochlear amplification. Otoacoustic emissions are a physiological byproduct of an active amplification mechanism when sound-induced vibrations are amplified in a frequency-specific manner by the OHCs of the organ of Corti [47, 48]. Compared to controls, the DPOAE responses of *Isl1CKO* were significantly reduced in the frequency range between 4 and 24 kHz (Fig. 4a). DPOAE amplitudes at frequencies of 26 kHz and higher were comparable between control and

Figure 4. Hearing impairment is detected in *Isl1CKO* mice. (**a**) Distortion product otoacoustic emissions (DPOAEs) show significantly reduced levels in the low and middle-frequency range (4-24 kHz). Data are expressed as mean ± SEM, two-way ANOVA with Bonferroni post-hoc test, *P < 0.05, **P < 0.01, ***P < 0.001 ****P < 0.0001. (**b**) The average auditory brainstem response (ABR) thresholds of *Isl1CKO* and control mice are analyzed by click-evoked ABR. Data are expressed as mean ± SD, two-way ANOVA with Bonferroni post-hoc test, **P < 0.01, ****P < 0.0001. (**c**) Averaged ABR response curves evoked by an 80 dB SPL click; (**e**) by a 90 dB SPL pure tone of 32 kHz frequency; and (**g**) by a 90 dB SPL pure tone of 8 kHz frequency are presented. (**d, f, h**) Averaged individual ABR wave amplitudes are shown for the corresponding peaks. Data are expressed as mean ± SEM, two-way ANOVA with Bonferroni post-hoc test, **P < 0.01 ***P < 0.001, ****P < 0.0001.

Isl1CKO mice. Based on the physiological place-frequency map in the normal mouse cochlea [49], frequencies above 26 kHz are located at the basal half of the cochlea. Thus, decreased DPOAE responses may be attributed to the more profound morphological abnormalities in the apex, disorganized innervation, and disorganized rows of OHCs in the apical end.

We evaluated auditory brainstem responses (ABRs), which measure electrical activity associated with the propagation of acoustic information through auditory nerve fibers. Measurements of ABR thresholds showed that all *Isl1CKO* animals displayed elevated thresholds indicative of hearing loss compared to age-matched control animals, except at frequencies 32 and 40 kHz, which were comparable to ABR thresholds in control mice (Fig. 4b). Using click-evoked ABR, we evaluated waveform characteristics associated with the propagation of acoustic information through the auditory nerve to higher auditory centers (Fig. 4c). Wave I reflects the synchronous firing of the auditory nerve. In contrast, waves II–V are attributed to the electrical activity of downstream circuits in the CN, superior olivary complex, lateral lemniscus, and inferior colliculus [50]. Waves IV and V are products of the sound-evoked neuronal activity in the lateral lemniscus and inferior colliculus [51, 52, 53, 54, 55]. The waves IV and V are often paired and superimposed in commonly used non-transgenic mouse strains [52, 55]. The ABR traces in all our panels were averaged traces from all measured animals. A part of the animals in both experimental groups had a hardly distinguishable wave V in response to the click and 8 kHz sound stimulation; therefore, to prevent bias, we did not include wave V. The amplitudes of ABR waves I-IV were significantly reduced in *Isl1CKO* (Fig. 4d). Since the ABR threshold for 32 kHz and above were comparable between age-matched controls and *Isl1CKO* mutants (Fig. 4b), we used the pure-tone stimuli of 32 kHz to evaluate ABR responses (Fig. 4e). A significant difference amongst the genotypes was found only for amplitude reduction of wave III (Fig. 4f), thus indicating preserved synchronized activities of peripheral and brainstem auditory processing. Although waves I and II amplitude for both mutant and control mice were similar, there were apparent differences in the ABR waveform morphology. The latency of ABR wave I was delayed, the relative interwave latency between peaks I and II was shortened, and the trough between wave I and II diminished, resulting in a fusion of both peaks in *Isl1CKO*. A delay of the leading peak of ABR wave I recovered towards ABR wave III. The wave I and II changes reflect abnormalities in the summated response from SGNs, auditory nerve fibers, and most likely the CN.

Additionally, we used a pure-tone stimulus of 8 kHz to evaluate ABR responses (Fig. 4g), as ABR thresholds for 8 kHz were significantly reduced for *Isl1CKO* mice. In contrast to ABR amplitudes at 32 kHz stimuli, ABR amplitudes for 8 kHz pure-tone stimuli were significantly reduced for I-III waves (Fig. 4h). The results indicate abnormalities in the cochlear auditory neurons and propagation of acoustic information through auditory nerve fibers to higher auditory centers. We observed a marked drop in the wave I growth function. Still, by comparing wave I and IV peaks, an increased central gain was noted in *Isl1CKO* (Supplementary Fig. 5), indicating compensatory plasticity at higher auditory circuits for cochlear damage with diminished afferent input [56].

Isl1CKO mice display abnormalities in the ascending auditory pathways

Having recognized uneven hearing function loss, as measured by ABRs, we further wanted to investigate the distribution of SGNs in the *Isl1CKO* cochlea and the morphology of the components of the central auditory pathway. We generated 3D visualization of the cochlea using light-sheet fluorescent microscopy (Fig. 5a-i, Supplementary Videos S1-S4). In the first cochlear preparation, neurons were visualized by *tdTomato* reporter expression and hair cells in the organ

Figure 5. Neurons in the *Isl1CKO* inner ear are abnormally distributed. (a-i) Microdissected cochleae of reporter control-*Ai14* and *Isl1CKO-Ai14* mice were cleared (CUBIC protocol), immunolabeled, imaged, and reconstructed in 3D using light-sheet fluorescence microscopy (see Supplementary Videos S1-S4). (**a**) In the control cochlea, tdTomato$^+$ SGNs form a coil of the spiral ganglion (SG) in the Rosenthal's canal and anti-Myo7a labeled hair cells show the spiral shape of the organ of Corti. Parts of the vestibular ganglion (VG) with tdTomato$^+$ neurons and the saccule with Myo7a$^+$ hair cells are shown. (**b, c**) Anti-NeuN (a nuclear marker of differentiated neurons) co-labeled tdTomato$^+$ neurons form SG and VG in the control cochlea; the modiolus contains the central processes, forming the auditory nerve (AN). (**d-f**) In *Isl1CKO-Ai14*, tdTomato$^+$ neurons are mainly located in the conical central part of the cochlea, the modiolus, in contrast to the spiral of the organ of Corti with *(Continued)*

Figure 5 *continued*

Myo7a[+] hair cells. Some *Isl1CKO* neurons form a segment of the spiral ganglion in the Rosenthal's canal (RC) in the mid-base region (arrowhead), shown in detail (**f**), the RC area is highlighted by dotted lines. (**g-h**) Abnormal and uneven distribution of cochlear tdTomato[+] neurons is shown by anti-NeuN co-labeling. Many neurons are misplaced in the modiolus, but some neurons are in their close-normal position in RC (arrowhead). (**j**) Schematic drawing of the mouse cochlea shows the cochlear place-frequency map for four cochlear regions termed the apex (corresponding to the best encoding frequencies ∼8 kHz at the distance from the basal end of 82 %), mid-apex (∼19 kHz at 49% length), mid-base (∼32 kHz at 32% length), and basal turn (∼62 kHz at 9% length) [49]. (**k**) Whole mount images of the P10 inner ear show distribution of tdTomato labeled neurons in superior (sVG) and inferior vestibular ganglia (iVG), and SG in the control-*Ai14*. (**l**) In the *Isl1CKO-Ai14* mutant, the VG is fused and enlarged, spatial segregation between VG and SG is diminished, forming an aberrant "spiro-vestibular" ganglion, and SG has lost its spiral shape. SGNs, spiral ganglion neurons. Scale bars: 200 μm, except for 100 μm (b, i).

277 of Corti were labeled using anti-Myo7a. In the second preparation, tdTomato[+] neurons were co-
278 labeled with anti-NeuN, a nuclear marker of differentiated neurons. Compared to the recognizable
279 coil of cochlear neurons of the spiral ganglion in the controls (Fig. 5a-c, Supplementary Videos
280 S1, S2), SGNs were aberrantly distributed in the *Isl1CKO* cochlea (Fig. 5d-i, Supplementary
281 Videos S3, S4). Many *Isl1CKO* neurons were in the center of the cochlea, the modiolus, that
282 normally contains only central processes of SGNs, as shown in the control cochlea (Fig. 5b, c).
283 Some *Isl1CKO* neurons were found in their proper position in the Rosenthal's canal, forming a
284 dense strip of neurons, suggesting a partially preserved organization of the spiral ganglion (Fig.
285 5f, i). This segment of the spiral ganglion formed in the mid-base of the *Isl1CKO* cochlea correlates
286 with the estimated characteristic place in the cochlear place-frequency map for the characteristic
287 frequency of 32 kHz [49] (Fig. 5j). A partial preservation of the spiral ganglion in the mid-base
288 corresponds to the retained high-frequency hearing function of *Isl1CKO*. Identical to the SGN
289 migration abnormalities recognized at P0, most SGNs were in the modiolus, outside of Rosenthal's
290 canal, in *Isl1CKO* adult mice (Supplementary Fig. 6). Apart from the changes in the cochlear
291 ganglion, the vestibular ganglion was fused and enlarged in *Isl1CKO* compared to distinct superior
292 and inferior vestibular ganglia in controls (Fig. 5k, l). The spatial segregation between the
293 vestibular and spiral ganglion was diminished in *Isl1CKO*, with some areas fused, and thus,
294 forming an aberrant "spiro-vestibular" ganglion.
295   Using dye tracing, we evaluated the segregation of central axons of the auditory nerve (the
296 cranial nerve VIII; Fig. 6a, schematic view of dye applications). In controls, the central axons
297 labeled by dyes applied into the base and apex are segregated in the auditory nerve and from the
298 vestibular ganglion, labeled by dye injected into the vestibular end-organs (Fig. 6b, Supplementary
299 Fig. 7). In contrast, the central axons from the cochlear base and apex virtually overlapped in the
300 auditory nerve (yellow fibers), and neurons labeled by cochlear dye applications were detected to
301 be intermingled with vestibular neurons in an aberrant "spiro-vestibular" ganglion in *Isl1CKO* (Fig.
302 6c). Unfortunately, the mixing of spiral and vestibular ganglion neurons in *Isl1CKO* mice
303 precluded a full quantitative assessment. The unusual fibers labeled by the apical dye application
304 were looping around the vestibular ganglion in *Isl1CKO*.
305   The CN is the first structure of the ascending auditory pathways, where the auditory nerve
306 fibers project. The auditory nerve bifurcated with one branch, synapsing in the posteroventral
307 (PVCN) and dorsal (DCN) CN and the other innervating the anteroventral CN (AVCN; Fig. 6a).

Figure 6. *Isl1CKO* neurons project unsegregated and disorganized central projections to the cochlear nucleus. (a) The schematic diagram visualizes dye tracing from the inner ear and its connections *(Continued)*

Figure 6 *continued*

to the auditory brainstem, using insertions of differently colored NeuroVue dyes into the vestibular end-organ (utricle, magenta), cochlear base (red), and apex (green). Axonal projections from cochlear neurons to the cochlear nucleus (CN) bifurcate with one branch synapsing in the dorsal (DCN) and the other innervating the anteroventral CN (AVCN). **(b)** Images of individual colors of the separate channels and a merged image show distinct and spatially restricted bundles of neuronal fibers of the auditory nerve (AN) projecting to the CN and the vestibular ganglion (VG) in controls, labeled by dye applications into the apex (green), base (red), and utricle (magenta). The only mixed bundle (yellow) is inner ear efferents (IEE). **(c)** In *Isl1CKO*, the segregation of central axons is lost, as fibers labeled from the apex (green) and base (red) are mainly overlapping in the AN (yellow fibers), and neurons labeled by dyes applied into the cochlear base and apex are mixed with VG neurons to form an aberrant ganglion, the "spiro-vestibular" ganglion. Note that fibers labeled from the apex (green) form an unusual fiber loop around the ganglion, and no IEE are recognizable in *Isl1CKO*. **(d)** The tonotopic organization of the CN subdivisions is shown with low-frequency afferents labeled from the apex (green) and high frequency from the base (red) and organized as parallel fibers in isofrequency bands in controls at E16.5. **(e)** In *Isl1CKO*, cochlear afferents enter AVCN as a single bundle instead of forming separate basal and apical projections. The branch synapsing in the AVCN is reduced and disorganized. Arrowheads indicate overlapping fibers. The DCN branch is represented by just a few fibers in *Isl1CKO* (inset in **e**). **(f, g)** Images of individual colors of the separate channels represent projections labeled from the apex (green) and base (red), shown as a merged image in e. **(h, i)** The confocal section shows segregated cochlear afferents from the base and apex, and vestibular afferents extend toward the control's lateral vestibular nucleus (LVN). In contrast, cochlear afferents of *Isl1CKO* overlap (yellow fibers), and some basal afferents extend toward the CN, and some project toward the LVN and loop back to the CN. **(j)** Graphic summary of aberrant location and projections of auditory neurons in *Isl1CKO*. In the control, the auditory system is tonotopically organized, including neurons of the spiral ganglion, the auditory nerve, and the CN subdivisions. In *Isl1CKO*, excluding some portion of neurons in the base, neurons are located outside of Rosenthal's canal. Some neurons projecting to the cochlea are mixed with vestibular neurons. The central axons from the apex and base are not segregated in the AN and AVCN. Projections entering the DCN are diminished in *Isl1CKO*. AC, anterior crista; D, dorsal; HC, horizontal crista; P, posterior axes; PVCN, posteroventral CN; PC, posterior crista; RB, restiform body; r1- r6, rhombomere 1 - 6; U, utricle; S, saccule; VN, vestibular nerve. Scale bars: 100 μm.

Dye tracing showed segregated projections of apical and basal cochlear afferents forming parallel
isofrequency bands in controls at E16.5 (Fig. 6d). In contrast, comparable injections in *Isl1CKO*
showed that axonal projections to the CN were reduced, restricted, disorganized, and lacked clear
apex- and base-projection segregations, as many fibers overlapped (Fig. 6e-g). Only a few fibers
can occasionally be seen expanding to the DCN in *Isl1CKO* at E16.5 (inset in Fig. 6e)*. Aberrant
central projections were further demonstrated by cochlear neurons projecting to the lateral
vestibular nucleus (LVN) in *Isl1CKO* (Fig. 6i). No such projections were found in control mice
with segregated vestibular projections to the vestibular nucleus, and basal and apical afferents
reaching the CN (Fig. 6h). Collectively, these data show disorganized central projections and a
loss of tonotopic organization of both the auditory nerve and the CN in *Isl1CKO,* as the apical and
basal projections are not completely segregated (see graphic summary in Fig. 6j).
Since the size and number of neurons in the CN depend on input from the auditory nerve
during a critical development period up to P9 [57], we also analyzed the volume of the CN of
Isl1CKO. The volume of the CN of adult mutants was reduced by approximately 50% compared
to controls at postnatal day P35 (Fig. 7a, b, g). As *Isl1* is not expressed in the CN [19], the reduced
size is likely a secondary effect associated with reduced afferent input consistent with the
previously reported impact of neonatal cochlear ablation [2]. The spherical and globular bushy cells
are principal cells that receive large auditory nerve endings, called "endbulbs of Held" and

"modified endbulbs", specialized for precise temporal firing [1, 58]. Using anti-parvalbumin to label
spherical bushy cells and anti-VGlut1 to label auditory-nerve endbulbs of Held [59, 60], we
demonstrated that auditory afferents of *Isl1CKO* and controls formed comparable clusters of
boutons that wrap the somas of their targets (Fig. 7c-f). Although the CN of *Isl1CKO* was smaller,
the density of bushy cells (number of parvalbumin$^+$ neurons per μm^2) was similar in controls and *Isl1CKO* (Fig. 7h).

Figure 7. The cochlear nucleus of *Isl1CKO* is reduced. (a, b) Coronal sections of a brain (immunostained with anti-NeuN) of adult controls and *Isl1CKO*, show the DCN and VCN; the dotted line indicates the boundaries of the CN. **(c-f)** Representative images of immunolabeled brain sections with anti-parvalbumin to label the spherical bushy cell soma and anti-VGlut1 to label auditory-nerve endbulbs of Held around the bushy cells. Higher-magnification images show the distribution of parvalbumin$^+$ neurons and the presence of VGlut1$^+$ auditory nerve synaptic terminals. **(g)** Quantification of the adult CN volume, adjusted to body weight (n = 5 mice per genotype) and **(h)** parvalbumin$^+$ neurons per μm^2 of the AVCN (n = 3 mice per genotype, 2-4 sections per mouse). Data are expressed as mean ± SD, unpaired *t-test* (***P < 0.001; ns, not significant). D, dorsal; M, medial axis; HS, Hoechst nuclear staining; DCN, dorsal cochlear nucleus; VCN, ventral cochlear nucleus. Scale bars: 200 μm (a-d); 20 μm (e,f).

Characteristics of inferior colliculus neurons are distorted in *Isl1CKO*

We evaluated inferior colliculus (IC) properties, which is the principal auditory structure of the midbrain for the ascending auditory pathways and descending inputs from the auditory cortex [61]. The IC allows for sound localization, integrates multisensory and non-auditory contributions to hearing, and plays an essential role in generating the startle response. We demonstrated no significant IC size reduction in *Isl1CKO* compared to controls (Fig. 8a, b). We compared neuronal characteristics in the central nucleus of the IC of *Isl1CKO* and control animals using multichannel electrodes. Extracellular electrophysiological recordings of neuronal activity in controls showed a well-defined narrow single-peaked profile of the excitatory receptive fields (Fig. 8c). Compared to simple narrow V-shape receptive fields in controls, we recorded the multipeaked broad tuning curves in the IC of *Isl1CKO* mice, suggesting multiple inputs from the lower levels of the auditory system and overall, the worsened tunning characteristics of IC neurons.

The investigation of the responsiveness of IC neurons (IC units) to different sound frequencies showed higher excitatory thresholds in *Isl1CKO* compared to control animals in measured frequencies between 6 to 28 kHz (Fig. 8d, e). However, responsiveness of IC units to recorded frequencies above 28 kHz was comparable between control and mutant mice. These comparable thresholds for high frequencies between control and *Isl1CKO* mice are consistent with the ABR measurements. At the low frequency range, below 6 kHz, we did not record any IC neurons in *Isl1CKO*. Compared to the

Figure 8. Characteristics of IC neurons (units) are affected in *Isl1CKO* mice. (a) Immunostaining of coronal brain sections for NeuN and nuclear staining Hoechst (HS). Scale bars, 200 µm. **(b)** Quantification of the adult control volume and *Isl1CKO* inferior colliculus (IC) adjusted to body weight. Data are expressed as mean ± SD (n = 5 mice per genotype), unpaired *t-test* (ns, not significant). **(c)** Representative examples of tuning curves recorded in the IC display impairments in tuning properties with broad and irregular receptive fields in *Isl1CKO* compared to control mice. **(d)** Scatter diagram shows the distribution of the excitatory thresholds of *(Continued)*

Figure 8 *(Continued)*

IC neurons in dependency on the characteristic frequency (CF) recorded in control and *Isl1CKO* mice. Note, no IC neurons are recorded below 6 kHz in *Isl1CKO* mutants. (**e**) Excitatory thresholds of the IC neurons at different CFs are shown as averages in 0.3-octave bins in control and *Isl1CKO* mice. Data are expressed as mean ± SEM. Two-way ANOVA with Bonferroni post hoc test. **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. (**f**) Comparison of the rate intensity function parameters between control (n = 8) and *Isl1CKO* mice (n = 12): broadband noise (BBN) threshold, dynamic range, spontaneous activity, and maximum response magnitude. Violin plots indicate median (middle line), 25th, and 75th percentile (dotted lines), unpaired *t*-test, ****$P < 0.0001$. (**g**) The sharpness of the neuronal tuning expressed by quality factor Q_{20} (the ratio between the CF and bandwidth at 20 dB above the minimum threshold) averaged in 0.3-octave bins is decreased in *Isl1CKO*. (**h**) Synchronization of units with click trains. Vector strength computed for different inter-click intervals. Data are expressed as mean ± SD, two-way ANOVA with Bonferroni post-hoc test, ****$P < 0.0001$.

controls, the IC neuronal responses in *Isl1CKO* had a higher broadband noise (BBN) threshold, a
narrower dynamic range, higher spontaneous activity, and a lower maximum response magnitude
(Fig. 8f). The results suggest a functional reduction in sensitivity to sound, audibility, intensity
discrimination, and increased excitability of IC neurons in *Isl1CKO*. A commonly used metric unit
of auditory tuning is the "quality factor", or Q, defined as the characteristic frequency (CF) divided
by the bandwidth, measured at 20 dB above the minimum threshold (Q_{20}). Results revealed a
significantly lower quality factor in the mutant mice (Fig. 8g), showing substantially worsened
frequency selectivity.
To evaluate the precise temporal representation of sound into the central auditory system,
we performed an acoustic stimulation of the IC units with trains of five clicks with different inter-
click intervals from 100 ms up to 15 ms. In control mice, increasing time interval between clicks
led to a better synchronization of neuronal responses with the individual clicks in the train,
implying precision and reliability of temporal sound discrimination ability (Fig. 8h). In contrast,
the precise temporal decoding in *Isl1CKO* was disrupted, as the synchronization of neuronal
responses was significantly lower for the whole range of inter-click intervals. In the case of the
Isl1CKO, the synchronization level of neuronal responses remains almost constant, suggesting a
lack of precise temporal sound processing.
391
The auditory behavior of *Isl1CKO* mice is altered
Next, we evaluated the behavioral responses of *Isl1CKO* mice to sound stimuli. Since the
vestibular dysfunctions might influence auditory behavioral responses, we first assessed the
vestibular and motor function of *Isl1CKO*. During open-field observations, *Isl1CKO* mice did not
display any signs of abnormal locomotor behavior, such as ataxia, difficulty maintaining balance,
wagging or vertical bobbing movements of the head, or circling movements. The air-righting test,
a basic vestibular function test, and motor function tests on the rotarod demonstrated comparable
motor coordination between control and *Isl1CKO* mice (Supplementary Fig. 8). Concurrently, the
size of the sensory epithelia of the vestibular organs and the size of the dorsal root ganglia were
unaffected in *Isl1CKO* (Supplementary Fig. 8). Thus, locomotor activities of *Isl1CKO* were
comparable to control mice. The acoustic startle response (ASR) is usually used as a behavioral
readout of hearing status mediated by a brainstem circuit linking cochlear root neurons to spinal
motoneurons. The structural basis of the ASR includes cochlear root neurons, CN neurons, the
nucleus of the lateral lemniscus, the caudal pontine reticular nucleus, spinal interneurons, and
spinal motor neurons [62, 63, 64]. Similar to the ABR thresholds, the ASR thresholds of *Isl1CKO*
significantly increased for startle tone stimuli of 8 kHz and BBN, but not for the high-frequency

Figure 9. The acoustic startle reflex (ASR) and prepulse inhibition (PPI) responses are altered in the *Isl1CKO* mutant. **(a)** ASR thresholds for broadband noise (BBN) bursts and tone pips at 8, 16, and 32 kHz in control and *Isl1CKO* mice are shown. Holm-Sidak method multiple comparison *t*-test. *P < 0.05, **P < 0.01. **(b)** Significantly increased ASR latency to BBN is found in *Isl1CKO* compared to control mice; unpaired *t*-test, **P < 0.01. **(c)** Amplitude-intensity ASR functions for BBN stimulation and **(d)** for tone pips of 8 kHz and **(e)** 32 kHz at different dB SPL intensities in control and *Isl1CKO* mice are shown. **(f)** Efficacy of the BBN, **(g)** 8 kHz, and **(h)** 32 kHz tone prepulse intensity on the relative ASR amplitudes are displayed; ASR ratio = 1 corresponds to the ASR amplitude without a prepulse (uninhibited ASR). Two-way ANOVA with Bonferroni post hoc tests. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. All data are expressed as mean ± SEM.

startle tones (Fig. 9a). The peak latency of the ASR to the BBN stimulation at the 110 dB SPL
intensity was prolonged in *Isl1CKO* (Fig. 9b), indicating a slower reaction to the acoustic stimuli.
We found significantly reduced ASR amplitudes for all tested sound stimuli at higher intensities,
showing deteriorated acoustic startle reactivity in *Isl1CKO* mice (Fig. 9c-e).
To further assess complex auditory discrimination behavior, we exposed control and
Isl1CKO mice to a prepulse inhibition (PPI) paradigm, i.e., the inhibition of the ASR induced by
presenting an acoustic stimulus shortly preceding the presentation of an acoustic stimulus, the
startling sound. The circuit mediating a prepulse on the startle reflex involves central structures of
the auditory pathway, including the IC and the auditory cortex [65, 66]. We used either BBN or pure
tone pips of 8 and 32 kHz at increasing intensities as a non-startling acoustic stimulus (prepulse)
that preceded the startle stimulus in a quiet background. The PPI with the prepulse of 32 kHz, the
well-preserved audible frequency in *Isl1CKO*, was comparable between control and mutant mice
(Fig. 9h). Interestingly, the prepulse of 8 kHz resulted in a more prominent inhibition of the startle
response in *Isl1CKO* than in controls (Fig. 9g), despite the significant hearing deficiency at 8 kHz,
as shown by ABR evaluations (Fig. 4). This indicates that the 8-kHz prepulse response was
enhanced in *Isl1CKO*, suggesting compensatory neural hyperactivity of the central auditory system
[56, 67]. Thus, the ASR and PPI of startle analyses indicate abnormalities of the acoustic behavior of
Isl1CKO mutants.
427

Discussion

Our study shows for the first time that LIM homeodomain transcription factor ISL1 regulates
neuronal development in the cochlea. Using RNA profiling, morphological, and physiological
analyses, we provide evidence that ISL1 coordinates genetic networks affecting the molecular
characteristics of SGNs, their pathfinding abilities, and auditory information processing. The
elimination of *Isl1* in neurons during inner ear development results in a migration defect of SGNs,
disorganized innervation in the cochlea, unsegregated and reduced central axons, and reduced size
of the CN. This neuronal phenotype of *Isl1CKO* was accompanied by hearing impairment,
abnormalities in sound processing in the IC, and aberrant auditory behavior.
ISL1 is critical for the development of multiple tissues, neuronal and non-neuronal cells [68,
69, 70, 71, 72]. Different aspects of neuronal development depend on ISL1, including specification of
motoneurons [68], sensory neurons [25, 70], axonal growth [73], and axonal pathfinding [74]. During inner
ear development, ISL1 is expressed in both neuronal and sensory precursors [18, 20, 21, 22, 75].
Transgenic modulations of *Isl1* indicate important roles of ISL1 in the maintenance and function
of neurons and hair cells and as a possible contributing factor in neurodegeneration [23, 76, 77, 78].
These studies suggest that ISL1 plays a role in developing neurons and sensory cells, but no direct
evaluation of ISL1 function has been performed. To circumvent the pleiotropic effects of *Isl1* in
embryonic development, in this study, we used *Neurod1^Cre* to delete *Isl1* specifically in the inner
ear neurons.
We established that ISL1 is necessary for neuronal differentiation programs in the cochlea
and the functional properties of the auditory system. Our RNA profiling of SGNs demonstrated
transcriptome changes induced by a loss of *Isl1* affecting the molecular characteristics of neurons
and pathfinding abilities, including neurotransmission, the structure of synapses, neuron migration,
axonogenesis, and expression of crucial guidance molecules. Consistent with the central role of
ISL1 in sensory neuron developmental programs [25], regulatory networks of signaling molecules
and transcription factors were affected in *Isl1CKO* neurons, such as proneural bHLH factors
(members of NeuroD, Olig, and Nscl families), LIM-only (*Lmo2*, *Lmo3*) and LIM homeodomain

transcription factors (*Lhx1, Lhx2, Isl2*), transcription activation complexes for coordination of particular differentiation programs Eyes absent (*Eya4* and *Eya2*) and Sine oculis (*Six2*), POU homeodomain trans-regulatory factors (*Pou3f2* and *Pou4f2*), and FGF signaling molecules (*Fgf10, Fgf11, Fgf13, Fgf14*) and their downstream targets (*Etv1, Etv4, Etv5*). Interestingly, the transcription factor *Gata3* was downregulated, suggesting that ISL1 is upstream of the Gata3 transcriptional network of neuronal differentiation programs [13]. Thus, ISL1 orchestrates a complex gene regulatory network driving multiple aspects of neuronal differentiation in the cochlea and defining neuronal features.

The most striking morphological features of the neuronal phenotype of *Isl1CKO* are the aberrant migration and pathfinding of SGNs. A similar migration deficit was reported for *ErbB2* null mutants [79]; however, the interpretation of the findings is convoluted by the compounded direct inner ear effects and the effects associated with neural crest-derived Schwann cells. Additionally, conditional deletion of *Sox10* produced by *Wnt1^Cre* resulted in abnormal migration of SGNs similar to the *Isl1CKO* phenotype [80]. Migration defects of SGNs in both *ErbB2* [79] and *Sox10* mutants [80] were attributed to the complete absence of Schwann cells in the entire inner ear ganglion. However, in our *Isl1CKO* mutant, Sox10$^+$ Schwann cells were found in a similar density in the spiral ganglion of both mutant and control mice, thus excluding any direct involvement of glial cells in the migration defects of *Isl1*-deficient neurons. We identified several genes with altered molecular profiles that are associated with neuronal phenotypes of projection and migration abnormalities. For instance, misplaced SGNs in the modiolus are in *Ntrk2* and *Ntrk3* neurotrophin receptor mutants [37]. Less severe phenotypes of misplaced SGNs have been reported for mice lacking *Dcc* [41] or mice with deficient Slit/Robo signaling [40]. All these reported SGN migration defects are also associated with abnormalities in projections. In line with the *Isl1CKO* phenotype of deficient neuronal migration and projections, *Isl1* elimination in SGNs resulted in major transcriptional changes of genes encoding members of all four classes of guidance molecules and their receptors, growth factors, neurotrophic factor receptors, and cell adhesion molecules (NCAM family, integrins, cadherins and protocadherins) among others that are known to participate in molecular networks coordinating neuronal migratory behaviors, axon guidance, and neural circuit assembly.

More profound disorganization of peripheral processes than defects found in *Isl1CKO* or both *ErbB2* null [79] and *Sox10* mutants [80] was reported for delayed conditional deletion of *Gata3* in SGNs [13]. Despite severe disorganization of cochlear wiring of this *Gata3* mutant, central projections maintained their overall tonotopic organization within the auditory nerve and the CN [12, 13]. In contrast, we provide compelling evidence that elimination of *Isl1* in SGNs affected the pathfinding abilities of neurons in the cochlea to form peripheral processes and establish central projections. Such profound disorganization of peripheral and central projections is known for *Neurod1* mutations [16, 18, 19]. Deletions of *Neurod1* result in miswired, and diminished SGNs, and loss of tonotopy [16, 19]. Somewhat similar disorganization of central axons with unsegregated auditory nerve fibers, reduced size of the CN, and missing tonotopic organization of synapsing branches in the CN subdivisions was found in our *Isl1CKO*. Although both *Neurod1CKO* and *Isl1CKO* demonstrated significant hearing loss, in contrast to the reduced sound frequency range of *Neurod1CKO* [19], responses for the entire measured frequency range were detected in *Isl1CKO*. The processing of high acoustic frequencies was broadly comparable between age-matched controls and *Isl1CKO*, indicating some preservation of peripheral neuronal activity. Specifically, high frequencies above 28 kHz were comparable between the control and *Isl1CKO* mice, as shown by ABR measurements and extracellular electrophysiological recordings of IC neuronal activity (Fig. 4b and 8d, e).

As a result of disorganized primary auditory neurons with derailed central projections, the characteristics of persistent auditory function in the IC were altered with worsened tuning capabilities of IC units and their increased spontaneous activity and threshold elevations and decreased dynamic range. The peripheral deficit in sound encoding results in abnormal auditory behavior of *Isl1CKO*. Although no significant differences of ABR thresholds at 32 kHz were observed between *Isl1CKO* and control mice, indicating retained hearing function, the startle reactions of *Isl1CKO* at 32 kHz were reduced. Plasticity of the startle response is also evident in the PPI responses of *Isl1CKO* mice, in which a weak prestimulus suppresses the response to a subsequent startling stimulus. *Isl1CKO* mice demonstrated PPI impairment for the pure tone of 8 kHz, reflecting abnormal sensorimotor gating due to compensatory hyperactivity of the central auditory system [67, 81].

Additionally, compared to control mice, DPOAE responses of *Isl1CKO* were reduced, indicating dysfunction of cochlear amplification. The OHCs of the organ of Corti play a central role in the active enhancement of sound-induced vibration. For given OHCs, amplification only occurs at a precise frequency, and thus, this mechanism provides a sharpening of the tuning curve and improves frequency selectivity [82]. Nevertheless, DPOAE analysis showed that some function was preserved in high-frequency OHCs in the *Isl1CKO* cochlea. Frequencies above 28 kHz are located at the basal half of the mouse cochlea, from the mid-base to the basal end [49], which correspond to the most preserved distribution of sensory neurons in the area of the spiral ganglion of *Isl1CKO* mice (Fig. 1, 5). Usually, decreased DPOAE amplitudes indicate loss and dysfunction of OHCs [76, 82, 83, 84]. Since *Neurod1^Cre* is not expressed in sensory cells in the cochlea, any changes in the development of OHCs represent secondary effects of *Isl1* elimination in neurons. Accordingly, cochlear amplification deficits in *Isl1CKO* correlated with the reduced and disorganized innervation of OHCs (Fig. 2). The medial olivocochlear efferents innervate OHCs from the brainstem, representing a sound-evoked negative feedback loop suppressing OHC activity [85]. The formation of efferent intraganglionic spiral bundle was altered in *Isl1CKO*, indicating changes in efferent innervation in the cochlear region. Besides efferents, OHCs are innervated by the type II SGNs [86, 87]. Although the function of type II SGNs remains obscure, it is clear that these neurons are involved in auditory nociception [88], and may also constitute the disputed sensory drive for the olivocochlear efferent reflex [89, 90]. As *Isl1* is expressed in both type I and type II SGNs during inner ear development [4], characteristics of both neuronal types might likely be affected in *Isl1CKO*.

The unexpected close-to-normal hearing function and auditory signal processing at high frequencies suggest some preservation of tonotopic organization in the *Isl1CKO* cochlea for the propagation of acoustic information. Correspondingly, our 3D images of the *Isl1CKO* cochlea show that some neurons in the base are in their proper position in the Rosenthal's canal, forming a segment of the spiral ganglion. These unexpected results may reflect a timeline of sequential neuronal differentiation from the base to the apex, as the first differentiated neurons are in the base and the last SGNs undergo terminal mitosis in the apex [91]. As the cochlea is extending, differentiating neurons migrate along the cochlea to form the spiral ganglion. This process is completely disrupted in the *Isl1CKO* cochlea, and many neurons are concentrated in a conical shaped central structure, the modiolus, with only a portion of neurons in the Rosenthal's canal. Notably, the functional and spatial organization of SGNs may differ in transcription factor networks required for their differentiation programs. For example, some *Neurod1* lacking neurons survive, form a rudimental cochlear ganglion, and establish bipolar connections to their targets in *Neurod1* null mice [11] or the otocyst-deleted *Neurod1* conditional mutant [17]. Consistent with these

findings, neurons forming a segment of the spiral ganglion in the Rosenthal's canal of *Isl1CKO*
may not require ISL1 for their differentiation and/or the migration mode. Another possibility is
that *Neurod1^Cre* may produce a delayed *Isl1* deletion, resulting in a close to-normal spatial
organization of SGNs and auditory system function at high frequencies. A different Cre driver
generating an earlier recombination of *Isl1* in the otocyst might be used to address the high-
frequency hearing phenotype of *Isl1CKO*.
Another secondary effect of *Isl1* elimination in neurons likely contributing to the hearing
deficit of *Isl1CKO* is a shortened cochlea accompanied by the disorganized apical epithelium.
Somewhat similar phenotypes of a shortened cochlea were previously reported for deletion
mutants of *Neurod1* and *Neurog1*, key transcription factors for inner ear neuronal development [11,
18, 19, 91]. A comparable effect of cochlear length reduction was observed following the loss of *Foxg1*
[92] and *Lmx1a* [93, 94]. Although mechanisms affecting the cochlear extension are unknown, it is clear
that these confounding features of the *Isl1CKO* phenotype would consequently impact mechanical
and neural tuning from the base to the apex of the cochlea and the ability to perform time-frequency
processing of sound.
A limitation of our study is that we only assessed the functional role of *Isl1* in the inner ear
neurons. The expression pattern of ISL1 indicates that ISL1 may control the development of
multiple inner ear progenitors [21]. Notably, the elimination of *Isl1* in developing neurons is only a
part of the story. ISL1 is highly expressed in inner ear neurons throughout the development and in
postnatal auditory neurons (Supplementary Figure 2) [4, 18, 21, 22, 95]. Consistent with our results, the
expression profile indicates an important role of ISL1 for the terminal differentiation of neurons,
and for the function and maintenance of differentiated neurons. In contrast, the expression of ISL1
in sensory precursors is downregulated as hair cell differentiation is initiated and thus, ISL1 is not
detected in the differentiated sensory hair and supporting cells in vestibular and cochlear epithelia
[21]. As the expression of ISL1 precedes the induction of ATOH1 and sensory cell differentiation, it
is conceivable that ISL1 may play a role in the specification of sensory fate and the regulation of
the initial sequential events in sensory precursor development. Further studies will be needed to
fully uncouple regulatory mechanisms in inner ear development by targeted eliminations of *Isl1* in
sensory precursors and in both neuronal and sensory precursor populations.
Altogether, our study provides compelling evidence that ISL1 is a critical regulator of SGN
development, affecting neuronal migration, pathfinding abilities to form cochlear wiring, and
central axonal projections. As such, ISL1 represents an essential factor in the regulation of
neuronal differentiation to produce functional neurons in cell-based therapies and stem cell
engineering [96, 97]. Additionally, this unique model contributes to our understanding of how
disorganization of the neuronal periphery affects information processing at higher centers of the
central auditory pathway at the physiological and behavioral levels.
583
584 ## Methods
585
Experimental animals
All experiments involving animals were performed according to the Guide for the Care and Use
of Laboratory Animals (National Research Council. Washington, DC. The National Academies
Press, 1996). The design of experiments was approved by the Animal Care and Use Committee of
the Institute of Molecular Genetics, Czech Academy of Sciences. The mice were housed in 12-
hour light/dark cycles and were fed *ad libitum*. To generate *Isl1CKO* (the genotype
Neurod1^Cre;Isl1^{loxP/loxP}), we cross-bred floxed *Isl1* (*Isl1^{loxP/loxP}*; *Isl1^{tm2Sev}/J,* # 028501, Jackson

Laboratory)[25] and *Neurod1[Cre]* transgenic mice (Tg(Neurod1-cre)1Able/J, # 028364, Jackson
Laboratory), which were generated by pronuclear injection of the *Neurod1-cre* BAC construct that
carries Cre-sequence downstream of the translational initiation codon ATG of the *Neurod1* gene
[24]. Heterozygous animals, *Neurod1[Cre];Isl1[+/loxP]* were viable, born in appropriate Mendelian ratios,
and were phenotypically indistinguishable from control (Cre negative) littermate mice. As control
mice, we used mice with the genotype Cre negative, *Isl1[loxP/loxP]*, and *Isl1[+/loxP]*. The mouse line
Neurod1[Cre] was also bred with Cre-reporter tdTomato line (*TomatoAi14*, B6.Cg-
Gt(ROSA)26Sor[tm14(CAG-tdTomato)Hze], # 7914 Jackson Laboratory). Using tdTomato reporter for our
analyses, we compared the reporter control-*Ai14* (genotype: *Neurod1[Cre];Isl1[+/loxP];TomatoAi14*)
and reporter *Isl1CKO-Ai14* (genotype: *Neurod1[Cre];Isl1[loxP/loxP];TomatoAi14*). We performed PCR
genotyping on tail DNA. We used both males and females for experiments. Lines are a mixed
C57BL/6/sv129 background. Phenotyping and data analyses were performed blind to the genotype
of the mice.
606
Morphological evaluation of the cochlea, the CN, and the IC
Dissected ears were fixed in 4% paraformaldehyde (PFA) in PBS. For vibratome sections, samples
were embedded in 4% agarose and sectioned at 80 µm using a Leica VT1000S vibratome.
Vibratome sections, whole inner ears, or whole embryos were defatted in 70% ethanol and then
rehydrated and blocked with serum, as described previously [20, 77]. Samples were then incubated
with primary antibodies at 4ºC for 72 hours. The primary antibodies used were: rabbit anti-Myosin
7a (Myo7a; Proteus BioSciences 25-6790, 1:500), mouse anti-acetylated α-tubulin (tubulin;
Sigma-Aldrich T6793, 1:400), rabbit anti-calretinin (Santa Cruz Biotechnology sc-50453, 1:100),
goat anti-prestin (Santa Cruz Biotechnology sc-22692, 1:50), rabbit anti-parvalbumin (Abcam
ab11427, 1:2000), mouse anti-VGLUT1 (Merck MAB5502, 1:200), rabbit anti-NeuN (Abcam
ab177487, 1:500), mouse anti-Isl1 (Developmental Hybridoma Bank 39.3F7 or 39.4D5, 1:130),
mouse anti-C-terminal binding protein 2 (CtBP2; BD Biosciences 612044, 1:200), rabbit anti-
cleaved Caspase-3 (Cell Signaling Technology 9661, 1:100), goat anti-Neurod1 (Santa Cruz
Biotechnology sc-1084, 1:100), anti-β tubulin III (Tuj1; BioLegend 801202, 1:500), and rabbit
anti-Sox10 (Abcam ab155279, 1:250). After several PBS washes, secondary antibodies were
added and incubated at 4ºC for 24 hours. The secondary antibodies Alexa Fluor® 488 AffiniPure
Goat Anti-Mouse IgG (Jackson ImmunoResearch Laboratories 115-545-146), Alexa Fluor® 594
AffiniPure Goat Anti-Rabbit IgG (Jackson ImmunoResearch Laboratories 111-585-144),
DyLight488-conjugated AffiniPure Mouse Anti-Goat IgG (Jackson ImmunoResearch
Laboratories 205-485-108), Alexa Fluor® 647-conjugated AffiniPure Donkey Anti-Goat IgG
(Jackson ImmunoResearch Laboratories 705-605-147), Alexa Fluor® 594-conjugated AffiniPure
Donkey Anti-Rabbit IgG (Jackson ImmunoResearch Laboratories 711-585-152), and Alexa
Fluor®488-conjugated AffiniPure Donkey Anti-Mouse IgG (Jackson ImmunoResearch
Laboratories 715-545-150) were used in 1:400 dilution. Nuclei were stained by Hoechst 33258
(Sigma-Aldrich 861405, 1:2000). Samples were mounted in Aqua-Poly/Mount (Polysciences
18606) or prepared Antifade medium. Images were taken on Zeiss LSM 880 NLO inverted
confocal microscope, Nikon CSU-W1 spinning disk confocal microscope, and Carl Zeiss
AxioZoomV16 macroscope. NIS-Elements, ImageJ, and ZEN software were used for image
processing.
The length of the organ of Corti was measured using the "Measure line" ImageJ plugin.
The volumes of the CN and IC were established by analyzing parallel, serial, equally spaced 80
µm coronal vibratome sections through the brain (n = 5 *Isl1CKO* and n = 5 control mice). The CN

and IC areas were determined in each section using ImageJ, and the volume of the organs was calculated. Volumes of organs were adjusted to the body weight. Whole-mount anti-tubulin labeling of the cochlea was used to measure the length and density of the radial fibers. The evaluation of the innervation was done separately for each region of the cochlea: the apex, mid-apex, mid-base, and base. Due to the disorganization of innervation in the apex, we only evaluated the cochlear base, mid-base, and mid-apex fiber density. The density of the radial fibers was expressed as the percentage of a positive area in the measured area of 152 x 66 μm^2 using the "Threshold" function of ImageJ. The length of the radial fibers was measured from the whole-mount anti-tubulin and Myo7a immunolabeled cochlea confocal images. For each genotype, we measured 3 samples, and in all four regions of the cochlea (the apex, mid-apex, mid-base, and base), we measured the length of the fibers in 3 radial fiber bundles from the IHCs to the IGSB. To compare how many neurons are correctly located in the Rosenthal's canal in *Isl1CKO*, we used NeuN immunolabeled whole-mount cochlea (n = 4 per genotype). Cells were manually counted in 40x magnification images in 5 Z-stacks using the "Cell counter" function of ImageJ; the area of the Rosenthal's canal with tdTomato$^+$ neurons was measured using "Threshold", "Polygon selection" and "Measure" functions of ImageJ to calculate the relative number of neurons for the control and *Isl1CKO* per Rosenthal's canal volume. For ribbon evaluations, IHCs were counted based on the calretinin-stained cell bodies and anti-CtBP2 labeled synapses. CtBP2$^+$ presynaptic ribbon structures were manually counted in 10 individual IHCs in each cochlear region: the apex, mid-apex, mid-base, and base. A total of 6 mutants and 6 control cochleae were evaluated except for 1 apex in the control group, which was lost during immunolabeling. The apoptosis of neurons was quantified using immunohistochemistry with anti-cleaved Caspase-3. Caspase-3 positive cells were manually counted in the inner ear ganglia labeled by the combination of anti-Cre and anti-Neurod1 in the whole-mounted head at 60x magnification at E10.5. For E12.5, Caspase-3$^+$/tdTomato$^+$ neurons were manually counted in the whole-mounted inner ear at 60x magnification. The area of E12.5 inner ear ganglia was measured using the "Threshold and polygon selection" function of ImageJ. For E14.5, the cochlea was sectioned, and Caspase-3$^+$/tdTomato$^+$ neurons were manually counted in all sections at 60x magnification. Three embryos per genotype were used for each embryonic day. All apoptotic cell counting was done using the "Cell counter" function of ImageJ. The density of spherical bushy cells in the VCN, 3 control and 3 *ISL1CKO* 2-month-old mice were used. Bushy cells were labeled by anti-parvalbumin in 80 μm vibratome sections, and manually counted in 2-4 sections/animal in 134.95 x 134.95 μm (63x objective). The size of vestibular sensory epithelia was measured in anti-Myo7a labeled wholemount preparations at P0. Myo7a$^+$ areas were measured in 5 control and 5 *Isl1CKO* mice using "Measure" ImageJ tool. The dorsal root ganglia (DRG) size was evaluated in the sagittal sections of E14.5 embryos with expression of tdTomato reporter. The area of DRG was quantified in 6 embryos per genotype and 2 sections per embryo using the "Threshold and polygon selection" function of ImageJ.

Light-sheet fluorescent microscopy (LFSM) and analysis of images
The cochleae were microdissected from 2 control-*Ai14* and 2 *Isl1CKO-Ai14* mice (postnatal day P0). We used an advanced CUBIC protocol[98] for tissue clearing to enable efficient imaging by light-sheet microscopy. Briefly, the microdissected cochlea was fixed in 4% PFA for 1 hrs, washed with PBS, and incubated in a clearing solution Cubic 1 for 7 days at 37°C. Before immunolabeling, samples were washed in PBT (0.5% Triton-X in PBS) 4 x for 30 min. In addition to *tdTomato* expression, cochlear samples were either immunolabeled using anti-NeuN (a nuclear marker of

differentiated neurons) or anti-Myo7a (a hair cell marker) antibodies. Samples were stored before
imaging in Cubic 2 at room temperature. Zeiss Lightsheet Z.1 microscope with illumination
objective Lightsheet Z.1 5x/0.1 and detection objective Dry objective Lightsheet Z.1 5x/0.16 was
used for imaging at the Light Microscopy Core Facility of the Institute of Molecular Genetics of
the Czech Academy of Sciences. IMARIS software v8.1.1 (Bitplane AG, CA, USA) was used for
image processing.
691
Isolation of genetically labeled neurons and library construction
Spiral ganglia were micro-dissected in Dulbecco's PBS on ice from E14.5 embryos of either sex
from four litters; each sample contains spiral ganglia from both inner ears of the individual embryo.
Spiral ganglia were incubated in 300 µl of lysis solution (0.05% trypsin, 0.53mM EDTA in
Dulbecco's PBS) in 37 ºC, shaking at 900 RPM for 5 min. The lysis was stopped by adding 600 µl
of FACS buffer (10mM EGTA, and 2% FBS in Dulbecco's PBS). After spinning down the samples
at 800 G, 4 ºC for 10 min, the supernatant was removed, and cell pellets were resuspended in 500
µl of ice-cold FACS buffer. Immediately before sorting, cells were passed through a 50 µm cell
sieve (CellTrics^{TM}, Sysmex Amercia Inc.) into a sterile 5 ml polystyrene round-bottom falcon to
remove clusters of cells and kept on ice. TdTomato$^+$ neurons were sorted using a flow cytometer
(BD FACSAria^{TM} Fusion), through a 100 µm nozzle in 20 psi, operated with BD FACSDiva™
Software (Supplementary Fig. 9). 100 sorted cells were collected into individual wells of 96-well
plate containing 5 µl of lysis buffer of NEB Next single-cell low input RNA library prep kit for
Illumina (New England Biolabs #E6420). Plates were frozen immediately on dry ice and stored at
−80 °C. The total time from euthanasia to cell collection was ~3 hrs.
RNAseq-libraries were prepared from 6 samples per each genotype, reporter control-*Ai14*
and *Isl1CKO-Ai14* mutant, and each sample contained 100 tdTomato$^+$ neurons. Following the
manufacturer's instructions, the NEB Next single-cell low input RNA library prep kit for
Illumina was used for cDNA synthesis, amplification, and library generation [99] at the Gene Core
Facility (Institute of Biotechnology CAS, Czechia). Fragment Analyzer assessed the quality of
cDNA libraries. The libraries were sequenced on an Illumina NextSeq 500 next-generation
sequencer. NextSeq 500/550 High Output kit 75 cycles (Illumina #200024906) were processed at
the Genomics and Bioinformatics Core Facility (Institute of Molecular Genetics CAS, Czechia).
715
Computational analysis of RNAseq data
RNA-Seq reads in FASTQ files were mapped to the mouse genome using STAR [version 2.7.0c
[100]] GRCm38 primary assembly and annotation version M8. The raw data of RNA sequencing
were processed with a standard pipeline. Using cutadapt v1.18[101], the number of reads
(minimum, 32 million; maximum, 73 million) was trimmed by Illumina sequencing adaptor and
of bases with reading quality lower than 20, subsequently reads shorter than 20 bp were filtered
out. TrimmomaticPE version 0.36 [102]. Ribosomal RNA and reads mapping to UniVec database
were filtered out using bowtie v1.2.2. with parameters -S -n 1 and SortMeRNA [103]. A count table
was generated by Rsubread v2.0.1 package using default parameters without counting multi
mapping reads. The raw RNAseq data were deposited at GEO: # GSE182575 study
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE182575). DESeq2 [v1.26.0 [104]] default
parameters were used to normalize data and compare the different groups. Differentially
expressed genes were identified based on an adjusted P-value $p_{adj} < 0.05$, FC > 2, and a base
mean ≥ 50 was applied to identify differentially expressed genes between *Isl1CKO* mutant and

control neurons. The functional annotation of the differentially expressed genes was performed
using GOTermFinder [105].
Enrichment mapping. The enrichment of the functional categories and functional annotation
clustering of the differentially expressed genes was performed using g: Profiler [106] input using
version e104_eg51_p15_3922dba with g: SCS multiple testing correction methods applying a
significance threshold of 0.05. In contrast, no electronic GO annotations were used. Only
Biological Processes (BP) data underwent further processing. Complete query details are available
in Query info tabs in Supplementary Data 2. The resulting GEM and combined GMT files were
loaded into Cytoscape [107] plugin "EnrichmentMap" [108] using 0.01 FDR q-value cutoff to generate
a network. Edge cutoff was set to 0.6, and nodes were filtered by gs_size<1800. Five GO terms
forming solitary nodes, or a pair of nodes, were excluded (listed in Supplementary Data 2). Further
adjustments were made in yFiles Layout Algorithms, Legend Creator (Cytoscape plugins), and
Inkscape (Inkscape Project, 2020).
743
Quantitative real-time PCR
Total RNA was isolated from both inner ears of the embryo at E14.5 using TRI Reagent (Sigma-
Aldrich T9424). We used 8 embryos for *Isl1CKO* and 7 embryos for the control group from three
litters. RNA from both inner ears of one embryo represented one sample. RNA samples (0.5 μg of
total RNA) were processed and analyzed as previously described [18, 77]. Briefly, following RT,
quantitative qPCR was performed with initial activation at 95 °C for 120 s, followed by 40 cycles
at 95 °C for 15 s, 60 °C for 30 s, and 72 °C for 30 s using the CFX384™ Real-Time PCR Detection
System (Bio-Rad Laboratories). The primer sequences (*Lhx1, Lhx2, Cdh7, Nhlh2, Ntrk2, Ntrk3,
Ntng1*) were obtained from (pga.mgh.harvard.edu/primerbank/) or designed using Primer3 tool
(*Dcc, Epha5, Gata3, Prdm8, Robo2, Slit2, Tbx3, Uncx*). The primer sequences are listed in
Supplementary Table 2. Relative mRNA expression was calculated using the $-\Delta\Delta Cq$ method with
Hprt1 as a reference gene. GraphPad Prism software was used for statistical analysis.
756
Lipophilic Dye Tracing
We studied the innervation pattern in whole or dissected ears using lipophilic dye tracing in
aldehyde-fixed tissues as previously described [109]. At least three mutants and similar numbers of
control littermates of both sexes were used for each evaluation. Filter strips loaded with NeuroVue
colored lipophilic dyes were inserted into the cochlear apex, base, and vestibular end-organ utricle.
After allowing appropriate time for diffusion of the lipophilic tracer (between 48-120 hours), we
prepared the ears as whole mounts in glycerol on a glass slide, using appropriate spacers to avoid
distortion, and imaged them using a Leica SP8 confocal microscope. Images were compiled into
plates to show the most pertinent details using Corel Draw. Only general image modifications such
as contrast, or brightness adjustments were used to enhance the visual appeal without affecting the
scientific content.
768
Hearing function evaluation
Auditory brainstem response (ABR) and distortion product otoacoustic emissions (DPOAEs) tests
were carried out on mice under general anesthesia with 35 mg/kg ketamine (Calypso 50 mg/ml)
and 6 mg/kg xylazine (Xylapan 20 mg/ml) in saline to give an application volume of 7 ml/kg body
weight via subcutaneous injection, maintained on a temperature-regulated blanket in a soundproof
room.

Distortion product otoacoustic emissions. For DPOAE recording were tested *Isl1CKO* (n = 10)
and control mice (n = 14). Cubic (2 F1–F2) distortion product otoacoustic emissions over an F2
frequency range from 4 to 39 kHz were recorded with a low-noise microphone system (Etymotic
probe ER-10B+, Etymotic Research). Acoustic stimuli (ratio F2/F1 = 1.21, F1 and F2 primary tone
levels of L1/L2 = 70/60 dB) were presented to the ear canal with two custom-made piezoelectric
stimulators connected to the probe with 10-cm-long silastic tubes. The signal from the microphone
was analyzed by the TDT System III (RP2 processor, sampling rate 100 kHz) using custom-made
MATLAB software. DPOAEs were recorded in the animals' ears at individual frequencies over
the frequency range 4–39 kHz with a resolution of ten points per octave. All the experiments and
analyses were done with no information on the genotype.
Auditory brainstem response. For auditory brainstem response (ABR) recording (n = 10 *Isl1CKO*
and n = 11 control mice), an active electrode was placed subcutaneously on the vertex and ground
and reference electrodes in the neck muscles. Responses to tone bursts (3 ms duration, 1 ms rise/fall
times, frequencies of 2, 4, 8, 16, 32, and 40 kHz) and clicks of different intensity were recorded.
Acoustic stimuli were conveyed to the animal in free-field conditions via a two-way loudspeaker
system (Selenium 6W4P woofer and RAAL70-20 tweeter) placed 70 cm in front of the animal's
head. The signal was processed with a TDT System III Pentusa Base Station and analyzed using
BioSig™ software. The response threshold to each frequency was determined as the minimal tone
intensity that still evoked a noticeable potential peak in the expected time window of the recorded
signal. The amplitude and latency of ABR peaks I-V were determined using BioSig software
(Tucker Davis Technologies). Central compensation of neuronal responsiveness (central gain) was
calculated using ABR wave IV to I amplitudes.
797
Extracellular recording of the neuronal activity in the IC
For extracellular recording in the IC, we evaluated *Isl1CKO* (n = 12, 624 units) and control mice
(n = 8, 395 units). The surgery and extracellular recording in the IC were performed in mice
anesthetized with 35 mg/kg ketamine (Calypsol 50 mg/ml) and 6 mg/kg xylazine (Xylapan 20
mg/ml) in saline via subcutaneous injection. Approximately every hour, supplement subcutaneous
injections of one-half of the original dose of the anesthetics were administered to keep a sufficient
level of anesthesia, judged by a positive pedal and palpebral (toe-pinch) reflex and movement of
the whiskers. Respiratory rate, and heart rate, were monitored. An incision was made through the
skull's skin to access the IC, and underlying muscles were retracted to expose the dorsal skull. A
holder was glued to the skull, and small holes were drilled over both ICs. Neuronal activity
(multiple units) in the IC was recorded using a 16-channel, single shank probe (NeuroNexus
Technologies) with 50 or 100 μm between the electrode spots. The obtained signal from the
electrode was amplified 10000 times, band-pass filtered over the range of 300 Hz to 10 kHz, and
processed by a TDT System III (Tucker Davis Technologies) using an RX5-2 Pentusa Base
Station. Individual spikes from the recorded signal were isolated online based on amplitude
discrimination and analyzed with BrainWare software (v. 8.12, Jan Schnupp, Oxford University).
Subsequent discrimination of spikes from the recorded data and their sorting according to the
amplitudes of the first positive and negative peaks were performed off-line and was used to sort
action potentials (spikes) among single units. The recorded data were processed and analyzed using
custom software based on MATLAB. The stimulation signals were generated using a TDT System
III with the RP 2.1 Enhanced Real-Time Processor. Acoustic stimuli were delivered in free-field
conditions via a two-driver loudspeaker system (Selenium 6W4P woofer and RAAL70-20 tweeter)
placed 70 cm in front of the animal's head.

Frequency-intensity mapping: To determine the neuronal receptive fields, pure tones (frequency 2
- 40 kHz with 1/8 octave step, 60 ms duration, 5 ms rise/fall times, various intensity with 5dB step)
were presented in a random order, each stimulus appearing three times. A discrete matrix
corresponding to the response magnitude evoked by each of the frequency-intensity combinations
was thereby obtained, smoothed using cubic spline interpolation, and used for extraction of the
basic parameters: the excitatory response threshold (the lowest stimulus intensity that excited the
neuron, measured in dB SPL), the characteristic frequency (CF) – the frequency with the minimal
response threshold, measured in Hz, and the bandwidth of the excitatory area 20 dB above the
excitatory threshold, expressed by quality factor Q (Q = CF/bandwidth).
Rate intensity function of the IC neurons: Neuronal responses to broadband noise (BBN) bursts of
variable intensity (10 dB steps, 50 repetitions) were used to construct the rate intensity function
(RIF). A 100% scale was assigned to the neuron's total range of response amplitudes, 0%
corresponding to spontaneous activity and 100% corresponding to its maximum response
magnitude [110]. The two points of interest are R10 and R90, which correspond to 10 and 90% of
this scale, respectively. R10, describing the starting point of the RIF's rise, was taken as the BBN
response threshold. RIFs were further used to evaluate the following parameters: the dynamic
range (DR) of the RIF: DR = S90–S10); and the maximum response magnitude. Spontaneous
activity of the IC neurons was determined at the 0dB SPL BBN stimulation.
Temporal properties of the IC neurons: We used trains of five clicks at an intensity of 70 dB SPL
for control and 80 dB SPL for *Isl1CKO* mice with various inter-click intervals (100, 50, 30, 20,
and 15 ms). We calculated the vector strength (VS) values and the Rayleigh statistics for each
spike pattern; only responses with at least 5.991 were significantly considered phase-locking (Zhou
and Merzenich, 2008). The VS quantifies how well the individual spikes are synchronized (phase-
locked) with a periodic signal.
845
Behavioral tests of motor coordination and balance
All testing was carried out during the light cycle and after a minimum 30-min acclimatization. *Air-*
righting reflex. The vestibular function was evaluated by the ability of the mice to right themselves
in the air when held supine and dropped onto a soft surface from a height of 50 cm[77].
Rotarod. Using the rotarod apparatus (Rota Rod 47600, Ugo Basile), the time (latency) to fall off
the rod rotating under continuous acceleration was recorded. During the acclimatization period,
mice with their heads in the direction of rotation were loaded on the rotarod at an initial speed of
4 rpm. This speed was maintained for 2 min and, if mice fell during this period, they were placed
back on the rotarod. The drum was slowly accelerated from 4 to 40 rpm for a maximum of 300 s
for each trial for the experimental measurements. The latency to fall off the rotarod within 300 s
was recorded. If the mouse clung to the grip of the rotating rod and failed to resume normal
performance for three consecutive revolutions, the sensor was manually triggered. Mice were
tested in three consecutive trials in one session per day with a 15-min rest period between each
trial.
860
Behavioral acoustic tests
Isl1CKO (n = 10) and control (n = 7) mice were used at 2-3 months of age. All behavioral tests
were performed in a sound-attenuated chamber (Coulbourn Habitest, model E10-21) located in a
soundproof room. Each mouse was placed in a wire mesh cage on a motion-sensitive platform
inside the box during the testing. The mouse's reflex movements to sound stimuli were detected
and transformed to a voltage signal by the load-cell response sensing platform. An amplified

voltage signal was acquired and processed using a TDT system 3 with a Real-Time Processor RP 2 (Tucker Davis Technologies, Alachua, Fl) and custom-made software in a Matlab environment. The startle responses were evaluated in 100 ms windows, beginning at the onset of the startling stimulus. The magnitude of the response was measured as the maximal peak-to-peak amplitude of transient voltage occurring in the response window. Acoustic stimuli were generated by the TDT system (Real-Time Processor RP 2), amplified and presented via a loudspeaker (SEAS, 29AF/W), and placed inside the chamber above the animal. Stimulus presentation and data acquisition were controlled by a custom-made application in a Matlab environment. Calibration of the apparatus was performed for frequencies between 4 kHz and 32 kHz by a 1/4-inch Brüel & Kjaer 4939 microphone connected to a Brüel & Kjaer ZC 0020 preamplifier and a B&K 2231 sound level meter. During the calibration, the calibrating microphone was positioned at the animal's head in the test cage.

Acoustic startle reflex (ASR) (a transient motor response to an intense, unexpected stimulus) was used to indicate the behavioral responsiveness to sound stimuli**.** The ASRs to 8, 16, and 32 kHz tone pips and BBN bursts (50 ms duration, 3 ms rise/fall times, varying intensity levels) were recorded. Each test session contained: a baseline trial (–10 dB SPL stimulus intensity) and 13 startle stimuli of different intensities (50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 115, and 120-dB SPL). The inter-trial interval varied from 15 to 50 s.

In *the prepulse inhibition (PPI)* procedure, 3 different trial types were used: a baseline trial without any stimulus, an acoustic startle pulse alone (white noise at 110 dB SPL, 50 ms, 3 ms rise/fall times), and a combination of the prepulse and startle pulse. The inter-stimulus interval between the prepulse and the startle stimulus was set to 50 ms; each trial type was presented three times. The inter-trial gap was randomized and varied from 15 to 50 s. The efficacy of the PPI of ASR was expressed as an ASR ratio in percentage, e.g., 100% corresponds to the amplitude of ASR without prepulse; smaller values of ASR ratio indicate stronger PPI. As a prepulse, we used either BBN bursts or tone pips (50 ms duration, 3 ms rise/fall time) at frequencies of 8 and 32 kHz at increasing intensities. It is expected that in the presence of prepulse, the amplitude of the following startle response decreases.

Experimental design and statistical analyses
All comparisons were made between animals with the same genetic background, typically littermates, and we used male and female mice. The number of samples (n) for each comparison can be found in the individual method descriptions and are given in the corresponding figures. Phenotyping and data analysis was performed blind to the genotype of the mice. All values are presented either as the mean ± standard deviation (SD) or standard error of the mean (SEM). For statistical analysis, GraphPad Prism software was used. To assess differences in the mean, one-way or two-way ANOVA with Bonferroni's multiple comparison test, multiple *t*-tests with Holm-Sidak comparison method, repeated measures ANOVA, multiple *t*-tests, and unpaired two-tailed *t*-tests were employed. Significance was determined as $P < 0.05$ (*), $P < 0.01$ (**), $P < 0.001$ (***) or $P < 0.0001$ (****). Complete results of the statistical analyses are included in the figure legends.

Ethics approval and consent to participate
Experiments were carried out following the animal welfare guidelines 2010/63/EC of the European Communities Council Directive, agreeing with the Guide for the Care and Use of Laboratory Animals (National Research Council. Washington, DC. The National Academies Press, 1996).

912 The design of experiments was approved by the Animal Care and Use Committee of the Institute
913 of Molecular Genetics, Czech Academy of Sciences (protocol # 104/2019).
914

915 Consent for publication
916 Not applicable.
917

918 Data availability
919 All data generated or analyzed during this study are included in this published article, and
920 Supplementary information. RNA-seq data are deposited in the publicly available Gene
921 Expression Omnibus (GEO) repository (# GSE182575 study). Source data are provided with this
922 paper.
923

924 Competing interests
925 The authors declare that they have no competing interests.
926

Funding
This research was supported by the Czech Science Foundation (20-06927S to GP), by the
institutional support of the Czech Academy of Sciences (RVO: 86652036 to GP), and NIH/NIA
(R01 AG060504, DC016099, AG051443 to BF and ENY).

Authors' contributions
GP, JS, BF designed and supervised the experiments. IF, KP, RB, MT, SV, MD, and BF performed
experiments and analyzed the data. JC, PH, and VF prepared samples and generated 3D images.
OS, SB, and LV carried RNAseq analyses. IF and KP prepared the first draft of the manuscript.
GP and BF wrote the manuscript, ENY and JS reviewed the manuscript. All authors read and
approved the final manuscript.
938

Acknowledgments
A. Pavlinek (King's College London) for editing the MS. We thank M. Anderova for providing
TomatoAi14 reporter line. We acknowledge Imaging Methods Core Facility at BIOCEV
supported by the MEYS CR (Large RI Project LM2018129 Czech-BioImaging) and ERDF
(project No. CZ.02.1.01/0.0/0.0/18_046/0016045) for its support with obtaining imaging data
and FACS experiments presented in this paper; Biocev GeneCore Facility for its support with
gene expression/transcriptome analyses, and Biocev Animal facility (LM 2018126 Czech Centre
for Phenogenomics by MEYS OP RDE CZ.02.1.01/0.0/0.0/18_046/0015861 CCP Infrastructure
Upgrade II by MEYS and ESIF).

References
1.   Muniak MA, Connelly CJ, Suthakar K, Milinkeviciute G, Ayeni FE, Ryugo DK. Central
Projections of Spiral Ganglion Neurons. In: *The Primary Auditory Neurons of the Mammalian*
Cochlea). Springer (2016).
953

954 2.   Rubel EW, Fritzsch B. Auditory system development: primary auditory neurons and their targets.
955   *Annu Rev Neurosci* **25**, 51-101 (2002).
956

957   3.   Shrestha BR, Chia C, Wu L, Kujawa SG, Liberman MC, Goodrich LV. Sensory Neuron Diversity
958   in the Inner Ear Is Shaped by Activity. *Cell* **174**, 1229-1246 e1217 (2018).

959
960   4.   Petitpre C*, et al.* Neuronal heterogeneity and stereotyped connectivity in the auditory afferent
961   system. *Nature communications* **9**, 3691 (2018).

962
963   5.   Sun S*, et al.* Hair Cell Mechanotransduction Regulates Spontaneous Activity and Spiral Ganglion
964   Subtype Specification in the Auditory System. *Cell* **174**, 1247-1263 e1215 (2018).

965
966   6.   Kandler K, Clause A, Noh J. Tonotopic reorganization of developing auditory brainstem circuits.
967   *Nat Neurosci* **12**, 711-717 (2009).

968
969   7.   Di Bonito M, Studer M. Cellular and Molecular Underpinnings of Neuronal Assembly in the
970   Central Auditory System during Mouse Development. *Front Neural Circuits* **11**, 18 (2017).

971
972   8.   Lopez-Poveda EA. Olivocochlear Efferents in Animals and Humans: From Anatomy to Clinical
973   Relevance. *Front Neurol* **9**, 197 (2018).

974
975   9.   Fritzsch B, Elliott KL. Evolution and Development of the Inner Ear Efferent System:
976   Transforming a Motor Neuron Population to Connect to the Most Unusual Motor Protein via
977   Ancient Nicotinic Receptors. *Front Cell Neurosci* **11**, 114 (2017).

978
979   10.   Ma Q, Anderson DJ, Fritzsch B. Neurogenin 1 null mutant ears develop fewer, morphologically
980   normal hair cells in smaller sensory epithelia devoid of innervation. *J Assoc Res Otolaryngol* **1**,
981   129-143 (2000).

982
983   11.   Kim W-Y*, et al.* NeuroD-null mice are deaf due to a severe loss of the inner ear sensory neurons
984   during development. *Development* **128**, 417-426 (2001).

985
986   12.   Duncan JS, Fritzsch B. Continued expression of GATA3 is necessary for cochlear neurosensory
987   development. *PloS one* **8**, e62046 (2013).

988
989   13.   Appler JM, Lu CC, Druckenbrod NR, Yu WM, Koundakjian EJ, Goodrich LV. Gata3 is a critical
990   regulator of cochlear wiring. *J Neurosci* **33**, 3679-3691 (2013).

991
992   14.   Huang EJ, Liu W, Fritzsch B, Bianchi LM, Reichardt LF, Xiang M. Brn3a is a transcriptional
993   regulator of soma size, target field innervation and axon pathfinding of inner ear sensory neurons.
994   *Development* **128**, 2421-2432 (2001).

995
996   15.   Liu M*, et al.* Essential role of BETA2/NeuroD1 in development of the vestibular and auditory
997   systems. *Genes & development* **14**, 2839-2854 (2000).

999   16.   Jahan I, Kersigo J, Pan N, Fritzsch B. Neurod1 regulates survival and formation of connections in
1000   mouse ear and brain. *Cell Tissue Res* **341**, 95-110 (2010).

1001
1002   17.   Filova I, Bohuslavova R, Tavakoli M, Yamoah EN, Fritzsch B, Pavlinkova G. Early Deletion of
1003   Neurod1 Alters Neuronal Lineage Potential and Diminishes Neurogenesis in the Inner Ear.
1004   *Frontiers in Cell and Developmental Biology* **10**, (2022).

1005
1006   18.   Filova I*, et al.* Combined Atoh1 and Neurod1 Deletion Reveals Autonomous Growth of Auditory
1007   Nerve Fibers. *Mol Neurobiol* **57**, 5307-5323 (2020).

1008
1009   19.   Macova I*, et al.* Neurod1 Is Essential for the Primary Tonotopic Organization and Related
1010   Auditory Information Processing in the Midbrain. *J Neurosci* **39**, 984-1004 (2019).

1011
1012   20.   Dvorakova M*, et al.* Incomplete and delayed Sox2 deletion defines residual ear neurosensory
1013   development and maintenance. *Sci Rep* **6**, 38253 (2016).

1014
1015   21.   Radde-Gallwitz K, Pan L, Gan L, Lin X, Segil N, Chen P. Expression of Islet1 marks the sensory
1016   and neuronal lineages in the mammalian inner ear. *J Comp Neurol* **477**, 412-421 (2004).

1017
1018   22.   Deng M, Yang H, Xie X, Liang G, Gan L. Comparative expression analysis of POU4F1,
1019   POU4F2 and ISL1 in developing mouse cochleovestibular ganglion neurons. *Gene Expr Patterns*
1020   **15**, 31-37 (2014).

1021
1022   23.   Chumak T*, et al.* Overexpression of Isl1 under the Pax2 Promoter, Leads to Impaired Sound
1023   Processing and Increased Inhibition in the Inferior Colliculus. *Int J Mol Sci* **22**, (2021).

1024
1025   24.   Li HJ, Kapoor A, Giel-Moloney M, Rindi G, Leiter AB. Notch signaling differentially regulates
1026   the cell fate of early endocrine precursor cells and their maturing descendants in the mouse
1027   pancreas and intestine. *Dev Biol* **371**, 156-169 (2012).

1028
1029   25.   Sun Y, Dykes IM, Liang X, Eng SR, Evans SM, Turner EE. A central role for Islet1 in sensory
1030   neuron development linking sensory and spinal gene regulatory programs. *Nat Neurosci* **11**,
1031   1283-1293 (2008).

1032
1033   26.   Jahan I, Pan N, Kersigo J, Fritzsch B. Neurod1 suppresses hair cell differentiation in ear ganglia
1034   and regulates hair cell subtype development in the cochlea. *PLoS One* **5**, e11661 (2010).

1035
1036   27.   Li C*, et al.* Comprehensive transcriptome analysis of cochlear spiral ganglion neurons at multiple
1037   ages. *Elife* **9**, (2020).

1038
1039   28.   Stoeckli ET. Understanding axon guidance: are we nearly there yet? *Development* **145**, (2018).

1040

1041 29. Fritzsch B, Kersigo J, Yang T, Jahan I, Pan N. Neurotrophic Factor Function During Ear
1042 Development: Expression Changes Define Critical Phases for Neuronal Viability. In: *The*
1043 *Primary Auditory Neurons of the Mammalian Cochlea* (eds Dabdoub A, Fritzsch B, Popper AN,
1044 Fay RR). Springer New York (2016).

1045
1046 30. Dennis DJ, Han S, Schuurmans C. bHLH transcription factors in neural development, disease,
1047 and reprogramming. *Brain Res* **1705**, 48-65 (2019).

1048
1049 31. Yang T, Kersigo J, Wu S, Fritzsch B, Bassuk AG. Prickle1 regulates neurite outgrowth of apical
1050 spiral ganglion neurons but not hair cell polarity in the murine cochlea. *PLoS One* **12**, e0183773
1051 (2017).

1052
1053 32. Bok J, Zenczak C, Hwang CH, Wu DK. Auditory ganglion source of Sonic hedgehog regulates
1054 timing of cell cycle exit and differentiation of mammalian cochlear hair cells. *Proc Natl Acad Sci*
1055 *U S A* **110**, 13869-13874 (2013).

1056
1057 33. Pauley S, Wright TJ, Pirvola U, Ornitz D, Beisel K, Fritzsch B. Expression and function of
1058 FGF10 in mammalian inner ear development. *Dev Dyn* **227**, 203-215 (2003).

1059
1060 34. Yu Y*, et al.* Sensorineural Hearing Loss and Mitochondrial Apoptosis of Cochlear Spiral
1061 Ganglion Neurons in Fibroblast Growth Factor 13 Knockout Mice. *Frontiers in Cellular*
1062 *Neuroscience* **15**, (2021).

1063
1064 35. Defourny J*, et al.* Ephrin-A5/EphA4 signalling controls specific afferent targeting to cochlear hair
1065 cells. *Nature communications* **4**, 1438 (2013).

1066
1067 36. Coate TM, Raft S, Zhao X, Ryan AK, Crenshaw EB, 3rd, Kelley MW. Otic mesenchyme cells
1068 regulate spiral ganglion axon fasciculation through a Pou3f4/EphA4 signaling pathway. *Neuron*
1069 **73**, 49-63 (2012).

1070
1071 37. Fritzsch B, Barbacid M, Silos-Santiago I. The combined effects of trkB and trkC mutations on the
1072 innervation of the inner ear. *Int J Dev Neurosci* **16**, 493-505 (1998).

1073
1074 38. Brors D*, et al.* Spiral ganglion outgrowth and hearing development in p75-deficient mice. *Audiol*
1075 *Neurootol* **13**, 388-395 (2008).

1076
1077 39. Coate TM, Spita NA, Zhang KD, Isgrig KT, Kelley MW. Neuropilin-2/Semaphorin-3F-mediated
1078 repulsion promotes inner hair cell innervation by spiral ganglion neurons. *Elife* **4**, (2015).

1079
1080 40. Wang SZ*, et al.* Slit/Robo signaling mediates spatial positioning of spiral ganglion neurons
1081 during development of cochlear innervation. *J Neurosci* **33**, 12242-12254 (2013).

1082

1083   41.   Kim YJ, Wang SZ, Tymanskyj S, Ma L, Tao HW, Zhang LI. Dcc Mediates Functional Assembly
1084   of Peripheral Auditory Circuits. *Sci Rep* **6**, 23799 (2016).

1085
1086   42.   Frisen J, Yates PA, McLaughlin T, Friedman GC, O'Leary DD, Barbacid M. Ephrin-A5 (AL-
1087   1/RAGS) is essential for proper retinal axon guidance and topographic mapping in the
1088   mammalian visual system. *Neuron* **20**, 235-243 (1998).

1089
1090   43.   Fujita Y, Nakanishi T, Ueno M, Itohara S, Yamashita T. Netrin-G1 Regulates Microglial
1091   Accumulation along Axons and Supports the Survival of Layer V Neurons in the Postnatal Mouse
1092   Brain. *Cell Rep* **31**, 107580 (2020).

1093
1094   44.   Ross SE*, et al.* Bhlhb5 and Prdm8 form a repressor complex involved in neuronal circuit
1095   assembly. *Neuron* **73**, 292-303 (2012).

1096
1097   45.   Palmesino E*, et al.* Foxp1 and lhx1 coordinate motor neuron migration with axon trajectory
1098   choice by gating Reelin signalling. *PLoS Biol* **8**, e1000446 (2010).

1099
1100   46.   Marcos-Mondejar P, Peregrin S, Li JY, Carlsson L, Tole S, Lopez-Bendito G. The lhx2
1101   transcription factor controls thalamocortical axonal guidance by specific regulation of robo1 and
1102   robo2 receptors. *J Neurosci* **32**, 4372-4385 (2012).

1103
1104   47.   Fettiplace R, Hackney CM. The sensory and motor roles of auditory hair cells. *Nature reviews
1105   Neuroscience* **7**, 19-29 (2006).

1106
1107   48.   Neely S, Kim DO. *Cochlear models incorporating active processes*. Springer New York (2007).

1108
1109   49.   Muller M, von Hunerbein K, Hoidis S, Smolders JW. A physiological place-frequency map of the
1110   cochlea in the CBA/J mouse. *Hear Res* **202**, 63-73 (2005).

1111
1112   50.   Zhou X, Jen PH, Seburn KL, Frankel WN, Zheng QY. Auditory brainstem responses in 10 inbred
1113   strains of mice. *Brain Res* **1091**, 16-26 (2006).

1114
1115   51.   Chumak T*, et al.* BDNF in Lower Brain Parts Modifies Auditory Fiber Activity to Gain Fidelity
1116   but Increases the Risk for Generation of Central Noise After Injury. *Molecular Neurobiology* **53**,
1117   5607-5627 (2016).

1118
1119   52.   Land R, Burghard A, Kral A. The contribution of inferior colliculus activity to the auditory
1120   brainstem response (ABR) in mice. *Hearing Research* **341**, 109-118 (2016).

1121
1122   53.   Liu Y*, et al.* Hearing loss is an early biomarker in APP/PS1 Alzheimer's disease mice. *Neurosci
1123   Lett* **717**, (2020).

1124

1125 54.   Melcher JR, Kiang NYS. Generators of the brainstem auditory evoked potential in cat III:
1126   identified cell populations. *Hearing Research* **93**, 52-71 (1996).

1127
1128 55.   Wolter S*, et al.* GC-B Deficient Mice With Axon Bifurcation Loss Exhibit Compromised
1129   Auditory Processing. *Frontiers in Neural Circuits* **12**, (2018).

1130
1131 56.   Chambers AR*, et al.* Central Gain Restores Auditory Processing following Near-Complete
1132   Cochlear Denervation. *Neuron* **89**, 867-879 (2016).

1133
1134 57.   Harris JA, Rubel EW. Afferent regulation of neuron number in the cochlear nucleus: cellular and
1135   molecular analyses of a critical period. *Hear Res* **216-217**, 127-137 (2006).

1136
1137 58.   Lauer AM, Connelly CJ, Graham H, Ryugo DK. Morphological characterization of bushy cells
1138   and their inputs in the laboratory mouse (Mus musculus) anteroventral cochlear nucleus. *PLoS*
1139   *One* **8**, e73308 (2013).

1140
1141 59.   Karmakar K*, et al.* Hox2 Genes Are Required for Tonotopic Map Precision and Sound
1142   Discrimination in the Mouse Auditory Brainstem. *Cell Rep* **18**, 185-197 (2017).

1143
1144 60.   Fujiyama T*, et al.* Inhibitory and excitatory subtypes of cochlear nucleus neurons are defined by
1145   distinct bHLH transcription factors, Ptf1a and Atoh1. *Development* **136**, 2049-2058 (2009).

1146
1147 61.   Gruters KG, Groh JM. Sounds and beyond: multisensory and other non-auditory signals in the
1148   inferior colliculus. *Front Neural Circuits* **6**, 96 (2012).

1149
1150 62.   Yeomans JS, Frankland PW. The acoustic startle reflex: neurons and connections. *Brain*
1151   *Research Reviews* **21**, 301-314 (1995).

1152
1153 63.   Koch M. The neurobiology of startle. *Progress in neurobiology* **59**, 107-128 (1999).

1154
1155 64.   Lee Y, Lopez DE, Meloni EG, Davis M. A primary acoustic startle pathway: obligatory role of
1156   cochlear root neurons and the nucleus reticularis pontis caudalis. *J Neurosci* **16**, 3775-3789
1157   (1996).

1158
1159 65.   Swerdlow NR, Geyer MA, Braff DL. Neural circuit regulation of prepulse inhibition of startle in
1160   the rat: current knowledge and future challenges. *Psychopharmacology* **156**, 194-215 (2001).

1161
1162 66.   Fitch RH, Threlkeld SW, McClure MM, Peiffer AM. Use of a modified prepulse inhibition
1163   paradigm to assess complex auditory discrimination in rodents. *Brain research bulletin* **76**, 1-7
1164   (2008).

1165
1166 67.   Hickox AE, Liberman MC. Is noise-induced cochlear neuropathy key to the generation of
1167   hyperacusis or tinnitus? *J Neurophysiol* **111**, 552-564 (2014).

1168
1169   68.   Pfaff SL, Mendelsohn M, Stewart CL, Edlund T, Jessell TM. Requirement for LIM homeobox
1170   gene Isl1 in motor neuron generation reveals a motor neuron-dependent step in interneuron
1171   differentiation. *Cell* **84**, 309-320 (1996).

1172
1173   69.   Yang L*, et al.* Isl1Cre reveals a common Bmp pathway in heart and limb development.
1174   *Development* **133**, 1575-1585 (2006).

1175
1176   70.   Huber K, Narasimhan P, Shtukmaster S, Pfeifer D, Evans SM, Sun Y. The LIM-Homeodomain
1177   transcription factor Islet-1 is required for the development of sympathetic neurons and adrenal
1178   chromaffin cells. *Dev Biol* **380**, 286-298 (2013).

1179
1180   71.   Ahlgren U, Pfaff SL, Jessell TM, Edlund T, Edlund H. Independent requirement for ISL1 in
1181   formation of pancreatic mesenchyme and islet cells. *Nature* **385**, 257-260 (1997).

1182
1183   72.   Cai CL*, et al.* Isl1 identifies a cardiac progenitor population that proliferates prior to
1184   differentiation and contributes a majority of cells to the heart. *Dev Cell* **5**, 877-889 (2003).

1185
1186   73.   Liang X*, et al.* Isl1 is required for multiple aspects of motor neuron development. *Molecular and
1187   cellular neurosciences* **47**, 215-222 (2011).

1188
1189   74.   Kania A, Jessell TM. Topographic motor projections in the limb imposed by LIM homeodomain
1190   protein regulation of ephrin-A:EphA interactions. *Neuron* **38**, 581-596 (2003).

1191
1192   75.   Yamashita T*, et al.* High-resolution transcriptional dissection of in vivo Atoh1-mediated hair cell
1193   conversion in mature cochleae identifies Isl1 as a co-reprogramming factor. *PLoS Genet* **14**,
1194   e1007552 (2018).

1195
1196   76.   Chumak T*, et al.* Deterioration of the Medial Olivocochlear Efferent System Accelerates Age-
1197   Related Hearing Loss in Pax2-Isl1 Transgenic Mice. *Mol Neurobiol* **53**, 2368-2383 (2016).

1198
1199   77.   Bohuslavova R*, et al.* Pax2-Islet1 Transgenic Mice Are Hyperactive and Have Altered Cerebellar
1200   Foliation. *Mol Neurobiol* **54**, 1352-1368 (2017).

1201
1202   78.   Huang M, Kantardzhieva A, Scheffer D, Liberman MC, Chen ZY. Hair cell overexpression of
1203   Islet1 reduces age-related and noise-induced hearing loss. *J Neurosci* **33**, 15086-15094 (2013).

1204
1205   79.   Morris JK*, et al.* A disorganized innervation of the inner ear persists in the absence of ErbB2.
1206   *Brain Res* **1091**, 186-199 (2006).

1207
1208   80.   Mao Y, Reiprich S, Wegner M, Fritzsch B. Targeted deletion of Sox10 by Wnt1-cre defects
1209   neuronal migration and projection in the mouse inner ear. *PLoS One* **9**, e94580 (2014).

1210

1211 81.   Geyer MA, McIlwain KL, Paylor R. Mouse genetic models for prepulse inhibition: an early
1212   review. *Mol Psychiatry* **7**, 1039-1053 (2002).

1213
1214 82.   LeMasurier M, Gillespie PG. Hair-cell mechanotransduction and cochlear amplification. *Neuron*
1215   **48**, 403-415 (2005).

1216
1217 83.   Dallos P*, et al.* Prestin-based outer hair cell motility is necessary for mammalian cochlear
1218   amplification. *Neuron* **58**, 333-339 (2008).

1219
1220 84.   Herranen A*, et al.* Deficiency of the ER-stress-regulator MANF triggers progressive outer hair
1221   cell death and hearing loss. *Cell death & disease* **11**, 100 (2020).

1222
1223 85.   Liberman LD, Liberman MC. Cochlear Efferent Innervation Is Sparse in Humans and Decreases
1224   with Age. *J Neurosci* **39**, 9560-9569 (2019).

1225
1226 86.   Hafidi A. Peripherin-like immunoreactivity in type II spiral ganglion cell body and projections.
1227   *Brain Res* **805**, 181-190 (1998).

1228
1229 87.   Elliott KL*, et al.* Developmental Changes in Peripherin-eGFP Expression in Spiral Ganglion
1230   Neurons. *Front Cell Neurosci* **15**, 678113 (2021).

1231
1232 88.   Zhang KD, Coate TM. Recent advances in the development and function of type II spiral
1233   ganglion neurons in the mammalian inner ear. *Seminars in cell & developmental biology* **65**, 80-
1234   87 (2017).

1235
1236 89.   Froud KE*, et al.* Type II spiral ganglion afferent neurons drive medial olivocochlear reflex
1237   suppression of the cochlear amplifier. *Nature communications* **6**, 7115 (2015).

1238
1239 90.   Maison S, Liberman LD, Liberman MC. Type II Cochlear Ganglion Neurons Do Not Drive the
1240   Olivocochlear Reflex: Re-Examination of the Cochlear Phenotype in Peripherin Knock-Out
1241   Mice. *eNeuro* **3**, (2016).

1242
1243 91.   Matei V*, et al.* Smaller inner ear sensory epithelia in Neurog 1 null mice are related to earlier hair
1244   cell cycle exit. *Dev Dyn* **234**, 633-650 (2005).

1245
1246 92.   Pauley S, Lai E, Fritzsch B. Foxg1 is required for morphogenesis and histogenesis of the
1247   mammalian inner ear. *Dev Dyn* **235**, 2470-2482 (2006).

1248
1249 93.   Nichols DH, Pauley S, Jahan I, Beisel KW, Millen KJ, Fritzsch B. Lmx1a is required for
1250   segregation of sensory epithelia and normal ear histogenesis and morphogenesis. *Cell Tissue Res*
1251   **334**, 339-358 (2008).

1252

1253   94.   Nichols DH, *et al.* Interaction with ectopic cochlear crista sensory epithelium disrupts basal
1254   cochlear sensory epithelium development in Lmx1a mutant mice. *Cell Tissue Res* **380**, 435-448
1255   (2020).

1256
1257   95.   Dvorakova M, Macova I, Bohuslavova R, Anderova M, Fritzsch B, Pavlinkova G. Early ear
1258   neuronal development, but not olfactory or lens development, can proceed without SOX2. *Dev*
1259   *Biol* **457**, 43-56 (2020).

1260
1261   96.   Pavlinkova G. Molecular Aspects of the Development and Function of Auditory Neurons. *Int J*
1262   *Mol Sci* **22**, (2020).

1263
1264   97.   Zine A, Messat Y, Fritzsch B. A human induced pluripotent stem cell-based modular platform to
1265   challenge sensorineural hearing loss. *Stem Cells*, (2021).

1266
1267   98.   Susaki EA, Tainaka K, Perrin D, Yukinaga H, Kuno A, Ueda HR. Advanced CUBIC protocols
1268   for whole-brain and whole-body clearing and imaging. *Nat Protoc* **10**, 1709-1727 (2015).

1269
1270   99.   Mamanova L, Miao Z, Jinat A, Ellis P, Shirley L, Teichmann SA. High-throughput full-length
1271   single-cell RNA-seq automation. *Nat Protoc* **16**, 2886-2915 (2021).

1272
1273   100.   Dobin A, *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

1274
1275   101.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011* **17**,
1276   3 (2011).

1277
1278   102.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
1279   *Bioinformatics* **30**, 2114-2120 (2014).

1280
1281   103.   Kopylova E, Noe L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in
1282   metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).

1283
1284   104.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
1285   data with DESeq2. *Genome Biol* **15**, 550 (2014).

1286
1287   105.   Boyle EI, *et al.* GO::TermFinder--open source software for accessing Gene Ontology information
1288   and finding significantly enriched Gene Ontology terms associated with a list of genes.
1289   *Bioinformatics* **20**, 3710-3715 (2004).

1290
1291   106.   Raudvere U, *et al.* g:Profiler: a web server for functional enrichment analysis and conversions of
1292   gene lists (2019 update). *Nucleic acids research* **47**, W191-W198 (2019).

1293
1294   107.   Shannon P, *et al.* Cytoscape: a software environment for integrated models of biomolecular
1295   interaction networks. *Genome Res* **13**, 2498-2504 (2003).

1296
1297   108.   Merico D, Isserlin R, Stueker O, Emili A, Bader GD. Enrichment map: a network-based method
1298   for gene-set enrichment visualization and interpretation. *PLoS One* **5**, e13984 (2010).

1299
1300   109.   Fritzsch B, Duncan JS, Kersigo J, Gray B, Elliott KL. Neuroanatomical Tracing Techniques in
1301   the Ear: History, State of the Art, and Future Developments. In: *B. Sokolowski, Ed: Auditory and*
1302   *Vestibular Research: Methods and Protocols*). Springer Science+Business Media New York
1303   (2016).

1304
1305   110.   Bures Z, Grécová J, Popelár J, Syka J. Noise exposure during early development impairs the
1306   processing of sound intensity in adult rats. *The European journal of neuroscience* **32**, 155-164
1307   (2010).

1308

1309