ISL1 is necessary for auditory neuron development and tonotopic organization

Authors: Iva Filova[1,7], Kateryna Pysanenko[2,7], Mitra Tavakoli[1], Simona Vochyanova[1], Martina Dvorakova[1], Romana Bohuslavova[1], Ondrej Smolik[1], Valeria Fabriciova[1], Petra Hrabalova[1], Sarka Benesova[3], Lukas Valihrach[3], Jiri Cerny[4], Ebenezer N. Yamoah[5], Josef Syka[2], Bernd Fritzsch[6], and Gabriela Pavlinkova[1*]

Affiliations:
[1]Laboratory of Molecular Pathogenetics, Institute of Biotechnology CAS, 25250 Vestec, Czechia
[2]Department of Auditory Neuroscience, Institute of Experimental Medicine CAS, 14220 Prague, Czechia
[3]Laboratory of Gene Expression, Institute of Biotechnology CAS, 25250 Vestec, Czechia
[4]Laboratory of Light Microscopy, Institute of Molecular Genetics CAS, 14220 Prague, Czechia
[5]Department of Physiology, School of Medicine, University of Nevada Reno, NV 89557, USA
[6]Department of Biology, Department of Otolaryngology, University of Iowa, Iowa City, IA 52242-1324, USA
[7]These authors contributed equally: Iva Filova, Kateryna Pysanenko.
[*] Email: Gabriela.Pavlinkova@ibt.cas.cz

Abstract

A cardinal feature of the auditory pathway is frequency selectivity, represented in a tonotopic map from the cochlea to the cortex. The molecular determinants of the auditory frequency map are unknown. Here, we discovered that the transcription factor ISL1 regulates the molecular and cellular features of auditory neurons, including the formation of the spiral ganglion and peripheral and central processes that shape the tonotopic representation of the auditory map. We selectively knocked out *Isl1* in auditory neurons using *Neurod1^{Cre}* strategies. In the absence of *Isl1*, spiral ganglion neurons migrate into the central cochlea and beyond, and the cochlear wiring is profoundly reduced and disrupted. The central axons of *Isl1* mutants lose their topographic projections and segregation at the cochlear nucleus. Transcriptome analysis of spiral ganglion neurons shows that *Isl1* regulates neurogenesis, axonogenesis, migration, neurotransmission-related machinery, and synaptic communication patterns. We show that peripheral disorganization in the cochlea affects the physiological properties of hearing in the midbrain and auditory behavior. Surprisingly, auditory processing features are preserved despite the significant hearing impairment, revealing central auditory pathway resilience and plasticity in *Isl1* mutant mice. Mutant mice have a reduced acoustic startle reflex, altered prepulse inhibition, and characteristics of compensatory neural hyperactivity centrally. Our findings show that ISL1 is one of the obligatory factors required to sculpt auditory structural and functional tonotopic maps. Still, upon *Isl1* deletion, the ensuing central compensatory plasticity of the auditory pathway does not suffice to overcome developmentally induced peripheral dysfunction of the cochlea.

Keywords

spiral ganglion neurons, inferior colliculus, auditory nuclei, inner ear, brain plasticity, *Isl1* mutation, auditory maps, auditory behavior

Introduction

Spiral ganglion neurons (SGNs) are bipolar, extending peripheral processes to the hair cells within the sensory epithelium (the organ of Corti) and central axons towards the cochlear nucleus (CN) complex, the first auditory nuclei in the brain. Sound-induced vibrations that reach the cochlea are amplified by the outer hair cells (OHCs) organized in three rows and innervated by type II SGNs. The inner hair cells (IHCs) receive, transduce, and transmit the auditory signal to type I SGNs that convey the signal via the vestibulocochlear cranial nerve to the CN of the brainstem. The auditory neurons are organized within the cochlea in an orderly fashion according to frequency, with high frequencies at the base and low frequencies at the apex [1, 2]. The tonotopic organization of type I SGNs corresponds to multiple diversities in their molecular profiles, connectivity patterns, and physiological features along the tonotopic axis [3, 4, 5]. The cochleotopic or tonotopic pattern is maintained throughout the auditory pathways in the brain [6]. The central auditory pathway transmits ascending acoustic information from the CN through the lateral lemniscus complex, the inferior colliculus in the midbrain, the medial geniculate nucleus of the thalamus to the auditory cortex [7]. The efferent motor neurons consist of the medial olivocochlear motor neurons, which modulate cochlear sound amplification by OHCs. In contrast, lateral olivocochlear motor neurons innervate afferent type I sensory neurons and regulate cochlear nerve excitability [7, 8, 9].

The cellular and molecular regulation of neuronal migration and the establishment of tonotopic connections to the hair cells or neurons of the hindbrain's first auditory nuclei, the CN, are not fully understood. Several transcription factors govern the development of inner ear neurons, including NEUROG1 [10], NEUROD1 [11], GATA3 [12, 13], and POU4F1 [14]. However, how the transcriptional networks regulate SGN development and how abnormalities in auditory neurons and neuronal wiring reshape the auditory central pathways' assembly and acoustic information processing remain unresolved.

The transcription factor *Neurod1* is vital for the differentiation and survival of inner ear neurons [15, 16, 17]. We previously demonstrated that *Isl1^Cre*-mediated *Neurod1* deletion (*Neurod1CKO*) results in a disorganized cochleotopic projection from SGNs [18], affecting acoustic information processing in the central auditory system of adult mice at the physiological and behavioral levels [19]. During inner ear development, ISL1 is expressed in the differentiating neurons and sensory precursors [18, 20, 21, 22, 23]. *Isl1* is expressed in all four populations of SGNs (type Ia, Ib, Ic, and type II) identified by single-cell RNA sequencing [4]. Studies suggest that ISL1 plays a role in developing neurons and sensory cells, but there has been no direct evaluation of ISL1 function in the inner ear. Using *Neurod1^Cre* [24], we conditionally deleted *Isl1* (*Isl1CKO*) in neurons without directly affecting the development of the inner ear sensory epithelium.

This work provides the first genetic and functional evidence for ISL1's role in establishing the spiral ganglion peripheral projection map and proper central auditory circuitry. Most *Isl1CKO* neurons migrated into the center of the cochlear modiolus. They extended outside the cartilaginous/bony otic capsule, in contrast, to control animals with an arrangement of SGNs in parallel to the spiraling cochlear duct. Additionally, we analyzed the transcriptome of neurons, hearing function, sound information processing in the inferior colliculus, and auditory behavior of *Isl1CKO* to demonstrate how abnormal SGN development affects the formation, wiring, and operation of the auditory sensory maps.

Results

ISL1 controls neuronal phenotype in the cochlea

To investigate the role of ISL1 in inner ear neuron development, we eliminated *Isl1* by crossing *Isl1*$^{loxP/loxP}$ mice [25] with *Neurod1*^{Cre} mice [24]. *Neurod1*^{Cre} is expressed in sensory neurons but not in the sensory epithelium, as visualized by *tdTomato* reporter expression (Supplementary Fig. 1). Analyses of ISL1 expression in *Isl1CKO* confirmed efficient recombination by *Neurod1*^{Cre} with virtually no expression of ISL1 during the differentiation of neurons in the inner ear ganglion as early as embryonic day E10.5, and later in the cochlear neurons, shown at E14.5 and postnatal day 0 (P0), in contrast to high levels of ISL1 in neurons in the control inner ear (Supplementary Fig. 2). It is worth mentioning that no difference was observed in the density of Schwann cells (Supplementary Fig. 2e, f).

The organization of the sensory epithelium in the organ of Corti of *Isl1CKO* was comparable to controls with three rows of OHCs and one row of IHCs (Fig. 1a-j), as shown by immunolabeling with the hair cell differentiation marker, myosin VIIA (Myo7a). Whole-mount anti-tubulin staining of innervation showed reduced and disorganized radial fibers. Significant gaps were found between radial fiber bundles, crisscrossing fibers, and an unusual dense innervation in the apex, with some large fiber bundles bypassing the organ of Corti and extending to the lateral wall in the *Isl1CKO* cochlea (arrowhead, Fig. 1j). Note, efferent fibers did not correctly form an intraganglionic spiral bundle (IGSB) in *Isl1CKO*.

Next, we evaluated the effects of *Isl1* deletion on the formation of the spiral ganglion. The SGN somas were well restricted to the Rosenthal's canal in control animals, arranged in a spiral parallel to the cochlear duct (Fig. 1k-m). In contrast, unevenly distributed neurons were found in *Isl1CKO* with many SGNs accumulated beyond the Rosenthal's canal, in the conical central part of the cochlea, the modiolus (Fig. 1n-p), which usually carries only afferent and efferent fibers. Reduced clusters of neurons were found in the apex (Fig. 1q, r). Noticeably, some neurons were located in the Rosenthal's canal. Higher-magnification images showed the aberrant distribution of tdTomato neurons$^+$ entangled with radial fibers near the organ of Corti (Fig. 1p, r). The unusual accumulation of cochlear neurons intertwined with the central axons in the modiolus of *Isl1CKO* compared to the control mice is depicted in the immunolabeled sections of the cochlea (Fig. 1s, t). A smaller number of neurons was positioned within the Rosenthal's canal of *Isl1CKO*, representing on average ~20% and 30% neurons in the basal and apical half of the control cochlea, respectively (Fig. 1u). To determine whether neuron number was affected by apoptosis during inner ear development, we performed immunostaining for cleaved Caspase-3 (Supplementary Fig. 3). We found moderately increased apoptosis of neurons in *Isl1CKO* at E12.5 but not at E10.5, although the size of inner ear ganglia was unaffected at E12.5. These results imply that missing neurons in the Rosenthal's canal are due to the altered migration of SGNs rather than neuronal death in the developing inner ear. Correspondingly, most apoptotic neurons were found outside the cochlea in the vestibular ganglion area in E14.5 *Isl1CKO*. In line with the unusual position of SGNs in the *Isl1CKO* cochlea, more elongated and disorganized radial fibers extended from the modiolus to the organ of Corti than the control cochlea. The radial fibers were significantly lengthened in all regions of the *Isl1CKO* cochlea (Supplementary Fig. 4), but the overall radial fiber density was reduced (Fig. 1w).

Interestingly, *Isl1* deficiency caused a shortening of the cochlea, which was on average ~20% and 25% shorter at P0 and in adults, respectively (Fig. 2a, b). Since *Neurod1*^{Cre} is expressed only in neurons, the shortening of the cochlea appears to be a secondary effect of the

abnormalities in neuronal development. A similar phenotype of truncated growth of the cochlea
was reported for *Neurod1* and *Neurog1* mutants because of alternations in spatiotemporal gene
expression [10, 18, 19, 26]. Correspondingly to the truncated cochlear phenotype [10, 18, 19, 26], detailed
analyses showed abnormalities in the epithelium at the apical end with disorganized rows of
OHCs and ectopic IHCs among OHCs in the *Isl1CKO* cochlea (Fig. 2c). The disorganized apical
epithelium represented ~11.8% of the length of the organ of Corti in adults (n = 6). Anti-tubulin
labeling revealed a reduced and disorganized innervation in the region of OHCs in *Isl1CKO* (Fig.
2d). In the controls, fibers extending through the tunnel of Corti were oriented towards the base
to form three parallel outer spiral bundles. In contrast, guiding defects in the extension of these
fibers in *Isl1CKO* were apparent, with neurites extending with random turns towards the base
and the apex (arrowheads in Fig. 1d). To further investigate cochlear histopathology in *Isl1CKO*,
we evaluated synaptic contacts from type I SGNs onto IHCs in 2-month-old mice (Fig. 2e).
CtBP2$^+$ ribbon synapse number was reduced in *Isl1CKO*, representing ~46%, 51%, 56%, and
69% ribbons of the control base, mid-base, mid-apex, and apex, respectively (Fig. 2f,
Supplementary Table 1).
Altogether, these results demonstrate the prominent effects of *Isl1* deletion on the
formation of the spiral ganglion and innervation patterns in the cochlea. The truncated cochlea in
Isl1CKO indicates an altered spatial base-to-apex organization of structural and physiological
characteristics of auditory neurons.
157
ISL1 regulates neuronal identity and differentiation programs of SGNs
To gain insight at the molecular level on how the *Isl1* elimination causes the neuronal phenotype
in the cochlea, we sought to identify potential ISL1 targets through global transcriptome
analysis. We opted to use Bulk-RNA sequencing to obtain sequencing depth and high-quality
data [27]. Six biological replicates were used per genotype, and each replicate contained a total of
100 tdTomato$^+$ SGNs isolated from the E14.5 cochlea. Spiral ganglia were dissected, dissociated
into single cells, and fluorescent tdTomato$^+$ cells were FACS-sorted (Fig. 3a). Compared to
controls, 650 protein-coding genes were differentially expressed in *Isl1CKO* neurons (adjusted p-
value < 0.05, and fold change > 2 cut-off values, see Methods), 332 genes down- and 318 genes
up-regulated (Fig. 3b, Supplementary Data 1). Gene ontology (GO) term enrichment analysis for
the GO term category biological process revealed highly enriched GO terms associated with
neuron development, including "neurogenesis", "neuron differentiation", and "nervous system
development", in both up- and down-differentially expressed genes (Fig. 3c, Supplementary Data
2). The most enriched and specific GO categories for downregulated genes were associated with
neurotransmission-related machineries, such as "transmembrane transporter", "voltage-gated
channel", "cation and ion transport", and "membrane potential regulation", indicating changes in
neuronal cell functions. The analysis identified enrichment of downregulated genes involved in
axon development, guidance, axonogenesis, and neuronal migration. These genes included
members of all four classes of axon guidance molecules and their cognate receptors, the ephrins-
Eph, semaphorins-plexin, netrin-unc5, and slit-roundabout [28], for instance, *Epha5, Epha4,*
Pdzrn3, Sema3e, Sema6d, Plxna4, Ntn3, Ntng1, Ntng2, Kirrel3, Unc5b, Unc5c, Slitrk3, and
Slitrk1. The neurotrophic tyrosine kinase receptors (*Ntrk2* and *Ntrk3*) and a G-protein-coupled
chemokine receptor (*Cxcr4*), important regulators of neuronal migration, were also
downregulated [29].
In contrast, upregulated genes particularly enriched in *Isl1CKO* neurons were associated
with "regulation of synapse organization, synapse structure or activity" and "regulation of

synapse assembly", indicating compensatory changes relating to the ability of neurons to form neuronal projections towards their targets. Upregulated genes encoding molecules critical for synapse formation and adhesion included several members of the cadherin superfamily (*Cdh7, Cdh10*, *Pcdh8, Pcdh9, Pcdh10, Pcdh18,* and *Pcdh19*), adhesion-related genes (*Ptprd, Ptprs, Cntnap2, Itga4, Itga5*), and the ephrin ligands (*Efna3*, *Efna5*). Interestingly, "synaptic signaling" was among the GO terms with significant representation in both up- and down-regulated genes, suggesting changes in synaptic circuits and neuronal activity. We also found many genes encoding transcription factors and signaling molecules, indicating that ISL1 may act through transcription networks instead of defined target genes. The expression of neural-specific bHLH factors important for differentiation and maturation of neurons was increased in *Isl1CKO*, including members of NeuroD (*Neurod6, Neurod2*), Nscl (*Nhlh2*), and Olig (*Bhlhe23*) families [30]. Genes encoding LIM domain proteins, the transcription factors *Lhx1, Lhx2*, *Lmo2*, *Lmo*3, a core member of the PCP signaling paradigm, *Prickle1*, and neuronal homeobox genes, *Emx2*, *Uncx*, and *Pknox2* were upregulated in *Isl1CKO* neurons. *Prickle1* is an essential regulator of neuron migration and neurons' distal and central projections in the cochlea [31]. Some of the identified genes encoding regulatory molecules shown to be essential for neuronal development, the formation of SGNs, and their projections were downregulated in *ISL1CKO* neurons, such as the signaling molecules *Shh* [32], *Wnt3* [3], and *Fgfs* (*Fgf10, Fgf11,* and *Fgf13*) [33, 34], and the transcription factors *Gata3* [12, 13], *Irx1, Irx2, Pou3f2, Pou4f2* [22], and *Runx1* [3, 4].

Consistent with the idea that ISL1 may control a gene expression program mediating different aspects of neuronal development in the cochlea, several identified genes are known to affect the formation of SGNs and their innervation patterns. For instance, a mouse model with EphA4 deficient signaling in SGNs results in aberrant mistargeting of type I projections to OHCs [35] and SGN fasciculation defects [36]. Targeted disruptions of the neurotrophin receptors, *Ntrk2* and *Ntrk3*, are associated with misplaced SGNs in the modiolus, disorganized projections, and reduced neuronal survival [37]. The neurotrophin receptor p75NTR (*Ngfr*)-deficient SGNs demonstrate enhanced neurite growth behavior [38]. Mutation of *Prickle1* results in disorganized apical afferents [31]. Reduced and aberrant cochlear innervation was shown in mice with *Gata3*-specific SGN deletion [13]. *Nrp2* expressed in SGNs is required for axon path finding and a normal pattern of cochlear innervation [39]. The axon guidance molecule SLIT2 and its receptor ROBO1/2 control precise innervation patterns, and their deletions induce misplacement of SGNs [40]. Mice lacking *Dcc* have the more severe phenotype of mislocated SGNs and disorganized central projections, although most SGNs of *Dcc*[-/-] remained within Rosenthal's canal [41].

Other significantly deregulated genes mediate several aspects of neural-circuit formation in the brain, including neurite elongation, axon guidance, and interactions between neurons and microglia. For instance, deletion of *Epha5* alters retinal axon guidance within the midbrain [42]; inhibition of *Ntng1* reduces microglial accumulation and loss of cortical neurons [43]; the absence of either *Bhlhb22* or *Prdm8* alters neural circuit assembly [44]; LHX1 controls neuronal migration and axonal projections of motor neurons [45]; and LHX2 influences axon guidance and the topographical sorting of axons by regulating ROBO1/2 [46]. These molecular differences dovetail well with abnormalities in the innervation pattern and SGN migration defects *Isl1CKO*. Selected differentially expressed genes from the RNAseq analysis, namely *Dcc, Epha5, Ntng1, Prdm8, Robo2, Slit2, Tbx3, Uncx, Gata3, Lhx1, Lhx2, Cdh7, Nhlh2, Ntrk2,* and *Ntrk3,* were further validated by RT-qPCR of RNA isolated from whole inner ears of E14.5 embryos (Supplementary Fig. 5).

Altogether, these changes indicate that ISL1 regulates transcriptional networks that
underlie neuronal identity and function during differentiation of SGNs and that *Isl1* elimination
results in a major impairment in the development, axonogenesis, migration, and molecular
characteristics of these neurons.

Isl1CKO mice have altered hearing function
Considering the substantial abnormalities in the formation of the spiral ganglion, innervation,
and molecular neuron features, we assessed the hearing function of *Isl1CKO* mice. We evaluated
distortion product otoacoustic emissions (DPOAE) to determine the robustness of OHC function
and cochlear amplification. Otoacoustic emissions are a physiological byproduct of an active
amplification mechanism when sound-induced vibrations are amplified in a frequency-specific
manner by the OHCs of the organ of Corti [47, 48]. Compared to controls, the DPOAE responses of
Isl1CKO were significantly reduced in the frequency range between 4 and 24 kHz (Fig. 4a).
DPOAE amplitudes at frequencies of 26 kHz and higher were comparable between control and
Isl1CKO mice. Based on the physiological place-frequency map in the normal mouse cochlea [49],
frequencies above 26 kHz are located at the basal half of the cochlea. Thus, decreased DPOAE
responses may be attributed to the more profound morphological abnormalities in the apex,
disorganized innervation, and disorganized rows of OHCs in the apical end.
We evaluated auditory brainstem responses (ABRs), which measure electrical activity
associated with the propagation of acoustic information through auditory nerve fibers.
Measurements of ABR thresholds showed that all *Isl1CKO* animals displayed elevated
thresholds indicative of hearing loss compared to age-matched control animals, except at
frequencies 32 and 40 kHz, which were comparable to ABR thresholds in control mice (Fig. 4b).
Using click-evoked ABR, we evaluated waveform characteristics associated with the
propagation of acoustic information through the auditory nerve to higher auditory centers (Fig.
4c). Wave I reflects the synchronous firing of the auditory nerve. In contrast, waves II–V are
attributed to the electrical activity of downstream circuits in the CN, superior olivary complex,
lateral lemniscus, and inferior colliculus [50]. Waves IV and V are products of the sound-evoked
neuronal activity in the lateral lemniscus and inferior colliculus [51, 52, 53, 54, 55]. The waves IV and
V are often paired and superimposed in commonly used non-transgenic mouse strains [52, 55]. The
ABR traces in all our panels were averaged traces from all measured animals. A part of the
animals in both experimental groups had a hardly distinguishable wave V in response to the click
and 8 kHz sound stimulation; therefore, to prevent bias, we did not include wave V. The
amplitudes of ABR waves I-IV were significantly reduced in *Isl1CKO* (Fig. 4d). Since the ABR
threshold for 32 kHz and above were comparable between age-matched controls and *Isl1CKO*
mutants (Fig. 4b), we used the pure-tone stimuli of 32 kHz to evaluate ABR responses (Fig. 4e).
A significant difference amongst the genotypes was found only for amplitude reduction of wave
III (Fig. 4f), thus indicating preserved synchronized activities of peripheral and brainstem
auditory processing. Although waves I and II amplitude for both mutant and control mice were
similar, there were apparent differences in the ABR waveform morphology. The latency of ABR
wave I was delayed, the relative interwave latency between peaks I and II was shortened, and the
trough between wave I and II diminished, resulting in a fusion of both peaks in *Isl1CKO*. A
delay of the leading peak of ABR wave I recovered towards ABR wave III. The wave I and II
changes reflect abnormalities in the summated response from SGNs, auditory nerve fibers, and
most likely the CN.

Additionally, we used a pure-tone stimulus of 8 kHz to evaluate ABR responses (Fig.
4g), as ABR thresholds for 8 kHz were significantly reduced for *Isl1CKO* mice. In contrast to
ABR amplitudes at 32 kHz stimuli, ABR amplitudes for 8 kHz pure-tone stimuli were
significantly reduced for I-III waves (Fig. 4h). The results indicate abnormalities in the cochlear
auditory neurons and propagation of acoustic information through auditory nerve fibers to higher
auditory centers. We observed a marked drop in the wave I growth function. Still, by comparing
wave I and IV peaks, an increased central gain was noted in *Isl1CKO* (Supplementary Fig. 6),
indicating compensatory plasticity at higher auditory circuits for cochlear damage with
diminished afferent input [56].

***Isl1CKO* mice display abnormalities in the ascending auditory pathways**
Having recognized uneven hearing function loss, as measured by ABRs, we further wanted to
investigate the distribution of SGNs in the *Isl1CKO* cochlea and the morphology of the
components of the central auditory pathway. We generated 3D visualization of the cochlea using
light-sheet fluorescent microscopy (Fig. 5a-i, Supplementary Videos S1-S4). In the first cochlear
preparation, neurons were visualized by *tdTomato* reporter expression and hair cells in the organ
of Corti were labeled using anti-Myo7a. In the second preparation, tdTomato$^+$ neurons were co-
labeled with anti-NeuN, a nuclear marker of differentiated neurons. Compared to the
recognizable coil of cochlear neurons of the spiral ganglion in the controls (Fig. 5a-c,
Supplementary Videos S1, S2), SGNs were aberrantly distributed in the *Isl1CKO* cochlea (Fig.
5d-i, Supplementary Videos S3, S4). Many *Isl1CKO* neurons were in the center of the cochlea,
the modiolus, that normally contains only central processes of SGNs, as shown in the control
cochlea (Fig. 5b, c). Some *Isl1CKO* neurons were found in their proper position in the
Rosenthal's canal, forming a dense strip of neurons, suggesting a partially preserved organization
of the spiral ganglion (Fig. 5f, i). This segment of the spiral ganglion formed in the mid-base of
the *Isl1CKO* cochlea correlates with the estimated characteristic place in the cochlear place-
frequency map for the characteristic frequency of 32 kHz [49] (Fig. 5j). A partial preservation of
the spiral ganglion in the mid-base corresponds to the retained high-frequency hearing function
of *Isl1CKO*. Identical to the SGN migration abnormalities recognized at P0, most SGNs were in
the modiolus, outside of Rosenthal's canal, in *Isl1CKO* adult mice (Supplementary Fig. 7). Apart
from the changes in the cochlear ganglion, the vestibular ganglion was fused and enlarged in
Isl1CKO compared to distinct superior and inferior vestibular ganglia in controls (Fig. 5k, l). The
spatial segregation between the vestibular and spiral ganglion was diminished in *Isl1CKO*, with
some areas fused, and thus, forming an aberrant "spiro-vestibular" ganglion.
Using dye tracing, we evaluated the segregation of central axons of the auditory nerve
(the cranial nerve VIII; Fig. 6a, schematic view of dye applications). In controls, the central
axons labeled by dyes applied into the base and apex are segregated in the auditory nerve and
from the vestibular ganglion, labeled by dye injected into the vestibular end-organs (Fig. 6b,
Supplementary Fig. 8). In contrast, the central axons from the cochlear base and apex virtually
overlapped in the auditory nerve (yellow fibers), and neurons labeled by cochlear dye
applications were detected to be intermingled with vestibular neurons in an aberrant "spiro-
vestibular" ganglion in *Isl1CKO* (Fig. 6c). Unfortunately, the mixing of spiral and vestibular
ganglion neurons in *Isl1CKO* mice precluded a full quantitative assessment. The unusual fibers
labeled by the apical dye application were looping around the vestibular ganglion in *Isl1CKO*.
The CN is the first structure of the ascending auditory pathways, where the auditory
nerve fibers project. The auditory nerve bifurcated with one branch, synapsing in the

posteroventral (PVCN) and dorsal (DCN) CN and the other innervating the anteroventral CN
(AVCN; Fig. 6a). Dye tracing showed segregated projections of apical and basal cochlear
afferents forming parallel isofrequency bands in controls at E16.5 (Fig. 6d). In contrast,
comparable injections in *Isl1CKO* showed that axonal projections to the CN were reduced,
restricted, disorganized, and lacked clear apex- and base-projection segregations, as many fibers
overlapped (Fig. 6e-g). Only a few fibers can occasionally be seen expanding to the DCN in
Isl1CKO at E16.5 (inset in Fig. 6e). Aberrant central projections were further demonstrated by
cochlear neurons projecting to the lateral vestibular nucleus (LVN) in *Isl1CKO* (Fig. 6i). No such
projections were found in control mice with segregated vestibular projections to the vestibular
nucleus, and basal and apical afferents reaching the CN (Fig. 6h). Collectively, these data show
disorganized central projections and a loss of tonotopic organization of both the auditory nerve
and the CN in *Isl1CKO,* as the apical and basal projections are not completely segregated (see
graphic summary in Fig. 6j).
Since the size and number of neurons in the CN depend on input from the auditory nerve
during a critical development period up to P9 [57], we also analyzed the volume of the CN of
Isl1CKO. The volume of the CN of adult mutants was reduced by approximately 50% compared
to controls at postnatal day P35 (Fig. 7a, b, g). As *Isl1* is not expressed in the CN [19], the reduced
size is likely a secondary effect associated with reduced afferent input consistent with the
previously reported impact of neonatal cochlear ablation [2]. The spherical and globular bushy
cells are principal cells that receive large auditory nerve endings, called "endbulbs of Held" and
"modified endbulbs", specialized for precise temporal firing [1, 58]. Using anti-parvalbumin to label
spherical bushy cells and anti-VGlut1 to label auditory-nerve endbulbs of Held [59, 60], we
demonstrated that auditory afferents of *Isl1CKO* and controls formed comparable clusters of
boutons that wrap the somas of their targets (Fig. 7c-f). Although the CN of *Isl1CKO* was
smaller, the density of bushy cells (number of parvalbumin$^+$ neurons per μm^2) was similar in
controls and *Isl1CKO* (Fig. 7h).
346
Characteristics of inferior colliculus neurons are distorted in *Isl1CKO*
We evaluated inferior colliculus (IC) properties, which is the principal auditory structure
of the midbrain for the ascending auditory pathways and descending inputs from the auditory
cortex [61]. The IC allows for sound localization, integrates multisensory and non-auditory
contributions to hearing, and plays an essential role in generating the startle response. We
demonstrated no significant IC size reduction in *Isl1CKO* compared to controls (Fig. 8a, b). We
compared neuronal characteristics in the central nucleus of the IC of *Isl1CKO* and control
animals using multichannel electrodes. Extracellular electrophysiological recordings of neuronal
activity in controls showed a well-defined narrow single-peaked profile of the excitatory
receptive fields (Fig. 8c). Compared to simple narrow V-shape receptive fields in controls, we
recorded the multipeaked broad tuning curves in the IC of *Isl1CKO* mice, suggesting multiple
inputs from the lower levels of the auditory system and overall, the worsened tunning
characteristics of IC neurons.
The investigation of the responsiveness of IC neurons (IC units) to different sound
frequencies showed higher excitatory thresholds in *Isl1CKO* compared to control animals in
measured frequencies between 6 to 28 kHz (Fig. 8d, e). However, responsiveness of IC units to
recorded frequencies above 28 kHz was comparable between control and mutant mice. These
comparable thresholds for high frequencies between control and *Isl1CKO* mice are consistent
with the ABR measurements. At the low frequency range, below 6 kHz, we did not record any

IC neurons in *Isl1CKO*. Compared to the controls, the IC neuronal responses in *Isl1CKO* had a higher broadband noise (BBN) threshold, a narrower dynamic range, higher spontaneous activity, and a lower maximum response magnitude (Fig. 8f). The results suggest a functional reduction in sensitivity to sound, audibility, intensity discrimination, and increased excitability of IC neurons in *Isl1CKO*. A commonly used metric unit of auditory tuning is the "quality factor", or Q, defined as the characteristic frequency (CF) divided by the bandwidth, measured at 20 dB above the minimum threshold (Q_{20}). Results revealed a significantly lower quality factor in the mutant mice (Fig. 8g), showing substantially worsened frequency selectivity.

To evaluate the precise temporal representation of sound into the central auditory system, we performed an acoustic stimulation of the IC units with trains of five clicks with different inter-click intervals from 100 ms up to 15 ms. In control mice, increasing time interval between clicks led to a better synchronization of neuronal responses with the individual clicks in the train, implying precision and reliability of temporal sound discrimination ability (Fig. 8h). In contrast, the precise temporal decoding in *Isl1CKO* was disrupted, as the synchronization of neuronal responses was significantly lower for the whole range of inter-click intervals. In the case of the *Isl1CKO,* the synchronization level of neuronal responses remains almost constant, suggesting a lack of precise temporal sound processing.

The auditory behavior of *Isl1CKO* mice is altered

Next, we evaluated the behavioral responses of *Isl1CKO* mice to sound stimuli. Since the vestibular dysfunctions might influence auditory behavioral responses, we first assessed the vestibular and motor function of *Isl1CKO*. During open-field observations, *Isl1CKO* mice did not display any signs of abnormal locomotor behavior, such as ataxia, difficulty maintaining balance, wagging or vertical bobbing movements of the head, or circling movements. The air-righting test, a basic vestibular function test, and motor function tests on the rotarod demonstrated comparable motor coordination between control and *Isl1CKO* mice (Supplementary Fig. 9). Concurrently, the size of the sensory epithelia of the vestibular organs and the size of the dorsal root ganglia were unaffected in *Isl1CKO* (Supplementary Fig. 9). Thus, locomotor activities of *Isl1CKO* were comparable to control mice. The acoustic startle response (ASR) is usually used as a behavioral readout of hearing status mediated by a brainstem circuit linking cochlear root neurons to spinal motoneurons. The structural basis of the ASR includes cochlear root neurons, CN neurons, the nucleus of the lateral lemniscus, the caudal pontine reticular nucleus, spinal interneurons, and spinal motor neurons [62, 63, 64]. Similar to the ABR thresholds, the ASR thresholds of *Isl1CKO* significantly increased for startle tone stimuli of 8 kHz and BBN, but not for the high-frequency startle tones (Fig. 9a). The peak latency of the ASR to the BBN stimulation at the 110 dB SPL intensity was prolonged in *Isl1CKO* (Fig. 9b), indicating a slower reaction to the acoustic stimuli. We found significantly reduced ASR amplitudes for all tested sound stimuli at higher intensities, showing deteriorated acoustic startle reactivity in *Isl1CKO* mice (Fig. 9c-e).

To further assess complex auditory discrimination behavior, we exposed control and *Isl1CKO* mice to a prepulse inhibition (PPI) paradigm, i.e., the inhibition of the ASR induced by presenting an acoustic stimulus shortly preceding the presentation of an acoustic stimulus, the startling sound. The circuit mediating a prepulse on the startle reflex involves central structures of the auditory pathway, including the IC and the auditory cortex [65, 66]. We used either BBN or pure tone pips of 8 and 32 kHz at increasing intensities as a non-startling acoustic stimulus (prepulse) that preceded the startle stimulus in a quiet background. The PPI with the prepulse of

32 kHz, the well-preserved audible frequency in *Isl1CKO*, was comparable between control and
mutant mice (Fig. 9h). Interestingly, the prepulse of 8 kHz resulted in a more prominent
inhibition of the startle response in *Isl1CKO* than in controls (Fig. 9g), despite the significant
hearing deficiency at 8 kHz, as shown by ABR evaluations (Fig. 4). This indicates that the 8-kHz
prepulse response was enhanced in *Isl1CKO*, suggesting compensatory neural hyperactivity of
the central auditory system [56, 67]. Thus, the ASR and PPI of startle analyses indicate
abnormalities of the acoustic behavior of *Isl1CKO* mutants.
419
Discussion
Our study shows for the first time that LIM homeodomain transcription factor ISL1 regulates
neuronal development in the cochlea. Using RNA profiling, morphological, and physiological
analyses, we provide evidence that ISL1 coordinates genetic networks affecting the molecular
characteristics of SGNs, their pathfinding abilities, and auditory information processing. The
elimination of *Isl1* in neurons during inner ear development results in a migration defect of
SGNs, disorganized innervation in the cochlea, unsegregated and reduced central axons, and
reduced size of the CN. This neuronal phenotype of *Isl1CKO* was accompanied by hearing
impairment, abnormalities in sound processing in the IC, and aberrant auditory behavior.
ISL1 is critical for the development of multiple tissues, neuronal and non-neuronal cells
[68, 69, 70, 71, 72]. Different aspects of neuronal development depend on ISL1, including specification
of motoneurons [68], sensory neurons [25, 70], axonal growth [73], and axonal pathfinding [74]. During
inner ear development, ISL1 is expressed in both neuronal and sensory precursors [18, 20, 21, 22, 75].
Transgenic modulations of *Isl1* indicate important roles of ISL1 in the maintenance and function
of neurons and hair cells and as a possible contributing factor in neurodegeneration [23, 76, 77, 78].
These studies suggest that ISL1 plays a role in developing neurons and sensory cells, but no
direct evaluation of ISL1 function has been performed. To circumvent the pleiotropic effects of
Isl1 in embryonic development, in this study, we used *Neurod1^Cre* to delete *Isl1* specifically in
the inner ear neurons.
We established that ISL1 is necessary for neuronal differentiation programs in the
cochlea and the functional properties of the auditory system. Our RNA profiling of SGNs
demonstrated transcriptome changes induced by a loss of *Isl1* affecting the molecular
characteristics of neurons and pathfinding abilities, including neurotransmission, the structure of
synapses, neuron migration, axonogenesis, and expression of crucial guidance molecules.
Consistent with the central role of ISL1 in sensory neuron developmental programs [25], regulatory
networks of signaling molecules and transcription factors were affected in *Isl1CKO* neurons,
such as proneural bHLH factors (members of NeuroD, Olig, and Nscl families), LIM-only
(*Lmo2*, *Lmo3*) and LIM homeodomain transcription factors (*Lhx1, Lhx2, Isl2*), transcription
activation complexes for coordination of particular differentiation programs Eyes absent (*Eya4*
and *Eya2*) and Sine oculis (*Six2*), POU homeodomain trans-regulatory factors (*Pou3f2* and
Pou4f2), and FGF signaling molecules (*Fgf10, Fgf11, Fgf13, Fgf14*) and their downstream
targets (*Etv1, Etv4, Etv5*). Interestingly, the transcription factor *Gata3* was downregulated,
suggesting that ISL1 is upstream of the Gata3 transcriptional network of neuronal differentiation
programs [13]. Thus, ISL1 orchestrates a complex gene regulatory network driving multiple aspects
of neuronal differentiation in the cochlea and defining neuronal features.
The most striking morphological features of the neuronal phenotype of *Isl1CKO* are the
aberrant migration and pathfinding of SGNs. A similar migration deficit was reported for *ErbB2*
null mutants [79]; however, the interpretation of the findings is convoluted by the compounded

direct inner ear effects and the effects associated with neural crest-derived Schwann cells.
Additionally, conditional deletion of *Sox10* produced by *Wnt1*[Cre] resulted in abnormal migration
of SGNs similar to the *Isl1CKO* phenotype [80]. Migration defects of SGNs in both *ErbB2* [79] and
Sox10 mutants [80] were attributed to the complete absence of Schwann cells in the entire inner ear
ganglion. However, in our *Isl1CKO* mutant, Sox10$^+$ Schwann cells were found in a similar
density in the spiral ganglion of both mutant and control mice, thus excluding any direct
involvement of glial cells in the migration defects of *Isl1*-deficient neurons. We identified
several genes with altered molecular profiles that are associated with neuronal phenotypes of
projection and migration abnormalities. For instance, misplaced SGNs in the modiolus are in
Ntrk2 and *Ntrk3* neurotrophin receptor mutants [37]. Less severe phenotypes of misplaced SGNs
have been reported for mice lacking *Dcc* [41] or mice with deficient Slit/Robo signaling [40]. All
these reported SGN migration defects are also associated with abnormalities in projections. In
line with the *Isl1CKO* phenotype of deficient neuronal migration and projections, *Isl1*
elimination in SGNs resulted in major transcriptional changes of genes encoding members of all
four classes of guidance molecules and their receptors, growth factors, neurotrophic factor
receptors, and cell adhesion molecules (NCAM family, integrins, cadherins and protocadherins)
among others that are known to participate in molecular networks coordinating neuronal
migratory behaviors, axon guidance, and neural circuit assembly.
More profound disorganization of peripheral processes than defects found in *Isl1CKO* or
both *ErbB2* null [79] and *Sox10* mutants [80] was reported for delayed conditional deletion of *Gata3*
in SGNs [13]. Despite severe disorganization of cochlear wiring of this *Gata3* mutant, central
projections maintained their overall tonotopic organization within the auditory nerve and the CN
[12, 13]. In contrast, we provide compelling evidence that elimination of *Isl1* in SGNs affected the
pathfinding abilities of neurons in the cochlea to form peripheral processes and establish central
projections. Such profound disorganization of peripheral and central projections is known for
Neurod1 mutations [16, 18, 19]. Deletions of *Neurod1* result in miswired, and diminished SGNs, and
loss of tonotopy [16, 19]. Somewhat similar disorganization of central axons with unsegregated
auditory nerve fibers, reduced size of the CN, and missing tonotopic organization of synapsing
branches in the CN subdivisions was found in our *Isl1CKO*. Although both *Neurod1CKO* and
Isl1CKO demonstrated significant hearing loss, in contrast to the reduced sound frequency range
of *Neurod1CKO* [19], responses for the entire measured frequency range were detected in
Isl1CKO. The processing of high acoustic frequencies was broadly comparable between age-
matched controls and *Isl1CKO*, indicating some preservation of peripheral neuronal activity.
Specifically, high frequencies above 28 kHz were comparable between the control and *Isl1CKO*
mice, as shown by ABR measurements and extracellular electrophysiological recordings of IC
neuronal activity (Fig. 4b and 8d, e).
As a result of disorganized primary auditory neurons with derailed central projections,
the characteristics of persistent auditory function in the IC were altered with worsened tuning
capabilities of IC units and their increased spontaneous activity and threshold elevations and
decreased dynamic range. The peripheral deficit in sound encoding results in abnormal auditory
behavior of *Isl1CKO*. Although no significant differences of ABR thresholds at 32 kHz were
observed between *Isl1CKO* and control mice, indicating retained hearing function, the startle
reactions of *Isl1CKO* at 32 kHz were reduced. Plasticity of the startle response is also evident in
the PPI responses of *Isl1CKO* mice, in which a weak prestimulus suppresses the response to a
subsequent startling stimulus. *Isl1CKO* mice demonstrated PPI impairment for the pure tone of 8

kHz, reflecting abnormal sensorimotor gating due to compensatory hyperactivity of the central
auditory system [67, 81].
Additionally, compared to control mice, DPOAE responses of *Isl1CKO* were reduced,
indicating dysfunction of cochlear amplification. The OHCs of the organ of Corti play a central
role in the active enhancement of sound-induced vibration. For given OHCs, amplification only
occurs at a precise frequency, and thus, this mechanism provides a sharpening of the tuning
curve and improves frequency selectivity [82]. Nevertheless, DPOAE analysis showed that some
function was preserved in high-frequency OHCs in the *Isl1CKO* cochlea. Frequencies above 28
kHz are located at the basal half of the mouse cochlea, from the mid-base to the basal end [49],
which correspond to the most preserved distribution of sensory neurons in the area of the spiral
ganglion of *Isl1CKO* mice (Fig. 1, 5). Usually, decreased DPOAE amplitudes indicate loss and
dysfunction of OHCs [76, 82, 83, 84]. Since *Neurod1^{Cre}* is not expressed in sensory cells in the
cochlea, any changes in the development of OHCs represent secondary effects of *Isl1* elimination
in neurons. Accordingly, cochlear amplification deficits in *Isl1CKO* correlated with the reduced
and disorganized innervation of OHCs (Fig. 1). The medial olivocochlear efferents innervate
OHCs from the brainstem, representing a sound-evoked negative feedback loop suppressing
OHC activity [85]. The formation of efferent intraganglionic spiral bundle was altered in *Isl1CKO*,
indicating changes in efferent innervation in the cochlear region. Besides efferents, OHCs are
innervated by the type II SGNs [86, 87]. Although the function of type II SGNs remains obscure, it
is clear that these neurons are involved in auditory nociception [88], and may also constitute the
disputed sensory drive for the olivocochlear efferent reflex [89, 90]. As *Isl1* is expressed in both type
I and type II SGNs during inner ear development [4], characteristics of both neuronal types might
likely be affected in *Isl1CKO*.
The unexpected close-to-normal hearing function and auditory signal processing at high
frequencies suggest some preservation of tonotopic organization in the *Isl1CKO* cochlea for the
propagation of acoustic information. Correspondingly, our 3D images of the *Isl1CKO* cochlea
show that some neurons in the base are in their proper position in the Rosenthal's canal, forming
a segment of the spiral ganglion. These unexpected results may reflect a timeline of sequential
neuronal differentiation from the base to the apex, as the first differentiated neurons are in the
base and the last SGNs undergo terminal mitosis in the apex [91]. As the cochlea is extending,
differentiating neurons migrate along the cochlea to form the spiral ganglion. This process is
completely disrupted in the *Isl1CKO* cochlea, and many neurons are concentrated in a conical
shaped central structure, the modiolus, with only a portion of neurons in the Rosenthal's canal.
Notably, the functional and spatial organization of SGNs may differ in transcription factor
networks required for their differentiation programs. For example, some *Neurod1* lacking
neurons survive, form a rudimental cochlear ganglion, and establish bipolar connections to their
targets in *Neurod1* null mice [11] or the otocyst-deleted *Neurod1* conditional mutant [17]. Consistent
with these findings, neurons forming a segment of the spiral ganglion in the Rosenthal's canal of
Isl1CKO may not require ISL1 for their differentiation and/or the migration mode. Another
possibility is that *Neurod1^{Cre}* may produce a delayed *Isl1* deletion, resulting in a close to-normal
spatial organization of SGNs and auditory system function at high frequencies. A different Cre
driver generating an earlier recombination of *Isl1* in the otocyst might be used to address the
high-frequency hearing phenotype of *Isl1CKO*.
Another secondary effect of *Isl1* elimination in neurons likely contributing to the hearing
deficit of *Isl1CKO* is a shortened cochlea accompanied by the disorganized apical epithelium.
Somewhat similar phenotypes of a shortened cochlea were previously reported for deletion

mutants of *Neurod1* and *Neurog1*, key transcription factors for inner ear neuronal development
[11, 18, 19, 91]. A comparable effect of cochlear length reduction was observed following the loss of
Foxg1 [92] and *Lmx1a* [93, 94]. Although mechanisms affecting the cochlear extension are unknown,
it is clear that these confounding features of the *Isl1CKO* phenotype would consequently impact
mechanical and neural tuning from the base to the apex of the cochlea and the ability to perform
time-frequency processing of sound.
A limitation of our study is that we only assessed the functional role of *Isl1* in the inner
ear neurons. The expression pattern of ISL1 indicates that ISL1 may control the development of
multiple inner ear progenitors [21]. Notably, the elimination of *Isl1* in developing neurons is only a
part of the story. ISL1 is highly expressed in inner ear neurons throughout the development and
in postnatal auditory neurons (Supplementary Figure 2) [4, 18, 21, 22, 95]. Consistent with our results,
the expression profile indicates an important role of ISL1 for the terminal differentiation of
neurons, and for the function and maintenance of differentiated neurons. In contrast, the
expression of ISL1 in sensory precursors is downregulated as hair cell differentiation is initiated
and thus, ISL1 is not detected in the differentiated sensory hair and supporting cells in vestibular
and cochlear epithelia [21]. As the expression of ISL1 precedes the induction of ATOH1 and
sensory cell differentiation, it is conceivable that ISL1 may play a role in the specification of
sensory fate and the regulation of the initial sequential events in sensory precursor development.
The ISL1-based combinatorial complex, together with bHLH factors, may control neuronal and
sensory competencies during inner ear development. We cannot exclude the possibility of
interactions between ISL1$^+$ neuronal and sensory progenitors, and the combined effects of *Isl1*
deletion in sensory and neuronal progenitors. Further studies will be needed to fully uncouple
regulatory mechanisms in inner ear development by targeted eliminations of *Isl1* in sensory
precursors and in both neuronal and sensory precursor populations.
Altogether, our study provides compelling evidence that ISL1 is a critical regulator of
SGN development, affecting neuronal migration, pathfinding abilities to form cochlear wiring,
and central axonal projections. As such, ISL1 represents an essential factor in the regulation of
neuronal differentiation to produce functional neurons in cell-based therapies and stem cell
engineering [96, 97]. Additionally, this unique model contributes to our understanding of how
disorganization of the neuronal periphery affects information processing at higher centers of the
central auditory pathway at the physiological and behavioral levels.
580
581 # Methods
582
Experimental animals
All experiments involving animals were performed according to the Guide for the Care and Use
of Laboratory Animals (National Research Council. Washington, DC. The National Academies
Press, 1996). The design of experiments was approved by the Animal Care and Use Committee
of the Institute of Molecular Genetics, Czech Academy of Sciences. The mice were housed in
12-hour light/dark cycles and were fed *ad libitum*. To generate *Isl1CKO* (the genotype
Neurod1^{Cre};Isl1$^{loxP/loxP}$), we cross-bred floxed *Isl1* (*Isl1$^{loxP/loxP}$*; *Isl1^{tm2Sev}/J*, # 028501, Jackson
Laboratory)[25] and *Neurod1^{Cre}* transgenic mice (Tg(Neurod1-cre)1Able/J, # 028364, Jackson
Laboratory), which were generated by pronuclear injection of the *Neurod1-cre* BAC construct
that carries Cre-sequence downstream of the translational initiation codon ATG of the *Neurod1*
gene [24]. Heterozygous animals, *Neurod1^{Cre};Isl1$^{+/loxP}$* were viable, born in appropriate Mendelian
ratios, and were phenotypically indistinguishable from control (Cre negative) littermate mice. As

control mice, we used mice with the genotype Cre negative, $Isl1^{loxP/loxP}$, and $Isl1^{+/loxP}$. The mouse line $Neurod1^{Cre}$ was also bred with Cre-reporter tdTomato line ($TomatoAi14$, B6.Cg-$Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze}$, # 7914 Jackson Laboratory). Using tdTomato reporter for our analyses, we compared the reporter control-$Ai14$ (genotype: $Neurod1^{Cre};Isl1^{+/loxP};TomatoAi14$) and reporter $Isl1CKO$-$Ai14$ (genotype: $Neurod1^{Cre};Isl1^{loxP/loxP};TomatoAi14$). We performed PCR genotyping on tail DNA. We used both males and females for experiments. Lines are a mixed C57BL/6/sv129 background. Phenotyping and data analyses were performed blind to the genotype of the mice.

Morphological evaluation of the cochlea, the CN, and the IC

Dissected ears were fixed in 4% paraformaldehyde (PFA) in PBS. For vibratome sections, samples were embedded in 4% agarose and sectioned at 80 µm using a Leica VT1000S vibratome. Vibratome sections, whole inner ears, or whole embryos were defatted in 70% ethanol and then rehydrated and blocked with serum, as described previously [20, 77]. Samples were then incubated with primary antibodies at 4ºC for 72 hours. The primary antibodies used were: rabbit anti-Myosin 7a (Myo7a; Proteus BioSciences 25-6790, 1:500), mouse anti-acetylated α-tubulin (tubulin; Sigma-Aldrich T6793, 1:400), rabbit anti-calretinin (Santa Cruz Biotechnology sc-50453, 1:100), goat anti-prestin (Santa Cruz Biotechnology sc-22692, 1:50), rabbit anti-parvalbumin (Abcam ab11427, 1:2000), mouse anti-VGLUT1 (Merck MAB5502, 1:200), rabbit anti-NeuN (Abcam ab177487, 1:500), mouse anti-Isl1 (Developmental Hybridoma Bank 39.3F7 or 39.4D5, 1:130), mouse anti-C-terminal binding protein 2 (CtBP2; BD Biosciences 612044, 1:200), rabbit anti-cleaved Caspase-3 (Cell Signaling Technology 9661, 1:100), goat anti-Neurod1 (Santa Cruz Biotechnology sc-1084, 1:100), anti-β tubulin III (Tuj1; BioLegend 801202, 1:500), and rabbit anti-Sox10 (Abcam ab155279, 1:250). After several PBS washes, secondary antibodies were added and incubated at 4ºC for 24 hours. The secondary antibodies Alexa Fluor® 488 AffiniPure Goat Anti-Mouse IgG (Jackson ImmunoResearch Laboratories 115-545-146), Alexa Fluor® 594 AffiniPure Goat Anti-Rabbit IgG (Jackson ImmunoResearch Laboratories 111-585-144), DyLight488-conjugated AffiniPure Mouse Anti-Goat IgG (Jackson ImmunoResearch Laboratories 205-485-108), Alexa Fluor® 647-conjugated AffiniPure Donkey Anti-Goat IgG (Jackson ImmunoResearch Laboratories 705-605-147), Alexa Fluor® 594-conjugated AffiniPure Donkey Anti-Rabbit IgG (Jackson ImmunoResearch Laboratories 711-585-152), and Alexa Fluor®488-conjugated AffiniPure Donkey Anti-Mouse IgG (Jackson ImmunoResearch Laboratories 715-545-150) were used in 1:400 dilution. Nuclei were stained by Hoechst 33258 (Sigma-Aldrich 861405, 1:2000). Samples were mounted in Aqua-Poly/Mount (Polysciences 18606) or prepared Antifade medium. Images were taken on Zeiss LSM 880 NLO inverted confocal microscope, Nikon CSU-W1 spinning disk confocal microscope, and Carl Zeiss AxioZoomV16 macroscope. NIS-Elements, ImageJ, and ZEN software were used for image processing.

The length of the organ of Corti was measured using the "Measure line" ImageJ plugin. The volumes of the CN and IC were established by analyzing parallel, serial, equally spaced 80 µm coronal vibratome sections through the brain (n = 5 $Isl1CKO$ and n = 5 control mice). The CN and IC areas were determined in each section using ImageJ, and the volume of the organs was calculated. Volumes of organs were adjusted to the body weight. Whole-mount anti-tubulin labeling of the cochlea was used to measure the length and density of the radial fibers. The evaluation of the innervation was done separately for each region of the cochlea: the apex, mid-apex, mid-base, and base. Due to the disorganization of innervation in the apex, we only

evaluated the cochlear base, mid-base, and mid-apex fiber density. The density of the radial fibers was expressed as the percentage of a positive area in the measured area of 152 x 66 μm^2 using the "Threshold" function of ImageJ. The length of the radial fibers was measured from the whole-mount anti-tubulin and Myo7a immunolabeled cochlea confocal images. For each genotype, we measured 3 samples, and in all four regions of the cochlea (the apex, mid-apex, mid-base, and base), we measured the length of the fibers in 3 radial fiber bundles from the IHCs to the IGSB. To compare how many neurons are correctly located in the Rosenthal's canal in *Isl1CKO*, we used NeuN immunolabeled whole-mount cochlea (n = 4 per genotype). Cells were manually counted in 40x magnification images in 5 Z-stacks using the "Cell counter" function of ImageJ; the area of the Rosenthal's canal with tdTomato$^+$ neurons was measured using "Threshold", "Polygon selection" and "Measure" functions of ImageJ to calculate the relative number of neurons for the control and *Isl1CKO* per Rosenthal's canal volume. For ribbon evaluations, IHCs were counted based on the calretinin-stained cell bodies and anti-CtBP2 labeled synapses. CtBP2$^+$ presynaptic ribbon structures were manually counted in 10 individual IHCs in each cochlear region: the apex, mid-apex, mid-base, and base. A total of 6 mutants and 6 control cochleae were evaluated except for 1 apex in the control group, which was lost during immunolabeling. The apoptosis of neurons was quantified using immunohistochemistry with anti-cleaved Caspase-3. Caspase-3 positive cells were manually counted in the inner ear ganglia labeled by the combination of anti-Cre and anti-Neurod1 in the whole-mounted head at 60x magnification at E10.5. For E12.5, Caspase-3$^+$/tdTomato$^+$ neurons were manually counted in the whole-mounted inner ear at 60x magnification. The area of E12.5 inner ear ganglia was measured using the "Threshold and polygon selection" function of ImageJ. For E14.5, the cochlea was sectioned, and Caspase-3$^+$/tdTomato$^+$ neurons were manually counted in all sections at 60x magnification. Three embryos per genotype were used for each embryonic day. All apoptotic cell counting was done using the "Cell counter" function of ImageJ. The density of spherical bushy cells in the VCN, 3 control and 3 *ISL1CKO* 2-month-old mice were used. Bushy cells were labeled by anti-parvalbumin in 80 µm vibratome sections, and manually counted in 2-4 sections/animal in 134.95 x 134.95 µm (63x objective). The size of vestibular sensory epithelia was measured in anti-Myo7a labeled wholemount preparations at P0. Myo7a$^+$ areas were measured in 5 control and 5 *Isl1CKO* mice using "Measure" ImageJ tool. The dorsal root ganglia (DRG) size was evaluated in the sagittal sections of E14.5 embryos with expression of tdTomato reporter. The area of DRG was quantified in 6 embryos per genotype and 2 sections per embryo using the "Threshold and polygon selection" function of ImageJ.

Light-sheet fluorescent microscopy (LFSM) and analysis of images
The cochleae were microdissected from 2 control-*Ai14* and 2 *Isl1CKO-Ai14* mice (postnatal day P0). We used an advanced CUBIC protocol[98] for tissue clearing to enable efficient imaging by light-sheet microscopy. Briefly, the microdissected cochlea was fixed in 4% PFA for 1 hrs, washed with PBS, and incubated in a clearing solution Cubic 1 for 7 days at 37 C. Before immunolabeling, samples were washed in PBT (0.5% Triton-X in PBS) 4 x for 30 min. In addition to *tdTomato* expression, cochlear samples were either immunolabeled using anti-NeuN (a nuclear marker of differentiated neurons) or anti-Myo7a (a hair cell marker) antibodies. Samples were stored before imaging in Cubic 2 at room temperature. Zeiss Lightsheet Z.1 microscope with illumination objective Lightsheet Z.1 5x/0.1 and detection objective Dry objective Lightsheet Z.1 5x/0.16 was used for imaging at the Light Microscopy Core Facility of the Institute of Molecular Genetics of the Czech Academy of Sciences. IMARIS software v8.1.1

(Bitplane AG, CA, USA) was used for image processing.

Isolation of genetically labeled neurons and library construction

Spiral ganglia were micro-dissected in Dulbecco's PBS on ice from E14.5 embryos of either sex

from four litters; each sample contains spiral ganglia from both inner ears of the individual

embryo. Spiral ganglia were incubated in 300 µl of lysis solution (0.05% trypsin, 0.53mM

EDTA in Dulbecco's PBS) in 37 □C, shaking at 900 RPM for 5 min. The lysis was stopped by

adding 600 µl of FACS buffer (10mM EGTA, and 2% FBS in Dulbecco's PBS). After spinning

down the samples at 800 G, 4 □C for 10 min, the supernatant was removed, and cell pellets were

resuspended in 500 µl of ice-cold FACS buffer. Immediately before sorting, cells were passed

through a 50 µm cell sieve (CellTrics^{TM}, Sysmex Amercica Inc.) into a sterile 5 ml polystyrene

round-bottom falcon to remove clusters of cells and kept on ice. TdTomato$^+$ neurons were sorted

using a flow cytometer (BD FACSAria^{TM} Fusion), through a 100 µm nozzle in 20 psi, operated

with BD FACSDiva™ Software (Supplementary Fig. 10). 100 sorted cells were collected into

individual wells of 96-well plate containing 5 µl of lysis buffer of NEB Next single-cell low

input RNA library prep kit for Illumina (New England Biolabs #E6420). Plates were frozen

immediately on dry ice and stored at −80□°C. The total time from euthanasia to cell collection

was ∼3 hrs.

RNAseq-libraries were prepared from 6 samples per each genotype, reporter control-*Ai14*

and *Isl1CKO-Ai14* mutant, and each sample contained 100 tdTomato$^+$ neurons. Following the

manufacturer's instructions, the NEB Next single-cell low input RNA library prep kit for

Illumina was used for cDNA synthesis, amplification, and library generation [99] at the Gene Core

Facility (Institute of Biotechnology CAS, Czechia). Fragment Analyzer assessed the quality of

cDNA libraries. The libraries were sequenced on an Illumina NextSeq 500 next-generation

sequencer. NextSeq 500/550 High Output kit 75 cycles (Illumina #200024906) were processed at

the Genomics and Bioinformatics Core Facility (Institute of Molecular Genetics CAS, Czechia).

Computational analysis of RNAseq data

RNA-Seq reads in FASTQ files were mapped to the mouse genome using STAR [version 2.7.0c

[100]] GRCm38 primary assembly and annotation version M8. The raw data of RNA sequencing

were processed with a standard pipeline. Using cutadapt v1.18[101], the number of reads

(minimum, 32 million; maximum, 73 million) was trimmed by Illumina sequencing adaptor and

of bases with reading quality lower than 20, subsequently reads shorter than 20 bp were filtered

out. TrimmomaticPE version 0.36 [102]. Ribosomal RNA and reads mapping to UniVec database

were filtered out using bowtie v1.2.2. with parameters -S -n 1 and SortMeRNA [103]. A count table

was generated by Rsubread v2.0.1 package using default parameters without counting multi

mapping reads. The raw RNAseq data were deposited at GEO: # GSE182575 study

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE182575). DESeq2 [v1.26.0 [104]] default

parameters were used to normalize data and compare the different groups. Differentially

expressed genes were identified based on an adjusted P-value $p_{adj} < 0.05$, FC > 2, and a base

mean ≥ 50 was applied to identify differentially expressed genes between *Isl1CKO* mutant and

control neurons. The functional annotation of the differentially expressed genes was performed

using GOTermFinder [105].

Enrichment mapping. The enrichment of the functional categories and functional annotation

clustering of the differentially expressed genes was performed using g: Profiler [106] input using

version e104_eg51_p15_3922dba with g: SCS multiple testing correction methods applying a

significance threshold of 0.05. In contrast, no electronic GO annotations were used. Only Biological Processes (BP) data underwent further processing. Complete query details are available in Query info tabs in Supplementary Data 2. The resulting GEM and combined GMT files were loaded into Cytoscape [107] plugin "EnrichmentMap" [108] using 0.01 FDR q-value cutoff to generate a network. Edge cutoff was set to 0.6, and nodes were filtered by gs_size<1800. Five GO terms forming solitary nodes, or a pair of nodes, were excluded (listed in Supplementary Data 2). Further adjustments were made in yFiles Layout Algorithms, Legend Creator (Cytoscape plugins), and Inkscape (Inkscape Project, 2020).

Quantitative real-time PCR

Total RNA was isolated from both inner ears of the embryo at E14.5 using TRI Reagent (Sigma-Aldrich T9424). We used 8 embryos for *Isl1CKO* and 7 embryos for the control group from three litters. RNA from both inner ears of one embryo represented one sample. RNA samples (0.5 µg of total RNA) were processed and analyzed as previously described [18, 77]. Briefly, following RT, quantitative qPCR was performed with initial activation at 95 °C for 120 s, followed by 40 cycles at 95 °C for 15 s, 60 °C for 30 s, and 72 °C for 30 s using the CFX384™ Real-Time PCR Detection System (Bio-Rad Laboratories). The primer sequences (*Lhx1, Lhx2, Cdh7, Nhlh2, Ntrk2, Ntrk3, Ntng1*) were obtained from (pga.mgh.harvard.edu/primerbank/) or designed using Primer3 tool (*Dcc, Epha5, Gata3, Prdm8, Robo2, Slit2, Tbx3, Uncx*). The primer sequences are listed in Supplementary Table 2. Relative mRNA expression was calculated using the $-\Delta\Delta Cq$ method with *Hprt1* as a reference gene. GraphPad Prism software was used for statistical analysis.

Lipophilic Dye Tracing

We studied the innervation pattern in whole or dissected ears using lipophilic dye tracing in aldehyde-fixed tissues as previously described [109]. At least three mutants and similar numbers of control littermates of both sexes were used for each evaluation. Filter strips loaded with NeuroVue colored lipophilic dyes were inserted into the cochlear apex, base, and vestibular end-organ utricle. After allowing appropriate time for diffusion of the lipophilic tracer (between 48-120 hours), we prepared the ears as whole mounts in glycerol on a glass slide, using appropriate spacers to avoid distortion, and imaged them using a Leica SP8 confocal microscope. Images were compiled into plates to show the most pertinent details using Corel Draw. Only general image modifications such as contrast, or brightness adjustments were used to enhance the visual appeal without affecting the scientific content.

Hearing function evaluation

Auditory brainstem response (ABR) and distortion product otoacoustic emissions (DPOAEs) tests were carried out on mice under general anesthesia with 35 mg/kg ketamine (Calypso 50 mg/ml) and 6 mg/kg xylazine (Xylapan 20 mg/ml) in saline to give an application volume of 7 ml/kg body weight via subcutaneous injection, maintained on a temperature-regulated blanket in a soundproof room.

Distortion product otoacoustic emissions. For DPOAE recording were tested *Isl1CKO* (n = 10) and control mice (n = 14). Cubic (2 F1–F2) distortion product otoacoustic emissions over an F2 frequency range from 4 to 39 kHz were recorded with a low-noise microphone system (Etymotic probe ER-10B+, Etymotic Research). Acoustic stimuli (ratio F2/F1 = 1.21, F1 and F2 primary tone levels of L1/L2 = 70/60 dB) were presented to the ear canal with two custom-made

piezoelectric stimulators connected to the probe with 10-cm-long silastic tubes. The signal from
the microphone was analyzed by the TDT System III (RP2 processor, sampling rate 100 kHz)
using custom-made MATLAB software. DPOAEs were recorded in the animals' ears at
individual frequencies over the frequency range 4–39 kHz with a resolution of ten points per
octave. All the experiments and analyses were done with no information on the genotype.
Auditory brainstem response. For auditory brainstem response (ABR) recording (n = 10 *Isl1CKO*
and n = 11 control mice), an active electrode was placed subcutaneously on the vertex and
ground and reference electrodes in the neck muscles. Responses to tone bursts (3 ms duration, 1
ms rise/fall times, frequencies of 2, 4, 8, 16, 32, and 40 kHz) and clicks of different intensity
were recorded. Acoustic stimuli were conveyed to the animal in free-field conditions via a two-
way loudspeaker system (Selenium 6W4P woofer and RAAL70-20 tweeter) placed 70 cm in
front of the animal's head. The signal was processed with a TDT System III Pentusa Base Station
and analyzed using BioSig™ software. The response threshold to each frequency was
determined as the minimal tone intensity that still evoked a noticeable potential peak in the
expected time window of the recorded signal. The amplitude and latency of ABR peaks I-V were
determined using BioSig software (Tucker Davis Technologies). Central compensation of
neuronal responsiveness (central gain) was calculated using ABR wave IV to I amplitudes.
796
Extracellular recording of the neuronal activity in the IC
For extracellular recording in the IC, we evaluated *Isl1CKO* (n = 12, 624 units) and control mice
(n = 8, 395 units). The surgery and extracellular recording in the IC were performed in mice
anesthetized with 35 mg/kg ketamine (Calypsol 50 mg/ml) and 6 mg/kg xylazine (Xylapan 20
mg/ml) in saline via subcutaneous injection. Approximately every hour, supplement
subcutaneous injections of one-half of the original dose of the anesthetics were administered to
keep a sufficient level of anesthesia, judged by a positive pedal and palpebral (toe-pinch) reflex
and movement of the whiskers. Respiratory rate, and heart rate, were monitored. An incision was
made through the skull's skin to access the IC, and underlying muscles were retracted to expose
the dorsal skull. A holder was glued to the skull, and small holes were drilled over both ICs.
Neuronal activity (multiple units) in the IC was recorded using a 16-channel, single shank probe
(NeuroNexus Technologies) with 50 or 100 µm between the electrode spots. The obtained signal
from the electrode was amplified 10000 times, band-pass filtered over the range of 300 Hz to 10
kHz, and processed by a TDT System III (Tucker Davis Technologies) using an RX5-2 Pentusa
Base Station. Individual spikes from the recorded signal were isolated online based on amplitude
discrimination and analyzed with BrainWare software (v. 8.12, Jan Schnupp, Oxford University).
Subsequent discrimination of spikes from the recorded data and their sorting according to the
amplitudes of the first positive and negative peaks were performed off-line and was used to sort
action potentials (spikes) among single units. The recorded data were processed and analyzed
using custom software based on MATLAB. The stimulation signals were generated using a TDT
System III with the RP 2.1 Enhanced Real-Time Processor. Acoustic stimuli were delivered in
free-field conditions via a two-driver loudspeaker system (Selenium 6W4P woofer and
RAAL70-20 tweeter) placed 70 cm in front of the animal's head.
Frequency-intensity mapping: To determine the neuronal receptive fields, pure tones (frequency
2 - 40 kHz with 1/8 octave step, 60 ms duration, 5 ms rise/fall times, various intensity with 5dB
step) were presented in a random order, each stimulus appearing three times. A discrete matrix
corresponding to the response magnitude evoked by each of the frequency-intensity
combinations was thereby obtained, smoothed using cubic spline interpolation, and used for

extraction of the basic parameters: the excitatory response threshold (the lowest stimulus
intensity that excited the neuron, measured in dB SPL), the characteristic frequency (CF) – the
frequency with the minimal response threshold, measured in Hz, and the bandwidth of the
excitatory area 20 dB above the excitatory threshold, expressed by quality factor Q (Q =
CF/bandwidth).
Rate intensity function of the IC neurons: Neuronal responses to broadband noise (BBN) bursts
of variable intensity (10 dB steps, 50 repetitions) were used to construct the rate intensity
function (RIF). A 100% scale was assigned to the neuron's total range of response amplitudes,
0% corresponding to spontaneous activity and 100% corresponding to its maximum response
magnitude [110]. The two points of interest are R10 and R90, which correspond to 10 and 90% of
this scale, respectively. R10, describing the starting point of the RIF's rise, was taken as the BBN
response threshold. RIFs were further used to evaluate the following parameters: the dynamic
range (DR) of the RIF: DR = S90–S10); and the maximum response magnitude. Spontaneous
activity of the IC neurons was determined at the 0dB SPL BBN stimulation.
Temporal properties of the IC neurons: We used trains of five clicks at an intensity of 70 dB
SPL for control and 80 dB SPL for *Isl1CKO* mice with various inter-click intervals (100, 50, 30,
20, and 15 ms). We calculated the vector strength (VS) values and the Rayleigh statistics for
each spike pattern; only responses with at least 5.991 were significantly considered phase-
locking (Zhou and Merzenich, 2008). The VS quantifies how well the individual spikes are
synchronized (phase-locked) with a periodic signal.
845
Behavioral tests of motor coordination and balance
All testing was carried out during the light cycle and after a minimum 30-min acclimatization.
Air-righting reflex. The vestibular function was evaluated by the ability of the mice to right
themselves in the air when held supine and dropped onto a soft surface from a height of 50 cm[77].
Rotarod. Using the rotarod apparatus (Rota Rod 47600, Ugo Basile), the time (latency) to fall off
the rod rotating under continuous acceleration was recorded. During the acclimatization period,
mice with their heads in the direction of rotation were loaded on the rotarod at an initial speed of
4 rpm. This speed was maintained for 2 min and, if mice fell during this period, they were placed
back on the rotarod. The drum was slowly accelerated from 4 to 40 rpm for a maximum of 300 s
for each trial for the experimental measurements. The latency to fall off the rotarod within 300 s
was recorded. If the mouse clung to the grip of the rotating rod and failed to resume normal
performance for three consecutive revolutions, the sensor was manually triggered. Mice were
tested in three consecutive trials in one session per day with a 15-min rest period between each
trial.
860
Behavioral acoustic tests
Isl1CKO (n = 10) and control (n = 7) mice were used at 2-3 months of age. All behavioral tests
were performed in a sound-attenuated chamber (Coulbourn Habitest, model E10-21) located in a
soundproof room. Each mouse was placed in a wire mesh cage on a motion-sensitive platform
inside the box during the testing. The mouse's reflex movements to sound stimuli were detected
and transformed to a voltage signal by the load-cell response sensing platform. An amplified
voltage signal was acquired and processed using a TDT system 3 with a Real-Time Processor RP
2 (Tucker Davis Technologies, Alachua, Fl) and custom-made software in a Matlab
environment. The startle responses were evaluated in 100 ms windows, beginning at the onset of
the startling stimulus. The magnitude of the response was measured as the maximal peak-to-peak

amplitude of transient voltage occurring in the response window. Acoustic stimuli were
generated by the TDT system (Real-Time Processor RP 2), amplified and presented via a
loudspeaker (SEAS, 29AF/W), and placed inside the chamber above the animal. Stimulus
presentation and data acquisition were controlled by a custom-made application in a Matlab
environment. Calibration of the apparatus was performed for frequencies between 4 kHz and 32
kHz by a 1/4-inch Brüel & Kjaer 4939 microphone connected to a Brüel & Kjaer ZC 0020
preamplifier and a B&K 2231 sound level meter. During the calibration, the calibrating
microphone was positioned at the animal's head in the test cage.
Acoustic startle reflex (ASR) (a transient motor response to an intense, unexpected stimulus) was
used to indicate the behavioral responsiveness to sound stimuli**.** The ASRs to 8, 16, and 32 kHz
tone pips and BBN bursts (50 ms duration, 3 ms rise/fall times, varying intensity levels) were
recorded. Each test session contained: a baseline trial (–10 dB SPL stimulus intensity) and 13
startle stimuli of different intensities (50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 115, and 120-
dB SPL). The inter-trial interval varied from 15 to 50 s.
In *the prepulse inhibition (PPI)* procedure, 3 different trial types were used: a baseline trial
without any stimulus, an acoustic startle pulse alone (white noise at 110 dB SPL, 50 ms, 3 ms
rise/fall times), and a combination of the prepulse and startle pulse. The inter-stimulus interval
between the prepulse and the startle stimulus was set to 50 ms; each trial type was presented
three times. The inter-trial gap was randomized and varied from 15 to 50 s. The efficacy of the
PPI of ASR was expressed as an ASR ratio in percentage, e.g., 100% corresponds to the
amplitude of ASR without prepulse; smaller values of ASR ratio indicate stronger PPI. As a
prepulse, we used either BBN bursts or tone pips (50 ms duration, 3 ms rise/fall time) at
frequencies of 8 and 32 kHz at increasing intensities. It is expected that in the presence of
prepulse, the amplitude of the following startle response decreases.
895
Experimental design and statistical analyses
All comparisons were made between animals with the same genetic background, typically
littermates, and we used male and female mice. The number of samples (n) for each comparison
can be found in the individual method descriptions and are given in the corresponding figures.
Phenotyping and data analysis was performed blind to the genotype of the mice. All values are
presented either as the mean ± standard deviation (SD) or standard error of the mean (SEM). For
statistical analysis, GraphPad Prism software was used. To assess differences in the mean, one-
way or two-way ANOVA with Bonferroni's multiple comparison test, multiple *t*-tests with
Holm-Sidak comparison method, repeated measures ANOVA, multiple *t*-tests, and unpaired
two-tailed *t*-tests were employed. Significance was determined as $P < 0.05$ (*), $P < 0.01$ (**), $P <$
0.001 (***) or $P < 0.0001$ (****). Complete results of the statistical analyses are included in the
figure legends.
908
Ethics approval and consent to participate
Experiments were carried out following the animal welfare guidelines 2010/63/EC of the
European Communities Council Directive, agreeing with the Guide for the Care and Use of
Laboratory Animals (National Research Council. Washington, DC. The National Academies
Press, 1996). The design of experiments was approved by the Animal Care and Use Committee
of the Institute of Molecular Genetics, Czech Academy of Sciences (protocol # 104/2019).
915
916 Consent for publication

917   Not applicable.
918
919   Data availability
920   All data generated or analyzed during this study are included in this published article, and
921   Supplementary information. RNA-seq data are deposited in the publicly available Gene
922   Expression Omnibus (GEO) repository (# GSE182575 study). Source data are provided with this
923   paper.
924

925   Competing interests
926   The authors declare that they have no competing interests.
927

928   Funding
929   This research was supported by the Czech Science Foundation (20-06927S to GP), by the
930   institutional support of the Czech Academy of Sciences (RVO: 86652036 to GP), and NIH/NIA
931   (R01 AG060504, DC016099, AG051443 to BF and ENY).
932

Authors' contributions
GP, JS, BF designed and supervised the experiments. IF, KP, RB, MT, SV, MD, and BF
performed experiments and analyzed the data. JC, PH, and VF prepared samples and generated
3D images. OS, SB, and LV carried RNAseq analyses. IF and KP prepared the first draft of the
manuscript. GP and BF wrote the manuscript, ENY and JS reviewed the manuscript. All authors
read and approved the final manuscript.
939

Acknowledgments
A. Pavlinek (King's College London) for editing the MS. We thank M. Anderova for providing
TomatoAi14 reporter line. We acknowledge Imaging Methods Core Facility at BIOCEV
supported by the MEYS CR (Large RI Project LM2018129 Czech-BioImaging) and ERDF
(project No. CZ.02.1.01/0.0/0.0/18_046/0016045) for its support with obtaining imaging data
and FACS experiments presented in this paper; Biocev GeneCore Facility for its support with
gene expression/transcriptome analyses, and Biocev Animal facility (LM 2018126 Czech Centre
for Phenogenomics by MEYS OP RDE CZ.02.1.01/0.0/0.0/18_046/0015861 CCP Infrastructure
Upgrade II by MEYS and ESIF).
949

950   **References**
951   1.   Muniak MA, Connelly CJ, Suthakar K, Milinkeviciute G, Ayeni FE, Ryugo DK. Central
952   Projections of Spiral Ganglion Neurons. In: *The Primary Auditory Neurons of the Mammalian*
953   *Cochlea*). Springer (2016).
954
955   2.   Rubel EW, Fritzsch B. Auditory system development: primary auditory neurons and their targets.
956   *Annu Rev Neurosci* **25**, 51-101 (2002).
957
958   3.   Shrestha BR, Chia C, Wu L, Kujawa SG, Liberman MC, Goodrich LV. Sensory Neuron Diversity
959   in the Inner Ear Is Shaped by Activity. *Cell* **174**, 1229-1246 e1217 (2018).
960

4. Petitpre C, *et al.* Neuronal heterogeneity and stereotyped connectivity in the auditory afferent
system. *Nature communications* **9**, 3691 (2018).

5. Sun S, *et al.* Hair Cell Mechanotransduction Regulates Spontaneous Activity and Spiral Ganglion
Subtype Specification in the Auditory System. *Cell* **174**, 1247-1263 e1215 (2018).

6. Kandler K, Clause A, Noh J. Tonotopic reorganization of developing auditory brainstem circuits.
Nat Neurosci **12**, 711-717 (2009).

7. Di Bonito M, Studer M. Cellular and Molecular Underpinnings of Neuronal Assembly in the
Central Auditory System during Mouse Development. *Front Neural Circuits* **11**, 18 (2017).

8. Lopez-Poveda EA. Olivocochlear Efferents in Animals and Humans: From Anatomy to Clinical
Relevance. *Front Neurol* **9**, 197 (2018).

9. Fritzsch B, Elliott KL. Evolution and Development of the Inner Ear Efferent System:
Transforming a Motor Neuron Population to Connect to the Most Unusual Motor Protein via
Ancient Nicotinic Receptors. *Front Cell Neurosci* **11**, 114 (2017).

10. Ma Q, Anderson DJ, Fritzsch B. Neurogenin 1 null mutant ears develop fewer, morphologically
normal hair cells in smaller sensory epithelia devoid of innervation. *J Assoc Res Otolaryngol* **1**,
129-143 (2000).

11. Kim W-Y, *et al.* NeuroD-null mice are deaf due to a severe loss of the inner ear sensory neurons
during development. *Development* **128**, 417-426 (2001).

12. Duncan JS, Fritzsch B. Continued expression of GATA3 is necessary for cochlear neurosensory
development. *PloS one* **8**, e62046 (2013).

13. Appler JM, Lu CC, Druckenbrod NR, Yu WM, Koundakjian EJ, Goodrich LV. Gata3 is a critical
regulator of cochlear wiring. *J Neurosci* **33**, 3679-3691 (2013).

14. Huang EJ, Liu W, Fritzsch B, Bianchi LM, Reichardt LF, Xiang M. Brn3a is a transcriptional
regulator of soma size, target field innervation and axon pathfinding of inner ear sensory neurons.
Development **128**, 2421-2432 (2001).

15. Liu M, *et al.* Essential role of BETA2/NeuroD1 in development of the vestibular and auditory
systems. *Genes & development* **14**, 2839-2854 (2000).

16. Jahan I, Kersigo J, Pan N, Fritzsch B. Neurod1 regulates survival and formation of connections in
mouse ear and brain. *Cell Tissue Res* **341**, 95-110 (2010).

17.   Filova I, Bohuslavova R, Tavakoli M, Yamoah EN, Fritzsch B, Pavlinkova G. Early Deletion of
Neurod1 Alters Neuronal Lineage Potential and Diminishes Neurogenesis in the Inner Ear.
Frontiers in Cell and Developmental Biology **10**, (2022).

1006
1007   18.   Filova I*, et al.* Combined Atoh1 and Neurod1 Deletion Reveals Autonomous Growth of Auditory
1008   Nerve Fibers. *Mol Neurobiol* **57**, 5307-5323 (2020).

1009
1010   19.   Macova I*, et al.* Neurod1 Is Essential for the Primary Tonotopic Organization and Related
1011   Auditory Information Processing in the Midbrain. *J Neurosci* **39**, 984-1004 (2019).

1012
1013   20.   Dvorakova M*, et al.* Incomplete and delayed Sox2 deletion defines residual ear neurosensory
1014   development and maintenance. *Sci Rep* **6**, 38253 (2016).

1015
1016   21.   Radde-Gallwitz K, Pan L, Gan L, Lin X, Segil N, Chen P. Expression of Islet1 marks the sensory
1017   and neuronal lineages in the mammalian inner ear. *J Comp Neurol* **477**, 412-421 (2004).

1018
1019   22.   Deng M, Yang H, Xie X, Liang G, Gan L. Comparative expression analysis of POU4F1,
1020   POU4F2 and ISL1 in developing mouse cochleovestibular ganglion neurons. *Gene Expr Patterns*
1021   **15**, 31-37 (2014).

1022
1023   23.   Chumak T*, et al.* Overexpression of Isl1 under the Pax2 Promoter, Leads to Impaired Sound
1024   Processing and Increased Inhibition in the Inferior Colliculus. *Int J Mol Sci* **22**, (2021).

1025
1026   24.   Li HJ, Kapoor A, Giel-Moloney M, Rindi G, Leiter AB. Notch signaling differentially regulates
1027   the cell fate of early endocrine precursor cells and their maturing descendants in the mouse
1028   pancreas and intestine. *Dev Biol* **371**, 156-169 (2012).

1029
1030   25.   Sun Y, Dykes IM, Liang X, Eng SR, Evans SM, Turner EE. A central role for Islet1 in sensory
1031   neuron development linking sensory and spinal gene regulatory programs. *Nat Neurosci* **11**,
1032   1283-1293 (2008).

1033
1034   26.   Jahan I, Pan N, Kersigo J, Fritzsch B. Neurod1 suppresses hair cell differentiation in ear ganglia
1035   and regulates hair cell subtype development in the cochlea. *PLoS One* **5**, e11661 (2010).

1036
1037   27.   Li C*, et al.* Comprehensive transcriptome analysis of cochlear spiral ganglion neurons at multiple
1038   ages. *Elife* **9**, (2020).

1039
1040   28.   Stoeckli ET. Understanding axon guidance: are we nearly there yet? *Development* **145**, (2018).

1041
1042   29.   Fritzsch B, Kersigo J, Yang T, Jahan I, Pan N. Neurotrophic Factor Function During Ear
1043   Development: Expression Changes Define Critical Phases for Neuronal Viability. In: *The*
1044   *Primary Auditory Neurons of the Mammalian Cochlea* (eds Dabdoub A, Fritzsch B, Popper AN,
1045   Fay RR). Springer New York (2016).

1046

1047   30.   Dennis DJ, Han S, Schuurmans C. bHLH transcription factors in neural development, disease,
1048   and reprogramming. *Brain Res* **1705**, 48-65 (2019).

1049

1050   31.   Yang T, Kersigo J, Wu S, Fritzsch B, Bassuk AG. Prickle1 regulates neurite outgrowth of apical
1051   spiral ganglion neurons but not hair cell polarity in the murine cochlea. *PloS one* **12**, e0183773
1052   (2017).

1053

1054   32.   Bok J, Zenczak C, Hwang CH, Wu DK. Auditory ganglion source of Sonic hedgehog regulates
1055   timing of cell cycle exit and differentiation of mammalian cochlear hair cells. *Proc Natl Acad Sci*
1056   *U S A* **110**, 13869-13874 (2013).

1057

1058   33.   Pauley S, Wright TJ, Pirvola U, Ornitz D, Beisel K, Fritzsch B. Expression and function of
1059   FGF10 in mammalian inner ear development. *Dev Dyn* **227**, 203-215 (2003).

1060

1061   34.   Yu Y*, et al.* Sensorineural Hearing Loss and Mitochondrial Apoptosis of Cochlear Spiral
1062   Ganglion Neurons in Fibroblast Growth Factor 13 Knockout Mice. *Frontiers in Cellular*
1063   *Neuroscience* **15**, (2021).

1064

1065   35.   Defourny J*, et al.* Ephrin-A5/EphA4 signalling controls specific afferent targeting to cochlear hair
1066   cells. *Nature communications* **4**, 1438 (2013).

1067

1068   36.   Coate TM, Raft S, Zhao X, Ryan AK, Crenshaw EB, 3rd, Kelley MW. Otic mesenchyme cells
1069   regulate spiral ganglion axon fasciculation through a Pou3f4/EphA4 signaling pathway. *Neuron*
1070   **73**, 49-63 (2012).

1071

1072   37.   Fritzsch B, Barbacid M, Silos-Santiago I. The combined effects of trkB and trkC mutations on the
1073   innervation of the inner ear. *Int J Dev Neurosci* **16**, 493-505 (1998).

1074

1075   38.   Brors D*, et al.* Spiral ganglion outgrowth and hearing development in p75-deficient mice. *Audiol*
1076   *Neurootol* **13**, 388-395 (2008).

1077

1078   39.   Coate TM, Spita NA, Zhang KD, Isgrig KT, Kelley MW. Neuropilin-2/Semaphorin-3F-mediated
1079   repulsion promotes inner hair cell innervation by spiral ganglion neurons. *Elife* **4**, (2015).

1080

1081   40.   Wang SZ*, et al.* Slit/Robo signaling mediates spatial positioning of spiral ganglion neurons
1082   during development of cochlear innervation. *J Neurosci* **33**, 12242-12254 (2013).

1083

1084   41.   Kim YJ, Wang SZ, Tymanskyj S, Ma L, Tao HW, Zhang LI. Dcc Mediates Functional Assembly
1085   of Peripheral Auditory Circuits. *Sci Rep* **6**, 23799 (2016).

1086

1087   42.   Frisen J, Yates PA, McLaughlin T, Friedman GC, O'Leary DD, Barbacid M. Ephrin-A5 (AL-
1088   1/RAGS) is essential for proper retinal axon guidance and topographic mapping in the
1089   mammalian visual system. *Neuron* **20**, 235-243 (1998).

1090
1091   43.   Fujita Y, Nakanishi T, Ueno M, Itohara S, Yamashita T. Netrin-G1 Regulates Microglial
1092   Accumulation along Axons and Supports the Survival of Layer V Neurons in the Postnatal Mouse
1093   Brain. *Cell Rep* **31**, 107580 (2020).

1094
1095   44.   Ross SE*, et al.* Bhlhb5 and Prdm8 form a repressor complex involved in neuronal circuit
1096   assembly. *Neuron* **73**, 292-303 (2012).

1097
1098   45.   Palmesino E*, et al.* Foxp1 and lhx1 coordinate motor neuron migration with axon trajectory
1099   choice by gating Reelin signalling. *PLoS Biol* **8**, e1000446 (2010).

1100
1101   46.   Marcos-Mondejar P, Peregrin S, Li JY, Carlsson L, Tole S, Lopez-Bendito G. The lhx2
1102   transcription factor controls thalamocortical axonal guidance by specific regulation of robo1 and
1103   robo2 receptors. *J Neurosci* **32**, 4372-4385 (2012).

1104
1105   47.   Fettiplace R, Hackney CM. The sensory and motor roles of auditory hair cells. *Nature reviews
1106   Neuroscience* **7**, 19-29 (2006).

1107
1108   48.   Neely S, Kim DO. *Cochlear models incorporating active processes*. Springer New York (2007).

1109
1110   49.   Muller M, von Hunerbein K, Hoidis S, Smolders JW. A physiological place-frequency map of the
1111   cochlea in the CBA/J mouse. *Hear Res* **202**, 63-73 (2005).

1112
1113   50.   Zhou X, Jen PH, Seburn KL, Frankel WN, Zheng QY. Auditory brainstem responses in 10 inbred
1114   strains of mice. *Brain Res* **1091**, 16-26 (2006).

1115
1116   51.   Chumak T*, et al.* BDNF in Lower Brain Parts Modifies Auditory Fiber Activity to Gain Fidelity
1117   but Increases the Risk for Generation of Central Noise After Injury. *Molecular Neurobiology* **53**,
1118   5607-5627 (2016).

1119
1120   52.   Land R, Burghard A, Kral A. The contribution of inferior colliculus activity to the auditory
1121   brainstem response (ABR) in mice. *Hearing Research* **341**, 109-118 (2016).

1122
1123   53.   Liu Y*, et al.* Hearing loss is an early biomarker in APP/PS1 Alzheimer's disease mice. *Neurosci
1124   Lett* **717**, (2020).

1125
1126   54.   Melcher JR, Kiang NYS. Generators of the brainstem auditory evoked potential in cat III:
1127   identified cell populations. *Hearing Research* **93**, 52-71 (1996).

1128

1129   55.   Wolter S, *et al.* GC-B Deficient Mice With Axon Bifurcation Loss Exhibit Compromised
1130   Auditory Processing. *Frontiers in Neural Circuits* **12**, (2018).

1131
1132   56.   Chambers AR, *et al.* Central Gain Restores Auditory Processing following Near-Complete
1133   Cochlear Denervation. *Neuron* **89**, 867-879 (2016).

1134
1135   57.   Harris JA, Rubel EW. Afferent regulation of neuron number in the cochlear nucleus: cellular and
1136   molecular analyses of a critical period. *Hear Res* **216-217**, 127-137 (2006).

1137
1138   58.   Lauer AM, Connelly CJ, Graham H, Ryugo DK. Morphological characterization of bushy cells
1139   and their inputs in the laboratory mouse (Mus musculus) anteroventral cochlear nucleus. *PLoS*
1140   *One* **8**, e73308 (2013).

1141
1142   59.   Karmakar K, *et al.* Hox2 Genes Are Required for Tonotopic Map Precision and Sound
1143   Discrimination in the Mouse Auditory Brainstem. *Cell Rep* **18**, 185-197 (2017).

1144
1145   60.   Fujiyama T, *et al.* Inhibitory and excitatory subtypes of cochlear nucleus neurons are defined by
1146   distinct bHLH transcription factors, Ptf1a and Atoh1. *Development* **136**, 2049-2058 (2009).

1147
1148   61.   Gruters KG, Groh JM. Sounds and beyond: multisensory and other non-auditory signals in the
1149   inferior colliculus. *Front Neural Circuits* **6**, 96 (2012).

1150
1151   62.   Yeomans JS, Frankland PW. The acoustic startle reflex: neurons and connections. *Brain*
1152   *Research Reviews* **21**, 301-314 (1995).

1153
1154   63.   Koch M. The neurobiology of startle. *Progress in neurobiology* **59**, 107-128 (1999).

1155
1156   64.   Lee Y, Lopez DE, Meloni EG, Davis M. A primary acoustic startle pathway: obligatory role of
1157   cochlear root neurons and the nucleus reticularis pontis caudalis. *J Neurosci* **16**, 3775-3789
1158   (1996).

1159
1160   65.   Swerdlow NR, Geyer MA, Braff DL. Neural circuit regulation of prepulse inhibition of startle in
1161   the rat: current knowledge and future challenges. *Psychopharmacology* **156**, 194-215 (2001).

1162
1163   66.   Fitch RH, Threlkeld SW, McClure MM, Peiffer AM. Use of a modified prepulse inhibition
1164   paradigm to assess complex auditory discrimination in rodents. *Brain research bulletin* **76**, 1-7
1165   (2008).

1166
1167   67.   Hickox AE, Liberman MC. Is noise-induced cochlear neuropathy key to the generation of
1168   hyperacusis or tinnitus? *J Neurophysiol* **111**, 552-564 (2014).

1169

1170 68. Pfaff SL, Mendelsohn M, Stewart CL, Edlund T, Jessell TM. Requirement for LIM homeobox
1171   gene Isl1 in motor neuron generation reveals a motor neuron-dependent step in interneuron
1172   differentiation. *Cell* **84**, 309-320 (1996).

1173
1174 69. Yang L*, et al.* Isl1Cre reveals a common Bmp pathway in heart and limb development.
1175   *Development* **133**, 1575-1585 (2006).

1176
1177 70. Huber K, Narasimhan P, Shtukmaster S, Pfeifer D, Evans SM, Sun Y. The LIM-Homeodomain
1178   transcription factor Islet-1 is required for the development of sympathetic neurons and adrenal
1179   chromaffin cells. *Dev Biol* **380**, 286-298 (2013).

1180
1181 71. Ahlgren U, Pfaff SL, Jessell TM, Edlund T, Edlund H. Independent requirement for ISL1 in
1182   formation of pancreatic mesenchyme and islet cells. *Nature* **385**, 257-260 (1997).

1183
1184 72. Cai CL*, et al.* Isl1 identifies a cardiac progenitor population that proliferates prior to
1185   differentiation and contributes a majority of cells to the heart. *Dev Cell* **5**, 877-889 (2003).

1186
1187 73. Liang X*, et al.* Isl1 is required for multiple aspects of motor neuron development. *Molecular and
1188   cellular neurosciences* **47**, 215-222 (2011).

1189
1190 74. Kania A, Jessell TM. Topographic motor projections in the limb imposed by LIM homeodomain
1191   protein regulation of ephrin-A:EphA interactions. *Neuron* **38**, 581-596 (2003).

1192
1193 75. Yamashita T*, et al.* High-resolution transcriptional dissection of in vivo Atoh1-mediated hair cell
1194   conversion in mature cochleae identifies Isl1 as a co-reprogramming factor. *PLoS Genet* **14**,
1195   e1007552 (2018).

1196
1197 76. Chumak T*, et al.* Deterioration of the Medial Olivocochlear Efferent System Accelerates Age-
1198   Related Hearing Loss in Pax2-Isl1 Transgenic Mice. *Mol Neurobiol* **53**, 2368-2383 (2016).

1199
1200 77. Bohuslavova R*, et al.* Pax2-Islet1 Transgenic Mice Are Hyperactive and Have Altered Cerebellar
1201   Foliation. *Mol Neurobiol* **54**, 1352-1368 (2017).

1202
1203 78. Huang M, Kantardzhieva A, Scheffer D, Liberman MC, Chen ZY. Hair cell overexpression of
1204   Islet1 reduces age-related and noise-induced hearing loss. *J Neurosci* **33**, 15086-15094 (2013).

1205
1206 79. Morris JK*, et al.* A disorganized innervation of the inner ear persists in the absence of ErbB2.
1207   *Brain Res* **1091**, 186-199 (2006).

1208
1209 80. Mao Y, Reiprich S, Wegner M, Fritzsch B. Targeted deletion of Sox10 by Wnt1-cre defects
1210   neuronal migration and projection in the mouse inner ear. *PLoS One* **9**, e94580 (2014).

1211

1212 81. Geyer MA, McIlwain KL, Paylor R. Mouse genetic models for prepulse inhibition: an early
1213   review. *Mol Psychiatry* **7**, 1039-1053 (2002).

1214
1215 82. LeMasurier M, Gillespie PG. Hair-cell mechanotransduction and cochlear amplification. *Neuron*
1216   **48**, 403-415 (2005).

1217
1218 83. Dallos P*, et al.* Prestin-based outer hair cell motility is necessary for mammalian cochlear
1219   amplification. *Neuron* **58**, 333-339 (2008).

1220
1221 84. Herranen A*, et al.* Deficiency of the ER-stress-regulator MANF triggers progressive outer hair
1222   cell death and hearing loss. *Cell death & disease* **11**, 100 (2020).

1223
1224 85. Liberman LD, Liberman MC. Cochlear Efferent Innervation Is Sparse in Humans and Decreases
1225   with Age. *J Neurosci* **39**, 9560-9569 (2019).

1226
1227 86. Hafidi A. Peripherin-like immunoreactivity in type II spiral ganglion cell body and projections.
1228   *Brain Res* **805**, 181-190 (1998).

1229
1230 87. Elliott KL*, et al.* Developmental Changes in Peripherin-eGFP Expression in Spiral Ganglion
1231   Neurons. *Front Cell Neurosci* **15**, 678113 (2021).

1232
1233 88. Zhang KD, Coate TM. Recent advances in the development and function of type II spiral
1234   ganglion neurons in the mammalian inner ear. *Seminars in cell & developmental biology* **65**, 80-
1235   87 (2017).

1236
1237 89. Froud KE*, et al.* Type II spiral ganglion afferent neurons drive medial olivocochlear reflex
1238   suppression of the cochlear amplifier. *Nature communications* **6**, 7115 (2015).

1239
1240 90. Maison S, Liberman LD, Liberman MC. Type II Cochlear Ganglion Neurons Do Not Drive the
1241   Olivocochlear Reflex: Re-Examination of the Cochlear Phenotype in Peripherin Knock-Out
1242   Mice. *eNeuro* **3**, (2016).

1243
1244 91. Matei V*, et al.* Smaller inner ear sensory epithelia in Neurog 1 null mice are related to earlier hair
1245   cell cycle exit. *Dev Dyn* **234**, 633-650 (2005).

1246
1247 92. Pauley S, Lai E, Fritzsch B. Foxg1 is required for morphogenesis and histogenesis of the
1248   mammalian inner ear. *Dev Dyn* **235**, 2470-2482 (2006).

1249
1250 93. Nichols DH, Pauley S, Jahan I, Beisel KW, Millen KJ, Fritzsch B. Lmx1a is required for
1251   segregation of sensory epithelia and normal ear histogenesis and morphogenesis. *Cell Tissue Res*
1252   **334**, 339-358 (2008).

1253

1254   94.   Nichols DH, *et al.* Interaction with ectopic cochlear crista sensory epithelium disrupts basal
1255   cochlear sensory epithelium development in Lmx1a mutant mice. *Cell Tissue Res* **380**, 435-448
1256   (2020).

1257
1258   95.   Dvorakova M, Macova I, Bohuslavova R, Anderova M, Fritzsch B, Pavlinkova G. Early ear
1259   neuronal development, but not olfactory or lens development, can proceed without SOX2. *Dev*
1260   *Biol* **457**, 43-56 (2020).

1261
1262   96.   Pavlinkova G. Molecular Aspects of the Development and Function of Auditory Neurons. *Int J*
1263   *Mol Sci* **22**, (2020).

1264
1265   97.   Zine A, Messat Y, Fritzsch B. A human induced pluripotent stem cell-based modular platform to
1266   challenge sensorineural hearing loss. *Stem Cells*, (2021).

1267
1268   98.   Susaki EA, Tainaka K, Perrin D, Yukinaga H, Kuno A, Ueda HR. Advanced CUBIC protocols
1269   for whole-brain and whole-body clearing and imaging. *Nat Protoc* **10**, 1709-1727 (2015).

1270
1271   99.   Mamanova L, Miao Z, Jinat A, Ellis P, Shirley L, Teichmann SA. High-throughput full-length
1272   single-cell RNA-seq automation. *Nat Protoc* **16**, 2886-2915 (2021).

1273
1274   100.   Dobin A, *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

1275
1276   101.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011* **17**,
1277   3 (2011).

1278
1279   102.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
1280   *Bioinformatics* **30**, 2114-2120 (2014).

1281
1282   103.   Kopylova E, Noe L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in
1283   metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).

1284
1285   104.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
1286   data with DESeq2. *Genome Biol* **15**, 550 (2014).

1287
1288   105.   Boyle EI, *et al.* GO::TermFinder--open source software for accessing Gene Ontology information
1289   and finding significantly enriched Gene Ontology terms associated with a list of genes.
1290   *Bioinformatics* **20**, 3710-3715 (2004).

1291
1292   106.   Raudvere U, *et al.* g:Profiler: a web server for functional enrichment analysis and conversions of
1293   gene lists (2019 update). *Nucleic acids research* **47**, W191-W198 (2019).

1294
1295   107.   Shannon P, *et al.* Cytoscape: a software environment for integrated models of biomolecular
1296   interaction networks. *Genome Res* **13**, 2498-2504 (2003).

1297
1298   108.   Merico D, Isserlin R, Stueker O, Emili A, Bader GD. Enrichment map: a network-based method
1299   for gene-set enrichment visualization and interpretation. *PLoS One* **5**, e13984 (2010).

1300
1301   109.   Fritzsch B, Duncan JS, Kersigo J, Gray B, Elliott KL. Neuroanatomical Tracing Techniques in
1302   the Ear: History, State of the Art, and Future Developments. In: *B. Sokolowski, Ed: Auditory and*
1303   *Vestibular Research: Methods and Protocols*). Springer Science+Business Media New York
1304   (2016).

1305
1306   110.   Bures Z, Grécová J, Popelár J, Syka J. Noise exposure during early development impairs the
1307   processing of sound intensity in adult rats. *The European journal of neuroscience* **32**, 155-164
1308   (2010).

1309

1310   **Figure legends**

1311

Figure 1. ***Isl1* deletion results in abnormalities in cochlear innervation and the formation of**
the spiral ganglion. (**a**) Diagram of the organization of the cochlea. Spiral ganglion neurons
(SGNs) are organized tonotopically from the apex to the base of the cochlea. High-frequency
sounds maximally stimulate the base of the cochlea, whereas the largest response to low-
frequency sounds occurs in the cochlear apex. (**b**) The top view diagram onto the sensory
epithelium shows outer hair cells (OHCs), inner hair cells (IHCs), and SGNs. Type I neurons
extend radial fibers (RF) toward IHCs (5-30 neurons innervate one IHC), and type II neurons
(representing 5% of SGNs) receive input from OHCs. The central processes of SGNs relay
acoustic information to the brain. Efferent axons from the superior olivary complex, forming the
intraganglionic spiral bundle (IGSB), innervate OHCs. (**c, d, g, h**) Representative images of
whole-mount immunolabeling of the base and apex with anti-Myo7a (hair cell marker) and anti-
α-tubulin (nerve fibers) show reduced and disorganized RF and missing or altered efferent fibers
forming IGSB (arrowhead) in *Isl1CKO* compared to control. (**e, f, i, j**) Higher-magnification
images show three rows of OHCs and one row of IHCs forming the organ of Corti (OC) and
connected to SGNs by RF. Abnormalities in innervation, including larger gaps between radial
fiber bundles, crisscrossing fibers (arrow), and fiber bundles bypassing OC (arrowhead in j), are
noticeable in *Isl1CKO*. (**k-m**) The shape of the spiral ganglion (SG) is shown in the basal and
apical half of the whole-mounted cochlea of control-*Ai14* reporter mice at P0. TdTomato[+]
neurons form a spiral located in the Rosenthal's canal and parallel to the OC (anti-Myo7a labeled
hair cells), nerve fibers are labeled with anti-tubulin. (**n, o**) In *Isl1CKO*, only some tdTomato[+]
neurons are in the Rosenthal's canal (delineated by white dotted lines), but many neurons are in
the centre of the cochlea, a conical-shaped structure, the modiolus (arrowhead). (**q**) The apex of
Isl1CKO contains a reduced number of neurons and disorganized innervation. (**p, r**) Higher-
magnification images show some neurons entangled with radial fibers near the OC. (**s, t**) The
vibratome sections of the cochlea labeled by anti-NeuN (a nuclear marker of differentiated
neurons) and anti-tubulin (nerve fibers) show the unusual position of cochlear neurons in the
modiolus, and neurons entwined in central axons in *Isl1CKO* compared to control (white lines
delineate a normal position of the SG). (**u**) NeuN[+] neurons were counted in the Rosenthal's canal
(n = 4 mice per genotype), and (**w**) fiber density was quantified in the base, mid-base, and mid-

apex (n = 3 mice per genotype). Error bars represent mean ± SD; unpaired *t*-test. Scale bars: 100 μm (c, d, g, h, k-t), 50 μm (e, f, i, j).

Figure 2. **The length of the organ of Corti is shortened, the apical end sensory epithelium is disorganized, and presynaptic ribbons are reduced in *Isl1CKO*.** (**a**) The organ of Corti of *Isl1CKO* is shorter than control, as shown by anti-Myo7a labeling of hair cells in the cochlea at P0. (**b**) The length of the organ of Corti was measured at postnatal day P0 and in 2-month-old mice. Error bars represent mean ± SD; unpaired *t*-test. (**c**) Immunohistochemistry for prestin (a marker for OHCs) and calretinin (a marker for IHCs) shows a comparable cytoarchitecture of the organ of Corti in the cochlear mid-base of controls and *Isl1CKO* but disorganized rows of OHCs in the apical end with multiple OHC rows and ectopic calretinin$^+$ IHCs among OHCs (arrowheads) in *Isl1CKO* at P35. (**d**) Higher-magnification images show a detail of anti-tubulin labeled innervation with fibers forming three parallel outer spiral bundles (arrows) that innervate multiple OHCs and turn toward the base in the control cochlea. Guiding defects in the extension of these fibers to OHCs are obvious in *Isl1CKO* with some fibers randomly turned toward the apex (arrowheads); note the disorganization of radial fiber bundles in *Isl1CKO*. (**e**) Representative images of whole-mount immunolabeling of the cochlear mid-base with calretinin (IHCs) and anti-CtBP2 (presynaptic ribbons) of 2-month-old mice. (**f**) CtBP2$^+$ synapses were counted in 10 IHCs per each cochlear region (n = 6 mice per genotype). Box plots indicate median (middle line), 25th, 75th percentile (box) and min to max (whiskers); multiple *t*-tests, ****P < 0.000001. HS, Hoechst nuclear staining; IHC, inner hair cell; OHC, outer hair cell. Scale bars: 100 μm (a), 50 μm (c), 20 μm (d), 10 μm (e).

Figure 3. **ISL1-mediated transcription signature in cochlear neurons. (a)** Image of the whole-mount cochlea with genetically labeled tdTomato neurons in the spiral ganglion at E14.5 [Hoechst nuclear staining (blue), Scale bar: 100 μm]; and workflow depicts microdissection, dissociation, FACS sorting of single tdTomato$^+$ spiral ganglion neurons for a bulk of 100 cells RNAseq analysis. **(b)** The volcano plot shows the change in protein-coding gene expression levels in the *Isl1CKO* compared to control spiral ganglion neurons (adjusted p-value < 0.05, and fold change > 2 cutoff values). The complete list of identified 332 down- and 318 up-differentially expressed genes is in Supplementary Data 1. **(c)** Enrichment map of down- and upregulated gene ontology (GO) sets visualized by the network. Each node represents a GO term; edges depict shared genes between nodes. Node size represents a number of genes of the mouse genome per the GO term, and node fill color represents a GO term significance. Each GO set cluster was assigned with representative keywords; a list of GO sets is available in Supplementary Data 2).

Figure 4. **Hearing impairment is detected in *Isl1CKO* mice.** (**a**) Distortion product otoacoustic emissions (DPOAEs) show significantly reduced levels in the low and middle-frequency range (4-24 kHz). Data are expressed as mean ± SEM, two-way ANOVA with Bonferroni post-hoc test, *P < 0.05, **P < 0.01, ***P < 0.001 ****P < 0.0001. (**b**) The average auditory brainstem response (ABR) thresholds of *Isl1CKO* and control mice are analyzed by click-evoked ABR. Data are expressed as mean ± SD, two-way ANOVA with Bonferroni post-hoc test, **P < 0.01, ****P < 0.0001. (**c**) Averaged ABR response curves evoked by an 80 dB SPL click; (**e**) by a 90 dB SPL pure tone of 32 kHz frequency; and (**g**) by a 90 dB SPL pure tone of 8 kHz frequency are presented. (**d, f, h**) Averaged individual ABR wave amplitudes are shown for the

1387 corresponding peaks. Data are expressed as mean ± SEM, two-way ANOVA with Bonferroni
1388 post-hoc test, **P < 0.01 ***P < 0.001, ****P < 0.0001.
1389
Figure 5. **Neurons in the *Isl1CKO* inner ear are abnormally distributed. (a-i)** Microdissected
cochleae of reporter control-*Ai14* and *Isl1CKO-Ai14* mice were cleared (CUBIC protocol),
immunolabeled, imaged, and reconstructed in 3D using light-sheet fluorescence microscopy (see
Supplementary Videos S1-S4). (**a**) In the control cochlea, tdTomato$^+$ SGNs form a coil of the
spiral ganglion (SG) in the Rosenthal's canal and anti-Myo7a labeled hair cells show the spiral
shape of the organ of Corti. Parts of the vestibular ganglion (VG) with tdTomato$^+$ neurons and
the saccule with Myo7a$^+$ hair cells are shown. (**b, c**) Anti-NeuN (a nuclear marker of
differentiated neurons) co-labeled tdTomato$^+$ neurons form SG and VG in the control cochlea;
the modiolus contains the central processes, forming the auditory nerve (AN). (**d-f**) In *Isl1CKO-*
Ai14, tdTomato$^+$ neurons are mainly located in the conical central part of the cochlea, the
modiolus, in contrast to the spiral of the organ of Corti with Myo7a$^+$ hair cells. Some *Isl1CKO*
neurons form a segment of the spiral ganglion in the Rosenthal's canal (RC) in the mid-base
region (arrowhead), shown in detail (**f**), the RC area is highlighted by dotted lines. (**g-h**)
Abnormal and uneven distribution of cochlear tdTomato$^+$ neurons is shown by anti-NeuN co-
labeling. Many neurons are misplaced in the modiolus, but some neurons are in their close-
normal position in RC (arrowhead). (**j**) Schematic drawing of the mouse cochlea shows the
cochlear place-frequency map for four cochlear regions termed the apex (corresponding to the
best encoding frequencies ∼8 kHz at the distance from the basal end of 82 %), mid-apex (∼19
kHz at 49% length), mid-base (∼32 kHz at 32% length), and basal turn (∼62 kHz at 9% length)
[49]. (**k**) Whole mount images of the P10 inner ear show distribution of tdTomato labeled neurons
in superior (sVG) and inferior vestibular ganglia (iVG), and SG in the control-*Ai14*. (**l**) In the
Isl1CKO-Ai14 mutant, the VG is fused and enlarged, spatial segregation between VG and SG is
diminished, forming an aberrant "spiro-vestibular" ganglion, and SG has lost its spiral shape.
SGNs, spiral ganglion neurons. Scale bars: 200 μm, except for 100 μm (b, i).
1414
Figure 6. ***Isl1CKO* neurons project unsegregated and disorganized central projections to the
cochlear nucleus. (a)** The schematic diagram visualizes dye tracing from the inner ear and its
connections to the auditory brainstem, using insertions of differently colored NeuroVue dyes into
the vestibular end-organ (utricle, magenta), cochlear base (red), and apex (green). Axonal
projections from cochlear neurons to the cochlear nucleus (CN) bifurcate with one branch
synapsing in the dorsal (DCN) and the other innervating the anteroventral CN (AVCN). (**b**)
Images of individual colors of the separate channels and a merged image show distinct and
spatially restricted bundles of neuronal fibers of the auditory nerve (AN) projecting to the CN
and the vestibular ganglion (VG) in controls, labeled by dye applications into the apex (green),
base (red), and utricle (magenta). The only mixed bundle (yellow) is inner ear efferents (IEE). (**c**)
In *Isl1CKO*, the segregation of central axons is lost, as fibers labeled from the apex (green) and
base (red) are mainly overlapping in the AN (yellow fibers), and neurons labeled by dyes applied
into the cochlear base and apex are mixed with VG neurons to form an aberrant ganglion, the
"spiro-vestibular" ganglion. Note that fibers labeled from the apex (green) form an unusual fiber
loop around the ganglion, and no IEE are recognizable in *Isl1CKO*. (**d**) The tonotopic
organization of the CN subdivisions is shown with low-frequency afferents labeled from the
apex (green) and high frequency from the base (red) and organized as parallel fibers in
isofrequency bands in controls at E16.5. (**e**) In *Isl1CKO*, cochlear afferents enter AVCN as a

single bundle instead of forming separate basal and apical projections. The branch synapsing in
the AVCN is reduced and disorganized. Arrowheads indicate overlapping fibers. The DCN
branch is represented by just a few fibers in *Isl1CKO* (inset in **e**). **(f, g)** Images of individual
colors of the separate channels represent projections labeled from the apex (green) and base
(red), shown as a merged image in e. **(h, i)** The confocal section shows segregated cochlear
afferents from the base and apex, and vestibular afferents extend toward the control's lateral
vestibular nucleus (LVN). In contrast, cochlear afferents of *Isl1CKO* overlap (yellow fibers), and
some basal afferents extend toward the CN, and some project toward the LVN and loop back to
the CN. (**j**) Graphic summary of aberrant location and projections of auditory neurons in
Isl1CKO. In the control, the auditory system is tonotopically organized, including neurons of the
spiral ganglion, the auditory nerve, and the CN subdivisions. In *Isl1CKO*, excluding some
portion of neurons in the base, neurons are located outside of Rosenthal's canal. Some neurons
projecting to the cochlea are mixed with vestibular neurons. The central axons from the apex and
base are not segregated in the AN and AVCN. Projections entering the DCN are diminished in
Isl1CKO. AC, anterior crista; D, dorsal; HC, horizontal crista; P, posterior axes; PVCN,
posteroventral CN; PC, posterior crista; RB, restiform body; r1- r6, rhombomere 1 - 6; U, utricle;
S, saccule; VN, vestibular nerve. Scale bars: 100 µm.
1450
Figure 7. **The cochlear nucleus of *Isl1CKO* is reduced. (a, b)** Coronal sections of a brain
(immunostained with anti-NeuN) of adult controls and *Isl1CKO*, show the DCN and VCN; the
dotted line indicates the boundaries of the CN. **(c-f)** Representative images of immunolabeled
brain sections with anti-parvalbumin to label the spherical bushy cell soma and anti-VGlut1 to
label auditory-nerve endbulbs of Held around the bushy cells. Higher-magnification images
show the distribution of parvalbumin$^+$ neurons and the presence of VGlut1$^+$ auditory nerve
synaptic terminals. (**g**) Quantification of the adult CN volume, adjusted to body weight (n = 5
mice per genotype) and (**h**) parvalbumin$^+$ neurons per µm^2 of the AVCN (n = 3 mice per
genotype, 2-4 sections per mouse). Data are expressed as mean ± SD, unpaired *t-test* (***P <
0.001; ns, not significant). D, dorsal; M, medial axis; HS, Hoechst nuclear staining; DCN, dorsal
cochlear nucleus; VCN, ventral cochlear nucleus. Scale bars: 200 µm (a-d); 20 µm (e,f).
1462
Figure 8 **Characteristics of IC neurons (units) are affected in *Isl1CKO* mice. (a)**
Immunostaining of coronal brain sections for NeuN and nuclear staining Hoechst (HS). Scale
bars, 200 µm. **(b)** Quantification of the adult control volume and *Isl1CKO* inferior colliculus (IC)
adjusted to body weight. Data are expressed as mean ± SD (n = 5 mice per genotype), unpaired *t-
test* (ns, not significant). (**c**) Representative examples of tuning curves recorded in the IC display
impairments in tuning properties with broad and irregular receptive fields in *Isl1CKO* compared
to control mice. (**d**) Scatter diagram shows the distribution of the excitatory thresholds of IC
neurons in dependency on the characteristic frequency (CF) recorded in control and *Isl1CKO*
mice. Note, no IC neurons are recorded below 6 kHz in *Isl1CKO* mutants. (**e**) Excitatory
thresholds of the IC neurons at different CFs are shown as averages in 0.3-octave bins in control
and *Isl1CKO* mice. Data are expressed as mean ± SEM. Two-way ANOVA with Bonferroni post
hoc test. **P < 0.01, ***P < 0.001, ****P < 0.0001. (**f**) Comparison of the rate intensity function
parameters between control (n = 8) and *Isl1CKO* mice (n = 12): broadband noise (BBN)
threshold, dynamic range, spontaneous activity, and maximum response magnitude. Violin plots
indicate median (middle line), 25th, and 75th percentile (dotted lines), unpaired *t*-test, ****P <
0.0001. (**g**) The sharpness of the neuronal tuning expressed by quality factor Q_{20} (the ratio

between the CF and bandwidth at 20 dB above the minimum threshold) averaged in 0.3-octave
bins is decreased in *Isl1CKO*. (**h**) Synchronization of units with click trains. Vector strength
computed for different inter-click intervals. Data are expressed as mean ± SD, two-way ANOVA
with Bonferroni post-hoc test, ****$P < 0.0001$.
1483
Figure 9. **The acoustic startle reflex (ASR) and prepulse inhibition (PPI) responses are**
altered in the *Isl1CKO* mutant. (**a**) ASR thresholds for broadband noise (BBN) bursts and tone
pips at 8, 16, and 32 kHz in control and *Isl1CKO* mice are shown. Holm-Sidak method multiple
comparison *t*-test. *$P < 0.05$, **$P < 0.01$. (**b**) Significantly increased ASR latency to BBN is
found in *Isl1CKO* compared to control mice; unpaired *t*-test, **$P < 0.01$. (**c**) Amplitude-intensity
ASR functions for BBN stimulation and (**d**) for tone pips of 8 kHz and (**e**) 32 kHz at different dB
SPL intensities in control and *Isl1CKO* mice are shown. (**f**) Efficacy of the BBN, (**g**) 8 kHz, and
(**h**) 32 kHz tone prepulse intensity on the relative ASR amplitudes are displayed; ASR ratio = 1
corresponds to the ASR amplitude without a prepulse (uninhibited ASR). Two-way ANOVA
with Bonferroni post hoc tests. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. All data
are expressed as mean ± SEM.

a Frequency high ← base, radial fibers, SGNs, apical end, low

b Central axons — afferents: Type II neurons, SGNs, IGSB, IHC, OHC; Type I neurons, radial fibers; efferents

c Control P0, base, OC, RF, IGSB — Myo7a, Tubulin

d apical end, IGSB, apex

e OHC, IHC

f OHC, IHC

g Isl1CKO P0, base, IGSB — Myo7a, Tubulin

h apical end, apex

i OHC, IHC

j OHC, IHC

k Control-Ai14 P0, base, OC — Myo7a, Tubulin, tdTomato

l SG, Central axons, RF

m apical end, apex

s Control P0, Tubulin, NeuN, Central axons, SG

n Isl1CKO-Ai14 P0, base — Myo7a, Tubulin, tdTomato

o modiolus

p

t Isl1CKO P0, Tubulin, NeuN, Central axons, modiolus

q Isl1CKO-Ai14 P0, apex, apical end — Myo7a, Tubulin, tdTomato

r apex

u Neurons located in the Rosenthal's canal
Number of neurons — P < 0.0001 (control/Isl1CKO Base), P = 0.0005 (control/Isl1CKO Apex)

w Density of radial fibers
% — P = 0.0082, control, Isl1CKO

a Control P0 / *Isl1CKO* P0 — Myo7a

b Length of the cochlea at P0 (P < 0.0001); Length of the adult cochlea (P < 0.0001) — mm; control / *Isl1CKO*

c Control mid-base P35 / Control apical end P35 / *Isl1CKO* mid-base P35 / *Isl1CKO* apical end P35 — Calretinin prestin HS; IHC OHC

d Control base P0 / Control apex P0 / *Isl1CKO* base P0 / *Isl1CKO* apex P0 — Tubulin; OHC3 OHC2 OHC1 IHC

e Control mid-base 2M / *Isl1CKO* mid-base 2M — Calretinin CtBP2 prestin HS

f CtBP2⁺synapses/IHC; Control / *Isl1CKO*; Base, Mid-base, Mid-apex, Apex; ****

a

E14.5 cochlea
Neurod1^Cre;TomatoAi14

E14.5

SG dissection
Cell dissociation

FACS sorting
100 Tomato⁺ cells

Library prep

RNAseq

Bioinformatic
analyses

Validation
(RT-qPCR)

b

332 down-regulated genes   318 up-regulated genes

Brinp3

Helt

Fgf10

Epha5

Lhx2

Sytl4

Ntrk2

Nhlh2

Lhx1

Uncx

Runx1

Pcdh18

Pcdh8

Shh

Sema3e

$-\log_{10}p_{adj}$

$\log_2FC$

c

DOWN-regulated gene set enrichment map

UP-regulated gene set enrichment map

other organ
development

neuron
migration

neuron
development

axonogenesis
axon guidance
axon development

transmembrane
ion transport

synaptic
signalling

cell
adhesion

synaptic
organization

synaptic
organization

synaptic
signalling

neuron development

Enrichment map legend
Node: GO term
Edge: gene overlap
Node fill color: adj. p-value

0.01   2.5E-03   7E-16

DOWN

0.01   5E-04   3.5E-08

UP

Node size: gene number

50 1025 2000

Control-*Ai14* P0

a tdTomato Myo7a
apex
organ of Corti
SG
VG
saccule
base

b tdTomato NeuN
apex
central projections
VG
AN
base

c tdTomato NeuN
apex
AN
VG
base

Isl1CKO-Ai14 P0

d tdTomato Myo7a
apex
modiolus
saccule
base

e tdTomato Myo7a
modiolus
saccule
base

f View of the mid-base tdTomato Myo7a
RC
base

Isl1CKO-Ai14 P0

g tdTomato NeuN
apex
modiolus
base

h tdTomato NeuN
base

i View of the mid-base tdTomato NeuN
RC

j
8 kHz
apex
20 kHz
mid-apex
mid-base
32 kHz
61.8 kHz
base

Control-*Ai14* P10
k apex SG tdTomato
base
iVG
sVG

Isl1CKO-Ai14 P10
l apex SGNs tdTomato
modiolus
"Spiro-vestibular" ganglion

a

inner ear
utricle
vestibular ganglion
VG
base
apex
auditory nerve (AN)

cochlear nucleus
AVCN
DCN
vestibular nerve(VN)

b Control E18.5

IEE
VG
AN
D
P

IEE
AN
NeuroVue-apex

IEE
AN
NeuroVue-base

VG
NeuroVue-utricle

c

Isl1CKO E18.5

AN

AN

AN

d Control E16.5

AVCN
DCN
VN
AN
NeuroVue-base
NeuroVue-apex
NeuroVue-utricle

h Control E18.5

LVN
RB
PVCN
VN
base
apex
Apex
Base
Utricle

i Isl1CKO E18.5

LVN
RB
PVCN
base
base
Apex
Base

e Isl1CKO E16.5

DCN
AVCN
DCN
VN
AN
Apex
Base
Utricle

f Isl1CKO E16.5

Apex

g Isl1CKO E16.5

Base

j Control

r1 r2 r3 r4 r5 r6
AVCN
DCN
HC
AC
U
VN
AN
VG
base
S PC
SGN
apex

Isl1CKO

AVCN
DCN
HC
AC
U
VN
AN
VG
base
S PC
SGN
apex

NeuN HS   VGlut1 Parvalbumin HS

a DCN / VCN — Control P35

b DCN / VCN — *Isl1CKO* P35

D / M

c VCN

d DCN / VCN

e

f

g

CN volume/BW (mm³/g)

***

Control   *Isl1CKO*

h

Parvalbumin⁺ neurons/µm²

ns

Control   *Isl1CKO*