Assessing the effectiveness of honey bee pollinators for cultivated

blueberries in South Africa

Keanu Martin[1*], Bruce Anderson[1], Corneile Minnaar[1], ML de Jager[1]

[1]Department of Botany and Zoology, Stellenbosch University, Matieland 7602, Cape Town,

Western Cape, South Africa

*Corresponding author: keanum@sun.ac.za

9 **Abstract**

Globally, agricultural crops are often extremely dependent on insect pollination. Blueberries

are an example of such a crop and owing to their proposed health benefits they are grown

around the world, including locations where their native bumble bee pollinators do not

occur. In the absence of bumble bees, blueberry pollination in South Africa and many other

parts of the non-native, commercial range is performed primarily by honey bees. Despite

this, the effectiveness of honey bee pollination on blueberries remains understudied. This

study determined the effect of honey bee pollination on the fruit characteristics of five

blueberry varieties that are extensively planted in South Africa. For each variety two metrics

were calculated: 1) the benefit of bees - a comparison of fruit yields after exposure to honey

bees and fruit yields after honey bee exclusion, 2) the pollination deficit - a comparison

between yields after hand pollination and yields after exposure to honey bee pollinators.

Honey bee pollination consistently resulted in improved yields, although the magnitude of

this improvement was dependant on the variety considered. Similarly, the pollination deficit

also varied considerably across varieties and while some varieties appeared to perform close

to maximum potential, others appeared to perform relatively poorly under honey bee

pollination. This study demonstrates that honey bees are functional pollinators of

blueberries in areas where native blueberry pollinators are absent. However, in such areas,

it is important that special focus be given to selecting blueberry varieties that perform well

with honey bees as their sole pollinator. Further research is necessary to determine how the

pollination deficit of blueberry varieties can be decreased as well as how to increase the

effectiveness of honey bee pollination.

Keywords: Honey bee, blueberries, pollination, agriculture, pollination ecosystem service,

yield

1. Introduction

Insect pollination is vital for numerous crops, not only to set fruit but also to increase fruit yield (Klein et al., 2007). Subsequently, this ecosystem service is said to be valued at €153 billion per year (Gallai et al., 2009). Although honey bees are globally the most widely used commercially available pollinator (Garibaldi et al., 2013), they are not always the most efficient pollinators and can be outperformed by wild insects (Garibaldi et al., 2013). In a comprehensive study using 41 crops, Garibaldi et al. (2013) found that wild insects were often more effective crop pollinators than honey bees. On average, fruit set doubled after increasing wild insect visitation compared to equivalent increases in honey bee visitation.

One crop where honey bees are thought to be less effective than other pollinators is blueberries. Wild and commercial blueberries are extremely reliant on insect pollination (Aras et al., 1996; MacKenzie, 1997; Dogterom et al., 2000; Fulton et al., 2015; Martin et al., 2021) and are typically pollinated by large bees. These include blueberry bees (Apidae) (Cane and Payne, 1993; Campbell et al., 2018), bumble bees (Apidae) (Mackenzie and Eickwort, 1996; Stubbs and Drummond, 2001; Javorek et al., 2002; Tuell et al., 2009; Campbell et al., 2017), and mining bees (Andrenidae) (Mackenzie and Eickwort, 1996; Javorek et al., 2002). These bees are thought to be effective pollinators of blueberry flowers due to their ability to buzz pollinate which causes the poricidal anthers to release their pollen (Cane and Payne, 1993). In comparison, honey bees cannot buzz pollinate and are less effective at releasing pollen from these specialised anthers. Consequently, bumble bees deposit about two-thirds more pollen per visit than honey bees (Javorek et al., 2002). However, recently Hoffmann et al. (2018) revealed that honey bees that foraged on blueberries had blueberry pollen on all body parts with the majority of blueberry pollen

being found on their legs and tarsi. These pollen-laden legs and tarsi regularly came into

contact with blueberry stigmas during landing, grooming, and foraging and have the ability

to effectively pollinate blueberry flowers. Moreover, Martin et al. (2021) showed that honey

bees effectively pollinated the common South African blueberry variety, Ventura,

significantly increasing both blueberry mass and diameter, as well as decreased ripening

time.

The perceived inefficiency of honey bees as pollinators has resulted in pressure to import

native blueberry pollinators, such as bumble bees. This is extremely problematic as

introducing bumble bees into novel areas has the potential to disrupt local ecosystems. For

example, introduced bumble bees can outcompete native pollinators, damage flowers, and

rob nectar (Morales et al., 2013; Schmid-Hempel et al., 2014; Aizen et al., 2018).

Consequently, it is important to quantify the benefits of honey bees for different blueberry

varieties so that objective decisions can be made about the perceived needs of importing

novel pollinators. This may also enable us to farm more intelligently with varieties that do

better under honey bee pollination.

We aim to determine the effects of honey bee pollination on the fruit characteristics of five

blueberry varieties that are extensively planted in South Africa. Particularly, we compare the

benefits (in terms of fruit yield i.e. fruit mass, fruit diameter, percentage fruit set, and

adjusted fruit mass) of honey bee pollination with situations when pollinators are absent.

Although honey bees are often viewed as ineffective pollinators of blueberries, they still

move pollen between flowers and often increase blueberry fruit production (Aras et al.,

1996b; Mackenzie, 1997; Sampson and Cane, 2000; Hoffman et al., 2018). Accordingly, we

hypothesize that honey bees considerably increase fruit yield (i.e. fruit mass, diameter,

percentage fruit set, adjusted fruit mass) and decrease fruit developmental period,

compared to fruit produced in the absence of pollinators. Additionally, we establish if the

pollination environment could be optimised so that fruit yield is improved relative to the

present conditions where honey bees are the sole pollinators. Despite honey bees moving

pollen between blueberry flowers (Hoffman et al., 2018), we predict considerable room for

improvement, with hand pollination resulting in substantially larger fruit yields than honey

bee pollination.

2. Methods

An experimental plot on Backsberg blueberry farm was used as the study site (Western

Cape, South Africa, 33°48'30.7"S 18°54'09.8"E). Farmlands surround the experimental plot

and there is also a nearby (2.98km) nature reserve consisting of Swartland shale

renosterveld and fynbos. Hives were stocked at 15 hives per hectare which correlates to

typical commercial blueberry farm hive density (Isaacs and Kirk, 2010; Gibbs et al., 2016;

Hoffman et al., 2018). Hive densities on blueberries are often an order of magnitude greater

than on crops such as apricots, peaches, and nectarines where one honey bee hive per

hectare provides sufficient pollination (Allsopp et al., 2008). This experiment comprised of

three treatments: pollinators excluded, open honey bee pollination, and hand pollination.

We determined the effects of honey bee pollination on blueberry production and also

whether there was scope to improve pollination conditions by comparing fruit yield (fruit

mass, fruit set, and developmental period) among these three treatments. Varieties used

were Emerald (*Vaccinium corymbosum* hybrid), Eureka (*Vaccinium corymbosum* hybrid),

Snowchaser (*Vaccinium sp. hybrid*), Suziblue (*Vaccinium corymbosum* hybrid), and Twilight

(*Vaccinium corymbosum* hybrid). Pollination combinations were repeated three times per

plant and across nine plants, resulting in 27 pollination combination replicates. Moreover,

we minimized any extrinsic factors which potentially influence fruit development by

conducting our experiments across varieties in an experimental plot of only 50 m X 115 m.

Fig 1: Hypothetical figure showing how the use of three treatments can be used to determine the benefits of

honey bee pollination as well as how yields may potentially be improved upon (originally produced by Martin

et al. 2021).

To determine pollination deficits and benefits of bees to the five varieties, we followed the

methods of Martin et al. (2021). Below we provide a brief description.

We determined the benefit of bees by comparing blueberry fruit production as a result of

honey bee pollination to blueberry fruit production in the absence of honey bees (pollinator

exclusion. Pollinators were denied access to flowers by covering 27 individual virgin flowers

with a fine mesh bag. Once flowers had wilted, these bags were removed to allow for

normal fruit maturation. We estimated the fruit yield through honey bee pollination by

exposing (unbagging) 27 flowers to bee pollinators. For the duration of the time spent at the

experimental plot (approximately 280 hours), honey bees were the only pollinator seen

visiting blueberry flowers.

We calculated the pollination deficit for each variety by comparing the honey bee treatment

to the optimised hand pollination treatment. Hand pollination is thought to result in the

best fruit quality as it enhances both pollen deposition and quality of pollen. Results from

Martin et al. (2019) were used to establish the most compatible pollen donor for each

variety, and this was used to pollinate flowers. We then followed hand pollination protocols

as set out by Martin et al. (2021).

2.3 Measurements of fruit quality

Following pollination treatments, fruit development was examined weekly to determine

whether fruit were ready to be picked. Fruit were considered ready for picking when the

entire berry had transformed to a uniform dark blue. Fruit were then harvested and the

weight, diameter, and developmental period (weeks from pollination to harvesting) were

determined on the harvest day. We also calculated the percentage fruit set per treatment.

Fruit mass and percentage fruit set were combined to calculate the adjusted fruit mass for

each treatment as follows:

$\quad Adjusted\ fruit\ mass$ = Proportion fruit set X mean fruit mass $\qquad$ (*Equation 1*)

2.4 Statistical analyses

Linear mixed-effects models were used with fruit mass, developmental period, and adjusted

fruit mass as functions of pollination treatment (fixed effect). To account for variation

between individual plants, plant ID was used as a random effect. This increased the

sensitivity of the model in identifying differences in fruit yield measures as a consequence of

treatment alone. Linear contrasts (Tukey) were used to detect treatment differences.

Nakagawa R^2 values (Nakagawa and Schielzeth, 2013) were used to assess model fit,

providing conditional variance (R^2c) and marginal variance (R^2m) estimates for linear mixed-

effects models, which can be equated to traditional R^2 values. The conditional R^2 values (R^2c)

explain the variance of the entire model (fixed effects and random effects), whereas

marginal R^2 values (R^2m) explain the variance of the fixed effects alone.

All statistical analyses were conducted in R (version 3.3.2) (R Core Team, 2017) using

the packages nlme (Pinheiro et al., 2017), multcomp (Hothorn et al., 2008), ggplot2

(Wickham, 2009), sjPlot (Lüdecke, 2017), car (Fox and Weisberg, 2011), lme4 (Bates et al.,

2015), grid (R Core Team, 2017), gridExtra (Auguie, 2017), lattice (Deepayan, 2008), MuMIn

(Barton, 2017), plyr (Wickham, 2011), and plotrix (Lemon, 2006).

3. Results

3.1 Emerald

3.1.1 Benefit of bees

We found strong statistical support indicating that honey bees increased blueberry production in Emerald compared to fruits produced in the absence of pollinators for three-quarters of the metrics measured. We provide further details on the four metrics below.

The conditional variance for the developmental period model was R^2c=0.471 and the marginal variance was R^2m=0.062. This shows that variation was mostly explained by the random effect (plant ID). The presence of honey bee pollinators appeared to decrease (statistically borderline) the developmental period of blueberry fruits by about 2 weeks (z=-2.085, p=0.089, Fig 2) compared to when they were absent.

Fig 2: Pollination deficit and the benefit of bees for Emerald fruit. A) Percentage fruit set. B) Developmental

period. C) Mean fruit mass. D) Mean fruit diameter. E) Adjusted fruit mass. Significance (p<0.05) of linear

contrasts (Tukey HSD) are indicated by letters. Error bars indicate standard error. BB = benefit of bees, the

difference (in percentage) between fruit produced in the absence of pollinators compared (n=27) to fruits

produced by honey bees (n=27). PD = pollination deficit, the difference (in percentage) between fruits

produced after honey bee exposure compared to fruits produced after hand pollination (n=27).

The conditional variance for the Emerald fruit mass model was R^2c=0.238 while the

marginal variance was R^2m=0.151. The conditional variance for the fruit diameter model

was R^2c=0.303 and the marginal variance was R^2m=0.210. This suggests that for both fruit

mass and diameter, variation is primarily explained by treatment. Fruit mass was

significantly increased by the presence of honey bees (mean ± SE fruit mass = 1.62g ± 0.12g,

z=2.799, p=0.013), resulting in a 156% mass increase compared to fruit produced in the

absence of pollinators (mean ± SE fruit mass = 0.64g ± 0.32g, Fig 2C). Likewise, honey bees

increased the fruit diameter by 4mm (33%) per fruit (mean ± SE = 15.51mm ± 0.41mm)

compared to treatments produced in the absence of pollinators (mean ± SE = 11.12mm ±

1.86mm, Fig 2D, z=3.412, p=0.002). There were significant differences in fruit set following

the three treatments (*Chi-square*=25.959, *df*=2, p<0.001, Fig 2A). Honey bees increased fruit

set by 63% relative to bagged plants. Similarly, honey bee pollination outperformed the

hand pollination treatment by 25%.

The conditional variance for the adjusted Emerald fruit mass model was R^2c=0.478

and marginal variance was R^2m=0.347. This suggests that adjusted fruit mass variation was

primarily explained by treatment. Honey bees led to a significant increase of adjusted fruit

mass (mean ± SE = 1.58g ± 0.11g, Fig 2E) of 153% compared to bagged treatments (mean ±

SE = 0.21g ± 0.11g, Fig 2E, z=4.639, p<0.001).

3.1.2 Pollination deficit

The pollination deficit metric indicated that for Emerald, honey bee pollinators performed

equally or better than hand pollinations. Below we present those results.

Hand pollination resulted in a one week decrease in the developmental period of

blueberry fruit, however, this was not significant (z=-0.073, p=0.997), from 13 ± 0.32 weeks

(mean ± SE) when open to honey bee pollination to 12 ± 0.76 (mean ± SE) when hand-

pollinated (Fig 2B). Mass of fruit produced from hand pollination (1.62g ± 0.16g, mean ± SE),

and fruit produced from honey bee pollination (1.64g ± 0.12g, mean ± SE, Fig 2C) did not

differ significantly (z=-0.110, p=0.993). The diameter of hand-pollinated fruit (mean ± SE =

15.52mm ± 0.54mm) and fruit pollinated by honey bees (mean ± SE = 15.51mm ± 0.41mm),

also did not differ significantly (z=-0.026, p=0.999). Honey bees outperformed hand

pollination with regards to fruit set by 25%.

The adjusted fruit mass was decreased significantly (z=-2.471, p=0.034) by 32% when

hand-pollinated (mean ± SE = 1.15g ± 0.11g, Fig 2E) in comparison to fruit produced from

honey bee pollination (mean ± SE = 1.58g ± 0.11g, Fig 2E).

3.2 Eureka

3.2.1 Benefit of bees

For all four metrics measured we found no statistical evidence suggesting that honey bees

improved Eureka fruit quality compared to fruits produced in the absence of pollinators. We

provide further details on each metric below.

For the developmental period model, the conditional variance was R^2c=0.405 and

the marginal variance was R^2m=0.370. The variation is therefore explained most by the fixed

effect (treatment). Honey bees did not change the developmental period substantially (z=-

0.042, p=0.999, Fig 3B).

Fig 3: Pollination deficit and the benefit of bees for Eureka fruit. A) Percentage fruit set. B) Developmental period. C) Mean fruit mass. D) Mean fruit diameter. E) Adjusted fruit mass. Significance (p<0.05) of linear contrasts (Tukey HSD) are indicated by letters. Error bars indicate standard error. BB = benefit of bees, the difference (in percentage) between fruit produced in the absence of pollinators compared (n=27) to fruits produced by honey bees (n=27). PD = pollination deficit, the difference (in percentage) between fruits produced after honey bee exposure compared to fruits produced after hand pollination (n=27).

The fruit mass model resulted in a conditional variance of R^2c=0.693 and a marginal

variance of R^2m=0.398. The conditional variance was R^2c=0.569 for the fruit diameter

model, while the marginal variance was R^2m=0.371. This indicated that variance is primarily

explained by treatment for fruit mass and diameter. Honey bee presence did not increase

fruit mass (mean ± SE fruit mass = 2.66g ± 0.21g) significantly (z=0.815, p=0.684), compared

to when pollinators were absent (mean ± SE fruit mass = 2.41g ± 0.20g, Fig 3C). Likewise,

honey bees did not significantly increase (z=0.232, p=0.970) fruit diameter (mean ± SE =

18.50mm ± 0.61mm) compared to when they were absent (mean ± SE = 18.33mm ±

0.52mm, Fig 3D,). Honey bee pollination appeared to increase fruit set by 14%, however the

differences between the three treatments were not significantly different (*Chi-*

square=2.253, *df*=2, *p*=0.324, Fig 3A).

The conditional variance for Eureka adjusted fruit mass was R^2c=0.678 and the

marginal variance was R^2m=0.383. This indicated that for the adjusted fruit mass model,

most of the variance was explained by pollination treatment. The adjusted fruit mass of

honey bee pollination (mean ± SE = 2.53g ± 0.20g, Fig 3E), did not differ significantly

(z=1.334, p=0.363) from the adjusted fruit mass of fruit produced in the absence of honey

bees (mean ± SE = 2.16g ± 0.18g, Fig 3E).

3.2.2 Pollination deficit

For all pollination deficit metrics measured for Eureka, hand pollinations outperformed

honey bee pollination. We provide further details on each metric below.

The developmental period of fruit produced by hand pollination was significantly

decreased by approximately two weeks (z=-4.521, p<0.001), from 10 ± 0.26 weeks (mean ±

SE) when honey bee pollinated to 8 ± 0.21 weeks (mean ± SE) when hand-pollinated (Fig

3B). Hand pollination significantly increased (z=3.239, p=0.003) the mass of fruit by 50%

(4.42g ± 0.36g, mean ± SE), compared to the fruit produced by flowers exposed to honey

bees (2.66g ± 0.21g, mean ± SE, Fig 3C). Furthermore, hand pollination also significantly

increased (z=3.657, p<0.001) fruit diameter (mean ± SE = 22.61mm ± 0.76mm) by 20% in

comparison to honey bee pollinated flowers (mean ± SE = 18.50mm ± 0.61mm, Fig 3D).

Hand pollination did not increase fruit set compared to honey bee pollination and in fact led

to a 4% decrease in fruit set.

The adjusted fruit mass increased significantly (z=3.008, p<0.007) by 45% after being

hand-pollinated (mean ± SE = 4.00g ± 0.33g, Fig 2E), compared to the adjusted fruit mass

after honey bee exposure (mean ± SE = 2.53g ± 0.20g, Fig 3E).

3.3 Snowchaser

3.3.1 Benefit of bees

For all four metrics, access by honey bees substantially improved Snowchaser fruit

production compared to when they were excluded. We provide further details on each

metric below.

The conditional variance of the Snowchaser developmental period model was

R^2c=0.482, while the marginal variance was R^2m=0.460, suggesting that the variance in the

developmental period was explained primarily by treatment (fixed effect). Honey bee

pollinators significantly shortened the developmental period of blueberry fruits (mean ± SE

fruit weeks = 12 weeks ± 0.20 weeks, z=-3.533, p=0.001, Fig 4B) by approximately one week

compared to fruits produced in the absence of honey bees (mean ± SE fruit weeks = 13

weeks ± 0.80 weeks, Fig 4B).

Fig 4: Pollination deficit and the benefit of bees for Snowchaser fruit. A) Percentage fruit set. B) Developmental period. C) Mean fruit mass. D) Mean fruit diameter. E) Adjusted fruit mass. Significance ($p<0.05$) of linear contrasts (Tukey HSD) are indicated by letters. Error bars indicate standard error. BB = benefit of bees, the difference (in percentage) between fruit produced in the absence of pollinators compared (n=27) to fruits produced by honey bees (n=27). PD = pollination deficit, the difference (in percentage) between fruits produced after honey bee exposure compared to fruits produced after hand pollination (n=27).

The conditional variance of the fruit mass model was R^2c=0.520 and the marginal

variance was R^2m=0.203. Similarly, the conditional variance of the fruit diameter model was

R^2c=0.469, while the marginal variance was R^2m=0.284. This suggests that variance in

Snowchaser fruit mass and diameter was primarily explained by treatment. Here, honey bee

presence significantly increased fruit mass (mean ± SE fruit mass = 1.52g ± 0.15g, z=2.362,

p=0.047), by 66% compared to treatments where honey bees were excluded (mean ± SE

fruit mass = 0.77g ± 0.25g, Fig 4C). Likewise, honey bee exposure resulted in a 4mm (33%)

mean increase in fruit diameter per fruit (mean ± SE = 14.70mm ± 0.59mm) in comparison

to fruit produced in the absence of honey bees (mean ± SE = 10.51mm ± 0.81mm, Fig 4D,

z=3.229, p=0.003). Honey bee pollination appeared to increase fruit set by 18%, however

the differences between the three treatments were not significantly different (*Chi-*

square=3.559, *df*=2, p=0.169, Fig 4A).

The adjusted fruit mass model resulted in a conditional variance of R^2c=0.576 and a

marginal variance of R^2m=0.273, suggesting that most of the variance in adjusted fruit mass

is explained by treatment. Honey bee pollination significantly (z=3.062, p=0.006) increased

adjusted fruit mass (mean ± SE = 1.30g ± 0.13g, Fig 4E), by 86% compared to the adjusted

fruit mass of treatments where pollinators were excluded (mean ± SE = 0.52g ± 0.17g, Fig

4E).

3.3.2 Pollination deficit

Statistics indicated that honey bee pollinators and hand pollinations performed similarly for

Snowchaser in three of the four metrics measured. We provide further details below.

Hand pollination resulted in a two week or 12% decrease in the developmental

period (z=-3.788, p<0.001), from 12 ± 0.20 weeks (mean ± SE) after honey bee exposure to

10 ± 0.16 weeks (mean ± SE) after hand pollination (Fig 4B). Hand-pollinated fruits had

similar fruit mass (mean ± SE fruit mass = 1.87g ± 0.14g, z=1.469, p=0.303) to fruit produced

after exposure to honey bees (1.52g ± 0.15g, mean ± SE, Fig 4C). Hand pollination slightly,

but not significantly (z=1.425, p=0.325) increased fruit diameter (mean ± SE = 16.10mm ±

0.54mm) by 9% in comparison to flowers exposed to honey bees (mean ± SE = 14.70mm ±

0.59mm, Fig 4D). Hand pollination did not lead to an increase in fruit set compared to honey

bee pollination.

The adjusted fruit mass appeared to increase by 20% for hand-pollinated flowers

(mean ± SE = 1.59g ± 0.12g, Fig 2E) compared to the adjusted fruit mass after honey bee

exposure (mean ± SE = 1.30g ± 0.13g, Fig 4E), however, this difference was not statistically

significant (z=1.486, p=0.295).

3.4 Suziblue

3.4.1 Benefit of bees

We found evidence suggesting that for Suziblue, honey bees do not improve blueberry fruit

quality compared to when pollinators were excluded. We provide further details on each

metric below.

318   Conditional variance for the developmental period model was R^2c=0.235 and the

319   marginal variance was R^2m=0.235, suggesting that the variation was mostly explained by the

320   fixed effect (treatment). Exposure to honey bees did not significantly affect the

321   developmental period of Suziblue fruits (z=-0.957, p=0.604, Fig 5B).

Fig 5: Pollination deficit and the benefit of bees for Suziblue fruit. A) Percentage fruit set. B) Developmental

period. C) Mean fruit mass. D) Mean fruit diameter. E) Adjusted fruit mass. Significance (p<0.05) of linear

contrasts (Tukey HSD) are indicated by letters. Error bars indicate standard error. BB = benefit of bees, the

difference (in percentage) between fruit produced in the absence of pollinators compared (n=27) to fruits

produced by honey bees (n=27). PD = pollination deficit, the difference (in percentage) between fruits

produced after honey bee exposure compared to fruits produced after hand pollination (n=27).

The conditional variance of the fruit mass model was R^2c=0.645 and the marginal

variance was R^2m=0.144. For the fruit diameter model, the conditional variance was

R^2c=0.492 and the marginal variance was R^2m=0.088. Plant ID explained most of the

variance for fruit mass and diameter. Honey bee pollination had a minimal effect on fruit

mass (mean ± SE fruit mass = 1.28g ± 0.14g, z=1.137, p=0.485), in comparison to bagged

treatments (mean ± SE fruit mass = 1.11g ± 0.09g, Fig 5C). Honey bees also had a minimal

effect on fruit diameter (mean ± SE = 13.91mm ± 0.56mm) in comparison to bagged

treatments (mean ± SE = 13.31mm ± 0.42mm, Fig 5D, z=1.018, p=0.562). Significant

differences were found in fruit set following the three treatments (*Chi-square*=6.003, *df*=2,

p=0.049, Fig 5A). Honey bees increased fruit set by 11% relative to bagged plants, however,

this still left a pollination deficit of 11% (honey bee treatment compared to hand

pollination).

The adjusted fruit mass model resulted in a conditional variance of R^2c=0.757 and a

marginal variance of R^2m=0.330. The variance in the adjusted fruit mass model was

explained mostly by treatment (fixed effect). Flowers open to honey bee pollination resulted

in similar adjusted fruit masses (mean ± SE fruit mass = 1.09g ± 0.12g, z=1.816, p=0.160),

compared to honey bee exclusion treatments (mean ± SE = 0.82g ± 0.07g, Fig 5E).

3.4.2 Pollination deficit

Below, we provide details showing that for all metrics measured, hand pollinations of

Suziblue flowers did not outperform honey bee pollination.

The developmental period of hand-pollinated fruit was similar (mean ± SE weeks =

12 ± 0.62 weeks, z=-2.047, p=0.101) to that of fruit produced from honey bee pollination (13

± 0.75, mean ± SE, Fig 5B). The fruit mass of hand-pollinated fruits was also similar (mean ±

SE fruit mass = 1.43g ± 0.14g, z=0.786, p=0.707), to fruit produced from honey bee

pollinated flowers (1.28g ± 0.14g, mean ± SE, Fig 5C). Hand pollination did not increase

(z=0.579, p=0.830) fruit diameter (mean ± SE = 14.30mm ± 0.59mm) in comparison to the

diameter of fruit produced by honey bee pollination (mean ± SE = 13.91mm ± 0.71mm, Fig

5D).

Adjusted fruit mass was similar and not significantly different (z=1.48, p=0.290) for

hand-pollinated flowers (mean ± SE = 1.38g ± 0.14g, Fig 5E) and for flowers exposed to

honey bees (mean ± SE = 1.09g ± 0.12g, Fig 5E).

3.5 Twilight

3.5.1 Benefit of bees

For three out of the four metrics measures, honey bee presence enhanced blueberry

production compared to when pollinators were excluded. We provide further details on

each metric below.

The developmental period model had a conditional variance of R^2c=0.360 and a

marginal variance of R^2m=0.320. The fixed effect (treatment) explained most of the

variation. Honey bee pollination decreased the developmental period (mean ± SE weeks =

11 weeks ± 0.43 weeks, Fig 6B) of blueberry fruits by two weeks or 17% compared to fruits

produced in the absence of honey bees (mean ± SE fruit weeks = 11 weeks ± 0.49 weeks, Fig

2C). This was borderline-significant (z=-2.170, p=0.076, Fig 6B).

Fig 6: Pollination deficit and benefit of bees for Twilight fruit. A) Percentage fruit set. B) Developmental period.

C) Mean fruit mass. D) Mean fruit diameter. E) Adjusted fruit mass. Significance (p<0.05) of linear contrasts

(Tukey HSD) are indicated by letters. Error bars indicate standard error. BB = benefit of bees, the difference (in

percentage) between fruit produced in the absence of pollinators compared (n=27) to fruits produced by

honey bees (n=27). PD = pollination deficit, the difference (in percentage) between fruits produced after honey

bee exposure compared to fruits produced after hand pollination (n=27).

The conditional variance for the Twilight fruit mass model was R^2c=0.337 and the

marginal variance was R^2m=0.170. For the fruit diameter model, the conditional variance

was R^2c=0.392, while the marginal variance was R^2m=0.369. This suggests that for both fruit

mass and diameter, the majority of the variance was explained by treatment. Fruit mass was

significantly increased after honey bee exposure (mean ± SE fruit mass = 2.45g ± 0.17g,

z=2.397, p=0.043), resulting in a 54% increase compared to bagged treatments (mean ± SE

fruit mass = 1.41g ± 0.17g, Fig 6C). Honey bee exposure similarly increased the mean fruit

diameter by 3mm (18%) per fruit (mean ± SE = 17.66mm ± 0.39mm) in comparison to

bagged treatments (mean ± SE = 14.73mm ± 0.47mm, Fig 6D, z=4.643, p<0.001). There were

significant differences in fruit set following the three treatments (*Chi-square*=13.176, *df*=2,

p=0.001, Fig 6A). Honey bees increased fruit set by 6% relative to bagged plants, however,

this still left a pollination deficit of 11% (honey bee treatment compared to hand

pollination).

The adjusted fruit mass model resulted in a conditional variance of R^2c=0.383 with

the marginal variance of R^2m=0.216 suggesting that fruit mass variance was primarily

explained by treatment. Honey bee presence significantly (z=2.368, p=0.046) increased

adjusted fruit mass (mean ± SE = 2.18g ± 0.15g, Fig 6E), resulting in a 60% increase

compared to pollinator exclusion treatments (mean ± SE = 1.18g ± 0.12g, Fig 6E).

3.5.2 Pollination deficit

Hand pollinations resulted in improved Twilight fruit quality compared to honey bee

pollinations. However, these were only statistically supported for developmental period.

Below, we explain the details relating to each of the metrics measured.

The developmental period of blueberry fruit was significantly decreased by approximately

one week or 10% (z=-2.824, p=0.0132), from 11 ± 0.43 weeks (mean ± SE) when flowers

were open to honey bee pollination to 10 ± 0.20 (mean ± SE) when flowers were pollinated

by hand (Fig 6B). Hand-pollinated fruits appeared to be about 24% heavier (3.11g ± 0.45g,

mean ± SE), than fruit produced from flowers exposed to honey bees (2.45g ± 0.17g, mean ±

SE, Fig 6C), however, this relationship was non-significant (z=1.134, p=0.493). Hand

pollination did not significantly increase (z=1.134, p=0.493) the fruit diameter (mean ± SE =

18.36mm ± 0.43mm) when compared to fruit produced by honey bee pollinated flowers

(mean ± SE = 17.66mm ± 0.39mm, Fig 6D).

Hand pollination appeared to improve the adjusted fruit mass by 35% (mean ± SE =

3.10g ± 0.45g, Fig 2E) when compared to flowers exposed to honey bees (mean ± SE = 2.18g

± 0.15g, Fig 2E), although this difference was not statistically significant (z=1.694, p=0.205).

4. Discussion

We provide a comprehensive comparison of honey bee effectiveness as pollinators of common commercial blueberry varieties in South Africa. Evidence suggests that the fruit quality of these southern highbush blueberry varieties are considerably affected by the pollination environment. Overall, we show that the absence of pollinators in highbush blueberries typically leads to reductions in fruit yield. This provides important information to blueberry growers as it shows that honey bees are usually extremely beneficial and can act as reliable blueberry pollinators, even in areas without access to native blueberry pollinators. Importantly, the study also highlights differences in the effectiveness of honey bees across different blueberry varieties. Finally, it also suggests that pollination conditions could be improved for certain varieties which appear not to be producing at their full production potential under honey bee pollination.

For comparative purposes, we have highlighted adjusted fruit mass as a summary table metric (Table 1) as it incorporates arguably the two most important yield metrics (fruit mass and fruit set), with fruit diameter being indirectly incorporated as it is highly correlated with fruit mass (Jorquera-fontena et al., 2017).

Table 1: Summary table comparing the benefit of bees and the pollination deficit for adjusted fruit mass in the

five blueberry varieties from this study and Ventura, which was published by Martin et al. (2021). Positive

benefit of bees imply that the addition of honey bees increases fruit yields compared to when pollinators are

absent. Positive pollination deficit indicates that plants pollinated by honey bees produce lower yields than

plants pollinated by hand. Statistical significance ($p < 0.05$) is indicated by bold fonts, while colour differences

(red vs black)

suggest differences

in trend direction.

Variety	Benefit of bees	Pollination deficit
Emerald	**153%**	**-32%**
Eureka	16%	**45%**
Snowchaser	**86%**	20%
Suziblue	28%	24%
Twilight	**60%**	35%
Ventura (Martin et al. 2021)	**134%**	**37%**

In all cases, honey bees increase yield (usually statistically significant) compared to

yield when pollinators were absent. This indicates that although honey bees lack the ability

to buzz pollinate, they do move pollen between blueberry flowers and typically lead to

increased blueberry yields. Martin et al. (2021) recently demonstrated this for Ventura,

another southern highbush variety used in South Africa. In this study honey bees led to a

134% increase of adjusted fruit mass compared to fruits formed in the absence of honey

bees, demonstrating that honey bees can effectively pollinate highbush blueberry varieties.

The addition of honey bees can result in very impressive yield increases for some

varieties (i.e. Emerald, Snowchaser, Twilight, and Ventura), but for two varieties (Eureka and

Suziblue), honey bee exposure had relatively small effects on fruit production. This may

occur if a variety is good at self-pollination and does not need pollinators. Good self-

pollinators should also have low pollination deficits because pollinators are not needed for

good fruit production. Suziblue appears to be a good self-pollinator because it is setting

close to its maximum possible yield (a small pollination deficit) and its yields are not strongly

increased by the presence of honey bees. We suggest that varieties such as Suziblue may be

good varieties to plant under conditions where pollinator visitation is extremely erratic.

Similarly, Eureka also did not benefit greatly (16%) from the presence of honey bees,

however, the high and significant pollination deficit (45%) for Eureka suggests that it is a

poor self-pollinator and that the marginal benefit of honey bee exposure is due to the fact

that this variety is extremely poorly adapted to honey bee pollination. Blueberry flowers

differ in terms of their morphological characters and it may be possible that it is difficult for

honey bees to access rewards in certain varieties. This may reduce pollinator visitation rates

to those varieties or possibly make it hard for pollen grains to be extracted. Studies on wild

pollinators have often found that traits such as floral tube length and diameter can have a

profound effect on the effectiveness of pollen transfer and also on whether pollinators are

able to access nectar or not (Campbell et al., 1996; Aigner, 2004; Fenster et al., 2006;

Martins and Johnson, 2013; Lázaro et al., 2015; Minnaar et al., 2019). We predict that

varieties with long tubes can exclude pollinators from accessing nectar, while varieties with

narrow entrances may exclude pollinators from accessing both nectar and pollen. Evidence

that some varieties have inaccessible or poor rewards for pollinators can be seen in studies

that show strong pollinator preferences for one blueberry variety over another (Dorr and

Matin, 1966; Courcelles et al., 2013).

The pollination deficit for Emerald was remarkable because it was so small (less than

zero), suggesting that honey bee pollination was better than hand pollination. The presence

of honey bees also had an enormous (153%) positive impact on yield, suggesting that this

variety is exceptionally well suited to honey bee pollination. Honey bees may be particularly

good at pollinating Emerald flowers due to the large flower openings (4.6 mm) which may

allow honey bees to more easily insert their bodies inside the flower. This may enhance the

probability of picking up pollen on each visit and it may also allow easier access to nectar,

which should make flowers more attractive to honey bees, potentially increasing visitation

rates. Emerald may therefore be an excellent variety to use in areas where there are no

bumble bees such as South Africa. Similarly, honey bee presence had a very positive impact

on Snowchaser yields and this, in combination with small (20%) pollination deficits suggests

that this variety is also quite well suited to honey bee pollination. Twilight and Ventura also

benefitted well from honey bee presence although the larger pollination deficits for these

varieties suggest that they are not as well suited to honey bee pollination as Emerald or

Snowchaser.

Honey bees are often thought to be poor pollinators of blueberries because of their

reluctance to forage in poor weather conditions (Tuell and Isaacs, 2010), slower foraging

times (Javorek et al., 2002), and decreased pollen deposition per visit (Javorek et al., 2002)

when compared to pollinators such as bumble bees. Despite this, we show that honey bees

are reliable pollinators of several blueberry varieties in regions that do not have access to

bumble bees. This is especially true for Emerald, where honey bee pollination is so effective

that it can potentially negate the need for other pollinators. Importantly the benefits of

honey bees can differ strongly between varieties, revealing the importance of conducting

research on the effectiveness of honey bee pollination on different highbush blueberry

varieties. Blueberry varieties Emerald, Snowchaser, and Twilight seem to do particularly well

under honey bee pollination and Eureka appears to be a very poor choice of variety for

environments that only have honey bee pollinators. However, it is also important to note

that without comparative studies on bumble bee pollination, it is difficult to say whether the

509 addition of bumble bees would significantly improve the pollination deficits observed. The

510 strong varietal differences in the effectiveness of honey bee pollination suggest that

511 considerable research is needed to identify which blueberry varieties do best under honey

512 bee pollination, what makes varieties successful or unsuccessful, which varieties can be

513 effectively pollinated by honey bees, and which varieties need other effective means to

514 increase fruit quality. Future research should therefore aim to determine how to decrease

515 the pollination deficit of blueberry varieties and increase the effectiveness of honey bees.

References

Aigner, P.A., 2004. Floral specialization without trade-offs: optimal corolla flare in contrasting pollination environments. Ecology 85, 2560–2569.

Aizen, M.A., Smith-Ramírez, C., Morales, C.L., Vieli, L., Sáez, A., Barahona-Segovia, R.M., Arbetman, M.P., Montalva, J., Garibaldi, L.A., Inouye, D.W., Harder, L.D., 2018. Coordinated species importation policies are needed to reduce serious invasions globally: The case of alien bumblebees in South America. J. Appl. Ecol. 1–7.

Aras, P., de Oliveira, D., Savoie, L., 1996a. Effect of a honey bee (Hymenoptera: Apidae) gradient on the pollination and yield of lowbush blueberry. J. Econ. Entomol. 89, 1080–1083.

Aras, P., de Oliveira, D., Savoie, L., 1996b. Effect of a honey bee (Hymenoptera: Apidae) gradient on the pollination and yield of lowbush blueberry. J. Econ. Entomol. 89, 1080–1083.

Auguie, B., 2017. GridExtra: miscellaneous functions for "grid" graphics.

Barton, K., 2017. MuMIn: multi-model inference.

Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. J. Stat. Softw. 67, 1–48.

Campbell, D.R., Waser, N.M., Price, M. V, 1996. Mechanisms of hummingbird-mediated selection for flower width in *Ipomopsis aggregata*. Ecology 77, 1463–1472.

Campbell, J.W., Brien, J.O., Irvin, J.H., Kimmel, C.B., Daniels, J.C., Ellis, J.D., 2017. Pollinator ecology and management managed bumble bees (*Bombus impatiens*) (Hymenoptera:

Apidae) caged with blueberry bushes at high density did not increase fruit set or fruit

weight compared to open pollination 46, 237–242.

Campbell, J.W., Kimmel, C.B., Bammer, M., Stanley-Stahr, C., Daniels, J.C., Ellis, J.D., 2018.

Managed and wild bee flower visitors and their potential contribution to pollination

services of low-chill highbush blueberry (*Vaccinium corymbosum* L.; Ericales:

Ericaceae). J. Econ. Entomol. 111, 2011–2016.

Cane, J.H., Payne, J.A., 1993. Regional, annual, and seasonal variation in pollinator guilds:

intrinsic traits of bees (Hymenoptera: Apoidea) underlie their patterns of abundance at

Vaccinium ashei (Ericaceae). Ann. Entomol. Soc. Am. 56, 577–588.

Courcelles, D.M.M., Button, L., Elle, E., 2013. Bee visit rates vary with floral morphology

among highbush blueberry cultivars (*Vaccinium corymbosum* L.). J. Appl. Entomol. 137,

Deepayan, S., 2008. Lattice: multivariate data visualization with R. Springer, New York.

Dogterom, M.H., Winston, M.L., Amy, M., 2000. Effect of pollen load size and source (self,

outcross) on seed and fruit production in highbush blueberry cv. "Bluecrop" (*Vaccinium*

corymbosum; Ericaceae). Am. J. Bot. 87, 1584–1591.

Dorr, J., Matin, E.C., 1966. Pollination studies on the high bush blueberry, *Vaccinium*

corymbosum LQ, Bulletin of Michigan State University Agricultural Experiment Station.

Fenster, C.B., Cheely, G., Dudash, M.R., Reynolds, R.J., 2006. Nectar reward and

advertisement in hummingbird-pollinated *Silene virginica* (Caryophyllaceae). Am. J. Bot.

93, 1800–1807.

Fox, J., Weisberg, S., 2011. An {R} companion to applied regression, Second Edi. ed. Sage,

California.

Fulton, M., Jesson, L.K., Bobiwash, K., Schoen, D.J., 2015. Mitigation of pollen limitation in

the lowbush blueberry agroecosystem: effect of augmenting natural pollinators.

Ecosphere 6, 1–19.

Gallai, N., Salles, J.M., Settele, J., Vaissière, B.E., 2009. Economic valuation of the

vulnerability of world agriculture confronted with pollinator decline. Ecol. Econ. 68,

810–821.

Garibaldi, L.A., Steffan-dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R., Cunningham,

S.A., Kremen, C., Carvalheiro, L.G., 2013. Wild pollinators enhance fruit set of crops

regardless of honey bee abundance. Science (80). 339, 1608–1611.

Hoffman, G.D., Lande, C., Rao, S., 2018. A novel pollen transfer mechanism by honey bee

foragers on highbush blueberry (Ericales: Ericaceae). Environ. Entomol. 47, 1465–1470.

Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric

models. Biometrical J. 50, 346–363.

Javorek, S.K., Mackenzie, K.E., Vander Kloet, S.P., 2002. Comparative pollination

effectiveness among bees (Hymenoptera: Apoidea) on lowbush blueberry (Ericaceae:

Vaccinium angustifolium). Ann. Entomol. Soc. Am. 95, 345–351.

Jorquera-fontena, E., Génard, M., Franck, N., 2017. Analysis of blueberry (V*accinium*

corymbosum L.) fruit water dynamics during growth using an ecophysiological model. J.

Hortic. Sci. Biotechnol. 92, 646–659.

Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C.,

Tscharntke, T., 2007. Importance of pollinators in changing landscapes for world crops.

Proc. R. Soc. London B Biol. Sci. 274, 303–313.

581 Lázaro, A., Vignolo, C., Santamaría, L., 2015. Long corollas as nectar barriers in *Lonicera*

582   *implexa*: interactions between corolla tube length and nectar volume. Evol. Ecol. 29,

583   419–435.

584 Lemon, J., 2006. Plotrix: a package in the red light district of R. R-News 6, 8–12.

585 Lüdecke, D., 2017. SjPlot: data visualization for statistics in social science.

586 MacKenzie, K., 1997. Pollination requirements of three highbush blueberry cultivars. J. Am.

587   Soc. Hortic. Sci.

588 Mackenzie, K.E., 1997. Pollination requirements of three highbush blueberry (*Vaccinium*

589   *corymbosum* L.) cultivars. J. Am. Soc. Hortic. Sci. 122, 891–896.

590 Mackenzie, K.E., Eickwort, G.C., 1996. Diversity and abundance of bees (Hymenoptera:

591   Apoidea) foraging on highbush blueberry (*Vaccinium corymbosum* L.) in central New

592   York. J. Kansas Entomol. Soc. 69, 185–194.

593 Martin, K., Minnaar, C., Jager, M. De, Anderson, B., 2019. Cross-compatibility of five

594   highbush blueberry varieties and ideal crossing combinations. bioRxiv 742114.

595 Martin, K., Anderson, B., Minnaar, C., Jager, M. De, 2021. Honey bees are important

596   pollinators of South African blueberries despite their inability to sonicate. South African

597   J. Bot. 137, 46–51.

598 Martins, D.J., Johnson, S.D., 2013. Interactions between hawkmoths and flowering plants in

599   East Africa: polyphagy and evolutionary specialization in an ecological context. Biol. J.

600   Linn. Soc. 110, 199–213.

601 Minnaar, C., de Jager, M.L., Anderson, B., 2019. Intraspecific divergence in floral-tube length

602 promotes asymmetric pollen movement and reproductive isolation. New Phytol. 224,

603 1160–1170.

604 Morales, C.L., Arbetman, M.P., Cameron, S.A., Aizen, M.A., 2013. Rapid ecological

605 replacement of a native bumble bee by invasive species. Front. Ecol. Environ. 11, 529–

606 534.

607 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R2 from

608 generalized linear mixed-effects models. Methods Ecol. Evol. 4, 133–142.

609 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Team, R.C., 2017. Nlme: linear and nonlinear

610 mixed effects models.

611 R Core Team, 2017. R: a language and environment for statistical computing.

612 Sampson, B.J., Cane, J.H., 2000. Pollination efficiencies of three bee (Hymenoptera:

613 Apoidea) species visiting rabbiteye blueberry. J. Econ. Entomol. 93, 1726–1731.

614 Schmid-Hempel, R., Eckhardt, M., Goulson, D., Heinzmann, D., Lange, C., Plischuk, S.,

615 Escudero, L.R., Salathé, R., Scriven, J.J., Schmid-Hempel, P., 2014. The invasion of

616 southern South America by imported bumblebees and associated parasites. J. Anim.

617 Ecol. 83, 823–837.

618 Stubbs, C.S., Drummond, F.A., 2001. *Bombus impatiens* (Hymenoptera: Apidae): an

619 alternative to *Apis mellifera* (Hymenoptera: Apidae) for lowbush blueberry pollination.

620 J. Econ. Entomol. 94, 609–616.

621 Tuell, J., Ascher, J., Isaacs, R., 2009. Wild Bees (Hymenoptera: Apoidea: Anthophila) of the

622 Michigan highbush blueberry agroecosystem. Ann. Entomol. Soc. Am. 102, 275–287.

623   Tuell, J.K., Isaacs, R., 2010. Weather during bloom affects pollination and yield of highbush

624   blueberry. J. Econ. Entomol. 103, 557–562.

625   Wickham, H., 2011. The split-apply-combine strategy for data analysis. J. Stat. Softw. 40, 1–

626   29.

627   Wickham, H., 2009. Ggplot2: elegant graphics for data analysis. Springer-Verlag, New York.