Article

Native Metabolomics Identifies the Rivulariapeptolide Family of Protease Inhibitors

Raphael Reher[1,3], Allegra T Aron[2], Pavla Fajtová[2], Chenxi Liu[2], Ido Y Ben Shalom[2], Wout Bittremieux[2], Mingxun Wang[2], Marie L Matos-Hernandez[4], Kelsey L Alexander[1,6], Eduardo J Caro-Diaz[4], C Benjamin Naman[5], Pieter C Dorrestein[2], Anthony J O'Donoghue[2], William H Gerwick[1,2,#] & Daniel Petras[1,2,7,#]

[1] Scripps Institution of Oceanography, University of California San Diego, USA
[2] Skaggs School of Pharmacy and Pharmaceutical Science, University of California San Diego, USA
[3] Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Halle (Saale), Germany.
[4] Department of Pharmaceutical Sciences, School of Pharmacy, University of Puerto Rico - Medical Sciences Campus, San Juan, Puerto Rico
[5] Li Dak Sum Yip Yio Chin Kenneth Li Marine Biopharmaceutical Research Center, Department of Marine Pharmacy, College of Food and Pharmaceutical Sciences, Ningbo University 315832, Ningbo, China
[6] Department of Chemistry and Biochemistry, University of California San Diego, USA
[7] CMFI Cluster of Excellence, Interfaculty Institute of Microbiology and Medicine, University of Tuebingen, Germany

[#] Correspondence: daniel.petras@uni-tuebingen.de (native metabolomics and MS-based structure elucidation), wgerwick@health.ucsd.edu (sampling and NMR-based structure elucidation)

Abstract

The identity and biological activity of most metabolites still remain unknown. A key bottleneck in the full exploration of this tremendous source of new structures and pharmaceutical activities is the compound purification needed for bioactivity assignments of individual compounds and downstream structure elucidation. To enable bioactivity-focused compound identification from complex mixtures, we developed a scalable native metabolomics approach that integrates non-targeted liquid chromatography tandem mass spectrometry, and simultaneous detection of protein binding via native mass spectrometry. While screening for new protease inhibitors from an environmental cyanobacteria community, native metabolomics revealed 30 cyclodepsipeptides as chymotrypsin binders. Mass spectrometry-guided purification then allowed for full structure elucidation of the most prevalent compounds via nuclear magnetic resonance spectroscopy, as well as orthogonal bioactivity studies. Together, these results identified the rivulariapeptolides as a family of serine protease inhibitors with nanomolar potency.

Main

Specialized metabolites, often referred to as natural products, are a tremendous pool of chemically diverse and pharmaceutically active organic compounds. By some estimates more than 50% of all current pharmaceuticals are based on or inspired by natural products[1]. Nevertheless, the vast majority of biological activities and pharmaceutical potential of specialized metabolites, still remain to be discovered, and their ecological functions to be understood[2].

While mining genome and meta-genome data starts to provide an overview of the biosynthetic potential of nature[3,4], most specialized metabolites remain inaccessible, as the living organism that produces these metabolites cannot be cultured or their gene clusters remain silent under laboratory culturing conditions. Natural product discovery and chemical ecology studies in environmental samples are hence becoming more and more attractive[5].

Along with next-generation sequencing technologies, recent instrument and computational advances in nuclear magnetic resonance (NMR) spectroscopy and non-

targeted liquid chromatography tandem mass spectrometry (LC-MS/MS) offer tremendous assistance to explore the uncharted metabolic space of nature[6]. These tools enable large scale compound dereplication, rapid identification of chemical analogs, and *de novo* annotation of molecular formulas, substructures, chemical classes and structures[7–12]. However, the assignment of bioactivity of newly identified metabolites, typically requires assays using pure compounds. Therefore, the isolation of specialized metabolites is typically guided by repetitive bioactivity assays. Together with full structure elucidation, this process usually takes months, and is therefore a major bottleneck for the systematic exploration of nature for novel pharmaceutically active compounds and comprehensive chemical ecology studies.

A logical next step is the development of *activity metabolomics*[13] or *functional metabolomics*[14] approaches, that aim to add functional information to the metabolites detected in a given system.

With native electrospray ionization (ESI), mass spectrometry (MS) is increasingly being used to analyze non-covalent binding of biomolecules, typically performed in direct infusion experiments[15–21]. While native MS workflows have been developed that can identify metabolites bound to proteins by MS/MS[22], combining the separation power of ultra-high-performance liquid chromatography (UHPLC) and the selectivity of native MS offer a great potential to decipher protein-metabolite interactions out of complex biological mixtures, such as environmental samples. However, typical mobile phase conditions disfavor non-covalent protein binding due to an acidic pH and high organic solvent content.

To perform native MS coupled to UHPLC, we developed an experimental setup that increases pH and water content of the mobile phase post-column and infuses a protein binding partner before entering the ESI interface (**Figure 1**). As the protein is constantly infused post-column, one can monitor the intact protein mass over the LC-MS run and observe mass shifts when eluting metabolites bind to the protein at a defined retention time. Using collision induced dissociation (e.g., *Higher Energy C-Trap Dissociation* (HCD) in our setup), the complex can be dissociated again in the mass spectrometer and a

"binding threshold" can be applied to distinguish between specific and non-specific

binding. In combination with parallel non-targeted MS/MS analyses, the mass and

compound ID or compound class can be assigned (level 2 or level 3 annotation[23,24]).

After our initial method validation, we screened for new protease inhibitors from an

environmental cyanobacteria biofilm as a first application. In general, cyanobacteria have

been a rich source of highly bioactive natural products[25–28], and in particular protease

inhibitors from numerous chemical classes[29–34]. Protease inhibitors are key compounds

used for treatment of viral infections (HIV and Hepatitis C)[35], cancer[36], diabetes[37],

hypertension[38], and as general anti-coagulants[39]. Several of the approved protease

inhibitors are analogs of natural products such as aliskiren, captopril, and carfilzomib that

target renin, angiotensin-converting enzyme, and proteasome, respectively[40].

In this study, we used chymotrypsin as the protease target to identify inhibitors from a

marine cyanobacteria community. Using our native metabolomics approach, we identified

30 chymotrypsin binders in the crude extract with a single LC-MS run. The masses and

MS/MS spectra of the binders were queried against structural and spectral databases,

revealing that most of them were unknown. This led to the targeted isolation and structure

elucidation of a family of new, and highly potent protease inhibitors, which we termed

"rivulariapeptolides".

Figure 1: (a) Native metabolomics setup. A crude extract is separated by μ-flow UHPLC. The pH is adjusted after chromatography with ammonium acetate to "native-like" conditions via the make-up pump. Orthogonally, protein of interest is infused, and the resulting protein-binder complexes are measured by FT-MS. The procedure is repeated as a metabolomics run (high resolution UHPLC-MS/MS acquisition without protein infusion). For the data analysis (b) the *Δ m/z* and retention time of the native MS run are correlated with *m/z* values and retention time of the metabolomics (LC-MS/MS) run and subsequently visualized using molecular networking and retention time pairing that links the observed mass differences of the protein in the native state vs bound states with the parent mass of MS/MS spectra of the small molecule.

Results

Development of the native metabolomics approach

In a crude extract, native metabolomics provides binding information about each compound towards a protein of interest. In our experimental set-up, we utilized a single 10-minute LC-MS run to discover compounds that bind to the serine protease, chymotrypsin. The workflow is as follows: A crude extract is analyzed using native ESI while the protein of interest is infused post-column throughout the entire LC gradient.

Binding of a small molecule to the protein of interest results in a peak with a mass
corresponding to the protein bound to the compound. The *m/z* difference between the
protein-ligand complex and the unbound protein reveals the molecular weight of the
ligand while the ratio of the intensity of the protein-ligand peaks relative to the unbound
protein peaks provides a qualitative indication of the binding affinity under these
conditions.

We first optimized the pH for native mass spectrometric acquisition of chymotrypsin and
confirmed that the enzyme remains active under our native metabolomics buffer
conditions. As a positive control for binding, we used molassamide[41], a known non-
covalent serine protease inhibitor of the 3-amino-6-hydroxy-2-piperidone (Ahp)-
cyclodepsipeptide family. We found that an ammonium acetate buffer of pH 4.5 showed
the highest peak intensity (**Figure S1a**). Next, we injected a serial dilution of molassamide
into our native metabolomics LC-MS setup where it was mixed post-column with a
constant concentration of chymotrypsin. The protein-ligand complex was detected at a
deconvoluted mass of 26195.1 Da and the unbound apoprotein at 25232.6 Da (**Figure
S1b**). The observed Δ *mass* of 962.5 Da matches the mass of molassamide
(962.4749 Da). After deconvolution and integrating the peak area of the protein-ligand
complex and plotting it against the molassamide concentration in the peak window, we
obtained a binding curve. The resulting curve depicts a concentration-dependent increase
of protein-ligand to unbound protein ratio with increasing ligand concentration (**Figure
S1c**). To test the specificity of these interactions, we evaluated binding of small molecule
QC standards and the peptidic cysteine protease inhibitor gallinamide A[27,42,43]. We could
not detect any binding to chymotrypsin by these negative controls under native MS
conditions (**Figure S2**).

To further test our native metabolomics conditions, chymotrypsin was assayed with crude
extract from the cyanobacterium *Rivularia* sp. using a fluorescence substrate competition
assay. The bioassay conditions were designed to mimic the pH and solvent composition
expected in the native mass spectrometry setup. Although chymotrypsin is optimally
active in near-neutral pH, it retains good activity in 10 mM ammonium acetate at pH 4.5.
Under these conditions, it was completely inhibited by 10 μg/mL of extract with 50%

inhibition at 0.84 µg/mL (**Figure S1d**). Chymotrypsin was then assayed in increasing

concentrations of acetonitrile (ACN) to determine if enzyme activity was retained in this

solvent. Activity was reduced by 9% to 34% in ACN concentrations up to 33.3% v/v. In

the presence of 41.7% v/v ACN which corresponds to the end of the UHPLC gradient,

activity was decreased by 70% (**Figure S1e**). These results confirmed that chymotrypsin

can be used as a target protease for native metabolomics as it retains activity at pH 4.5

in ACN concentrations up to 42% and binds to compounds from an inhibitory crude

extract.

Native metabolomics reveals chymotrypsin binders

Following the successful proof-of-concept experiment with molassamide, we next

screened for potential chymotrypsin binders from a crude extract of a biofilm from the

marine cyanobacterium *Rivularia* sp, collected from coral sediments at Carlos Rosario

Beach in Culebra, Puerto Rico, U.S. The methanol extract was separated by reversed-

phase UHPLC and ammonium acetate buffer and chymotrypsin were infused post-

column, prior to native ESI and acquisition of mass spectrometry data in the high m/z

range (2500-5000 *m/z*). The crude extract was subsequently re-injected, without infusion

of chymotrypsin to obtain high-resolution LC-MS/MS data of compounds in the extract in

the low m/z range (300-2000 *m/z*).

As a first step of data analysis, we plotted the total ion current (TIC) of the crude extract

metabolomics data (**Figure 2a,** left) and extracted ion chromatogram (XIC) of the apo-

chymotrypsin from charge state deconvoluted native mass spectrometry data (**Figure 2a,**

right) which shows several negative peaks in the range of 4.5 - 5.5 minutes. The decrease

in signal of the apo-protein in that retention time range is due to the emergence of larger

masses that correspond to protein-ligand complexes. After feature finding of the

deconvoluted masses and matching of the parallel metabolomics LC-MS/MS data of the

crude extracts by retention time and exact mass matching, we could identify more than

30 potential small molecule-protein complexes. To display the family of small molecules

that form protein ligand complexes and to show their structural relations, we visualized

them in a correlation molecular network (**Figure 2b**) that is based on their MS/MS

similarity (grey line), retention time, and mass matching between protein and small

molecules through the red dashed lines.

The molecular weight of these chymotrypsin binding compounds ranged from 900 to

1,250 Da (**Figure S3**), the most abundant being *m/z* 1186 and *m/z* 1156 that also show

perfect overlap of the chromatographic profiles between intact protein and metabolomics

LC-MS/MS data (**Figure 2c**). Based on their highest relative abundance we targeted them

for further purification, structure determination by NMR and orthogonal protease inhibition

assays.

193
Figure 2: (a) Left panel: Full mass spectrum (*m/z* 300 - 2,000) of cyanobacterial crude extract and blank.
The retention times (RT = 4.5 - 5.5 min) of major Ahp-cyclodepsipeptides are highlighted. Right panel:
Deconvoluted extracted ion chromatogram (*m/z* 25,233 ± 5) of alpha-chymotrypsin screened against the
cyanobacterial crude extract under native MS conditions and negative control. (b) Correlation molecular
network of deconvoluted chymotrypsin and putatively new small molecule inhibitors binders by native MS.
(c) Upper left panel: Extracted ion chromatogram (*m/z* 1,186.6284 - 1,186.6522) and MS² spectrum of
putative new chymotrypsin-binder rivulariapeptolide 1185 at RT = 5.5 min. Upper right panel: Extracted ion
chromatogram (*m/z* 26,420 ± 5) of a putative chymotrypsin-binder complex. Mass difference between the
putative chymotrypsin-binder complex (*m/z* 26,420 ± 5) and apo chymotrypsin (25,233 ± 5) suggests a
molecular weight of 1,187 ± 5 Da for the putative chymotrypsin binder rivulariapeptolide 1185. Lower left
panel: Extracted ion chromatogram (*m/z* 1,156.5819 - 1,156.6051) and MS² spectrum of putative

chymotrypsin-binder rivulariapeptolide 1185 at RT = 5.1 min. Lower right panel: Extracted ion
chromatogram (*m/z* 26,385 ± 5) of a putative chymotrypsin-binder complex. Mass difference between the
putative chymotrypsin-binder complex (*m/z* 26,385 ± 5) and apo chymotrypsin (25,233 ± 5) suggests a
molecular weight of 1,152 ± 5 Da for the putative chymotrypsin binder rivulariapeptolide 1155.
209

Rivulariapeptolides, a family of new Ahp-cyclodepsipeptides

The potential chymotrypsin binder with the nominal *m/z* 1186, identified by native

metabolomics, was next targeted for isolation and structure elucidation. Hence, we first

separated the *Rivularia* crude extract into four fractions of decreasing polarity via solid

phase extraction. SMART NMR[11,44] analysis was applied to the most hydrophilic fraction

and all but one structure of the top 10 SMART results were predicted as cyclic

depsipeptides (**Figures 3a, S4**), including 6 of the top 10 as Ahp-cyclodepsipeptides

(three from marine filamentous cyanobacteria: somamide B, molassamide, lyngyastatin

6)[41,45–47]. Complementarily, MS/MS-based molecular networking analysis of the *Rivularia*

crude fractions helped with the annotation of the known Ahp-peptides molassamide,

kurahamide, and loggerpeptin A along with several putatively new ones (**Figure 3b, S5a**).

Next, the SIRIUS and ZODIAC[48,49] tools were applied to determine the molecular formula

of the chymotrypsin-binding features, with exact mass *m/z* 1186.6400 [M+H]$^+$, as

$C_{61}H_{87}N_9O_{15}$ (0.5 ppm). Subsequently, we classified the MS/MS spectrum indicative for a

'cyclic depsipeptide' based on the classification with CANOPUS[9]. Further substructures

of the molecule were predicted as benzene, hydroxy-benzene, and proline/ N-acyl-

pyrrolidine derivatives, as well as piperidinone/delta-lactam for the Ahp-family defining

moiety (**Figure 3c**).

To unambiguously determine the structure, we isolated nominal *m/z* 1186, named

rivulariapeptolide 1185 (**1**), and performed 1D and 2D NMR experiments and manual

MS/MS interpretation (**Figure 3d, Figures S6-S11, Table S1**). Subsequently, we targeted

the isolation of further rivulariapeptolides by preparative HPLC, based on their protein-

ligand complex ratios from the native metabolomics experiments as well as their relative

abundance. In that way, we isolated and elucidated the planar structures of the

rivulariapeptolides 1185, 1155, 1121, and 989 (**1**, **2**, **3**, **4**) with the exact masses

1186.6400 [M+H]$^+$ ($C_{61}H_{88}N_9O_{15}$, 0.5 ppm), 1156.5923 [M+H]$^+$ ($C_{59}H_{82}N_9O_{15}$, - 0.2 ppm),

1122.6080 [M+H]$^+$ ($C_{56}H_{84}N_9O_{15}$, - 0.1 ppm) and 989.4978 [M+H]$^+$ ($C_{50}H_{69}N_8O_{13}$, - 0.1

ppm). In addition to rivulariapetolides, we identified the already known molassamide (**5**)
as well as new derivative we termed "molassamide B" (**6**) with *m/z* 1041.3924 [M+H]$^+$
(C$_{48}$H$_{66}$BrN$_8$O$_{13}$, -0.3 ppm), which is ortho-brominated (**Scheme S1, Table S2-S3,**
Figures S12-S34).

Finally, the chymotrypsin inhibitory activities of the purified Ahp-cyclodepsipeptides **1**-**6**
were assessed by specific biochemical assays and confirmed the results of our native
metabolomic protein infusion MS experiments (**Figure 4b**). All six compounds were found
to be nanomolar chymotrypsin inhibitors with compound **1** being the most potent (**Figure**
4, IC$_{50}$ = 8.5 ± 0.54 nM). Intriguingly, a single ortho-bromination in the *N*-methyltyrosine
moiety led to a 30-fold increase in potency for the new compound **6** (IC$_{50}$ = 16.54 ± 1.02
nM), when compared to the known compound **5** (IC$_{50}$ = 503.2 ± 33.6 nM). These
promising results led us to test these six Ahp-cyclodepsipeptides against two other serine
proteases, elastase and proteinase K, which are of biomedical and ecological relevance,
respectively.

While **2** was found to be the most potent elastase inhibitor (IC$_{50}$ = 13.95 ± 0.36 nM), **6**
was the most potent proteinase K inhibitor (IC$_{50}$ = 3.83 ± 0.37 nM). The isolated
compounds were docked by induced-fit, inside the binding pocket of alpha-chymotrypsin
(PDBID 4Q2K) and all were found to have a similar binding mode (**Figure S35**) that was
revealed by crystal structures of Ahp-cyclodepsipeptides in complex with serine
proteases[50,51].

Figure 3: Structure elucidation workflow based on NMR and MS/MS data (a). The workflow combined automated *in-silico* MS/MS and NMR annotation tools for fast compound class identification and dereplication of known natural products exemplified for the new natural product rivulariapeptolide 1185. (b) Molecular networking and SMART analysis suggested the presence of a AHP-cyclodepsipeptide molecular family. (c) In depth MS and MS/MS analysis of the new natural products with the SIRIUS tool framework helped to establish the molecular formula and to gain substructural information about the *N*-acylated proline residues, characteristic for the rivulariapeptolide subfamily. (d) Unambiguous structure elucidation by various 1D/2D NMR and MS/MS experiments led to the planar structure of rivulariapeptolide 1185. Selected 2D NMR-derived correlations and MS² fragmentations are depicted.

a

b

c

Compound		R_1	R_2	R_3	R_4	IC$_{50}$ Chymo-trypsin [nM]	IC$_{50}$ Elastase [nM]	IC$_{50}$ Proteinase K [nM]
Rivulariapeptolide 1185	(**1**)	*N*-Ba-Pro-Thr-	-CH-CH$_2$-iPr	Ph		8.50 ± 0.5	19.6 ± 1.3	40.5 ± 3.9
Rivulariapeptolide 1155	(**2**)	*N*-Ba-Pro-Thr-	-C=CH-CH$_3$	Ph		28.2 ± 2.2	3.95 ± 0.4	39.8 ± 4.1
Rivulariapeptolide 1121	(**3**)	*N*-Ba-Pro-Thr-	-C=CH-CH$_3$	iPr		26.3 ± 3.1	10.9 ± 1.4	32.6 ± 3.5
Rivulariapeptolide 988	(**4**)	H	-C=CH-CH$_3$	Ph		63.6 ± 2.6	13.4 ± 2.2	58.8 ± 5.6
Molassamide	(**5**)				H	503.2 ± 33.6	31.3 ± 2.4	15.7 ± 0.6
Molassamide B	(**6**)				Br	16.5 ± 1.0	9.1 ± 0.8	3.8 ± 0.4

Figure 4: (a) Structures of the isolated rivulariapeptolides 1185 (**1**), 1155 (**2**), 1121 (**3**), 988 (**4**). (b) Structures of the isolated known molassamide (**5**), and the new molassamide B (**6**). (b) Potency of isolated compounds for selected serine proteases following 40 min pre-incubations. Data are presented as the mean ± SD, n = 3.

Discussion

Here we describe the use of native metabolomics protein infusion MS to simultaneously detect protein-metabolite binding and annotate their molecular structures. This approach can be used for rapid screening of small molecule modulators for proteins of interest, directly from crude extracts. In our case study, we identified 30 chymotrypsin binding natural products in a 10 min LC-MS run from a few µg of crude extract. This process would have taken month to years via traditional bioactivity-guided natural product discover approaches, and most likely would not have been possible to this extent.

The amounts of protein needed for native metabolomics high-throughput screening are a

few mg for a 24 h screen of 96 samples with a 10 min LC-MS method, which is both

achievable and affordable with commercially available proteins or in house heterologous

protein expression.

Besides the ad-hoc experimental determination of binding information, we systematically

organized this information in the public GNPS spectral library through spectrum tags.

Hence, native metabolomics derived properties are accessible in future experiments and

can provide biological context to complex metabolomes.

While we primarily used the method for an initial screening approach and assigned binary

binding information (binder/non-binder), titration experiments using native mass

spectrometry can be used to determine relative dissociation constants (K_d) by fitting the

intensity ratio of bound and unbound protein as a function of the added ligand. This

method first assumes that no dissociation takes place during the transmission through

the mass spectrometer, and second, that the observed gas phase intensity ratio

correlates with the solution ratio. These assumptions imply that ESI titration

measurements can deliver a relative "snapshot" of the solution concentrations to reflect

solution-phase binding affinities. Nevertheless, as the experimental environment is

inherently different (gas phase vs. solution) absolute binding affinities might differ from

solution based orthogonal assays[52].

From a natural product discovery perspective, it is very interesting that 30 putative

bioactive molecules were discovered in a single extract. This indicates that the chemical

space for certain bioactive molecular families can often be underestimated when

compared to traditional bioactivity-guided approaches, as they are typically biased

towards the most abundant or most active compounds. At least for Ahp-

cyclodepsipeptides, recent biosynthetic studies suggest that the high structural diversity

of these compounds is mainly driven by the hypervariability of amino acids in the positions

proceeding and following the Ahp-unit (see **Figure 4a** for the definitions of residues R_3

and R_2, respectively)[53]. The events impacting R_2 can be explained by high-frequency

point mutations. This is sought to provide an evolutionary platform to iteratively test

combinations while maintaining the central activity. However, the amino acid substitutions

at R_3 most likely occur via recombination events, thereby allowing for evolutionary

shortcuts[53]. These biosynthetic hypotheses are supported by the compounds isolated in

this study and add to a better understanding of structure-activity relationships for Ahp-

cyclodepsipeptides. Comparing rivulariapeptolide 1185 (**1**) to rivulariapeptolide 1121 (**3**),

a leucine residue is swapped for an Abu unit at R_2, and phenylalanine is replaced by

leucine at R_3. The most surprising structure-activity relationship gained from this study,

however, was that a single substitution of bromine (molassamide B, **6**) for hydrogen

(molassamide, **5**) led to a thirty-, three-, and four-fold increase in potency towards

chymotrypsin, elastase, and proteinase K, respectively. This finding, together with the fact

that several additional brominated Ahp-peptides but no chlorinated Ahp-peptides could

be identified (**Figure S36**), indicates the need for further structure-activity investigations

as well as biosynthetic studies that characterize the responsible halogenases.

The protease inhibition of the compounds discovered with the native metabolomics

workflow was confirmed with an orthogonal fluorescence assay against three proteases.

At nanomolar concentration their IC_{50} show high potency and exhibit distinct selectivity

(**Figure 4c, Tables S4 and S5**). For example, molassamide (**5**) is the second most potent

inhibitor screened against proteinase K but is the least potent inhibitor for chymotrypsin

and elastase. Rivulariapeptolide 1155 (**2**), on the other hand, is the most potent elastase

inhibitor but shows much lower inhibition against both chymotrypsin and proteinase K.

Together, these findings highlight the utility of the native metabolomics approach

presented herein. Beyond the discovery of novel protease inhibitors, we anticipate that

native metabolomics will be applied for the screening of a broad variety of non-covalent

interactions of biomolecules from complex mixtures and could be of central importance

for the development of activity/ functional metabolomics workflows, both for drug

discovery as well as chemical ecology studies.

Methods

Cyanobacterial collection and taxonomy

Marine cyanobacteria biofilm samples were collected in an intertidal zone growing on rock/reef substrate on May 14th, 2019, and at $0.5 − 2.0$ m of water at Carlos Rosario Beach in Culebra, Puerto Rico, U.S (GPS coordinates: 18°19'30.0"N 65°19'48.0"W) on April 6th, 2019. Biomass for both samples was hand collected (DRNA Permit O-VS-PVS15-SJ-01165-15102020). Microscopic examination indicated that this collection was morphologically consistent with the genus *Rivularia*. 16S rDNA analysis confirmed the identity as *Rivularia* spp. PCC 7116. Voucher specimen available from E.C.D. as collection no. MAP14MAY19-1, and from W. H. G. as collection no. CUR6APR19-1.

Micro-flow LC-MS/MS data acquisition

For micro-flow UHPLC-MS/MS analysis 2 µL were injected into vanquish UHPLC system coupled to a Q-Exactive quadrupole orbitrap mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) with an Agilent 1260 quaternary HPLC pump (Agilent, Santa Clara, USA) as a make-up pump. For reversed-phase chromatographic, a C18 core-shell microflow column (Kinetex C18, 150 x 1 mm, 1.8 um particle size, 100 A pore size, Phenomenex, Torrance, USA) was used. The mobile phase consisted of solvent A (H_2O + 0.1 % formic acid (FA)) and solvent B (acetonitrile (ACN) + 0.1 % FA). The flow rate was set to 150 µL/min. A linear gradient from 5-50 % B between 0-4 min and 50-99 % B between 4 and 5 min, followed by a 2 min washout phase at 99% B and a 3 min re-equilibration phase at 5 % B. Data-dependent acquisition (DDA) of MS/MS spectra was performed in positive mode. Electrospray ionization (ESI) parameters were set to 40 arbitrary units (AU) sheath gas flow, auxiliary gas flow was set to 10 AU and sweep gas flow was set to 0 AU. Auxiliary gas temperature was set to 400 °C. The spray voltage was set to 3.5 kV and the inlet capillary was heated to 320 °C. S-lens level was set to 70 V applied. MS scan range was set to 200-2000 m/z with a resolution at m/z 200 ($R_{m/z\ 200}$) of 70,000 with one micro-scan. The maximum ion injection time was set to 100 ms with automatic gain control (AGC) target of 5E5. Up to two MS/MS spectra per duty cycle were acquired at $R_{m/z\ 200}$ 17,000 with one micro-scan. The maximum ion injection time for MS/MS scans was set to 100 ms with an AGC target of 5.0E5 ions and a minimum 5%

AGC. The MS/MS precursor isolation window was set to m/z 1. The normalized collision

energy was stepped from 20 to 30 to 40% with $z = 1$ as the default charge state. MS/MS

scans were triggered at the apex of chromatographic peaks within 2 to 15 s from their first

occurrence. Dynamic precursor exclusion was set to 5 s. Ions with unassigned charge

states were excluded from MS/MS acquisition as well as isotope peaks. For native

metabolomics experiments, the same chromatographic parameters were used and in

addition 150 μL/min ammonium acetate buffer was infused post-column through a make-

up pump and a PEEKT-splitter and enzyme solution was infused with 2 μL/min flow rate

via a syringe pump. ESI settings were set to 40 arbitrary units (AU) sheath gas flow,

auxiliary gas flow was set to 10 AU and sweep gas flow was set to 0 AU. Auxiliary gas

temperature was set to 400 °C. The spray voltage was set to 3.5 kV and the inlet capillary

was heated to 300 °C. S-lens level was set to 80 V applied. MS scan range was set to

2000-4000 m/z with a resolution $R_{m/z\ 200}$ 140,000 with 2 microscans. MS acquisition was

performed in all ion fragmentation (AIF) mode with $R_{m/z\ 200}$ with 20% HCD collision energy

and an isolation window of 2000 - 4000 m/z. To estimate binding strength a second MS

data acquisition method was used that ramped HCD collision energy in 4 iterative scans

with 0%, 20%, 40% and 60% per duty cycle.

Native metabolomics data analysis

For native LC-MS data, multiple charged spectra were deconvoluted using the xtract

algorithm in Qualbrowser, part of the Xcalibur software (Thermo Scientific). Both

deconvoluted native LC-MS and metabolomics LC-MS/MS .raw files were converted to

centroid .mzML file format using MSconvert of the proteowizard software package.

Feature finding of both file types was performed using a modified version of MZmine2.37

(corr.17.7). Feature tables from both intact protein mass and metabolomics data were

matched by their retention time (RT) and an m/z offset corresponding to the mass of

chymotrypsin (25234 Da) with an RT tolerance of 0.2 min and a mass tolerance of 4 Da.

Feature tables (.csv), MS/MS spectra files (.mgf), and ion identity networking results (.csv)

were exported and uploaded to the MassIVE repository. LC-MS/MS data was submitted

to GNPS for feature-based molecular networking analysis. Downstream combined

Molecular-Networks and chymotrypsin small molecule binding were visualized as
networks in cytoscape (3.8.2).

pH dependency of native metabolomics

pH values over the entire LC gradient and peak intensities were both assessed using
three different make-up solvents. Make-up solvent A was water, make-up solvent B was
10 mM ammonium acetate buffer, and make-up solvent C was 10 mM ammonium acetate
buffer + 0.2% ammonium hydroxide, v/v. Molassamide was prepared as 100 µM solutions
from a 10 mM stock solution in DMSO by preparing a 1:100 dilution into solvent mixture
1 (water + 10% acetonitrile + 0.1% formic acid). Samples were analyzed as described in
micro-flow LC-MS/MS data acquisition; 2 µL of each solution were injected into the mass
spectrometer, while chymotrypsin (Sigma),dissolved in water to a final concentration of 2
mg/mL, was injected through the syringe pump at a flow rate of 2 µL/min. pH values were
assessed by disconnecting the flow to the source and collecting ~30 µL of solvent every
minute then testing this value on pH paper.

Titration of ligands and concentration dependency

Molassamide, Riv 1155, and Riv 1185 were prepared as 100 µM solutions from a 10 mM
stock solution in DMSO by preparing a 1:100 dilution into solvent mixture 1 (water + 10%
acetonitrile + 0.1% formic acid). From this 100 µM solution, dilutions were prepared at 10
µM, 1 µM, and 0.1 µM into solvent mixture 1. 2 µL of each solution were injected into the
mass spectrometer, then 5 µL and 10 µL of the 100 µM solution were injected to yield
final concentrations of 250 µM and 500 µM, respectively. Samples were analyzed as
described in Micro-Flow LC-MS/MS data acquisition, while chymotrypsin (Sigma Aldrich)
was dissolved in water to a final concentration of 2 mg/mL and injected through the
syringe pump at a flow rate of 2 µL/min. The ratio of bound to unbound protein was plotted
against the ligand concentration in a given HPLC peak window. Data points were fitted
using the solver function.

Chymotrypsin activity assays in native mass spectrometry buffer

Cyanobacteria extract in (1 mg/ml, methanol) was diluted in 10 mM ammonium acetate pH 4.5 to 30 ug/mL and then sequentially diluted 1.5-fold to 0.52 µg/mL in the same buffer. Bovine chymotrypsin (Sigma Aldrich) and Z-Ala-Ala-Pro-Phe-AMC (Calbiochem, 230914) were diluted to 300 nM and 150 µM, respectively in 10 mM ammonium acetate pH 4.5. In a 384-well black microplate, 10 µL of enzyme, substrate and cyanobacteria extract were combined (30 µL final volume) such that the concentrations in the reaction were 100 nM chymotrypsin, 50 µM of Z-Ala-Ala-Pro-Phe-AMC and 10 µM to 0.17 nM of cyanobacteria extract. For solvent compatibility assays, chymotrypsin (100 nM) was assayed with 10 µg/mL and 1 µg/mL extract in 10 mM ammonium acetate, pH 4.5 containing 8.3 to 41.7% acetonitrile. A control assay lacked acetonitrile and cyanobacteria extract. All assays were performed in triplicate wells at 25°C in a Synergy HTX Multi-Mode Microplate Reader (BioTek, Winooski, VT) with excitation and emission wavelengths of 360 and 460 nm, respectively. The initial reaction velocity in each well was recorded and the dose response curve was generated using GraphPad Prism 9 software.

Calculation of Km and inhibition constants

To determine Km value of Z-Ala-Ala-Pro-Phe-AMC substrate, chymotrypsin was assayed with 1.95 to 500 µM of the substrate in Dulbecco's phosphate buffered saline, pH 7.4 containing 0.01% Tween-20 in a final volume of 30 µL. Kinetic measurements were performed in triplicate wells at 25°C. The result Vmax values were plotted against substrate concentration and Km was calculated using the Michaelis-Menten model in GraphPad Prism 9. For inhibition studies, chymotrypsin (1 nM) was preincubated with 0 to 3 µM of each compound for 40 min in Dulbecco's phosphate buffered saline, pH 7.4 containing 0.01% Tween-20. The reaction was initiated by addition of 25 µM of Z-Ala-Ala-Pro-Phe-AMC (Calbiochem, 230914) in a final volume of 30 µL, respectively. The release of the AMC fluorophore was recorded in a Synergy HTX multi-mode reader (BioTek Instruments, Winooski, VT) with excitation and emission wavelengths at 340 nm and 460 nm, respectively. The maximum velocity was calculated in RFU/sec over 10 sequential points on the linear part of the progress curve. The IC50 values were determined by

nonlinear regression and Ki values were determined using the Morrison equation in
GraphPad Prism 9.

NMR spectroscopy

Deuterated NMR solvents were purchased from Cambridge Isotope Laboratories. 1H
NMR and 2D NMR spectra were collected on a Bruker Avance III DRX-600 NMR with a
1.7 mm dual tune TCI cryoprobe (600 and 150 MHz for 1H and 13C NMR, respectively)
and a JEOL ECZ 500 NMR spectrometer equipped with a 3 mm inverse detection probe.
NMR spectra were referenced to residual solvent DMSO signals (δH 2.50 and δC 39.5
as internal standards). The NMR spectra were processed using MestReNova (Mnova
12.0, Mestrelab Research) or TopSpin 3.0 (Bruker Biospin) software.

Extraction and isolation

The preserved cyanobacterial biomass from collection no. CUR6APR19-1 was filtered
through cheesecloth, and then (98.7 g dry wt) was extracted repeatedly by soaking in 500
mL of 2:1 CH_2Cl_2 /MeOH with warming (<30 °C) for 30 min to afford 1.44 g of dried extract.
A portion of the extract was fractionated by reverse-phase solid phase extraction (C_{18}-
SPE) using a stepwise gradient solvent system of decreasing polarity (Fr. 1-1 35%
ACN/H_2O, 124.4 mg; Fr. 1-2 70% ACN/H_2O, 76.1 mg; Fr. 1-3 100% ACN, 77 mg; Fr. 1-4
100% MeOH, 254.9 mg. Fr. 1-2 was dissolved in 70% ACN/H2O and purified by
preparative HPLC using a Kinetex 5 µm RP 100 Å column (21.00 × 150mm) and isocratic
elution using 50% ACN/H2O for 8 minutes then ramping up to 100% in 14 minutes at the
flow rate of 20 mL/min, yielding 56 subfractions. Rivulariapeptolide 1185 (compound **1**)
and 1155 (compound **2**) were isolated from subfractions 1-2-20 to 1-2-23 that were
combined (4.5 mg) and further purified by semi-preparative HPLC using a Synergi 4 µm
Hydro-RP 80 Å column (10.00 × 250 mm) and isocratic elution gradient elution using 35%
ACN / 65% H2O isocratic at the flow rate of 3.5 mL/min for 3 minutes the ramping up to
55% ACN in 22 minutes, then ramping up to 100% ACN in one minute and holding the
gradient at 100% ACN for another 5 minutes yielding **1** (1.6 mg, RT = 23.3 min) and **2**
(1.3 mg, RT =21.8 min) as a colorless, amorphous solid. The same HPLC conditions were
used to isolate compounds **3** (rivulariapeptolide 1121, from subfraction 1-2-10 and 1-2-

11, 1.1 mg, RT= 18.5min) and **4** (rivulariapeptolide 988, from subfraction 1-2-7, 1.3 mg, RT = 12.6 min). Fr. 1-1 was dissolved in 30% ACN/H2O and purified by preparative HPLC using a Kinetex 5 µm RP 100 Å column (21.00 × 150mm) and isocratic elution using 30% ACN/H2O for 10 minutes then ramping up to 50% in 10 minutes and then to 95% in 2 min at the flow rate of 20 mL/min, yielding 29 subfractions. Molassamide (compound **5**) was isolated from subfractions 1-1-6 and 2-1-10 were combined (3.3 mg) and further purified by semi-preparative HPLC using a Synergi 4 µm Hydro-RP 80 Å column (10.00 × 250 mm) and isocratic elution gradient elution using 35% ACN / 65% H2O isocratic at the flow rate of 3.5 mL/min for 3 minutes the ramping up to 55% ACN in 22 minutes, then ramping up to 100% ACN in one minute and holding the gradient at 100% ACN for another 5 minutes yielding **5** as a colorless, amorphous solid (1.8 mg) at RT = 10.8 min. The same HPLC conditions were used to isolate compound **6** (Molassamide B, from subfraction 1-1-4 and 1-1-5, 1.8 mg, RT = 13.7 min).

Data Availability

All raw (.raw), deconvoluted (xtract.raw) and centroided (.mzXML or .mzML) mass spectrometry data as well as processed data feature table (.csv) and MS/MS spectra (.mgf) are available through the MassIVE repository (massive.ucsd.edu) with the identifier MSV000087964 and can be accessed under the following link: https://massive.ucsd.edu/ProteoSAFe/dataset.jsp?task=7ab346a09ca64aa1bd19bdc03 5801c15. The MS/MS spectra of the new discovered derivatives, including tags as protease inhibitors, have been added to the GNPS library (gnps.ucsd.edu) with the following IDs: rivulariapeptolide 1185 (**1**): CCMSLIB00005723387; rivulariapeptolide 1155 (**2**): CCMSLIB00005723986, CCMSLIB00005720236; rivulariapeptolide 1121 (**3**): CCMSLIB00005723398; rivulariapeptolide 988 (**4**): CCMSLIB00005723393; molassamide (**5**): CCMSLIB00005723404; molassamide B (**6**): CCMSLIB00006710020. Raw NMR data for compounds **1** -**6** has been deposited to Zenodo (zenodo.org) and can be accessed under the following link: https://sandbox.zenodo.org/record/905199.

Code Availability

The modified version of MZmine2.37 (corr.17.7) used in this study is available at https://github.com/robinschmid/mzmine2/releases. The code for the mass-offset matching for native metabolomics data analysis is available under https://github.com/Functional-Metabolomics-Lab/Native-Metabolomics.

References

1. Newman, D. J. & Cragg, G. M. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* **83**, 770–803 (2020).
2. Pye, C. R., Bertin, M. J., Lokey, R. S., Gerwick, W. H. & Linington, R. G. Retrospective analysis of natural products provides insights for future discovery trends. *PNAS* **114**, 5601–5606 (2017).
3. Gavriilidou, A. *et al. A global survey of specialized metabolic diversity encoded in bacterial genomes*. http://biorxiv.org/lookup/doi/10.1101/2021.08.11.455920 (2021) doi:10.1101/2021.08.11.455920.
4. Medema, M. H., de Rond, T. & Moore, B. S. Mining genomes to illuminate the specialized chemistry of life. *Nat Rev Genet* **22**, 553–571 (2021).
5. Hooft, J. J. J. van der *et al.* Linking genomics and metabolomics to chart specialized metabolic diversity. *Chem. Soc. Rev.* **49**, 3297–3314 (2020).
6. Peisl, B. Y. L., Schymanski, E. L. & Wilmes, P. Dark matter in host-microbiome metabolomics: Tackling the unknowns–A review. *Analytica Chimica Acta* **1037**, 13–27 (2018).
7. Scheubert, K. *et al.* Significance estimation for large scale metabolomics annotations by spectral matching. *Nat Commun* **8**, 1494 (2017).
8. Schmid, R. *et al.* Ion identity molecular networking for mass spectrometry-based metabolomics in the GNPS environment. *Nat Commun* **12**, 3832 (2021).
9. Dührkop, K. *et al.* Systematic classification of unknown metabolites using high-resolution fragmentation mass spectra. *Nat Biotechnol* **39**, 462–471 (2021).
10. Ludwig, M. *et al.* Database-independent molecular formula annotation using Gibbs sampling through ZODIAC. *Nat Mach Intell* **2**, 629–641 (2020).
11. Reher, R. *et al.* A Convolutional Neural Network-Based Approach for the Rapid Annotation of Molecularly Diverse Natural Products. *J. Am. Chem. Soc.* **142**, 4114–4120 (2020).
12. Stravs, M. A., Dührkop, K., Böcker, S. & Zamboni, N. *MSNovelist: De novo structure generation from mass spectra*. 2021.07.06.450875 https://www.biorxiv.org/content/10.1101/2021.07.06.450875v1 (2021) doi:10.1101/2021.07.06.450875.

13. Rinschen, M. M., Ivanisevic, J., Giera, M. & Siuzdak, G. Identification of bioactive
metabolites using activity metabolomics. *Nat Rev Mol Cell Biol* **20**, 353–367 (2019).
14. Yan, M. & Xu, G. Current and future perspectives of functional metabolomics in
disease studies–A review. *Analytica Chimica Acta* **1037**, 41–54 (2018).
15. Su, H. *et al.* Anti-SARS-CoV-2 activities in vitro of Shuanghuanglian preparations
and bioactive ingredients. *Acta Pharmacol Sin* **41**, 1167–1177 (2020).
16. Bolla, J. R., Agasid, M. T., Mehmood, S. & Robinson, C. V. Membrane Protein–Lipid
Interactions Probed Using Mass Spectrometry. *Annu. Rev. Biochem.* **88**, 85–111
(2019).
17. Liu, M., Van Voorhis, W. C. & Quinn, R. J. Development of a target identification
approach using native mass spectrometry. *Sci Rep* **11**, 2387 (2021).
18. Gault, J. *et al.* High-resolution mass spectrometry of small molecules bound to
membrane proteins. *Nat Methods* **13**, 333–336 (2016).
19. Yen, H.-Y. *et al.* PtdIns(4,5)P2 stabilizes active states of GPCRs and enhances
selectivity of G-protein coupling. *Nature* **559**, 423–427 (2018).
20. Yen, H.-Y. *et al.* Ligand binding to a G protein–coupled receptor captured in a mass
spectrometer. *Science Advances* **3**, e1701016 (2017).
21. Vu, H. *et al.* Plasmodium Gametocyte Inhibition Identified from a Natural-Product-
Based Fragment Library. *ACS Chem. Biol.* **8**, 2654–2659 (2013).
22. Gault, J. *et al.* Combining native and 'omics' mass spectrometry to identify
endogenous ligands bound to membrane proteins. *Nat Methods* **17**, 505–508
(2020).
23. Sumner, L. W. *et al.* Proposed minimum reporting standards for chemical analysis.
Metabolomics **3**, 211–221 (2007).
24. Schymanski, E. L. *et al.* Identifying Small Molecules via High Resolution Mass
Spectrometry: Communicating Confidence. *Environ. Sci. Technol.* **48**, 2097–2098
(2014).
25. Hong, J. & Luesch, H. Largazole: From discovery to broad-spectrum therapy. *Nat.*
Prod. Rep. **29**, 449–456 (2012).
26. Gerwick, W. H. *et al.* Structure of Curacin A, a Novel Antimitotic, Antiproliferative
and Brine Shrimp Toxic Natural Product from the Marine Cyanobacterium Lyngbya
majuscula. *The Journal of Organic Chemistry* **59**, 1243–1245 (1994).
27. Boudreau, P. D. *et al.* Design of Gallinamide A Analogs as Potent Inhibitors of the
Cysteine Proteases Human Cathepsin L and *Trypanosoma cruzi* Cruzain. *J. Med.*
Chem. **62**, 9026–9044 (2019).
28. Luesch, H., Yoshida, W. Y., Moore, R. E., Paul, V. J. & Corbett, T. H. Total Structure
Determination of Apratoxin A, a Potent Novel Cytotoxin from the Marine
Cyanobacterium *Lyngbya m ajuscula*. *Journal of the American Chemical Society*
123, 5418–5423 (2001).

29. Köcher, S. *et al.* From dolastatin 13 to cyanopeptolins, micropeptins, and
lyngbyastatins: the chemical biology of Ahp-cyclodepsipeptides. *Nat. Prod. Rep.*
(2019) doi:10.1039/C9NP00033J.
30. Lodin-Friedman, A. & Carmeli, S. Microginins from a Microcystis sp. Bloom Material
Collected from the Kishon Reservoir, Israel. *Marine Drugs* **16**, 78 (2018).
31. Murakami, M. *et al.* Microviridins, elastase inhibitors from the cyanobacterium
Nostoc minutum (NIES-26). *Phytochemistry* **45**, 1197–1202 (1997).
32. Nostocyclamide: A new macrocyclic, thiazole-containing allelochemical from Nostoc
sp. 31 (cyanobacteria) | The Journal of Organic Chemistry.
https://pubs.acs.org/doi/pdf/10.1021/jo00129a032.
33. Issac, M. *et al.* Cyclotheonellazoles A–C, Potent Protease Inhibitors from the Marine
Sponge Theonella aff. swinhoei. *J. Nat. Prod.* **80**, 1110–1116 (2017).
34. Adiv, S. & Carmeli, S. Protease Inhibitors from Microcystis aeruginosa Bloom
Material Collected from the Dalton Reservoir, Israel. *ACS Publications*
https://pubs.acs.org/doi/pdf/10.1021/np4006844 (2013) doi:10.1021/np4006844.
35. Anderson, J., Schiffer, C., Lee, S.-K. & Swanstrom, R. Viral Protease Inhibitors. in
Antiviral Strategies (eds. Kräusslich, H.-G. & Bartenschlager, R.) 85–110 (Springer,
2009). doi:10.1007/978-3-540-79086-0_4.
36. Ito, S. Proteasome Inhibitors for the Treatment of Multiple Myeloma. *Cancers* **12**,
265 (2020).
37. Deacon, C. F. A review of dipeptidyl peptidase-4 inhibitors. Hot topics from
randomized controlled trials. *Diabetes, Obesity and Metabolism* **20**, 34–46 (2018).
38. Messerli, F. H., Bangalore, S., Bavishi, C. & Rimoldi, S. F. Angiotensin-Converting
Enzyme Inhibitors in Hypertension. *Journal of the American College of Cardiology*
71, 1474–1482 (2018).
39. Lee, C. J. & Ansell, J. E. Direct thrombin inhibitors. *British Journal of Clinical*
Pharmacology **72**, 581–592 (2011).
40. Al-Awadhi, F. H. & Luesch, H. Targeting eukaryotic proteases for natural products-
based drug development. *Nat. Prod. Rep.* **37**, 827–860 (2020).
41. Molassamide, a Depsipeptide Serine Protease Inhibitor from the Marine
Cyanobacterium Dichothrix utahensis | Journal of Natural Products.
https://pubs.acs.org/doi/10.1021/np900603f.
42. Linington, R. G. *et al.* Symplocamide A, a Potent Cytotoxin and Chymotrypsin
Inhibitor from the Marine Cyanobacterium Symploca sp. *J. Nat. Prod.* **71**, 22–27
(2008).
43. Miller, B. *et al.* The Marine Cyanobacterial Metabolite Gallinamide A Is a Potent and
Selective Inhibitor of Human Cathepsin L. *Journal of Natural Products* **77**, 92–99
(2014).
44. H. Medema, M. The year 2020 in natural product bioinformatics: an overview of the
latest tools and databases. *Natural Product Reports* **38**, 301–306 (2021).

45. Nogle, L. M., Williamson, R. T. & Gerwick, W. H. Somamides A and B, Two New
Depsipeptide Analogues of Dolastatin 13 from a Fijian Cyanobacterial Assemblage
of Lyngbya majuscula and Schizothrix Species. *J. Nat. Prod.* **64**, 716–719 (2001).
46. Al-Awadhi, F. H., Paul, V. J. & Luesch, H. Structural Diversity and Anticancer
Activity of Marine-Derived Elastase Inhibitors: Key Features and Mechanisms
Mediating the Antimetastatic Effects in Invasive Breast Cancer. *ChemBioChem* **19**,
815–825 (2018).
47. Taori, K., Matthew, S., Rocca, J. R., Paul, V. J. & Luesch, H. Lyngbyastatins 5–7,
Potent Elastase Inhibitors from Floridian Marine Cyanobacteria, Lyngbya spp. *J.
Nat. Prod.* **70**, 1593–1600 (2007).
48. Dührkop, K. *et al.* SIRIUS 4: a rapid tool for turning tandem mass spectra into
metabolite structure information. *Nat Methods* **16**, 299–302 (2019).
49. Ludwig, M. *et al.* Database-independent molecular formula annotation using Gibbs
sampling through ZODIAC. *Nat Mach Intell* **2**, 629–641 (2020).
50. Salvador, L. A. *et al.* Potent Elastase Inhibitors from Cyanobacteria: Structural Basis
and Mechanisms Mediating Cytoprotective and Anti-Inflammatory Effects in
Bronchial Epithelial Cells. *J. Med. Chem.* **56**, 1276–1290 (2013).
51. Keller, L. *et al.* Tutuilamides A–C: Vinyl-Chloride-Containing Cyclodepsipeptides
from Marine Cyanobacteria with Potent Elastase Inhibitory Properties. *ACS Chem.
Biol.* **15**, 751–757 (2020).
52. Leney, A. C. & Heck, A. J. R. Native Mass Spectrometry: What is in the Name? *J.
Am. Soc. Mass Spectrom.* **28**, 5–13 (2017).
53. Baunach, M., Chowdhury, S., Stallforth, P. & Dittmann, E. The Landscape of
Recombination Events That Create Nonribosomal Peptide Diversity. *Molecular
Biology and Evolution* **38**, 2116–2130 (2021).
668

Acknowledgment

We thank the Deutsche Forschungsgemeinschaft for the support of D.P. through a
postdoctoral research fellowship (PE 2600/1-1) and through the CMFI Cluster of
Excellence (EXC 2124). R.R., P.C.D., and W.H.G. were supported by the Gordon and
Betty Moore Foundation (GBMF7622) and by the US National Institutes of Health (R01
GM107550, P41 GM103484 and R03 CA211211). WB was supported in part by the
Research Foundation – Flanders (12W0418N).

677
678
679
680

Author information

Affiliations

Scripps Institution of Oceanography, University of California San Diego, USA

Raphael Reher, Kelsey L Alexander, William H Gerwick, Daniel Petras

Skaggs School of Pharmacy and Pharmaceutical Science, University of California San Diego, USA

Allegra T Aron, Pavla Fajtova, Chenxi Liu, Ido Y Ben Shalom, Wout Bittremieux, Mingxun Wang, Pieter C Dorrestein, Anthony J O'Donoghue, William H Gerwick, Daniel Petras

Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Halle (Saale), Germany

Raphael Reher

Department of Pharmaceutical Sciences, School of Pharmacy, University of Puerto Rico - Medical Sciences Campus, San Juan, Puerto Rico

Marie L Matos-Hernandez, Eduardo J Caro-Diaz

Li Dak Sum Yip Yio Chin Kenneth Li Marine Biopharmaceutical Research Center, Department of Marine Pharmacy, College of Food and Pharmaceutical Sciences, Ningbo University 315832, Ningbo, China

C Benjamin Naman

Department of Chemistry and Biochemistry, University of California San Diego, USA

Kelsey L Alexander

CMFI Cluster of Excellence, Interfaculty Institute of Microbiology and Medicine, University of Tuebingen, Germany

Daniel Petras

Contributions

R.R., W.H.G., and D.P. conceived the study. R.R., K.L.A., C.B.N., E.J.C., and W.H.G. collected and extracted environmental samples. A.T.A., P.C.D., and D.P. developed the native metabolomics approach. W.B. wrote software code. M.W. aided in integration with GNPS tags. R.R., A.T.A., and D.P. performed MS experiments. R.R. and M.L.M. performed compound isolation and R.R. carried out NMR experiments. P.F., C.L., and A.J.O. performed activity assays. I.B.S. performed docking studies. R.R., W.H.G., and D.P. wrote the manuscript. All authors edited and approved the final manuscript.

Corresponding authors

Correspondence should be addressed to Daniel Petras (daniel.petras@uni-tuebingen.de) for questions regarding native metabolomics and MS-based structure elucidation, and to William Gerwick (wgerwick@health.ucsd.edu) for question regarding cyanobacteria sampling and NMR-based structure elucidation.

Ethics declarations

Competing interests

P.C.D. and W.H.G. are scientific advisors of Sirenas. P.C.D. is a scientific advisor of Galileo, Cybele, and scientific advisor and co-founder of Ometa Labs LLC and Enveda with approval by the UC San Diego. M.W. is a founder of Ometa Labs LLC.

Supplementary information

Supplementary Results, Methods, Figures, and NMR tables.