A Causal Perspective for Batch Effects: when is no answer better than a wrong answer?

Eric W. Bridgeford[1,†], Michael Powell[1], Gregory Kiar[2], Stephanie Noble[3], Jaewon Chung[1], Sambit Panda[1], Ross Lawrence[1], Ting Xu[2], Michael Milham[2], Brian Caffo[1], Joshua T. Vogelstein[1]

Abstract. Batch effects, undesirable sources of variance across multiple experiments, present significant challenges for scientific and clinical discoveries. Specifically, batch effects can (i) produce spurious, and/or (ii) obscure genuine signals, contributing to the ongoing reproducibility crisis. Typically, batch effects are modeled as classical, rather than causal, statistical effects. This model choice renders the methods unable to differentiate between biological or experimental sources of variability, leading to unnecessary false positive and negative effect detections and overconfidence. We formalize batch effects as causal effects to address these concerns, and augment existing batch effect detection and correction approaches with causal machinery. Simulations illustrate that our causal approaches mitigate spurious findings and reveal otherwise obscured signals as compared to non-causal approaches. Applying our causal methods to a large neuroimaging mega-study reveals instances where prior art confidently asserts that the data do not support the presence of batch effects when we expect to detect them. On the other hand, our causal methods correctly discern that there exists irreducible confounding in the data, so it is unclear whether differences are due to batches or not. Moreover, we illustrate that causal and non-causal batch effect correction techniques produce radically different subsequent inferences, underlying the importance of causal modeling of batches. This work therefore provides a framework for understanding the potential capabilities and limitations of analysis of multi-site data.

1 Introduction

The 21[st] century has seen the advent of high-throughput techniques for acquiring data on an unprecedented scale. Collection of these datasets often occurs through consortia across different sites, requiring post-hoc data aggregation approaches. These "mega-studies" comprising numerous individual studies rendering samples sizes substantially larger than any individual study, addressing the small-sample size woes associated with modern big biomedical data [36].

Unfortunately, aggregating data across diverse datasets introduces a source of undesirable variability known as a batch effect. Lazar et al. [32] provides a recent consensus definition of batch effect: "the batch effect represents the systematic technical differences when samples are processed and measured in different batches and which are unrelated to any biological variation." Unfortunately, this unified description provides limited information about how batch effects can be detected or mitigated.

Broadly, many approaches model the batch collection or measurement process as a nuisance variable which is to be removed [28, 33, 34, 46, 48, 62, 67], which implies batch effect is a particular component of an associational or conditional model. These approaches typically make strong, potentially unjustified, and often inappropriate parametric assumptions about the data. Two of the most prominent examples of these techniques are `ComBat` and Conditional `ComBat` (`cComBat`) batch effect correction procedures [28]. These approaches have demonstrated empirical utility in various genomics and neuroimaging contexts [48, 68]; however, it remains unclear when these approaches will be successful and when they will fail. Specifically, it is unclear when they remove biofidelic variability or fail to remove nuisance variability [3].

In this work, we develop causal approach to define, detect, estimate, and mitigate batch effects. Our main conceptual advance is modeling batch effects as causal, rather than associational or conditional, effects. Given this understanding, we introduce a formal definition of causal batch effects. This formal definition reveals the limitations of (typically inappropriate) assumptions implicit in existing approaches [50, 51, 55], and provides a theoretical explanation for many limitations of batch harmonization noted in Bayer et al. [3].

Methodologically, we introduce a simple pre-processing strategy that one can apply to existing

---

[1] Johns Hopkins University [2] Child Mind Institute, [3] Yale University, Northeastern University, [†] Corresponding author: Eric W. Bridgeford (ebridge2@jhu.edu), 3400 N. Charles Street Clark Hall, Suite 200. Baltimore, MD, USA 21218.

techniques for batch effect detection and mitigation which addresses this new perspective. First, to detect and estimate batch effects, we introduce CAUSAL cDcORR [7, 63]—building on modern, non-parametric statistical methods. —to estimate and detect the presence of batch effects. Second, to mitigate batch effects, we introduce Causal cComBat—an augmentation of the ComBat procedure [28]—which uses causal methods to remove batch effects while limiting the removal of variation due to observed potentially confounding variables.

Our main empirical contribution is to apply these methods to simulations and a large neuroimaging mega-study assembled by the Consortium for Reliability and Reproducibility (CoRR) [69], consisting of more than $1,700$ individuals across 27 disparate studies featuring different measurement protocols. Our investigations reveal that previous strategies fail to differentiate between veridical signal and sources of variability due to other experimental design factors. Further, our simulations demonstrate that existing strategies can, under many realistic use-cases, either remove more apparent veridical biological signal, or fail to mitigate the batch effect entirely, than *Causal* ComBat.

A conceptual illustration of the value of causal modeling of batch effects Consider a simple example in Figure 1. We observe $n = 300$ measurements in one of two batches, where one batch (orange) tends to sample younger people, and the other batch (blue) tends to sample older people. The solid lines indicate the true data-generating distributions, which is unknown by batch effect correction strategies in practice. That the data generating distributions differ (red box) indicates that a batch effect is present. Instead, techniques must learn how to remove the batch effect given only the covariates and the outcomes (points). In this simplified scenario, age is the only covariate. In Figure 1**(A)**, we illustrate an example with almost no covariate overlap. ComBat style strategies learn from each batch, and then *extrapolate* trends (across age) to infer a relationship between the two batches. The problem is that this extrapolation approach requires, and is strongly sensitive to, model selection. That is, how does one perform the extrapolation?

As the true data generating distribution is unknown at the time of analysis, most ComBat style approaches [8, 17, 28, 48] tend to perform a linear extrapolation. When covariate overlap is imperfect, however, this misspecification can be disasterous. When covariate overlap is low in Figure 1**(A)**, cComBat strategies are subject to strong biases. While before correction the true data-generating distribution for the orange batch exceeds the blue batch, after correction the blue batch exceeds the orange batch (the batch effect was *over-corrected*; e.g., too much correction was applied). When covariate overlap is higher, but still imperfect in Figure 1**(B)** and Figure 1**(C)**, cComBat strategies are still unable to properly remove the batch effect, and may over or under-correct for batch effects. As a result, "batch effect corrected data" from cComBat may not actually be corrected, and in many situations may be more different after "correction" than before. This is particularly problematic in the era of big data mega-studies, in which determining the appropriateness (or lackthereof) of parametric techniques is nearly impossible to perform when the data may feature hundreds or thousands of dimensions.

In contrast, causal strategies avoid inference on data with extremely low overlap entirely because these situations have what is called 'irrecoverable confounding', as in Figure 1**III.(A)**. That is, there is no way to discern whether differences between the two samples arise from batch effects or veridical biological differences without making strong assumptions, which causal strategies correctly identify. On the other hand in Figure 1**III.(B)** or Figure 1**III.(C)**, causal techniques instead focus on deriving conclusions within a range of covariate overlap in the data, and therefore are not subject to the same extrapolatory biases. The true data-generating distributions (and the points themselves) almost perfectly align after batch effect correction across the range of overlapping covariates, indicating that the batch effect has been removed, despite no foreknowledge of the true data-generating model.

Figure 1: **Non-causal procedures are subject to model misspecifications and confounding biases**. **I.** shows the observed data (points), where color indicates batch. The orange line and the blue line indicated the expected outcome per-batch, and the "batch effect" describes the observed difference (red box) between the expected outcomes. Ideally, after batch effect correction, the blue and orange lines should overlap. The orange batch tends to oversample people with younger ages, and the blue batch tends to oversample people with higher ages. **(A)** a scenario where the covariate distributions are heavily confounded and do not overlap, and differentiating biological variability from batch variability is difficult without foreknowledge of the sigmoidal relationship. **(B)** a scenario where the covariate distributions are moderately confounded and partially overlap, and the orange batch tends to see higher outcomes than the blue batch. **(C)** a scenario where the covariate distributions are moderately confounded and partially overlap, and the orange batch tends to see lower outcomes than the blue batch. **II.** and **III.** indicate the corrected data after correction, via `cComBat` and `Causal cComBat` respectively. If the batch effect is removed, the orange and blue lines should be approximately equal.

In this manuscript, we posit that the most principled causal procedure is often to do *nothing*, and avoid inference entirely if the datasets do not share similar demographics. Informally, these results can be summarized succinctly: when upstream covariate distributions of measured individuals vary widely, and those covariates are known to impact both the measurements and the selection of individuals to particular batches (the covariates are confounding), traditional techniques can yield imprecise conclusions regarding batch effects unless first passed them through a causal lens. Leveraging this perspective, we illustrate how augmentation of traditional approaches via our proposed causal procedures, CAUSAL CDCORR for batch effect detection and `Causal cComBat` for batch effect correction, can overcome many of these limitations. Our simulations and real data analyses reveals two salient points:

1. Traditional strategies to detect or correct batch effects from mega-studies may struggle to parse differences between batch effects and real demographic variability when demographic variables are insufficiently similar.
2. Caution is advisable while interpreting results post batch effect mitigation, because the details of how the mitigation is performed can strongly impact the results.

These issues highlight the primary challenges of performing valid statistical inference while pooling data across studies. This work therefore contributes to the ongoing effort to improve the validity and reliability of inferences in past and future mega-studies.

2 Methods

2.1 The CoRR Studies have disparate demographic characteristics

The motivation for our development of a modified framework for defining, detecting, and mitigating batch effects is the neuroimaging mega-study produced by the Consortium for Reliability and Reproducibility [69], a collection of over $3{,}600$ functional neuroimaging measurements from over $1{,}700$ individuals spanning 27 separate datasets. Figure 2A explores the demographic characteristics for the individuals in the CoRR mega-study. Many of the studies have a narrow age range, and several studies only include females. Because sex [25, 53, 64], age [21, 52, 59], and continent [19, 38] are variables that have been associated with brain connectivity, they serve as demographic covariates used in our investigation.

Figure 2B illustrates the level of demographic overlap in the CoRR mega-study, using the distribution-free overlapping index [44], as discussed in Appendix E. The CoRR mega-study includes many pairs of datasets with varying degrees of overlap, from high (near 1) to low (near 0). Further, many of the datasets do not overlap at all (overlap of 0), making inference about batch effects (e.g., via detection or mitigation) impossible without making strong assumptions. This is particularly troublesome, as the covariate records for the CoRR mega-study common to all sites include only three covariates: age, sex, and continent of measurement. The addition of successive covariates can only reduce the estimated overlap between the pairs of datasets, so having poor overlap on such a coarse set of covariates indicates that the actual demographic overlap may be even lower.

2.2 Detecting and mitigating batch effects

At present, literature approaches for detecting and mitigating batch effects tend to focus attention heavily on effects which could be classified as associational or conditional, which for the above listed reasons, fail to adequately account for confounding biases. To this end, we propose a simple technique to augment associational and conditional strategies and derive stronger causal claims that test or correct for actual batch effects. We propose the following strategy, illustrated graphically in Figure 2(C):

1. Use classical causal procedures to re-weight the measurements from the batches so that the demographic distributions across all batches to be considered overlapping or are approximately equal. For rendering the covariate distributions approximately overlapping for the detection problem, we leverage vector matching [35]. For rendering the covariate distributions approximately equal for correction, we leverage propensity trimming followed by nearest neighbor matching [49, 55]. Intuitively, vector matching and propensity trimming followed by nearest neighbor matching assign weights w_i to each sample i which take a value of 0 (removing the sample from subsequent analysis) or 1 (including the sample for subsequent analysis), and conceptually "rebalance" the observed data such that the covariate distributions are more alike.

2. Apply classical procedures for detection or removal of conditional effects. For the purposes of elucidating the benefits of this perspective, we choose to apply our techniques to cDcorr for batch effect detection [7, 63], and cComBat for batch effect removal [28, 33, 34, 46, 48, 62, 67].

For a more technical description of the causal procedures employed in this manuscript, see Appendix A. It is important to note that we do *not* make the claim that the specific choices for re-weighting measurements (propensity trimming followed by nearest-neighbor matching, or vector matching) are necessarily the best possible approach to balancing the demographic distributions across the batches. Indeed, there are almost two decades of research dedicated towards making these design choices for observational studies. Rather, we chose simple, principled, and straightforward approaches for illustrative purposes. Our simulations contained herein and in our related manuscript [7] elucidate that even these possibly simplistic balancing approaches are sufficient to highlight substantial shortcomings in the existing literature for detecting or correcting for disparities between multiple batches of data in the face of covariate imbalance, as well as the substantial benefits of a causal perspective.

3 Results

A Empirical Distribution of Demographics

B Disparities in Demographic Distributions

C Demographic Balancing Procedure

I Unadjusted Data II Propensity Trim/Match III Adjusted Data IV Downstream Analysis

Causal cDCorr

Causal cComBat

Figure 2: **Demographic data for each of the** $N = 27$ **studies from the CoRR Mega-Study**. **(A)** Each point represents the age of a participant corresponding to a single measurement, rows are studies, boxes are continents, and color indicates sex. **(B)** Even with only 3 covariates made available (sex, age, continent of measurement) the CoRR studies generally show extremely poor covariate overlap [44]. This makes inference regarding batch effects difficult. **(C)** The demographic balancing procedure, to demographically align poorly balanced datasets using CAUSAL cDcorr or Causal cComBat. (C.I) The unadjusted datasets are imbalanced in covariate distributions. The smallest dataset, the *reference*, has a narrower age distribution than the controls. (C.II) Propensity trimming (shaded boxes) and matching (lines) are conducted to identify reasonable matches between the reference and the control datasets. (C.III) The adjusted datasets have far improved demographic overlap after propensity trimming, matching, and removing reference subjects who do not have suitable matches in the control datasets. (C.IV) Downstream batch effect detection via CAUSAL cDcorr or correction using Causal cComBat are performed on the data using conditional strategies on the adjusted data.

3.1 A causal framework for batch effects

To contextualize the importance of modeling batch effects as causal effects, consider the following example. We have two batches of people, where for each person, we measured some outcome (such as disease state). We want to know whether the differences between the batches are due to experimental variability, or veridical differences in outcomes across the two batches. In the ideal case where the same people are measured across the two batches (a *crossover* scenario), subsequent variability in the observed outcomes can be trivially argued to be due to batch effects, but this crossover property is often not met and we usually have different people measured in each batch. The simplest approach to handle this case would be to directly compare the outcomes across batches. This approach is susceptible to false positives and negatives. If the two batches differ because they have sampled individuals in different age groups, and the outcome is a function of age, then this approach might falsely conclude there exists experimental variability when it is not present. On the other hand, if experimental variability is truly present, but somehow another covariate cancels out this variability, then this approach could falsely conclude there does not exist experimental variability when it is present.

To mitigate these issues, one could instead compare the distribution of outcomes across batches, but conditioned on age. Naive conditional approaches can however, also be susceptible to both false positive and negative results. Imagine that the two batches have covariates that partially overlap; for example, one sampled adolescents and adults, but not geriatrics, and the other sampled adults and geriatrics, but not adolescents. Conditional approaches make implicit assumptions about how to interpolate the covariate distributions across batches. When those assumptions do not reflect the real data, the results can be erroneous.

A causal modeling approach can address this issue. In simple terms, the causal modeling approach discards all data across batches that are not "comparable". For example, in the above example, it might discard all the adolescents and geriatrics, and simply focus on modeling the adults, for which we have data in both batches. While this reduces the sample size, it avoids the pitfalls of confidently concluding the presence or absence of a batch, even when it is not clear.

Here, we introduce formal definitions of batch effects, which to our knowledge are not yet otherwise available in the literature. Specifically, we define a batch effect in Definition 1, which in general cannot be directly tested in observed samples unless assumptions are made. We elucidate these assumptions through *associational effects* in Definition 2 which can be practically tested via DCORR [56], *conditional effects* in Definition 4 which can be practically tested via cDCORR [63], *adjusted effects* in Definition 6 which can be practically tested via CAUSAL cDCORR [7], and *crossover effects* in Definition 8 and Definition 9. To mitigate each of these effects, we define associational batch effect correction in Definition 3 which can be applied via ComBat [28], conditional batch effect correction in Definition 5 which can be applied via cComBat [17, 28, 48], and adjusted batch effect correction in Definition 7 which can be applied via the procedures delineated in Section 2 for Causal cComBat.

A   Associational Procedure Directed Acyclic Graph (DAG)

B   Conditional (Non-Causal) Procedure DAG

C   Adjusted (Causal) Procedure DAG

D   Crossover (Causal) Procedure DAG

Figure 3: **Causal Graph of Study Covariates**. Causal Directed Acyclic Graphs (DAG) illustrating the underlying assumptions under which various procedures to detect or correct for batch effects are reasonable. Boxes represent variables of interest, and arrows indicate "cause and effect" type of relationships between variables. The causal estimand is the impact of the exposure (the batch) on the outcome (the participant's measurement), and is a cumulative effect of effect modifiers (black variables) both known and unknown that yield batch-specific differences. Variables are *confounding* if they effect both the exposure and the outcome. **(A)** associational procedures and **(B)** conditional procedures are reasonable when measured or unmeasured covariates are non-confounding. **(C)** adjusted causal procedures are reasonable when confounding variables are measured. **(D)** crossover procedures are reasonable under many forms of potential confounding, both measured or unmeasured, so long as participant states are non-confounding. If participant states are confounding, the confounding states must be measured (red arrow).

3.2   Batch effects in biological datasets

The effects discussed above have immediate practical implications on measuring batch effects in modern biological datasets. To this end, we demonstrate the applicability of these effect types in the context of a the CoRR mega-study, in which measurements are collected from individuals across various batches (the *exposures*). The logical process, however, extends beyond this context to many other biological data modalities, such as genomics. Directionality

of arrows (e.g., $T \to Y$) indicates that the variable T can influence the variable Y. For instance, aging may cause changes to the brain, which are reflected in the measurements. The exposure is the batch in which participants' data are collected, and the outcomes are the measurements themselves. We seek to estimate the *causal estimand*, which is the causal effect of the exposure to a particular batch on the outcome (here, the connectome). The causal estimand is a cumulative effect of numerous factors (both known and unknown) relating to the measurement procedure for each batch, and are typically standardized across measurements in a single study, as illustrated in Figure 3**(A)**. For instance, one study conducted at one facility may have access to a particular scanner, and another may have access to a different scanner, and the scanner model may fundamentally change the observed measurements (e.g., the batch effects the outcome via the scanner disparity). It is important to convey that in many biological datasets (such as the CoRR study [69]), these factors are *completely determined* by the batch, and are not truly "random" given the batch label. Therefore, we cannot more precisely describe a "batch effect" (for instance, delineating the portion of a batch effect due to a particular scanner setting), without additional assumptions. Numerous covariates, both known and unknown, impart potentially confounding biases, in which the variables can influence both the exposure and the outcome (e.g., $T \to Y$, but $X \to T$ and $X \to Y$, in which case X is a confounder). For instance, one batch collected in a children's hospital may tend to over-sample younger individuals relative to other batches, and younger children may have distinct patterns of brain connectivity compared to older children (age is confounding). For this investigation, we use age and sex (which are commonly used for adjusting for batch effects) as well as continent, as measured covariates in subsequent analyses. We depart from many previous batch effect correction analyses by leveraging continent due to the growing literature [19, 43] which suggest differences in brain function for individuals across cultures. Unaccounted confounding via culture/race (for which continent serves an imperfect surrogate) may yield confounding biases in subsequent analysis, in which batch effects may be conflated with veridical demographic differences.

Unmeasured covariates are often prominent confounders in biological datasets, making proper causal analyses troublesome. *Participant history* involves variables related to specific life experiences, both environmental and societal, that the individual has been exposed to. *Participant traits* are characteristics of a person that remain relatively constant over time. *Participant states* refer to characteristics of a person that change over an individual's lifetime. *Exposure determinants* are factors which impact batch membership for a particular individual, but do not impact the measurement itself. *Outcome determinants* are factors which do not impact the batch assignment, but impact the outcome. In this particular study, people are not explicitly recruited on the basis of traits or histories which may be affected by their measurements (e.g., neurodiversities). This yields that the causal graph is also acyclic, or a DAG.

Figure 3**(A)-(D)** indicates graphical illustrations of the assumptions which are sufficient to render associational, conditional (non-causal), adjusted (causal), or crossover procedures as-defined in Section B equivalent to batch effects. An immediate implication is that the associational, conditional, adjusted, and/or cross-over estimation procedures are valid under those assumptions, and not necessarily valid otherwise. For example, if the batch and the outcome are dependent upon any covariates, which they often are in real mega-studies, then an estimated associational effect is not a valid estimate of a batch effect (Figure 3**(A)**). If there are any confounding covariates, then an estimated conditional (non-causal) effect is not a valid estimate of a batch effect (Figure 3**(B)**). If there are unobserved and confounded covariates, then adjusted (causal) procedures do not produce a valid estimate of a batch effect (Figure 3**(C)**).

Crossover effects can account for many types of measured and unmeasured covariates which are potentially confounding (Figure 3**(D)**). Because of the crossover property, the batches will be balanced across certain measured *and* unmeasured covariates, specifically patient histories and traits. Hence, many of the unmeasured variables from Figure 3**(D)** are deconfounded by design of the data collection procedure (the dashed black arrows do not exist in a cross-over setting). However, unmeasured changes in participant state between different exposure groups (such as measuring individuals only

during the daytime in one batch and only during the night time in another batch) remain potential confounders if not carefully balanced across exposure groups through either randomization or direct experimental intervention. If these states are confounding, the participant states must be measured, and controlled via adjusted (causal) procedures.

3.3 Traditional approaches fail to adequately remove batch effects when covariate overlap is imperfect

We constructed simulations to help us understand the differences between conditional (non-causal) reasoning and adjusted (causal) reasoning. To this end, we evaluate the performance of our proposed adjusted (causal) technique, `Causal cComBat`, in comparison with two conditional (non-causal) approaches one may use for batch effect correction, including `cComBat` [28] and `cComBatGAM` [48], the latter of which leverages generalized additive models (GAMs) to capture non-linear trends in covariate/outcome relationships. For the purposes of this experiment, we assume a linear predictor for age for both `Causal cComBat` and `cComBat`, and a non-linear predictor for `cComBatGAM` using GAMs. For this dataset (featuring two dimensions) we could theoretically manually construct what visually appears to be a reasonable predictor for the age/outcome relationship. In real biological datasets, however, the data is far larger in scale (connectomics data, for instance, typically features tens of thousands of dimensions). This makes it infeasible for an investigator to manually select suitable relationships across all dimensions, especially when the data features numerous possibly interacting covariates, resulting in linear techniques or GAMs being leveraged for most biological applications.

Figure 4**A** illustrates the procedure for our simulations. In Figure 4**(A)I** and Figure 4**(A)II**, we illustrate the covariate distributions and expected outcome conditional on the covariate value for each batch. The "covariate overlap" in Figure 4**(A)I** describes the cumulative density shared the two conditional covariate distributions. Conceptually, this denotes the level of "covariate similarity" between the covariates of the two batches (illustrated is a covariate overlap of about 0.15). When the "covariate overlap" is closer to 1, the batches have similar covariate distributions; and when closer to 0, the batches have more dissimilar covariate distributions. Figure 4**(A)II** illustrates the expected outcomes for the non-linear (sigmoidal) regime. The conditional average treatment effect (CATE) describes the difference between the expected outcomes blue and orange batches for each covariate level (red box; here, -0.5). That the expected outcomes for each batch differ indicate the presence of a batch effect. The average treatment effect (ATE) is the expected difference (over all covariates) and is therefore also -0.5. The expected outcomes for each batch serve as the *ground truth* for our investigation. In Figure 4**(A)III**, $n = 500$ samples are generated with equal probability from either of the two batches, and then covariate values are sampled given the batch. Finally, the observed outcome is generated as a random offset from the expected outcome given the covariate and batch of the particular sample.

The observed outcomes are leveraged in Figure 4**(A)IV** to learn a batch effect correction using one of the three techniques of interest (`cComBat`, `cComBatGAM`, or `Causal cComBat`). In Figure 4**(A)V**, we apply the learned batch effect correction to the ground truths for each batch. If the batch effect is "removed" successfully, the ground truths (after correction) should be identical. We measure how similar/dissimilar the ground truths after correction are using the ATE(after), which will be non-zero if the ground truths after correction differ (the batch effect was not removed). In the illustration, the original batch effect (the -0.5 magnitude offset between the blue and orange batches) has been overcorrected, and now has a magnitude of 0.5. Figure 4**B** applies the above procedure over data with batch effects before correction of varying magnitude (a positive ATE(before) in Figure 4**(B)1**, a null simulation where no batch effect is present in Figure 4**(B)2**, and a negative ATE(before) in Figure 4**(B)3**), across multiple possible covariate/outcome relationships (linear in Figure 4**(B)I**, non-linear/sigmoidal in Figure 4**(B)II**, and non-monotone/bump in Figure 4**(B)III**). We repeat the procedure described in Figure 4**A** over $R = 1,000$ trials, and compute the mean ATE(after) over all trials. As-described above, ideally, all procedures for batch effect correction will leave the data with no batch effect (ATE(after) = 0.0, dotted black line). In general, `Causal cComBat` fails to run more than 50% of the time when the level

of covariate overlap is below 0.1 (dotted red line). When covariate overlap is high, all strategies have reasonably strong performance. As covariate overlap drops, however, naive conditional (non-causal) strategies perform arbitrarily poorly across all but the linear regime; when the level of covariate overlap is 0.1, for instance, the ATE(after) has a magnitude of about 1 for both `cComBat` and `cComBatGAM`, which indicates that the ground truths are a factor of about two ($1/0.5 = 2$) more dissimilar after correction than before. On the other hand, `Causal cComBat` maintains high performance even at modest levels of covariate overlap across all regimes. As covariate balance declines, `Causal cComBat` tends to fail to execute at an increasing rate due to no points being retained across both batches after matching. On the other hand, unless the relationship between the covariates and the outcome is perfectly linear, `cComBat` and `cComBatGAM` perform arbitrarily poorly in low-overlap regimes. The "corrected" data is as much as a factor of about five times more different between the batches than before any "correction" was applied (ATE(after) reaches a maximal value around 2.5, but the original ATE(before) in Figure 4**(B)1** and Figure 4**(B)3** had magnitudes of 0.5). Further, in regimes where no batch effect was present in Figure 4**(B)2**, application of `cComBat` and `cComBatGAM` have introduced batch effects (ATE(after) is non-zero). Together, these results illustrate that if batch effect correction is performed without causal considerations and the batches do not feature near perfect covariate alignment or omniscient knowledge of the covariate/outcome relationship, which cannot be practically achieved in most contexts, subsequent batch effect correction often does not actually "correct" the data, and can even worsen or introduce new disparities between the batches.

Conceptually, researchers may feel as though they are left with a tradeoff:

1. When datasets do not have high overlap demographically, the (implied or explicit) limitations of a model leveraged by a batch effect detection or correction technique can impart substantial bias on the conclusions of the investigation. While the limitations or shortcomings may *seem* inconsequential, the scale of the residual signal disparities imparted can dwarf the original batch effect, and yield measurements which are *more dissimilar* than prior to batch effect adjustment, as we saw in Figure 4(E).II. and (E).III.
2. Enforcing demographic overlap mechanistically through propensity trimming and/or matching may *seem* like it yields fewer samples for analysis, or narrows the scope of conclusions.

In practice, the reality is that with biological data, there are *no* ground truths. As a result, there is no way to evaluate the introduction of imparted residual bias, and no way to know whether downstream conclusions are a product of imparted residual biases. This means that any statistical inference we derive may be arbitrarily true or false, and we have no way of qualifying in which pool our inference stands.

When we mechanistically re-weight the data to impose demographic overlap, proper statistical inference tempers conclusions as a direct consequence of the smaller demographic region under consideration. Stated another way, this means that when we balance *prior to* detecting or correcting for batch effects, we have tools which allow us to make internally valid conclusions supported by the data in regions in which the data *are able*. This means that, although our conclusions may not apply to the *entire* region of covariate support in the observed data, they will be *statistically and methodologically valid* with respect to the re-weighted population (albeit, in a possibly narrower set of covariate values) that we *did* analyze.

A Procedure for assessing batch effect correction methods

B Performance of batch effect correction techniques across simulated data

Figure 4: **Non-causal procedures are subject to strong biases without covariate matching**. **A** the procedure for assessing batch effect correction techniques. **(A)I** using the covariate distribution and **(A)II** the data-generating model for each covariate level, **(A)III** $n = 500$ samples are generated from each batch with equal probability. **(A)II** the batch-specific data-generating models have a conditional (on the covariates) average treatment effect of -0.5 across all covariate levels. Therefore, the average treatment effect (ATE) before correction is -0.5. **(A)IV** the observed data is used to learn how to "correct" the batch effect between the batches. **(A)V** using the data-generating model (the ground truth), we apply the batch effect "correction" to determine the effectiveness. In the ideal case, the ATE after correction will be 0 (the learned batch effect "correction" removed the batch effect), but this is not necessarily the case. In the illustrative example, the average difference between the blue and orange lines (the ATE(after)) is greater than 0. **B** evaluating the effectiveness of different batch effect correction techniques for (Row **1.** and **3.**) removing batch effects when present, or (Row **2.**) avoiding the introduction of batch effects when none are present *a priori*. (Column **I**) illustrates the performance when the data-generating model is a linear relationship, (Column **II**) for a non-linear (sigmoidal) relationship, and (Column **III**) for a non-monotone (bump-shaped) relationship, over $R = 1,000$ trials. Techniques with high performance will have a mean ATE after correction at or near 0. Across all regimes, `Causal cComBat` struggles to execute more than 50% of the time when covariate overlap is insufficient (approximately 0.1). Across linear regimes, all techniques are effective for batch effect removal, regardless of covariate overlap. In both non-linear and non-monotone regimes, both `cComBat` and `cComBatGAM` fail to remove batch effects when present (Row **1** and **3**) and introduce batch effects when none are present *a priori* (Row **2**). Note, in particular for Row **1** and Row **3** that the magnitude of the ATE after correction exceeds the magnitude of the ATE before any correction was applied at all (0.5), indicating that the data for each batch are *more dissimilar* after correction than before correction when using `cComBat` or `cComBatGAM`.

Figure 5: **Comparison of types of effects between datasets from the CoRR study**. **(A)** Heatmap of different types of effects (conditional and adjusted causal procedures) that can be used to detect differences between each pair of datasets in the CoRR study. Whereas most (25.8%) non-confounded conditional effects are not significant, most (66.1%) non-confounded adjusted effects are significant. **(B).I** delineation of how one possible source of batch effects, scanner model, impacts significance rates of batch effects. Almost all pairs of studies conducted on different scanners with high covariate overlap ($> .05$) have discernable batch effects. The rate of detected effects is lower when the scanner model is the same. **(B).II** and **(B).III** When the level of estimated covariate overlap is lower ($< .05$) or zero (different continent), conditional effects never detect a difference across datasets. On the other hand, adjusted causal procedures instead report that the data is too confounded for subsequent inference, and avoid running entirely.

3.4 Detecting Batch Effects in the CoRR mega-study

Figure 5**A** focuses on discerning the viability of different types of effects one could use to test for batch effects from Section 2. For each pair of datasets in the CoRR study, we perform a hypothesis test for whether (or not) a discernable effect is present, while controlling for age, sex, and continent differences between the individuals within each dataset. Intuitively, we would like to believe that if such a test rejects the null hypothesis in favor of the alternative, the data supports that a batch effect is present. When we account for demographic covariates using conditional (non-causal) approaches (orange squares, top left), discrepancies between the datasets are detected 25.8% of the time (checkboxes, conditional distance correlation two-sample test, BH correction, $\alpha = 0.05$). As explained in Section 3.2**B**, the validity of these approaches as tests for batch effects is the assumption that the measured covariates are overlapping and non-confounding, which is overwhelmingly false for many of these comparisons, as shown in Figure 2**B**.

We perform similar tests using adjusted (causal) procedures in Figure 5**A** (blue squares, bottom right). Many pairs of studies could not have an adjusted effect estimated due to poor demographic alignment (282 of 406 pairs of studies). Colloquially, when covariate overlap is moderate or limited, we cannot tell whether discernable differences are due to batch effects or biological heterogeneities between the batches (e.g., due to age, biological sex, or culture/continent) without heavy assumption-based extrapolation as those made by conditional procedures, and therefore a conservative strategy would be to avoid inference entirely. In this way, causal strategies pre-pended to batch effect detection

and mitigation strategies are *more conservative in their application*, in that they prevent the experimenter from potentially deriving spurious or erroneous conclusions (either detections or non-detections) due to confounding biases. Notably, adjusted (causal) procedures can be used to estimate effects between all pairs of the American Clique (a group of separate studies consisting of similar demographics collected in North America, consisting of NYU1, NYU2, IBATRT, UWM, MRN1). After adjustment for the sample characteristics of the demographics associated with individual studies, a majority of adjusted (causal) effects (66.1%) remain significant (causal conditional distance correlation two-sample test, BH correction, $\alpha = 0.05$).

Note that the rate of effect detection in Figure 5A for conditional effects extremely closely resembles the levels of covariate overlap, as the checkboxes tend to correspond to the pairs of datasets which feature high covariate overlap in Figure 2**B**. We attempt to investigate the possible causes of a failure to reject for conditional effects in Figure 5**B**. A possible source of a batch effect could be different types of scanners were used to collect the data [8, 28, 48], since the scanner model could be a causal effect modifier as in Figure 3**A**. Therefore, a pair of studies differing in scanner may yield the potential for a true batch effect to be present. To this end, we aggregate the fraction of comparisons which report the indicated test outcome for different levels of covariate overlap (rows) when the scanner model is the same (right column) or different (left column). Figure 5**(B)I** shows that when the scanners are different and the covariates overlap, both causal and non-causal methods reliably detect batch effects a majority of the time, and they do so at a higher rate than when the scanner is the same.

When the covariate overlap is low, or when the continent of measurement is different entirely (and the overlap is zero), conditional (non-causal) procedures almost always fail to reject. On the other hand, adjusted (causal) procedures overwhelmingly report that a given pair of studies are demographically confounded, and are too different for any effect to be investigated all together. Taken together, this illustrates that, when demographic overlap is high and the scanner differs, batch effects are nearly always detectable using conditional (non-causal) or adjusted (causal) procedures. On the other hand, when demographic overlap is lower, even if the scanner still differs, effects are entirely undetectable by conditional (non-causal) procedures. In a situation where we expect batch effects (different scanners), we go from almost always detecting effects to never detecting effects simply by looking at pairs of comparisons with less demographic overlap. On the other hand, adjusted (causal) procedures do not suffer from this limitation, and explicitly delineate when the datasets are overly confounded by demographic covariates.

Appendix Figure 13 delineates similar plots for the other two effect types (associational and causal crossover effects).

3.5 Traditional Approaches Produce Disparate Inference from Techniques which Leverage Causality

At this point, we have demonstrated, through simulation and real examples, if a covariate is a *confounder*, failure to properly account for demographic balancing can, potentially, yield improper detection and mitigation of batch effects and may, in turn, fundamentally alter true signals in the dataset or detect spurious demographic noise. We saw that, for many of the CoRR [69] datasets, demographic balance was extremely poorly aligned, which allows only a fraction of the original possible comparisons between datasets to be analyzed in a theoretically, and empirically, reasonable manner. The final question we have is: does it matter? Even if we disrupt or misrepresent veridical signals in the dataset, can we still detect them, irrespective of whether the signal is maintained in its unadulterated form?

To this end, we investigate the strength of the sex effect, conditional on age, in the connectomes before and after batch effect correction (Figure 6A). `ComBat` and `cComBat` are performed on the entire CoRR study (top row) with age and sex covariates, and subsequently all techniques are restricted to the subset of connectomes upon which `Causal cComBat` was executed (the *American Clique*, bottom row). This amounts to the individuals within the study who were retained after matching on the basis of age, sex, and continent within the American Clique. We also compared these techniques to the

approach developed by the NeuroHarmonize package (Conditional and Causal NeuroH) [17, 48], an implementation of `cComBatGAM` for neuroimaging data. The edges which show a significant, sex effect conditional on age (partial distance correlation, BH correction, $\alpha = .05$) are colored from smallest test statistic to largest test statistic (rank = 1, dark purple), and edges without a significant conditional sex effect are colored white. Causal approaches produce vastly different statistical inference compared to other correction techniques on the American Clique. Causal approaches were unable to be executed on the set of all studies, because many of the studies in the CoRR dataset do not overlap in terms of their covariate distributions (the red square indicates the analysis is not possible on the basis of continent, age, and sex). Using non-causal strategies for batch effect removal prior to restricting analysis to the matched produces substantially different inference than restricting analysis to the matches samples, as inference would suggest no sex effects versus many sex effects (bottom row).

Next, we seek to answer whether statistical inference for the entire CoRR study (top row) is similar to inference that would be obtained on the matched American Clique using only `Causal cComBat` strategies. This investigation stems from the observation that the heatmaps appear, at face value, to be approximately similar in terms of the number of significant edges, and the visual patterns observed, between the four left-most plots of the top row and the two right-most plots of the bottom row. Perhaps it is the case that while causal approaches provide more informative inference on a smaller matched sample, the inference is still the same if we just ignored matching to achieve demographic balance all together. We compare the DICE overlap of the top n edges (by effect size) from each of the four non-causal approaches applied to all the data with the top n edges of `Causal cComBat` on the matched American Clique in Figure 6(B) A high DICE overlap indicates that the subsets of edges are similar. None of the techniques show high overlap with the edges that are indicated to have a large effect after correction with `Causal cComBat`. Finally, using the raw connectomes, `ComBat`, Conditional NeuroH, and `cComBat` tend to produce similar inferences, as all have high DICE overlaps near one. In contrast, they all have low overlap with `Causal cComBat` (Figure 6C), with DICE overlap with `Causal cComBat` nearly zero. On the other hand, Causal NeuroH has a high overlap with `Causal cComBat`. This suggests that causal and conditional strategies for batch effect correction produce somewhat orthogonal subsequent inference.

4 Discussion Our primary conceptual contribution is establishing that batch effects can be formalized as causal effects. Given this realization, we propose a framework for quantifying and removing batch effects in high-dimensional datasets featuring non-Euclidean measurements. We propose a straightforward approach to augment existing strategies for proper batch effect correction and removal, by first prepending existing techniques using causal balancing procedures. We explore the advantages of these procedures by demonstrating that causal approaches (such as `Causal cComBat` and CAUSAL cDCORR) yield empirically valid approaches to handling batch effects, or report that no inference can be made with insufficient covariate overlap. This represents a principled approach over alternative strategies, which we show may introduce spurious biases and dramatically under-estimate or over-estimate batch effects. We illustrated how, ignorant of potential confounders, otherwise principled strategies may dramatically under-estimate the prevalence of batch effects. This illustrates that causal adjustment procedures can augment hypothesis testing by indicating when the data may be unsuitable to provide answers to a particular question due to irrevocable confounding, rather than permitting false positives or false negatives. Finally, our neuroimaging use-case shows the utility of the methods developed herein for estimating and removing batch-related effects from large biological datasets. Specifically, our methods identify different demographic effects as compared to prior approaches, each of which identifies features similar to a completely unadjusted approach, which are well established in the causal literature to permit numerous false positives and false negatives [13, 16]. Together, our work rigorously defines and studies measurement bias decoupled from demographic confounders in complex, non-Euclidean data.

Figure 6: **Significant Edges Before and After Batch Effect Removal**. **(A)** The presence of a sex effect (conditional on individual age) is investigated for each edge in the connectome. The top 100 significant edges are shown in rank-order from largest (rank = 1) to smallest sex effect ($\alpha = .05$, Benjamini Hochberg [5] correction). Analysis is either performed over *all* individuals (top row) or only upon the individuals for which sufficient demographic overlap could be obtained, the *American clique* (bottom row). `Causal cComBat` and Causal NeuroH can only be executed on the American Clique (red square denotes Causal approaches are not possible with *all* studies). **(B)** The DICE overlap of the top n edges of each batch effect removal technique on all the data (green squares, top row) as compared to those from `Causal cComBat` (green square, bottom row). The red line (random) indicates the overlap to be obtained for a particular choice of n to be expected as a result of random chance, and the gray ribbon indicates the 99% confidence interval of the estimate (1000 repetitions). The DICE overlap increases as a function of n, regardless of whether any signal is actually present. **(C)** The DICE overlap of the top 100 edges for all pairs of correction techniques indicated with a green square from **(A)**. Overlap of the top signal edges is high between no correction (raw), `ComBat`, `cComBat`, and Conditional NeuroH, but all techniques have low overlap with `Causal cComBat` and Causal NeuroH, indicating that Causal approaches yield measurements with signal that the associational approaches fail to identify.

The conclusions of our work bring into focus the pervasive question of how to properly interpret neuroimaging—or more generally, multi-site scientific—studies.This concern is at the root of any statistical inference task: a conflict between the internal and external validity. Internal validity refers to how well the conclusions generalize from the *sample* population to the *study* population; that is, from the observed samples to other samples that are distributed the same as the observed samples. External validity refers to the degree to which the conclusions generalize to the *target* population; that is, the population to which we wish to apply the conclusions of our inferences. The target population is typically much broader than the study population, to include, for example, all members of the species, or possibly even multiple species.

Through simulation, we demonstrated that minor misspecifications of the statistical model employed for batch effect removal can be disastrous in the presence of confounding. Specifically, a minor change of the relationship between the confounder (the covariate) and the measurement from a straight line to a (still monotonic) nonlinearity, the sigmoid, not only inhibited `cComBat` from removing a batch ef-

fect, but also introduced an artifact that dwarfed the magnitude of the original batch effect itself (an *overcorrection*). This observation extends directly to other forms of model misspecification, such as non-monotonicities (which resulted in an *undercorrection*). As discussed in the Appendix D.2 and expanded on via further simulations, the resulting estimate of the batch effect can, by construction, be considered neither statistically unbiased nor consistent, thus failing two preconditions for internal validity [65]. Colloquially, it is almost impossible to know whether models employed for batch effect correction will be reasonable to apply to out-of-sample data (e.g., areas of the covariate distribution for which there is no covariate overlap). That is, we cannot tell whether the data we have will fall into Appendix Figure 10 or Appendix Figure 7 and the batch effect correction technique will work reasonably given the underlying true signal, or Figure 4, Appendix Figure 8, or Appendix Figure 9 where the assumptions of the batch effect correction technique will be entirely unreasonable. On the other hand, causal strategies can help us to apply far more conservative logic, indicating to us when batch effects are unreasonable to estimate as is often the case with limited overlap in our simulations. Alternatively, they can aid us to first balance the covariate distributions so that we can avoid the pitfalls of the simulations which illustrate poor performance by `cComBat` in moderate or limited overlap regimes. Stated another way, in the absence of knowledge as to the appropriateness of the model employed for batch effect corrections, causal strategies make fewer assumptions about the covariate/outcome relationship to produce reasonable estimates.

Given the major shortcomings theoretically and empirically of traditional approaches for detecting and mitigating batch effects, we therefore chose to first prioritize the development of strategies which sub-sample individuals into demographically-aligned subsets using causal techniques. In so doing, we prioritized internal over external validity, as internal validity is a prerequisite for external validity. Colloquially, one cannot generalize results of an experiment to a target population if the results are, at best, dubiously applicable to the study population. The trade-off of subsampling individuals is that sample size, sample diversity, and external validity are exchanged for internal validity. We saw this tradeoff materialize dramatically in the CoRR dataset, where over half of all possible dataset pairs could not be assessed for batch effects due to a lack of demographic alignment. This led to a much higher failure to reject rate in the conditional strategy than the adjusted causal strategy for batch effect detection, even in situations in which it would tend to reject (high overlap, different scanner model) with sufficient covariate overlap. This illustrates dramatically the limitation of a "reject" or "failure to reject" mindset; prevalence rates for batch effects may directly be a function of simply not having sufficient data to support whether (or not) a batch effect is present. Causal techniques tend to provide further granularity to these two outcomes of a hypothesis test. Particularly, an outcome of "confounded" indicates that, given even infinite samples from the respective study populations, in some situations we may still be unable to make any determinations about the presence of batch effects due to the fact that the study populations are simply orthogonal.

The final hurdle of our analysis, therefore, is to address the question: does the internal validity of batch effect removal techniques actually impact downstream statistical inference? A failure to remove batch effects in an internally valid manner, we believe, presents a substantial hurdle for producing internally valid statistical inference downstream, since downstream statistical inference is a product of the corrected measurements. Specifically, one can endeavor to mitigate batch effects in one of two complementary ways. First, one can filter individuals by ensuring demographic balancing, and then run the entire analysis only on those individuals (as causal approaches, like `Causal cComBat`, do). Second, one can run an algorithm for batch effect correction (and potentially benefit from a richer number of samples), and then filter to ensure demographic balancing only when measuring downstream demographic-specific effect sizes (as we did by looking for a sex effect only considering individuals from the Raw, `ComBat`, or `cComBat` corrected connectomes in the American Clique). If internally valid, causal batch effect correction techniques may substantially restrict the sample size. Perhaps, even if the batch effect correction technique is not internally valid, measurements corrected using it may still have

utility for downstream analysis. We anticipated that the most significant features for the inference task would be, at least, ordered similarly between the two analyses. However, our finding that the analyses produce radically different downstream inferences, even when we restrict the non-causal techniques to the same datasets, suggests that there is a link between downstream conclusions and the internal validity of batch effect correction techniques. Stated another way, when we perform batch correction using as few assumptions as possible about the study population through causal techniques, we get extremely disparate downstream scientific conclusions from state-of-the-art conditional techniques. This suggests to us that further work is needed to better understand how the removal of batch effects can be achieved while mitigating the deleterious effects on either internal or external validity through careful consideration of the study populations of analyses.

Limitations and Future Work The most prominent limitation of our work is that it appears as though we argue in favor of using strategies such as matching to pre-align measurements in terms of the covariate distributions. Indeed, this assumption is extremely restrictive: data will, inevitably, be discarded in the pursuit of internally reasonable estimates. Other techniques we could have leveraged, such as inverse propensity reweighting [55], subvert this limitation, but introduce the related issue that we are still estimating batch effects by upweighting or downweighting particular samples from the data. On the contrary, we do not advocate that the techniques proposed are the *best* approaches one could use for batch effect correction. We posit that the techniques (e.g., prepending matching) or variations thereof (e.g., via inverse propensity reweighting) represent the only context in which naive `ComBat`, `cComBat`, or related strategies leveraging potentially misspecified linear models could be applicable without rigorous justification of linear modeling choices in high-dimensional datasets with non-overlapping covariates as are often encountered in multi-site or multi-experiment mega-studies.

This limitation can be subverted by the development of more robust approaches for estimating the covariate/outcome relationship, building off the `cComBatGAM` approach [48] of incorporating non-linear modeling approaches (such as Generalized Additive Models, or GAMs) into the estimation process for the relationships between the covariate and the outcome instead of only between the covariates. With robust techniques to non-linearities for estimating the covariate/outcome relationship, the restrictive assumption of alignment of the propensity distributions can be exchanged with overlap of the propensity distributions, which informally means that the datasets must have a similar "range" of covariates (though, their distributions may differ, which is more realistic for observational data) [35]. Informally, you can "guess" the covariate/outcome relationship incorrectly and obtain reasonable estimates of batch effects as long as the covariate distributions are similar, or you can "guess" (or estimate) it correctly and need only covariates to be sufficiently represented in each batch. This is illustrated directly in our manuscript specifically for the testing problem [7]. Unfortunately, directly estimating non-linear effects in high-dimensionality, low sample-size data (such as neuroimaging connectomes) has proven difficult, and most techniques approach this problem with an assumption of sparsity in the data [24, 37], whereas the connectomes of this application (and numerous others, such as in genomics) are often dense. Strategies which can reasonably (explicitly or implicitly) estimate the covariate/outcome relationship provide an exciting avenue for future work.

Biomedical data science, including neuroimaging and connectomics, are plagued by a lack of generalizability and replicability [6, 57]. Many investigations posit that this may be a consequence of small sample sizes or overly simplistic datasets [40, 57] lacking sufficient subject-specific measurements. We argue here that a lack of understanding regarding batch effects, and more fundamentally *causality*, further exacerbates these issues. Indeed, these two perspectives may be considered complementary, in that with insufficient subject-specific measurements both causal and non-causal techniques will run into unmeasured confounding biases. A limitation of our proposed methods are that their adequacy to overcome confounding pre-supposes the measurement of a rich set of subject-specific covariates, which may often not be the case (and, likely, is not in the CoRR mega study). In this way, our real

data investigation serves as illustrative of the wide disparities that can be found when leveraging causal techniques. On the other hand, with sufficient subject-specific covariates (e.g., more than age, biological sex, and continent as a loose proxy for race and culture) we are still likely to run into confounding issues which can be ameliorated by causal techniques such as those proposed herein.

To subvert this limitation, this work suggests that harmonized analyses should be conducted with both harmonized measurement parameters *and* demographic distributions. This represents a departure from the manner in which many mega-studies, including so-called harmonized effects, are currently generated [11, 12, 66, 69], in which potentially demographically-disparate (or diagnostically) populations are analyzed and only loosely harmonized on the basis of the measurement protocols. We saw that post-hoc batch effect detection and removal (and consequently, downstream statistical inference) presents both theoretical and conceptual inconsistencies with regard to internal validity when not first passed through the lens of causality. This problem is highlighted by looking at the poor demographic overlap for popular neuroimaging studies such as CoRR in Figure 2, and in the ABIDE study [11, 12] detailed in Appendix Figure 12. While the SRBPS [66] has greater overlap on the basis of age, sex, and likely race (same continent) for many pairs of datasets (but still has two datasets that are dissimilar from the others), both ABIDE and SRBPS add the additional hurdle of neurodivergences being used in participant recruitment to batches. If the connectome is a proxy for brain function, neurodivergent brain features captured by the connectome may cause symptoms of neurodivergence [22, 31, 60], introducing a potential causal mediation effect if the neurodivergence is leveraged directly in participant recruitment [1, 47]. Intuitively, this can be conceptualized as participant illness status creating a *cycle* in a system-level directed graph (such as Figure 3), which yields a violation of the ignorability assumption that is critical for conditional or causal augmentations to yield stable estimates of batch effects [7]. While this problem has been defined for the wider problem of data fusion [2], is unclear the extent to which causal mediation impacts batch effect detection or correction, nor the suitability of alternative strategies, in high-dimensional biological studies.

Future work will focus on applying the methods developed here to correct for batch effects in mega-studies such as the Adolescent Brain Cognitive Development (ABCD) [29], which includes $N > 12,000$ demographically diverse individuals from across the United States, and measurements were produced with a consistent, harmonized measurement protocol. Further, future studies may seek to build upon the NKI-RS and the recent work by Noble et al. [41] by collecting measurements from the same expansive and diverse group of people across multiple neuroimaging sites or scanning protocols. Our work provides a theoretical foundation for evaluating existing neuroimaging studies, such as Noble et al. [41] and [66], which advocate for the aggregation of studies featuring individuals measured at a wide array of sites (so called *travelling subject* datasets), so that the nature of batch effects can be explored with minimal assumptions needed due to the crossover property. In such datasets, scientists can appreciate internally valid demographic-specific effects on the measurement, and learn more reasonable modeling assumptions, so that perhaps extrapolatory pre- or post-hoc batch effect removal techniques (such as in Appendix Figure 10) can be employed to mitigate the batch effects in other non-demographic-aligned datasets. Failure to attain a comprehensive understanding of batch effects and the extent to which they pervade statistical inferences, we believe, calls into question the validity of existing results in scientific meta-analyses.

Code Availability The code used for analysis within this manuscript is available at github.com/ebridge2/batch_effects, and a docker container with all software versions is available at hub.docker.com/r/neurodata/batch_effects.

Data Availability The raw data analyzed in this manuscript can be obtained at CoRR Mega-Study. The pre-processed data analyzed in this manuscript is available at neurodata.io/mri.

Contributions EWB wrote the manuscript; EWB and JTV revised the manuscript; EWB, JTV, and MP conducted study conception and design; EWB, JTV, BC, and MP conducted interpretation of results; EWB, JC, SP, and RL analyzed raw neuroimaging and demographic data; EWB and JTV devised the statistical methods; EWB performed statistical analyses; MP, GK, SN, TX, MM, BC, and JTV provided valuable feedback and discussion throughout the work.

Acknowledgements The authors are grateful for the support from the National Science Foundation (NSF) administered through NSF Career Award NSF 17-537, the National Institute of Health (NIH) through National Institute of Mental Health (NIMH) Research Projects 1R01MH120482-01 and RF1MH128696, and the NIH through Research Project RO1AG066184-01. We also wanted to thank Betsy Ogburn for numerous insightful conversations regarding the presentation and framing of this work.

References

1. Elias Bareinboim and Judea Pearl. Controlling Selection Bias in Causal Inference. In Artificial Intelligence and Statistics, pages 100–108. PMLR, March 2012. URL https://proceedings.mlr.press/v22/bareinboim12.html.

2. Elias Bareinboim and Judea Pearl. Causal inference and the data-fusion problem. Proceedings of the National Academy of Sciences, 113(27):7345–7352, July 2016. doi: 10.1073/pnas.1510507113.

3. Johanna M. M. Bayer, Paul M. Thompson, Christopher R. K. Ching, Mengting Liu, Andrew Chen, Alana C. Panzenhagen, Neda Jahanshad, Andre Marquand, Lianne Schmaal, and Philipp G. Sämann. Site effects how-to and when: An overview of retrospective techniques to accommodate site effects in multi-site neuroimaging analyses. Frontiers in Neurology, 13:923988, October 2022. ISSN 1664-2295. doi: 10.3389/fneur.2022.923988.

4. Yashar Behzadi, Khaled Restom, Joy Liau, and Thomas T. Liu. A Component Based Noise Correction Method (CompCor) for BOLD and Perfusion Based fMRI. Neuroimage, 37(1):90, Aug 2007. doi: 10.1016/j.neuroimage.2007.04.042.

5. Yoav Benjamini and Yosef Hochberg. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society, 1995.

6. Eric W. Bridgeford, Shangsi Wang, Zhi Yang, Zeyi Wang, Ting Xu, Cameron Craddock, Jayanta Dey, Gregory Kiar, William Gray-Roncal, Carlo Colantuoni, Christopher Douville, Stephanie Noble, Carey E. Priebe, Brian Caffo, Michael Milham, Xi-Nian Zuo, , and Joshua T. Vogelstein. Eliminating accidental deviations to minimize generalization error and maximize reliability: applications in connectomics and genomics. bioRxiv, page 802629, Dec 2020. URL https://doi.org/10.1101/802629.

7. Eric W. Bridgeford, Jaewon Chung, Brian Gilbert, Sambit Panda, Adam Li, Cencheng Shen, Alexandra Badea, Brian Caffo, and Joshua T. Vogelstein. Learning sources of variability from high-dimensional observational studies. ArXiv e-prints, July 2023. doi: 10.48550/arXiv.2307.13868.

8. Andrew A. Chen, Joanne C. Beer, Nicholas J. Tustison, Philip A. Cook, Russell T. Shinohara, Haochang Shou, and Alzheimer's Disease Neuroimaging Initiative. Mitigating site effects in covariance for machine learning in neuroimaging data. Human Brain Mapping, 43(4):1179–1195, March 2022. ISSN 1097-0193. doi: 10.1002/hbm.25688.

9. Stephen R. Cole and Constantine E. Frangakis. The Consistency Statement in Causal Inference: A Definition or an Assumption? Epidemiology, 20(1):3, January 2009. ISSN 1044-3983. doi: 10.1097/EDE.0b013e31818ef366.

10. Cameron Craddock, Sikka Sharad, Cheung Brian, Khanuja Ranjeet, Ghosh Satrajit, Yan Chaogan, Li Qingyang, Lurie Daniel, Joshua Vogelstein, Randal Burns, Colcombe Stanley, Maarten Mennes, Kelly Clare, Di Adriana, Francisco Castellanos, and Milham Michael. Towards Automated Analysis of Connectomes: The Configurable Pipeline for the Analysis of Connectomes (C-PAC). Front. Neuroinf., 7, Jan 2013. doi: 10.3389/conf.fninf.2013.09.00042.

11. Adriana Di Martino, Chao-Gan Yan, Qingyang Li, Erin Denio, Francisco X Castellanos, Kaat Alaerts,

Jeffrey S Anderson, Michal Assaf, Susan Y Bookheimer, Mirella Dapretto, et al. The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. Molecular psychiatry, 19(6):659–667, 2014.

12. Adriana Di Martino, David O'connor, Bosi Chen, Kaat Alaerts, Jeffrey S Anderson, Michal Assaf, Joshua H Balsters, Leslie Baxter, Anita Beggiato, Sylvie Bernaerts, et al. Enhancing studies of the connectome in autism using the autism brain imaging data exchange ii. Scientific data, 4(1):1–15, 2017.

13. Peng Ding and Luke W. Miratrix. To Adjust or Not to Adjust? Sensitivity Analysis of M-Bias and Butterfly-Bias. Journal of Causal Inference, 3(1):41–57, Mar 2015. ISSN 2193-3685. doi: 10.1515/jci-2013-0021.

14. Bradley Efron. Large-Scale Simultaneous Hypothesis Testing. J. Am. Stat. Assoc., 99(465):96–104, Mar 2004. ISSN 0162-1459. doi: 10.1198/016214504000000089.

15. V. S. Fonov, A. C. Evans, R. C. McKinstry, C. R. Almli, and D. L. Collins. Unbiased nonlinear average age-appropriate brain templates from birth to adulthood. Neuroimage, 47:S102, Jul 2009. ISSN 1053-8119. doi: 10.1016/S1053-8119(09)70884-5.

16. Wolfgang Forstmeier, Eric-Jan Wagenmakers, and Timothy H. Parker. Detecting and avoiding likely false-positive findings – a practical guide. Biol. Rev., 92(4):1941–1968, Nov 2017. ISSN 1464-7931. doi: 10.1111/brv.12315.

17. Jean-Philippe Fortin, Nicholas Cullen, Yvette I. Sheline, Warren D. Taylor, Irem Aselcioglu, Philip A. Cook, Phil Adams, Crystal Cooper, Maurizio Fava, Patrick J. McGrath, Melvin McInnis, Mary L. Phillips, Madhukar H. Trivedi, Myrna M. Weissman, and Russell T. Shinohara. Harmonization of cortical thickness measurements across scanners and sites. Neuroimage, 167:104, February 2018. doi: 10.1016/j.neuroimage.2017.11.024.

18. K. J. Friston, S. Williams, R. Howard, R. S. Frackowiak, and R. Turner. Movement-related effects in fMRI time-series. Magn. Reson. Med., 35(3):346–355, Mar 1996. ISSN 0740-3194. doi: 10.1002/mrm.1910350312.

19. Jianqiao Ge, Guoyuan Yang, Meizhen Han, Sizhong Zhou, Weiwei Men, Lang Qin, Bingjiang Lyu, Hai Li, Haobo Wang, Hengyi Rao, Zaixu Cui, Hesheng Liu, Xi-Nian Zuo, and Jia-Hong Gao. Increasing diversity in connectomics with the Chinese Human Connectome Project. Nature Neuroscience, 26(1):163–172, January 2023. ISSN 1546-1726. doi: 10.1038/s41593-022-01215-1.

20. Douglas N. Greve and Bruce Fischl. Accurate and robust brain image alignment using boundary-based registration. Neuroimage, 48(1):63–72, Oct 2009. ISSN 1095-9572. doi: 10.1016/j.neuroimage.2009.06.060.

21. Michelle Hampson, Fuyuze Tokoglu, Xilin Shen, Dustin Scheinost, Xenophon Papademetris, and R. Todd Constable. Intrinsic Brain Connectivity Related to Age in Young and Middle Aged Adults. PLoS One, 7(9):e44067, Sep 2012. ISSN 1932-6203. doi: 10.1371/journal.pone.0044067.

22. Sheema Hashem, Sabah Nisar, Ajaz A. Bhat, Santosh Kumar Yadav, Muhammad Waqar Azeem, Puneet Bagga, Khalid Fakhro, Ravinder Reddy, Michael P. Frenneaux, and Mohammad Haris. Genetics of structural and functional brain changes in autism spectrum disorder. Translational Psychiatry, 10, 2020. doi: 10.1038/s41398-020-00921-3.

23. Daniel E Ho, Kosuke Imai, Gary King, and Elizabeth A Stuart. MatchIt: Nonparametric preprocessing for parametric causal inference. Journal of Statistical Software, 42(8):1–28, 2011. URL https://www.jstatsoft.org/v42/i08/.

24. Jian Huang, Joel L. Horowitz, and Fengrong Wei. VARIABLE SELECTION IN NONPARAMETRIC ADDITIVE MODELS. Annals of statistics, 38(4):2282, August 2010. doi: 10.1214/09-AOS781.

25. Madhura Ingalhalikar, Alex Smith, Drew Parker, Theodore D. Satterthwaite, Mark A. Elliott, Kosha Ruparel, Hakon Hakonarson, Raquel E. Gur, Ruben C. Gur, and Ragini Verma. Sex differences in the structural connectome of the human brain. Proc. Natl. Acad. Sci. U.S.A., 111(2):823–828, Jan 2014. ISSN 0027-8424. doi: 10.1073/pnas.1316909110.

26. Mark Jenkinson, Peter Bannister, Michael Brady, and Stephen Smith. Improved optimization for the robust and accurate linear registration and motion correction of brain images. Neuroimage, 17 (2):825–841, Oct 2002. ISSN 1053-8119. doi: $10.1016/\mathrm{s}1053\text{-}8119(02)91132\text{-}8$.

27. Mark Jenkinson, Christian F. Beckmann, Timothy E. J. Behrens, Mark W. Woolrich, and Stephen M. Smith. FSL. Neuroimage, 62(2):782–790, Aug 2012. ISSN 1095-9572. doi: $10.1016/\mathrm{j}.\mathrm{neuroimage}.2011.09.015$.

28. W. Evan Johnson, Cheng Li, and Ariel Rabinovic. Adjusting batch effects in microarray expression data using empirical Bayes methods. Biostatistics, 8(1):118–127, Jan 2007. ISSN 1465-4644. doi: $10.1093/\mathrm{biostatistics}/\mathrm{kxj}037$.

29. Nicole R. Karcher and Deanna M. Barch. The ABCD study: understanding the development of risk for mental and physical health outcomes - Neuropsychopharmacology. Neuropsychopharmacology, 46(1):131–142, Jan 2021. ISSN 1740-634X. doi: $10.1038/\mathrm{s}41386\text{-}020\text{-}0736\text{-}6$.

30. Gregory Kiar, Eric W. Bridgeford, William R. Gray Roncal, , Vikram Chandrashekhar, Disa Mhembere, Sephira Ryman, Xi-Nian Zuo, Daniel S. Margulies, R. Cameron Craddock, Carey E. Priebe, Rex Jung, Vince D. Calhoun, Brian Caffo, Randal Burns, Michael P. Milham, and Joshua T. Vogelstein. A High-Throughput Pipeline Identifies Robust Connectomes But Troublesome Variability. bioRxiv, page 188706, Apr 2018. URL https://doi.org/10.1101/188706.

31. Kerstin Konrad and Simon B. Eickhoff. Is the ADHD brain wired differently? A review on structural and functional connectivity in attention deficit hyperactivity disorder. Human Brain Mapping, 31(6): 904, June 2010. doi: $10.1002/\mathrm{hbm}.21058$.

32. Cosmin Lazar, Stijn Meganck, Jonatan Taminau, David Steenhoff, Alain Coletta, Colin Molter, David Y. Weiss-Solís, Robin Duque, Hugues Bersini, and Ann Nowé. Batch effect removal methods for microarray gene expression data integration: a survey. Briefings Bioinf., 14(4):469–490, Jul 2013. ISSN 1477-4054. doi: $10.1093/\mathrm{bib}/\mathrm{bbs}037$.

33. Jeffrey T Leek and Roger D Peng. Statistics: P values are just the tip of the iceberg. http://www.nature.com/news/statistics-p-values-are-just-the-tip-of-the-iceberg-1.17412, April 2015. Accessed: 2020-10-12.

34. Jeffrey T Leek, Robert B Scharpf, Héctor Corrada Bravo, David Simcha, Benjamin Langmead, W Evan Johnson, Donald Geman, Keith Baggerly, and Rafael A Irizarry. Tackling the widespread and critical impact of batch effects in high-throughput data. Nat. Rev. Genet., 11(10):733–739, October 2010.

35. Michael J. Lopez and R. Gutman. Matching to estimate the causal effects from multiple treatments, 2014. URL https://www.semanticscholar.org/paper/Matching-to-estimate-the-causal-effects-from-Lopez-Gutman/11ca8c8fa81498fe75349cff0001552c89115c8c. [Online; accessed 9. Feb. 2023].

36. Scott Marek, Brenden Tervo-Clemmens, Finnegan J. Calabro, David F. Montez, Benjamin P. Kay, Alexander S. Hatoum, Meghan Rose Donohue, William Foran, Ryland L. Miller, Timothy J. Hendrickson, Stephen M. Malone, Sridhar Kandala, Eric Feczko, Oscar Miranda-Dominguez, Alice M. Graham, Eric A. Earl, Anders J. Perrone, Michaela Cordova, Olivia Doyle, Lucille A. Moore, Gregory M. Conan, Johnny Uriarte, Kathy Snider, Benjamin J. Lynch, James C. Wilgenbusch, Thomas Pengo, Angela Tam, Jianzhong Chen, Dillan J. Newbold, Annie Zheng, Nicole A. Seider, Andrew N. Van, Athanasia Metoki, Roselyne J. Chauvin, Timothy O. Laumann, Deanna J. Greene, Steven E. Petersen, Hugh Garavan, Wesley K. Thompson, Thomas E. Nichols, B. T. Thomas Yeo, Deanna M. Barch, Beatriz Luna, Damien A. Fair, and Nico U. F. Dosenbach. Reproducible brain-wide association studies require thousands of individuals. Nature, 603:654–660, March 2022. ISSN 1476-4687. doi: $10.1038/\mathrm{s}41586\text{-}022\text{-}04492\text{-}9$.

37. Lukas Meier, Sara van de Geer, and Peter Bühlmann. High-dimensional additive modeling. Annals of Statistics, 37(6B):3779–3821, December 2009. ISSN 0090-5364. doi: $10.1214/09\text{-}AOS692$.

38. Maria B. Misiura, J. Christina Howell, Junjie Wu, Deqiang Qiu, Monica W. Parker, Jessica A. Turner, and William T. Hu. Race modifies default mode connectivity in Alzheimer's disease. Transl. Neurodegener., 9, 2020. doi: 10.1186/s40035-020-0186-4.

39. N N Tzourio-Mazoyer, B Landeau, D Papathanassiou, F Crivello, O Etard, N Delcroix, B Mazoyer, and M Joliot. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. Neuroimage, Jan 2002. doi: 10.1006/nimg.2001. 0978.

40. Derek Evan Nee. fMRI replicability depends upon sufficient individual-level data. Commun. Biol., 2 (130):1–4, Apr 2019. ISSN 2399-3642. doi: 10.1038/s42003-019-0378-6.

41. Stephanie Noble, Dustin Scheinost, Emily S. Finn, Xilin Shen, Xenophon Papademetris, Sarah C. McEwen, Carrie E. Bearden, Jean Addington, Bradley Goodyear, Kristin S. Cadenhead, Heline Mirzakhanian, Barbara A. Cornblatt, Doreen M. Olvet, Daniel H. Mathalon, Thomas H. McGlashan, Diana O. Perkins, Aysenil Belger, Larry J. Seidman, Heidi Thermenos, Ming T. Tsuang, Theo G. M. van Erp, Elaine F. Walker, Stephan Hamann, Scott W. Woods, Tyrone D. Cannon, and R. Todd Constable. Multisite reliability of MR-based functional connectivity. Neuroimage, 146:959–970, Feb 2017. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2016.10.020.

42. Kate Brody Nooner, Stanley Colcombe, Russell Tobe, Maarten Mennes, Melissa Benedict, Alexis Moreno, Laura Panek, Shaquanna Brown, Stephen Zavitz, Qingyang Li, Sharad Sikka, David Gutman, Saroja Bangaru, Rochelle Tziona Schlachter, Stephanie Kamiel, Ayesha Anwar, Caitlin Hinz, Michelle Kaplan, Anna Rachlin, Samantha Adelsberg, Brian Cheung, Ranjit Khanuja, Chaogan Yan, Cameron Craddock, Vincent Calhoun, William Courtney, Margaret King, Dylan Wood, Christine Cox, Clare Kelly, Adriana DiMartino, Eva Petkova, Philip Reiss, Nancy Duan, Dawn Thompsen, Bharat Biswal, Barbara Coffey, Matthew Hoptman, Daniel C. Javitt, Nunzio Pomara, John Sidtis, Harold Koplewicz, Francisco Xavier Castellanos, Bennett Leventhal, and Michael Milham. The NKI-Rockland Sample: A Model for Accelerating the Pace of Discovery Science in Psychiatry. Front. Neurosci., 6, Oct 2012. ISSN 1662-453X. doi: 10.3389/fnins.2012.00152.

43. Denise C. Park and Chih-Mao Huang. Culture Wires the Brain: A Cognitive Neuroscience Perspective. Perspectives on psychological science : a journal of the Association for Psychological Science, 5(4):391, July 2010. doi: 10.1177/1745691610374591.

44. Massimiliano Pastore and Antonio Calcagnì. Measuring Distribution Similarities Between Samples: A Distribution-Free Overlapping Index. Front. Psychol., 0, 2019. ISSN 1664-1078. doi: 10.3389/fpsyg.2019.01089.

45. Judea Pearl. Causal inference in statistics: An overview. ssu, 3(none):96–146, Jan 2009. ISSN 1935-7516. doi: 10.1214/09-SS057.

46. Judea Pearl. On measurement bias in causal inference. AUAI Press, Arlington, TX, USA, Jul 2010. ISBN 978-0-97490396-5. doi: 10.5555/3023549.3023599.

47. Judea Pearl. Interpretation and identification of causal mediation. Psychological Methods, 19(4): 459–481, December 2014. ISSN 1939-1463. doi: 10.1037/a0036434.

48. Raymond Pomponio, Guray Erus, Mohamad Habes, Jimit Doshi, Dhivya Srinivasan, Elizabeth Mamourian, Vishnu Bashyam, Ilya M. Nasrallah, Theodore D. Satterthwaite, Yong Fan, Lenore J. Launer, Colin L. Masters, Paul Maruff, Chuanjun Zhuo, Henry Völzke, Sterling C. Johnson, Jurgen Fripp, Nikolaos Koutsouleris, Daniel H. Wolf, Raquel Gur, Ruben Gur, John Morris, Marilyn S. Albert, Hans J. Grabe, Susan M. Resnick, R. Nick Bryan, David A. Wolk, Russell T. Shinohara, Haochang Shou, and Christos Davatzikos. Harmonization of large MRI datasets for the analysis of brain imaging patterns throughout the lifespan. Neuroimage, 208:116450, Mar 2020. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2019.116450.

49. Michael Powell, Allison Koenecke, James Brian Byrd, Akihiko Nishimura, Maximilian F. Konig, Ruoxuan Xiong, Sadiqa Mahmood, Vera Mucaj, Chetan Bettegowda, Liam Rose, Suzanne Tamang, Adam Sacarny, Brian Caffo, Susan Athey, Elizabeth A. Stuart, and Joshua T. Vogelstein. A how-to

guide for conducting retrospective analyses: example COVID-19 study. OSF Preprints, Sep 2020. doi: 10.31219/osf.io/3drch.

50. Paul R. Rosenbaum and Donald B. Rubin. The central role of the propensity score in observational studies for causal effects. Biometrika, 70(1):41–55, Apr 1983. ISSN 0006-3444. doi: 10.1093/biomet/70.1.41.

51. Paul R. Rosenbaum and Donald B. Rubin. Constructing a control group using multivariate matched sampling methods that incorporate the propensity score. The American Statistician, 39(1):33–38, 1985. ISSN 00031305. URL http://www.jstor.org/stable/2683903.

52. Roser Sala-Llonch, David Bartrés-Faz, and Carme Junqué. Reorganization of brain networks in aging: a review of functional connectivity studies. Front. Psychol., 0, 2015. ISSN 1664-1078. doi: 10.3389/fpsyg.2015.00663.

53. Theodore D. Satterthwaite, Daniel H. Wolf, David R. Roalf, Kosha Ruparel, Guray Erus, Simon Vandekar, Efstathios D. Gennatas, Mark A. Elliott, Alex Smith, Hakon Hakonarson, Ragini Verma, Christos Davatzikos, Raquel E. Gur, and Ruben C. Gur. Linked Sex Differences in Cognition and Functional Connectivity in Youth. Cerebral Cortex (New York, NY), 25(9):2383, Sep 2015. doi: 10.1093/cercor/bhu036.

54. Cencheng Shen, Carey E Priebe, and Joshua T Vogelstein. From Distance Correlation to Multiscale Generalized Correlation. Journal of American Statistical Association, October 2017. URL http://arxiv.org/abs/1710.09768.

55. Elizabeth A. Stuart. Matching Methods for Causal Inference: A Review and a Look Forward. Statist. Sci., 25(1):1–21, Feb 2010. ISSN 0883-4237. doi: 10.1214/09-STS313.

56. Gábor J. Székely, Maria L. Rizzo, and Nail K. Bakirov. Measuring and testing dependence by correlation of distances. Ann. Stat., 35(6):2769–2794, Dec 2007. ISSN 0090-5364. doi: 10.1214/009053607000000505.

57. Benjamin O. Turner, Erick J. Paul, Michael B. Miller, and Aron K. Barbey. Small sample sizes reduce the replicability of task-based fMRI studies. Commun. Biol., 1(62):1–10, Jun 2018. ISSN 2399-3642. doi: 10.1038/s42003-018-0073-z.

58. A. W. van der Vaart. Asymptotic Statistics. Cambridge University Press, Cambridge, England, UK, October 1998. ISBN 978-0-51180225-6. doi: 10.1017/CBO9780511802256.

59. Eleanna Varangis, Christian G. Habeck, Qolamreza R. Razlighi, and Yaakov Stern. The Effect of Aging on Resting State Connectivity of Predefined Networks in the Brain. Front. Aging Neurosci., 0, 2019. ISSN 1663-4365. doi: 10.3389/fnagi.2019.00234.

60. Joshua T. Vogelstein, Eric W. Bridgeford, Benjamin D. Pedigo, Jaewon Chung, Keith Levin, Brett Mensh, and Carey E. Priebe. Connectal coding: discovering the structures linking cognitive phenotypes to individual histories. Current Opinion in Neurobiology, 55:199–212, April 2019. ISSN 0959-4388. doi: 10.1016/j.conb.2019.04.005.

61. Joshua T Vogelstein, Eric W Bridgeford, Qing Wang, Carey E Priebe, Mauro Maggioni, and Cencheng Shen. Discovering and deciphering relationships across disparate data modalities. Elife, 8, January 2019. URL http://dx.doi.org/10.7554/eLife.41690.

62. Christian Wachinger, Anna Rieckmann, and Sebastian Pölsterl. Detect and Correct Bias in Multi-Site Neuroimaging Datasets. arXiv, Feb 2020. doi: 10.1016/j.media.2020.101879.

63. Xueqin Wang, Wenliang Pan, Wenhao Hu, Yuan Tian, and Heping Zhang. Conditional Distance Correlation. J. Am. Stat. Assoc., 110(512):1726, 2015. doi: 10.1080/01621459.2014.993081.

64. Susanne Weis, Kaustubh R. Patil, Felix Hoffstaedter, Alessandra Nostro, B. T. Thomas Yeo, and Simon B. Eickhoff. Sex Classification by Resting State Brain Connectivity. Cereb. Cortex, 30(2):824–835, Mar 2020. ISSN 1047-3211. doi: 10.1093/cercor/bhz129.

65. Daniel Westreich, Jessie K. Edwards, Catherine R. Lesko, Stephen R. Cole, and Elizabeth A. Stuart. Target Validity and the Hierarchy of Study Designs. Am. J. Epidemiol., 188(2):438, February 2019. doi: 10.1093/aje/kwy228.

66. Ayumu Yamashita, Noriaki Yahata, Takashi Itahashi, Giuseppe Lisi, Takashi Yamada, Naho Ichikawa, Masahiro Takamura, Yujiro Yoshihara, Akira Kunimatsu, Naohiro Okada, Hirotaka Yamagata, Koji Matsuo, Ryuichiro Hashimoto, Go Okada, Yuki Sakai, Jun Morimoto, Jin Narumoto, Yasuhiro Shimada, Kiyoto Kasai, Nobumasa Kato, Hidehiko Takahashi, Yasumasa Okamoto, Saori C. Tanaka, Mitsuo Kawato, Okito Yamashita, and Hiroshi Imamizu. Harmonization of resting-state functional MRI data across multiple imaging sites via the separation of site differences into sampling bias and measurement bias. PLoS Biol., 17(4):e3000042, Apr 2019. ISSN 1545-7885. doi: 10.1371/journal.pbio.3000042.

67. Meichen Yu, Kristin A. Linn, Philip A. Cook, Mary L. Phillips, Melvin McInnis, Maurizio Fava, Madhukar H. Trivedi, Myrna M. Weissman, Russell T. Shinohara, and Yvette I. Sheline. Statistical harmonization corrects site effects in functional connectivity measurements from multi-site fMRI data. Hum. Brain Mapp., 39(11):4213–4227, Nov 2018. ISSN 1065-9471. doi: 10.1002/hbm.24241.

68. Yuqing Zhang, Giovanni Parmigiani, and W. Evan Johnson. ComBat-seq: batch effect adjustment for RNA-seq count data. NAR Genomics Bioinf., 2(3), Sep 2020. ISSN 2631-9268. doi: 10.1093/nargab/lqaa078.

69. Xi-Nian Zuo, Jeffrey S. Anderson, Pierre Bellec, Rasmus M. Birn, Bharat B. Biswal, Janusch Blautzik, John C. S. Breitner, Randy L. Buckner, Vince D. Calhoun, F. Xavier Castellanos, Antao Chen, Bing Chen, Jiangtao Chen, Xu Chen, Stanley J. Colcombe, William Courtney, R. Cameron Craddock, Adriana Di Martino, Hao-Ming Dong, Xiaolan Fu, Qiyong Gong, Krzysztof J. Gorgolewski, Ying Han, Ye He, Yong He, Erica Ho, Avram Holmes, Xiao-Hui Hou, Jeremy Huckins, Tianzi Jiang, Yi Jiang, William Kelley, Clare Kelly, Margaret King, Stephen M. LaConte, Janet E. Lainhart, Xu Lei, Hui-Jie Li, Kaiming Li, Kuncheng Li, Qixiang Lin, Dongqiang Liu, Jia Liu, Xun Liu, Yijun Liu, Guangming Lu, Jie Lu, Beatriz Luna, Jing Luo, Daniel Lurie, Ying Mao, Daniel S. Margulies, Andrew R. Mayer, Thomas Meindl, Mary E. Meyerand, Weizhi Nan, Jared A. Nielsen, David O'Connor, David Paulsen, Vivek Prabhakaran, Zhigang Qi, Jiang Qiu, Chunhong Shao, Zarrar Shehzad, Weijun Tang, Arno Villringer, Huiling Wang, Kai Wang, Dongtao Wei, Gao-Xia Wei, Xu-Chu Weng, Xuehai Wu, Ting Xu, Ning Yang, Zhi Yang, Yu-Feng Zang, Lei Zhang, Qinglin Zhang, Zhe Zhang, Zhiqiang Zhang, Ke Zhao, Zonglei Zhen, Yuan Zhou, Xing-Ting Zhu, and Michael P. Milham. An open science resource for establishing reliability and reproducibility in functional connectomics. Sci. Data, 1(140049):1–13, Dec 2014. ISSN 2052-4463. doi: 10.1038/sdata.2014.49.

Appendix A. Procedures for detecting and mitigating batch effects.

A.1 Detecting batch effects with CAUSAL CDCORR Many of the more direct types of detectable effects, such as associational and conditional effects, fail to adaequately account for confounding biases present in the data. We instead propose the use of CAUSAL cDcorr, in which a conditional K-sample test [63] is performed on samples with the same "range" of covariate values after propensity trimming via a strategy known as vector matching [35]. This strategy is the focus of a complementary theoretical manuscript, in which we illustrate the theoretical and empirical (via simulations) benefits of this technique over competing approaches for detecting causal effects between potential outcomes. This strategy maintains both high sensitivity and specificity under traditionally problematic data regimes (high-dimensionality, non-monotonicities, and non-linearities) in which other methods typically fail [7], making it a natural choice for causal discrepancy testing (of which "batch effect" detection, termed *causal unconditional discrepancy testing* in [7], as-defined herein is a special case).

Figure 2C illustrates visually the causal procedures employed for adjusting the batches. Rather than fully matching to produce the adjusted batches, samples are retained only such that they have approximately overlapping covariate distributions (propensity trimming, shaded boxes). A full schematic illustrating the use of vector matching is detailed in Appendix E of [7]. In the event that the batch assignment mechanism is ignorable given the covariates and that effects between datasets are in the same direction across all covariate levels, the adjusted effect detected by CAUSAL cDcorr is a causal batch effect, as proven in [7]. That the effects between datasets are in the same "direction" across all covariate levels can be best intuited via example. Consider the case where a batch effect exists, such that there is a signal disparity (a difference in the expected connectivity) in a particular edge of a connectome between two batches. If this signal disparity is a positive difference between batches 1 and 2 across all covariate levels (or, a negative difference across all covariate levels), CAUSAL cDcorr will detect a causal batch effect. In the event that the assignment mechanism is ignorable but that effects between datasets are not in the same direction across all covariate levels, the adjusted effect detected by CAUSAL cDcorr is a causal effect, but may reflect a *causal conditional discrepancy* (e.g., there are batch-specific differences, but they are isolated to particular covariate levels). In the event that the batch assignment mechanism is not ignorable, the effect may not reflect any causal effects (e.g., it may reflect unmeasured demographic differences between the batches).

A.2 Mitigating batch effects using Causal ComBat Unfortunately, many existing techniques for the removal of batch effects fail to adequately account for confounding biases that may be present in the data. We propose Causal ComBat, in which Conditional ComBat is performed on a subset of observational studies in which all pairs of studies are balanced against measured demographic covariates. Causal ComBat is performed as follows. Given measurements and covariates from n individuals across K batches, each of which has n_k individuals:

1. Identify the smallest dataset k' as the exposure dataset and all $k \neq k'$ datasets as control datasets.
2. Propensity trim and then match the greatest integer less than $n_k/n_{k'}$ control individuals using nearest-neighbor matching to individuals in the exposure dataset. Methods B discusses the adjustment procedure in more detail.
3. Discard individuals from the exposure dataset who do not have matches in all K control datasets, yielding the matched exposure cohort.
4. Discard individuals from the control datasets who are not matched to the matched exposure cohort, yielding the matched control cohorts.
5. Perform Conditional ComBat [28] on the measurements of the matched exposure and matched control individuals across the K batches, conditioned on the measured covariates X.

In the event that the conditioning set closes backdoor paths [45, 46], Causal ComBat yields the removal of an internally valid causal effect and does not require extrapolation assumptions, unlike Conditional

`ComBat` [23, 50, 51, 55]. If the conditioning set does not close backdoor paths, the effect removed is a conditional effect and may potentially yield the removal of demographic effects, as we saw in Figure 4. Appendix E depicts the impact on the empirical covariate distribution of the adjustment procedure.

Appendix B. Methods (Expanded). For a formal description of the below definitions, see Bridgeford et al. [7]. Informally, we adopt the following notation to describe batch effect correction. We assume that Y_i represents an observed measurement of interest (e.g., a connectome), T_i represents the batch in which the measurement is collected, X_i represents observed covariates, (e.g., age or biological sex), and Z_i represents unobserved covariates (e.g., height). The data (Y_i, T_i, X_i, Z_i) are n independent and identical samples from some true and unknown data generating distribution. The quantity $Y_i^{(t)}$ represents the measurement that would have been observed, had the measurement been collected in a given batch t. The key distinction is that Y_i represents the *actual* measurement, which is collected and studied. On the other hand, $Y_i^{(t)}$ is a potentially hypothetical measurement, which is only actually observed for the case where $T_i = t$ in most experimental contexts [9]. That is, individuals have a potential measurement $Y_i^{(t)}$ for every possible batch, but only a single of these potential measurements are studied under standard observational contexts. This is known as the *consistency assumption*, and can be written mathematically as:

$$Y_i = \sum_{t \in \mathcal{T}} \mathbf{1}_{\{T_i = t\}} Y_i^{(t)} = Y_i^{(T_i)},$$

where $\mathcal{T}$ is the set of batch labels. That the observed measurements Y_i given the batch (or measured covariates) and the potential measurements $Y_i^{(t)}$ do not, in general, have similar distributions is known as the "fundamental problem of causal inference." This problem requires resorting to additional assumptions to make conclusions on the basis of the observed data.

Using this convention, a batch effect is defined in Definition 1.

Definition 1 (Batch Effect). *A batch effect exists between two batches t and t' if $Y_i^{(t)}$ and $Y_i^{(t')}$ have different distributions.*

In light of this definition, a batch effect can be conceptualized as the *potential* measurements having different distributions between a given pair of batches. A batch effect is present if an individual's potential measurements (which are random quantities) would differ (in terms of their distributions) *by virtue of them being measured* in batches t and t'.

The major conceptual leap is that, under the manner in which most mega-studies are collected (where individuals are *unique* to each batch), Definition 1 is about *potential* measurements rather than *realized* measurements. Causal claims regarding potential measurements will only be valid on the basis of the observed (realized) data insofar as the reasonableness of the assumptions upon which they rest.

Associational Effects In an associational context, we observe measurements y_i and batches t_i for each individual $i \in [n]$, so effects can only be determined from realizations of Y_i and T_i.
Definition 2 (Associational Effect). *An associational effect exists between batches t and t' if $Y_i | T_i = t$ and $Y_i | T_i = t'$ have different distributions.*

Likewise, we can use this intuition to develop a suitable definition for associational batch effect correction:
Definition 3 (Associational Effect Correction). *Associational effect correction is a function g, where:*

$$g\left(\{(Y_i, T_i)\}_{i \in [n]}\right) = \left\{\tilde{Y}_i\right\}_{i \in [n]}$$

such that for all pairs of batches t and t', $\tilde{Y}_i | T_i = t$ and $\tilde{Y}_i | T_i = t'$ have the same distribution.

Whereas Y_i represents the measurement for an individual, the intuition is that $\tilde{Y}_i$ can be thought of as the "corrected" measurement, which does not have an associational effect for any pair of batches in the observed data.

Associational effects will often poorly characterize whether a batch effect is present. Consider, for instance, if in one batch t, we tend to measure older individuals, whereas in another batch t', we tend to measure younger individuals. If age is related to the measurement we obtained, then the differences between $Y_i | T_i = t$ and $Y_i | T_i = t'$ could be due to age or batch identity, and we have no way of differentiating whether the effect is a *bona fide* batch effect versus merely an associational effect. A sufficient condition for an associational effect to facilitate detecting or estimating a batch effect would be that individuals are randomized to each batch, in that individuals are randomly assigned to be measured in particular batches *pre-hoc*. Associational effect detection can be facilitated via DCORR [56], and associational effect correction can be facilitated via ComBat [28].

Conditional Effects In a conditional context, we observe measurements y_i, batches t_i, and covariates x_i for all individuals $i \in [n]$, so effects can be determined from realizations of Y_i, T_i, and X_i.

Definition 4 (Conditional Effect). *A conditional effect exists between batches t and t' if for some covariate x, $Y_i | T_i = t, X_i = x$ and $Y_i | T_i = t', X_i = x$ have different distributions.*

We can define conditional effect correction using this logic:

Definition 5 (Conditional effect correction). *Conditional effect correction is a function g, where:*

$$g\Big(\{(Y_i, T_i, X_i)\}_{i \in [n]} \Big) = \Big\{ \tilde{Y}_i \Big\}_{i \in [n]}$$

such that for all pairs of batches t and t' and for all covariate levels x, $\tilde{Y}_i | T_i = t, X_i = x$ and $\tilde{Y}_i | T_i = t', X_i = x$ have the same distribution.

Conceptually, a conditional effect is equivalent to a batch effect if two conditions hold:

1. the measured covariates overlap in distribution between the batches (propensity overlap), and
2. the measured covariates provide all of the information regarding the mechanism about how people ended up in one batch versus the other (strong ignorability).

The former condition denotes that both batches must contain similar groups of people (in terms of measured covariates), and the latter condition specifies that the measured covariates X_i tell us all of the information needed to "exchange" measurements from one batch to the other. Borrowing the preceding example, even if we observe more young people in a batch t, we must still observe *some* young people in the other batch t'. In this sense, the measured covariates contain the information needed to "deconfound" disparities that might be batch effects or veridical effects due to upstream covariates. Therefore, when we make subsequent comparisons, we do not need to guess what people with similar covariates would have looked like in the other batch, and vice versa.

In this fashion, our comparisons can be thought of as locally (with respect to the covariates) exchanging a realized measurement Y_i in batch t with a realized measurement Y_i in batch t', where both individuals have similar covariates x. Intuitively, these comparisons can therefore be conceptualized as synthetically comparing $Y_i^{(t)}$ and $Y_i^{(t')}$ (the target estimand for establishing a batch effect) by using observed measurements Y_i from individuals who are similar on the basis of the covariates X_i between the two batches. Condition 1 ensures that we can make this intuitive step over the entire span of covariates in our batches. Conditional effect detection can be facilitated via cDcorr [63], and conditional effect correction can be facilitated via cComBat [28].

In practice, we never know whether the propensity distributions overlap; we can only estimate them from the data. If our estimated propensities do not overlap given finite data, we again cannot reasonably differentiate between differences in the two groups being due to *bona fide* batch effects or empirical differences in the propensity distributions. This motivates a third approach.

Adjusted Effects As before, we observe measurements y_i, batches t_i, and covariates x_i for all individuals $i \in [n]$, and we determine effects from realizations of Y_i, T_i, and X_i. Prior to assessing the equality of any distributions, however, we weight the observations such that the observed covariate distributions are rendered approximately overlapping.

Definition 6 (Adjusted Effect). *An adjusted effect exists between batches t and t' if after re-weighting samples such that $X_i|T_i = t$ and $X_i|T_i = t'$ are approximately overlapping (or, alternatively, approximately equal) in distribution, $Y_i|T_i = t, X_i = x$ and $Y_i|T_i = t', X_i = x$ have different distributions.*

We can similarly define adjusted effect correction as:

Definition 7 (Adjusted Effect Correction). *Assume that the samples are re-weighted via weights w_i (possibly 0 or 1) such that after re-weighting, $X_i|T_i = t$ and $X_i = t'$ are approximately overlapping (or alternatively, approximately equal) in distribution for all $w_i \neq 0$.*
Adjusted effect correction is a function g, where:

$$g\Big(\{(Y_i, T_i, X_i, w_i)\}_{i\in[n]}\Big) = \Big\{\tilde{Y}_i\Big\}_{i\in[n]}$$

such that for all pairs of batches t and t' and for all covariate levels x, $\tilde{Y}_i|T_i = t, X_i = x$ and $\tilde{Y}_i|T_i = t', X_i = x$ have the same distribution.

Adjusted effects, by default, satisfy the first criterion for a conditional effect to be a batch effect. This is because rendering measured covariate distributions approximately equal intuitively is a more strict criterion than simply ensuring that they approximately overlap. The reason that we believe that re-weighting to ensure approximate covariate distribution equality is desirable for effect correction, versus simply approximate covariate overlap, is discussed in Section 3.3 and Appendix D.2. Adjusted effect detection can be facilitated via CAUSAL CDCORR [7], and adjusted effect correction can be facilitated with Causal cComBat (described in Section 2.2).

We still must satisfy the latter criterion for a conditional effect to be a batch effect; that is, given the measured covariates, we can ignore how people ended up in one batch versus the other. This assumption has the same interpretation as before.

Crossover Effects We observe measurements $y_i^{(t)}$ and covariates $x_i^{(t)}$, for all individuals $i \in [n]$ and for all batches $t \in \mathcal{T}$. In this case, we can make statements on the basis of $Y_i^{(t)}$ itself, because we actually observe outcomes for each possible batch. Therefore, we typically will not need to resort to local exchangeability (similar individuals/covariates across batches) as before, unless there are aspects of the individuals changing from one batch to another. *Crossover effects in general require the fewest assumptions to derive causal conclusions, since we directly observe all possible potential measurements for each individual.*

Definition 8 (Crossover Effect, constant states). *A crossover effect exists between batches t and t' if, given that $\left(X_i^{(t)}, Z_i^{(t)}\right)$ and $\left(X_i^{(t')}, Z_i^{(t')}\right)$ are sufficiently similar, $Y_i^{(t)}$ and $Y_i^{(t')}$ have the same distribution.*

We are certain that any *traits* of the participants (i.e., variables that are constant for a given participant, such as genome) are the same across the two groups since the group members are identical (even if we did not measure these traits). However, *states* may differ as they may be functions of location or time. For example, if being measured impacts subsequent states, then a crossover effect may not be indicative of a batch effect without further assumptions and/or experimental design considerations (such as randomizing exposure order across participants, or resorting to adjusted effect strategies as before if these states are measured).

In the case where participant states are unchanging or are randomized, we can resort directly to associational strategies for subsequent detection or correction, and can make causal conclusions about batch effects with few additional considerations. In the case where participant states are changing and

are not randomized but are measured, we can resort to adjusted strategies for adjusted effect detection or correction, via a crossover effect for non-constant states:

Definition 9 (Crossover Effect, non-constant states). *A crossover effect exists between batches t and t' if, after re-weighting samples such that $X_i^{(t)}$ and $X_i^{(t')}$ are approximately overlapping (or, alternatively, approximately equal) in distribution, $Y_i^{(t)}$ and $Y_i^{(t')}$ have the same distribution.*

For this effect to be a true batch effect, we need to make the same assumptions as before; that is, that given the measured covariates (which include the changing states), we can ignore how people ended up in one batch versus the other.

B.1 Covariate Adjustment The exposed group t is selected to be the smaller of the two groups, and the unexposed group t' is selected to be the larger of the two groups, where n_t is the number of individuals in the exposed group and $n_{t'}$ is the number of individuals in the unexposed group. The "covariate overlap" procedure attempts to ensure positivity of the propensity distribution for the unexposed group (i.e., $e(t'|x) > 0$). Intuitively, we exclude individuals from the unexposed group who do not appear "similar" to individuals in the exposed group. In this work, covariate overlap is established through logistic regression by excluding individuals in the control group with a propensity for exposure lower than 0.01. Similarly, the "covariate balancing" procedure attempts to re-weight observations in the per-batch covariate distributions such that the covariate distributions are approximately equal (i.e., $f(x|t) \approx f(x|t')$). In this work, we perform $k : 1$ nearest-neighbor matching using the `MatchIt` package [23]. The number of matches $k = \lfloor \frac{n_{t'}}{n_t} \rfloor$ is chosen to be the largest number of unexposed matches possible. Individuals are balanced on the basis of individual sex, individual age, and continent of study. As there are likely many other categories with which brain connectivity may be confounded, we do not believe this covariate set is necessarily sufficient to identify a Causal Batch Effect 1, as we would need to be confident that these covariates exhibited the covariate sufficiency property. We perform exact matching on the basis of individual sex and individual continent of measurement, and use a 0.1-standard deviation caliper on the propensity score to obtain at most k matched participants for each treated individual [49].

B.2 Hypothesis Testing Recall that statements of the form $f(y) = g(y)$ against $f(y) \neq g(y)$ are equivalent to $P_f = P_g$ against $P_f \neq P_g$, as probability densities uniquely define distribution functions. Therefore, hypotheses for the effects described in Section 2 require two-sample and conditional two-sample testing procedures. A natural test statistic for the two-sample testing procedure is the Distance Correlation [56], which is a non-parametric test for testing whether two variables are correlated. A simple augmentation of the distance correlation procedure [54, 61] shows that `DCorr` can be used for the two-sample test, or a test of whether two samples are drawn from different distributions. `DCorr` is exactly equivalent in this context to the Maximum Mean Discrepency (MMD), which embeds points in a reproducing kernel Hilbert Space (RKHS) and looks for functions over the unit ball in the RKHS that maximize the difference of the means of the embedded points. When we instead consider the conditional two-sample test (i.e., a test of $f(y|x) = g(y|x)$ against $f(y|x) \neq g(y|x)$), we instead use the conditional distance correlation [63], a kernel-based approach in which the points are embedded in a new, non-linear Hilbert Space, which augments the traditional linear Hilbert Space used in distance correlation to allow the definition of the squared distance covariance. Below, we let $\mathbf{Y} = (y_i) \in \mathcal{Y}^n$ denote realizations across both samples and $\vec{t} = (t_i) \in \{t, t'\}^n$ indicate from which sample each realization is drawn. $\mathbf{X} = (x_i) \in \mathcal{X}^n$ denotes covariates which are known about the objects of interest.

Associational Effect We have the following null and alternative hypotheses:

$$H_0 : f(y|t) = f(y|t') \text{ against } H_A : f(y|t) \neq f(y|t')$$

A test of the preceding hypotheses is performed using the distance correlation, and the natural test statistic is $\text{DCorr}(\mathbf{Y}, \vec{t})$.

Conditional Effect We have the following null and alternative hypotheses:

$$H_0 : f(y|t,x) = f(y|t',x) \text{ against } H_A : f(y|t,x) \neq f(y|t',x)$$

A test of the preceding hypotheses is performed using the conditional distance correlation, and the natural test statistic is $\texttt{cDCorr}(\mathbf{Y}, \vec{t}|\mathbf{X})$.

Adjusted Conditional Effect We have the following null and alternative hypotheses:

$$H_0 : \tilde{f}(y|t,x) = \tilde{f}(y|t',x) \text{ against } H_A : \tilde{f}(y|t,x) \neq \tilde{f}(y|t',x)$$

Unlike the preceding tests, we instead consider the data $(\tilde{\mathbf{Y}}, \vec{\tilde{t}}, \tilde{\mathbf{X}})$, which are the measurements, sample indicators, and covariates of the n realizations after covariate adjustment. A test of the preceding hypotheses is performed using the conditional distance correlation, and the natural test statistic is $\texttt{cDCorr}(\tilde{\mathbf{Y}}, \vec{\tilde{t}}|\tilde{\mathbf{X}})$.

Causal Crossover Effect We have the following null and alternative hypotheses:

$$H_0 : f\left(y^{(t)}|t, x^{(t)}\right) = f\left(y^{(t')}|t', x^{(t')}\right) \text{ against } H_A : H_0 : f\left(y^{(t)}|t, x^{(t)}\right) \neq f\left(y^{(t')}|t', x^{(t')}\right)$$

If the known covariates are identical between batches t and t', we test the preceding hypotheses using the distance correlation, and the natural test statistic is $\texttt{DCorr}(\mathbf{Y}, \vec{t})$. If the known covariates are not identical between batches t and t', we test the preceding hypotheses using the conditional distance correlation, and the natural test statistic is $\texttt{cDCorr}(\mathbf{Y}, \vec{t}|\mathbf{X})$.

More Than Two Batches The above approaches generalize sufficiently to K batches using K-sample testing approaches [7]. With f_k for $k \in [K]$ denoting the densities associated with K sites or batches, this motivates hypotheses of the form:

$$H_0 : f_k = f_l \text{ for all } k,l \text{ against } H_A : f_k \neq f_l \text{ for some } k \neq l,$$

which can be tested using the distance correlation or the conditional distance correlation as above, with the caveat that $\vec{t}$ becomes the matrix $\mathbf{T} = (t_{i,k}) \in \{0,1\}^{n \times K}$. Each entry $t_{ik} = 1$ if sample i is in batch k, and 0 otherwise. For the purposes of this manuscript, we focus on the two-batch case.

p-values and Multiple Hypothesis Correction p-values in this manuscript are estimated using permutation testing, which is an approach to obtain the distribution of the test statistic under the null with minimal assumptions and approximations [14]. All p-values are estimated using $N = 1,000$ (detection, Figure 5) or $N = 10,000$ (correction, Figure 6) permutations. Across all figures associated with this work, we are concerned with obtaining a proper estimate of the rate at which we detect effects (*discoveries*). Therefore, we control the false discovery rate (FDR) with the Benjamini-Hochberg Correction [5].

B.3   Batch effect correction via z-scoring We compare z-scoring per-feature within each batch [32] and traditional $\texttt{ComBat}$ to Causal $\texttt{ComBat}$. For a given edge $y_i^{(t)}(k,l)$ (k,l) for an individual i in batch t, we define the z-score corrected edge-weight to be:

$$z_i^{(t)}(k,l) = \frac{y_i^{(t)}(k,l) - \bar{y}^{(t)}(k,l)}{s^{(t)}(k,l)},$$

where $\bar{y}^{(t)}(k,l)$ is the sample mean and $s^{(t)}(k,l)$ is the sample standard deviation for edge (k,l) in batch t. These strategies reduce known topological properties of human connectomes; particularly, homotopy, as shown in Figure 16 and Figure 15 and are therefore not considered for more careful analysis.

B.4 Control Numerical Experiments Control experiments are performed to ensure that after batch effect correction, the resulting data maintains interpretability and utility for scientific inquiry. Even if the data is devoid of batch effects, it must still be useful for downstream inference. For our connectome data, we identify two key control scenarios, the preservation of demographic effects and the preservation topological effects, which are desirable to preserve after batch effect correction.

Testing Topological Properties

Demographic Effect Demographic effects are investigated across both the subset of connectomes upon which Causal `ComBat` is executed (the American Clique) and the full dataset. We observe the tuple $(y_i(k,l), s_i, a_i, t_i)$ for $i \in [n]$, and $k, l \in [V]$, where $V = 116$ denotes the number of parcels in the Automated Anatomical Labelling (AAL) parcellation [39]. We suppose that $Y(k,l)$ is the $[0,1]$-valued random variable denoting the weight of edge (k,l), S is the binary-valued random variable denoting the biological sex, A is the positive real-valued random variable denoting age, and T is the $[K]$-valued random variable denoting the batch. We let $\vec{y}(k,l) = (y_i(k,l)) \in [0,1]^n$ denote the realized edge weights, $\vec{s} = (s_i) \in \{0,1\}^n$ denote the realized biological sexes, $\vec{a} = (a_i) \in \mathbb{R}^n$ denote the realized biological ages, and $\vec{t} = (t_i) \in [K]^n$ denote the realized batches. We say that a **demographic sex effect** exists when:

$$f(y|a,s) \neq f(y|a,s')$$

To test for a demographic sex effect, we have the following null and alternative hypotheses:

$$H_0 : f(y|a,s) = f(y|a,s') \text{ against } f(y|a,s) \neq f(y|a,s')$$

We are able to test the preceding hypothesis using the partial distance correlation, and the test statistic is $\text{pDCorr}(\vec{y}(k,l), \vec{s}|\vec{a})$.

B.5 Datasets The CoRR Mega-Study [69] is an aggregate dataset consisting of 27 studies collected with a similar goal: assessing the reliability and reproducibility of neuroimaging data. The mega-study consists of 1313 individuals, most of whom are measured numerous times, for a total of $N = 3597$ connectomes. All connectomes are estimated using the `m2g` (MRI to Graphs) pipeline [30], which provides a wrapper for the CPAC Pipeline [10]. fMRI scans for each individual are first processed to remove motion artifacts using `mcflirt` [26]. The fMRI scans are then registered to the corresponding individual's anatomical scan using FSL's boundary-based registration (BBR) via `epireg` [20]. A non-linear transformation from the individual's anatomical scan to the MNI152 [15] template is estimtaed using `FNIRT` [27]. Nuisance artifacts are removed by fitting the voxelwise timeseries to a regression model incorporating regressors for the Friston 24-parameter model [18], the top five principal components of the voxelwise timeseries in cerebrospinal fluid `aCompCor` [4], and a quadratic drift term. The adjusted voxelwise timeseries is downsampled to the regions of interest (ROIs) of the Automated Anatomical Labelling (AAL) parcellation [39] by taking the spatial mean signal for each timepoint across voxels within the region of interest. Functional connectivity is estimated using the pairwise correlation between all pairs of ROI timeseries within the AAL parcellation. Parcels are sorted throughout the manuscript according to hemispheric order, in which the parcels are aligned with left hemisphere parcels followed by right hemisphere parcels. Within hemisphere, parcels are sorted by AAL parcel number. For each study, we have baseline covariates for the continent, sex, and the age of participants.

The American Clique The "American Clique" describes a subset of the CoRR Mega-Study in which the sample populations share similarities in sample demographic characteristics. These studies share a demographic focusing on males and females (in roughly equal proportions) of individuals across a wide age range, and include the "NYU2", "IBATRT", "MRN1", "UWM", and "NYU1" studies. The 833 connectomes comprising the studies of the American clique are reduced to the $N = 321$ connectomes with maximal demographic overlap identified through covariate adjustment B.1.

The NKI Rockland Sample [42] is a single study from the CoRR Mega-Study consisting of 24 individuals, each of whom is measured two times across three functional MRI acquisition protocols, which vary in the repetition time for each slice of the sequence (TR). The data was collected with the intention of investigating the impact of the different MRI protocols in a crossover-randomized approach. Due to the crossover property, evidence in favor of an effect provides strong evidence of a causal batch effect. Images with a TR of 645 millisecond (ms), 1400 ms, and 2500 ms are measured, with the prompt for each subject remaining identical.

Appendix C. Correction for Batch-Related Effects.

Appendix D. Simulations.

D.1   Batch Effect Detection Simulations Simulations illustrating the sensitivity (high testing power when a causal effect is present) and specificity (tests which do not falsely detect effects) of CAUSAL CDCORR for causal effect detection are in Bridgeford et al. [7].

D.2   Batch Effect Removal Simulations $n = 400$ points are sampled from Batch 0 or Batch 1 with probability 0.5; e.g. $T_i \overset{iid}{\sim} \mathrm{Bern}(0.5)$.

Covariate distributions For each row, the covariate distributions are:

I. Overlap The covariate distribution is $X_i \overset{iid}{\sim} 2\mathrm{Beta}(2,2) - 1$.

II. Moderate Overlap The covariate distribution is:

$$X_i | T_i = t \overset{d}{\sim} \begin{cases} 2\mathrm{Beta}(2,6) - 1 & t = 0 \\ 2\mathrm{Beta}(6,2) - 1 & t = 1 \end{cases}$$

III. Limited Overlap The covariate distribution is:

$$X_i | T_i = t \overset{d}{\sim} \begin{cases} 2\mathrm{Beta}(2,12) - 1 & t = 0 \\ 2\mathrm{Beta}(12,2) - 1 & t = 1 \end{cases}$$

Simulation contexts We investigate these in four simulation contexts, where for all simulations, $\epsilon_i \overset{iid}{\sim} \mathcal{N}(0,1)$:

Non-linearity A sigmoidal relationship between the covariate and the outcome. The outcome is:

$$Y_i = 4\mathsf{sigmoid}(8X_i) + T_i + \frac{1}{2}\epsilon_i,$$

where $\mathsf{sigmoid}(x)$ is the non-linear sigmoid function; e.g.:

$$\mathsf{sigmoid}(x) = \frac{1}{1 + \exp(-x)}.$$

The effectiveness of batch effect correction methods for the non-linearity setting are shown in Figure 4.

Symmetric non-monotonicity A gaussian non-monotonic relationship between the covariate and the outcome. The outcome is:

$$Y_i = \varphi\left(X_i, \mu = 0, \sigma = \frac{1}{2}\right) + T_i + \frac{1}{2}\epsilon_i,$$

where $\varphi(x, \mu, \sigma)$ is the probability density function of the Normal distribution:

$$\varphi(x, \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{ \left(\frac{x - \mu}{\sigma}\right)^2 \right\}$$

Algorithm 1 Causal ComBat. The Causal `ComBat` approach for batch effect correction. Conditional `ComBat` is performed on demographically matched batches to eliminate confounding biases in observational studies. The smallest batch is arbitrarily chosen as the treated batch, and then successive batches are matched against the treated batch. Treated individuals (as well as their matched controls) with no suitable matches in one (or more) batches are eliminated from the analysis. `CovariateMatch()` is a function which takes the treatment and control covariates and matches control individuals to treatment individuals. $\mathcal{I}_{mat}^k$ is a list of the treated individuals with at least one match, and $\mathcal{M}_{mat}^k$ is a list of the treated individual that each control individual was (potentially) matched to. `BalanceCovariates()` uses these lists to identify the treatment individuals who have matches across all control batches and then excludes control individuals who are not matched to treatment individuals with matches across all batches.

Input: The tuples $(\mathbf{Y}^{(k)}, \mathbf{X}^{(k)})$ for $k = 1, \ldots, K$, where $\mathbf{Y}^{(k)} \in \mathbb{R}^{n_k \times p}$ denotes the measurements from batch id k, and $\mathbf{X}^{(k)} \in \mathbb{R}^{n_k \times d}$ denotes the covariates from batch id k, where $N = \sum_k n_k$ is the total number of measurements.

Output: The $N' \times p$ batch effect corrected data, where $N' \leq N$.

1: **function** CAUSALCOMBAT($\mathbf{Y}, \mathbf{X}, \vec{t}$)
2:      $\tilde{\mathbf{Y}}, \tilde{\mathbf{X}}, \tilde{n} = $ BalanceCovariates($\mathbf{Y}, \mathbf{X}$)
3:      Let $\vec{t} = (1_{\tilde{n}_1}, 2_{\tilde{n}_2}, \ldots, K_{\tilde{n}_K})$ ▷ vector indicating the batch id of each individual
4:      $\tilde{\mathbf{Y}}_n = $ ComBat($\tilde{\mathbf{Y}}, \tilde{\mathbf{X}}, \vec{t}$) ▷ Remove the effect due to the batch $\vec{t}$, conditional on the covariates $\mathbf{X}$
5:      **return** $\tilde{\mathbf{Y}}_n$ ▷ Return the batch
6: **end function**
7: **function** BALANCECOVARIATES($\mathbf{Y}, \mathbf{X}$)
8:      $k' = \operatorname{argmin}_k n_k$ ▷ Select treatment study to be the smallest study
9:      **for** $k \neq k'$ **do**
10:          $\mathcal{M}_{mat}^k, \mathcal{I}_{mat}^k = $ CovariateMatch($\mathbf{X}^{(k')}, \mathbf{X}^{(k)}$)   ▷ Covariate match the two studies
11:      **end for**
12:      $\mathcal{I}_{mat} = \bigcap_{k \neq k'} \mathcal{I}_{mat}^k$ ▷ Identify treatment individuals with a control match across all control groups
13:      $\mathbf{Y}^{(k')} = \mathbf{Y}^{(k')}[\mathcal{I}_{mat}, :]$ ▷ Remove other treatment individuals
14:      $\mathbf{X}^{(k')} = \mathbf{X}^{(k')}[\mathcal{I}_{mat}, :]$
15:      $n_{k'} = |\mathcal{I}_{mat}|$ ▷ Number of treated individuals remaining
16:      **for** $k \neq k$ **do**
17:          $\mathcal{J}_{mat}^k = $ which($\mathcal{M}_{mat}^k$ %in% $\mathcal{I}_{mat}$)
                 ▷ Control individuals matched to treatment individuals with matches across all groups
18:          $\mathbf{Y}^{(k)} = \mathbf{Y}^{(k)}[\mathcal{J}_{mat}^k, :]$ ▷ Remove other control individuals
19:          $\mathbf{X}^{(k)} = \mathbf{X}^{(k)}[\mathcal{J}_{mat}^k, :]$
20:          $n_k = |\mathcal{J}_{mat}^k|$ ▷ Number of control individuals remaining
21:      **end for**
22:      $\mathbf{Y} = [\mathbf{Y}^{(1)}; \ldots; \mathbf{Y}^{(K)}]$ ▷ raw measurements across batches
23:      $\mathbf{X} = [\mathbf{X}^{(1)}; \ldots; \mathbf{X}^{(K)}]$ ▷ covariates across batches
       return $\mathbf{Y}, \mathbf{X}, n$
24: **end function**

We call it "symmetric" because $\mu = 0$, which leads to the both the covariate/outcome relationship and the covariates themselves being symmetric about $x = 0$. The effectiveness of batch effect correction methods for the symmetric non-monotonicity setting are shown in Figure 7.

Symmetric non–monotonicity

Figure 7: A symmetric non-monotonicity about $x = 0$. `cComBat` provides reasonable results when the non-monotonicity and the covariate distributions (for each batch) are symmetric about the same value (here, $x = 0$). In high balance the MAAD of `cComBat` and `Causal cComBat` are 0.076 and 0.051 (`Causal cComBat` better 67.6% of the time), in moderate balance are 0.179 and 0.173 (`Causal cComBat` better 50.9% of the time), and in limited overlap are 0.358 and 0.289 (`Causal cComBat` better 66.7% of the time) with `Causal cComBat` only executing in 36 simulations.

Asymmetric non-monotonicity A gaussian non-monotonic relationship between the covariate and the outcome. The outcome is:

$$Y_i = \varphi\left(X_i, \mu = -0.5, \sigma = \frac{1}{2}\right) + T_i + \frac{1}{2}\epsilon_i.$$

We call it "asymmetric" because $\mu = -0.5$, which leads to the effect not being symmetric about $x = 0$ (whereas the covariates are, by construction, symmetric about $x = 0$). The effectiveness of batch effect correction methods for the asymmetric non-monotonicity setting are shown in Figure 8.

Symmetric non-monotonicity, asymmetric covariates A gaussian non-monotonic relationship between the covariate and the outcome, and a different covariate distribution from the rest of the simulations. The covariate distribution is:

$$X_i | T_i = t \overset{d}{\sim} \begin{cases} 2\text{Beta}(2, 2) - 1 & t = 0 \\ 2\text{Beta}(b, 2) - 1 & t = 1 \end{cases}$$

where $b = 2$, $b = 6$, and $b = 12$ for the overlap, moderate, and limited conditions, respectively.

The outcome is:

$$Y_i = \varphi\left(X_i, \mu = 0, \sigma = \frac{1}{2}\right) + T_i + \frac{1}{2}\epsilon_i.$$

This time, the non-monotonicity is symmetric, but the covariate distributions are increasingly asymmetric. The effectiveness of batch effect correction methods for the asymmetric non-monotonicity setting are shown in Figure 9.

Asymmetric non−monotonicity

Figure 8: An asymmetric non-monotonicity in which the non-monotonicity is not symmetric about $x = 0$, while the covariate distributions (for each batch) are symmetric about $x = 0$. cComBat struggles and worsens the batch effect as the covariate balance decreases, while Causal cComBat is able to work as long as there are moderate amounts of covariate overlap. In high balance the MAAD of cComBat and Causal cComBat are 0.0497 and 0.0421 (Causal cComBat better 59.5% of the time), in moderate balance are 0.074 and 0.130 (Causal cComBat better 99.7% of the time), and in limited overlap are 2.61 and 0.407 (Causal cComBat always better) with Causal cComBat only executing in 33 simulations.

Linear A linear relationship between the covariate and the outcome. The outcome is:

$$Y_i = -2(X_i - 1) + T_i + \frac{1}{2}\epsilon_i.$$

The effectiveness of batch effect correction methods for the linear setting are shown in Figure 10.

D.3 Mean Average Absolute Difference (Mean AAD) To evaluate the effectiveness of each batch effect correction technique on simulated data, we compute the true data expected signal for each covariate level; e.g., $\mathbb{E}\left[Y_i^{(t)}|x\right]$ for each batch t and each covariate level x. Since ϵ_i has mean 0, this would be the quantity:

$$\mathbb{E}\left[Y_i^{(t)}|x\right] = f(x) + t,$$

where f is the covariate/outcome relationship (possibly incorrectly modeled). Since x is continuous, we compute this value for x across $B = 200$ evenly spaced breakpoints for evaluation. Given a set of samples, we train a batch effect correction model, and fit the trained model to the expected signal for each covariate level, leaving us with corrected expected signal for each batch, which we denote by $v(t, x)$. In theory, if the batch effect correction technique removes the batch effect as modeled, $v(1, x) \approx v(0, x)$ for all x. To evaluate each technique, we consider the average absolute difference for each trial r of 1000 trials:

$$d_r = \frac{1}{B}\sum_{x \in \mathcal{X}} |v(1, x) - v(0, x)|,$$

Figure 9: A symmetric non-monotonicity about $x = 0$, but the covariate distribution is no longer symmetric about $x = 0$. `cComBat` struggles to remove the batch effect and leaves a residual signal disparity, but `Causal cComBat` is able to address the batch effect within a range of covariate overlap. In high balance the MAAD of `cComBat` and `Causal cComBat` are 0.075 and 0.0506 (`Causal cComBat` better 65.8% of the time), in moderate balance are 0.181 and 0.0751 (`Causal cComBat` better 96.0% of the time), and in limited overlap are 0.84 and 0.106 (`Causal cComBat` always better). Due to the fact that the covariate distribution for batches 0 and 1 overlap for this simulation, `Causal cComBat` always executes.

and the magnitudes of d_r are annotated in the plots (shaded red boxes) for a single simulation. We compute the mean AAD as:

$$d = \frac{1}{r} \sum_{r \in [R]} d_r.$$

A value of 0 corresponds to the batch effect being completely eliminated (the expected signal for each batch after correction is identical), a value of 1 would equate to the residual disparity between the expected signals being the same as before batch effect correction, and a value > 1 corresponds to the residual disparity between the expected signals *exceeding* the original batch effect.

D.4 Causal `ComBat` outperforms Conditional `ComBat` on simulated data Across all simulations, simulated measurements are collected from an additive gaussian model with either a linear (Figure 10), non-linear (Figure 4), or non-monotonic (Figure 7, Figure 8, and Figure 9) covariate/outcome relationship, with a batch-specific offset. We performed these simulations for cases where the covariate distributions overlap (row I), moderately overlap (row II), and have limited overlap (row III). When the assumptions between the covariate and the measurements are correct, covariate overlap is irrelevant, as the covariate-specific signal can be extrapolated successfully across the entire range of the covariate. When the assumptions between the covariate and the measurements are incorrect, the batch effect can still be removed, as shown in column (D), when further conditions are satisfied, such as:

1. covariate distributions are matched: can be conceptualized as the inaccurate assumptions of the model, in effect, being uniformly *misrepresented* by the model in each of the two batches, which *still* enables a consistent estimate of the batch effect to be estimated and removed. `Causal cComBat` attempts to enforce this approach through matching. We explored the implications of this in Figure 4 and Figure 8.

Figure 10: A linear simulation in which the assumptions of `cComBat` are perfectly satisfied. In high balance the MAAD of `cComBat` and `Causal cComBat` are 0.035 and 0.0375 (`Causal cComBat` better 42.5% of the time), in moderate balance are 0.074 and 0.113 (`Causal cComBat` better 33.1% of the time), and in limited overlap are 0.145 and 0.292 (`Causal cComBat` better 23.1% of the time) with `Causal cComBat` only executing in 39 simulations.

2. the covariate distributions (and the covariate/outcome relationship) are symmetric about the same point: again, the idea here is that intuitively, the the inaccurate assumptions of the model must be uniformly misrepresented by the model in the two batches. This is a pre-hoc condition and relates directly to the specific underlying true veridical signal, and is difficult to exploit practically. We explore the implications of this in Figure 7 and Figure 9.

As shown in the simulations in Figure 4 and Figure 8, these artifacts can be arbitrarily large depending on the nature of the incorrect assumption. In these examples, the residual signal disparity introduced by `cComBat` in limited overlap cases (row III) dwarfs the magnitude of the original batch effect, and is of similar magnitude with moderate covariate overlap (row II).

On the other hand, `Causal cComBat` first pre-aligns the covariate distributions, and even with model misspecifications, is able to successfully remove a batch effect for the subset of the covariate space with sufficient covariate overlap. Importantly, causal strategies *explicitly prohibit* learning about data which falls outside of the region of demographic overlap. When overlap conditions are not met for a portion of the sample space, causal strategies do not do anything, as is typically the case for most of the illustrated similations in the "Limited Overlap" condition. In this sense, causal techniques are therefore more conservative for batch effect detection and mitigation. We believe this to be favorable for batch effect detection and mitigation as, in practice, it is effectively impossible to tell whether the implicit or explicit assumptions of the model for batch effect correction are sufficient or not, and almost impossible for our model to account for all ways a batch effect could manifest in high-dimensional data (such as higher order correlations, non-linearities, variance differences) given a finite sample. This presents an enormous point of caution when applying batch effect removal techniques to high-dimensional datasets with confounding, as it is infeasible to explore whether extrapolatory modeling assumptions are within reason on a dimension-by-dimension (or, higher-order) basis.

Addressing the batch effect One way to characterize the batch effect in univariate contexts (and within our simulations) would be analogous to the average treatment effect [50]; e.g.:

$$\text{ATE} = \mathbb{E}[Y(1) - Y(0)]$$

For all simulations, by construction the ATE is 1 and does not depend on the covariates.

By definition, a difference in the expectations implies that $Y(1) \overset{\mathcal{D}}{\neq} Y(0)$, and a batch effect is present by Definition 1.

It is well understood in the statistical literature that for models of the form $Y_i = g(X_i) + \beta T_i + Z_i$ where $Z_i \overset{iid}{\sim} \mathcal{N}(\mu, \Sigma)$, that as long as either the strong ignorability condition [50] or f is known, the adjustment factor $\hat{\beta}_1 - \hat{\beta}_0$ will be a consistent and unbiased estimate of the ATE [50, 58]. For a non-technical explanation in words, we can obtain reasonable estimates of the ATE in one of three ways (and consequently, account for the batch effect):

1. Extrapolation: if we know $g(x)$, then we can "guess" what $\mathbb{E}[Y_i(t)|x]$ looks like for values of x that do not exist in our batch, by appropriate estimation of g via the elements of our dataset that we *do* have. In this case, where the model is linear, extrapolation can be performed by linear regression. Once we have identified $g(x)$, we can extrapolate what $g(x)$ looks like for values of x which fall outside the range of our dataset, as in Figure 10; particularly, noting cells (C).II and (C).III.
2. Use asymptotics: if we do not know $g(x)$ but we can reasonably assume that the strong ignorability condition holds, we can leverage asymptotic statistics to conclude that we can still adequately estimate the CATE. Proofs of this result can be found in [50, 51]. In this particular case, $X_i \perp\!\!\!\perp T_i$, which implies the strong ignorability condition [50]. For more trivial reading specific to the case where $X_i \perp\!\!\!\perp T_i$, see [58]. This result is exhibited in the simulations for Figure 4 and Appendix Figure 8 looking at column (D) compared to column (C), and noting that causal pre-processing procedures tend to produce a reasonable result (while non-causal conditional procedures with a model misspecification do not).
3. The covariate distribution is symmetric about the same point that a non-monotonicity is symmetric: in this extremely special case, we have that $f(x|1) = f(-x|0)$, and $f(y(0)|x) = f(y(1)|-x)$ for every x. This result is exhibited by way of comparison of Figures 7 and Figure 8, and noting that while Figure 7(C).III is able to arrive at a reasonable solution, Figure 8(C).III does not.

By a similar argument, the estimated batch disparity can also be consistently estimated in either of these contexts. In these cases, removal of the estimated offset associated with the particular batch of a measurement is a reasonable approach to removal of the estimated batch effect.

Simulations in which the assumption about $f(x)$ is incorrect and X_i is distributed conditionally on T_i (and consequently, the strong ignorability condition will not hold), will not necessarily yield a reasonable estimate of the ATE.

Appendix E. Overlap of Empirical Covariates.

The empirical overlap of the covariate distributions is difficult to compute in the case of data without making heavy parametric assumptions. For this reason, we turn to the distribution-free overlapping index [44]. The distribution free overlapping index, $\hat{v}_{kl}$, is computed by first approximating the density of the distribution of the measured covariates for each dataset d, $X = (A, S, C)$, where A is a random variable whose realizations $a \in \mathcal{A}$ are ages, S is a random variable whose realizations $s \in \mathcal{S}$ are sexes (M or F), and C is a random variable whose realizations $c \in \mathcal{C}$ denote continent, using the base R function `stats::density`. The random variable D has realizations $d \in \mathcal{D}$ whose realizations denote dataset. The density $f_d(a|S = s, C = c)$ is the conditional density of age, conditional on the individual's sex being s, continent of measurement being c for a given dataset d. The mass $\mathbb{P}_d(S = s|C = c)$ is the conditional mass of sex, conditional on the individual's continent of measurement being c, for dataset d.

Finally, the mass $\mathbb{P}_d(C = c)$ represents the mass of an individual's continent of measurement being c, for dataset d (0 or 1 for all d, since all individuals from dataset d are either measured on continent c or not). An estimate of the overlap between the two densities, $\hat{v}_{kl}$ between datasets k and l, is computed using the formula:

$$\hat{v}_{kl} = \sum_{c \in \mathcal{C}} \min\Big(\hat{\mathbb{P}}_k(C = c), \hat{\mathbb{P}}_l(C = c)\Big) \sum_{s \in \{\mathsf{M},\mathsf{F}\}} \min\Big(\hat{\mathbb{P}}_k(S = s | C = c), \hat{\mathbb{P}}_l(S = s | C = c)\Big) \Bigg[$$

$$\int_{\mathcal{A}} \min\Big(\hat{f}_d(a | S = s, C = c), \hat{f}_l(a | S = s, C = c)\Big) \, \mathrm{d}a \Bigg]$$

which is obtained via numerical quadrature.

Intuitively, this can be conceptualized as representing the mass of the "area under the curve" which is shared by the two densities for datasets k and l.

Figure 11: **The overlap of the empirical covariate distributions for the American Clique, before and after adjustment**. **(A)** The empirical distributions of the covariates before adjustment. **(B)** The empirical overlap of the covariate distributions after the adjustment procedure is applied, as discussed in Appendix C.

Associational and Crossover effect detection As for conditional and adjusted effects in Figure 5, we can perform a similar investigation for associational or causal crossover effects, which we add in Appendix Figure 13. Associational effects can almost always be detected (405 of 406 pairs of datasets), regardless of whether or not any demographic overlap is present or any comparison seems reasonable (distance correlation, BH correction, $\alpha = 0.05$). This highlights that associational effects, perhaps, can be best thought of as determining whether or not the data is simply different across a given pair of batches, and are not particularly useful for attribution of the source of a detected difference (such as to batch effects, or demographic effects). The NKI Rockland Sample takes this a step further, allowing a crossover effect to be determined in a few pairs of batches (purple squares). This study was, in fact,

Figure 12: **The overlap of the empirical covariate distributions for two mega-studies. (A)** The empirical distribution of covariates. **(B)** The overlap of covariate distributions given by the distribution-free overlapping index for ABIDE mega-study [11, 12]. **(C)** The overlap of covariate distributions given by the distribution-free overlapping index for the SRPBS mega-study. [66]. Like for the CoRR mega-study, while several pairs of sites have overlapping demographic distributions, many of the sites have extremely poor overlap in both mega-studies. In these cases, attempts to normalize for batch effects using model-based approaches like Conditional ComBat would be subject to the pitfalls of Figure 4, Appendix 8, or Appendix Figure 9 if modeling assumptions are not reasonable.

collected on an identical set of participants, in the same location, with the same research technicians, with data varying only in the repetition time for the magnet used in the scanner. Causal crossover effects, which require the fewest assumptions to be bonified causal batch effects, are present for 2 of

Figure 13: **Comparison of associational and causal crossover effects between pairs of datasets from the CoRR study**. **(A)** A heatmap analogous to Figure 5(A), but for associational and causal crossover effects. Most associational effects are significant (405 of 406) and most causal crossover effects are siginficant (2 of 3 pairs of datasets with the same individuals measured multiple times). **(B)** analogous to Figure 5(B), but for associational and conditional effects.

3 possible combinations of datasets (distance correlation, BH correction, $\alpha = 0.05$). Unfortunately, this requires a very specific recruitment setup (that is, the same individuals measured across the batches) which is not met for the 403 remaining possible pairs of datasets.

Batch Effects are reduced with Causal `ComBat` Here we investigate batch effects before and after correction. Figure 14 shows average connectomes from the MRN1 and IBATRT studies. With no correction, while the average connectomes appear similar between the two studies, a difference between the average connectomes is present (bottom row). Three different batch effect mitigation strategies (Z-scoring across features within a single batch, `ComBat`, and Causal `ComBat`) all appear to eliminate the differences between the average connectomes. Z-scoring has disrupted fundamental topological properties of the human connectome, as explored in detail in Appendix F. For this reason, we do not consider Z-scoring going forward, and focus instead on variations of the `ComBat` procedure.

Appendix F. `ComBat` **Preserves Topological Properties of fMRI Connectomes.** The presence of two prominent topological *signal effects* in human connectomes, homophily and homotopy, are investigated in Figure 15. The left column of Figure 15A looks at the edges comprising these two disparate community structures. When edges are sorted by hemisphere, the homophilic effect is characterized by a modular two-block structure, with higher connectivity in the lower left and upper right blocks (dark red) than in the two off-diagonal blocks (pink). The homotopic effect is characterized by edges in the off-diagonal band (dark blue) showing higher connectivity than other edges within the connectome (light blue). The right column of Figure 15A looks at a single individual's connectome from the NYU2 study before and after correction. The homotopic property appears to be preserved in only the `ComBat` connectomes, whereas Z-scored connectomes do not appear to preserve this property. In

Average connectomes across 2 datasets with similar covariate distribution

Figure 14: **Distribution of Detected Effects Before and After Correction**. A comparison of the average connectomes from two studies from the same continent, with similar sex and age distributions conditional on sex. While the *Raw* average connectomes appear similar, the absolute difference between the two is appreciable. All of the three adjustment strategies reduce the absolute difference between the average connectomes for the two batches.

Figure 15B, these two properties are examined statistically for the individuals from the American Clique (Mann-Whitney U Statistic, see Methods B.4 for details). Points falling along the diagonal line $y = x$ tend to have a similar signal effect before and after batch correction. Whereas `ComBat` strategies tend to preserve the homophilic and homotopic effects, Z-scoring tends to decrease these two topological properties of functional connectomes. This is noted by the fact that the majority of the points for Z-scoring across both effects tend to fall below the $y = x$ line, indicating that the effect size after Z-scoring is lower than the effect size in the raw connectomes.

Figure 16 explores the distribution of edge weights within these two community structures before and after correction. The conclusions are similar in that Causal `ComBat` and `ComBat` preserve the property that the homotopic edges have far higher connectivity than non-homotopic edges.

Figure 15: **Preservation of Topological Properties of Connectomes after Batch Correction**. The presence of topological effects is explored in the context of the homophilic and homotopic differences in connectivity in the American Clique. **(A)** (*Left*) A map indicating which edges are associated with homophilic/non-homophilic or homotopic/non-homotopic edge communities. (*Right*) The edge-weight normalized connectome associated with a single individual from the NYU2 study using each removal strategy. **(B)** A scatter plot, showing the effect size (Mann-Whitney U Statistic) before (x axis) and after (y axis) the indicated Batch Effect Correction Method.

Figure 16: Histogram and rug plot indicating the empirical distribution of the normalized edge weights associated with each of the four edge communities for a single individual's connectome.