1   **Wheat *EARLY FLOWERING3* is a dawn-expressed circadian oscillator**

2   **component that regulates heading date**

Lukas Wittern[1][¶], Gareth Steed[1][¶], Laura J. Taylor[1], Keith Gardner[2], Andy Greenland[2],

Matthew A. Hannah[3], Alex A. R. Webb[1]*

[1]Department of Plant Sciences, University of Cambridge, Downing Street,

Cambridge, CB2 3EA, UK

[2]BASF, BBCC – Innovation Center Gent, Technologiepark-Zwijnaarde, Gent,

Belgium

[3]NIAB, Huntingdon Road, Cambridge, UK

*Corresponding author

Email: aarw2@cam.ac.uk (AARW)

[¶]These authors contributed equally to this work

Abstract

Optimising the seasonal control of flowering in the major crops is an important component of breeding to match crop adaptation to the target environment. Using an eight parent Multiparent Advanced Generation Inter-Cross (MAGIC) population we investigated the contribution of variation at circadian clock-associated genes to the regulation of heading date in UK and European winter wheat varieties. We identified homoeologues of *EARLY FLOWERING* 3 (*ELF3*) as candidate genes for the *Earliness per se* (*Eps*) *D1* and *B1 loci* in field conditions. We confirmed that a SNP within the coding region of *TaELF3-B1* is the likely causal polymorphism underlying the *Eps-B1 locus.* We also identified that a reported deletion at the *Eps-D1 locus* encompassing *TaELF3-D1,* is in fact a novel allele that lies within an introgression region that contains an inversion relative to the Chinese Spring D genome. Our findings that *ELF3* might be associated with the regulation of heading date prompted us to investigate whether *ELF3* is a circadian oscillator gene in wheat, as it is in Arabidopsis. Using *T. turgidum cv. Kronos* carrying loss of function alleles for both copies of *TtELF3,* we found that circadian rhythms were severely disrupted. Furthermore, in *T. aestivum*, we found that loss of functional *LUX ARRHYTHMO* (*LUX*), an orthologue of the protein partner of ELF3 in Arabidopsis, also severely disrupted circadian rhythms. Whilst these data suggest a function for both *ELF3* and *LUX* in the wheat circadian oscillator, that oscillator might be structured differently to that of Arabidopsis because wheat *ELF3* and *LUX* transcripts are maximal at the end of the night and day respectively, rather than co-expressed at dusk as they are in Arabidopsis. We conclude that there is sufficient allelic diversity within the three wheat *ELF3* homoeologues for selection to delay or advance heading, and that this can be achieved without pleiotropic deleterious alterations to circadian rhythms.

Introduction

Circadian clocks are endogenous timing mechanisms with a near 24 hour period that sequence biological events, provide anticipation of environmental rhythms and permit measurement of daylength to regulate seasonal adjustments (McClung 2019). Plant breeders have indirectly selected for variation at circadian-associated *loci* in many of the world's major crops when breeding to increase yield and improve crop performance by adapting genotypes to the local environment (Steed et al. 2021). Here we investigated whether variation at circadian-associated *loci* contributes to wheat yield traits in the field using a Multiparent Advanced Generation Inter-Cross (MAGIC) mapping population (Mackay *et al.* 2014). Wheat is a major crop, providing approximately 20% of the dietary calories and protein for the world's population (Hawkesford *et al.* 2013). Since the green revolution wheat yields have continued to increase but with the world's population estimated to reach 9.6 billion in 2050 (Gerland *et al.* 2014) it is likely that demand will outpace predicted yield increases (Hawkesford *et al.* 2013). Identification of novel genetic variation contributing to yield is critical to increasing the rate of yield gains in wheat.

Understanding of the nature of plant circadian oscillators is based primarily on investigation of Arabidopsis. In that model, the circadian oscillator contains a series of interlocking transcriptional feedback loops. The circadian oscillator cycle can be considered to start at dawn when expression of the transcriptional repressors *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) and *LATE ELONGATED HYPOCOTYL* (*LHY*) peaks, followed sequentially by peaks of expression of *PSEUDO REPSONSE REGULATOR 9* (*PRR9*)*, PRR7, PRR5, PRR3* and finally *PRR1* (also known as *TIMING OF CAB EXPRESSION 1* (*TOC1*)) at dusk (Hsu and Harmer 2014). PRR9

and PRR7 repress the expression of *CCA1* and *LHY* which releases repression of

evening phased genes including, *TOC1, GIGANTEA* (*GI*)*, EARLY FLOWERING 3*

(*ELF3*)*, ELF4* and *LUX ARRHYTHMO* (*LUX*) (Hsu and Harmer 2014). LUX is a MYB-

like transcription factor which associates with ELF3 and ELF4 to form the tripartite

evening complex (EC), which represses expression of *GI, TOC1, PRR7, PRR9* and

LUX (Dixon *et al.* 2011, Nusinow *et al.* 2011, Herrero *et al.* 2012)*.* Mutation of any of

the three EC components causes arrhythmia of the Arabidopsis circadian oscillator in

continuous light and temperature (Nagel and Kay 2012).

Orthologues of *ELF3* underlie *loci* regulating heading date in multiple crops, including

barley *early maturity8* (*eam8*) (Faure *et al.* 2012), rice *heading date17* (*Hd17*)

(Matsubara *et al.* 2012), *Earliness per se* (*Eps*)-A^m1 in *T. monococcum* and *Eps-D1* in

T. aestivum (Zikhali *et al.* 2016). In hexaploid wheat, the magnitude of the *Eps-D1*

effect on heading date increases with decreasing temperature (Ochagavía *et al.* 2019).

Orthologues of *TOC1* are associated with heading date, plant height and thousand

grain weight (Sun *et al.* 2020). An orthologue of *LUX* is a candidate for *Eps-3A^m* in *T.*

monococcum (Gawronski *et al.* 2014) and *HvLUX* is a candidate gene for *eam10* in

barley (Campoli *et al.* 2013). Furthermore, the *Photoperiod-1* (*Ppd-1*) genes*,* which

contribute to photoperiodic regulation of ear emergence, are orthologues of

Arabidopsis *PRRs* (Beales *et al.* 2007, Shaw *et al.* 2012, Bentley *et al.* 2013, Turner

et al. 2013).

We used the allelic diversity of a wheat MAGIC population to investigate the role of

circadian clock genes in regulating yield component traits in the field. The founder

varieties for the MAGIC population used in this study (Alchemy, Brompton, Claire,

Hereward, Rialto, Robigus, Soissons and Xi-19) are representative of UK and

European winter wheat varieties, capturing >80% of the polymorphism found across

UK wheat varieties from the last 70 years (Gardner *et al.* 2016). Our data suggest that

breeding for modified *ELF3* function can affect the regulation of flowering time but

potentially without compromising yield through disrupted circadian oscillations. We

find evidence of differences in the role for ELF3 in the circadian system between wheat

and Arabidopsis by concluding that in wheat, *ELF3* is expressed in the morning, rather

than the evening.

Results

Earliness per se *loci Eps-B1* and *Eps-D1* segregate in an eight-parent wheat MAGIC population

We performed QTL analysis on a set of replicated trial data for 784 recombinant inbred

lines (RILs) for the NIAB Elite MAGIC population from the 2012/2013 and 2013/2014

growing seasons at the NIAB experimental farms in Cambridge, UK. Growth stage 55

(GS55, equivalent to heading date) (Zadoks *et al.* 1974) was chosen for analysis and

QTL mapping to enable comparisons with previous studies (Griffiths *et al.* 2009). In

both 2013 and 2014, Soissons, which contains the *Ppd-D1a* (photoperiod insensitive)

allele, reached GS55 6-8 days earlier than any of the other MAGIC parents, which

contain the *Ppd-D1b* allele (photoperiod sensitive) (Fig 1). Heading date of the other

MAGIC parents varied over 5.5 days in 2013 and 7 days in 2014.

We performed the QTL analysis using MPWGAIM (Verbyla *et al.* 2014) for GS55

phenotypes in both years and tested whether any circadian clock genes co-locate

with the identified QTLs (< 2 cM to nearest 90 k marker or < 5 Mb to gene). Because

we are interested in the effects of circadian clock on yield, we report only QTLs

associated with circadian clock genes. We found that four circadian gene

orthologues (*TaELF3-B1, TaELF3-D1, Ppd-1, TaZTL-7A*) fell within 0.20, 0.27, 0.58

and 4.31 Mb respectively of one of the QTL flanking markers (Table 1). The absence

of significant heading date QTLs linked to vernalisation genes such as *VRN1*

confirmed that complete vernalisation had occurred, ensuring that observed

variations in heading date were true *Earliness per-se* (*Eps*) effects rather than a

variation in the vernalisation requirements between the MAGIC parents (Bentley *et*

al. 2013, Camargo *et al.* 2016). A previous study identified a single GS55 QTL in the

NIAB eight parent MAGIC population (Camargo *et al.* 2016) but this difference in

findings is likely a result of different growth conditions (greenhouse compared to

field), lower replicate number (208 RILs in pots compared to 784 RILs in field plots)

and the use of a different mapping methodology. Further validation of our analysis

pipeline and approaches are described in the methods and reported in (S1 Table).

A QTL on the distal end of chromosome 1B, delineated by the markers

Ra_c109187_371 and RAC875_c37075_283, mapped closely to the physical position

of *TaELF3-B1* (TraesCS1B02G477400) (Table 1). The 1B locus also had a

homoeologous QTL on chromosome 1D delineated by the markers Kukri_c29687_369

and Kukri_rep_c69829_307 identifying *TaELF3-D1* (TraesCS1D02G451200) as a

candidate gene (Table 1, Fig 2A). These QTLs co-locate with the previously described

heading date QTLs *Eps-B1* and *Eps-D1* (Griffiths *et al.* 2009). In our data, *Eps-B1* had

a significant effect in both years, explaining 3.4% of genetic variance in GS55 in 2013

and 2% in 2014 (Table 1). *Eps-D1* was also detected in both years but only had an

overall significant effect ($-\log10(p) = 2.32$) in 2014, accounting for 4.1% of genetic

variance in GS55 in that year (Table 1). The *Eps-B1* and *Eps-D1 loci* were chosen for

further in-depth analysis because of their proximity to *ELF3* homoeologues.

TaELF3-B1 is a candidate gene for the *Eps-B1* QTL

At the 1B QTL, the alleles derived from the MAGIC founders Soissons, Alchemy,

Robigus and Claire caused a relative reduction in the time to GS55, indicated by the

negative sign of the QTL, whilst alleles from Rialto, Hereward, Xi-19 and Brompton

caused a delay to GS55, indicated by the positive sign of the QTL (Table 2). The

interval between the markers Ra_c109187_371 (296.34cM / 685.7Mb) and

RAC875_c37075_283 (297.00cM / 687.7Mb) is 0.66cM, corresponding to a physical

length of ca. 2 Mb (Fig 2A).

The IWGSCv1.1 gene annotation suggests there are 26 high confidence gene models

within the interval of Ra_c109187_371 and RAC875_c37075_283. Based on current

gene annotation and orthology to cloned rice genes (Yao *et al.* 2018) there are no

novel candidates for the *Eps-B1* locus in addition to those described previously

(Faricelli *et al.* 2010, Alvarez *et al.* 2016, Zikhali *et al.* 2016). Compared to the *T.*

monococcum physical map of the *Eps-A^m1* region (Alvarez *et al.* 2016) *TmADK* (similar

to rice putative kinase *ADK1)* and *TmCK1-like* (*CASEIN KINASE I ISOFORM DELTA-*

LIKE) are not syntenic and the locus containing *TaELF3*, *TaGRIK1-like*

(*GEMINIVIRUS REP INTERACTING KINASE 1 LIKE*) and *TaPWWP1* (*PWWP*

domain-containing protein) is inverted (Fig 2A). Nineteen genes had homoeologues

on the syntenic 1D chromosome (Fig 2A).

Publicly available sequences for *TaELF3-B1* in Claire, Robigus (earlier heading

alleles) (Clavijo *et al.* 2017) and Rialto and Cadenza – the parents of Xi-19 - (later

heading alleles) (Zikhali *et al.* 2016) differed by one non-synonymous SNP within exon

4 at position chr1B:685645813. No further nonsynonymous SNPs were identified from

the 820k SNP dataset from CerealsDB (Wilkinson *et al.* 2012), which contains all

MAGIC parents.

The SNP (Adenine at position 2020 of CDS to Guanine; A2020G) causes a predicted

deleterious amino acid substitution from the ancestral serine to a glycine residue at

position 674 of the predicted primary amino acid sequence (Ser674Gly; SIFT 0.01)

(Fig 2B). A KASP marker was designed for this SNP and used to genotype the MAGIC

population.   The   KASP   marker   mapped   to   the   same   genetic   position   as

Ra_c109187_371 and segregated with the predicted founder effects, confirming that

TaELF3-B1 is part of the 1B GS55 QTL.

A predicted glycine at residue 674 of *TaELF3-B1*, associated with late heading, is

globally rare in wheat. In the 1000Exome data (He *et al.* 2019) 4.87% of lines have

the predicted glycine residue and in the wheat HapMap panel (Jordan *et al.* 2015),

only Rialto has the predicted glycine residue out of the 62 globally diverse lines

analysed (S2 Table). Similarly, of the 14 hexaploid varieties currently available from

the 10+ Wheat Genome project only three (Cadenza, Paragon, ArinaLrFor) possess

the non-ancestral glycine residue (S2 Table). However, analysis of multiple datasets

(see methods) shows the non-ancestral glycine674 is more common in wheat varieties

with a UK pedigree (9/13 varieties) (S3 Table). Alignment of *TaELF3* orthologues

shows that the predicted serine residue is also highly conserved across monocots (S1 Fig).

When looking for other potentially causal sequence variation we found that within the *ELF3-B1* promoter regions of Claire, Robigus, Paragon and Cadenza there are two sites polymorphic for the number of repeated cytosines ($(C)_n$) located 1967 bp and 1151 bp upstream of the transcriptional start site (TSS) (Fig 2B). As Robigus and Claire have an early heading phenotype at the *Eps-B1* locus and Cadenza a late heading phenotype (Alvarez *et al.* 2016) it is unlikely that the promoter $(C)_n$ polymorphisms cause the changes in heading dates because there is only a difference in two cytosines at the 1967 bp $(C)_n$ site between Cadenza and Robigus/Claire (Fig 2B). Furthermore *TaELF3-B1* on Julius, Lancer, Mace, Norin61 and Sy Mattis seem to be inverted compared to ArinaLrFor, CDC Landmark, CDC Stanley, Chinese Spring and Jagger. However, as none of these lines share the same haplotype with ArinaLrFor and Paragon (which both carry the Ser674Gly SNP) at *ELF3-B1* it is unlikely that an inversion is the cause behind the *Eps-B1* QTL. Thus, we conclude that the Ser674Gly SNP in *ELF3-B1* is the best candidate to underlie *Eps-B1*.

Subtelomeric introgression and inversion might underlie the *Eps-D1* heading date QTL

A subtelomeric deletion in Cadenza and Spark was previously described as the causal polymorphism underlying the *Eps-D1* locus using Spark × Rialto, Avalon × Cadenza mapping populations (Zikhali *et al.* 2016, Ochagavía *et al.* 2019). Our QTL analysis for 2012 (S1 Table) and 2014 (Table 1) shows a minor QTL overlapping this locus. In both years the Xi-19 founder genotype is associated with earlier flowering (2012: -1.26

days; p = 0.015; 2014; -0.79 days; p = 0.053). However correct genetic map

construction and resulting assignment of founder genotypes at this locus might be

more difficult given the high-density marker region previously identified at the end of

1D. These high-density marker regions were hypothesised to be associated with

introgressions (Gardner et al., 2016). Based on the previously reported deletion in

Cadenza and Spark (Zikhali, Wingen, and Griffiths 2016) and the pedigree of Xi-19

(Cadenza/Rialto//Cadenza) we hypothesised that the Cadenza allele inherited from

Tonic was a likely candidate for the observed QTL effects. By analysing data from

recently published wheat assemblies (Walkowiak *et al.* 2020) we suggest the

polymorphism is due to a subtelomeric inversion derived from introgressed material

rather than a deletion. A BLAST search of the Chinese Spring *TaELF3-D1* genomic

sequence against the 10+ Wheat Genome Project varieties identified high confidence

matches in all varieties including Cadenza. Interestingly, the sequence retrieved from

the variety Jagger was identical to Cadenza. However, the sequence identity of the

Jagger and Cadenza coding sequence for TraesJAG1D01G492900.1 and

TraesCAD_scaffold_026352_01G000400.1 compared to the Chinese Spring gene is

much lower (96.72%) than for the remaining 1D coding sequences (>99.9%). The

Jagger and Cadenza 1D coding sequences have a greater similarity to the Chinese

Spring *TaELF3-A1* (97.50%) and Chinese Spring *TaELF3-B1* (97.88%)

homoeologues than to Chinese Spring *TaELF3-D1*.

Visualisation of the 1D chromosome for the 10+ Wheat Genome Project varieties using

the Crop Haplotypes viewer (Brinton *et al.* 2020) showed a shared haplotype block at

the distal end of chromosome 1D between Cadenza (Haplotype viewer coordinates:

482.8 Mbp – 495.4 Mbp) and Jagger (480.0 Mbp to 493.5 Mbp) where *TaELF3-D1* is

found.

The Cadenza v1.1 reference sequence has yet to be assembled into chromosomes,

but since the Crop Haplotypes viewer (Brinton *et al.* 2020) data indicated identical by

state (IBS) haplotypes between Jagger and Cadenza, we compared the distal end

structure of the Jagger 1D chromosome with Chinese Spring. The end of the Jagger

chromosome contains an inversion of approximately 2.6 Mbp relative to Chinese

Spring which includes *TaELF3-D1* (Fig 3A). Alignment of the Jagger/Cadenza

TaELF3-D1 genomic sequence with Chinese Spring *TaELF3-D1* showed some

sequence differences including a deletion within intron 2 (Fig 3B, S4 Table). We

developed a subgenome specific PCR assay detailed in Fig 3B that amplifies the

region of *TaELF3-D1* containing intron 2 and confirmed that of the eight MAGIC

parents, only Xi-19 carried the Cadenza/Jagger *TaELF3-D1* allele (Fig 3C and D). As

hypothesised Spark and Tonic also carry the Cadenza *TaELF3-D1* allele (Fig 3C, D

and S2 Fig) as well as the varieties Maris Fundin and Cordiale. The intron 2 deletion

is also present in *Aegilops speltoides* and *Triticum timopheevii* (S3 Fig).

Given the low sequence conservation of the *TaELF3-D1* gene in Cadenza and Jagger

and the presence of the intronic deletion in *Aegilops speltoides* and *Triticum*

timopheevii, we investigated the wider phylogeny of the 1D Jagger/ Cadenza region

using the 820k SNP dataset from CerealsDB. This showed that Cadenza, Xi-19 and

Spark as well as several other Tonic descendants (KWS-Podium, Cordiale, Gallant,

Duxford) and five other cultivars (Moisson, Bacanora, Tuerkis, Badger, Pavon)

clustered separately from the remaining *T. aestivum* cultivars (S4 Fig). These lines

also cluster with six Watkins lines from India (Watkins34), Yugoslavia (Watkins352),

China (Watkins141), UK (Watkins103), Cyprus (Watkins292) and Turkey

(Watkins299). *T. timopheevii* clusters closely with the above lines. Gene alignment of

ELF-D1 orthologues and alignment of 300kb surrounding *ELF3-D1* also supports an

origin outside the D genome for the Jagger/ Cadenza *ELF3-D1* gene (S5 Fig, S5

Table).

Allelic variation in *TaELF3* had no detectable effect on circadian rhythms in the MAGIC founder varieties

We investigated if the variations at *TaELF3-B1* and *TaELF3-D1* in the MAGIC parents,

or other *loci,* might affect circadian rhythms in a manner that might explain the effects

on heading date. We quantified circadian rhythms in the MAGIC founder varieties by

measuring the period and amplitude of rhythms of delayed chlorophyll fluorescence

(DF) in otherwise constant light (Gould *et al.* 2009). We detected robust circadian

rhythms of DF in all the MAGIC parent lines (RAE < 0.5) (Fig 4A). Hereward had the

shortest circadian period (25.0 h) (Fig 4C) and Xi-19 the longest period (26.2 h) but

the difference between these periods and the global mean was not significant ($p =$

0.132) (Fig 4C). Thus, the allelic diversity in homoeologues of *TaELF3* did not cause

detectable changes in circadian rhythms.

Loss of function mutation of *TtELF3* disrupts wheat circadian clocks

296 Because we found that variation in the MAGIC parents at *TaELF3* was not correlated

297 with a detectable effect on circadian rhythms, we examined whether *ELF3* is a

circadian oscillator component in wheat. We obtained a tetraploid TILLIING mutant line carrying point mutations encoding stop codons in both *TtELF3-A1* and *TtELF3-B1* (Alvarez *et al.* 2016) herein designated *Ttelf3*-null (S6 Fig). We quantified circadian rhythms of "prompt" chlorophyll *a* fluorescence (PF) (Dakhiya *et al.* 2017) and leaf temperature under continuous light in both the *Ttelf3*-null and the respective *TtELF3*-WT obtained from the initial mutant TILLING line (Alvarez *et al.* 2016). We detected circadian rhythms in *TtELF3*-WT but *Ttelf3*-null was arrhythmic for all measures of PF (Fig 5C and S7 Fig) including $F_{m'}$ (*TtELF3*-WT RAE 0.36 ± 0.03; *Ttelf3*-null RAE 0.82 ± 0.08) (Fig 5C), leaf temperature (*TtELF3*-WT RAE 0.20 ± 0.04; *Ttelf3*-null RAE 0.56 ± 0.10) (Fig 5D-F) and DF (S8 Fig). The mean peak phase of relative leaf temperature for *TtELF3*-WT (Fig 5D) was 20.0 ± 0.2 h after subjective dawn, indicating, as expected, that stomata were most closed during the middle of the subjective night.

Loss of functional *TtELF3* perturbs circadian oscillator gene expression in light/dark cycles and constant light

To further investigate the function of *ELF3* in the wheat circadian system we measured the abundance of transcripts of orthologues of the Arabidopsis circadian oscillator components *LHY, TOC1, Ppd-1* (*PRR37*)*, PRR73, GI, ELF3* and *LUX* in light dark (LD) and continuous light (LL) cycles. In LD cycles all transcripts had rhythmic expression (Fig 6A-G). Consistent with previous findings in Arabidopsis, *TtLHY* had peak expression at dawn (Fig 6A), the *PRR* orthologues peaked during the photoperiod (Fig 6B-C) with *TtTOC1* (Fig 6D), *TtGI* (Fig 6E) and *TtLUX* (Fig 6F) peaking around dusk. However, the peak in *TtELF3* abundance was at the end of the night just prior to dawn (Fig 6G), approximately antiphase to the expression of *TtLUX* (Fig 6F). This differs markedly from Arabidopsis in which *ELF3* and *LUX* are co-

expressed near dusk (Nusinow *et al.* 2011, Huang *et al.* 2016, Ezer *et al.* 2017). For

all transcripts, except *Ppd-1* (Fig 6B), we detected oscillations in transcript abundance

in constant light, with the phasing in LL consistent with that in LD. These oscillations

of circadian transcript abundance were abolished, or of very low amplitude, in *Ttelf3-*

null (Fig 6A-G). There was a phase advance in peak transcript abundance of

approximately 3 h in LD in *Ttelf3*-null compared to *TtELF3-WT* (Fig 6A-G) except for

TtELF3 transcript abundance where the peak phase was at dawn in both *TtELF3*-WT

and *Ttelf3*-null (Fig 6G). The amplitude of transcript abundance under diel conditions

was also less in the *Ttelf3*-null line compared to *ELF3*-WT (Fig 6). Together, these

data demonstrate that functional *TtELF3* is required for the maintenance of robust

oscillations of circadian oscillator gene expression in LL and for normal circadian

timing in LD.

LUX loss of function disrupts the wheat circadian clock

The antiphase expression of wheat *ELF3* and *LUX* raises the possibility that the EC

might not be formed, or functions differently to that in Arabidopsis. Previous studies in

diploid wheat have identified *TmLUX* as a circadian oscillator gene (Gawronski *et al.*

2014), so we investigated whether *LUX* also contributes to circadian rhythms in

hexaploid wheat, and whether functionality of one homoeologue is sufficient to

maintain robust oscillations. We obtained three hexaploid Japanese cultivars,

Chogokuwase, Minaminokomugi and Geurumil that have variation at *TaLUX*

associated with flowering time (Mizuno *et al.* 2012). Chogokuwase is an extra early

flowering variety and is the offspring of Geurumil and Minaminokomugi. Chogokuwase

carries predicted non-functional mutations in all three sub-genome copies of *TaLUX*

(S9 Fig). Chogokuwase also carries a *TaVRN-D1* spring allele and a photoperiod

insensitive *Ppd-D1a* allele. Geurumil has the same predicted non-functional alleles of

TaLUX and *Ppd-D1a* but has a winter growth habit due to lacking the *VRN-D1* spring

allele. Minaminokomugi carries the non-functional *Talux-a1* allele but wild-type alleles

of *TaLUX-B1* and *TaLUX-D1* in addition to carrying the *TaVRN-D1* and *Ppd-D1a*

alleles (Mizuno *et al.* 2016).

Minaminokomugi maintained robust oscillations of chlorophyll fluorescence (RAE 0.16

± 0.01) and leaf temperature (RAE 0.26 ± 0.02) (Fig 7A-C). In contrast Chogokuwase,

which lacks functional copies of *TaLUX,* was arrhythmic for PF (RAE 0.56 ± 0.06) and

Geurumil, which is also lacking functional *TaLUX*, had reduced robustness of PF

oscillations (RAE 0.33 ± 0.05) compared to Minaminokomugi (Fig 7A-C). Similarly,

there was a decreased robustness of oscillations in leaf surface temperature for

Chogokuwase and Geurumil (RAE 0.48 ± 0.09 and RAE 0.46 ± 0.11) which appeared

to dampen to arrhythmia after the first two true LL cycles (Fig 7D-F). We conclude that

functional *TaLUX* is required for robust circadian rhythms in hexaploid wheat.

Discussion

Wheat *ELF3* contributes to heading date and circadian

rhythms

The eight MAGIC parent varieties provided the opportunity to measure the likely

contribution of the *ELF3* homoeologues to heading date and circadian rhythms. Of the

MAGIC parents Brompton, Hereward, Rialto and Xi-19 have a G|G substitution variant

of *TaELF3-B1* and in addition Xi-19 has an inverted copy of *TaELF3-D1,* likely arising

from a past introgression event. These variations in *TaELF3* alleles had no discernible

impact on circadian rhythms (Fig 4) but had measurable effects on heading date in the

field (S10 Fig).

Consistent with previous studies, we conclude that *ELF3* underlies *Eps-B1* and is a

candidate for *Eps-D1 loci* (Zikhali *et al.* 2016). We have extended the findings

concerning the *Eps-D1* locus by demonstrating that a copy of *ELF3-D1* is present in

Cadenza (as well as Xi-19, Spark, Cordiale, Tonic and Maris Fundin) but lies within an

introgression region that contains an inversion relative to the Chinese Spring D

genome (Fig 3). Based on analysis of the 820k SNP data we find that a number of

descendants of the cultivar Tonic carry the putative introgression, as do several other

unrelated cultivars and six Watkins lines from India, China, UK, Cyprus, Turkey and

Yugoslavia. The introgression is also likely present within Chinese elite germplasm

(Wang *et al.* 2016). As a result, we hypothesise that this region predates modern

breeding efforts. The same genomic region has been identified as being under

selection in a collection of bread wheat accessions dating from 1790 to 1930, though

in that study it was presumed to be a deletion rather than introgression (Przewieslik-

Allen *et al.* 2021). While the origin of the introgression is uncertain, the presence of

Jagger/Cadenza *TaELF3-D1* intronic deletion in *Ae. speltoides* and *T. timopheevii* (S3

Fig) and the clustering outside of *Aegilops tauschii* lines (S4 and S5 Figs) supports an

origin outside of the D genome, more closely related to the G or S genome. The

increase in quantity and quality of genomic resources for diverse accessions and

wheat relative species will help in determining the origin of this introgression.

Our data also identify *TaELF3-B1* as a candidate gene underlying the *Eps-B1* locus.

We found that a predicted glycine674 in *TaELF3-B1,* which segregates with *Eps-B1,*

caused a delay in heading and flowering compared to the ancestral serine674 variant

(S10 Fig), in agreement with previous observations in Avalon (Zikhali *et al.* 2016). It

was previously concluded that G|G is the ancestral variant (Zikhali *et al.* 2016) but our

evidence suggests that it is more likely that A|A is ancestral because the G|G allele is

globally rare, present in 4.87% of lines included in the 1000 Exome dataset, in one of

the 62 wheat HapMap panel lines (S2 Table) and in three of the 14 10+ Wheat

Genome project lines (S2 Table) and the ancestral serine residue is highly conserved,

present in all sequenced grasses. However, within UK wheat germplasm (as defined

by locality on wheat GRIS) the G|G allele is more common and is found in nine of the

13 UK lines we have investigated (S3 Table). The later flowering G|G allele may be

more favoured in northern regions (Langer *et al.* 2014).

We found that the allelic variation at *TaELF3* had no detectable effect on circadian

rhythms in the MAGIC founders. Epistasis between *ELF3* and *Ppd-D1* has previously

been reported for flowering time (Faure *et al.* 2012, Alvarez *et al.* 2016). However, we

find that variation at *Ppd-A1* has no obvious effects on circadian rhythms or

modification of circadian phenotypes in *Ttelf3* nulls (S11 Fig). We also find no evidence

of epistasis between *eps-B1* and *eps-D1* in regulation of circadian rhythms because

Xi-19 has both *eps-B1* and *eps-D1* but was not distinguishable from the rest of the

population in terms of circadian behaviour.

Only complete loss of functional *TtELF3* caused arrhythmia of PF (Fig 5A-C & S7 Fig),

leaf temperature (Fig 5D-F) and gene expression in LL (Fig 6), and this loss of function

caused advanced circadian phase in LD. This demonstrates that one role of *ELF3* is

to contribute to the correct alignment of internal phase to external phase, which is a

primary function of circadian oscillators. The earlier phasing of gene expression in LD

cycles is similar to barley containing the *eam8*.w allele where the peak phase of

HvCCA1, HvTOC1 and *HvGI* expression was also earlier (Faure *et al.* 2012). Similarly,

the same study reported loss of rhythmic oscillations in gene expression upon

transition to LL in barley. We found that functional *TtELF3* is required for high

amplitude *TtLHY* rhythms, and that *TtELF3* loss of function mutants have increased

trough transcript abundance of *TtGI* (Fig 6E). The similarity of these observations to

those in barley and rice lines carrying *ELF3* mutations (Zhao *et al.* 2012, Boden *et al.*

2014) suggests that *ELF3* genes might have similar functions across grasses.

An altered role for the evening complex (EC) in wheat

Binding of the Arabidopsis EC to target promoters requires LUX as both ELF3 and

ELF4 lack DNA binding domains, and it is thought this might explain the similar timing

of the transcript accumulation (Nusinow *et al.* 2011). However, in wheat the maximal

abundance of *ELF3* and *LUX* transcripts is separated by 12 hours (Fig 6F and G). This

difference in the peak time of *ELF3* and *LUX* transcript abundance also occurs in *T.*

monococcum (Alvarez *et al.* 2016) and rice (Zhao *et al.* 2012) and therefore may be a

general feature of cereal circadian oscillators. Wheat also lacks a clear orthologue of

AtELF4 and instead contains *ELF4-like* genes (Calixto *et al.* 2015). *ELF4-like* genes

might have a conserved role because expression of a barley *ELF4-like* gene restored

circadian rhythms to an Arabidopsis *elf4-2* mutant (Kolmos *et al.* 2009). There is

further evidence for conservation of EC function in monocots because Brachypodium

ELF3 binds Arabidopsis ELF4 and LUX, and can complement Arabidopsis *elf3-2*

mutants (Huang *et al.* 2017). The temporal separation of wheat *ELF3* and *LUX*

transcript abundance does not preclude the formation of an EC complex, but could

suggest one is not formed, or occurs at a different time in the diel cycle to Arabidopsis.

Whether through an EC or not, our data demonstrate that functional *ELF3* and *LUX*

are required for robust circadian rhythms in wheat (Figs 6- 8).

The difference in timing of *TaELF3* and *TaLUX* expression compared to their

Arabidopsis orthologues might arise as a consequence of promoter sequence

differences. Within 5000 bp upstream of the three *TaELF3* homoeologues transcription

start sites the most common motifs are linked to morning phased expression (37

morning, 20 evening and 7 midnight) (S6 Table). The general composition of the

TaELF3 promoters is similar to *TaLHY* (22 morning, 9 evening and 3 midnight) (S7

Table) but different to the evening expressed *TaLUX* promoters where the most

abundant motifs are associated with evening phased expression (32) compared to

morning (23) and midnight (12) (S8 Table). In particular*, TaELF3* promoters contain

Morning Element 1 (ME1) and Morning Element 2 (ME2) motifs, associated with

morning phased expression (Michael *et al.* 2008), which are also found in the promoter

region of the dawn expressed *TaLHY* homoeologues (S7 Table) but are absent in the

AtELF3 promoter (S6 Table). The *TaELF3* promoters also contain further morning-

related motifs such as the multiple Hormone up at dawn 1 (HUD1) and Hormone up

at dawn 2 (HUD2) elements (Michael *et al.* 2008) which are less frequent in the *AtELF3*

promoter (S6 Table).

Wheat *ELF3* might be a promising breeding target

ELF3 might be a suitable target for breeding new lines with heading date better

adapted to the local environment in the practice of chronoculture (Steed *et al.* 2021).

In other species such as pea, lentil and barley, allelic variation in *ELF3* has permitted

the cultivation of these crops at different latitudes (Weller *et al.* 2012, Zakhrabekova

et al. 2012, Lundqvist 2014). Whilst complete loss of functional *ELF3* in wheat

significantly advances flowering time, severely disrupts circadian rhythms and

decreases the spikelet number (Alvarez *et al.* 2016), we have demonstrated that

altering the function of particular homoeologues of *ELF3* can affect heading without a

measurable disruption to the circadian oscillator and the potential associated yield

penalties (Dodd *et al.* 2005) if some functional copies of *ELF3* are present. It might be

possible that by combining different alleles of *ELF3* with alleles of other circadian clock

genes, flowering time can be precisely tuned to suit the growing environment. This

could be tested by measuring flowering time in selected Cadenza TILLING lines

(Krasileva *et al.* 2017) containing mutations in other circadian clock genes as Cadenza

contains both the *TaELF3-B1* Ser674Gly SNP and the 1D introgression. In addition to

the *ELF3* alleles described in this work, we have also identified further *ELF3* alleles

including the absence of *ELF3-A1* in Julius (S9 Table) highlighting a range of *ELF3*

alleles within modern wheat cultivars which could be incorporated into breeding

programmes. The variation at *ELF3* might have occurred because of the positioning

of *ELF3* within the subtelomeric region (Aguilar and Prieto 2020) which has enhanced

levels of recombination which can lead to increased number of sequence and

structural variants. This has been observed for other regions, recent studies have

reported that a deletion at the distal end of chromosome 4AL is associated with altered

heat tolerance (Zhai *et al.* 2021) and a deletion at the distal end of 5AL encompasses

the *Rht12* locus which is associated with dwarfing (Sun *et al.* 2019). Secondly,

variation at *ELF3* may have been selected by past breeding efforts, leading to a

change in allele frequencies, due to advantageous adaptation of flowering time to local

climates similar to that seen for selection of *Ppd-1* alleles based on latitudinal cline

(Cockram *et al.* 2007) and altitude (Guo *et al.* 2020). A more systematic understanding

of circadian clock gene alleles and their combinations will allow for a more targeted

approach in optimising flowering time without impacting the functioning of the circadian

clock.

Materials and methods

Plant material

The eight MAGIC parents; Alchemy, Brompton, Claire, Hereward, Rialto, Robigus,

Soissons and Xi-19 were obtained from NIAB (National Institute of Agricultural Botany,

UK). The *Ttelf3*-null and *TtELF3*-WT lines were donated by Jorge Dubcovksy (Alvarez

et al. 2016) where the *Ttelf3*-null line is the progeny of bulked *elf3-null/Ppd-A1b* and

the *TtELF3*-WT line is progeny of bulked *ELF3WT/Ppd-A1b* (Table 4). The

Chogokuwase, Minaminokomugi and Geurumil lines were donated by Hidetaka

Nishida (Mizuno *et al.* 2012).

Plant growth conditions

Seeds were surface sterilised by washing once in 70% (v/v) ethanol for one minute

(Fischer Scientific, UK), once in autoclaved water for one minute, followed by 15

minutes in 10% (v/v) sodium hypochloride solution (Fischer Scientific, UK) and then

washed a further three times in autoclaved deionised water. Seeds were sown directly

onto 96 well modular trays (P G Horticulture Ltd, UK), containing a 1:1 mix of Levington

M2 potting compost (Levington, UK) and vermiculite (fine, Willian Sinclair Horticulture,

UK) and treated with the insecticide Intercept 70W (0.02 g L-1 soil, Bayer, Germany).

Plants were grown in growth chambers (Conviron, Canada) under long days and

constant temperature (16 h L: 8 h D; 200 μmol m^{-2} s^{-1}; 20 °C).

Field trials

In 2012/2013 a fully replicated yield trial of 784 F_5 MAGIC lines at the NIAB

experimental farm in Cambridge, UK was scored by NIAB for the timing of Growth

Stages on the Zadok's scale (Zadoks *et al.* 1974, Mackay *et al.* 2014). Each plot was

assessed every two days for its growth stage based on representative plants at the

centre of each plot. Once a plot reached GS55 (half of ear emerged above flag leaf

ligule) the date was logged. The phenotype was assessed with growth stage

measurements scored as days after sowing. In the 2013/2014 field season the same

growth stage phenotypes were collected from a fully replicated yield trial of 784 F_6

MAGIC lines.

Analysis of field phenotypes

Asreml-R 3.0 (Gilmour *et al.* 1997) was used to minimise or remove spatial effects in

phenotype data due to field variation.

Marker genotypes and their respective chromosomal groupings from (Gardner *et al.*

2016) were used. The genetic map was reordered visually using the R package

mpMapInteractive (Shah 2013) to improve ordering and consider physical ordering

information from IWGSCRefSeqv1. Recombination fractions were recalculated in

mpMap (Huang and George 2011) for each chromosome using mpestrf().

Recalculating recombination fractions on a per chromosome basis with the existing

marker groupings significantly reduced computation resources required and enables

the use of finer recombination bins (r <- c(0:5/600, 2:20/200, 11:50/100) compared to

the default values. Genetic map lengths were recalculated using the Kosambi mapping

function in computemap().

To reduce computational effort a skimmed genetic map with only unique mapping

locations was used to calculate the probability that each location on a genome was

inherited from each founder and conduct QTL mapping in MPWGAIM only (Verbyla *et*

al. 2014). A marker for each unique coordinate was chosen which had the lowest

missing calls and preferably a clear BLAST hit to the IWGSCRefSeqv1 reference

sequence, which was used to assign physical mapping coordinates. Out of 18089

mapped markers 4458 were chosen for founder probability calculations.

To test and validate our pipeline and analysis tools we used them to interrogate a data

set describing flowering time in a 2011/2012 nursery of the NIAB eight Parent MAGIC

population that had previously been analysed using Bayesian Networks (Scutari *et al.*

2014). Reanalysis of this data using MPWGAIM (Verbyla *et al.* 2014) detected 13

significant QTLs for flowering time with significant founder effects (S1 Table).

This analysis identified a major QTL (18.3% genetic variation explained, LOGP 18.33)

between the markers Kukri_c27309_590 (48.57 cM) and BS00064538_51 (56.64 cM),

defining a maximum mapping interval of 31.81 Mb – 34.23 Mb between the two

mapping bins as several markers map to the 56.64 cM bin. The mean founder effects

of the QTL predict a significant acceleration of flowering time for Soissons, which is

the only founder carrying the photoperiod insensitive *Ppd-D1a* allele (Bentley *et al.*

2013) and furthermore *Ppd-D1* maps within the QTL (33.95 Mb). This analysis of the

effect of *Ppd-D1a* on flowering time validated MPWGAIM and our QTL mapping

pipeline in MAGIC.

Synteny analysis between Chinese Spring and Jagger 1D

A conserved haplotype block between Cadenza and Jagger at the distal end of

chromosome 1D was identified using the Crop Haplotypes web browser (Crop

Haplotypes (crop-haplotypes.com) (Brinton *et al.* 2020). Physical locations of genes

on the Chinese Spring 1D chromosome (IWGSCRefSeqv1.1) and Jagger 1D

chromosome (PGSBv2.0) were mapped using the mapviewer v2.6.1

(https://10wheatgenomes.plantinformatics.io/mapview) and then confirmed manually

using EnsemblPlants.

SNP mining and analysis

820k and 35k SNP array data was obtained from the CerealsDB website (Wilkinson

et al. 2012) and the 1000Exome data from the 1000 Exomes project website

(http://wheatgenomics.plantpath.ksu.edu/1000EC/). Mapping of 35k and 820k SNPs

to the IWGSCv1.1 Ref sequence were obtained from EnsemblPlantsv51. SNPs for the

Cadenza/Jagger haplotype (chr1D: 482.3Mb – end of chromosome 1D) were

extracted and heterozygous calls set as missing. Distance matrices and phylogenetic

trees were constructed using the R package "ape" (Paradis and Schliep 2019) 100

bootstrap permutations were run also using the R package "ape".

To investigate the prevalence of the Ser674Gly SNP in a broad range of germplasm

we analysed variation data from multiple sources including the 820k SNP genotyping

array (Winfield *et al.* 2016), the wheat HapMap panel (Jordan *et al.* 2015), the 1000

wheat Exome project (He *et al.* 2019) and the 10+ Wheat Genome project (Walkowiak

et al. 2020). We found that the Ser674Gly SNP was also present in the HapMap panel

as chr1B_685645813 and within the 1000 Exome data as scaffold48561_337271.

Phylogenetic tree construction

ELF3-D1 orthologous groups were retrieved from public repositories. Coding

sequences were aligned using the MCoffee aligner on default settings

(https://pubmed.ncbi.nlm.nih.gov/16556910/). Maximum Likelihood (ML) Phylogenetic

trees were build using CLC Genomics Workbench 21.0.3. The best substitution model

was determined using the Model testing function before running Molecular

Phylogenetic analysis by Maximum Likelihood method with 100 permutations for

Bootstrap values. The *ELF3* gene from *Brachypodium distachyon* was used as an

outgroup.

Quantitative Real-Time PCR

The *TtELF3*-WT and *Ttelf3*-null lines were sown and grown in two antiphased LED

growth chambers (Conviron) under long day conditions (16h L: 8h D; 250 µmol m^{-2} s^{-1}

, 20°C day: 16°C night). Sampling commenced 16 days post-sowing, the first true

leaf was sampled every 3 hours for 96 hours. After the first 24 hours of sampling the

cabinet was switched to constant light and temperature (250 µmol m^{-2} s^{-1} , 20°C). Total

RNA was extracted from leaf samples using the KingFisher Flex purification system

(Thermofisher scientific) in conjunction with the Maxwell® HT simplyRNA Kit

(Promega). RNA concentration was determined using the Nanodrop ND-1000. DNA

contamination was removed using the Arcticzyme HL-DNAse kit. RNA integrity was

checked using the Fragment Analyzer™ (AATI, USA) before cDNA was synthesised

from extracted RNA using the High-Capacity cDNA reverse transcription kit (Applied

Biosystems). Three technical replicates of gene specific products were amplified in 10

µl reactions using Power SYBR Green Master mix on a CFX384 Touch Real-Time

PCR detection system. Gene expression levels were determined relative to the

expression of two housekeeping genes, *TtRP15* and *TtRPT5A,* that were selected

using the GeNorm alogorithm (Vandesompele *et al.* 2002) (Table 5).

PCR of *ELF3-D1*

DNA extracted using the DNeasy plant mini extraction kit (Qiagen, UK). The presence

of *ELF3-D1* was confirmed by PCR using the *TaELF3-D1* primers listed in Table 5.

PCR was performed using the DreamTaq Green PCR kit (Thermo Scientific, UK) on

a ProFlex PCR system (Thermo Scientific, UK). PCR reaction conditions were as

follows: 3 minutes at 95ºC; 40 cycles of 30 s at 95ºC, 30 s at 68ºC, 90 s at 72ºC; 15

minutes at 72ºC.

Delayed chlorophyll fluorescence and prompt chlorophyll *a* fluorescence imaging

Leaf samples were taken from the upper half of the first true leaf of 16 day old

seedlings (non-vernalised) and were cut into 3 mm by 3 mm pieces and placed on

solid 0.8% (w/v) bactoagar, 0.5 x MS Vitamin medium (M3900, Sigma-Aldrich, UK)

supplemented with 0.5 µM benzyl-aminopurine (BAP), poured in a black 96 well

imaging plates (Greiner, UK). Delayed fluorescence (Gould *et al.* 2009) was measured

for 60 s in an LB985 Nightshade (Berthold, UK) camera. Between measurements of

DF light was supplied from blue (470 nm) red (660 nm) and LED arrays at 100 μmol

m^{-2} s^{-1} during the experiment. Temperature was maintained at 20°C.

Chlorophyll *a* fluorescence and calculations of parameters were carried out using a

CFimager (Technologica Ltd, Colchester, UK) and the image processing scripts

provided by the manufacturer. Chlorophyll fluorescence images were captured using

a Stingray F145B ASG camera (Allied Vision Technologies, UK) through an RG665

long pass filter to exclude blue light from the LEDs. When imaging under continuous

light, leaf fragments were exposed to 40 minutes of 100 μmol m^{-2} s^{-1} of blue light

followed by a saturating pulse of 6172 μmol m^{-2} s^{-1} blue light for 800 ms then 20

minutes of darkness (with non-actinic measuring light on) followed by a second

saturating pulse of 6172 μmol m^{-2} s^{-1} for 800 ms.

Circadian measures of amplitude, phase, period and relative amplitude error were

calculated using the FFT-NLLS method in Biodare2 (www.biodare2.ed.ac.uk)

(Zielinski *et al.* 2014). Where necessary data was normalised to between -1 and 1

using the normalisation feature of Biodare2.

Direct contact leaf temperature measurements

Leaf surface temperature relative to background temperature was measured using an

in-house built device and provides an alternative measure of circadian rhythms caused

by circadian regulated changes in the stomatal aperture (Dakhiya and Green 2019).

The temperature logger was validated by measuring circadian rhythms of "in-phase"

and "anti-phase" wheat leaves where peak temperature, and therefore maximum

closure, occurred during the subjective night (S12 Fig). Leaf temperature

measurements were taken from fully expanded flag leaf of 4-week-old plants under

continuous light and temperature (200 µmol m^{-2} s^{-1}, 20°C) that were previously grown

under long days (16h L: 8h D; 200 µmol m^{-2} s^{-1}, 20°C day: 15°C night) in a growth

cabinet (Conviron, UK).

Wheat was grown in 10 cm pots. Circadian measurements of leaf temperature were

taken when the pants were four-weeks old. Type-K thermocouples were attached to

the leaf such the end of the thermocouple was pressing onto the surface of the fully

expanded flag leaf. This was kept in position by positioning a paperclip onto the leaf

and fixing the paperclip to a stake. The thermocouples were connected to an Arduino

Mega using the Adafruit 1778, a 10 bit D-A converter and x20 amplifier giving a

resolution of ± 0.1°C with a DS3231 RTC added to keep time, temperature was logged

every 30 s. Background temperature was measured using unattached thermocouples.

Leaf temperatures were averaged using a 60-point moving average, background

temperature was then subtracted from leaf temperature to give leaf temperature

relative to background.

Bibliography

Aguilar, M. and Prieto, P., 2020. Sequence analysis of wheat subtelomeres reveals a high polymorphism among homoeologous chromosomes. *The Plant Genome*, 13 (3), e20065.

Alvarez, M.A., Tranquilli, G., Lewis, S., Kippes, N., and Dubcovsky, J., 2016. Genetic and physical mapping of the earliness *per se* locus *Eps-A^m1* in *Triticum monococcum* identifies *EARLY FLOWERING 3* (*ELF3*) as a candidate gene. *Functional & Integrative Genomics*, 16 (4), 365–82.

Beales, J., Turner, A., Griffiths, S., Snape, J.W., and Laurie, D.A., 2007. A *Pseudo-Response Regulator* is misexpressed in the photoperiod insensitive *Ppd-D1a* mutant of wheat (*Triticum aestivum L.*). *Theoretical and Applied Genetics*, 115 (5), 721–733.

Bentley, A.R., Horsnell, R., Werner, C.P., Turner, A.S., Rose, G.A., Bedard, C., Howell, P., Wilhelm, E.P., Mackay, I.J., Howells, R.M., Greenland, A., Laurie, D.A., and Gosman, N., 2013. Short, natural, and extended photoperiod response in BC2F4 lines of bread wheat with different *Photoperiod-1* (*Ppd-1*) alleles. *Journal of Experimental Botany*, 64 (7), 1783–1793.

Boden, S.A., Weiss, D., Ross, J.J., Davies, N.W., Trevaskis, B., Chandler, P.M., and Swain, S.M., 2014. *EARLY FLOWERING3* Regulates Flowering in Spring Barley by Mediating Gibberellin Production and *FLOWERING LOCUS T* Expression. *The Plant Cell*, 26 (4), 1557–1569.

Brinton, J., Ramirez-Gonzalez, R.H., Simmonds, J., Wingen, L., Orford, S., Griffiths, S., Haberer, G., Spannagl, M., Walkowiak, S., Pozniak, C., and Uauy, C., 2020. A haplotype-led approach to increase the precision of wheat breeding. *Communications Biology*, 3 (1), 1–11.

Calixto, C.P.G., Waugh, R., and Brown, J.W.S., 2015. Evolutionary Relationships

Among Barley and Arabidopsis Core Circadian Clock and Clock-Associated

Genes. *Journal of Molecular Evolution*, 80 (2), 108–119.

Camargo, A. V., Mott, R., Gardner, K.A., Mackay, I.J., Corke, F., Doonan, J.H., Kim,

J.T., and Bentley, A.R., 2016. Determining Phenological Patterns Associated

with the Onset of Senescence in a Wheat MAGIC Mapping Population. *Frontiers*

in Plant Science, 7, 1540.

Campoli, C., Pankin, A., Drosse, B., Casao, C.M., Davis, S.J., and Von Korff, M.,

2013. *HvLUX1* is a candidate gene underlying the *early maturity 10* locus in

barley: Phylogeny, diversity, and interactions with the circadian clock and

photoperiodic pathways. *New Phytologist*, 199 (4), 1045–1059.

Clavijo, B.J., Venturini, L., Schudoma, C., Accinelli, G.G., Kaithakottil, G., Wright, J.,

Borrill, P., Kettleborough, G., Heavens, D., Chapman, H., Lipscombe, J., Barker,

T., Lu, F.H., McKenzie, N., Raats, D., Ramirez-Gonzalez, R.H., Coince, A.,

Peel, N., Percival-Alwyn, L., Duncan, O., Trösch, J., Yu, G., Bolser, D.M.,

Namaati, G., Kerhornou, A., Spannagl, M., Gundlach, H., Haberer, G., Davey,

R.P., Fosker, C., Di Palma, F., Phillips, A.L., Millar, A.H., Kersey, P.J., Uauy, C.,

Krasileva, K. V., Swarbreck, D., Bevan, M.W., and Clark, M.D., 2017. An

improved assembly and annotation of the allohexaploid wheat genome identifies

complete families of agronomic genes and provides genomic evidence for

chromosomal translocations. *Genome Research*, 27 (5), 885–896.

Cockram, J., Jones, H., Leigh, F.J., O'Sullivan, D., Powell, W., Laurie, D.A., and

Greenland, A.J., 2007. Control of flowering time in temperate cereals: genes,

domestication, and sustainable productivity. *Journal of Experimental Botany*, 58

(6), 1231–1244.

Dakhiya, Y. and Green, R.M., 2019. Thermal imaging as a non-invasive technique

for analyzing circadian rhythms in plants. *New Phytologist*, 224, 1685–1696.

Dakhiya, Y., Hussien, D., Fridman, E., Kiflawi, M., and Green, R., 2017. Correlations

between Circadian Rhythms and Growth in Challenging Environments. *Plant*

Physiology, 173 (3), 1724–1734.

Dixon, L.E., Knox, K., Kozma-Bognar, L., Southern, M.M., Pokhilko, A., and Millar,

A.J., 2011. Temporal repression of core circadian genes is mediated through

EARLY FLOWERING 3 in Arabidopsis. *Current Biology*, 21 (2), 120–125.

Dodd, A.N., Salathia, N., Hall, A., Kévei, E., Tóth, R., Nagy, F., Hibberd, J.M., Millar,

A.J., and Webb, A. a R., 2005. Plant circadian clocks increase photosynthesis,

growth, survival, and competitive advantage. *Science (New York, N.Y.)*, 309

(5734), 630–633.

Ezer, D., Jung, J.H., Lan, H., Biswas, S., Gregoire, L., Box, M.S., Charoensawan, V.,

Cortijo, S., Lai, X., Stöckle, D., Zubieta, C., Jaeger, K.E., and Wigge, P.A., 2017.

The evening complex coordinates environmental and endogenous signals in

Arabidopsis. *Nature Plants*, 3 (7).

Faricelli, M.E., Valárik, M., and Dubcovsky, J., 2010. Control of flowering time and

spike development in cereals: The *earliness per se Eps-1* region in wheat, rice,

and *Brachypodium. Functional and Integrative Genomics*, 10 (2), 293–306.

Faure, S., Turner, a. S., Gruszka, D., Christodoulou, V., Davis, S.J., von Korff, M.,

and Laurie, D. a., 2012. Mutation at the circadian clock gene *EARLY*

MATURITY 8 adapts domesticated barley (*Hordeum vulgare*) to short growing

seasons. *Proceedings of the National Academy of Sciences*, 109 (21), 8328–

8333.

Gardner, K.A., Wittern, L.M., and Mackay, I.J., 2016. A highly recombined, high-

density, eight-founder wheat MAGIC map reveals extensive segregation

distortion and genomic locations of introgression segments. *Plant Biotechnology*

Journal, 14 (6), 1406–1417.

Gawronski, P., Ariyadasa, R., Himmelbach, A., Poursarebani, N., Kilian, B., Stein,

N., Steuernagel, B., Hensel, G., Kumlehn, J., Sehgal, S.K., Gill, B.S., Gould, P.,

Hall, A., and Schnurbusch, T., 2014. A Distorted Circadian Clock Causes Early

Flowering and Temperature-Dependent Variation in Spike Development in the

Eps-3^{A}m Mutant of Einkorn Wheat. *Genetics*, 196 (4), 1253–1261.

Gendron, J.M., Pruneda-Paz, J.L., Doherty, C.J., Gross, A.M., Kang, S.E., and Kay,

S.A., 2012. Arabidopsis circadian clock protein, TOC1, is a DNA-binding

transcription factor. *Proceedings of the National Academy of Sciences*, 109 (8),

3167–3172.

Gerland, P., Raftery, A.E., Sevcikova, H., Li, N., Gu, D., Spoorenberg, T., Alkema,

L., Fosdick, B.K., Chunn, J., Lalic, N., Bay, G., Buettner, T., Heilig, G.K., and

Wilmoth, J., 2014. World population stabilization unlikely this century. *Science*,

346 (6206), 234–237.

Gilmour, A.R., Cullis, B.R., and Verbyla, A.P., 1997. Accounting for Natural and

Extraneous Variation in the Analysis of Field Experiments. *Journal of*

Agricultural, Biological, and Environmental Statistics, 2 (3), 269–293.

Gould, P.D., Diaz, P., Hogben, C., Kusakina, J., Salem, R., Hartwell, J., and Hall, A.,

2009. Delayed fluorescence as a universal tool for the measurement of

circadian rhythms in higher plants. *Plant Journal*, 58 (5), 893–901.

Griffiths, S., Simmonds, J., Leverington, M., Wang, Y., Fish, L., Sayers, L., Alibert,

L., Orford, S., Wingen, L., Herry, L., Faure, S., Laurie, D., Bilham, L., and

Snape, J., 2009. Meta-QTL analysis of the genetic control of ear emergence in

793  elite European winter wheat germplasm. *Theoretical and Applied Genetics*, 119

794  (3), 383–395.

795 Guo, W., Xin, M., Wang, Z., Yao, Y., Hu, Z., Song, W., Yu, K., Chen, Y., Wang, X.,

796  Guan, P., Appels, R., Peng, H., Ni, Z., and Sun, Q., 2020. Origin and adaptation

797  to high altitude of Tibetan semi-wild wheat. *Nature Communications*, 11 (1), 1–

798  12.

799 Hawkesford, M.J., Araus, J.L., Park, R., Calderini, D., Miralles, D., Shen, T., Zhang,

800  J., and Parry, M. a. J., 2013. Prospects of Doubling Global Wheat Yields. *Food*

801  *and Energy Security*, 2 (1), 34–48.

802 He, F., Pasam, R., Shi, F., Kant, S., Keeble-Gagnère, G., Kay, P., Forrest, K., Fritz,

803  A., Hucl, P., Wiebe, K., Know, R., Cuthbert, R., Pozniak, C., Akhunova, A.,

804  Morrell, P.L., Davies, J.P., Webb, S.R., Spangenberg, G., Hayes, B., Daetwyler,

805  H., Tibbits, J., Hayden, M., and Akhunov, E., 2019. Exome sequencing

806  highlights the role of wild-relative introgression in shaping the adaptive

807  landscape of the wheat genome. *Nature genetics*, 51 (5), 896–904.

808 Herrero, E., Kolmos, E., Bujdoso, N., Yuan, Y., Wang, M., Berns, M.C., Uhlworm, H.,

809  Coupland, G., Saini, R., Jaskolski, M., Webb, A., Goncalves, J., and Davis, S.J.,

810  2012. EARLY FLOWERING4 Recruitment of EARLY FLOWERING3 in the

811  Nucleus Sustains the *Arabidopsis* Circadian Clock. *the Plant Cell Online*, 24 (2),

812  428–443.

813 Hsu, P.Y. and Harmer, S.L., 2014. Wheels within wheels: The plant circadian

814  system. *Trends in Plant Science*, 19 (4), 240–249.

815 Huang, B.E. and George, A.W., 2011. R/mpMap: A computational platform for the

816  genetic analysis of multiparent recombinant inbred lines. *Bioinformatics*, 27 (5),

817  727–729.

Huang, H., Alvarez, S., Bindbeutel, R., Shen, Z., Naldrett, M.J., Evans, B.S., Briggs,

S.P., Hicks, L.M., Kay, S.A., and Nusinow, D.A., 2016. Identification of evening

complex associated proteins in *Arabidopsis* by affinity purification and mass

spectrometry. *Molecular and Cellular Proteomics*, 15 (1), 201–217.

Huang, H., Gehan, M.A., Huss, S.E., Alvarez, S., Lizarraga, C., Gruebbling, E.L.,

Gierer, J., Naldrett, M.J., Bindbeutel, R.K., Evans, B.S., Mockler, T.C., and

Nusinow, D.A., 2017. Cross-species complementation reveals conserved

functions for *EARLY FLOWERING 3* between monocots and dicots. *Plant*

Direct, 00, 1–14.

Huang, W., Perez-Garcia, P., Pokhilko, A., Millar, A.J., Antoshechkin, I., Riechmann,

J.L., and Mas, P., 2012. Mapping the Core of the *Arabidopsis* Circadian Clock

Defines the Network Structure of the Oscillator. *Science*, 336 (6077), 75–79.

Jordan, K.W., Wang, S., Lun, Y., Gardiner, L.-J., MacLachlan, R., Hucl, P., Wiebe,

K., Wong, D., Forrest, K.L., Sharpe, A.G., Sidebottom, C.H., Hall, N., Toomajian,

C., Close, T., Dubcovsky, J., Akhunova, A., Talbert, L., Bansal, U.K., Bariana,

H.S., Hayden, M.J., Pozniak, C., Jeddeloh, J.A., Hall, A., and Akhunov, E.,

2015. A haplotype map of allohexaploid wheat reveals distinct patterns of

selection on homoeologous genomes. *Genome Biology*, 16 (1), 48.

Kolmos, E., Nowak, M., Werner, M., Fischer, K., Schwarz, G., Mathews, S., Schoof,

H., Nagy, F., Bujnicki, J.M., and Davis, S.J., 2009. Integrating *ELF4* into the

circadian system through combined structural and functional studies. *HFSP*

Journal, 3 (5), 350–366.

Krasileva, K. V, Vasquez-gross, H.A., Howell, T., Bailey, P., Paraiso, F., and

Clissold, L., 2017. Uncovering hidden variation in polyploid wheat. *PNAS*, E913–

E921.

Langer, S.M., Longin, C.F.H., and Würschum, T., 2014. Flowering time control in

European winter wheat. *Frontiers in Plant Science*, 5 (OCT), 1–11.

Lundqvist, U., 2014. Scandinavian mutation research in barley - a historical review.

Hereditas, 151 (6), 123–131.

Mackay, I.J., Bansept-Basler, P., Bentley, A.R., Cockram, J., Gosman, N.,

Greenland, A.J., Horsnell, R., Howells, R., O'Sullivan, D.M., Rose, G.A., and

Howell, P.J., 2014. An eight-parent multiparent advanced generation inter-cross

population for winter-sown wheat: Creation, properties, and validation. *G3:*

Genes, Genomes, Genetics, 4 (9), 1603–1610.

Matsubara, K., Ogiso-Tanaka, E., Hori, K., Ebana, K., Ando, T., and Yano, M., 2012.

Natural variation in *Hd17*, a homolog of arabidopsis *ELF3* that is involved in rice

photoperiodic flowering. *Plant and Cell Physiology*, 53 (4), 709–716.

McClung, C., 2019. The Plant Circadian Oscillator. *Biology*, 8 (1), 14.

Michael, T.P., Mockler, T.C., Breton, G., McEntee, C., Byer, A., Trout, J.D., Hazen,

S.P., Shen, R., Priest, H.D., Sullivan, C.M., Givan, S.A., Yanovsky, M., Hong, F.,

Kay, S.A., and Chory, J., 2008. Network discovery pipeline elucidates conserved

time-of-day-specific cis-regulatory modules. *PLoS Genetics*, 4 (2), 14.

Mizuno, N., Kinoshita, M., Kinoshita, S., Nishida, H., Fujita, M., Kato, K., Murai, K.,

and Nasuda, S., 2016. Loss-of-function mutations in three homoeologous

PHYTOCLOCK 1 genes in common wheat are associated with the extra-early

flowering phenotype. *PLoS ONE*, 11 (10), 1–19.

Mizuno, N., Nitta, M., Sato, K., and Nasuda, S., 2012. A wheat homologue of

PHYTOCLOCK 1 is a candidate gene conferring the early heading phenotype to

einkorn wheat. *Genes & genetic systems*, 87 (6), 357–67.

Nagel, D.H. and Kay, S.A., 2012. Complexity in the wiring and regulation of plant

circadian networks. *Current Biology*, 22 (16), 648–657.

Nusinow, D.A., Helfer, A., Hamilton, E.E., King, J.J., Imaizumi, T., Schultz, T.F.,

Farré, E.M., and Kay, S.A., 2011. The ELF4–ELF3–LUX complex links the

circadian clock to diurnal control of hypocotyl growth. *Nature*, 475 (7356), 398–

402.

Ochagavía, H., Prieto, P., Zikhali, M., Griffiths, S., and Slafer, G.A., 2019. Earliness

Per Se by Temperature Interaction on Wheat Development. *Scientific Reports*, 9

(1), 2584.

Paradis, E. and Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics

and evolutionary analyses in R. *Bioinformatics*, 35 (3), 526–528.

Przewieslik-Allen, A.M., Wilkinson, P.A., Burridge, A.J., Winfield, M.O., Dai, X.,

Beaumont, M., King, J., Yang, C., Griffiths, S., Wingen, L.U., Horsnell, R.,

Bentley, A.R., Shewry, P., Barker, G.L.A., and Edwards, K.J., 2021. The role of

gene flow and chromosomal instability in shaping the bread wheat genome.

Nature Plants, 7 (2), 172–183.

Scutari, M., Howell, P., Balding, D.J., and Mackay, I., 2014. Multiple Quantitative

Trait Analysis Using Bayesian Networks. *Genetics*, 198 (1), 129–137.

Shah, R., 2013. mpMapinteractive: Interactive components for package mpMap. R

package version 0.2.

Shaw, L.M., Turner, A.S., and Laurie, D.A., 2012. The impact of photoperiod

insensitive *Ppd-1a* mutations on the photoperiod pathway across the three

genomes of hexaploid wheat (*Triticum aestivum*). *The Plant Journal*, 71 (1), 71–

84.

Steed, G., Cano-Ramirez, D.L., Hannah, M.A., and Webb, A.A., 2021.

Chronoculture, harnessing the circadian clock to improve crop yield and

sustainability. *Science*, 372 (April).

Sun, H., Zhang, W., Wu, Y., Gao, L., Cui, F., Zhao, C., Guo, Z., and Jia, J., 2020.

The Circadian Clock Gene, *TaPRR1*, Is Associated With Yield-Related Traits in

Wheat (*Triticum aestivum L.*). *Frontiers in Plant Science*, 11, 285.

Sun, L., Yang, W., Li, Y., Shan, Q., Ye, X., Wang, D., Yu, K., Lu, W., Xin, P., Pei, Z.,

Guo, X., Liu, D., Sun, J., Zhan, K., Chu, J., and Zhang, A., 2019. A wheat

dominant dwarfing line with *Rht12* , which reduces stem cell length and affects

gibberellic acid synthesis, is a 5AL terminal deletion line. *The Plant Journal*, 97

(5), 887–900.

Turner, A.S., Faure, S., Zhang, Y., and Laurie, D.A., 2013. The effect of day-neutral

mutations in barley and wheat on the interaction between photoperiod and

vernalization. *Theoretical and Applied Genetics*, 126 (9), 2267–2277.

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A.,

and Speleman, F., 2002. Accurate normalization of real-time quantitative RT-

PCR data by geometric averaging of multiple internal control genes. *Genome*

biology, 3 (7), RESEARCH0034.

Verbyla, A.P., George, A.W., Cavanagh, C.R., and Verbyla, K.L., 2014.

Whole-genome QTL analysis for MAGIC. *Theoretical and Applied Genetics*, 127

(8), 1753–1770.

Walkowiak, S., Gao, L., Monat, C., Haberer, G., Kassa, M.T., Brinton, J., Ramirez-

Gonzalez, R.H., Kolodziej, M.C., Delorean, E., Thambugala, D., Klymiuk, V.,

Byrns, B., Gundlach, H., Bandi, V., Siri, J.N., Nilsen, K., Aquino, C.,

Himmelbach, A., Copetti, D., Ban, T., Venturini, L., Bevan, M., Clavijo, B., Koo,

D.H., Ens, J., Wiebe, K., N'Diaye, A., Fritz, A.K., Gutwin, C., Fiebig, A., Fosker,

C., Fu, B.X., Accinelli, G.G., Gardner, K.A., Fradgley, N., Gutierrez-Gonzalez, J.,

Halstead-Nussloch, G., Hatakeyama, M., Koh, C.S., Deek, J., Costamagna,

A.C., Fobert, P., Heavens, D., Kanamori, H., Kawaura, K., Kobayashi, F.,

Krasileva, K., Kuo, T., McKenzie, N., Murata, K., Nabeka, Y., Paape, T.,

Padmarasu, S., Percival-Alwyn, L., Kagale, S., Scholz, U., Sese, J., Juliana, P.,

Singh, R., Shimizu-Inatsugi, R., Swarbreck, D., Cockram, J., Budak, H.,

Tameshige, T., Tanaka, T., Tsuji, H., Wright, J., Wu, J., Steuernagel, B., Small,

I., Cloutier, S., Keeble-Gagnère, G., Muehlbauer, G., Tibbets, J., Nasuda, S.,

Melonek, J., Hucl, P.J., Sharpe, A.G., Clark, M., Legg, E., Bharti, A., Langridge,

P., Hall, A., Uauy, C., Mascher, M., Krattinger, S.G., Handa, H., Shimizu, K.K.,

Distelfeld, A., Chalmers, K., Keller, B., Mayer, K.F.X., Poland, J., Stein, N.,

McCartney, C.A., Spannagl, M., Wicker, T., and Pozniak, C.J., 2020. Multiple

wheat genomes reveal global variation in modern breeding. *Nature*, 588 (7837),

277–283.

Wang, J., Wen, W., Hanif, M., Xia, X., Wang, H., Liu, S., Liu, J., Yang, L., Cao, S.,

and He, Z., 2016. *TaELF3-1DL*, a homolog of *ELF3*, is associated with heading

date in bread wheat. *Molecular Breeding*, 36 (12), 1–9.

Wang, Z.Y., Kenigsbuch, D., Sun, L., Harel, E., Ong, M.S., and Tobin, E.M., 1997. A

Myb-related transcription factor is involved in the phytochrome regulation of an

Arabidopsis Lhcb gene. *The Plant Cell*, 9 (4), 491–507.

Weller, J.L., Liew, L.C., Hecht, V.F.G., Rajandran, V., Laurie, R.E., Ridge, S.,

Wenden, B., Schoor, J.K.V., Jaminon, O., Blassiau, C., Dalmais, M., Rameau,

C., Bendahmane, A., Macknight, R.C., and Lejeune-Hénaut, I., 2012. A

conserved molecular basis for photoperiod adaptation in two temperate

legumes. *Proceedings of the National Academy of Sciences of the United States*

of America, 109 (51), 21158–21163.

Wilkinson, P.A., Winfield, M.O., Barker, G.L.A., Allen, A.M., Burridge, A., Coghill,

J.A., and Edwards, K.J., 2012. CerealsDB 2.0: an integrated resource for plant

breeders and scientists. *BMC Bioinformatics*, 13 (1), 219.

Winfield, M.O., Allen, A.M., Burridge, A.J., Barker, G.L.A., Benbow, H.R., Wilkinson,

P.A., Coghill, J., Waterfall, C., Davassi, A., Scopes, G., Pirani, A., Webster, T.,

Brew, F., Bloor, C., King, J., West, C., Griffiths, S., King, I., Bentley, A.R., and

Edwards, K.J., 2016. High-density SNP genotyping array for hexaploid wheat

and its secondary and tertiary gene pool. *Plant Biotechnology Journal*, 14 (5),

1195–1206.

Yao, W., Li, G., Yu, Y., and Ouyang, Y., 2018. funRiceGenes dataset for

comprehensive understanding and application of rice functional genes.

GigaScience, 7 (1), 1–9.

Zadoks, J.C., Chang, T.T., and Konzak, C.F., 1974. A decimal code for the growth

stages of cereals. *Weed Research*, 14 (6), 415–421.

Zakhrabekova, S., Gough, S.P., Braumann, I., Muller, A.H., Lundqvist, J., Ahmann,

K., Dockter, C., Matyszczak, I., Kurowska, M., Druka, A., Waugh, R., Graner, A.,

Stein, N., Steuernagel, B., Lundqvist, U., and Hansson, M., 2012. Induced

mutations in circadian clock regulator *Mat-a* facilitated short-season adaptation

and range extension in cultivated barley. *Proceedings of the National Academy*

of Sciences, 109 (11), 4326–4331.

Zhai, H., Jiang, C., Zhao, Y., Yang, S., Li, Y., Yan, K., Wu, S., Luo, B., Du, Y., Jin,

H., Liu, X., Zhang, Y., Lu, F., Reynolds, M., Ou, X., Qiao, W., Jiang, Z., Peng,

T., Gao, D., Hu, W., Wang, J., Gao, H., Yin, G., Zhang, K., Li, G., and Wang, D.,

2021. Wheat heat tolerance is impaired by heightened deletions in the distal end

of 4AL chromosomal arm. *Plant Biotechnology Journal*, 19 (5), 1038–1051.

Zhao, J., Huang, X., Ouyang, X., Chen, W., Du, A., and Zhu, L., 2012. *OsELF3-1* ,

an Ortholog of *Arabidopsis EARLY FLOWERING 3* , Regulates Rice Circadian

Rhythm and Photoperiodic Flowering. *PLoS ONE*, 7 (8), 1–10.

Zielinski, T., Moore, A.M., Troup, E., Halliday, K.J., and Millar, A.J., 2014. Strengths

and limitations of period estimation methods for circadian data. *PLoS ONE*, 9

(5).

Zikhali, M., Wingen, L.U., and Griffiths, S., 2016. Delimitation of the *Earliness per se*

D1 (*Eps-D1*) flowering gene to a subtelomeric chromosomal deletion in bread

wheat (*Triticum aestivum*). *Journal of Experimental Botany*, 67 (1), 287–299.

992 **Figures**

994 **Fig 1**. **Variation in time to reach GS55 across a UK wheat MAGIC population**.

995 Time in days for individual MAGIC RILs (recombinant inbred lines) to reach GS55 for the (A) 2012/2013

996 growing season and (B) 2013/2014 growing season. The number of days after sowing to reach GS55

997 for the eight MAGIC parents is indicated by arrows.

1000

Fig 2. *ELF3* **is a candidate gene for** *Eps-B1***.**

(A) Syntenic relationships of the *Eps-B1* QTL to the *Eps-D1* QTL and the *T. monococcum Eps-A^m1*

QTL. *ELF3* highlighted in green. (B) *TaELF3-B1* gene model with exons represented by filled

rectangles, UTR's as white rectangles. The two promoter cytosine repeat polymorphisms with location

relative to the transcription start site (TSS) are indicated with the cytosine (C) repeat number for Claire,

Robigus, Cadenza and Paragon. The Ser674Gly SNP is situated within exon 4 of *ELF3-B1.*

1017

Fig 3. A subtelomeric chromosomal introgression and inversion containing *ELF3-D1* is likely to be the causal polymorphism underlying the *Eps-D1* QTL.

(A) Syntenic relationships between the distal end of Chinese Spring (CS) 1D chromosome (IWGSCv1.1) and Jagger (Jag) 1D (PGSBv2.0), *TaELF3-D1* highlighted in green, each black arrow corresponds to a gene in the forward (pointing right) or reverse (pointing left) strand. (B) the Cadenza/Jagger *ELF3-D1*

1023   contains a deletion within intron 2 which is not present in CS, F and R refer to relative primer binding

1024   sites for PCR amplification shown in (C). (D) Agarose gel separation of PCR products from (C). A PCR

1025   product at 776bp is indicative of the presence of a subtelomeric chromosomal introgression and

1026   inversion.

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

Fig 4. Circadian rhythms of delayed chlorophyll fluorescence in the MAGIC parents.

(A) Mean DF luminescence in counts per second (cps) normalised to -1 to 1 using Biodare2 with error bars representing standard deviation (n = 27 - 33). Circadian amplitude (B), period (C) and RAE (D) calculated using FFT-NLLS (Biodare2). Upper and lower hinges represent the first and third quartiles (25th and 75th percentiles), the middle hinge represents the median value, red triangle represents mean value and black dots represent individual replicates.

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

Fig 5. *Ttelf3*-null is arrhythmic in continuous light.

(A) Mean *Fm'* of *Ttelf3*-null (red) and *TtELF3-WT* (black) in LL (n = 24) (B) Mean period length (hours)

and (C) RAE values for plot (A). Upper and lower hinges represent the first and third quartiles (25th

and 75th percentiles), the middle hinge represents the median value, yellow triangle represents mean

value and black dots represent individual replicates. (D) Mean leaf surface temperature relative to

1080   background and normalised to the mean in LL and constant environmental temperature (20°C) for

1081   *Ttelf3-null* (red) *TtELF3-WT* (black) and *Ttelf3-null* (red) (n = 7) (E) Mean period length and (F) RAE

1082   for plot (D). For clarity every 20th point plotted in (D). Error bars represent SEM.

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

Fig 6. Abundance of wheat circadian clock transcripts in the *TtELF3-WT* and *ttelf3* lines in LD and LL.

(A-G) Mean abundance of circadian oscillator transcripts (n= 3 - 5) in *TtELF3*-WT (black) and *Ttelf3*-null (red) in a 24 h LD cycle in long day conditions (16 h light at 250 μmol m^{-2} s^{-1}, 20°C: 8 h dark 16°C, represented by vertical grey bar) followed by constant light and temperature (20°C) from time 24 to 96 h. Transcript abundance (ΔΔCq) is relative to *TtRP15* and *TtRPT5A* (A) *TtLHY,* (B) *Ppd-1,* (C) *TtPRR73,* (D) *TtTOC1,* (E) *TtGI,* (F) *TtLUX* and (G) *TtELF3* . White bar represents light, black bar represents darkness and grey bar represents subjective night.

1118

Fig 7. Functional *LUX* is required for maintenance of robust circadian oscillations under LL

conditions in hexaploid wheat.

(A) Mean $F_{m'}$ for Chogokuwase (blue), Geurumil (green) and Minaminokomugi (orange) in LL (n = 16)

(B) Mean period and (C) RAE for (A) calculated using FFT-NLLS (Biodare2). Upper and lower hinges

1123 represent the first and third quartiles (25th and 75th percentiles), the middle hinge represents the median

1124 value, red triangle represents mean value and black dots represent individual replicates. (D) Mean

1125 Chogokuwase, Geurumil and Minaminokomugi leaf temperature relative to background in LL (n = 4).

1126 (E) mean period and (F) RAE for plot (D). Error bars are SEM, for (D) every 20th point plotted for clarity.

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

Table 1: **Summary of circadian clock associated QTLs and their mapping positions.** Output from

univariate QTL model for the NIAB 2013 and 2014 GS55 data. Gene (Mb) refers to the BLAST hit

starting coordinate of the gene's cDNA against the IWGSCRefSeqv1.1. Dist (cM) refers to the markers

genetic mapping distance in centiMorgan. Dist (Mb) refers to the BLAST hit starting coordinate of the

marker sequence against the IWGSCRefSeqv1.1 LOGP is –log10(p) which corresponds to the overall

significance of the QTLs as a measure of its strength. % var is the percentage of genetic variance

contributed by QTL. Further description of methodology and column headers can be found in (Verbyla

et al. 2014).

Circadian clock gene	Gene (Mb)	Chr	L. Marker	dist (cM)	dist (Mb)	R. Marker	dist (cM)	dist (Mb)	Year	LOGP	% var
TaELF3	685,646,575	1B	Ra_c109187_371	296.34	685,741,261	RAC875_c37075_283	297	687,715,336	2013	4.48	3.4
		1B	Ra_c109187_371	296.34	685,741,261	RAC875_c37075_284	297	687,715,336	2014	3.2	2
TaELF3	493,485,612	1D	Kukri_c29687_369	100.95	493,212,138	Kukri_rep_c69829_307	101.95	488,577,079	2014	2.32	4.1
Ppd-1	33,952,591	2D	Kukri_c27309_590	48.57	31,806,310	BS00064538_51	56.64	33,370,535	2013	20.04	17.2
		2D	Kukri_c27309_590	48.57	31,806,310	BS00064538_51	56.64	33,370,535	2014	43.92	30.2
TaZTL	625,638,942	7A	Kukri_c6676_172	207.62	610,934,248	Ku_c2990_1939	208.12	621,322,194	2013	2.87	2.6

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

 Table 2. 2013 and 2014 GS55 QTL summary for the *eps-B1 locus*. Estimated parental haplotype

effects on RIL BLUPs. Abbreviations as per (Verbyla *et al.* 2014): Dist (cM) refers to the markers

genetic mapping distance in centiMorgans (cM). LOGP is –log10(p) and corresponds to the overall

significance of the QTL as a measure of its strength. Founder Prob: probability QTL is from founder

shown (p). Strength of associations can thus be reported at the overall or founder level. Size: Size of

effect of each founder allele (days to GS55).

QTL	Year	Founder	Size	Founder Prob	% var	LOGP	KASP (chr1B:685645813)
Eps-B1	2013	Alchemy	-0.555	0.054	3.4	4.48	A:A
		Brompton	0.317	0.123			G:G
		Claire	-0.076	0.415			A;A
		Hereward	0.623	0.008			G:G
		Rialto	0.331	0.118			G:G
		Robigus	-0.102	0.376			A:A
		Soissons	-0.683	0.003			A:A
		Xi-19	0.053	0.426			G:G
Eps-B1	2014	Alchemy	-0.448	0.196	2	3.2	A:A
		Brompton	0.395	0.178			G:G
		Claire	-0.442	0.202			A:A
		Hereward	0.591	0.072			G:G
		Rialto	0.39	0.182			G:G
		Robigus	-0.313	0.264			A:A
		Soissons	-1.085	0.002			A:A
		Xi-19	0.453	0.15			G:G

Table 3. 2012 and 2014 GS55 QTL summary for the *eEps-D1 locus*. Estimated parental haplotype

effects on RIL BLUPs. Abbreviations as per (Verbyla *et al.* 2014): Dist (cM) refers to the markers

genetic mapping distance in centiMorgans (cM). LOGP is –log10(p) and corresponds to the overall

significance of the QTL as a measure of its strength. Founder Prob: probability QTL is from founder

shown (p). Strength of associations can thus be reported at the overall or founder level. Size: Size of

effect of each founder allele (days to GS55).

1197

QTL	Year	Founder	Size	Founder Prob	% var	LOGP	PCR (Figure 3D)
Eps-D1	2012	Alchemy	0.552	0.254	0.9	0.56	1097bp
		Brompton	0.105	0.446			1097bp
		Claire	-0.11	0.447			1097bp
		Hereward	0.021	0.49			1097bp
		Rialto	-0.044	0.477			1097bp
		Robigus	0.003	0.498			1097bp
		Soissons	0.734	0.134			1097bp
		Xi-19	-1.261	0.015			776bp
Eps-D1	2014	Alchemy	-0.058	0.472	4.1	2.32	1097bp
		Brompton	0.869	0.104			1097bp
		Claire	-1.412	0.051			1097bp
		Hereward	-0.582	0.252			1097bp
		Rialto	-0.071	0.457			1097bp
		Robigus	-0.265	0.38			1097bp
		Soissons	1.244	0.013			1097bp
		Xi-19	-0.79	0.053			776bp

1198

1199

1200

1201

1202

1203

1204

1205

1206   **Table 4: Wheat lines used in this study.** GRU refers to the germplasm resource unit at the John Innes

1207   Centre, Norwich.

Species	Cultivar	Genotype	Reference
Ae. speltoides			GRU 2140023
T. aestivum	Alchemy		(Mackay *et al.* 2014)
T. aestivum	Brompton		(Mackay *et al.* 2014)
T. aestivum	Claire		(Mackay *et al.* 2014)
T. aestivum	Hereward		(Mackay *et al.* 2014)
T. aestivum	Rialto		(Mackay *et al.* 2014)
T. aestivum	Robigus		(Mackay *et al.* 2014)
T. aestivum	Soissons		(Mackay *et al.* 2014)
T. aestivum	Xi-19		(Mackay *et al.* 2014)
T. aestivum	Cadenza		GRU WBCDB0013-PG1
T. aestivum	Jagger		GRU PANG0006
T. aestivum	Spark		GRU WBCDB0052
T. aestivum	Chinese Spring		
T. aestivum	Chogokuwase		(Mizuno *et al.* 2016)
T. aestivum	Minaminokomugi		(Mizuno *et al.* 2016)
T. aestivum	Geurumil		(Mizuno *et al.* 2016)
T. aestivum	Axona		GRU W3580
T. aestivum	Cordiale		GRU W10003
T. aestivum	Maris Fundin		GRU W1357
T. aestivum	Tonic		GRU W3572
T. monococcum			GRU T1040057
T. sphaerococcum			GRU 1210007
T. timopheevii			GRU 116002
T. turgidum	Kronos	*Ttelf3*-null line	(Alvarez *et al.* 2016)
T. turgidum	Kronos	*TtELF3*-WT	(Alvarez *et al.* 2016)

1208

1209

1210

1211

1212

1213

1214

1215

1216   **Table 5: Primers used in this study**

Gene	Sequence 5'-3'
TtCCA1	F: CCTGGAATTGGAGATGGAGA
	R: TGAGCATGGCTTCTGATTTG
TtELF3	F: TCTCCAGATGATGTTGTCGGT
	R: CTCGAACACTTGGACAGCAAA
TtGI	F: GGTAGGTGATAGACGGCACTT
	R: GTGCTACAGATGGGATGCTTG
TtLUX	F: ACAAGCGGTTCGTGGAGG
	R: CCTGCATCCGCTTGACGTA
TtPpd-1	F: CCTGTGGACTGTCGATCTCAA
	R: CAAGGGATGGCAGCGATAATG
TtPRR73	F: TCCCGAAGTTCCTCTCTTTCC
	R: AGCGGTAGTGGCAATGACA
TtTOC1	F: GGCATGGCACTTCATTCAGTT
	R: GCACATTCATACCAGCAGGAC
TtRP15	F: GCACACGTGCTTTGCAGATAAG
	R: GCCCTCAAGCTCAACCATAACT
TtRPT5A	F: GCTGGCTCGTTCAACTGATG
	R: GGACCAAGCGTTCTGATTACTC
TaELF3-D1	F: CAACGGCGCTTCTAATCTG
	R: AAACCCGCTGCTGACTGTATA

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1 Supporting information

3 **Figure S1 The Ser674 is highly conserved in monocots.** MCoffee alignment visualised in CLC
4 Genomics Workbench, Amino acids coloured based on RasMol colour scheme.

S2 Fig. The Cadenza/Jagger *ELF3-D1* intronic deletion is present within Cordiale, Maris Fundin and Tonic.
(A) Sequence alignment for PCR fragments between exon 2 and 3 of *ELF3-D1* (see Fig 3) and (B) Agarose gel separation of PCR products. A PCR product at 776bp is indicative of the presence of the introgressed and inverted region.

A

Chinese Spring

Ae. speltoides

T. timopheevii

B

Ladder | *Ae. speltoides* | *T. monococcum* | *T. dicoccum* | *T. sphaerococcum* | *T. timopheevii*

1097 bp ◀

776 bp ◀

S3 Fig. The Cadenza/Jagger *ELF3-D1* intronic deletion is present within *Ae. speltoides* and *T. timopheevii*.
(A) Sequence alignment for PCR fragments between exon 2 and 3 of *ELF3-D1* (see Fig 3) and (B) Agarose gel separation of PCR products. A PCR product at 776bp is indicative of the presence of the introgressed and inverted region.

S4 Fig. Phylogeny of the 1D Jagger/ Cadenza shared haplotype using the 820k SNP dataset from CerealsDB.
Neighbour Joining Tree and bootstraps created from 820k SNPs using R package "ape". All SNPs mapping to the CS coordinates (1D:482.83Mb until 1D end) of the shared Cadenza and Jagger haplotype were used. Heterozygotes were set as missing.

S5 Fig. Phylogeny of *ELF3-D1* gene.

Maximum Likelihood phylogeny based on T-coffee alignment of publicly available coding sequences.

90 **S6 Fig. The location of mutations within the A and B sub-genome copies of *TtELF3*.**

91 (A-B) Scale diagram of (A) *TtELF3-A1* and (B) *TtELF3-B1* where black rectangles represent exons
92 ordered 5'-3' (left-to-right), horizontal line above represents length of gene from start to stop codon (bp),
93 space between exons represents introns. Yellow triangles denote the position of premature stop codons
94 denoting the source TILLING line as annotated in (Alvarez *et al.* 2016).

S7 Fig. *Ttelf3-null* is arrhythmic for PF parameters in LL.

(A) Mean F_v/F_m of *TtELF3*-WT (black) and *Ttelf3*-null (red) in LL (n = 24) (B) Mean period length (hours)
and (C) RAE values for plot (A). (D) Mean NPQ of *TtELF3-WT* (black) and *Ttelf3*-null (red) in LL (n =
24) (E) Mean period length (hours) and (F) RAE values for plot (D). Error bars represent SEM. For plots
(B), (C), (D) & (E) upper and lower hinges represent the first and third quartiles (25th and 75th
percentiles), the middle hinge represents the median value, yellow triangle represents mean value and
black dots represent individual replicates.

S8 Fig. *Ttelf3-null* is arrhythmic for DF in LL.

(A) Mean DF luminescence in counts per second (cps) normalised to -1 to 1 using Biodare2 with error bars representing standard deviation (n = *TtELF3*-WT 63, *Ttelf3*-null *72*). (B) Amplitude and (C) RAE for *TtELF3*-WT (black) and *Ttelf3*-null (red) in (A) calculated using FFT-NLLS fit in Biodare2 where upper and lower hinges represent the first and third quartiles (25[th] and 75[th] percentiles), the middle hinge represents the median value, yellow triangle represents mean value and black dots represent individual replicates.

134
135
136

S9 Fig. The location of mutations in the different sub-genome copies of *LUX*.

(A-C) Diagram of the A sub-genome (A), B sub-genome (B) and D sub-genome (C) copies of *LUX* where black rectangle represents the *LUX* exon drawn 5' to 3' (left-to-right) and the embedded white rectangle denotes the highly conserved MYB-domain. Yellow triangles denote the location of amino acid substitution within the SHAQKYF motif (A, C). (B) line and annotation show the location of 142 bp deletion in the *lux-B1* allele (Mizuno *et al.* 2016).

143
144
145
146

S10 Fig. The Ser674Gly SNP is a candidate for the causal *Eps-B1* SNP.

KASP Genotype association with MAGIC founder haplotype effects at the *Eps-B1* locus. Founder haplotype effects are calculated as the mean of the 2013 and 2014 founder effect. Genotype of A|A corresponds to homozygous for adenine, encoding the ancestral Serine, whilst G|G corresponds to homozygous guanine encoding the non-ancestral Glycine residue.

Figure S11 There is no significant difference between the circadian rhythms of kronos and

***Ttelf3-null* lines containing the *Ppd-A1a* and *Ppd-A1b* alleles.**

(A) Mean DF luminescence in counts per second (cps) normalised to -1 to 1 using Biodare2 with error
bars representing standard deviation (n = *Ppd-A1a* 68, *Ppd-A1b* 62). (B) Amplitude and (C) RAE for
Ppd-A1a (purple) and *Ppd-A1b* (black) in (A). (D) Mean DF luminescence in counts per second (cps)
normalised to -1 to 1 using Biodare2 with error bars representing standard deviation (n = *Ttelf3-null*
Ppd-A1a 66, *Ttelf3-null Ppd-A1b* 22) and (E) RAE for *Ttelf3-null Ppd-A1a* (blue) and *Ttelf3-null Ppd-*
A1b (red). Amplitude and RAE calculated using FFT-NLLS fit in Biodare2 where upper and lower hinges
represent the first and third quartiles (25th and 75th percentiles), the middle hinge represents the median

value, yellow triangle represents mean value and black dots represent individual replicates for plants
where estimation was possible.

200
201
202 **S12 Fig. In-phase and anti-phase leaf temperature rhythms in continuous light.**

203 (A) Mean leaf temperature relative to background (n = 3) of four week old cadenza plants entrained to
204 16 h light:8 h dark cycles in-phase (yellow) or 16 h light: 8 h dark cycles that were 12 hours anti-phased
205 (purple). (B) Phase plot of data shown in (A).

 S1 Table: GS55 2012 QTLs

222 Output from the reanalysis of the NIAB 2012 GS55 data using MPWGAIM, originally analysed using Bayesian Networks (Scutari *et al.* 2014). Dist (cM) refers
223 to the markers genetic mapping distance in centiMorgan. Founder corresponds to founder variety in alphabetical order (1 Alchemy, 2 Brompton, 3 Claire, 4
224 Hereward, 5 Rialto, 6 Robigus, 7 Soissons and 8 Xi-19). Size: Size of effect of each founder allele (days to GS55). LOGP is –log10(p) and corresponds to the
225 overall significance of the QTL as a measure of its strength. Prob: probability QTL is from all founders shown (p). Founder LOGP and Founder Prob are
226 LOGP and Prob for individual founders. % var is the percentage of genetic variance explained by each QTL.

Chr	Left Marker	dist (cM)	Right Marker	dist (cM)	Founder	Size	Founder Prob	Founder LOGP	Prob	% var	LOGP
1B	IAAV5516	297.84	BS00001128_51	298.34	1	-0.207	0.399	0.4	0.048	1.6	**1.32**
					2	0.946	0.077	1.11			
					3	0.112	0.445	0.35			
					4	0.318	0.304	0.52			
					5	0.698	0.148	0.83			
					6	-1.119	0.07	1.15			
					7	-1.185	0.021	1.67			
					8	0.437	0.259	0.59			
1D	Kukri_c2464_592	18.04	D_contig57829_235	18.21	1	-0.81	0.113	0.95	0.003	2.5	**2.59**
					2	-0.569	0.238	0.62			
					3	1.282	0.042	1.37			
					4	-1.024	0.066	1.18			
					5	0.631	0.27	0.57			
					6	0.849	0.12	0.92			
					7	-1.18	0.036	1.45			
					8	0.82	0.21	0.68			
1D	CAP11_c1157_168	100.79	Kukri_c29687_369	100.95	1	0.552	0.254	0.6	0.275	0.9	**0.56**
					2	0.105	0.446	0.35			
					3	-0.11	0.447	0.35			
					4	0.021	0.49	0.31			
					5	-0.044	0.477	0.32			

Chr	Left Marker	dist (cM)	Right Marker	dist (cM)	Founder	Size	Founder Prob	Founder LOGP	Prob	% var	LOGP
					6	0.003	0.498	0.3			
					7	0.734	0.134	0.87			
					8	-1.261	0.015	1.82			
2A	RAC875_c20273_823	35.38	BS00094817_51	35.88	1	0.45	0.218	0.66	0.035	1.6	**1.45**
					2	0.771	0.163	0.79			
					3	0.133	0.418	0.38			
					4	-1.126	0.032	1.5			
					5	-0.193	0.404	0.39			
					6	-1.262	0.019	1.71			
					7	0.739	0.11	0.96			
					8	0.487	0.241	0.62			
2B	BS00060618_51	317.61	BS00000012_51	318.61	1	0.176	0.417	0.38	0.049	1.6	**1.31**
					2	0.748	0.175	0.76			
					3	1.643	0.022	1.66			
					4	-0.88	0.107	0.97			
					5	-0.565	0.206	0.69			
					6	-0.392	0.267	0.57			
					7	-0.326	0.304	0.52			
					8	-0.404	0.265	0.58			
2D	Kukri_c27309_590	48.57	BS00064538_51	56.64	1	0.669	0.293	0.53	0	18.5	**18.33**
					2	2.727	0.009	2.03			
					3	0.215	0.44	0.36			
					4	1.112	0.211	0.68			
					5	0.759	0.276	0.56			
					6	0.897	0.245	0.61			
					7	-6.194	0	7.11			
					8	-0.185	0.437	0.36			
3A	wsnp_Ex_rep_c106152_90334299	78.8	Kukri_c51247_322	83.81	1	0.671	0.184	0.74	0.001	3	**2.84**
					2	-1.839	0.006	2.25			

Chr	Left Marker	dist (cM)	Right Marker	dist (cM)	Founder	Size	Founder Prob	Founder LOGP	Prob	% var	LOGP
					3	-0.671	0.187	0.73			
					4	0.716	0.155	0.81			
					5	-0.885	0.102	0.99			
					6	0.215	0.374	0.43			
					7	0.247	0.364	0.44			
					8	1.546	0.01	2.01			
3D	RAC875_c48773_253	31.11	RFL_Contig2471_119	31.28	1	-0.133	0.442	0.35	0.417	0.7	**0.38**
					2	0.159	0.431	0.37			
					3	-0.351	0.351	0.45			
					4	-0.525	0.276	0.56			
					5	-0.025	0.489	0.31			
					6	-0.443	0.244	0.61			
					7	1.191	0.021	1.68			
					8	0.128	0.423	0.37			
4A	BS00064140_51	133.18	BS00101512_51	134.18	1	0.521	0.273	0.56	0.002	2.4	**2.75**
					2	0.176	0.415	0.38			
					3	0.277	0.376	0.42			
					4	0.266	0.36	0.44			
					5	0.118	0.447	0.35			
					6	0.019	0.488	0.31			
					7	0.727	0.187	0.73			
					8	-2.104	0	3.63			
5B	BobWhite_c22572_782	196.48	Kukri_c836_513	196.65	1	-3.398	0.001	3.11	0	6	**3.85**
					2	0.725	0.232	0.63			
					3	0.886	0.158	0.8			
					4	1.621	0.088	1.05			
					5	-0.001	0.5	0.3			
					6	-1.505	0.037	1.43			
					7	0.704	0.206	0.69			

Chr	Left Marker	dist (cM)	Right Marker	dist (cM)	Founder	Size	Founder Prob	Founder LOGP	Prob	% var	LOGP
					8	0.968	0.141	0.85			
6B	wsnp_Ex_c12450_19850827	215.75	tplb0045b09_1555	216.42	1	0.089	0.461	0.34	0.269	1.1	**0.57**
					2	0.723	0.117	0.93			
					3	-0.552	0.264	0.58			
					4	0.368	0.333	0.48			
					5	-0.111	0.447	0.35			
					6	0.247	0.388	0.41			
					7	0.467	0.299	0.53			
					8	-1.232	0.02	1.71			
6D	BS00094895_51	124.37	GENE_4168_925	125.04	1	-0.044	0.472	0.33	0.169	1	**0.77**
					2	-0.093	0.449	0.35			
					3	0.85	0.082	1.09			
					4	-0.622	0.216	0.66			
					5	-0.674	0.198	0.7			
					6	0.592	0.169	0.77			
					7	-0.56	0.152	0.82			
					8	0.551	0.187	0.73			
7B	BS00036788_51	58.06	CAP7_c10566_170	58.22	1	-0.73	0.18	0.74	0.008	2.5	**2.09**
					2	-0.247	0.402	0.4			
					3	0.422	0.312	0.51			
					4	1.264	0.029	1.53			
					5	0.22	0.411	0.39			
					6	1.249	0.065	1.19			
					7	-0.704	0.173	0.76			
					8	-1.473	0.012	1.91			

229 **S2 Table. The chr1B_685645813/ Ser674Gly SNP is globally rare.** Wheat HapMap data of a diverse
230 germplasm collection and the 10+ Wheat genome project (italics). UK varieties highlighted in bold.
231 Genotype of A|A corresponds to homozygous for adenine at the Ser674Gly SNP encoding the ancestral
232 serine whilst G|G corresponds to homozygous guanine encoding the non-ancestral glycine (red).
233

Variety	Genotype (forward strand)	Variety	Genotype (forward strand)	Variety	Genotype (forward strand)
93	A|A	*Jagger*	*A|A*	PI278297	A|A
102	A|G	*Julius*	*A|A*	PI349512	A|A
403	A|A	klein_chamaco	A|A	PI366716	A|A
407-IV_60	A|A	*Lancer*	*A|A*	PI366905	A|A
AC_Barrie	A|A	M6	A|A	PI382150	A|A
acc1	A|A	*Mace*	*A|A*	PI406517	A|A
acc2	A|A	Marquis	A|A	PI445736	A|A
acc3	A|G	Neepawa	A|A	PI470817	A|A
acc4	A|G	*Norin61*	*A|A*	PI477870	A|A
acc5	A|A	Opata	A|A	PI481718	A|A
Alabasskaja	A|A	***Paragon***	***G|G***	PI481923	A|A
Arina	*G|G*	pavon	A|A	PI565213	A|A
Avalon	**A|A**	pbw343	A|A	PI82469	A|A
Cadenza	***G|G***	PI153785	A|A	PI8813	A|A
CDC Landmark	*A|A*	PI166180	A|A	PR267	A|A
CDC Stanley	*A|A*	PI166333	A|A	rac875	A|A
chakwat86	A|A	PI177943	A|A	**Rialto**	**G|G**
cham6	A|A	PI185715	A|A	***Robigus***	***A|A***
Chinese Spring	A|A	*PI190962*	*A|A*	*Sy Mattis*	*A|A*
Claire	***A|A***	PI192001	A|A	**Taxi**	**A|A**
clear_white	A|A	PI192569	A|A	Truman	A|A
dharwar_dry	A|A	PI222669	A|A	Utmost	A|A
Estacao	A|A	PI245368	A|A	vorobey	A|A
hidhab	A|A	PI262611	A|A	*Weebil*	*A|A*

239 **S3 Table. The Ser674Gly SNP segregates with heading date in UK varieties.** Additional genotyping
240 and heading information from Zikhali *et al.*, 2016 (Zikhali *et al.* 2016) TGAC reference sequences
241 (Clavijo *et al.* 2017). MAGIC founder varieties highlighted in bold, non-ancestral glycine in red.

Variety	Allele	Heading
Alchemy	A|A	early
Claire	A|A	early
Robigus	A|A	early
Avalon	A|A	early
Brompton	G|G	late
Hereward	G|G	late
Rialto	G|G	late
Xi-19	G|G	late
Paragon	G|G	NA
Cadenza	G|G	late
Spark	G|G	NA
Badger	G|G	NA
Savannah	G|G	NA

ELF3-1D region	Exon 2	Intron 2																									
Variant location	T1832C	C1933T	C1982T	A2028G	C2040A	A2046G	T2056G	A2084G	A2102G	G2104A	G2109C	C2128T	A2161G	2167-2471	T2483C	2504-2505	A2521G	2555	T2562C	2609	2613-2614	2616-2622	2625-2627	2633	2636-2637	G2646C	2651-2653
Cultivar / Haplotype	Var 1	Var 2	Var 3	Var 4	Var 5	Var 6	Var 7	Var 8	Var 9	Var 10	Var 11	Var 12	Var 13	Var 14	Var 15	Var 16	Var 17	Var 18	Var 19	Var 20	Var 21	Var 22	Var 23	Var 24	Var 25	Var 26	Var 27
Alchemy / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
ArinaLrFor / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Brompton / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Cadenza / H2	C	T	T	G	A	G	G	G	G	A	C	T	G	Del	C	AG	G	T	C	-	-	Del	Del	-	-	C	TGG
Chinese Spring / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Claire / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Hereward / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Jagger / H2	C	T	T	G	A	G	G	G	G	A	C	T	G	Del	C	AG	G	T	C	-	-	Del	Del	-	-	C	TGG
Julius / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Landmark / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Mace / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Norin61 / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Paragon / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Rialto / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Robigus / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Soissons / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Spark / H2	C	T	T	G	A	G	G	G	G	A	C	T	G	Del	C	AG	G	T	C	-	-	Del	Del	-	-	C	TGG
Stanley / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
SyMattis / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Weebil / H1	T	C	C	A	C	A	T	A	A	G	G	C	A	Ins	T	-	A	-	T	G	GT	Ins	Ins	A	CT	G	CAA
Xi-19 / H2	C	T	T	G	A	G	G	G	G	A	C	T	G	Del	C	AG	G	T	C	-	-	Del	Del	-	-	C	TGG

267 **S5 Table. Alignment of 300kb surrounding *ELF3-1D* in 10+ wheat genome lines.** Average
268 nucleotide identity scores from alignment of 300kb sequence surrounding *ELF3-1D* in the 10+ wheat
269 genome cultivars. Alignment produced using the whole genome alignment tool from CLC workbench
270 21.0.3 using a 15-base seed with no mismatches.
271

Cultivar	ArinaLrFor	CDC_Landmark	CDC_Stanley	Lancer	Mace	Norin61	CSv1.0	Julius	SYMattis	Jagger	Cadenza
ArinaLrFor	100.00	99.82	99.94	99.90	99.95	99.81	99.89	99.93	99.97	91.00	91.44
CDC_Landmark	99.82	100.00	100.00	99.89	99.85	99.93	99.82	99.90	99.94	90.57	91.44
CDC_Stanley	99.94	100.00	100.00	99.89	99.95	99.97	99.89	99.92	99.94	90.76	91.44
Lancer	99.90	99.89	99.89	100.00	99.94	99.92	99.91	99.94	99.93	91.00	91.42
Mace	99.95	99.85	99.95	99.94	100.00	99.90	99.86	99.98	99.98	90.51	91.44
Norin61	99.81	99.93	99.97	99.92	99.90	100.00	99.95	99.93	99.95	91.01	91.43
CSv1.0	99.89	99.82	99.89	99.91	99.86	99.95	100.00	99.93	99.83	91.00	91.44
Julius	99.93	99.90	99.92	99.94	99.98	99.93	99.93	100.00	99.99	91.00	91.43
SYMattis	99.97	99.94	99.94	99.93	99.98	99.95	99.83	99.99	100.00	91.00	91.43
Jagger	91.00	90.57	90.76	91.00	90.51	91.01	91.00	91.00	91.00	100.00	99.94
Cadenza	91.44	91.44	91.44	91.42	91.44	91.43	91.44	91.43	91.43	99.94	100.00

294 **S6 Table. Promoter motif analysis of *TtELF3.*** Column A refers to the A sub-genome, B sub-genome
295 and D to the D sub-genome. Motifs and corresponding sequences taken from (Wang *et al.* 1997,
296 Gendron *et al.* 2012, Huang *et al.* 2012).
297

Motif	Abbreviation	Sequence	A	B	D	Arabidopsis
Evening element	EE	AAAATATCT	0	1	0	0
Evening Like element	ELE	AATATCT	1	0	0	1
CCA1 binding site (A variant)	/	AAAAATCT	0	0	0	0
CCA1 binding site (C variant)	/	AACAATCT	0	0	0	2
Morning element 1	ME 1	CCACAC	0	1	1	0
Morning element 2	ME 2	GTGTGG	1	1	2	0
Hormone Up at Dawn 1	HUD 1	CACATG	2	3	4	2
Hormone Up at Dawn 2	HUD 2	CATGTG	3	9	6	0
G BOX (morning)	/	CACGTG	1	0	3	0
GATA (evening)	/	GGATA	4	8	6	6
Starch box (midnight)	/	AAGCCC	1	0	1	0
Telo-Box (midnight)	/	AAACCCT	2	0	0	0
Protein Box (midnight)	/	ATGGCC	0	2	1	0

S7 Table. Promoter motif analysis of *TtLHY.* Column A refers to the A sub-genome, B sub-genome
and D to the D sub-genome. Motifs and corresponding sequences taken from (Wang *et al.* 1997,
Gendron *et al.* 2012, Huang *et al.* 2012).

Motif	Abbreviation	Sequence	A	B	D	Arabidopsis
Evening element	EE	AAAATATCT	0	0	0	0
Evening Like element	ELE	AATATCT	0	0	0	0
CCA1 binding site (A variant)	/	AAAAATCT	0	1	0	0
CCA1 binding site (C variant)	/	AACAATCT	1	0	0	0
Morning element 1	ME 1	CCACAC	1	1	1	1
Morning element 2	ME 2	GTGTGG	2	0	0	0
Hormone Up at Dawn 1	HUD 1	CACATG	1	1	1	1
Hormone Up at Dawn 2	HUD 2	CATGTG	4	3	3	0
G BOX (morning)	/	CACGTG	1	1	0	1
GATA (evening)	/	GGATA	3	3	3	3
Starch box (midnight)	/	AAGCCC	0	0	1	1
Telo-Box (midnight)	/	AAACCCT	0	0	0	0
Protein Box (midnight)	/	ATGGGCC	0	1	1	0

344 **S8 Table. Promoter motif analysis of *TtLUX*.** Column A refers to the A sub-genome, B sub-genome
345 and D to the D sub-genome. Motifs and corresponding sequences taken from (Wang *et al.* 1997,
346 Gendron *et al.* 2012, Huang *et al.* 2012).

Motif	Abbreviation	Sequence	A	B	D	Arabidopsis
Evening element	EE	AAAATATCT	9	4	4	1
Evening Like element	ELE	AATATCT	3	1	1	2
CCA1 binding site (A variant)	/	AAAAATCT	1	0	0	2
CCA1 binding site (C vairiant)	/	AACAATCT	0	1	0	0
Morning element 1	ME 1	CCACAC	1	1	1	0
Morning element 2	ME 2	GTGTGG	1	1	0	0
Hormone Up at Dawn 1	HUD 1	CACATG	2	1	2	0
Hormone Up at Dawn 2	HUD 2	CATGTG	2	2	0	1
G BOX (morning)	/	CACGTG	2	3	2	2
GATA (evening)	/	GGATA	3	5	2	2
Starch box (midnight)	/	AAGCCC	0	0	2	0
Telo-Box (midnight)	/	AAACCCT	3	2	1	2
Protein Box (midnight)	/	ATGGGCC	1	1	2	0

S9 Table. Total number of SNPs within the coding sequence of *ELF3* in the 10+ Wheat
Genomes project cultivars. A SNP was recorded if the base was different to the Chinese Spring
IWGSCRefseqv1.1 sequence in the alignment. SNPs within the same codon were recorded
individually. It was not possible to create clear alignments between *ELF3-1D* Chinese Spring, Jagger
and Cadenza and so the coding sequence for the three cultivars was downloaded from ENSEMBL
plants, aligned and the number of SNPs manually counted. NS = non-synonymous SNP, S=
synonymous SNP, PAV = presence/absence variation. Cultivars were considered to share the same
allele if the CDS was identical. Zavitan is a tetraploid variety and so lacks a D sub genome.

	Chromosome					
	1A		1B		1D	
Cultivar	**NS**	**S**	**NS**	**S**	**NS**	**S**
ArinaLrFor	2	2	1	0	0	0
CDC Stanley	2	2	0	0	0	0
CDC Landmark	2	2	0	0	0	0
Cadenza	0	1	1	0	35	23
Claire	0	1	0	0	0	0
Jagger	2	3	0	1	35	23
Julius	PAV	PAV	0	0	1	0
Lancer	2	2	0	0	0	0
Norin61	2	2	0	0	0	0
Mace	2	2	0	0	0	0
PI190962	0	0	0	0	0	0
Paragon	0	1	1	0	0	0
Robigus	0	0	0	0	0	0
SY Mattis	0	1	0	0	0	0
Zavitan	7	2	1	3	*NA*	*NA*
Total No. SNP's	21	21	3	4	71	46
Total No. Alleles	5		3		3	