1 **Wheat *EARLY FLOWERING3* is a dawn-expressed circadian oscillator**

2 **component that regulates heading date**

Lukas Wittern[1][¶], Gareth Steed[1][¶], Laura J. Taylor[1], Dora Cano Ramirez[1], Gabriela

Pingarron-Cardenas,[1] Keith Gardner[2], Andy Greenland[2], Matthew A. Hannah[3], Alex

A. R. Webb[1]*

[1] Department of Plant Sciences, University of Cambridge, Downing Street,

Cambridge, CB2 3EA, UK

[2] NIAB, Huntingdon Road, Cambridge, UK

[3] BASF, BBCC – Innovation Center Gent, Technologiepark-Zwijnaarde, Gent,

Belgium

*Corresponding author

Email: aarw2@cam.ac.uk (AARW)

[¶]These authors contributed equally to this work

Abstract

Using an eight-parent Multiparent Advanced Generation Inter-Cross (MAGIC) population we investigated how variation at circadian clock-associated genes contributes to the regulation of heading date in UK and European winter wheat varieties. We identified homoeologues of *EARLY FLOWERING* 3 (*ELF3*) as candidates for the *Earliness per se* (*Eps*) *D1* and *B1 loci* in field conditions. We confirmed that a SNP within the coding region of *TaELF3-B1* is a candidate polymorphism underlying the *Eps-B1 locus.* We found that a reported deletion at the *Eps-D1 locus* encompassing *TaELF3-D1,* is instead a novel allele that lies within an introgression region containing an inversion relative to the Chinese Spring D genome. Using *T. turgidum cv. Kronos* carrying loss of function alleles of *TtELF3* we show that *ELF3* does regulate heading by demonstrating that the loss of a single *ELF3* homoeologue was sufficient to alter heading date. These studies demonstrated that *ELF3* forms part of the circadian oscillator but loss of all homoeologues was required to affect circadian rhythms. Similarly, loss of functional *LUX ARRHYTHMO* (*LUX*) in *T. aestivum*, an orthologue of a protein partner of Arabidopsis ELF3, severely disrupted circadian rhythms. *ELF3* and *LUX* transcripts are not co-expressed at dusk suggesting the structure of the wheat circadian oscillator might differ to that of Arabidopsis. Our demonstration that alteration to *ELF3* homoeologues can affect heading date separate from effects on the circadian oscillator suggests a role for *ELF3* in cereal photoperiodic responses that could be selected for, without pleiotropic deleterious alterations to circadian rhythms.

Introduction

Circadian clocks are endogenous timing mechanisms with a near 24 hour period that sequence biological events, provide anticipation of environmental rhythms and permit measurement of daylength to regulate seasonal adjustments (McClung 2019). Plant breeders have indirectly selected for variation at circadian-associated *loci* in many of the world's major crops when breeding to increase yield and improve crop performance by adapting genotypes to the local environment (Steed et al. 2021). Here we investigated whether variation at circadian-associated *loci* contributes to wheat yield traits in the field using a Multiparent Advanced Generation Inter-Cross (MAGIC) mapping population (Mackay *et al.* 2014). Wheat is a major crop, providing approximately 20% of the dietary calories and protein for the world's population (Hawkesford *et al.* 2013). Since the green revolution wheat yields have continued to increase but with the world's population estimated to reach 9.6 billion in 2050 (Gerland *et al.* 2014) it is likely that demand will outpace predicted yield increases (Hawkesford *et al.* 2013). Identification of novel genetic variation contributing to yield is critical to increasing the rate of yield gains in wheat.

Understanding of the nature of plant circadian oscillators is based primarily on investigation of Arabidopsis. In that model, the circadian oscillator contains a series of interlocking transcriptional feedback loops. The circadian oscillator cycle can be considered to start at dawn when expression of the transcriptional repressors *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) and *LATE ELONGATED HYPOCOTYL* (*LHY*) peaks, followed sequentially by peaks of expression of *PSEUDO RESPONSE REGULATOR 9* (*PRR9*)*, PRR7, PRR5, PRR3* and finally *PRR1* (also known as *TIMING OF CAB EXPRESSION 1* (*TOC1*)) at dusk (Hsu and Harmer 2014). PRR9

and PRR7 repress the expression of *CCA1* and *LHY* which releases repression of

evening phased genes including, *TOC1, GIGANTEA* (*GI*)*, EARLY FLOWERING 3*

(*ELF3*)*, ELF4* and *LUX ARRHYTHMO* (*LUX*) (Hsu and Harmer 2014). LUX is a MYB-

like transcription factor which associates with ELF3 and ELF4 to form the tripartite

evening complex (EC), which represses expression of *GI, TOC1, PRR7, PRR9* and

LUX (Dixon *et al.* 2011, Nusinow *et al.* 2011, Herrero *et al.* 2012)*.* Mutation of any of

the three EC components causes arrhythmia of the Arabidopsis circadian oscillator in

continuous light and temperature (Nagel and Kay 2012).

Orthologues of *ELF3* underlie *loci* regulating heading date in multiple crops, including

barley *early maturity8* (*eam8*) (Faure *et al.* 2012), rice *heading date17* (*Hd17*)

(Matsubara *et al.* 2012), *Earliness per se* (*Eps*)-A^m1 in *T. monococcum* and *Eps-D1* in

T. aestivum (Zikhali *et al.* 2016). In hexaploid wheat, the magnitude of the *Eps-D1*

effect on heading date increases with decreasing temperature (Ochagavía *et al.* 2019).

Orthologues of *TOC1* are associated with heading date, plant height and thousand

grain weight (Sun *et al.* 2020). An orthologue of *LUX* is a candidate for *Eps-3A^m* in *T.*

monococcum (Gawronski *et al.* 2014) and *HvLUX* is a candidate gene for *eam10* in

barley (Campoli *et al.* 2013). Furthermore, the *Photoperiod-1* (*Ppd-1*) genes*,* which

contribute to photoperiodic regulation of ear emergence, are orthologues of

Arabidopsis *PRRs* (Beales *et al.* 2007, Shaw *et al.* 2012, Bentley *et al.* 2013, Turner

et al. 2013).

We used the allelic diversity of a wheat MAGIC population to investigate the role of

circadian clock genes in regulating yield component traits in the field. The parent

varieties for the MAGIC population used in this study (Alchemy, Brompton, Claire,

Hereward, Rialto, Robigus, Soissons and Xi-19) are representative of UK and

European winter wheat varieties, capturing >80% of the polymorphism found across

UK wheat varieties from the last 70 years (Gardner *et al.* 2016). Our data demonstrate

that modified *ELF3* function can affect the regulation of flowering time without

disrupting circadian oscillations and thus potentially avoid associated yield penalties

(Dodd *et al.* 2005). Together, this suggests *ELF3* is a possible breeding target for crop

improvement. We find evidence that in wheat *ELF3* is expressed in the morning, rather

than the evening as it is in Arabidopsis and conclude there are likely differences in the

role for *ELF3* in the circadian system between wheat and Arabidopsis. Our finding of

separable roles for *ELF3* in the circadian oscillator and the regulation of heading date

suggest a direct role for ELF3 in photoperiodic regulation in cereals.

115 Results

116 Earliness per se *loci Eps-B1* and *Eps-D1* segregate in an

117 eight-parent wheat MAGIC population

We performed QTL analysis on a set of replicated trial data for 784 recombinant inbred

lines (RILs) for the NIAB Elite MAGIC population from the 2012/2013 and 2013/2014

growing seasons at the NIAB experimental farms in Cambridge, UK. Growth stage 55

(GS55, equivalent to heading date) (Zadoks *et al.* 1974) was chosen for analysis and

QTL mapping to enable comparisons with previous studies (Griffiths *et al.* 2009). In

both 2013 and 2014, Soissons, which contains the *Ppd-D1a* (photoperiod insensitive)

allele, reached GS55 6-8 days earlier than any of the other MAGIC parents, which

contain the *Ppd-D1b* allele (photoperiod sensitive) (Figure 1). Heading date of the

other MAGIC parents varied over 5.5 days in 2013 and 7 days in 2014.

We performed the QTL analysis using MPWGAIM (Verbyla *et al.* 2014) for GS55

phenotypes in both years and tested whether any circadian clock genes co-locate

with the identified QTLs (< 2 cM to nearest 90 k marker or < 5 Mb to gene). Because

we are interested in the effects of the circadian clock on yield traits, we report only

QTLs associated with circadian clock genes. We found that four circadian gene

orthologues (*TaELF3-B1, TaELF3-D1, Ppd-1, TaZTL-7A*) fell within 0.20, 0.27, 0.58

and 4.31 Mb respectively of one of the QTL flanking markers (Table 1). The absence

of significant heading date QTLs linked to vernalisation genes such as *VRN1*

confirmed that complete vernalisation had occurred, ensuring that observed

variations in heading date were true *Earliness per-se* (*Eps*) effects rather than a

variation in the vernalisation requirements between the MAGIC parents (Bentley *et*

al. 2013, Camargo *et al.* 2016). A previous study identified a single GS55 QTL in the

NIAB eight-parent MAGIC population (Camargo *et al.* 2016) but this difference in

findings is likely a result of different growth conditions (greenhouse compared to

field), lower replicate number (208 RILs in pots compared to 784 RILs in field plots)

and the use of a different mapping methodology. Further validation of our analysis

pipeline and approaches are described in the methods and reported in

(Supplemental Table S1).

A QTL on the distal end of chromosome 1B, delineated by the markers

Ra_c109187_371 and RAC875_c37075_283, mapped closely to the physical position

of *TaELF3-B1* (TraesCS1B02G477400) (Table 1). The 1B locus also had a

homoeologous QTL on chromosome 1D delineated by the markers Kukri_c29687_369

and Kukri_rep_c69829_307 identifying *TaELF3-D1* (TraesCS1D02G451200) as a

candidate gene (Table 1, Figure 2A). These QTLs co-locate with the previously

described heading date QTLs *Eps-B1* and *Eps-D1* (Griffiths *et al.* 2009). In our data,

Eps-B1 had a significant effect in both years, explaining 3.4% of genetic variance in

GS55 in 2013 and 2% in 2014 (Table 1). *Eps-D1* was also detected in both years but

only had an overall significant effect (–log10(p) = 2.32) in 2014, accounting for 4.1%

of genetic variance in GS55 in that year (Table 1). The *Eps-B1* and *Eps-D1 loci* were

chosen for further in-depth analysis because of their proximity to *ELF3* homoeologues.

TaELF3-B1 is a candidate gene for the *Eps-B1* QTL

At the 1B QTL, the alleles derived from the MAGIC parents Soissons, Alchemy,

Robigus and Claire are associated with a relative reduction in the time to GS55,

indicated by the negative sign of the QTL, whilst alleles from Rialto, Hereward, Xi-19

and Brompton are associated with a delay to GS55, indicated by the positive sign of

the QTL (Table 2). The interval between the markers Ra_c109187_371 (296.34cM /

685.7Mb) and RAC875_c37075_283 (297.00cM / 687.7Mb) is 0.66cM, corresponding

to a physical length of ca. 2 Mb (Figure 2A).

The IWGSCv1.1 gene annotation suggests there are 26 high confidence gene models

within the interval of Ra_c109187_371 and RAC875_c37075_283. Based on current

gene annotation and orthology to cloned rice genes (Yao *et al.* 2018) there are no

novel candidates for the *Eps-B1* locus in addition to those described previously

(Faricelli *et al.* 2010, Alvarez *et al.* 2016, Zikhali *et al.* 2016). Compared to the *T.*

monococcum physical map of the *Eps-A^{m}1* region (Alvarez *et al.* 2016) *TmADK* (similar

to rice putative kinase *ADK1)* and *TmCK1-like* (*CASEIN KINASE I ISOFORM DELTA-*

LIKE) are not syntenic and the locus containing *TaELF3*, *TaGRIK1-like*

(*GEMINIVIRUS REP INTERACTING KINASE 1 LIKE*) and *TaPWWP1* (*PWWP*

domain-containing protein) is inverted (Figure 2A). Nineteen genes had

homoeologues on the syntenic 1D chromosome (Figure 2A).

Publicly available sequences for *TaELF3-B1* in Claire, Robigus (earlier heading

alleles) (Clavijo *et al.* 2017) and Rialto and Cadenza – the parents of Xi-19 - (later

heading alleles) (Zikhali *et al.* 2016) differed by one nonsynonymous SNP within exon

4 at position chr1B:685645813. No further nonsynonymous SNPs were identified from

the 820k SNP dataset from CerealsDB (Wilkinson *et al.* 2012), which contains all

MAGIC parents.

The SNP (Adenine at position 2020 of CDS to Guanine; A2020G) causes a predicted

deleterious amino acid substitution from the ancestral serine to a glycine residue at

position 674 of the predicted primary amino acid sequence (Ser674Gly; SIFT 0.01)

(Figure 2B). A KASP marker was designed for this SNP and used to genotype the

MAGIC population. The KASP marker mapped to the same genetic position as

Ra_c109187_371 and segregated with the predicted founder effects, confirming that

TaELF3-B1 is part of the 1B GS55 QTL.

This predicted glycine at residue 674 of *TaELF3-B1*, associated with late heading, is

globally rare in wheat. In the 1000Exome data (He *et al.* 2019) 4.87% of lines have

the predicted glycine residue and in the wheat HapMap panel (Jordan *et al.* 2015),

only Rialto has the predicted glycine residue out of the 62 globally diverse lines

analysed (Supplemental Table S2). Similarly, of the 14 hexaploid varieties currently

available from the 10+ Wheat Genome project only three (Cadenza, Paragon,

ArinaLrFor) possess the non-ancestral glycine residue (Supplemental Table S2).

However, analysis of multiple datasets (see methods) found the non-ancestral

glycine674 is more common in wheat varieties with a UK pedigree (9/13 varieties)

(Supplemental Table S3). Alignment of *TaELF3* orthologues shows that the predicted

serine residue is also highly conserved across monocots (Supplemental Figure S1).

When looking for other potentially candidate sequence variation we found that within

the *ELF3-B1* promoter regions of Claire, Robigus, Paragon and Cadenza there are

two sites polymorphic for the number of repeated cytosines $((C)_n)$ located 1967 bp and

1151 bp upstream of the transcriptional start site (TSS) (Figure 2B). Variation in CT

repeat number within the 5'-UTR region of the wheat *cellulose synthase like* (*Csl*) gene

is associated with *Csl* expression and tiller number (Hyles *et al.* 2017). As Robigus

and Claire have an early heading phenotype at the *Eps-B1* locus and Cadenza a late

heading phenotype (Alvarez *et al.* 2016) it is unlikely that the promoter $(C)_n$

polymorphisms cause the changes in heading dates because there is only a difference

in two cytosines at the 1967 bp $(C)_n$ site between Cadenza and Robigus/Claire (Figure

2B). This conclusion is supported by our finding that there is no correlation between

the $(C)_n$ genotype and *TaELF3* expression (ANOVA F = 0.484, p = 0.748)

(Supplemental Figure S2). It was not possible to quantify the relative transcript

abundance of the *TaELF3-B1* homoeologue due to a lack of homoeologous SNPs in

the coding region of *TaELF3* as previously reported (Zikhali *et al.* 2016). Furthermore

TaELF3-B1 in Julius, Lancer, Mace, Norin61 and Sy Mattis seems to be inverted

compared to ArinaLrFor, CDC Landmark, CDC Stanley, Chinese Spring and Jagger.

However, as none of these lines share the same haplotype with ArinaLrFor and

Paragon (which both carry the Ser674Gly SNP) at *ELF3-B1* it is unlikely that an

inversion is the cause behind the *Eps-B1* QTL. Thus, we conclude that the Ser674Gly

SNP in *ELF3-B1* is the best candidate to underlie *Eps-B1*.

Subtelomeric introgression and inversion might underlie the *Eps-D1* heading date QTL

A subtelomeric deletion in Cadenza and Spark was previously described as the causal polymorphism underlying the *Eps-D1* locus using Spark × Rialto, Avalon × Cadenza mapping populations (Zikhali *et al.* 2016, Ochagavía *et al.* 2019). Our QTL analysis for 2012 (Supplemental Table S1) and 2014 (Table 1) shows a minor QTL overlapping this locus. In both years the Xi-19 founder genotype is associated with earlier heading (2012: -1.26 days; p = 0.015; 2014; -0.79 days; p = 0.053). However correct genetic map construction and resulting assignment of founder genotypes at this locus might be more difficult given the high-density marker region previously identified at the end of 1D. These high-density marker regions were hypothesised to be associated with introgressions (Gardner et al., 2016). Based on the previously reported deletion in Cadenza and Spark (Zikhali, Wingen, and Griffiths 2016) and the pedigree of Xi-19 (Cadenza/Rialto//Cadenza) we hypothesised that the Cadenza allele inherited from Tonic was a likely candidate for the observed QTL effects. By analysing data from recently published wheat assemblies (Walkowiak *et al.* 2020) we suggest the polymorphism is due to a subtelomeric inversion derived from introgressed material rather than a deletion. A BLAST search of the Chinese Spring *TaELF3-D1* genomic sequence against the 10+ Wheat Genome Project varieties identified high confidence matches in all varieties including Cadenza. Interestingly, the sequence retrieved from the variety Jagger was identical to Cadenza. However, the sequence identity of the Jagger and Cadenza coding sequence for TraesJAG1D01G492900.1 and TraesCAD_scaffold_026352_01G000400.1 compared to the Chinese Spring gene is much lower (96.72%) than for the remaining 1D coding sequences (>99.9%). The

Jagger and Cadenza 1D coding sequences have a greater similarity to the Chinese

Spring *TaELF3-A1* (97.50%) and Chinese Spring *TaELF3-B1* (97.88%)

homoeologues than to Chinese Spring *TaELF3-D1*.

Visualisation of the 1D chromosome for the 10+ Wheat Genome Project varieties using

the Crop Haplotypes viewer (Brinton *et al.* 2020) showed a shared haplotype block at

the distal end of chromosome 1D between Cadenza (Haplotype viewer coordinates:

482.8 Mbp – 495.4 Mbp) and Jagger (480.0 Mbp to 493.5 Mbp) where *TaELF3-D1* is

found.

The Cadenza v1.1 reference sequence has yet to be assembled into chromosomes,

but since the Crop Haplotypes viewer (Brinton *et al.* 2020) data indicated identical by

state (IBS) haplotypes between Jagger and Cadenza, we compared the distal end

structure of the Jagger 1D chromosome with Chinese Spring. The end of the Jagger

chromosome contains an inversion of approximately 2.6 Mbp relative to Chinese

Spring which includes *TaELF3-D1* (Figure 3A). Alignment of the Jagger/Cadenza

TaELF3-D1 genomic sequence with Chinese Spring *TaELF3-D1* showed some

sequence differences including a deletion within intron 2 (Figure 3B, Supplemental

Table S4). We developed a subgenome specific PCR assay detailed in Figure 3B that

amplifies the region of *TaELF3-D1* containing intron 2 and used this as a simple

marker for the 1D introgression. We confirmed that of the eight MAGIC parents, only

Xi-19 carried the Cadenza/Jagger *TaELF3-D1* allele (Figure 3C and D). As

hypothesised Spark and Tonic also carry the Cadenza *TaELF3-D1* allele (Figure 3C,

D and Supplemental Figure S3) as well as the varieties Maris Fundin and Cordiale

which show introgression signatures on the distal end of chromosome 1D (Scott *et al.*

2021). A simplified pedigree diagram shows how some of the cultivars are related

(Supplemental Figure S4). The intron 2 deletion is also present in *Aegilops speltoides*

and *Triticum timopheevii* (Supplemental Figure S5).

Given the low sequence conservation of the *TaELF3-D1* gene in Cadenza and Jagger

and the presence of the intronic deletion in *Aegilops speltoides* and *Triticum*

timopheevii, we investigated the wider phylogeny of the 1D Jagger/Cadenza region

using the 820k SNP dataset from CerealsDB. This showed that Cadenza, Xi-19 and

Spark as well as several other Tonic descendants (KWS-Podium, Cordiale, Gallant,

Duxford) and five other cultivars (Moisson, Bacanora, Tuerkis, Badger, Pavon)

clustered separately from the remaining *T. aestivum* cultivars (Supplemental Figure

S6). These lines also cluster with six Watkins lines from India (Watkins34), Yugoslavia

(Watkins352), China (Watkins141), UK (Watkins103), Cyprus (Watkins292) and

Turkey (Watkins299). *T. timopheevii* clusters closely with the above lines. Gene

alignment of *ELF3-D1* orthologues and alignment of 300 kb surrounding *ELF3-D1* also

supports an origin outside the D genome for the Jagger/Cadenza *ELF3-D1* gene

(Supplemental Figure S7, Supplemental Table S5).

Allelic variation in *TaELF3* had no detectable effect on circadian rhythms in the MAGIC parent varieties

299 We investigated if the variations at *TaELF3-B1* and *TaELF3-D1* in the MAGIC parents,

300 or other *loci,* might be associated with alterations of circadian rhythms in a manner

301 that might explain the effects on heading date. We quantified circadian rhythms in the

302 MAGIC parent varieties by measuring the period and amplitude of rhythms of delayed

303 chlorophyll fluorescence (DF) in otherwise constant light (Gould *et al.* 2009). There

were robust circadian rhythms of DF in all the MAGIC parent lines (RAE < 0.5) (Figure 4A). Hereward had the shortest circadian period (25.0 h) (Figure 4C) and Xi-19 the longest period (26.2 h) but the difference between these periods and the mean of all parents was not significant (p = 0.132) (Figure 4C). To determine whether the lack of an altered circadian phenotype in the MAGIC parent lines was the result of masking by compensatory alleles at other *loci,* we selected some of the earliest and latest heading MAGIC RILs from 2014 and used chlorophyll *a* fluorescence (PF) to measure circadian rhythms in continuous light (Supplemental Figure S8). There was no significant relationship between heading (early or late) and circadian period (p = 0.4216) (Supplemental Figure S8B) or between heading and circadian rhythm amplitude (p = 0.7385) (Supplemental Figure S8C). We detected robust rhythms of PF in all lines (RAE < 0.5) (Supplemental Figure S8D) indicating that in these lines there was no effect on circadian rhythm robustness. We then investigated whether there was any significant relationship between circadian rhythm parameters and the *ELF3-B1, ELF3-D1* or *Ppd-1* alleles associated with altered heading date. There was no significant difference between circadian period and *ELF3-B1* alleles (p = 0.542), *ELF3-D1* alleles (p = 0.521) or *Ppd-1* alleles (0.507). Similarly, there was no significant difference between circadian amplitude and *ELF3-B1* alleles (p = 0.942), *ELF3-D1* alleles (p = 0.1302) and *Ppd-1* alleles (0.966). Thus, the allelic diversity in homoeologues of *TaELF3* that was associated with alterations in heading date did not cause detectable changes in circadian rhythms.

TtELF3 has separable effects on heading date and the wheat circadian clock

Because allelic variation of *TaELF3* was associated with heading date but not

circadian rhythms in the MAGIC parents and RILs, we examined directly whether *ELF3*

affects heading and is a circadian clock component in wheat by examining tetraploid

TILLING mutant lines carrying point mutations encoding stop codons in both *TtELF3-*

A1, TtELF3-B1 or both (Alvarez *et al.* 2016) (Supplemental Figure S9). The *Ttelf3*-null

line headed first (48.7 days) with *TtELF3*-WT heading significantly (p < 0.0001) later

(56.4 days) as previously reported (Alvarez *et al.* 2016). There was no significant

difference (p = 0.98) in heading date between *Ttelf3*-Anull (51.0 days) and *Ttelf3*-Bnull

(52.4 days) suggesting that the *TtELF3-A* and *TtELF3-B* homoeologues contribute

equally to heading date. However, they were both significantly later heading than

Ttelf3-null (p < 0.0001) and significantly earlier than *TtELF3*-WT (p < 0.0001).

We found that functional *ELF3* is required for robust circadian rhythms in continuous

light and that single wild type *ELF3* homoeologues can maintain circadian function in

the absence of other functional homoeologues. There were robust circadian rhythms

of $F_{m'}$ in *TtELF3*-WT (RAE 0.16 ± 0.05), *Ttelf3*-Anull (RAE 0.14 ± 0.04) and *Ttelf3*-

Bnull (RAE 0.15 ± 0.07) but no robust circadian rhythms in *Ttelf3*-null (RAE 0.53 ±

0.19) (Figure 5B-C). The arrhythmic phenotype of *Ttelf3*-null was confirmed using DF

and leaf temperature measurements (Supplemental Figure S10A-F) and was not the

result of reduced maximum quantum yield of photosystem 2 (F_v/F_m) as this was higher

for *Ttelf3*-null for the duration of measurement (Supplemental Figure S10G-I). There

was no significant difference in circadian period of $F_{m'}$ (p > 0.05) between *TtELF3*-WT

(24.4 h ± 1.2), *Ttelf3*-Anull (24.3 h ± 0.7) and *Ttelf3*-Bnull (24.9 h ± 1.0). Similarly, there

was no significant difference (p > 0.05) in the amplitude of $F_{m'}$ oscillations between

TtELF3-WT (241 ± 53), *Ttelf3*-Anull (218 ± 50) and *Ttelf3*-Bnull (236 ± 57).

Our results demonstrate that allelic variation at individual *ELF3* homoeologues is

sufficient to alter heading date without affecting circadian rhythms, indicating that *ELF3*

has separable effects on heading date and the circadian clock in wheat.

Loss of functional *TtELF3* perturbs circadian oscillator gene expression in light/dark cycles and constant light

To further investigate the function of *ELF3* in the wheat circadian system we measured

the relative transcript abundance of orthologues of the Arabidopsis circadian oscillator

components *LHY, TOC1, Ppd-1* (*PRR37*)*, PRR73, GI, ELF3* and *LUX* in light dark

(LD) and continuous light (LL) cycles. In LD cycles all transcripts had rhythmic

expression (Figure 6A-G). Consistent with previous findings in Arabidopsis, *TtLHY* had

peak expression at dawn (Figure 6A), the *PRR* orthologues peaked during the

photoperiod (Figure 6B-C) with *TtTOC1* (Figure 6D), *TtGI* (Figure 6E) and *TtLUX*

(Figure 6F) peaking around dusk. However, the peak in *TtELF3* abundance was at the

end of the night just prior to dawn (Figure 6G), approximately antiphase to the

expression of *TtLUX* (Figure 6F). This differs markedly from Arabidopsis in which *ELF3*

and *LUX* are co-expressed near dusk (Nusinow *et al.* 2011, Huang *et al.* 2016, Ezer

et al. 2017). For all transcripts, except *Ppd-1* (Figure 6B), we detected oscillations in

relative transcript abundance in constant light, with the phasing in LL consistent with

that in LD. These oscillations of circadian transcript abundance were abolished, or of

very low amplitude, in *Ttelf3*-null (Figure 6A-G). Interestingly, the expression of *Ppd-1*

was consistently higher in *Ttelf3*-null in both LD and the first LL cycle (Figure 6B) and

the minimum level of the *Ppd-1* homologue *TtPRR73* was also higher in *Ttelf3*-null in

LD (Figure 6C) which suggests that ELF3 might act as negative regulator of *Ppd-1*

and *TtPRR73*.

There was a phase advance in peak transcript abundance of approximately 3 h in LD

in *Ttelf3*-null compared to *TtELF3-WT* (Figure 6A-G) except for *TtELF3* transcript

abundance where the peak phase was at dawn in both *TtELF3*-WT and *Ttelf3*-null

(Figure 6G). The amplitude of transcript abundance under diel conditions was also

less in the *Ttelf3*-null line compared to *ELF3*-WT (Figure 6). Together, these data

demonstrate that functional *TtELF3* is required for the maintenance of robust

oscillations of circadian oscillator gene expression in LL and for normal circadian

timing in LD.

390   *LUX* loss of function disrupts the wheat circadian clock

The antiphase expression of wheat *ELF3* and *LUX* raises the possibility that the EC

might not be formed, or functions differently to that in Arabidopsis. Previous studies in

diploid wheat have identified *TmLUX* as a circadian oscillator gene (Gawronski *et al.*

2014), so we investigated whether *LUX* also contributes to circadian rhythms in

hexaploid wheat, and whether functionality of one homoeologue is sufficient to

maintain robust oscillations. We obtained three hexaploid Japanese cultivars,

Chogokuwase, Minaminokomugi and Geurumil that have variation at *TaLUX*

associated with flowering time (Mizuno *et al.* 2012). Chogokuwase is an extra early

flowering variety and is the offspring of Geurumil and Minaminokomugi. Chogokuwase

carries predicted non-functional mutations in all three sub-genome copies of *TaLUX*

(Supplemental Figure S11). Chogokuwase also carries a *TaVRN-D1* spring allele and

a photoperiod insensitive *Ppd-D1a* allele. Geurumil has the same predicted non-

functional alleles of *TaLUX* and *Ppd-D1a* but has a winter growth habit due to lacking

the *VRN-D1* spring allele. Minaminokomugi carries the non-functional *Talux-a1* allele

but wild-type alleles of *TaLUX-B1* and *TaLUX-D1* in addition to carrying the *TaVRN-*

D1 and *Ppd-D1a* alleles (Mizuno *et al.* 2016).

Minaminokomugi maintained robust oscillations of chlorophyll fluorescence (RAE 0.16

± 0.01) and leaf temperature (RAE 0.26 ± 0.02) (Figure 7A-C). In contrast

Chogokuwase, which lacks functional copies of *TaLUX,* was arrhythmic for PF (RAE

0.56 ± 0.06) and Geurumil, which is also lacking functional *TaLUX*, had reduced

robustness of PF oscillations (RAE 0.33 ± 0.05) compared to Minaminokomugi (Figure

7A-C). Similarly, there was a decreased robustness of oscillations in leaf surface

temperature for Chogokuwase and Geurumil (RAE 0.48 ± 0.09 and RAE 0.46 ± 0.11)

which appeared to dampen to arrhythmia after the first two true LL cycles (Figure 7D-

F). We conclude that functional *TaLUX* is required for robust circadian rhythms in

hexaploid wheat.

Discussion

Wheat *ELF3* contributes to heading date and circadian

rhythms independently

The eight MAGIC parent varieties provided the opportunity to measure the likely

contribution of the *ELF3* homoeologues to heading date and circadian rhythms. Of the

MAGIC parents Brompton, Hereward, Rialto and Xi-19 have a G|G substitution variant

of *TaELF3-B1* and in addition Xi-19 has an inverted copy of *TaELF3-D1,* likely arising

from a past introgression event. These variations in *TaELF3* alleles had no discernible

impact on circadian rhythms (Figure 4) but were associated with measurable effects

on heading date in the field (Supplemental Figure S12).

Consistent with previous studies, *ELF3* is a candidate for *Eps-D1 loci* (Zikhali *et al.*

2016). We have extended the findings concerning the *Eps-D1* locus by demonstrating

that rather than being within a deletion, a copy of *ELF3-D1* is present in Cadenza (as

well as Xi-19, Spark, Cordiale, Tonic and Maris Fundin) but does carry candidate

sequence variations. *ELF3-D1* lies within an introgression region that contains an

inversion relative to the Chinese Spring D genome (Figure 3) and current data cannot

rule out other candidate genes within the introgression and/or inversion region. Based

on analysis of the 820k SNP data we find that several descendants of the cultivar

Tonic carry the putative introgression, as do several other unrelated cultivars and six

Watkins lines from India, China, UK, Cyprus, Turkey, and Yugoslavia. The

introgression is also likely present within Chinese elite germplasm (Wang *et al.* 2016).

As a result, we hypothesise that this region predates modern breeding efforts. The

same genomic region has been identified as being under selection in a collection of

bread wheat accessions dating from 1790 to 1930, though in that study it was

presumed to be a deletion rather than introgression (Przewieslik-Allen *et al.* 2021).

While the origin of the introgression is uncertain, the presence of the Jagger/Cadenza

TaELF3-D1 intronic deletion in *Ae. speltoides* and *T. timopheevii* (Supplemental

Figure S5) and the clustering outside of *Aegilops tauschii* lines (Supplemental Figure

S6 and Supplemental Figure S7) supports an origin outside of the D genome, more

closely related to the G or S genome. The increase in quantity and quality of genomic

resources for diverse accessions and wheat relative species will help in determining

the origin of this introgression.

Our data also identify *TaELF3-B1* as a candidate gene underlying the *Eps-B1* locus.

We found that a predicted glycine674 (G|G) in *TaELF3-B1,* which segregates with

Eps-B1, caused a delay in heading and flowering compared to the ancestral serine674

variant (A|A) (Supplemental Figure S12), in agreement with previous observations in

Avalon (Zikhali *et al.* 2016). It was previously concluded that G|G is the ancestral

variant (Zikhali *et al.* 2016) but our evidence suggests that it is more likely that A|A is

ancestral because the G|G allele is globally rare, present in 4.87% of lines included in

the 1000 Exome dataset, in one of the 62 wheat HapMap panel lines (Supplemental

Table S2) and in three of the 14 10+ Wheat Genome project lines (Supplemental Table

S2) and the ancestral serine residue is highly conserved, present in all sequenced

grasses. However, within UK wheat germplasm (as defined by locality on wheat GRIS)

the G|G allele is more common and is found in nine of the 13 UK lines we have

investigated (Supplemental Table S3). The later flowering G|G allele may be more

favoured in northern regions (Langer *et al.* 2014).

We found that the allelic variation at *TaELF3* had no detectable effect on circadian

rhythms in the MAGIC parents or early and late heading RILs. Epistasis between *ELF3*

and *Ppd-D1* has previously been reported for flowering time (Faure *et al.* 2012, Alvarez

et al. 2016). However, we find that variation at *Ppd-A1* has no effects on circadian

rhythms or modification of circadian phenotypes in *Ttelf3*-nulls (Supplemental Figure

S13). We also find no evidence of epistasis between *eps-B1* and *eps-D1* in the

regulation of circadian rhythms because Xi-19 has both *eps-B1* and *eps-D1* but was

not distinguishable from the rest of the population in terms of circadian behaviour.

Using loss of function lines, we demonstrate that *ELF3* regulates heading and is

required for circadian rhythms in wheat. Whilst complete loss of functional *TtELF3*

caused arrhythmia of PF (Figure 5B-E & Supplemental Figure S10G-I), leaf

temperature (Supplemental Figure S10D-F) and gene expression in LL (Figure 6), loss

of function of either the A or B homoeologue was without effect on circadian rhythms

if the other homoeologue was functional (Figure 5B-E). This contrasts with the effect

on heading, where loss of either the A or B homoeologue function resulted in an

advance in heading date relative to wildtype and the magnitude of this advance was

intermediate relative to the double null mutant (Figure 5A). These results, coupled with

the increased expression in *Ttelf3*-null of the floral promoter *Ppd-1* (Figure 6B) and

FT1 (Supplemental Figure S14)*,* suggest that wheat *ELF3* functions as a core

circadian oscillator component whilst also having a direct effect on flowering

independent of its role in the circadian oscillator (Figure 8).

Loss of functional *TtELF3* advanced circadian phase in LD. This demonstrates that

one role of *ELF3* is to contribute to the correct alignment of internal phase to external

phase, which is a primary function of circadian oscillators. The earlier phasing of gene

expression in LD cycles is similar to barley containing the *eam8*.w allele where the

peak phase of *HvCCA1, HvTOC1* and *HvGI* expression was also earlier (Faure *et al.*

2012). Similarly, the same study reported loss of rhythmic oscillations in gene

expression upon transition to LL in barley. We found that functional *TtELF3* is required

for high amplitude *TtLHY* rhythms, and that *TtELF3* loss of function mutants have

increased trough transcript abundance of *TtGI* (Figure 6E). The similarity of these

observations to those in barley and rice lines carrying *ELF3* mutations (Zhao *et al.*

2012, Boden *et al.* 2014) suggests that *ELF3* genes might have similar functions

across grasses.

Wheat PHYTOCHROME B (PHYB) and PHYC are required for accelerated heading

in long days (Chen *et al.* 2014, Pearce *et al.* 2016, Kippes *et al.* 2020, Bouché *et al.*

2022) and PHYB and PHYC are required for the induction of *Ppd*-1 expression during

night-breaks in wheat (Pearce *et al.* 2017). In Arabidopsis PHYB co-locates at sites

bound by the EC (Ezer *et al.* 2017) and in Brachypodium ELF3 appears to affect PHYC

activation of *Ppd-1* physical interaction (Gao *et al.* 2019). Together these results

support the hypothesis that wheat ELF3 may act as an integrator of light signalling

downstream of the phytochromes and upstream of Ppd-1 (Kippes *et al.* 2020) placing

it within one of the wheat photoperiod pathways (Shaw *et al.* 2020) (Figure 8).

An altered role for the evening complex (EC) in wheat

Binding of the Arabidopsis EC to target promoters requires LUX, as both ELF3 and

ELF4 lack DNA binding domains, and it is thought this might explain the similar timing

of the transcript accumulation (Nusinow *et al.* 2011). However, in wheat the maximal

abundance of *ELF3* and *LUX* transcripts is separated by 12 hours (Figure 6F and G).

This difference in the peak time of *ELF3* and *LUX* transcript abundance also occurs in

T. monococcum (Alvarez *et al.* 2016), rice (Zhao *et al.* 2012) and between *ELF3-like1*

and *LUX* in maize, sorghum and foxtail millet (Lai *et al.* 2020). This suggests that

temporal separation of *ELF3* and *LUX* expression might be a general feature of cereal

circadian oscillators. Wheat also lacks a clear orthologue of *AtELF4* and instead

contains *ELF4-like* genes (Calixto *et al.* 2015). *ELF4-like* genes might have a

conserved role because expression of a barley *ELF4-like* gene restored circadian

rhythms to an Arabidopsis *elf4-2* mutant (Kolmos *et al.* 2009). There is further

evidence for conservation of EC function in monocots because Brachypodium ELF3

binds Arabidopsis ELF4 and LUX, and can complement Arabidopsis *elf3-2* mutants

(Huang *et al.* 2017). The temporal separation of wheat *ELF3* and *LUX* transcript

abundance does not preclude the formation of an EC complex, but could suggest one

is not formed, or occurs at a different time in the diel cycle to Arabidopsis. Whether

through an EC or not, our data demonstrate that functional *ELF3* and *LUX* are required

for robust circadian rhythms in wheat (Figures 6- 8).

Whilst it is notoriously difficult to attribute plant promoter *cis*-elements to specific

patterns of expression (Harmer and Kay 2005), the difference in timing of *TaELF3* and

TaLUX expression compared to their Arabidopsis orthologues might arise because of

promoter sequence differences. Within 5000 bp upstream of the three *TaELF3*

homoeologues transcription start sites the most common motifs are linked to morning

phased expression (37 morning, 20 evening and 7 midnight) (Supplemental Table S6).

The general composition of the *TaELF3* promoters is similar to *TaLHY* (22 morning, 9

evening and 3 midnight) (Supplemental Table S7) but different to the evening

expressed *TaLUX* promoters where the most abundant motifs are associated with

evening phased expression (32) compared to morning (23) and midnight (12)

(Supplemental Table S8). In particular*, TaELF3* promoters contain Morning Element

1 (ME1) and Morning Element 2 (ME2) motifs, associated with morning phased

expression (Michael *et al.* 2008), which are also found in the promoter region of the

dawn expressed *TaLHY* homoeologues (Supplemental Table S7) but are absent in

the *AtELF3* promoter (Supplemental Table S6). The *TaELF3* promoters also contain

further morning-related motifs such as the multiple Hormone up at dawn 1 (HUD1) and

Hormone up at dawn 2 (HUD2) elements (Michael *et al.* 2008) which are less frequent

in the *AtELF3* promoter (Supplemental Table S6).

Wheat *ELF3* might be a promising breeding target

ELF3 might be a suitable target for breeding new lines with heading date better

adapted to the local environment in the practice of chronoculture (Steed *et al.* 2021).

In other species such as pea, lentil and barley, allelic variation in *ELF3* has permitted

the cultivation of these crops at different latitudes (Weller *et al.* 2012, Zakhrabekova

et al. 2012, Lundqvist 2014). Whilst complete loss of functional *ELF3* in wheat

significantly advances flowering time, severely disrupts circadian rhythms and

decreases the spikelet number (Alvarez *et al.* 2016), we have demonstrated that

altering the function of particular homoeologues of *ELF3* can affect heading without a

measurable disruption to the circadian oscillator (Figure 5) and the potential

associated yield penalties (Dodd *et al.* 2005), providing some functional copies of

ELF3 are present. It might be possible by combining different alleles of *ELF3* with

alleles of other circadian clock genes, to precisely tune flowering time suited to the

growing environment. This could be tested by measuring flowering time in selected

Cadenza TILLING lines (Krasileva *et al.* 2017) containing mutations in other circadian

clock genes as Cadenza contains both the *TaELF3-B1* Ser674Gly SNP and the 1D

introgression. In addition to the *ELF3* alleles described in this work, we have also

identified further *ELF3* alleles including the absence of *ELF3-A1* in Julius

(Supplemental Table S9) highlighting a range of *ELF3* alleles within modern wheat

cultivars which could be incorporated into breeding programmes. The variation at

ELF3 might have occurred because of the positioning of *ELF3* within the subtelomeric

region (Aguilar and Prieto 2020) which has enhanced levels of recombination which

can lead to increased number of sequence and structural variants. This has been

observed for other regions, recent studies have reported that a deletion at the distal

end of chromosome 4AL is associated with altered heat tolerance (Zhai *et al.* 2021)

and a deletion at the distal end of 5AL encompasses the *Rht12* locus which is

associated with dwarfing (Sun *et al.* 2019). Secondly, variation at *ELF3* may have been

selected by past breeding efforts, leading to a change in allele frequencies, due to

advantageous adaptation of flowering time to local climates similar to that seen for

selection of *Ppd-1* alleles based on latitudinal cline (Cockram *et al.* 2007) and altitude

(Guo *et al.* 2020). A more systematic understanding of circadian clock gene alleles

and their combinations will allow for a more targeted approach in optimising flowering

time without impacting the functioning of the circadian clock.

Materials and methods

Plant material

The eight MAGIC parents; Alchemy, Brompton, Claire, Hereward, Rialto, Robigus,

Soissons and Xi-19 were obtained from NIAB (National Institute of Agricultural Botany,

UK). The *Ttelf3*-null and *TtELF3*-WT lines were donated by Jorge Dubcovksy (Alvarez

et al. 2016) where the *Ttelf3*-null line is the progeny of bulked *elf3-null/Ppd-A1b* and

the *TtELF3*-WT line is progeny of bulked *ELF3WT/Ppd-A1b* (Table 4). By outcrossing

we selected lines that contained just the *Ttelf3-A1* mutation (herein *Ttelf3*-Anull) or just

the *Ttelf3-B1* mutation (*Ttelf3*-Bnull). The Chogokuwase, Minaminokomugi and

Geurumil lines were donated by Hidetaka Nishida (Mizuno *et al.* 2012).

Plant growth conditions

Seeds were surface sterilised by washing once in 70% (v/v) ethanol for one minute (Fischer Scientific, UK), once in autoclaved water for one minute, followed by 15 minutes in 10% (v/v) sodium hypochloride solution (Fischer Scientific, UK) and then washed a further three times in autoclaved deionised water. Seeds were sown directly onto 96 well modular trays (P G Horticulture Ltd, UK), containing a 1:1 mix of Levington M2 potting compost (Levington, UK) and vermiculite (fine, Willian Sinclair Horticulture, UK) and treated with the insecticide Intercept 70W (0.02 g L-1 soil, Bayer, Germany). Plants were grown in growth chambers (Conviron, Canada) under long days and constant temperature (16 h L: 8 h D; 200 µmol m^{-2} s^{-1}; 20 °C or 16 h L: 8 h D; 400 µmol m^{-2} s^{-1}; 22 °C) or in growth cabinets (see specific methods).

Field trials

In 2012/2013 a fully replicated yield trial of 784 F_5 MAGIC lines at the NIAB experimental farm in Cambridge, UK was scored by NIAB for the timing of Growth Stages on the Zadok's scale (Zadoks *et al.* 1974, Mackay *et al.* 2014). Each plot was assessed every two days for its growth stage based on representative plants at the centre of each plot. Once a plot reached GS55 (half of ear emerged above flag leaf ligule) the date was logged. The phenotype was assessed with growth stage measurements scored as days after sowing. In the 2013/2014 field season the same growth stage phenotypes were collected from a fully replicated yield trial of 784 F_6 MAGIC lines.

Analysis of field phenotypes

Asreml-R 3.0 (Gilmour *et al.* 1997) was used to minimise or remove spatial effects in

phenotype data due to field variation.

Marker genotypes and their respective chromosomal groupings from (Gardner *et al.*

2016) were used. The genetic map was reordered visually using the R package

mpMapInteractive (Shah 2013) to improve ordering and consider physical ordering

information from IWGSCRefSeqv1. Recombination fractions were recalculated in

mpMap (Huang and George 2011) for each chromosome using mpestrf().

Recalculating recombination fractions on a per chromosome basis with the existing

marker groupings significantly reduced computation resources required and enables

the use of finer recombination bins (r <- c(0:5/600, 2:20/200, 11:50/100) compared to

the default values. Genetic map lengths were recalculated using the Kosambi mapping

function in computemap().

To reduce computational effort a skimmed genetic map with only unique mapping

locations was used to calculate the probability that each location on a genome was

inherited from each parent and conduct QTL mapping in MPWGAIM only (Verbyla *et*

al. 2014). A marker for each unique coordinate was chosen which had the lowest

missing calls and preferably a clear BLAST hit to the IWGSCRefSeqv1 reference

sequence, which was used to assign physical mapping coordinates. Out of 18089

mapped markers 4458 were chosen for founder probability calculations.

To test and validate our pipeline and analysis tools we used them to interrogate a data

set describing flowering time in a 2011/2012 nursery of the NIAB eight Parent MAGIC

population that had previously been analysed using Bayesian Networks (Scutari *et al.*

2014). Reanalysis of this data using MPWGAIM (Verbyla *et al.* 2014) detected 13

significant QTLs for flowering time with significant founder effects (Supplemental Table

S1).

This analysis identified a major QTL (18.3% genetic variation explained, LOGP 18.33)

between the markers Kukri_c27309_590 (48.57 cM) and BS00064538_51 (56.64 cM),

defining a maximum mapping interval of 31.81 Mb – 34.23 Mb between the two

mapping bins as several markers map to the 56.64 cM bin. The mean founder effects

of the QTL predict a significant acceleration of flowering time for Soissons, which is

the only parent carrying the photoperiod insensitive *Ppd-D1a* allele (Bentley *et al.*

2013) and furthermore *Ppd-D1* maps within the QTL (33.95 Mb). This analysis of the

effect of *Ppd-D1a* on flowering time validated MPWGAIM and our QTL mapping

pipeline in MAGIC.

662 Synteny analysis between Chinese Spring and Jagger 1D

A conserved haplotype block between Cadenza and Jagger at the distal end of

chromosome 1D was identified using the Crop Haplotypes web browser (Crop

Haplotypes (crop-haplotypes.com) (Brinton *et al.* 2020). Physical locations of genes

on the Chinese Spring 1D chromosome (IWGSCRefSeqv1.1) and Jagger 1D

chromosome (PGSBv2.0) were mapped using the mapviewer v2.6.1

(https://10wheatgenomes.plantinformatics.io/mapview) and then confirmed manually

using EnsemblPlants.

671 SNP mining and analysis

820k and 35k SNP array data was obtained from the CerealsDB website (Wilkinson

et al. 2012) and the 1000Exome data from the 1000 Exomes project website

(http://wheatgenomics.plantpath.ksu.edu/1000EC/). Mapping of 35k and 820k SNPs

to the IWGSCv1.1 Ref sequence were obtained from EnsemblPlantsv51. SNPs for the

Cadenza/Jagger haplotype (chr1D: 482.3Mb – end of chromosome 1D) were

extracted and heterozygous calls set as missing. Distance matrices and phylogenetic

trees were constructed using the R package "ape" (Paradis and Schliep 2019) 100

bootstrap permutations were run also using the R package "ape".

To investigate the prevalence of the Ser674Gly SNP in a broad range of germplasm

we analysed variation data from multiple sources including the 820k SNP genotyping

array (Winfield *et al.* 2016), the wheat HapMap panel (Jordan *et al.* 2015), the 1000

wheat Exome project (He *et al.* 2019) and the 10+ Wheat Genome project (Walkowiak

et al. 2020). We found that the Ser674Gly SNP was also present in the HapMap panel

as chr1B_685645813 and within the 1000 Exome data as scaffold48561_337271.

Phylogenetic tree construction

ELF3-D1 orthologous groups were retrieved from public repositories. Coding

sequences were aligned using the MCoffee aligner on default settings

(https://pubmed.ncbi.nlm.nih.gov/16556910/). Maximum Likelihood (ML) Phylogenetic

trees were build using CLC Genomics Workbench 21.0.3. The best substitution model

was determined using the Model testing function before running Molecular

Phylogenetic analysis by Maximum Likelihood method with 100 permutations for

Bootstrap values. The *ELF3* gene from *Brachypodium distachyon* was used as an

outgroup.

Quantitative Real-Time PCR

The *TtELF3*-WT and *Ttelf3*-null lines were sown and grown in two antiphased LED

growth chambers (Conviron) under long day conditions (16h L: 8h D; 250 µmol m^{-2} s^{-}

^{1}, 20°C day: 16°C night). Sampling commenced 16 days post-sowing, the first true

leaf was sampled every 3 hours for 96 hours. After the first 24 hours of sampling the

cabinet was switched to constant light and temperature (250 µmol m^{-2} s^{-1}, 20°C). Total

RNA was extracted from leaf samples using the KingFisher Flex purification system

(Thermofisher scientific) in conjunction with the Maxwell® HT simplyRNA Kit

(Promega). RNA concentration was determined using the Nanodrop ND-1000. DNA

contamination was removed using the Arcticzyme HL-DNAse kit. RNA integrity was

checked using the Fragment AnalyzerTM (AATI, USA) before cDNA was synthesised

from extracted RNA using the High-Capacity cDNA reverse transcription kit (Applied

Biosystems). Three technical replicates of gene specific products were amplified in 10

µl reactions using Power SYBR Green Master mix on a CFX384 Touch Real-Time

PCR detection system. Gene expression levels were determined relative to the

expression of two housekeeping genes, *RP15* and *RPT5A,* that were selected using

the GeNorm alogorithm (Vandesompele *et al.* 2002) (Table 5).

For quantification of *TaELF3* transcript abundance Cadenza, Chinese Spring, Claire,

Paragon and Robigus seed was sown and grown in a Conviron growth room (16h L:

8h D; 400 µmol m^{-2} s^{-1}, continuous 22°C. After 14-days leaf samples were taken and

RNA extracted using the RNAeasy plant mini kit (Qiagen, UK) with an on column

DNAse digest (Qiagen, UK). RNA concentration was determined on the Nanodrop ND-

1000 and cDNA synthesised from 500 ng RNA using the RevertAid First Strand cDNA

synthesis kit (Thermo Scientific, UK). Technical duplicates of gene-specific products

were synthesised in 10 µl reactions using the Rotor-Gene SYBR Green PCR kit

(Qiagen, UK) on a Rotor-Gene 6000 Real-Time PCR machine (Qiagen, UK).

726   PCR of *ELF3-D1*

DNA extracted using the DNeasy plant mini extraction kit (Qiagen, UK). The presence

of *ELF3-D1* was confirmed by PCR using the *TaELF3-D1* primers listed in Table 5.

PCR was performed using the DreamTaq Green PCR kit (Thermo Scientific, UK) on

a ProFlex PCR system (Thermo Scientific, UK). PCR reaction conditions were as

follows: 3 minutes at 95ºC; 40 cycles of 30 s at 95ºC, 30 s at 68ºC, 90 s at 72ºC; 15

minutes at 72ºC.

734   Delayed chlorophyll fluorescence and prompt chlorophyll *a*

735   fluorescence imaging

Leaf samples were taken from the upper half of the first true leaf of 16-day old

seedlings (non-vernalised) grown and entrained in Conviron growth cabinets and were

cut into 3 mm by 3 mm pieces and placed on solid 0.8% (w/v) bactoagar, 0.5 x MS

Vitamin medium (M3900, Sigma-Aldrich, UK) supplemented with 0.5 µM benzyl-

aminopurine (BAP), poured in a black 96 well imaging plates (Greiner, UK). Delayed

fluorescence (Gould *et al.* 2009) was measured for 60 s in an LB985 Nightshade

(Berthold, UK) camera. Between measurements of DF light was supplied from blue

(470 nm) red (660 nm) and LED arrays at 100 µmol m^{-2} s^{-1} during the experiment.

Temperature was maintained at 20ºC.

Chlorophyll *a* fluorescence and calculations of parameters were carried out using a

CFimager (Technologica Ltd, Colchester, UK) and the image processing scripts

provided by the manufacturer. Chlorophyll fluorescence images were captured using

a Stingray F145B ASG camera (Allied Vision Technologies, UK) through an RG665

long pass filter to exclude blue light from the LEDs. When imaging under continuous

light, leaf fragments were exposed to 40 minutes of 100 µmol m^{-2} s^{-1} of blue light

followed by a saturating pulse of 6172 µmol m^{-2} s^{-1} blue light for 800 ms then 20

minutes of darkness (with non-actinic measuring light on) followed by a second

saturating pulse of 6172 µmol m^{-2} s^{-1} for 800 ms.

Circadian measures of amplitude, phase, period and relative amplitude error were

calculated using the FFT-NLLS method in Biodare2 (www.biodare2.ed.ac.uk)

(Zielinski *et al.* 2014). Where necessary data was normalised to between -1 and 1

using the normalisation feature of Biodare2.

Heading date measurements in *TtELF3* kronos lines

Seeds sown as described above and moved to long day conditions (Conviron growth

cabinet, 16 h light: 8 h dark; 400 µmol m^{-2} s^{-1}, 22°C). After two weeks seedlings

transplanted to 10 cm pots. Heading (GS55) was scored once per day.

Direct contact leaf temperature measurements

Leaf surface temperature relative to background temperature was measured using an

in-house built device and provides an alternative measure of circadian rhythms caused

by circadian regulated changes in the stomatal aperture (Dakhiya and Green 2019).

The temperature logger was validated by measuring circadian rhythms of "in-phase"

and "anti-phase" wheat leaves where peak temperature, and therefore maximum

closure, occurred during the subjective night (Supplemental Figure S15). Leaf

temperature measurements were taken from fully expanded flag leaf of 4-week-old

plants under continuous light and temperature (200 µmol m^{-2} s^{-1}, 20°C) that were

previously grown under long days (16h L: 8h D; 200 µmol m^{-2} s^{-1}, 20°C day: 15°C night)

in a growth cabinet (Conviron, UK).

Wheat was grown in 10 cm pots. Circadian measurements of leaf temperature were

taken when the pants were four-weeks old. Type-K thermocouples were attached to

the leaf such the end of the thermocouple was pressing onto the surface of the fully

expanded flag leaf. This was kept in position by positioning a paperclip onto the leaf

and fixing the paperclip to a stake. The thermocouples were connected to an Arduino

Mega using the Adafruit 1778, a 10-bit D-A converter and x20 amplifier giving a

resolution of ± 0.1°C with a DS3231 RTC added to keep time, temperature was logged

every 30 s. Background temperature was measured using unattached thermocouples.

Leaf temperatures were averaged using a 60-point moving average, background

temperature was then subtracted from leaf temperature to give leaf temperature

relative to background.

790 Acknowledgements and Funding

791 LW, GS and LT were supported by BBSRC

792 studentships BB/K011790/1, BB/M015416/1 and BB/M011194/1 awarded to MAH

793 and AARW, GS is currently supported by UKRI BBSRC grant BB/S006370/1 awarded

794   to AARW. DCR is supported by UKRI BBSRC grant BB/S002251/1 and GP-C is

795   supported by UKRI BBSRC grant BB/W001209/1, both awarded to AARW.

Contributions

798   AARW, MAH, AG, KG, LW and GS conceived this study and planned the experiments.

799   LW, GS and LJT performed the experiments and analysed data. DCR and GP-C

800   assisted with scoring heading date and sampling for qRT-PCR. AARW, LW and GS

801   wrote the first draft of the manuscript. AARW, MAH, KG, LW, GS and LJT contributed

802   to the revision of the manuscript.

Bibliography

Aguilar, M. and Prieto, P., 2020. Sequence analysis of wheat subtelomeres reveals a

high polymorphism among homoeologous chromosomes. *The Plant Genome*,

13 (3), e20065.

Alvarez, M.A., Tranquilli, G., Lewis, S., Kippes, N., and Dubcovsky, J., 2016. Genetic

and physical mapping of the earliness per se locus Eps-A m 1 in Triticum

monococcum identifies EARLY FLOWERING 3 (ELF3) as a candidate gene.

Functional & Integrative Genomics, 16 (4), 365–82.

Beales, J., Turner, A., Griffiths, S., Snape, J.W., and Laurie, D.A., 2007. A Pseudo-

Response Regulator is misexpressed in the photoperiod insensitive Ppd-D1a

mutant of wheat (Triticum aestivum L.). *Theoretical and Applied Genetics*, 115

(5), 721–733.

Bentley, A.R., Horsnell, R., Werner, C.P., Turner, A.S., Rose, G.A., Bedard, C.,

Howell, P., Wilhelm, E.P., Mackay, I.J., Howells, R.M., Greenland, A., Laurie,

D.A., and Gosman, N., 2013. Short, natural, and extended photoperiod

response in BC2F4 lines of bread wheat with different Photoperiod-1 (Ppd-1)

alleles. *Journal of Experimental Botany*, 64 (7), 1783–1793.

Boden, S.A., Weiss, D., Ross, J.J., Davies, N.W., Trevaskis, B., Chandler, P.M., and

Swain, S.M., 2014. EARLY FLOWERING3 Regulates Flowering in Spring Barley

by Mediating Gibberellin Production and FLOWERING LOCUS T Expression.

The Plant Cell, 26 (4), 1557–1569.

Bouché, F., Woods, D.P., Linden, J., Li, W., Mayer, K.S., Amasino, R.M., and

Périlleux, C., 2022. EARLY FLOWERING 3 and Photoperiod Sensing in

Brachypodium distachyon. *Frontiers in Plant Science*, 12 (January), 1–14.

Brinton, J., Ramirez-Gonzalez, R.H., Simmonds, J., Wingen, L., Orford, S., Griffiths,

S., Haberer, G., Spannagl, M., Walkowiak, S., Pozniak, C., and Uauy, C., 2020.

A haplotype-led approach to increase the precision of wheat breeding.

Communications Biology, 3 (1), 1–11.

Calixto, C.P.G., Waugh, R., and Brown, J.W.S., 2015. Evolutionary Relationships

Among Barley and Arabidopsis Core Circadian Clock and Clock-Associated

Genes. *Journal of Molecular Evolution*, 80 (2), 108–119.

Camargo, A. V., Mott, R., Gardner, K.A., Mackay, I.J., Corke, F., Doonan, J.H., Kim,

J.T., and Bentley, A.R., 2016. Determining Phenological Patterns Associated

with the Onset of Senescence in a Wheat MAGIC Mapping Population. *Frontiers*

in Plant Science, 7, 1540.

Campoli, C., Pankin, A., Drosse, B., Casao, C.M., Davis, S.J., and Von Korff, M.,

2013. HvLUX1 is a candidate gene underlying the early maturity 10 locus in

barley: Phylogeny, diversity, and interactions with the circadian clock and

photoperiodic pathways. *New Phytologist*, 199 (4), 1045–1059.

Chen, A., Li, C., Hu, W., Lau, M.Y., Lin, H., Rockwell, N.C., Martin, S.S., Jernstedt, J.

a, Lagarias, J.C., and Dubcovsky, J., 2014. PHYTOCHROME C plays a major

role in the acceleration of wheat flowering under long-day photoperiod.

Proceedings of the National Academy of Sciences of the United States of

America, 111 (28), 10037–10044.

Clavijo, B.J., Venturini, L., Schudoma, C., Accinelli, G.G., Kaithakottil, G., Wright, J.,

Borrill, P., Kettleborough, G., Heavens, D., Chapman, H., Lipscombe, J., Barker,

T., Lu, F.H., McKenzie, N., Raats, D., Ramirez-Gonzalez, R.H., Coince, A.,

Peel, N., Percival-Alwyn, L., Duncan, O., Trösch, J., Yu, G., Bolser, D.M.,

Namaati, G., Kerhornou, A., Spannagl, M., Gundlach, H., Haberer, G., Davey,

R.P., Fosker, C., Di Palma, F., Phillips, A.L., Millar, A.H., Kersey, P.J., Uauy, C.,

Krasileva, K. V., Swarbreck, D., Bevan, M.W., and Clark, M.D., 2017. An

improved assembly and annotation of the allohexaploid wheat genome identifies

complete families of agronomic genes and provides genomic evidence for

chromosomal translocations. *Genome Research*, 27 (5), 885–896.

Cockram, J., Jones, H., Leigh, F.J., O'Sullivan, D., Powell, W., Laurie, D.A., and

Greenland, A.J., 2007. Control of flowering time in temperate cereals: genes,

domestication, and sustainable productivity. *Journal of Experimental Botany*, 58

(6), 1231–1244.

Dakhiya, Y. and Green, R.M., 2019. Thermal imaging as a non-invasive technique

for analyzing circadian rhythms in plants. *New Phytologist*, 224, 1685–1696.

Dakhiya, Y., Hussien, D., Fridman, E., Kiflawi, M., and Green, R., 2017. Correlations

between Circadian Rhythms and Growth in Challenging Environments. *Plant*

Physiology, 173 (3), 1724–1734.

Dixon, L.E., Knox, K., Kozma-Bognar, L., Southern, M.M., Pokhilko, A., and Millar,

A.J., 2011. Temporal repression of core circadian genes is mediated through

EARLY FLOWERING 3 in Arabidopsis. *Current Biology*, 21 (2), 120–125.

Dodd, A.N., Salathia, N., Hall, A., Kévei, E., Tóth, R., Nagy, F., Hibberd, J.M., Millar,

A.J., and Webb, A. a R., 2005. Plant circadian clocks increase photosynthesis,

growth, survival, and competitive advantage. *Science (New York, N.Y.)*, 309

(5734), 630–633.

Ezer, D., Jung, J.H., Lan, H., Biswas, S., Gregoire, L., Box, M.S., Charoensawan, V.,

Cortijo, S., Lai, X., Stöckle, D., Zubieta, C., Jaeger, K.E., and Wigge, P.A., 2017.

The evening complex coordinates environmental and endogenous signals in

Arabidopsis. *Nature Plants*, 3 (7).

Faricelli, M.E., Valárik, M., and Dubcovsky, J., 2010. Control of flowering time and

spike development in cereals: The earliness per se Eps-1 region in wheat, rice,

and Brachypodium. *Functional and Integrative Genomics*, 10 (2), 293–306.

Faure, S., Turner, a. S., Gruszka, D., Christodoulou, V., Davis, S.J., von Korff, M.,

and Laurie, D. a., 2012. Mutation at the circadian clock gene EARLY

MATURITY 8 adapts domesticated barley (Hordeum vulgare) to short growing

seasons. *Proceedings of the National Academy of Sciences*, 109 (21), 8328–

8333.

Gao, M., Geng, F., Klose, C., Staudt, A.-M., Huang, H., Nguyen, D., Lan, H.,

Mockler, T.C., Nusinow, D.A., Hiltbrunner, A., Schäfer, E., Wigge, P.A., and

Jaeger, K.E., 2019. Phytochromes measure photoperiod in Brachypodium.

bioRxiv.

Gardner, K.A., Wittern, L.M., and Mackay, I.J., 2016. A highly recombined, high-

density, eight-founder wheat MAGIC map reveals extensive segregation

distortion and genomic locations of introgression segments. *Plant Biotechnology*

Journal, 14 (6), 1406–1417.

Gawronski, P., Ariyadasa, R., Himmelbach, A., Poursarebani, N., Kilian, B., Stein,

N., Steuernagel, B., Hensel, G., Kumlehn, J., Sehgal, S.K., Gill, B.S., Gould, P.,

Hall, A., and Schnurbusch, T., 2014. A Distorted Circadian Clock Causes Early

Flowering and Temperature-Dependent Variation in Spike Development in the

Eps-3Am Mutant of Einkorn Wheat. *Genetics*, 196 (4), 1253–1261.

Gerland, P., Raftery, A.E., Sevcikova, H., Li, N., Gu, D., Spoorenberg, T., Alkema,

L., Fosdick, B.K., Chunn, J., Lalic, N., Bay, G., Buettner, T., Heilig, G.K., and

Wilmoth, J., 2014. World population stabilization unlikely this century. *Science*,

346 (6206), 234–237.

Gilmour, A.R., Cullis, B.R., and Verbyla, A.P., 1997. Accounting for Natural and

Extraneous Variation in the Analysis of Field Experiments. *Journal of*

Agricultural, Biological, and Environmental Statistics, 2 (3), 269–293.

Gould, P.D., Diaz, P., Hogben, C., Kusakina, J., Salem, R., Hartwell, J., and Hall, A.,

2009. Delayed fluorescence as a universal tool for the measurement of

circadian rhythms in higher plants. *Plant Journal*, 58 (5), 893–901.

Griffiths, S., Simmonds, J., Leverington, M., Wang, Y., Fish, L., Sayers, L., Alibert,

L., Orford, S., Wingen, L., Herry, L., Faure, S., Laurie, D., Bilham, L., and

Snape, J., 2009. Meta-QTL analysis of the genetic control of ear emergence in

elite European winter wheat germplasm. *Theoretical and Applied Genetics*, 119

(3), 383–395.

Guo, W., Xin, M., Wang, Z., Yao, Y., Hu, Z., Song, W., Yu, K., Chen, Y., Wang, X.,

Guan, P., Appels, R., Peng, H., Ni, Z., and Sun, Q., 2020. Origin and adaptation

to high altitude of Tibetan semi-wild wheat. *Nature Communications*, 11 (1), 1–

12.

Harmer, S.L. and Kay, S.A., 2005. Positive and negative factors confer phase-

specific circadian regulation of transcription in Arabidopsis. *Plant Cell*, 17 (7),

1926–1940.

Hawkesford, M.J., Araus, J.L., Park, R., Calderini, D., Miralles, D., Shen, T., Zhang,

J., and Parry, M. a. J., 2013. Prospects of Doubling Global Wheat Yields. *Food*

and Energy Security, 2 (1), 34–48.

He, F., Pasam, R., Shi, F., Kant, S., Keeble-Gagnère, G., Kay, P., Forrest, K., Fritz,

A., Hucl, P., Wiebe, K., Know, R., Cuthbert, R., Pozniak, C., Akhunova, A.,

Morrell, P.L., Davies, J.P., Webb, S.R., Spangenberg, G., Hayes, B., Daetwyler,

H., Tibbits, J., Hayden, M., and Akhunov, E., 2019. Exome sequencing

highlights the role of wild-relative introgression in shaping the adaptive

landscape of the wheat genome. *Nature genetics*, 51 (5), 896–904.

Herrero, E., Kolmos, E., Bujdoso, N., Yuan, Y., Wang, M., Berns, M.C., Uhlworm, H.,

Coupland, G., Saini, R., Jaskolski, M., Webb, A., Goncalves, J., and Davis, S.J.,

2012. EARLY FLOWERING4 Recruitment of EARLY FLOWERING3 in the

Nucleus Sustains the Arabidopsis Circadian Clock. *the Plant Cell Online*, 24 (2),

428–443.

Hsu, P.Y. and Harmer, S.L., 2014. Wheels within wheels: The plant circadian

system. *Trends in Plant Science*, 19 (4), 240–249.

Huang, B.E. and George, A.W., 2011. R/mpMap: A computational platform for the

genetic analysis of multiparent recombinant inbred lines. *Bioinformatics*, 27 (5),

727–729.

Huang, H., Alvarez, S., Bindbeutel, R., Shen, Z., Naldrett, M.J., Evans, B.S., Briggs,

S.P., Hicks, L.M., Kay, S.A., and Nusinow, D.A., 2016. Identification of evening

complex associated proteins in arabidopsis by affinity purification and mass

spectrometry. *Molecular and Cellular Proteomics*, 15 (1), 201–217.

Huang, H., Gehan, M.A., Huss, S.E., Alvarez, S., Lizarraga, C., Gruebbling, E.L.,

Gierer, J., Naldrett, M.J., Bindbeutel, R.K., Evans, B.S., Mockler, T.C., and

Nusinow, D.A., 2017. Cross-species complementation reveals conserved

functions for EARLY FLOWERING 3 between monocots and dicots. *Plant

Direct*, 00, 1–14.

Hyles, J., Vautrin, S., Pettolino, F., MacMillan, C., Stachurski, Z., Breen, J., Berges,

H., Wicker, T., and Spielmeyer, W., 2017. Repeat-length variation in a wheat

cellulose synthase-like gene is associated with altered tiller number and stem

cell wall composition. *Journal of Experimental Botany*, 68 (7), 1519–1529.

Jordan, K.W., Wang, S., Lun, Y., Gardiner, L.-J., MacLachlan, R., Hucl, P., Wiebe,

K., Wong, D., Forrest, K.L., Sharpe, A.G., Sidebottom, C.H., Hall, N., Toomajian,

C., Close, T., Dubcovsky, J., Akhunova, A., Talbert, L., Bansal, U.K., Bariana,

H.S., Hayden, M.J., Pozniak, C., Jeddeloh, J.A., Hall, A., and Akhunov, E.,

2015. A haplotype map of allohexaploid wheat reveals distinct patterns of

selection on homoeologous genomes. *Genome Biology*, 16 (1), 48.

Kippes, N., VanGessel, C., Hamilton, J., Akpinar, A., Budak, H., Dubcovsky, J.,

Dubcovsky, J., and Pearce, S., 2020. Effect of phyB and phyC loss-of-function

mutations on the wheat transcriptome under short and long day photoperiods.

BMC Plant Biology, 20 (1), 1–17.

Kolmos, E., Nowak, M., Werner, M., Fischer, K., Schwarz, G., Mathews, S., Schoof,

H., Nagy, F., Bujnicki, J.M., and Davis, S.J., 2009. Integrating ELF4 into the

circadian system through combined structural and functional studies. *HFSP*

Journal, 3 (5), 350–366.

Krasileva, K. V, Vasquez-gross, H.A., Howell, T., Bailey, P., Paraiso, F., and

Clissold, L., 2017. Uncovering hidden variation in polyploid wheat. *PNAS*, E913–

E921.

Lai, X., Bendix, C., Yan, L., Zhang, Y., Schnable, J.C., and Harmon, F.G., 2020.

Interspecific analysis of diurnal gene regulation in panicoid grasses identifies

known and novel regulatory motifs. *BMC genomics*, 21 (1), 428.

Langer, S.M., Longin, C.F.H., and Würschum, T., 2014. Flowering time control in

European winter wheat. *Frontiers in Plant Science*, 5 (OCT), 1–11.

Lundqvist, U., 2014. Scandinavian mutation research in barley - a historical review.

Hereditas, 151 (6), 123–131.

Mackay, I.J., Bansept-Basler, P., Bentley, A.R., Cockram, J., Gosman, N.,

Greenland, A.J., Horsnell, R., Howells, R., O'Sullivan, D.M., Rose, G.A., and

Howell, P.J., 2014. An eight-parent multiparent advanced generation inter-cross

population for winter-sown wheat: Creation, properties, and validation. *G3:*

Genes, Genomes, Genetics, 4 (9), 1603–1610.

Matsubara, K., Ogiso-Tanaka, E., Hori, K., Ebana, K., Ando, T., and Yano, M., 2012.

Natural variation in Hd17, a homolog of arabidopsis ELF3 that is involved in rice

photoperiodic flowering. *Plant and Cell Physiology*, 53 (4), 709–716.

McClung, C., 2019. The Plant Circadian Oscillator. *Biology*, 8 (1), 14.

Michael, T.P., Mockler, T.C., Breton, G., McEntee, C., Byer, A., Trout, J.D., Hazen,

S.P., Shen, R., Priest, H.D., Sullivan, C.M., Givan, S.A., Yanovsky, M., Hong, F.,

Kay, S.A., and Chory, J., 2008. Network discovery pipeline elucidates conserved

time-of-day-specific cis-regulatory modules. *PLoS Genetics*, 4 (2), 14.

Mizuno, N., Kinoshita, M., Kinoshita, S., Nishida, H., Fujita, M., Kato, K., Murai, K.,

and Nasuda, S., 2016. Loss-of-function mutations in three homoeologous

PHYTOCLOCK 1 genes in common wheat are associated with the extra-early

flowering phenotype. *PLoS ONE*, 11 (10), 1–19.

Mizuno, N., Nitta, M., Sato, K., and Nasuda, S., 2012. A wheat homologue of

PHYTOCLOCK 1 is a candidate gene conferring the early heading phenotype to

einkorn wheat. *Genes & genetic systems*, 87 (6), 357–67.

Nagel, D.H. and Kay, S.A., 2012. Complexity in the wiring and regulation of plant

circadian networks. *Current Biology*, 22 (16), 648–657.

Nusinow, D.A., Helfer, A., Hamilton, E.E., King, J.J., Imaizumi, T., Schultz, T.F.,

Farré, E.M., and Kay, S.A., 2011. The ELF4–ELF3–LUX complex links the

circadian clock to diurnal control of hypocotyl growth. *Nature*, 475 (7356), 398–

402.

Ochagavía, H., Prieto, P., Zikhali, M., Griffiths, S., and Slafer, G.A., 2019. Earliness

Per Se by Temperature Interaction on Wheat Development. *Scientific Reports*, 9

(1), 2584.

Paradis, E. and Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics

and evolutionary analyses in R. *Bioinformatics*, 35 (3), 526–528.

Pearce, S., Kippes, N., Chen, A., Debernardi, J.M., and Dubcovsky, J., 2016. RNA-

seq studies using wheat PHYTOCHROME B and PHYTOCHROME C mutants

reveal shared and specific functions in the regulation of flowering and shade-

avoidance pathways. *BMC Plant Biology*, 16 (1), 1–19.

Pearce, S., Shaw, L.M., Lin, H., Cotter, J.D., Li, C., and Dubcovsky, J., 2017. Night-

break experiments shed light on the Photoperiod 1-mediated flowering. *Plant*

Physiology, pp.00361.2017.

Przewieslik-Allen, A.M., Wilkinson, P.A., Burridge, A.J., Winfield, M.O., Dai, X.,

Beaumont, M., King, J., Yang, C., Griffiths, S., Wingen, L.U., Horsnell, R.,

Bentley, A.R., Shewry, P., Barker, G.L.A., and Edwards, K.J., 2021. The role of

gene flow and chromosomal instability in shaping the bread wheat genome.

Nature Plants, 7 (2), 172–183.

Scott, M.F., Fradgley, N., Bentley, A.R., Brabbs, T., Corke, F., Gardner, K.A.,

Horsnell, R., Howell, P., Ladejobi, O., Mackay, I.J., Mott, R., and Cockram, J.,

2021. Limited haplotype diversity underlies polygenic trait architecture across 70

years of wheat breeding. *Genome Biology*, 22 (1), 1–30.

Scutari, M., Howell, P., Balding, D.J., and Mackay, I., 2014. Multiple Quantitative

Trait Analysis Using Bayesian Networks. *Genetics*, 198 (1), 129–137.

Shah, R., 2013. mpMapinteractive: Interactive components for package mpMap. R

package version 0.2.

Shaw, L.M., Li, C., Woods, D.P., Alvarez, M.A., Lin, H., Lau, M.Y., Chen, A., and

Dubcovsky, J., 2020. Epistatic interactions between PHOTOPERIOD1,

CONSTANS1 and CONSTANS2 modulate the photoperiodic response in wheat.

PLoS Genetics, 16 (7), 1–28.

Shaw, L.M., Turner, A.S., and Laurie, D.A., 2012. The impact of photoperiod

insensitive Ppd-1a mutations on the photoperiod pathway across the three

genomes of hexaploid wheat (Triticum aestivum). *The Plant Journal*, 71 (1), 71–

84.

Steed, G., Cano-Ramirez, D.L., Hannah, M.A., and Webb, A.A., 2021.

Chronoculture, harnessing the circadian clock to improve crop yield and

sustainability. *Science*, 372 (April).

Sun, H., Zhang, W., Wu, Y., Gao, L., Cui, F., Zhao, C., Guo, Z., and Jia, J., 2020.

The Circadian Clock Gene, TaPRR1, Is Associated With Yield-Related Traits in

Wheat (Triticum aestivum L.). *Frontiers in Plant Science*, 11, 285.

Sun, L., Yang, W., Li, Y., Shan, Q., Ye, X., Wang, D., Yu, K., Lu, W., Xin, P., Pei, Z.,

Guo, X., Liu, D., Sun, J., Zhan, K., Chu, J., and Zhang, A., 2019. A wheat

dominant dwarfing line with Rht12 , which reduces stem cell length and affects

gibberellic acid synthesis, is a 5AL terminal deletion line. *The Plant Journal*, 97

(5), 887–900.

Turner, A.S., Faure, S., Zhang, Y., and Laurie, D.A., 2013. The effect of day-neutral

mutations in barley and wheat on the interaction between photoperiod and

vernalization. *Theoretical and Applied Genetics*, 126 (9), 2267–2277.

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A.,

and Speleman, F., 2002. Accurate normalization of real-time quantitative RT-

PCR data by geometric averaging of multiple internal control genes. *Genome*

biology, 3 (7), RESEARCH0034.

Verbyla, A.P., George, A.W., Cavanagh, C.R., and Verbyla, K.L., 2014.

Whole-genome QTL analysis for MAGIC. *Theoretical and Applied Genetics*, 127

(8), 1753–1770.

Walkowiak, S., Gao, L., Monat, C., Haberer, G., Kassa, M.T., Brinton, J., Ramirez-

Gonzalez, R.H., Kolodziej, M.C., Delorean, E., Thambugala, D., Klymiuk, V.,

Byrns, B., Gundlach, H., Bandi, V., Siri, J.N., Nilsen, K., Aquino, C.,

Himmelbach, A., Copetti, D., Ban, T., Venturini, L., Bevan, M., Clavijo, B., Koo,

D.H., Ens, J., Wiebe, K., N'Diaye, A., Fritz, A.K., Gutwin, C., Fiebig, A., Fosker,

C., Fu, B.X., Accinelli, G.G., Gardner, K.A., Fradgley, N., Gutierrez-Gonzalez, J.,

Halstead-Nussloch, G., Hatakeyama, M., Koh, C.S., Deek, J., Costamagna,

A.C., Fobert, P., Heavens, D., Kanamori, H., Kawaura, K., Kobayashi, F.,

Krasileva, K., Kuo, T., McKenzie, N., Murata, K., Nabeka, Y., Paape, T.,

Padmarasu, S., Percival-Alwyn, L., Kagale, S., Scholz, U., Sese, J., Juliana, P.,

Singh, R., Shimizu-Inatsugi, R., Swarbreck, D., Cockram, J., Budak, H.,

Tameshige, T., Tanaka, T., Tsuji, H., Wright, J., Wu, J., Steuernagel, B., Small,

I., Cloutier, S., Keeble-Gagnère, G., Muehlbauer, G., Tibbets, J., Nasuda, S.,

Melonek, J., Hucl, P.J., Sharpe, A.G., Clark, M., Legg, E., Bharti, A., Langridge,

P., Hall, A., Uauy, C., Mascher, M., Krattinger, S.G., Handa, H., Shimizu, K.K.,

Distelfeld, A., Chalmers, K., Keller, B., Mayer, K.F.X., Poland, J., Stein, N.,

McCartney, C.A., Spannagl, M., Wicker, T., and Pozniak, C.J., 2020. Multiple

wheat genomes reveal global variation in modern breeding. *Nature*, 588 (7837),

277–283.

Wang, J., Wen, W., Hanif, M., Xia, X., Wang, H., Liu, S., Liu, J., Yang, L., Cao, S.,

and He, Z., 2016. TaELF3-1DL, a homolog of ELF3, is associated with heading

date in bread wheat. *Molecular Breeding*, 36 (12), 1–9.

Weller, J.L., Liew, L.C., Hecht, V.F.G., Rajandran, V., Laurie, R.E., Ridge, S.,

Wenden, B., Schoor, J.K.V., Jaminon, O., Blassiau, C., Dalmais, M., Rameau,

C., Bendahmane, A., Macknight, R.C., and Lejeune-Hénaut, I., 2012. A

conserved molecular basis for photoperiod adaptation in two temperate

legumes. *Proceedings of the National Academy of Sciences of the United States*

of America, 109 (51), 21158–21163.

Wilkinson, P.A., Winfield, M.O., Barker, G.L.A., Allen, A.M., Burridge, A., Coghill,

J.A., and Edwards, K.J., 2012. CerealsDB 2.0: an integrated resource for plant

breeders and scientists. *BMC Bioinformatics*, 13 (1), 219.

Winfield, M.O., Allen, A.M., Burridge, A.J., Barker, G.L.A., Benbow, H.R., Wilkinson,

P.A., Coghill, J., Waterfall, C., Davassi, A., Scopes, G., Pirani, A., Webster, T.,

Brew, F., Bloor, C., King, J., West, C., Griffiths, S., King, I., Bentley, A.R., and

Edwards, K.J., 2016. High-density SNP genotyping array for hexaploid wheat

and its secondary and tertiary gene pool. *Plant Biotechnology Journal*, 14 (5),

1195–1206.

Yao, W., Li, G., Yu, Y., and Ouyang, Y., 2018. funRiceGenes dataset for

comprehensive understanding and application of rice functional genes.

GigaScience, 7 (1), 1–9.

Zadoks, J.C., Chang, T.T., and Konzak, C.F., 1974. A decimal code for the growth

stages of cereals. *Weed Research*, 14 (6), 415–421.

Zakhrabekova, S., Gough, S.P., Braumann, I., Muller, A.H., Lundqvist, J., Ahmann,

K., Dockter, C., Matyszczak, I., Kurowska, M., Druka, A., Waugh, R., Graner, A.,

Stein, N., Steuernagel, B., Lundqvist, U., and Hansson, M., 2012. Induced

mutations in circadian clock regulator Mat-a facilitated short-season adaptation

and range extension in cultivated barley. *Proceedings of the National Academy*

of Sciences, 109 (11), 4326–4331.

Zhai, H., Jiang, C., Zhao, Y., Yang, S., Li, Y., Yan, K., Wu, S., Luo, B., Du, Y., Jin,

H., Liu, X., Zhang, Y., Lu, F., Reynolds, M., Ou, X., Qiao, W., Jiang, Z., Peng,

T., Gao, D., Hu, W., Wang, J., Gao, H., Yin, G., Zhang, K., Li, G., and Wang, D.,

2021. Wheat heat tolerance is impaired by heightened deletions in the distal end

of 4AL chromosomal arm. *Plant Biotechnology Journal*, 19 (5), 1038–1051.

Zhao, J., Huang, X., Ouyang, X., Chen, W., Du, A., and Zhu, L., 2012. OsELF3-1 ,

an Ortholog of Arabidopsis EARLY FLOWERING 3 , Regulates Rice Circadian

Rhythm and Photoperiodic Flowering. *PLoS ONE*, 7 (8), 1–10.

Zielinski, T., Moore, A.M., Troup, E., Halliday, K.J., and Millar, A.J., 2014. Strengths

and limitations of period estimation methods for circadian data. *PLoS ONE*, 9

(5).

Zikhali, M., Wingen, L.U., and Griffiths, S., 2016. Delimitation of the Earliness per se

D1 (Eps-D1) flowering gene to a subtelomeric chromosomal deletion in bread

wheat (Triticum aestivum). *Journal of Experimental Botany*, 67 (1), 287–299.

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144 # Figures

1145

1146 **Figure 1**. **Variation in time to reach GS55 across a UK wheat MAGIC population**.

1147 Time in days for individual MAGIC RILs (recombinant inbred lines) to reach GS55 for the (A) 2012/2013

1148 growing season and (B) 2013/2014 growing season. The number of days after sowing to reach GS55

1149 for the eight MAGIC parents is indicated by arrows.

1150

1151

1152

1153

1154

1155

Figure 2. *ELF3* **is a candidate gene for** *Eps-B1*.

(A) Syntenic relationships of the *Eps-B1* QTL to the *Eps-D1* QTL and the *T. monococcum Eps-A^m1*

QTL. *ELF3* highlighted in green. (B) *TaELF3-B1* gene model with exons represented by filled

rectangles, UTR's as white rectangles. The two promoter cytosine repeat polymorphisms with location

relative to the transcription start site (TSS) are indicated with the cytosine (C) repeat number for Claire,

Robigus, Cadenza and Paragon. The Ser674Gly SNP is situated within exon 4 of *ELF3-B1.*

1162

1163

1164

1165

1166

1167

1168

1169

Figure 3. A subtelomeric chromosomal introgression and inversion containing *ELF3-D1* is likely

to be the causal polymorphism underlying the *Eps-D1* QTL.

(A) Syntenic relationships between the distal end of Chinese Spring (CS) 1D chromosome (IWGSCv1.1)

and Jagger (Jag) 1D (PGSBv2.0), *TaELF3-D1* highlighted in green, each black arrow corresponds to a

gene in the forward (pointing right) or reverse (pointing left) strand. (B) the Cadenza/Jagger *ELF3-D1*

contains a deletion within intron 2 which is not present in CS, F and R refer to relative primer binding sites for PCR amplification shown in (C). (D) Agarose gel separation of PCR products from (C). A PCR product at 776bp is indicative of the presence of a subtelomeric chromosomal introgression and inversion.

1208

Figure 4. Circadian rhythms of delayed chlorophyll fluorescence in the MAGIC parents.

(A) Mean DF luminescence in counts per second (cps) normalised to -1 to 1 using Biodare2 with error bars representing standard deviation (n = 27 - 33). Circadian amplitude (B), period (C) and RAE (D) calculated using FFT-NLLS (Biodare2). Upper and lower hinges represent the first and third quartiles (25th and 75th percentiles), the middle hinge represents the median value, red triangle represents mean value and black dots represent individual replicates.

1226

1227

Figure 5. Mutation of single *ELF3* homoeologues affects heading date but not circadian

rhythms of chlorophyll *a* fluorescence but *Ttelf3*-null is arrhythmic in continuous light.

(A) GS55 for *Ttelf3*-null (red), *Ttelf3*-Anull (blue), *Ttelf3*-Bnull (purple) and *TtELF3*-WT (grey) defined

as the days after sowing (DAS) at which half of the first ear has emerged past the ligule. Upper and

lower hinges represent the first and third quartiles (25th and 75th percentiles), the middle hinge

represents the median value, yellow triangle represents mean value and black dots represent

individual replicates (n = 5). (B) Mean *Fm'* of lines shown in (A) in LL with SEM bars (n = 24). (C) RAE

where points below dashed line considered rhythmic (D) circadian period length (hours)and (E)

amplitude for genotypes inplot (B). Box jitter plots as described above, note plotted points in (C-E)

correspond to those where data was successfully fitted by FFT-NLLS model in Biodare2. Significant

differences ($p < 0.05$) calculated in R using Kruskal-Wallis test followed by post-hoc Dunn's test.

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

Figure 6. Abundance of wheat circadian clock transcripts in the *TtELF3-WT* and *ttelf3* lines in

LD and LL.

(A-G) Mean abundance of circadian oscillator transcripts (n= 3 - 5) in *TtELF3*-WT (black) and *Ttelf3*-

null (red) in a 24 h LD cycle in long day conditions (16 h light at 250 μmol m^{-2} s^{-1}, 20°C: 8 h dark 16°C,

represented by horizontal grey bar) followed by constant light and temperature (20°C) from time 24 to

96 h. Transcript abundance (ΔΔCq) is relative to *TtRP15* and *TtRPT5A* (A) *TtLHY,* (B) *Ppd-1,* (C)

TtPRR73, (D) *TtTOC1,* (E) *TtGI,* (F) *TtLUX* and (G) *TtELF3* . White bar represents light, black bar

represents darkness and grey bar represents subjective night.

Figure 7. Functional *LUX* is required for maintenance of robust circadian oscillations under LL

conditions in hexaploid wheat.

(A) Mean $F_{m'}$ for Chogokuwase (blue), Geurumil (green) and Minaminokomugi (orange) in LL (n = 16)

(B) Mean period and (C) RAE for (A) calculated using FFT-NLLS (Biodare2). Upper and lower hinges

represent the first and third quartiles (25^{th} and 75^{th} percentiles), the middle hinge represents the median

value, red triangle represents mean value and black dots represent individual replicates. (D) Mean

Chogokuwase, Geurumil and Minaminokomugi leaf temperature relative to background in LL (n = 4).

(E) mean period and (F) RAE for plot (D). Error bars are SEM, for (D) every 20^{th} point plotted for clarity.

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

Figure 8: ELF3 functions in the circadian oscillator and regulates heading of wheat

1309

Genes associated with the circadian oscillator are positioned and coloured relative to the time of day

1310

at which their transcript abundance is highest in LD. The progression of the circadian oscilaltor is

1311

indicated by the circle. Lines indicate potential regulatory links with outputs and other pathways.

1312

Pointed arrow heads represent activation and blunt arrowhead represents repression.

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

Table 1: **Summary of circadian clock associated QTLs and their mapping positions.** Output from

univariate QTL model for the NIAB 2013 and 2014 GS55 data. Gene (Mb) refers to the BLAST hit

starting coordinate of the gene's cDNA against the IWGSCRefSeqv1.1. Dist (cM) refers to the markers

genetic mapping distance in centiMorgan. Dist (Mb) refers to the BLAST hit starting coordinate of the

marker sequence against the IWGSCRefSeqv1.1 LOGP is –log10(p) which corresponds to the overall

significance of the QTLs as a measure of its strength. % var is the percentage of genetic variance

contributed by QTL. Further description of methodology and column headers can be found in (Verbyla

et al. 2014).

Circadian clock gene	Gene (Mb)	Chr	L. Marker	dist (cM)	dist (Mb)	R. Marker	dist (cM)	dist (Mb)	Year	LOGP	% var
TaELF3	685,646,575	1B	Ra_c109187_371	296.34	685,741,261	RAC875_c37075_283	297	687,715,336	2013	4.48	3.4
		1B	Ra_c109187_371	296.34	685,741,261	RAC875_c37075_284	297	687,715,336	2014	3.2	2
TaELF3	493,485,612	1D	Kukri_c29687_369	100.95	493,212,138	Kukri_rep_c69829_307	101.95	488,577,079	2014	2.32	4.1
Ppd-1	33,952,591	2D	Kukri_c27309_590	48.57	31,806,310	BS00064538_51	56.64	33,370,535	2013	20.04	17.2
		2D	Kukri_c27309_590	48.57	31,806,310	BS00064538_51	56.64	33,370,535	2014	43.92	30.2
TaZTL	625,638,942	7A	Kukri_c6676_172	207.62	610,934,248	Ku_c2990_1939	208.12	621,322,194	2013	2.87	2.6

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

 Table 2. 2013 and 2014 GS55 QTL summary for the *eps-B1 locus*. Estimated parental haplotype

1349 effects on RIL BLUPs. Abbreviations as per (Verbyla *et al.* 2014): Dist (cM) refers to the markers

1350 genetic mapping distance in centiMorgans (cM). LOGP is –log10(p) and corresponds to the overall

1351 significance of the QTL as a measure of its strength. Founder Prob: probability QTL is from founder

1352 shown (p). Strength of associations can thus be reported at the overall or founder level. Size: Size of

1353 effect of each founder allele (days to GS55).

1354

QTL	Year	Founder	Size	Founder Prob	% var	LOGP	KASP (chr1B:685645813)
Eps-B1	2013	Alchemy	-0.555	0.054	3.4	4.48	A:A
		Brompton	0.317	0.123			G:G
		Claire	-0.076	0.415			A;A
		Hereward	0.623	0.008			G:G
		Rialto	0.331	0.118			G:G
		Robigus	-0.102	0.376			A:A
		Soissons	-0.683	0.003			A:A
		Xi-19	0.053	0.426			G:G
Eps-B1	2014	Alchemy	-0.448	0.196	2	3.2	A:A
		Brompton	0.395	0.178			G:G
		Claire	-0.442	0.202			A:A
		Hereward	0.591	0.072			G:G
		Rialto	0.39	0.182			G:G
		Robigus	-0.313	0.264			A:A
		Soissons	-1.085	0.002			A:A
		Xi-19	0.453	0.15			G:G

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

Table 3. 2012 and 2014 GS55 QTL summary for the *eEps-D1 locus*. Estimated parental haplotype

effects on RIL BLUPs. Abbreviations as per (Verbyla *et al.* 2014): Dist (cM) refers to the markers

genetic mapping distance in centiMorgans (cM). LOGP is –log10(p) and corresponds to the overall

significance of the QTL as a measure of its strength. Founder Prob: probability QTL is from founder

shown (p). Strength of associations can thus be reported at the overall or founder level. Size: Size of

effect of each founder allele (days to GS55).

QTL	Year	Founder	Size	Founder Prob	% var	LOGP	PCR (Figureure 3D)
Eps-D1	2012	Alchemy	0.552	0.254	0.9	0.56	1097bp
		Brompton	0.105	0.446			1097bp
		Claire	-0.11	0.447			1097bp
		Hereward	0.021	0.49			1097bp
		Rialto	-0.044	0.477			1097bp
		Robigus	0.003	0.498			1097bp
		Soissons	0.734	0.134			1097bp
		Xi-19	-1.261	0.015			776bp
Eps-D1	2014	Alchemy	-0.058	0.472	4.1	2.32	1097bp
		Brompton	0.869	0.104			1097bp
		Claire	-1.412	0.051			1097bp
		Hereward	-0.582	0.252			1097bp
		Rialto	-0.071	0.457			1097bp
		Robigus	-0.265	0.38			1097bp
		Soissons	1.244	0.013			1097bp
		Xi-19	-0.79	0.053			776bp

1381 **Table 4: Wheat lines used in this study.** GRU refers to the germplasm resource unit at the John Innes

1382 Centre, Norwich.

Species	Cultivar	Genotype	Reference
Ae. speltoides			GRU 2140023
T. aestivum	Alchemy		(Mackay *et al.* 2014)
T. aestivum	Brompton		(Mackay *et al.* 2014)
T. aestivum	Claire		(Mackay *et al.* 2014)
T. aestivum	Hereward		(Mackay *et al.* 2014)
T. aestivum	Rialto		(Mackay *et al.* 2014)
T. aestivum	Robigus		(Mackay *et al.* 2014)
T. aestivum	Soissons		(Mackay *et al.* 2014)
T. aestivum	Xi-19		(Mackay *et al.* 2014)
T. aestivum	Cadenza		GRU WBCDB0013-PG1
T. aestivum	Jagger		GRU PANG0006
T. aestivum	Spark		GRU WBCDB0052
T. aestivum	Chinese Spring		
T. aestivum	Chogokuwase		(Mizuno *et al.* 2016)
T. aestivum	Minaminokomugi		(Mizuno *et al.* 2016)
T. aestivum	Geurumil		(Mizuno *et al.* 2016)
T. aestivum	Axona		GRU W3580
T. aestivum	Cordiale		GRU W10003
T. aestivum	Maris Fundin		GRU W1357
T. aestivum	Tonic		GRU W3572
T. monococcum			GRU T1040057
T. sphaerococcum			GRU 1210007
T. timopheevii			GRU 116002
T. turgidum	Kronos	*Ttelf3*-null line	(Alvarez *et al.* 2016)
T. turgidum	Kronos	*TtELF3*-WT	(Alvarez *et al.* 2016)
T. turgidum	Kronos	*Ttelf3*-Anull	In house
T. turgidum	Kronos	*Ttelf3*-Bnull	In house

1383

1384

1385

1386

1387

1388

1389

1390

1391 **Table 5: Primers used in this study**

Gene	Sequence 5'-3'
TtCCA1	F: CCTGGAATTGGAGATGGAGA
	R: TGAGCATGGCTTCTGATTTG
TtELF3	F: TCTCCAGATGATGTTGTCGGT
	R: CTCGAACACTTGGACAGCAAA
TtGI	F: GGTAGGTGATAGACGGCACTT
	R: GTGCTACAGATGGGATGCTTG
TtFT1	F: TGAGGACCTTCTACACACTCG
	R: ACCGGGGATATCTGTCACAAG
TtLUX	F: ACAAGCGGTTCGTGGAGG
	R: CCTGCATCCGCTTGACGTA
TtPpd-1	F: CCTGTGGACTGTCGATCTCAA
	R: CAAGGGATGGCAGCGATAATG
TtPRR73	F: TCCCGAAGTTCCTCTCTTTCC
	R: AGCGGTAGTGGCAATGACA
TtTOC1	F: GGCATGGCACTTCATTCAGTT
	R: GCACATTCATACCAGCAGGAC
TtRP15	F: GCACACGTGCTTTGCAGATAAG
	R: GCCCTCAAGCTCAACCATAACT
TtRPT5A	F: GCTGGCTCGTTCAACTGATG
	R: GGACCAAGCGTTCTGATTACTC
TaELF3-D1	F: CAACGGCGCTTCTAATCTG
	R: AAACCCGCTGCTGACTGTATA

1392