Sex Chromosome Heteromorphism and the Fast-X Effect in Poeciliids

Iulia Darolti[1], Lydia J. M. Fong[1], Benjamin A. Sandkam[2], David C. H. Metzger[1], Judith E.

Mank[1,3]

1 Department of Zoology and Biodiversity Research Centre, University of British Columbia,

Vancouver, Canada

2 Department of Neurobiology and Behavior, Cornell University, Ithaca, USA

3 Centre for Ecology and Conservation, College of Life and Environmental Sciences,

University of Exeter, Cornwall, United Kingdom

Corresponding author: Iulia Darolti, e-mail: darolti@zoology.ubc.ca

Abstract

Accelerated rates of sequence evolution on the X chromosome compared to autosomes,

known as Fast-X evolution, have been observed in a range of heteromorphic sex

chromosomes. However, it remains unclear how early in the process of sex chromosome

differentiation the Fast-X effect becomes detectible. Recently, we uncovered an extreme

variation in sex chromosome heteromorphism across Poeciliid fish species. The common

guppy, *Poecilia reticulata*, Endler's guppy, *P. wingei*, swamp guppy, *P. picta*, and para guppy,

P. parae, appear to share the same XY system and exhibit a remarkable range of

heteromorphism. This sex chromosome system is absent in recent outgroups. We combined

analyses of sequence divergence and polymorphism data across Poeciliids to investigate X

chromosome evolution as a function of hemizygosity and reveal the causes for Fast-X effects.

Consistent with the extent of Y degeneration in each species, we detect higher rates of

divergence on the X relative to autosomes, a signal of Fast-X evolution, in *P. picta* and *P. parae*,

while no change in the rate of evolution of X-linked relative to autosomal genes in *P. reticulata*.

In *P. wingei*, the species with intermediate sex chromosome differentiation, we see an

increase in the rate of nonsynonymous substitutions on the older stratum of divergence only.

We also use our comparative approach to test different models for the origin of the sex

chromosomes in this clade. Taken together, our study reveals an important role of

hemizygosity in Fast-X and suggests a single, recent origin of the sex chromosome system in

this clade.

Keywords: sex chromosome evolution, heteromorphy, hemizygosity

Introduction

Owing to their unusual inheritance pattern and hemizygosity in males, X chromosomes display many distinct evolutionary properties compared to the rest of the genome (Charlesworth et al. 1987; Vicoso and Charlesworth 2006). The strength of selection and genetic drift, as well as the role of dominance and effective population size, are expected to differ between sex chromosomes and autosomes (Kirkpatrick and Hall 2004; Mank et al. 2010; Meisel and Connallon 2013). Comparing differences in the evolution of sex-linked and autosomal loci is important for understanding patterns of mutation and selection acting across the genome.

An elevated rate of coding sequence evolution on the X chromosome relative to the autosomes, referred to as the Fast-X effect (or Fast-Z in the case of ZW systems) has been observed in a diversity of organisms, including humans (Lu and Wu 2005), primates (Stevenson et al. 2007), mice (Kousathanas et al. 2014), birds (Mank et al. 2007; Mank et al. 2010; Wright et al. 2015), *Drosophila* (Mank et al. 2010; Ávila et al. 2014; Charlesworth et al. 2018), aphids (Jaquiéry et al. 2018), spiders (Bechsgaard et al. 2019), and lepidoptera (Pinharanda et al., 2019; Sackton et al., 2014). However, the magnitude of this effect can vary substantially across species due to demographic factors, differences in mating system or regulatory mechanisms acting on the sex chromosomes (Mank et al. 2010). Notably, all the cases of Fast-X above were observed in highly heteromorphic sex chromosomes, where very few genes, if any, remain on the Y chromosome and the X is largely hemizygous in the heterogametic sex. As such, it remains unclear how early in the process of sex chromosome differentiation the Fast-X effect becomes detectible.

There are two potential causes of Fast-X. The single functional copy of the X chromosome in males leads to hemizygous exposure of genes and therefore stronger purifying selection against recessive deleterious mutations and positive selection for recessive beneficial ones expressed in males (Charlesworth et al. 1987). This adaptive cause of Fast-X is mainly expected in species with heteromorphic sex chromosomes and highly degenerated Y chromosomes, as a large proportion of X-linked genes will be hemizygous in males. Alternatively, in every male and female pair, there are only three X chromosomes to four copies of each autosome, although this varies substantially based on mating system and type of heterogamety (Vicoso and Charlesworth 2009; Mank et al. 2010; Wright and Mank 2013; Wright et al. 2015). This

reduced effective population size of the X relative to the autosomes diminishes the relative

power of selection on the X, potentially leading to both greater genetic drift and fixation of

weakly deleterious mutations on the X chromosome (Charlesworth et al. 1987). This non-

adaptive cause of Fast-X does not necessarily require male hemizygosity, just recombination

suppression between the X and Y chromosomes and differences in effective population size

between X-linked and autosomal loci, and could apply to X loci where the Y copy is expressed

and remains functional.

Recently, we uncovered a case of extreme variation across Poeciliids in the rate of sex

chromosome degeneration and dosage compensation. The same chromosome pair acts as the

XY system in the common guppy, *Poecilia reticulata*, its sister species Endler's guppy, *P. wingei*,

the more distantly related swamp guppy, *P. picta,* and the para guppy, *P. parae*, all of which

last shared a common ancestor roughly 20 million years ago (Darolti et al. 2019; Sandkam et

al. 2021). Notably, *P. latipinna* and *G. holbrooki*, recent outgroup species to this clade, do not

show evidence of a sex chromosome system on the same chromosome (Darolti et al. 2019),

and therefore we hypothesized based on parsimony that the sex chromosome system arose

once, roughly 20 mya (supplementary Fig. S1, Parsimony Model). Since this origin, the sex

chromosomes of *P. reticulata* and *P. wingei* have remained largely homomorphic, although *P.*

wingei shows greater divergence in a limited region (Darolti et al. 2019). In contrast, the *P.*

picta and *P. parae* sex chromosomes are completely nonrecombining (with the exception of a

small pseudoautosomal region at the distal end, 20-21Mb, which comprises ~6% of the

chromosome) and the Y chromosome has undergone substantial degeneration, leaving most

of the X chromosome in males effectively hemizygous (Darolti et al. 2019; Sandkam et al.

2021). Moreover, *P. picta* and *P. parae* exhibit complete dosage compensation (Darolti et al.

2019; Metzger et al. 2021), which is predicted to accentuate the Fast-X effect (Charlesworth

et al. 1987; Mank et al. 2010). The range of sex chromosome differentiation present in this

clade allows us to investigate patterns of X chromosome coding sequence evolution and to

test causes underlying Fast-X effects.

In contrast to the Parsimony Model, it has recently been proposed that the sex chromosomes

of *P. reticulata* represent a recent turnover event, and that the highly diverged *P. picta* and *P.*

parae system is in fact ancient (Charlesworth et al. 2021). This Turnover Model

(supplementary Fig. S1) posits that the *P. picta* X chromosome arose well before the

immediate ancestor of *P. parae, P. picta*, *P. reticulata* and *P. wingei* (Fig. 1 from Charlesworth
et al. 2021), and that the almost complete loss of Y gene content and evolution of dosage
compensation observed in *P. picta* occurred well before the species split. This model supposes
that the homomorphism observed in *P. reticulata* and *P. wingei* represents a turnover event,
where the current guppy X and Y chromosomes arose from the ancestral X of *P. picta*. Sex
chromosome turnovers are frequent in many animal groups (Bachtrog et al. 2014), so the
model is plausible. Disentangling the evolutionary origin of this sex chromosome pair is
important for understanding the causes of the observed heterogeneity in sex chromosome
differentiation patterns and dosage compensation in this clade.

The unique molecular and evolutionary signatures that accumulate on sex chromosomes can
be detected even after they revert to being autosomes. For example, the *Drosophila* dot
chromosome was ancestrally a highly differentiated sex chromosome that reverted to an
autosome with the emergence of Drosophilidae, and yet it still maintains many of the unique
characteristics of a differentiated X chromosome, such as a feminizing effect, non-random
gene content and a chromosome-specific gene expression regulatory mechanism (Vicoso and
Bachtrog 2013). This means that if the Turnover Model is true, Fast-X patterns, evolving over
long periods of time in the distant ancestor of *P. picta* would be evident in internal branches
of the phylogeny and should also remain detectible on the X chromosome in *P. reticulata* and
P. wingei after turnover. Furthermore, the Turnover Model requires additional turnover
events in several other taxa within the clade that do not share the same sex chromosome as
P. picta (supplementary Fig. S1), and we might also expect these species to exhibit ancient
patterns of Fast-X on the chromosome, which is fully autosomal. Our comparative dataset
allows us to also test these alternative hypotheses for the origin of the guppy sex
chromosomes.

Using a combination of sequence divergence, polymorphism and expression data analyses
across Poeciliid species, we estimate rates of gene sequence evolution across the genome and
assess the presence of Fast-X evolution in each system. Consistent with the extent of Y
degeneration, we find significantly higher rates of X coding sequence evolution compared to
the autosomes, a signal of Fast-X effect, in *P. picta* and *P. parae*, potentially accelerated by the
evolution of chromosome-wide dosage compensation in these species. The absence of a

consistent Fast-X evolution in *P. reticulata*, *P. wingei*, *P. latipinna*, and *G. holbrooki*, as well as
in the ancestral branch to *P. reticulata* and *P. picta*, supports the Parsimony Model.

Results

Fast-X evolution in the heteromorphic sex chromosomes of *P. picta* and *P. parae*

We first assessed the strength of Fast-X in our study species with heteromorphic sex
chromosomes. The extensive X chromosome hemizygosity of *P. picta* and *P. parae* males is
expected to accentuate the strength of Fast-X evolution in these species (Vicoso and
Charlesworth 2006), as is the mechanism of complete X chromosome dosage compensation
(Mank et al. 2010). Indeed, our analysis in *P. picta* and *P. parae* revealed a higher rate of
nonsynonymous substitutions for X-linked loci, excluding genes in the pseudoautosomal
region (PAR) (see methods), compared to the autosomes (1,000 replicates permutation test,
$p < 0.001$; supplementary Table S1). Both species also exhibit a significantly elevated rate of
divergence (d_N/d_S) on the X chromosome relative to the rest of the genome (Fig. 1C, D),
consistent with a signal of Fast-X evolution (Fig. 2; supplementary Fig. S2) measured as the
ratio of the rate of nonsynonymous over synonymous substitutions for the X chromosome
relative to that for the autosomes. Our results thus indicate that the Fast-X signal is present in
both *P. picta* and *P. parae*.

Absence of Fast-X effect in species with homomorphic sex chromosomes

We next asked whether patterns of coding sequence evolution are different between the
species with heteromorphic sex chromosomes, *P. picta* and *P. parae*, and the species with
homomorphic sex chromosomes, *P. reticulata* and *P. wingei*. Previous work has indicated that
P. reticulata and *P. wingei* share the same male-heterogametic sex chromosome system
(Nanda et al. 2014; Morris et al. 2018; Darolti et al. 2019). Analyses of coverage differences
between males and females indicate that Y degeneration is restricted to the distal end of the
sex chromosomes, in a region ancestral to *P. reticulata* and *P. wingei* (designated as Stratum
I) (Wright et al. 2017; Darolti et al. 2019; Darolti et al. 2020; Fraser et al. 2020; Qiu et al. 2022),
and suggest that Y degeneration is slightly more exaggerated in *P. wingei* compared to *P.
reticulata*. This nonrecombining region coincides with the previously mapped location of the
sex determining region in *P. reticulata* (Winge 1922; Winge 1927; Winge and Ditlevsen 1947;

Traut and Winking 2001; Tripathi et al. 2009) and is also found across six natural guppy

populations from Trinidad (Almeida et al. 2021).

We therefore assessed the signal of Fast-X evolution for genes in Stratum I (20-26Mb on *P.*

reticulata chromosome 12; 17-20Mb on *P. wingei* chromosome syntenic to the guppy sex

chromosome). In *P. reticulata*, the rate of divergence for X-linked genes in Stratum I was not

significantly different than that for autosomal genes (Fig. 1A; supplementary Table S1). To

exclude the possibility that we were lacking power in our analysis due to the small number of

genes identified in Stratum I, we reanalyzed *P. reticulata* using Ensembl coding sequences

instead of our *de novo* generated transcripts. We were able to recover more than twice as

many X-linked loci, however the estimates of divergence remain the same as those based on

de novo transcripts (supplementary Fig. S3).

In *P. wingei*, genes in Stratum I show significantly higher rates of nonsynonymous and of

synonymous substitutions (supplementary Table S1), however the overall rate of divergence

is similar between the autosomes and Stratum I (Fig. 1B). *P. wingei* therefore shows an

intermediate pattern of Fast-X evolution, not as accentuated as in *P. picta* but more evident

than in *P. reticulata* (Fig. 2; supplementary Fig. S2). Recent work has suggested that the *P.*

wingei X and Y chromosomes are somewhat more differentiated from each other compared

to those of *P. reticulata* (Darolti et al. 2019), and there is also evidence for this from previous

cytogenetic work (Nanda et al. 1992; Nanda et al. 2014). This greater divergence might explain

the difference in d_N/d_S estimates for X-linked genes in Stratum I that we observe between

these two species, though the effect is negligible. However, it should be noted that following

filtering (see methods), only a handful of *P. wingei* genes remained in Stratum I, and as such

our statistical power to detect a significant Fast-X signal in this species is reduced.

If the ancestor of the guppy and *P. picta* had an almost completely degenerated sex

chromosome system and had evolved dosage compensation (as suggested in the model by

Charlesworth et al. 2021), then a signal of Fast-X evolution would have also accumulated prior

to the species split. The molecular and evolutionary signatures that accumulate on sex

chromosomes remain observable even when they revert to being autosomes (Vicoso and

Bachtrog 2013). As such, if the guppy sex chromosomes represent a turnover event, we would

expect the pattern of Fast-X that we observed in *P. picta* and *P. parae* to still be observed

throughout the *P. reticulata* and *P. wingei* sex chromosomes outside of Stratum I. Thus, we

next estimated rates of sequence divergence for the remainder of the X chromosome,

excluding genes in Stratum I and genes on the PAR (see methods), as the high recombination

events in the PAR can alter rates of evolution and polymorphism (Otto et al. 2011). Although

the Fast-X effect was >1 in *P. wingei* (Fig. 2; supplementary Fig. S2), the rate of mutation and

mean d_N/d_S on the X chromosome were not different from those on the autosomes in either

P. reticulata or *P. wingei* (Fig. 1A, B; supplementary Table S1). This finding is consistent with

the limited Y degeneration observed in these species (Wright et al. 2017; Darolti et al. 2019),

and does not support the Turnover Model.

Previous comparisons of replicate *P. reticulata* natural population have shown evidence for

greater X-Y divergence in replicate upstream, low predation populations relative to their

downstream, high predation populations pair within Quare, Aripo and Yarra rivers (Wright et

al. 2017; Almeida et al. 2021). This pattern of divergence has occurred over a relatively short

time span, at some point after the colonization of Trinidad during the last glacial maximum.

Therefore, we investigated whether we can detect higher rates of X chromosome to autosome

coding sequence evolution in upstream compared to downstream populations. We found a

significantly elevated d_N/d_S on the X chromosome, excluding genes on the PAR and those on

Stratum I, compared to the autosomes in the Yarra upstream population only (supplementary

Tables S2-4), and across all rivers we did not find a consistently stronger Fast-X effect in the

upstream relative to the downstream populations (Fig. 3). Compared to the other rivers, the

upstream population of the Yarra river showed the greatest increase in Y divergence relative

to the downstream population, measured by both male:female F_{ST} and male-specific SNPs

(Almeida et al. 2021). X-Y divergence between upstream and downstream populations is

greater in Yarra compared to Quare and Aripo, and this might explain why a pattern of Fast-X

effect in the upstream population was recovered in Yarra alone.

Ancestral Fast-X

If sex chromosome divergence and dosage compensation occurred before the split of *P. picta*

from the *P. latipinna* or the *Gambusia* clade, we would expect that an elevated rate of

sequence divergence would be seen for X-linked relative to autosomal loci in the branch

immediately ancestral to the *P. picta, P. parae, P. reticulata* and *P. wingei* clade. We ran

separate phylogenetic analyses with different outgroup species, however neither recovered

any difference in the overall rate of divergence or the rates of either nonsynonymous or
synonymous substitutions between the autosomal and sex-linked gene categories (Table 1,
Fig. 2). This indicates the absence of Fast-X signal in the branch that is ancestral to *P. reticulata*,
P. wingei and *P. picta*, which is inconsistent with the Turnover model and the hypothesis of an
ancestral degenerated sex chromosome system.

It has been proposed that the large-scale Y gene loss observed in *P. picta* would have taken
considerable amounts of time, much more than the time since *P. picta* split from *P. reticulata*
(Charlesworth et al. 2021), estimated at ~20 mya (Meredith et al. 2010; Rabosky et al. 2018).
If that is the case, the Turnover Model also requires additional sex chromosome turnover
events in *P. latipinna*, an outgroup species to the *P. reticulata*–*P. wingei* which has split from
P. parae only ~26 mya (Meredith et al. 2010; Rabosky et al. 2018), if not also in the more
distantly related *G. holbrooki* (supplementary Fig. S2). We therefore estimated sequence
divergence in the outgroups, *P. latipinna* and *G. holbrooki*, both species in which the guppy
chromosome 12 has not been implicated as the sex chromosome, and found no evidence of
elevated rates of evolution on the chromosome syntenic to the guppy X relative to the rest of
the genome in either species (supplementary Table S1). This result, together with our findings
in *P. reticulata* and *P. wingei* supports the Parsimony Model.

Polymorphism and the underlying causes of Fast-X
A higher d_N/d_S on the X chromosome relative to the rest of the genome could be the result of
both increased genetic drift (Charlesworth et al. 1987; Mank et al. 2010) and increased efficacy
of selection (Vicoso and Charlesworth 2009). Therefore, we used sequence and polymorphism
data together to test for the causes of elevated Fast-X evolution in *P. picta* and *P. parae*. The
McDonald-Kreitman test contrasts the number of nonsynonymous and synonymous
substitutions with polymorphisms, where an excess of nonsynonymous substitutions relative
to polymorphisms is indicative of positive selection (McDonald and Kreitman 1991). Using this
test, we detected no X-linked genes with signatures of positive selection in any of the Poeciliid
species. However, the McDonald-Kreitman test is very conservative, as it is restricted to genes
with sufficient numbers of substitutions and polymorphisms (Begun et al. 2007; Andolfatto
2008), and as such our analysis was limited to a few X-linked contigs.

To increase our statistical power to detect signatures of positive selection, we also used the
direction of selection (DoS) test which is less sensitive to low counts (Stoletzki and Eyre-Walker
2011). For each contig, DoS calculates the difference between the proportion of
nonsynonymous substitutions and the proportion of nonsynonymous polymorphisms, where
a positive DoS indicates adaptive evolution (Stoletzki and Eyre-Walker 2011). Using this
approach, we recovered more genes with signatures of positive selection, however the X
chromosome was not enriched in these genes compared to the rest of the genome in any of
the species (supplementary Table S6).

Since our results did not identify positive selection as the main force underlying the observed
Fast-X effect in *P. picta* and *P. parae*, we next used polymorphism data alone to test for a signal
of purifying selection acting on the X chromosome. A higher rate of nonsynonymous to
synonymous polymorphisms on the X relative to the autosomes is indicative of reduced
efficacy of selection to remove mildly deleterious mutations, while a lower rate suggests
purifying selection resulting from hemizygous exposure of deleterious mutations in males. Our
results show a lower rate of nonsynonymous polymorphisms on the X chromosome in *P. picta*
and *P. parae* only (Table 2, 1,000 replicates permutation tests $p = 0.04$ for *P. picta* and $p = 0.03$
for *P. parae*; supplementary Table S5), suggesting that the observed Fast-X effect in these
species is influenced by purifying selection acting to remove deleterious mutations on the X
(Charlesworth et al. 1987; Vicoso and Charlesworth 2009).

Discussion

The role of hemizygosity in Fast-X
We observe a clear pattern of Fast-X in *P. picta* and *P. parae*, as shown through the
significantly higher rate of nonsynonymous substitutions for X-linked loci (supplementary
Table S1) and significantly elevated rate of divergence on the X chromosome relative to the
rest of the genome (Fig. 1C, D; Fig. 2), consistent with the extensive Y chromosome
degeneration in this species. In contrast, we did not observe significant patterns of Fast-X
evolution in either *P. reticulata* or *P. wingei*. There is little evidence of loss of Y gene coding
sequence in either of these species (Darolti et al. 2019; Darolti et al. 2020), and very few, if
any, genes are hemizygously expressed in males, even in the older Stratum I. Interestingly,
the *P. wingei* Y is somewhat more diverged than the Y in *P. reticulata* (Nanda et al. 1992;

Nanda et al. 2014; Darolti et al. 2019), and we observe a slight, though non-significant Fast-X

effect in Stratum I in the former. We also observe an effect in the Yarra upstream population,

which shows the greatest X-Y divergence across natural populations in Trinidad (Almeida et

al. 2021). All this points to the key role of sex chromosome hemizygosity in driving Fast-X

evolution.

Some previous work has failed to detect X-Y divergence based on male:female coverage

differences in Stratum I of *Poecilia reticulata* (Bergero et al. 2019; Charlesworth et al. 2020;

Kirkpatrick et al. 2022). However, studies that have used similar genomic methods across a

range of datasets all recovered patterns consistent with this region (Darolti et al 2019; Wright

et al. 2017; Almeida et al. 2020; Fraser et al. 2020; Sigeman et al. 2021), as have studies using

male-specific sequence (Morris et al. 2018; Almeida et al. 2020; Darolti et al. 2020). A recent

methodological analysis revealed that the detection of Stratum I in *P. reticulata* is largely

dependent upon the reference genome used (Darolti et al. 2022), with studies using reference

genomes from distant populations or even different species less able to detect subtle patterns

of divergence that characterize the Y chromosome in this species. Our detection of a slight,

but non-significant pattern of Fast-X in Stratum I of the Yarra *P. reticulata* population as well

as in the same region of the *P. wingei* X chromosome gives further weight to the presence of

this Stratum.

There are two potential causes of Fast-X evolution. Male heterogamety leads to hemizygous

exposure of genes resulting in stronger purifying selection against recessive deleterious

mutations and positive selection for recessive beneficial ones expressed in males

(Charlesworth et al. 1987). Alternatively, the reduced effective population size of the X relative

to the autosomes diminishes the relative power of selection on the X, potentially leading to

non-adaptive causes of Fast-X (Charlesworth et al. 1987) which do not necessarily require

male hemizygosity, just recombination suppression between the X and Y chromosomes.

In addition to hemizygosity and effective population size, theory predicts that complete

dosage compensation, whereby balance in expression between the sex chromosomes and the

autosomes is restored, may facilitate a more pronounced Fast-X effect (Charlesworth et al.

1987; Mank et al. 2010). Under this theory, in systems with incomplete dosage compensation,

where only a subset of the genes are compensated for but overall expression for X-linked

genes is reduced compared to the autosomes in males, X-linked beneficial mutations may have lower expression and weaker phenotypic effects in males, potentially limiting the Fast-X effect (Charlesworth et al. 1987; Mank et al. 2010). *P. picta* and *P. parae* have evolved chromosome-wide dosage compensation, seemingly through a mechanism involving the hyperexpression of the single X in males (Darolti et al. 2019; Metzger et al. 2021), and this may have contributed to the observed signature of Fast-X evolution.

The role of hemizygosity and dosage compensation we observe in *P. picta* and *P. parae* Fast-X suggest an adaptive mechanism, and our polymorphism analysis is consistent with greater purifying selection acting in males to remove recessive deleterious variation. However, although we were unable to differentiate adaptive and non-adaptive causes of Fast-X in our polymorphism data, polymorphism estimates are sensitive to demographic fluctuations (Tajima 1989; Pool and Nielsen 2007) and, thus, it is difficult to determine to what extent the Fast-X pattern in *P. picta* is adaptive using polymorphism data alone. Future work identifying true X-hemizygous loci in males and analyzing their patterns of sequence evolution may prove more revealing.

Differentiating between models of sex chromosome origin

Two models have been proposed for the origin of the sex chromosome system in this group. The Parsimony Model posits that the sex chromosomes arose once roughly 20 mya (supplementary Fig. S1) in the ancestor of the species that all share it and experienced different rates of Y decay in different sub-clades (Darolti et al. 2019), possibly exacerbated by the evolution of complete dosage compensation in the immediate ancestor of *P. picta* and *P. parae* (Lenormand et al. 2020; Metzger et al. 2021; Sandkam et al. 2021). The Turnover Model proposes that the sex chromosomes of *P. reticulata* represent a recent turnover event, and that the highly diverged *P. picta* and *P. parae* system arose well before the immediate ancestor of *P. parae, P. picta*, *P. reticulata* and *P. wingei*, and large-scale Y degeneration and complete dosage compensation occurred before the most recent common ancestor of the four species (Charlesworth et al. 2021). Because the molecular signatures that accumulate on sex chromosomes remain observable even when they revert to being autosomes (Vicoso and Bachtrog 2013), we were able to critically test these alternative models with several different approaches.

First, if the guppy sex chromosomes represent a turnover event, we would expect the pattern

of Fast-X in *P. picta* and *P. parae* to also be observed throughout the *P. reticulata* and *P. wingei*

sex chromosomes outside of Stratum I, as the majority of it would be expected to have

accumulated prior to turnover. We failed to recover signatures of Fast-X in *P. reticulata* and *P.*

wingei that would be consistent with the Turnover Model, even after investigating multiple

datasets (based on *de novo* transcripts of a lab population, replicate wild populations, and

Ensembl coding sequences).

Second, the Turnover Model also requires additional sex chromosome turnover events in the

outgroup species to the *P. reticulata–P. wingei* clade, and we would expect a Fast-X signal at

least in the close outgroup *P. latipinna* if not also in *G. holbrooki*, where the proposed ancient

X chromosome is presumably now fully autosomal. We did not observe Fast-X in either of

these species.

Finally, if the sex chromosomes were ancient, we would expect a branch-specific signature of

Fast-X in the ancestral branch preceding the split of *P. picta, P. parae, P. reticulata* and *P.*

wingei. Instead, our results support the Parsimony Model, and suggest that the sex

chromosomes originated in the immediate ancestor of the *P. picta–P. parae–P. reticulata–P.*

wingei clade and divergence occurred more rapidly in the common ancestor of *P. picta* and *P.*

parae, potentially accelerated by the evolution of complete X chromosome dosage

compensation in this clade, which reduces selection to maintain Y chromosome expression for

dosage sensitive genes (Metzger et al. 2021).

Concluding Remarks

Taken together, our comparative analyses of divergence and polymorphism data across

poeciliids reveal that the degree of Fast-X evolution follows the extent of X hemizygosity in

males, emphasizing the important role of male hemizygosity in this evolutionary process. Our

results across the broader clade are consistent with a recent origin of the sex chromosome

system, as opposed to an ancient origin with turnovers. This indicates that patterns of Fast-X

evolution can accumulate rapidly in the evolution of sex chromosomes.

Materials and Methods

Sample collection and sequencing

We have previously obtained tissue samples, extracted and sequenced RNA from the tails of three males and three females of *P. reticulata*, *P. wingei*, *P. picta*, *P. parae, P. latipinna* and *G. holbrooki* (BioProject IDs PRJNA353986, PRJNA528814, PRJNA741270) (Wright et al. 2017; Darolti et al. 2019; Metzger et al. 2021). *P. reticulata* samples were obtained from our outbred laboratory population originating from the Quare River in Trinidad (Kotrschal et al. 2013). *P. wingei* samples were collected from our laboratory population established from a strain maintained by a UK fish fancier. *P. picta* and *P. parae* samples were acquired from Guyana and Suriname, while *P. latipinna* and *G. holbrooki* samples were obtained in Florida. All samples were collected in accordance with national and institutional ethical guidelines.

We extracted RNA from each sample using the Qiagen RNeasy Kit, following the instructions of the manufacturer. Library preparation and sequencing were performed at the University of Oxford Wellcome Centre for Human Genetics and Genome Québec, following standard Illumina protocols and using the Illumina HiSeq 4000 and NextSeq platforms. The data were quality assessed using FastQC v0.11.3 (www.bioinformatics.babraham.ac.uk/projects/fastqc) and trimmed with Trimmomatic v0.36 (Bolger 2014), removing adaptor sequences, reads with an average Phred score < 15 in a sliding window of four bases, reads with leading or trailing bases with a Phred score < 3 and reads shorter than 50bp following trimming.

Identifying orthogroups

For each species, we mapped RNA-seq reads to a previously constructed species-specific female *de novo* genome assembly (Wright et al. 2017; Darolti et al. 2019; Sandkam et al. 2021) using HISAT2 v2.0.4 (Kim et al. 2015), with the exception of *P. latipinna* for which a male genome assembly was used, and a non-redundant set of transcripts in GTF file format was constructed using StringTie v1.2.4 (Pertea et al. 2015). We filtered the transcripts for non-coding RNA (ncRNA) by extracting transcript sequences with BEDtools getfasta (Quinlan and Hall 2010) and removing transcripts with a BLAST hit to ncRNA sequences from *Poecilia formosa* (PoeFor_5.1.2), *Oryzias latipes* (MEDAKA1), *Gasterosteus aculeatus* (BROADS1) and *Danio rerio* (GRCz10) from Ensembl 104 (Flicek et al. 2014). Only genes with positional information on chromosomal fragments were kept for further analyses and the longest

isoform for each target gene was selected. Lastly, we applied a minimum expression filter of

2 RPKM in at least half of the samples of each sex (minimum of two out of three individuals

of either sex), resulting in 13,306 *P. reticulata*, 15,089 *P. wingei,* 13,156 *P. picta*, 24,446 *P.*

parae, 14,468 *P. latipinna* and 21,861 *G. holbrooki* genic sequences.

We obtained coding sequences from the outgroup species *P. formosa* (PoeFor_5.1.2),

Xiphophorus maculatus (Xipmac4.4.2) and *O. latipes* (MEDAKA1) from Ensembl 104 and

extracted the longest isoform for each gene. Separately for each of our target species, we

determined orthology across target and outgroup sequences using reciprocal BLASTn v2.7.1

(Altschul et al. 1990) with an e-value cut-off of $10e^{-10}$ and a minimum percentage identity of

30%. For genes with multiple blast hits, we chose the top hit based on the highest BLAST

score. Our analysis resulted in 7,296 *P. reticulata*, 7,253 *P. wingei*, 7,786 *P. picta*, 9,171 *P.*

parae, 7,251 *P. latipinna* and 6,477 *G. holbrooki* orthogroups (four-way 1:1 orthologs).

We tested the robustness of the 1:1 ortholog datasets resulting from the reciprocal BLASTn

approach by separately inferring orthologous groups using OrthoFinder (Emms and Kelly

2015). More than half of the orthogroups recovered using the OrthoFinder approach are

shared with the reciprocal BLASTn approach (supplementary Fig. S4). Restricting the

sequence divergence analysis (detailed below) to only orthogroups that are shared between

the two approaches revealed similar estimates of rates of divergence to the original estimates

based on all orthogroups identified using the reciprocal BLASTn approach (supplementary Fig.

S5). We therefore concluded that the reciprocal best-hit datasets were appropriate to use in

this case.

Estimating sequence divergence across orthogroups

For each contig of each orthogroup, we obtained open reading frames using *O. latipes*

(MEDAKA1) protein-coding sequences from Ensembl 104 and BLASTx v2.3.0 with an e-value

cut-off of $10e^{-10}$ and a minimum percentage identity of 30%, excluding orthogroups without

BLASTx hits or valid protein-coding sequences. We aligned orthologous gene sequences with

PRANK v170427 (Löytynoja and Goldman 2008), using the rooted tree (((Target species, *P.*

formosa), *X. maculatus*), *O. latipes*), where the target species was in turn *P. reticulata*, *P.*

wingei, *P. picta*, *P. parae*, *P. latipinna* and *G. holbrooki*. Alignments were then filtered to

remove gaps.

To avoid false positive signals of adaptive evolution, poorly aligned or error-rich regions were masked with SWAMP (Harrison et al. 2014). We ran SWAMP twice, first using a threshold of six nonsynonymous substitutions in a window size of 15 codons, and second using a threshold of two and a window size of five. This approach eliminates sequencing errors that cause short stretches of nonsynonymous substitutions as well as alignment errors that cause longer stretches of nonhomologous sequence due to variation in exon splicing or misannotation (Harrison et al. 2014). To select these thresholds, we first applied a range of masking criteria on our datasets. We then ran the branch-site test for positive selection on the target species branches for both the unmasked and masked datasets. We visually inspected the alignments of genes with the highest likelihood ratios and chose the masking criteria that was most efficient at reducing false positive rates. Finally, we discarded orthologs for which the alignment length was shorter than 300 bp following gap removal and masking, as these likely represent incomplete sequences.

To obtain divergence estimates for each orthogroup and calculate mean d_N/d_S across the target species branch, we used branch model (model=2, nssites=0) in the CODEML package in PAML v4.8 (Yang 2007), using the phylogeny ((Target species#1, *P. formosa*), *X. maculatus*, *O. latipes*), where the target species was successively *P. reticulata*, *P. wingei*, *P. picta*, *P. parae*, *P. latipinna* and *G. holbrooki*. To avoid inaccurate divergence estimates dues to mutational saturation and double hits, orthologous genes with $d_S > 2$ were removed from subsequent analyses (Axelsson et al. 2008).

For each of the target species, we divided orthologs into autosomal and sex-linked categories based on their chromosomal location. The sex-linked category excluded genes in the previously identified pseudoautosomal regions (0-5Mb and >26Mb of *P. reticulata* chromosome 12; >20Mb of *P. wingei*, *P. picta*, *P. parae*, *P. latipinna* and *G. holbrooki* chromosomes that are syntenic to the guppy chromosome 12 (Darolti et al. 2019; Sandkam et al. 2021)). In addition, it has recently been discovered that the *P. reticulata* reference genome, which served as reference for scaffold ordering and orientation for our *P. reticulata* de novo genome assembly, has a large inversion on the X chromosome that is specific to the guppy strain on which the reference genome assembly was built and is not present in any of our lab or wild-caught samples (Darolti et al. 2020; Almeida et al. 2021). We thus corrected

for this inversion when excluding genes from the PAR and assigning genes to the sex-linked

category in *P. reticulata*.

Separately for each genomic category, we extracted the number of nonsynonymous

substitutions (D_N), the number of nonsynonymous sites (N), the number of synonymous

substitutions (D_S) and the number of synonymous sites (S). Taking into account alignment

length, we calculated mean d_N and mean d_S as the ratio of the number of substitutions across

all orthologs in that group divided by the number of sites ($d_N = D_N / N$; $d_S = D_S / S$), thus avoiding

bias from short sequences and the issue of infinitely high d_N/d_S estimates due to very low d_S

(Mank et al. 2007). We identified significant differences in d_N, d_S and d_N/d_S between genomic

categories by subsampling without replacement the autosomal dataset at the size of the

smaller dataset (e.g., X chromosome dataset; Stratum I dataset) using 1,000 replicates, and

we used bootstrapping with 1,000 replicates to determine 95% confidence intervals for each

divergence estimate. The strength of the Faster-X effect was calculated for each target species

as the ratio of the rate of divergence for the X chromosome over the rate of divergence for

the autosomes X(d_N/d_S):A(d_N/d_S).

Additional sequence divergence analyses for *P. reticulata*

For comparison to our estimates of rates of divergence based on *de novo* transcripts, we also

estimated rates of sequence evolution in *P. reticulata* using publicly available coding

sequences from Ensembl 104 (Guppy_female_1.0_MT). For this analysis, we followed the

same steps outlined above in terms of extracting the longest isoform for each gene,

identifying orthogroups, aligning target and outgroup sequences, masking and obtaining

divergence estimates.

In addition, we have previously constructed high quality *P. reticulata* female *de novo* genome

assemblies for replicate upstream and downstream populations of three rivers in Trinidad,

Quare, Aripo, and Yarra (Almeida et al. 2021). To each genome assembly, we mapped *P.*

reticulata RNA-seq reads from our laboratory population and generated a non-redundant set

of transcripts for each wild population following the methods detailed above. For each

upstream and downstream population, we then estimated rates of coding sequence

evolution for the X chromosome, excluding the PAR, and the autosomes by following the

same pipeline described in the previous section.

Estimating rates of divergence for the ancestral branch to *P. reticulata* and *P. picta*

We performed additional sequence divergence analyses following the same methodology

described above but using different phylogenies that allowed us to estimate rates of evolution

on the internal branch that is ancestral to *P. reticulata* and *P. picta*. In addition to the *X.*

maculatus coding sequences, we obtained sequences for *G. affinis* (ASM309773v1) and *P.*

latipinna (P_latipinna-1.0) from Ensembl 105. We used the phylogenies ((*P. reticulata*, *P.*

picta)#1, *P. latipinna*, *X. maculatus*) and ((*P. reticulata*, *P. picta*)#1, *G. affinis*, *X. maculatus*) to

obtain divergence estimates in PAML. Similarly, we estimated rates of evolution for all the

other internal branches of the tree (Fig. 2).

Polymorphism data

To obtain polymorphism data, for each target species we first mapped female RNA-seq reads

to the genome assembly using the STAR v2.4.2a aligner in two-pass mode (Dobin et al. 2013),

and then called SNPs using SAMtools v1.3.1 mpileup (Li et al. 2009) with a minimum base

quality of 20 and VarScan v2.3.9 mpileup2snp (Koboldt et al. 2012) with a minimum coverage

of two, a minimum average quality of 20, a minimum variant allele frequency of 0.1, *p* value

of 0.05 and strand filter set to "on". We next imposed that the polymorphism dataset also

passes the filtering criteria used for calculating rates of sequence divergence. As such, we

identified codons for which all sites pass the minimum coverage threshold of 20 in at least

half of the individuals (in this case, a minimum of two out of three individuals), there are no

alignment gaps following alignment with PRANK, and no ambiguity data (Ns) following

masking with SWAMP. To make our dataset compatible for the McDonald-Kreitman test of

selection, we only kept genes with both divergence and polymorphism data. We have

therefore used the same number of nonsynonymous and synonymous sites in our divergence

and polymorphism calculations. We identified whether SNPs were synonymous or

nonsynonymous by matching them to the reading frame using custom scripts (available at

https://github.com/manklab/Darolti_etal_2022_FastX_evolution). For each genomic

category, we calculated mean p_N and mean p_S as the ratio of the number of polymorphisms

across all orthologs in that group divided by the number of sites ($p_N=P_N/N$; $p_S=P_S/S$).

Testing for selection using divergence and polymorphism data

For each species, we estimated the number of genes evolving under neutral and adaptive

evolution using the McDonald-Kreitman test (McDonald and Kreitman 1991). Neutral theory

predicts that the ratio of nonsynonymous to synonymous changes within species (p_N/p_S)

should be equal to that between species (d_N/d_S) (McDonald and Kreitman 1991). As such, the

McDonald-Kreitman test identifies signatures of positive selection, where there is an excess

of nonsynonymous substitutions relative to polymorphisms ($d_N/d_S > p_N/p_S$), and of relaxed

purifying selection, where there is a deficit of nonsynonymous substitutions compared to

polymorphisms ($d_N/d_S < p_N/p_S$). To test for deviations from neutrality, for each contig, we used

a 2 x 2 contingency table and a Fisher's Exact Test in R v3.6.2 (R Core Team 2015). The power

of the McDonald-Kreitman test is limited with low table counts, therefore we restricted this

analysis to genes for which the sum of each row and column in the contingency table was

equal to or greater than six (Begun et al. 2007; Andolfatto 2008).

As the McDonald-Kreitman test is a very conservative test, we also used the divergence and

polymorphism data to calculate the direction of selection statistic (DoS) (Stoletzki and Eyre-

Walker 2011). For each gene, we calculated the difference between the proportion of

nonsynonymous substitutions and polymorphisms (DoS=$D_N(D_N+D_S)-P_N(P_N+P_S)$), where

positive DoS values are indicative of positive selection. We used Fisher's Exact test in R to test

for significant differences in the proportion of genes under positive selection between the

autosomes and the X chromosome in each species.

Lastly, we used the polymorphism data alone to test for an excess or deficit of

nonsynonymous polymorphisms on the X chromosome compared to the autosomes, which

would suggest a relaxed constraint or purifying selection, respectively. For this, we

concatenated P_N and P_S estimates within each species and used Fisher's Exact test in R to test

for significant differences in P_N/P_S between the autosomes and the X chromosome.

Acknowledgements

532 This work was supported by the European Research Council (grant agreement 680951), a

533 Canada 150 Research Chair, and an NSERC Discovery Grant to J.E.M. We thank C. Lacy for the

534 fish illustrations and members of the Mank lab for helpful discussions and suggestions.

Author Contributions

536 J.E.M. and I.D. designed the research; I.D., L.F., B.A.S., D.C.H.M. and J.E.M. performed the data

537 analysis and contributed to the writing of the manuscript.

Data availability

Scripts used for data processing and analysis are available at https://github.com/manklab/Darolti_etal_2022_FastX_evolution.

References

Almeida P, Sandkam BA, Morris J, Darolti I, Breden F, Mank JE. 2021. Divergence and remarkable diversity of the Y chromosome in guppies. Mol Biol Evol. 38:619–633.

Altschul SF, Gish W, Miller W, Myers E, Lipman DJ. 1990. Basic local alignment search tool. J Mol Biol. 215:403–410.

Andolfatto P. 2008. Controlling Type-I Error of the McDonald–Kreitman Test in Genomewide Scans for Selection on Noncoding DNA. Genetics. 180:1767–1771.

Ávila V, De Procé SM, Campos JL, Borthwick H, Charlesworth B, Betancourt AJ. 2014. Faster-X effects in two drosophila lineages. Genome Biol Evol. 6:2968–2982.

Axelsson E, Hultin-Rosenberg L, Brandström M, Zwahlén M, Clayton DF, Ellegren H. 2008. Natural selection in avian protein-coding genes expressed in brain. Mol Ecol. 17:3008–3017.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J, Mayrose I, Ming R, et al. 2014. Sex determination: why so many ways of doing it? PLoS Biol. 12:e1001899.

Bechsgaard J, Schou MF, Vanthournout B, Hendrickx F, Knudsen B, Settepani V, Schierup MH, Bilde T. 2019. Evidence for faster X chromosome evolution in spiders. Mol Biol Evol. 36:1281–1293.

Begun DJ, Holloway AK, Stevens K, Hillier LW, Poh YP, Hahn MW, Nista PM, Jones CD, Kern AD, Dewey CN, et al. 2007. Population genomics: whole-genome analysis of polymorphism and divergence in *Drosophila simulans*. PLoS Biol. 5:2534–2559.

Bergero R, Gardner J, Bader B, Yong L, Charlesworth D. 2019. Exaggerated heterochiasmy in a fish with sex-linked male coloration polymorphisms. PNAS 116:6924-6931.

Bolger AMLMUB. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30:2114–2120.

Charlesworth B, Campos JL, Jackson BC. 2018. Theory and evidence from *Drosophila*. Mol Ecol. 27:3753–3771.

Charlesworth B, Coyne JA, Barton NH. 1987. The relative rates of evolution of sex chromosomes and autosomes. Am Nat. 130:113–146.

Charlesworth D, Bergero R, Graham C, Gardner J, Keegan H. 2021. How did the guppy Y chromosome evolve? PLoS Genet. 17:e1009704.

Charlesworth D, Bergero R, Graham C, Gardner J, Yong L. 2020. Locating the sex determining region of linkage group 12 of guppy (Poecilia reticulata). G3 10:3639:3649.

Darolti I, Almeida P, Wright AE, Mank JE. 2022. A comparison of methodological approaches to the study of young sex chromosomes: A case study in Poecilia. J. Evol. Biol.

https://doi.org/10.1111/jeb.14013.

Darolti I, Wright AE, Mank JE. 2020. Guppy Y chromosome integrity maintained by
incomplete recombination suppression. Genome Biol Evol. 12:965–977.

Darolti I, Wright AE, Sandkam BA, Morris J, Bloch NI, Farré M, Fuller RC, Bourne GR, Larkin
DM, Breden F, et al. 2019. Extreme heterogeneity in sex chromosome differentiation and
dosage compensation in livebearers. Proc Natl Acad Sci USA. 116:19031–19036.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29:15–21.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. Evol. 20:1-14.

Flicek P, Amode MR, Barrell D, Beal K, Billis K, Brent S, …, Searle SMJ. 2014. Ensembl 2014.
Nucleic Acids Res. 42:749–755.

Fraser BA, Whiting JR, Paris JR, Weadick CJ, Parsons PJ, Charlesworth D, Bergero R, Bemm F,
Hoffmann M, Kottler VA, et al. 2020. Improved reference genome uncovers novel sex-linked
regions in the guppy (*Poecilia reticulata*). Genome Biol Evol. 12:1789–1805.

Harrison PW, Jordan GE, Montgomery SH. 2014. SWAMP : Sliding Window Alignment
Masker for PAML. Evol Bioinform. 10:EBO-S18193.

Jaquiéry J, Peccoud J, Ouisse T, Legeai F, Prunier-Leterme N, Gouin A, Nouhaud P, Brisson
JA., Bickel R, Purandare S, et al. 2018. Disentangling the Causes for Faster-X Evolution in
Aphids. Genome Biol Evol. 10:507–520.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: A fast spliced aligner with low memory
requirements. Nat Methods. 12:357–360.

Kirkpatrick M, Hall DW. 2004. Male-biased mutation, sex linkage, and the rate of adaptive
evolution. Evol. 58:437–440.

Kirkpatrick M, Sardell JM, Pinto BJ, Dixon G, Peichel CL, Schartl M. 2022. Evolution of the
canonical sex chromosomes of the guppy and relatives. G3 12:jkab435.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L,
Wilson RK. 2012. VarScan 2: Somatic mutation and copy number alteration discovery in
cancer by exome sequencing. Genome Res. 22:568–576.

Kotrschal A, Rogell B, Bundsen A, Svensson B, Zajitschek S, Immler S, Maklakov AA, Kolm N.
2013. Experimental evidence for costs and benefits of evolving a larger brain. Curr Biol.
23:168–171.

Kousathanas A, Halligan DL, Keightley PD. 2014. Faster-X adaptive protein evolution in
house mice. Genetics. 196:1131–1143.

Lenormand T, Fyon F, Sun E, Roze D. 2020. Sex chromosome degenration by regulatory
evolution. Curr Biol. 30:3001-3006.e3005.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.
2009. The sequence alignment / map format and SAMtools. Bioinformatics. 25:2078–2079.

Löytynoja A, Goldman N. 2008. Phylogeny-aware gap replacement prevents errors in

sequence alignment and evolutionary analysis. Science. 320:1632–1635.

Lu J, Wu C-I. 2005. Weak selection revealed by the whole-genome comparison of the X
chromosome and autosomes of human and chimpanzee. PNAS. 102:4063–4067.

Mank JE, Axelsson E, Ellegren H. 2007. Fast-X on the Z: rapid evolution of sex-linked genes in
birds. Genome Res. 17:618–624.

Mank JE, Vicoso B, Berlin S, Charlesworth B. 2010. Effective population size and the Faster-X
effect: empirical results and their interpretation. Evolution. 64:663–674.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the ADH locus in *Drosophila*.
Nature. 351:652–654.

Meisel RP, Connallon T. 2013. The faster-X effect: integrating theory and data. Trends
Genet. 29:537–544.

Meredith RW, Pires MN, Reznick DN, Springer MS. 2010. Molecular phylogenetic
relationships and the evolution of the placenta in Poecilia (Micropoecilia) (Poeciliidae:
Cyprinodontiformes). *Mol. Phylogenet. Evol.* 55:631-639.

Metzger DCH, Sandkam BA, Darolti I, Mank JE. 2021. Rapid evolution of complete dosage
compensation in *Poecilia*. Genome Biol Evol. 13:evab155.

Morris J, Darolti I, Bloch NI, Wright AE, Mank JE. 2018. Shared and Species-Specific Patterns
of Nascent Y Chromosome Evolution in Two Guppy Species. Genes. 9:238.

Nanda I, Schartl M, Feichtinger W, Epplen JT, Schmid M. 1992. Early stags of sex
chromosome differentiation in fish as analysed by simple repetitive DNA sequences.
Chromosoma. 101:301–310.

Nanda I, Schories S, Tripathi N, Dreyer C, Haaf T, Schmid M, Schartl M. 2014. Sex
chromosome polymorphism in guppies. Chromosoma. 123:373–383.

Otto SP, Pannell JR, Peichel CL, Ashman T-L, Charlesworth D, Chippindale AK, Delph LF,
Guerrero RF, Scarpino S V, McAllister BF. 2011. About PAR: the distinct evolutionary
dynamics of the pseudoautosomal region. Trends Genet. 27:358–367.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol.
33:290–295.

Pinharanda A, Rousselle M, Martin SH, Hanly JJ, Davey JW, Kumar S, Galtier N, Jiggins CD.
2019. Sexually dimorphic gene expression and transcriptome evolution provide mixed
evidence for a fast-Z effect in *Heliconius*. J Evol Biol. 32:194–204.

Pool JE, Nielsen R. 2007. Population size changes reshape genomic patterns of diversity. J
Org Evol. 61:30001–3006.

Qiu S, Yong L, Wilson A, Croft DP, Graham C, Charlesworth D. 2022. Partial sex linkage and
linkage disequilibrium on the guppy sex chromosome. BioRxiv doi:
https://doi.org/10.1101/2022.01.14.476360.

Quinlan AR, Hall IM. 2010. Genome analysis BEDTools: A flexible suite of utilities for
comparing genomic features. Bioinformatics. 26:841–842.

R Core Team. 2015. R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing. Available from: https//www.r-project.org.
Accessed August 25, 2021.

Rabosky DL, Chang J, Cowman PF, Sallan L, Friedman M, Kascher K, Garilao C, Near TJ, Coll
M, Alfaro ME. 2018. An inverse latitudinal gradient in speciation rate for marine fishes.
Nature. 559:392-395.

Rousselle M, Faivre N, Ballenghien M, Galtier N, Nabholz B. 2016. Hemizygosity enhances
purifying selection: Lack of fast-Z evolution in two satyrine butterflies. Genome Biol Evol.
8:3108–3119.

Sackton TB, Corbett-Detig RB, Nagaraju J, Vaishna L, Arunkumar KP, Hartl DL. 2014. Positive
selection drives faster-Z evolution in silkmoths. Evolution. 68:2331–2342.

Sandkam BA, Almeida P, Darolti I, Furman BLS, van der Bijl W, Morris J, Bourne GR, Breden
F, Mank JE. 2021. Extreme Y chromosome polymorphism corresponds to five male
reproductive morphs of a freshwater fish. Nat Ecol Evol. 5:939–948.

Sigeman H, Sinclair B, Hansson B. 2022. FindZX: an automated pipeline for detecting and
visualising sex chromosomes using whole-genome sequencing data. BMC Genomics 23:328.

Stevenson BJ, Iseli C, Panji S, Zahn-Zabal M, Hide W, Old LJ, Simpson AJ, Jongeneel CV. 2007.
Rapid evolution of cancer/testis genes on the X chromosome. BMC Genomics. 8:1–11.

Stoletzki N, Eyre-Walker A. 2011. Estimation of the neutrality index. Mol Biol Evol. 28:63–70.

Tajima F. 1989. The effect of change in population size on DNA polymorphism. Genetics.
123:597–601.

Traut W, Winking H. 2001. Meiotic chromosomes and stages of sex chromosome evolution
in fish: zebrafish, platyfish and guppy. Chromosom Res. 9:659–672.

Tripathi N, Hoffmann M, Willing EM, Lanz C, Weigel D, Dreyer C. 2009. Genetic linkage map
of the guppy, *Poecilia reticulata*, and quantitative trait loci analysis of male size and colour
variation. Proc Biol Sci. 276:2195–2208.

Vicoso B, Bachtrog D. 2013. Reversal of an ancient sex chromosome to an autosome in
Drosophila. Nature. 499:332–335.

Vicoso B, Charlesworth B. 2006. Evolution on the X chromosome: unusual patterns and
processes. Nat Rev Genet. 7:645.

Vicoso B, Charlesworth B. 2009. The deficit of male-biased genes on the *D. melanogaster* X
chromosome is expression-dependent: A consequence of dosage compensation? J Mol Evol.
68:576–583.

Winge Ö. 1922. One-sided masculine and sex-linked inheritance in *Lebistes reticulatus*. J
Genet. 12:145–162.

Winge Ö. 1927. The location of eighteen genes in *Lebistes reticulatus*. J Genet. 18:1–43.

Winge Ö, Ditlevsen E. 1947. Colour inheritance and sex determination in *Lebistes*. Heredity.
1:65–83.

Wright AE, Darolti I, Bloch NI, Oostra V, Sandkam B, Buechel SD, Kolm N, Breden F, Vicoso B,

Mank JE. 2017. Convergent recombination suppression suggests role of sexual selection in
guppy sex chromosome formation. Nat Commun. 8:14251.

Wright AE, Harrison PW, Zimmer F, Montgomery SH, Pointer MA, Mank JE. 2015. Variation
in promiscuity and sexual selection drives avian rate of Faster-Z evolution. Mol Ecol.
24:1218–1235.

Wright AE, Mank JE. 2013. The scope and strength of sex-specific selection in genome
evolution. J Evol Biol. 26:1841–1853.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol.
24:1586–1591.

Figure Legends

Figure 1. Estimates of the rate of divergence (d_N/d_S) for autosomal and X-linked genes in (A)
***P. reticulata*, (B) *P. wingei*, (C) *P. picta*, (D) *P. parae*, (E) *P. latipinna* and (F) *G. holbrooki*.** For
all species, the X chromosome category excludes genes on the PAR. Additionally, in *P. reticulata* and *P. wingei*, the X chromosome category excludes genes in Stratum I. In *P. latipinna* and *G. holbrooki*, the X chromosome represents the chromosome syntenic to guppy chromosome 12. 95% confidence intervals are based on bootstrapping with 1,000 replicates. Differences between autosomal and X-linked loci are based on 1,000 replicate permutation tests. Only significant differences are shown (p value < 0.01).

Figure 2. Fast-X effect, calculated as the ratio of d_N/d_S for the X chromosome (excluding the PAR) to that of the autosomes, across the poeciliids. In *P. latipinna* and *G. holbrooki*, chromosome 12 has not been implicated as the sex chromosome. Numbers on each branch represent the estimated Fast-X effect. Phylogeny based on Meredith et al. 2010 and Rabosky et al. 2018.

Figure 3. Estimates of rates of evolution in wild *P. reticulata* populations. (A) Fast-X effect across *P. reticulata* upstream, low predation populations (orange) and downstream, high predation populations (black) in the Quare, Aripo and Yarra rivers. The Fast-X effect is calculated as the ratio of the rate of nonsynonymous over synonymous substitutions for the X chromosome, excluding the PAR, relative to that for the autosomes.

723 **Tables**

Table 1. Divergence estimates for autosomal and X-linked genes on the ancestral branch to *P. reticulata* and *P. picta*. The X chromosome category excludes genes on the PAR. 95% confidence intervals are based on bootstrapping with 1,000 replicates.

Phylogeny	Category	No. genes	d_N (95% CI)	d_S (95% CI)	d_N/d_S (95% CI)
(((*P. reticulata*, *P. picta*), *P. latipinna*), *X. maculatus*)	Autosomes	2,213	0.0013 (0.0013-0.0014)	0.0094 (0.0092-0.0097)	0.1422 (0.1338-0.1507)
	X chromosome	82	0.0014 (0.0010-0.0018)	0.0096 (0.0085-0.0110)	0.1416 (0.1016-0.1937)
(((*P. reticulata*, *P. picta*), *G. affinis*), *X. maculatus*)	Autosomes	3,654	0.0049 (0.0048-0.0051)	0.0451 (0.0444-0.0457)	0.1093 (0.1055-0.1130)
	X chromosome	127	0.0052 (0.0044-0.0061)	0.0477 (0.0441-0.0510)	0.1096 (0.0924-0.1299)

Table 2. Polymorphism estimates for autosomal and X-linked genes across Poeciliid species. Only genes with both divergence and polymorphism data were included in this analysis. For all species, the X chromosome category excludes genes on the PAR. 95% confidence intervals are based on bootstrapping with 1,000 replicates. Differences between autosomal and sex-linked categories are based on 1,000 replicate permutation tests and significant differences are indicated by bold values ($p < 0.01$).

Species	Category	No. genes	p_N (95% CI)	p_S (95% CI)	p_N/p_S (95% CI)
P. reticulata	Autosomes	2,211	0.0012 (0.0011-0.0012)	0.0110 (0.0105-0.0113)	0.1054 (0.0980-0.1134)
	X chromosome	74	0.0012 (0.0008-0.0016)	0.0131 (0.0109-0.0159)	0.0883 (0.0591-0.1225)
P. wingei	Autosomes	934	0.0009 (0.0008-0.0010)	0.0077 (0.0071-0.0082)	0.1155 (0.1036-0.1275)
	X chromosome	34	0.0008 (0.0004-0.0012)	0.0103 (0.0080-0.0130)	0.0791 (0.0411-0.1257)
P. picta	Autosomes	1,485	0.0006 (0.0006-0.0007)	0.0051 (0.0048-0.0053)	0.1191 (0.1082-0.1311)
	X chromosome	40	0.0003 (0.0002-0.0006)	0.0043 (0.0030-0.0061)	0.0790 (0.0438-0.1293)
P. parae	Autosomes	4,167	0.0010 (0.0009-0.0011)	0.0078 (0.0076-0.0080)	0.1307 (0.1249-0.1365)
	X chromosome	123	0.0007 (0.0006-0.0009)	**0.0059 (0.0049-0.0068)**	0.1301 (0.0983-0.1776)
P. latipinna	Autosomes	1,041	0.0010 (0.0009-0.0011)	0.0095 (0.0089-0.0100)	0.1081 (0.1082-0.1311)
	Chromosome syntenic to guppy X	35	0.0016 (0.0010-0.0024)	0.0111 (0.0081-0.0147)	0.1452 (0.0832-0.2469)
G. holbrooki	Autosomes	1,464	0.0014 (0.0013-0.0015)	0.0155 (0.0146-0.0163)	0.0872 (0.0800-0.0947)
	Chromosome syntenic to guppy X	42	0.0013 (0.0008-0.0018)	0.0191 (0.0153-0.0234)	0.0659 (0.0410-0.0990)

 Figures

 Fig. 1

729   **Fig. 2**

731 **Fig. 3**