Sex Chromosome Heteromorphism and the Fast-X Effect in Poeciliids

Iulia Darolti[1], Lydia J. M. Fong[1], Judith E. Mank[1,2]

1 Department of Zoology and Biodiversity Research Centre, University of British Columbia,

Vancouver, Canada

2 Centre for Ecology and Conservation, College of Life and Environmental Sciences,

University of Exeter, Cornwall, United Kingdom

Corresponding author: Iulia Darolti, e-mail: darolti@zoology.ubc.ca

Abstract

An accelerated rate of sequence evolution on the X chromosome compared to autosomes,

known as Fast-X evolution, has been observed in a range of heteromorphic sex chromosomes.

However, it remains unclear how early in the process of sex chromosome differentiation the

Fast-X effect becomes detectible. Recently, we uncovered an extreme variation in sex

chromosome heteromorphism across Poeciliid fish species. The common guppy, *Poecilia*

reticulata, Endler's guppy, *P. wingei*, and the swamp guppy, *P. picta*, appear to share the same

XY system and exhibit a remarkable range of heteromorphism. The sex chromosome system

is absent in recent outgroups, including *P. latipinna* and *Gambusia holbrooki*. We combined

analyses of sequence divergence and polymorphism data across Poeciliids to investigate X

chromosome evolution as a function of hemizygosity and reveal the causes for Fast-X effects.

Consistent with the extent of Y degeneration in each species, we detect higher rates of

divergence on the X relative to autosomes and a strong Fast-X effect in *P. picta*, while no

change in the rate of evolution of X-linked relative to autosomal genes in *P. reticulata*. In *P.*

wingei, the species with intermediate sex chromosome differentiation, we see an increase in

the rate of nonsynonymous substitutions on the older stratum of divergence only. We also

use our comparative approach to test different models for the origin of the sex chromosomes

in this clade. Taken together, our study reveals an important role of hemizygosity in Fast-X

and suggests a single, recent origin of the sex chromosome system in this clade.

Keywords: sex chromosome evolution, heteromorphy, hemizygosity

Introduction

Owing to their unusual inheritance pattern and hemizygosity in males, X chromosomes display many distinct evolutionary properties compared to the rest of the genome (Charlesworth et al. 1987; Vicoso and Charlesworth 2006). The strength of selection and genetic drift, as well as the role of dominance and effective population size, are expected to differ between sex chromosomes and autosomes (Kirkpatrick and Hall 2004; Mank et al. 2010; Meisel and Connallon 2013). Comparing differences in the evolution of sex-linked and autosomal loci is important for understanding patterns of mutation and selection acting across the genome.

An elevated rate of coding sequence evolution on the X chromosome relative to the autosomes, referred to as the Fast-X effect (or Fast-Z in the case of ZW systems) has been observed in a diversity of organisms, including humans (Lu and Wu 2005), primates (Stevenson et al. 2007), mice (Kousathanas et al. 2014), birds (Mank et al. 2007; Mank et al. 2010; Wright et al. 2015), *Drosophila* (Mank et al. 2010; Ávila et al. 2014; Charlesworth et al. 2018), aphids (Jaquiéry et al. 2018), spiders (Bechsgaard et al. 2019), and lepidoptera (Pinharanda et al., 2019; Sackton et al., 2014). However, the magnitude of this effect can vary substantially across species due to demographic factors, differences in mating system or regulatory mechanisms acting on the sex chromosomes (Mank et al. 2010). Notably, all the cases of Fast-X above were observed in highly heteromorphic sex chromosomes, where very few genes, if any, remain on the Y chromosome and the X is largely hemizygous in the heterogametic sex.

There are two potential causes of Fast-X. The single functional copy of the X chromosome in males leads to hemizygous exposure of genes and therefore stronger purifying selection against recessive deleterious mutations and positive selection for recessive beneficial ones expressed in males (Charlesworth et al. 1987). This adaptive cause of Fast-X is mainly expected in species with heteromorphic sex chromosomes and highly degenerated Y chromosomes, as a large proportion of X-linked genes will be hemizygous in males. Alternatively, in every male and female pair, there are only three X chromosomes to four copies of each autosome, and this varies substantially based on mating system and type of heterogamety (Vicoso and Charlesworth 2009; Mank et al. 2010; Wright and Mank 2013; Wright et al. 2015). This reduced effective population size of the X relative to the autosomes diminishes the relative power of selection on the X, potentially leading to both greater genetic drift and fixation of weakly

deleterious mutations on the X chromosome (Charlesworth et al. 1987). This non-adaptive

cause of Fast-X does not necessarily require male hemizygosity, just recombination

suppression between the X and Y chromosomes and differences in effective population size

between X-linked and autosomal loci, and could apply to X loci where the Y copy is expressed

and remains functional.

Recently, we uncovered an extreme variation across Poeciliids in the rate of sex chromosome

degeneration and dosage compensation. The same chromosome pair acts as the XY system in

the common guppy, *Poecilia reticulata*, its sister species Endler's guppy, *P. wingei*, and the

more distantly related swamp guppy, *P. picta,* all of which last shared a common ancestor

roughly 20 million years ago (Darolti et al. 2019). Notably, *P. latipinna* and *G. holbrooki*, recent

outgroup species to this clade, do not show evidence of a sex chromosome system on the

same chromosome (Darolti et al. 2019), and therefore we hypothesized based on parsimony

that the sex chromosome system arose roughly 20 mya (supplementary Fig. S1, Parsimony

Model). Since this origin, the sex chromosomes of *P. reticulata* and *P. wingei* have remained

largely homomorphic, with little sequence differentiation between the X and the Y. In

contrast, the *P. picta* sex chromosomes are completely nonrecombining (with the exception

of a small pseudoautosomal region at the distal end, >20 Mb) and the Y chromosome has

undergone substantial degeneration, leaving most of the X chromosome in males effectively

hemizygous (Darolti et al. 2019). Moreover, *P. picta* exhibits complete dosage compensation,

a pattern that has also been observed in its sister taxon, *P. parae* (Metzger et al. 2021;

Sandkam et al. 2021), which is predicted to accentuate the Fast-X effect (Charlesworth et al.

1987; Mank et al. 2010). The range of sex chromosome differentiation present in this clade

allows us to investigate patterns of X chromosome coding sequence evolution and to test

causes underlying Fast-X effects.

In contrast to the Parsimony Model, it has recently been proposed that the sex chromosomes

of *P. reticulata* represent a recent turnover event, and that the highly diverged *P. picta* and *P.*

parae system is in fact ancient (Charlesworth et al. 2021). This Turnover Model

(supplementary Fig. S1) posits that the *P. picta* X chromosome arose well before the

immediate ancestor of *P. parae, P. picta*, *P. reticulata* and *P. wingei*, and slowly diverged over

time resulting in the heteromorphism observed*.* This model supposes that the

homomorphism observed in *P. reticulata* and *P. wingei* represents a turnover event, where

the current guppy X and Y chromosomes arose from the ancestral X of *P. picta*. Sex
chromosome turnovers are frequent in many animal groups (Bachtrog et al. 2014), so the
model is plausible.

However, the unique molecular and evolutionary signatures that accumulate on sex
chromosomes remain observable even when they revert to being autosomes (Vicoso and
Bachtrog 2013). That means that if the Turnover Model is true, Fast-X patterns, evolving over
long periods of time in the distant ancestor of *P. picta*, would also remain detectible on the X
chromosome in *P. reticulata* and *P. wingei* after turnover. Furthermore, the Turnover Model
requires additional turnover events in several other taxa within the clade that do not share
the same sex chromosome as *P. picta* (supplementary Fig. S1), and we would also expect these
species to exhibit ancient patterns of Fast-X on the chromosome, which is fully autosomal.
Our comparative dataset allows us to also test these alternative hypotheses for the origin of
the guppy sex chromosomes.

Using a combination of sequence divergence, polymorphism and expression data analyses
across Poeciliid species, we estimate rates of gene sequence evolution across the genome and
assess the presence of Fast-X evolution in each system. Consistent with the extent of Y
degeneration, we find significantly higher rates of X coding sequence evolution compared to
the autosomes and a strong signal of Fast-X effect in *P. picta*, potentially accelerated by the
evolution of chromosome wide dosage compensation in this system. The absence of a
consistent Fast-X evolution in *P. reticulata* and *P. wingei*, as well as in the outgroup species *P.
latipinna* and *G. holbrooki*, argues against the Turnover Model.

Results

Strong signal of Fast-X evolution in the heteromorphic sex chromosomes of *P. picta*
We first assessed the strength of Fast-X in our study species with heteromorphic sex
chromosomes. The extensive X chromosome hemizygosity of *P. picta* males is expected to
accentuate the strength of Fast-X evolution in this species (Vicoso and Charlesworth 2006), as
is the mechanism of complete X chromosome dosage compensation (Mank et al. 2010).
Indeed, our analysis in *P. picta* revealed a greater mutation rate on the X chromosome,
excluding genes in the pseudoautosomal region (PAR) (see methods), compared to the
autosomes, as shown through the significantly higher rate of nonsynonymous substitutions

for X-linked loci (1,000 replicates permutation test, $p < 0.001$; supplementary Table S1). *P. picta* also exhibits a significantly elevated rate of divergence (d_N/d_S) on the X chromosome relative to the rest of the genome (Fig. 1C) and a strong Fast-X effect (Fig. 2), calculated as the ratio of the rate of nonsynonymous over synonymous substitutions for the X chromosome relative to that for the autosomes.

Absence of Fast-X effect in species with homomorphic sex chromosomes

We next asked whether patterns of coding sequence evolution are different between *P. picta* and the species with homomorphic sex chromosomes, *P. reticulata* and *P. wingei*. Previous work has indicated that *P. reticulata* and *P. wingei* share the same male-heterogametic sex chromosome system (Nanda et al. 2014; Morris et al. 2018; Darolti et al. 2019). Analyses of coverage differences between males and females indicate that Y degeneration is restricted to the distal end of the sex chromosomes, in a region ancestral to *P. reticulata* and *P. wingei* (designated as Stratum I) (Wright et al. 2017; Darolti et al. 2019; Darolti et al. 2020; Fraser et al. 2020), and suggest that Y degeneration is slightly more exaggerated in *P. wingei* compared to *P. reticulata*. This nonrecombining region coincides with the previously mapped location of the sex determining region in *P. reticulata* (Winge 1922; Winge 1927; Winge and Ditlevsen 1947; Traut and Winking 2001; Tripathi et al. 2009) and is also found across six natural guppy populations (Almeida et al. 2021).

We therefore assessed the signal of Fast-X for genes in Stratum I (20-26Mb on *P. reticulata* chromosome 12; 17-20Mb on *P. wingei* syntenic chromosome to guppy sex chromosome). In *P. reticulata*, the rate of divergence for X-linked genes in Stratum I was not significantly different than that for autosomal genes (Fig. 1A; supplementary Table S1). To exclude the possibility that we were lacking power in our analysis due to the small number of genes identified in Stratum I, we redid the analysis for *P. reticulata* using Ensembl coding sequences instead of our *de novo* generated transcripts. We were able to recover more than twice as many X-linked loci, however the estimates of divergence remain the same as those based on *de novo* transcripts (supplementary Fig. S2).

In *P. wingei*, genes in Stratum I show significantly higher rates of nonsynonymous and of synonymous substitutions (supplementary Table S1), however the overall rate of divergence is similar between the autosomes and Stratum I (Fig. 1B) and the Fast-X effect is not as

accentuated as in *P. picta* (Fig. 2). Recent work has suggested that the *P. wingei* X and Y

chromosomes are somewhat more differentiated from each other compared to those of *P.*

reticulata (Darolti et al. 2019), and there is also evidence for this from previous cytogenetic

work (Nanda et al. 1992; Nanda et al. 2014). This greater divergence might explain the

difference in d_N/d_S estimates for X-linked genes in Stratum I that we observe between these

two species, though the effect is negligible. However, note that following filtering (see

methods), only a handful of *P. wingei* genes remained in Stratum I, and as such our statistical

power to detect a significant Fast-X signal in this species is reduced.

If the guppy sex chromosomes represent a turnover event, we would expect the pattern of

Fast-X in *P. picta* to still be observed throughout the *P. reticulata* and *P. wingei* sex

chromosomes outside of Stratum I, as the majority of it accumulated prior to turnover. Thus,

we next estimated rates of sequence divergence for the remainder of the X chromosome,

excluding genes in Stratum I and genes on the PAR (see methods), as the high recombination

events in the PAR can alter rates of evolution and polymorphism (Otto et al. 2011). The rate

of mutation and mean d_N/d_S on the X chromosome were not different from that on the

autosomes in either *P. reticulata* or *P. wingei* (Fig. 1A, B; Fig. 2; supplementary Table S1). This

finding is consistent with the limited Y degeneration observed in these species (Wright et al.

2017; Darolti et al. 2019), and contradicts the Turnover Model. In addition, the Turnover

Model also requires sex chromosome turnover events in the outgroup species to the *P.*

reticulata–P. wingei clade (supplementary Fig. S2). We therefore estimated sequence

divergence in the outgroups, *P. latipinna* and *G. holbrooki*, both species in which the guppy

chromosome 12 has not been implicated as the sex chromosome. Our data fails to show

elevated rates of evolution on the chromosome syntenic to the guppy X relative to the rest of

the genome in either *P. latipinna* or *G. holbrooki* (supplementary Table S1). This result,

together with our findings in *P. reticulata* and *P. wingei* argues against the Turnover Model.

Previous comparisons of replicate *P. reticulata* natural population have shown evidence for

greater X-Y divergence in replicate upstream, low predation populations relative to their

downstream, high predation populations pair within Quare, Aripo and Yarra rivers (Wright et

al. 2017; Almeida et al. 2021). This pattern of divergence has occurred over a relatively short

time span, at some point after the colonization of Trinidad at the last glacial maximum.

Therefore, we investigated whether we can detect higher rates of X chromosome to autosome

coding sequence evolution in upstream compared to downstream populations. We find a significantly elevated d_N/d_S on the X chromosome, excluding genes on the PAR and those on Stratum I, compared to the autosomes in the Yarra upstream population only (supplementary Tables S2-4), and across all rivers we do not find a consistently stronger Fast-X effect in the upstream relative to the downstream populations (Fig. 3). Compared to the other rivers, the upstream population of the Yarra river shows the greatest increase in Y divergence relative to the downstream population, measured by both male:female F_{ST} and male-specific SNPs (Almeida et al. 2021). X-Y divergence between upstream and downstream populations is greater in Yarra compared to Quare and Aripo, and this might explain why a pattern of Fast-X effect in the upstream population is recovered in Yarra alone.

Polymorphism and the underlying causes of Fast-X

A higher d_N/d_S on the X chromosome relative to the rest of the genome could be the result of both an increased genetic drift (Charlesworth et al. 1987; Mank et al. 2010) and an increased efficacy of selection (Vicoso and Charlesworth 2009). Therefore, we used sequence and polymorphism data together to test for the causes of elevated Fast-X evolution in *P. picta*. The McDonald-Kreitman test contrasts the number of nonsynonymous and synonymous substitutions with polymorphisms, where an excess of nonsynonymous substitutions relative to polymorphisms is indicative of positive selection (McDonald and Kreitman 1991). Using this test, we detected no X-linked genes with signatures of positive selection in any of the Poeciliid species. However, the McDonald-Kreitman test is very conservative, as it is restricted to genes with sufficient numbers of substitutions and polymorphisms (Begun et al. 2007; Andolfatto 2008), and as such our analysis was limited to a few X-linked contigs.

To increase our statistical power to detect signatures of positive selection, we also used the direction of selection (DoS) test which is less sensitive to low cell counts (Stoletzki and Eyre-Walker 2011). For each contig, DoS calculates the difference between the proportion of nonsynonymous substitutions and the proportion of nonsynonymous polymorphisms, where a positive DoS indicates adaptive evolution (Stoletzki and Eyre-Walker 2011). Using this approach, we recovered more genes with signatures of positive selection, however the X chromosome was not enriched in these genes compared to the rest of the genome in any of the species (supplementary Table S6).

Next, we used polymorphism data alone to test for deviations from neutrality. A higher rate
of nonsynonymous to synonymous polymorphisms on the X relative to the autosomes is
indicative of reduced efficacy of selection to remove mildly deleterious mutations, while a
lower rate suggests purifying selection resulting from hemizygous exposure of deleterious
mutations in males. Our results show a lower, though not significantly, rate of nonsynonymous
polymorphisms in *P. picta* only (Table 1, supplementary Table S5), which may suggest that the
observed Fast-X effect in *P. picta* is influenced by purifying selection acting to remove
deleterious mutations on the X (Charlesworth et al. 1987; Vicoso and Charlesworth 2009).

Discussion

The role of hemizygosity in Fast-X
We observe a clear pattern of Fast-X in *P. picta*, as shown through the significantly higher rate
of nonsynonymous substitutions for X-linked loci (supplementary Table S1) and significantly
elevated rate of divergence on the X chromosome relative to the rest of the genome (Fig. 1C,
Fig. 2), consistent with the extensive Y chromosome degeneration in this species. In contrast,
we did not observe significant patterns of Fast-X evolution in either *P. reticulata* or *P. wingei*.
There is little evidence of loss of Y gene coding sequence in either of these species (Darolti et
al. 2019; Darolti et al. 2020), and very few, if any, genes are hemizygously expressed in males,
even in the older Stratum I. Interestingly, the *P. wingei* Y is somewhat more diverged than the
Y in *P. reticulata* (Nanda et al. 1992; Nanda et al. 2014; Darolti et al. 2019), and we observe a
slight, though non-significant Fast-X effect in Stratum I in the former. We also observe an
effect in the Yarra upstream population, which shows the greatest X-Y divergence across
natural populations in Trinidad (Almeida et al. 2021). All this points to the key role of sex
chromosome hemizygosity in driving Fast-X evolution.

Theory predicts that complete dosage compensation, whereby balance in expression between
the sex chromosomes and the autosomes is restored, may facilitate a more pronounced Fast-
X effect (Charlesworth et al. 1987; Mank et al. 2010). Under this theory, in systems with
incomplete dosage compensation, where only a subset of the genes is compensated for but
overall expression for X-linked genes is reduced compared to the autosomes in males, X-linked
beneficial mutations may have lower expression and weaker phenotypic effects in males,
potentially limiting the Fast-X effect (Charlesworth et al. 1987; Mank et al. 2010). *P. picta* has

evolved chromosome-wide dosage compensation, seemingly through a mechanism involving

the hyperexpression of the single X in males (Darolti et al. 2019), and this may have

contributed to the observed signature of Fast-X evolution.

There are two potential causes of Fast-X. Male heterogamety leads to hemizygous exposure

of genes and therefore stronger purifying selection against recessive deleterious mutations

and positive selection for recessive beneficial ones expressed in males (Charlesworth et al.

1987). Alternatively, the reduced effective population size of the X relative to the autosomes

diminishes the relative power of selection on the X, potentially leading to non-adaptive causes

of Fast-X (Charlesworth et al. 1987) which do not necessarily require male hemizygosity, just

recombination suppression between the X and Y chromosomes.

The role of hemizygosity and dosage compensation we observe in *P. picta* Fast-X suggest an

adaptive mechanism, and our polymorphism analysis is consistent with greater purifying

selection acting in males to remove recessive deleterious variation. However, although we

were unable to differentiate adaptive and non-adaptive causes of Fast-X in our polymorphism

data, polymorphism estimates are sensitive to demographic fluctuations (Tajima 1989; Pool

and Nielsen 2007) and, thus, it is difficult to determine to what extent the Fast-X pattern in *P.*

picta is adaptive using polymorphism data alone. Future work identifying true X-hemizygous

loci in males and analyzing their patterns of sequence evolution may prove more revealing.

Differentiating different models of sex chromosome origin

Two models have been proposed for the origin of the sex chromosome system in this group.

The Parsimony Model posits that the sex chromosomes arose roughly 20 mya (supplementary

Fig. S1) in the ancestor of the species that all share it and experienced different rates of Y

decay in different sub-clades (Darolti et al. 2019), possibly exacerbated by the evolution of

complete dosage compensation in *P. picta* and close relatives (Metzger et al. 2021; Sandkam

et al. 2021). The Turnover Model proposes that the sex chromosomes of *P. reticulata*

represent a recent turnover event, and that the highly diverged *P. picta* and *P. parae* system

arose well before the immediate ancestor of *P. parae, P. picta*, *P. reticulata* and *P. wingei*,

slowly diverged over time resulting in the heteromorphism observed (Charlesworth et al.

2021). Because the molecular signatures that accumulate on sex chromosomes remain

observable even when they revert to being autosomes (Vicoso and Bachtrog 2013), we were

able to critically test these alternative models. If the guppy sex chromosomes represent a
turnover event, we would expect the pattern of Fast-X in *P. picta* to still be observed
throughout the *P. reticulata* and *P. wingei* sex chromosomes outside of Stratum I, as the
majority of it accumulated prior to turnover. In addition, the Turnover Model also requires sex
chromosome turnover events in the outgroup species to the *P. reticulata–P. wingei* clade, and
we would expect Fast-X in *P. latipinna* or *G. holbrooki*, where the ancient X chromosome is
presumably now fully autosomal.

We do not observe signatures of Fast-X in *P. reticulata* and *P. wingei* that would be consistent
with the Turnover Model, even after investigating multiple datasets (based on *de novo*
transcripts of a lab population and replicate wild populations, and Ensembl coding sequences).
Similarly, the lack of Fast-X in *P. latipinna* and *G. holbrooki*, the near outgroups to *P. picta, P.*
reticulata, and *P. wingei* (Darolti et al. 2019), which would have also experienced sex
chromosome turnover to alternative autosomes if the *P. picta* sex chromosomes are ancient,
also argues against an ancient origin and turnover. Instead, our results support the Parsimony
Model, and suggest that the sex chromosomes originated in the immediate ancestor of the *P.*
picta–P. reticulata–P. wingei clade and divergence occurred more rapidly in *P. picta,*
potentially accelerated by the evolution of complete X chromosome dosage compensation in
this species, which reduces selection to maintain Y chromosome expression for dosage
sensitive genes (Metzger et al. 2021).

Materials and Methods

Sample collection and sequencing
We have previously obtained tissue samples, extracted and sequenced RNA from the tails of
three males and three females of *P. reticulata*, *P. wingei*, *P. picta*, *P. latipinna* and *G. holbrooki*
(BioProject ID PRJNA353986, PRJNA528814) (Wright et al. 2017; Darolti et al. 2019). *P.*
reticulata samples were obtained from our outbred laboratory population originating from
the Quare River in Trinidad (Kotrschal et al. 2013). *P. wingei* samples were collected from our
laboratory population established from a strain maintained by a UK fish fancier. *P. picta*
samples were acquired from Guyana, while *P. latipinna* and *G. holbrooki* samples were
obtained in Florida. All samples were collected in accordance with national and institutional
ethical guidelines.

We extracted RNA from each sample using the Qiagen RNeasy Kit, following the instructions of the manufacturer. Library preparation and sequencing were performed at the University of Oxford Wellcome Centre for Human Genetics, following standard Illumina protocols and using the Illumina HiSeq 4000 platform. The data were quality assessed using FastQC v0.11.3 (www.bioinformatics.babraham.ac.uk/projects/fastqc) and trimmed with Trimmomatic v0.36 (Bolger 2014), removing adaptor sequences, reads with an average Phred score < 15 in a sliding window of four bases, reads with leading or trailing bases with a Phred score < 3 and reads shorter than 50bp following trimming.

Identifying orthogroups

For each species, we mapped RNA-seq reads to a previously constructed species-specific female *de novo* genome assembly (Wright et al. 2017; Darolti et al. 2019) using HISAT2 v2.0.4 (Kim et al. 2015), with the exception of *P. latipinna* for which a male genome assembly was used, and a non-redundant set of transcripts in GTF file format was constructed using StringTie v1.2.4 (Pertea et al. 2015). We filtered the transcripts for non-coding RNA (ncRNA) by extracting transcript sequences with BEDtools getfasta (Quinlan and Hall 2010) and removing transcripts with a BLAST hit to ncRNA sequences from *Poecilia formosa* (PoeFor_5.1.2), *Oryzias latipes* (MEDAKA1), *Gasterosteus aculeatus* (BROADS1) and *Danio rerio* (GRCz10) from Ensembl 104 (Flicek et al. 2014). Only genes with positional information on chromosomal fragments were kept for further analyses and the longest isoform for each target gene was selected. Lastly, we applied a minimum expression filter of 2 RPKM in at least half of the samples of each sex (minimum of two out of three individuals of either sex), resulting in 13,306 *P. reticulata*, 15,089 *P. wingei,* 13,156 *P. picta*, 14,468 *P. latipinna* and 21,861 *G. holbrooki* genic sequences.

We obtained coding sequences from the outgroup species *P. formosa* (PoeFor_5.1.2), *Xiphophorus maculatus* (Xipmac4.4.2) and *O. latipes* (MEDAKA1) from Ensembl 104 and extracted the longest isoform for each gene. Separately for each of our target species, we determined orthology across target and outgroup sequences using reciprocal BLASTn v2.7.1 (Altschul et al. 1990) with an e-value cut-off of $10e^{-10}$ and a minimum percentage identity of 30%. For genes with multiple blast hits, we chose the top hit based on the highest BLAST score. Our analysis resulted in 7,296 *P. reticulata*, 7,253 *P. wingei*, 7,786 *P. picta*, 7,251 *P. latipinna* and 6,477 *G. holbrooki* orthogroups (four-way 1:1 orthologs).

Estimating sequence divergence across orthogroups

For each contig of each orthogroup, we obtained open reading frames using *O. latipes* (MEDAKA1) protein-coding sequences from Ensembl 104 and BLASTx v2.3.0 with an e-value cut-off of $10e^{-10}$ and a minimum percentage identity of 30%, excluding orthogroups without BLASTx hits or valid protein-coding sequences. We aligned orthologous gene sequences with PRANK v170427 (Löytynoja and Goldman 2008), using the rooted tree (((Target species, *P. formosa*), *X. maculatus*), *O. latipes*), where the target species was in turn *P. reticulata*, *P. wingei*, *P. picta*, *P. latipinna* and *G. holbrooki*. Alignments were then filtered to remove gaps.

To avoid false positive signals of adaptive evolution, poorly aligned or error-rich regions were masked with SWAMP (Harrison et al. 2014). We ran SWAMP twice, first using a threshold of six nonsynonymous substitutions in a window size of 15 codons, and second using a threshold of two and a window size of five. This approach eliminates sequencing errors that cause short stretches of nonsynonymous substitutions as well as alignment errors that cause longer stretches of nonhomologous sequence due to variation in exon splicing or misannotation (Harrison et al. 2014). To select these thresholds, we first applied a range of masking criteria on our datasets. We then ran the branch-site test for positive selection on the target species branches for both the unmasked and masked datasets. We visually inspected the alignments of genes with the highest likelihood ratios and chose the masking criteria that was most efficient at reducing false positive rates. Finally, we discarded orthologs for which the alignment length was shorter than 300 bp following gap removal and masking, as these likely represent incomplete sequences.

To obtain divergence estimates for each orthogroup and calculate mean d_N/d_S across the target species branch, we used branch model (model=2, nssites=0) in the CODEML package in PAML v4.8 (Yang 2007), using the phylogeny ((Target species#1, *P. formosa*), *X. maculatus*, *O. latipes*), where the target species was successively *P. reticulata*, *P. wingei*, *P. picta*, *P. latipinna* and *G. holbrooki*. To avoid inaccurate divergence estimates dues to mutational saturation and double hits, orthologous genes with $d_S > 2$ were removed from subsequent analyses (Axelsson et al. 2008).

For each of the target species, we divided orthologs into autosomal and sex-linked categories based on their chromosomal location. The sex-linked category excluded genes in the

previously identified pseudoautosomal regions (0-5Mb and >26Mb of *P. reticulata* chromosome 12; >20Mb of *P. wingei*, *P. picta*, *P. latipinna* and *G. holbrooki* chromosomes that are syntenic to the guppy chromosome 12 (Darolti et al. 2019)). In addition, it has recently been discovered that the *P. reticulata* reference genome, which served as reference for scaffold ordering and orientation for our *P. reticulata* de novo genome assembly, has a large inversion on the X chromosome that is specific to the guppy strain on which the reference genome assembly was built and is not present in any of our lab or wild-caught samples (Darolti et al. 2020; Almeida et al. 2021). We thus corrected for this inversion when excluding genes from the PAR and assigning genes to the sex-linked category in *P. reticulata*.

Separately for each genomic category, we extracted the number of nonsynonymous substitutions (D_N), the number of nonsynonymous sites (N), the number of synonymous substitutions (D_S) and the number of synonymous sites (S). Taking into account alignment length, we calculated mean d_N and mean d_S as the ratio of the number of substitutions across all orthologs in that group divided by the number of sites ($d_N = D_N/N$; $d_S = D_S/S$), thus avoiding bias from short sequences and the issue of infinitely high d_N/d_S estimates due to very low d_S (Mank et al. 2007). We identified significant differences in d_N, d_S and d_N/d_S between genomic categories using 1,000 permutation test replicates and we used bootstrapping with 1,000 replicates to determine 95% confidence intervals. The strength of the Faster-X effect was calculated for each target species as the ratio of the rate of divergence for the X chromosome over the rate of divergence for the autosomes X(d_N/d_S):A(d_N/d_S).

Additional sequence divergence analyses for *P. reticulata*

For comparison to our estimates of rates of divergence based on *de novo* transcripts, we also estimated rates of sequence evolution in *P. reticulata* using publicly available coding sequences from Ensembl 104 (Guppy_female_1.0_MT). For this analysis, we followed the same steps outlined above in terms of extracting the longest isoform for each gene, identifying orthogroups, aligning target and outgroup sequences, masking and obtaining divergence estimates.

In addition, we have previously constructed high quality *P. reticulata* female *de novo* genome assemblies for replicate upstream and downstream populations of three rivers in Trinidad, Quare, Aripo, and Yarra (Almeida et al. 2021). To each genome assembly, we mapped *P.*

391   *reticulata* RNA-seq reads from our laboratory population and generated a non-redundant set

392   of transcripts for each wild population following the methods detailed above. For each

393   upstream and downstream population, we then estimated rates of coding sequence

394   evolution for the X chromosome, excluding the PAR, and the autosomes by following the

395   same pipeline described in the previous section.

Polymorphism data

To obtain polymorphism data, for each target species we first mapped female RNA-seq reads

to the genome assembly using the STAR v2.4.2a aligner in two-pass mode (Dobin et al. 2013),

and then called SNPs using SAMtools v1.3.1 mpileup (Li et al. 2009) with a minimum base

quality of 20 and VarScan v2.3.9 mpileup2snp (Koboldt et al. 2012) with a minimum coverage

of two, a minimum average quality of 20, a minimum variant allele frequency of $1e^{-1}$, *p* value

of 0.05 and strand filter set to on. We next imposed that the polymorphism dataset also

passes the filtering criteria used for calculating rates of sequence divergence. As such, we

identified codons for which all sites pass the minimum coverage threshold of 20, there are no

alignment gaps following alignment with PRANK, and no ambiguity data (Ns) following

masking with SWAMP. To make our dataset compatible for the McDonald-Kreitman test of

selection, we only kept genes with both divergence and polymorphism data. Finally, for each

genomic category, we calculated mean p_N and mean p_S as the ratio of the number of

polymorphisms across all orthologs in that group divided by the number of sites ($p_N=P_N/N$;

$p_S=P_S/N$).

Testing for selection using divergence and polymorphism data

For each species, we estimated the number of genes evolving under neutral and adaptive

evolution using the McDonald-Kreitman test (McDonald and Kreitman 1991). Neutral theory

predicts that the ratio of nonsynonymous to synonymous changes within species (p_N/p_S)

should be equal to that between species (d_N/d_S) (McDonald and Kreitman 1991). As such, the

McDonald-Kreitman test identifies signatures of positive selection, where there is an excess

of nonsynonymous substitutions relative to polymorphisms ($d_N/d_S>p_N/p_S$), and of relaxed

purifying selection, where there is a deficit of nonsynonymous substitutions compared to

polymorphisms ($d_N/d_S<p_N/p_S$). To test for deviations from neutrality, for each contig, we used

a 2 x 2 contingency table and a Fisher's Exact Test in R v3.6.2 (R Core Team 2015). The power

of the McDonald-Kreitman test is limited with low table counts, therefore we restricted this

analysis to genes for which the sum of each row and column in the contingency table was

equal to or greater than six (Begun et al. 2007; Andolfatto 2008).

As the McDonald-Kreitman test is a very conservative test, we also used the divergence and

polymorphism data to calculate the direction of selection statistic (DoS) (Stoletzki and Eyre-

Walker 2011). For each gene, we calculated the difference between the proportion of

nonsynonymous substitutions and polymorphisms ($\text{DoS}=D_N(D_N+D_S)-P_N(P_N+P_S)$), where

positive DoS values are indicative of positive selection. We used Fisher's Exact test in R to test

for significant differences in the proportion of genes under positive selection between the

autosomes and the X chromosome in each species.

Lastly, we used the polymorphism data alone to test for an excess or deficit of

nonsynonymous polymorphisms on the X chromosome compared to the autosomes, which

would suggest a relaxed constraint or purifying selection, respectively. For this, we

concatenated P_N and P_S estimates within each species and used Fisher's Exact test in R to test

for significant differences in P_N/P_S between the autosomes and the X chromosome.

Acknowledgements

437   This work was supported by the European Research Council (grant agreement 680951), a

438   Canada 150 Research Chair, and an NSERC Discovery Grant to J.E.M. We thank C. Lacy for the

439   fish illustrations and members of the Mank lab for helpful discussions and suggestions.

Author Contributions

441   J.E.M. and I.D. designed the research; I.D. and L.F. performed the research; I.D., L.F. and J.E.M.

442   analyzed the data; I.D., L.F. and J.E.M. wrote the manuscript.

References

Almeida P, Sandkam BA, Morris J, Darolti I, Breden F, Mank JE. 2021. Divergence and
remarkable diversity of the Y chromosome in guppies. Mol Biol Evol. 38:619–633.

Altschul SF, Gish W, Miller W, Myers E, Lipman DJ. 1990. Basic local alignment search tool. J
Mol Biol. 215:403–410.

Andolfatto P. 2008. Controlling Type-I Error of the McDonald–Kreitman Test in Genomewide
Scans for Selection on Noncoding DNA. Genetics. 180:1767–1771.

Ávila V, De Procé SM, Campos JL, Borthwick H, Charlesworth B, Betancourt AJ. 2014. Faster-
X effects in two drosophila lineages. Genome Biol Evol. 6:2968–2982.

Axelsson E, Hultin-Rosenberg L, Brandström M, Zwahlén M, Clayton DF, Ellegren H. 2008.
Natural selection in avian protein-coding genes expressed in brain. Mol Ecol. 17:3008–3017.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J,
Mayrose I, Ming R, et al. 2014. Sex determination: why so many ways of doing it? PLoS Biol.
12:e1001899.

Bechsgaard J, Schou MF, Vanthournout B, Hendrickx F, Knudsen B, Settepani V, Schierup
MH, Bilde T. 2019. Evidence for faster X chromosome evolution in spiders. Mol Biol Evol.
36:1281–1293.

Begun DJ, Holloway AK, Stevens K, Hillier LW, Poh YP, Hahn MW, Nista PM, Jones CD, Kern
AD, Dewey CN, et al. 2007. Population genomics: whole-genome analysis of polymorphism
and divergence in *Drosophila simulans*. PLoS Biol. 5:2534–2559.

Bolger AMLMUB. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 30:2114–2120.

Charlesworth B, Campos JL, Jackson BC. 2018. Theory and evidence from *Drosophila*. Mol
Ecol. 27:3753–3771.

Charlesworth B, Coyne JA, Barton NH. 1987. The relative rates of evolution of sex
chromosomes and autosomes. Am Nat. 130:113–146.

Charlesworth D, Bergero R, Graham C, Gardner J, Keegan H. 2021. How did the guppy Y
chromosome evolve? PLoS Genet. 17:e1009704.

Darolti I, Wright AE, Mank JE. 2020. Guppy Y chromosome integrity maintained by
incomplete recombination suppression. Genome Biol Evol. 12:965–977.

Darolti I, Wright AE, Sandkam BA, Morris J, Bloch NI, Farré M, Fuller RC, Bourne GR, Larkin
DM, Breden F, et al. 2019. Extreme heterogeneity in sex chromosome differentiation and
dosage compensation in livebearers. Proc Natl Acad Sci USA. 116:19031–19036.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29:15–21.

Flicek P, Amode MR, Barrell D, Beal K, Billis K, Brent S, …, Searle SMJ. 2014. Ensembl 2014.
Nucleic Acids Res. 42:749–755.

Fraser BA, Whiting JR, Paris JR, Weadick CJ, Parsons PJ, Charlesworth D, Bergero R, Bemm F,
Hoffmann M, Kottler VA, et al. 2020. Improved reference genome uncovers novel sex-linked
regions in the guppy (*Poecilia reticulata*). Genome Biol Evol. 12:1789–1805.

Harrison PW, Jordan GE, Montgomery SH. 2014. SWAMP : Sliding Window Alignment
Masker for PAML. Evol Bioinform. 10:EBO-S18193.

Jaquiéry J, Peccoud J, Ouisse T, Legeai F, Prunier-Leterme N, Gouin A, Nouhaud P, Brisson
JA., Bickel R, Purandare S, et al. 2018. Disentangling the Causes for Faster-X Evolution in
Aphids. Genome Biol Evol. 10:507–520.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: A fast spliced aligner with low memory
requirements. Nat Methods. 12:357–360.

Kirkpatrick M, Hall DW. 2004. Male-biased mutation, sex linkage, and the rate of adaptive
evolution. Evol. 58:437–440.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: Somatic mutation and copy number alteration discovery in cancer by exome sequencing. Genome Res. 22:568–576.

Kotrschal A, Rogell B, Bundsen A, Svensson B, Zajitschek S, Immler S, Maklakov AA, Kolm N. 2013. Experimental evidence for costs and benefits of evolving a larger brain. Curr Biol. 23:168–171.

Kousathanas A, Halligan DL, Keightley PD. 2014. Faster-X adaptive protein evolution in house mice. Genetics. 196:1131–1143.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The sequence alignment / map format and SAMtools. Bioinformatics. 25:2078–2079.

Löytynoja A, Goldman N. 2008. Phylogeny-aware gap replacement prevents errors in sequence alignment and evolutionary analysis. Science. 320:1632–1635.

Lu J, Wu C-I. 2005. Weak selection revealed by the whole-genome comparison of the X chromosome and autosomes of human and chimpanzee. PNAS. 102:4063–4067.

Mank JE, Axelsson E, Ellegren H. 2007. Fast-X on the Z: rapid evolution of sex-linked genes in birds. Genome Res. 17:618–624.

Mank JE, Vicoso B, Berlin S, Charlesworth B. 2010. Effective population size and the Faster-X effect: empirical results and their interpretation. Evolution. 64:663–674.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the ADH locus in *Drosophila*. Nature. 351:652–654.

Meisel RP, Connallon T. 2013. The faster-X effect: integrating theory and data. Trends Genet. 29:537–544.

Metzger DCH, Sandkam BA, Darolti I, Mank JE. 2021. Rapid evolution of complete dosage compensation in *Poecilia*. Genome Biol Evol. 13:evab155.

Morris J, Darolti I, Bloch NI, Wright AE, Mank JE. 2018. Shared and Species-Specific Patterns of Nascent Y Chromosome Evolution in Two Guppy Species. Genes. 9:238.

Nanda I, Schartl M, Feichtinger W, Epplen JT, Schmid M. 1992. Early stags of sex chromosome differentiation in fish as analysed by simple repetitive DNA sequences. Chromosoma. 101:301–310.

Nanda I, Schories S, Tripathi N, Dreyer C, Haaf T, Schmid M, Schartl M. 2014. Sex chromosome polymorphism in guppies. Chromosoma. 123:373–383.

Otto SP, Pannell JR, Peichel CL, Ashman T-L, Charlesworth D, Chippindale AK, Delph LF, Guerrero RF, Scarpino S V, McAllister BF. 2011. About PAR: the distinct evolutionary dynamics of the pseudoautosomal region. Trends Genet. 27:358–367.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol. 33:290–295.

Pinharanda A, Rousselle M, Martin SH, Hanly JJ, Davey JW, Kumar S, Galtier N, Jiggins CD. 2019. Sexually dimorphic gene expression and transcriptome evolution provide mixed evidence for a fast-Z effect in *Heliconius*. J Evol Biol. 32:194–204.

Pool JE, Nielsen R. 2007. Population size changes reshape genomic patterns of diversity. J
Org Evol. 61:30001–3006.

Quinlan AR, Hall IM. 2010. Genome analysis BEDTools: A flexible suite of utilities for
comparing genomic features. Bioinformatics. 26:841–842.

R Core Team. 2015. R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing. Available from: https//www.r-project.org.
Accessed August 25, 2021.

Rousselle M, Faivre N, Ballenghien M, Galtier N, Nabholz B. 2016. Hemizygosity enhances
purifying selection: Lack of fast-Z evolution in two satyrine butterflies. Genome Biol Evol.
8:3108–3119.

Sackton TB, Corbett-Detig RB, Nagaraju J, Vaishna L, Arunkumar KP, Hartl DL. 2014. Positive
selection drives faster-Z evolution in silkmoths. Evolution. 68:2331–2342.

Sandkam BA, Almeida P, Darolti I, Furman BLS, van der Bijl W, Morris J, Bourne GR, Breden
F, Mank JE. 2021. Extreme Y chromosome polymorphism corresponds to five male
reproductive morphs of a freshwater fish. Nat Ecol Evol. 5:939–948.

Stevenson BJ, Iseli C, Panji S, Zahn-Zabal M, Hide W, Old LJ, Simpson AJ, Jongeneel CV. 2007.
Rapid evolution of cancer/testis genes on the X chromosome. BMC Genomics. 8:1–11.

Stoletzki N, Eyre-Walker A. 2011. Estimation of the neutrality index. Mol Biol Evol. 28:63–70.

Tajima F. 1989. The effect of change in population size on DNA polymorphism. Genetics.
123:597–601.

Traut W, Winking H. 2001. Meiotic chromosomes and stages of sex chromosome evolution
in fish: zebrafish, platyfish and guppy. Chromosom Res. 9:659–672.

Tripathi N, Hoffmann M, Willing EM, Lanz C, Weigel D, Dreyer C. 2009. Genetic linkage map
of the guppy, *Poecilia reticulata*, and quantitative trait loci analysis of male size and colour
variation. Proc Biol Sci. 276:2195–2208.

Vicoso B, Bachtrog D. 2013. Reversal of an ancient sex chromosome to an autosome in
Drosophila. Nature. 499:332–335.

Vicoso B, Charlesworth B. 2006. Evolution on the X chromosome: unusual patterns and
processes. Nat Rev Genet. 7:645.

Vicoso B, Charlesworth B. 2009. The deficit of male-biased genes on the *D. melanogaster* X
chromosome is expression-dependent: A consequence of dosage compensation? J Mol Evol.
68:576–583.

Winge Ö. 1922. One-sided masculine and sex-linked inheritance in *Lebistes reticulatus*. J
Genet. 12:145–162.

Winge Ö. 1927. The location of eighteen genes in *Lebistes reticulatus*. J Genet. 18:1–43.

Winge Ö, Ditlevsen E. 1947. Colour inheritance and sex determination in *Lebistes*. Heredity.
1:65–83.

Wright AE, Darolti I, Bloch NI, Oostra V, Sandkam B, Buechel SD, Kolm N, Breden F, Vicoso B,
Mank JE. 2017. Convergent recombination suppression suggests role of sexual selection in

guppy sex chromosome formation. Nat Commun. 8:14251.

Wright AE, Harrison PW, Zimmer F, Montgomery SH, Pointer MA, Mank JE. 2015. Variation
in promiscuity and sexual selection drives avian rate of Faster-Z evolution. Mol Ecol.
24:1218–1235.

Wright AE, Mank JE. 2013. The scope and strength of sex-specific selection in genome
evolution. J Evol Biol. 26:1841–1853.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol.
24:1586–1591.

 Figure Legends

 Figure 1. Estimates of the rate of divergence (d_N/d_S) for autosomal and X-linked genes in (A)
P. reticulata, (B) P. wingei, (C) P. picta, (D) P. latipinna and (E) G. holbrooki. For all species,
the X chromosome category excludes genes on the PAR. Additionally, in *P. reticulata* and *P.
wingei*, the X chromosome category excludes genes in Stratum I. In *P. latipinna* and *G.
holbrooki*, the X chromosome represents the chromosome syntenic to guppy chromosome
12. 95% confidence intervals are based on bootstrapping with 1,000 replicates. Differences
between autosomal and X-linked loci are based on 1,000 replicates permutation tests. Only
significant differences are shown (*p* value < 0.01).

**Figure 2. Fast-X effect, calculated as the ratio of d_N/d_S for the X chromosome (excluding
the PAR) to that of the autosomes, across the poeciliids.** In *P. latipinna* and *G. holbrooki*,
the X chromosome represents the chromosome syntenic to guppy X. 95% confidence
intervals are based on bootstrapping with 1,000 replicates. Green color shading represents
the extent of sex chromosome degeneration (mild degeneration in light green to extreme
degeneration in dark green), while species in which the guppy chromosome 12 has not been
implicated as the sex chromosome are shown in grey.

Figure 3. Estimates of rates of evolution in wild *P. reticulata* populations. (A) Fast-X effect
across *P. reticulata* upstream, low predation populations (orange) and downstream, high
predation populations (black) in the Quare, Aripo and Yarra rivers. The Fast-X effect is
calculated as the ratio of the rate of nonsynonymous over synonymous substitutions for the
X chromosome, excluding the PAR, relative to that for the autosomes.

 Tables

Table 1. Differences between nonsynonymous and synonymous polymorphisms on the autosomes and X chromosome of each species. For all species, the X chromosome category excludes genes on the PAR.

Species	Category	P_N	P_S	p value[a]
P. reticulata	Autosomes	1,890	5,566	p = 0.76
	X chromosome	59	184	
P. wingei	Autosomes	650	1,783	p = 0.69
	X chromosome	18	56	
P. picta	Autosomes	837	2,178	p = 0.24
	X chromosome	12	47	
P. latipinna	Autosomes	1,165	3,265	p = 0.24
	Chromosome syntenic to guppy X	52	107	
G. holbrooki	Autosomes	1,212	4270	p = 0.08
	Chromosome syntenic to guppy X	28	144	

[a]*p* values from Fisher's Exact test

606 **Fig. 1**

608 **Fig. 2**

610 **Fig. 3**