An Extended Motif in the SARS-CoV-2 Spike Modulates Binding and Release of Host

Coatomer in Retrograde Trafficking

Debajit Dey[1,†], Suruchi Singh[1,†], Saif Khan[2,‡] , Matthew Martin[1,§], Nicholas J. Schnicker[2], Lokesh

Gakhar[2,3,¶], Brian G. Pierce[4,5], and S. Saif Hasan[1,6,7,*]

[1]Department of Biochemistry and Molecular Biology, University of Maryland School of Medicine,

Baltimore MD, USA

[2]Protein and Crystallography Facility, Carver College of Medicine, University of Iowa, Iowa City

IA, USA

[3]Department of Biochemistry, Carver College of Medicine, University of Iowa, Iowa City IA, USA

[4]W.M. Keck Laboratory for Structural Biology, University of Maryland Institute for Bioscience

and Biotechnology Research, Rockville MD, USA

[5]Department of Cell Biology and Molecular Genetics, University of Maryland, College Park MD,

USA

[6]University of Maryland Marlene and Stewart Greenebaum Cancer Center, University of

Maryland Medical Center, Baltimore MD, USA

[7]Center for Biomolecular Therapeutics, University of Maryland School of Medicine, Rockville

MD, USA

[†]These authors contributed equally

[‡]Current address: Bridge Institute, USC Michelson Center for Convergent Bioscience, University

of Southern California, Los Angeles CA, USA

[§]Current address: University of Pittsburgh Kenneth P. Dietrich School of Arts and Sciences,

Pittsburgh PA, USA

[¶]Current address: PAQ Therapeutics, Cambridge MA, USA

*Correspondence: Department of Biochemistry and Molecular Biology, University of Maryland

School of Medicine, Baltimore MD, USA; sshasan@som.umaryland.edu

Abstract

β-Coronaviruses such as SARS-CoV-2 hijack coatomer protein-I (COPI) for spike protein retrograde trafficking to the progeny assembly site in endoplasmic reticulum-Golgi intermediate compartment (ERGIC). However, limited residue-level details are available into how the spike interacts with COPI. Here we identify an extended COPI binding motif in the spike that encompasses the canonical K-x-H dibasic sequence. This motif demonstrates selectivity for αCOPI subunit. Guided by an *in silico* analysis of dibasic motifs in the human proteome, we employ mutagenesis and binding assays to show that the spike motif terminal residues are critical modulators of complex dissociation, which is essential for spike release in ERGIC. αCOPI residues critical for spike motif binding are elucidated by mutagenesis and crystallography and found to be conserved in the zoonotic reservoirs, bats, pangolins, camels, and in humans. Collectively, our investigation on the spike motif identifies key COPI binding determinants with implications for retrograde trafficking.

Introduction

β-Coronaviruses have been responsible for major human respiratory diseases in the last two decades. In 2002, the severe acute respiratory syndrome coronavirus (SARS-CoV) was implicated in an epidemic first reported in China before spreading to 27 countries, which resulted in 774 deaths [1]. A decade later, Middle East respiratory syndrome (MERS) was reported in Saudi Arabia in 2012 with over 30% fatality in patients [2,3]. Most recently, the novel SARS-CoV-2 has been implicated in the COVID-19 global pandemic that has claimed over four million lives. Current efforts to contain the pandemic are focused primarily on vaccinations using the viral spike protein that is responsible for SARS-CoV-2 entry into host cells [4,5]. Fundamental insights into spike biogenesis will advance the understanding of how β-coronaviruses exploit host resources during viral infection and may potentially lead to the development of novel therapeutics.

The trimeric β-coronavirus spike is organized into an ecto-domain, a transmembrane domain, and a cytosolic domain [6,7] (**Figure 1**). In infected cells, the newly synthesized spike protein in ER is transported first to the Golgi, and then from Golgi to the ERGIC compartment, which is the site of β-coronavirus progeny assembly [8-10]. This retrograde trafficking of the post-translationally modified spike from Golgi to ERGIC involves a cytosolic dibasic motif, K-x-H-x-x (Lys-x-His, where x is any amino acid) [9,10] (**Figure 1**). Such C-terminal dibasic motifs and variants such as K-x-K-x-x and K-K-x-x are widely reported in the cytosolic tail of host membrane proteins that undergo retrograde trafficking [11-14]. As such, the β-coronavirus spike demonstrates molecular mimicry of dibasic trafficking motifs [9,10]. This recycling of spike protein has been suggested to enhance interactions with the viral membrane (M) protein localized in ERGIC during progeny assembly and is crucial for spike maturation [9,10,15]. These observations establish a key role of the dibasic motif in SARS-CoV and SARS-CoV-2 infection and propagation cycles. Interestingly, the spike dibasic motif and adjacent residues are completely conserved in sarbecoviruses, i.e., SARS-CoV and SARS-CoV-2, although sequence divergence in residues neighboring the dibasic motif is noted in MERS-CoV [9,10].

On the host side, retrograde trafficking is mediated by the interactions of dibasic motifs with the coatomer protein-I (COPI) complex [11,16]. Seven subunits, namely, α, β, β', γ, δ, ε and ζ, assemble into a COPI complex on retrograde trafficking vesicles that carry cargo [17-24]. Prior genetic, biochemical, biophysical, and structural investigations have shown that the binding site for host cargo dibasic motifs maps to the N-terminal β-propeller WD40 domains of α and β'COPI subunits, which are structural homologs [11,25-29]. Mutagenesis analyses of α and β' subunit N-

73   terminal WD40 domains have identified residues critical for binding of host protein dibasic motifs
74   [27]. This and another study [28] provided important structural details of how dibasic host and viral
75   peptides bind to αCOPI-WD40 and β'COPI-WD40 domains.

76   Cellular and biochemical investigations in recent years and during the ongoing COVID-19
pandemic have suggested a role of COPI interactions in sarbecovirus spike trafficking,
maturation, glycan processing, and syncytia formation during infection [10,15,30]. These studies have
established a platform to investigate the underlying chemistry of spike-COPI interaction. For
instance, it is not known whether the K-x-H motif is sufficient to determine the strength of this
interaction, whether adjacent residues in the spike cytosolic tail play a role in this binding, and
what are the sequence determinants of spike-COPI disassembly crucial for spike release in
ERGIC. On the host side, it is presently not known which COPI residues are critical for spike
interactions. As such, key facets of this initial binding event in spike trafficking remain largely
unknown for SARS-CoV-2 as well as SARS-CoV. In the present investigation, we address these
questions using a combination of bio-layer interferometry (BLI), molecular modeling,
mutagenesis, X-ray crystallography, and an *in silico* analysis of the human membrane proteome.
Employing a sarbecovirus spike hepta-peptide corresponding to the K-x-H-x-x motif, we identify
critical residues in αCOPI-WD40 for hepta-peptide binding and demonstrate structural alterations
in an αCOPI-WD40 mutant. Amino acid propensity is described in human dibasic motifs and
adjacent downstream residues, and mutagenesis experiments driven by this analysis provide
insights into how sarbecovirus spike modulates strength of binding to COPI. Collectively, our
study advances the structural and biophysical understanding of how the dibasic motif hijacks
COPI for spike retention in endo-membranes and trafficking to the plasma membrane during
sarbecovirus infections.

Results

Direct binding of sarbecovirus spike hepta-peptide is selective for αCOPI-WD40 domain
98   In this investigation, we heterologously expressed and purified the N-terminal WD-40
domain of β'COPI-WD40 (residues 1-301) from *Saccharomyces cerevisiae* and αCOPI-WD40
(residues 1-327) from *Schizosaccharomyces pombe* (**Supplementary Figures S1, S2**). Although
SARS-CoV-2 infects mammals, we chose these yeast constructs for two reasons. First, the
putative interaction interface for dibasic peptides is conserved between these constructs and
theαCOPI and β'COPI orthologs in humans, COPA and COPB2, respectively [27,28]. Second, these
domains have been previously crystallized and structurally characterized [27,28]. This is consistent

with our aim of understanding the structural basis of spike-COPI interactions. Although β'COPI-

WD40 was expressed in *E. coli* as described previously, αCOPI-WD40 has reportedly presented

challenges in protein production [27,28]. Hence, the αCOPI-WD40 construct was expressed in

Expi293 cells. This generated yields of purified αCOPI-WD40 of ~4 mg per 100 ml cell culture

volume. For comparison with a previously published construct of αCOPI-WD40 expressed in *E.*

coli [28], the crystal structure of the purified αCOPI-WD40 domain was determined to 1.8Å resolution

(**Supplementary Table T1**). The αCOPI-WD40 domain is organized into a β-propeller and is

consistent with previously described structures of αCOPI-WD40 (Cα root-mean-square-deviation

is <0.5Å) [28]. However, a peripheral loop and a short α-helix (Gly^{168}-Ala^{188}, shown with an arrow in

Figure 2a) demonstrate substantial differences from previously described αCOPI-WD40

structures likely due to altered crystal packing. An N-terminal acetylation of the αCOPI-WD40

polypeptide was identified in the structure. Importantly, the αCOPI-WD40 domain interface for

putative interactions with dibasic motifs is similar between the structure determined here and

previously published structures [28].

119   Recent investigations of SARS-CoV-2 spike and previously of SARS-CoV spike with COPI

have employed cellular lysates [10,15,30]. Hence, we asked if there is direct interaction between the

purified components. To address this question, we first established a BLI assay to test this

interaction. Monomeric constructs of the spike cytosolic domain demonstrate similar interactions

with COPI as the trimeric cytosolic domain [10]. Hence, a hepta-peptide representing a monomeric

dibasic motif in the sarbecovirus spike (^{1267}Gly-Val-Lys-Leu-His-Tyr-Thr^{1273}) was synthesized with

an N-terminal biotin tag attached via a linker. The hepta-peptide C-terminus has a free carboxylate

to mimic the C-terminus of a polypeptide. This hepta-peptide (or its sequence variants) was

immobilized on a streptavidin biosensor for BLI analysis. The purified α or β'COPI-WD40 domain

was provided as the analyte in the BLI assay (**Figure 2b-h, Table 1**). It was observed that the

spike hepta-peptide binds directly to the purified αCOPI-WD40 domain with an equilibrium

dissociation constant (K_D)=4.17±0.04 μM and a kinetic K_D=2.75±0.09 μM at pH 7.5 (**Figure 2b,**

Table 1). A scrambled sequence of this hepta-peptide showed no detectable interaction with

αCOPI-WD40 (**Figure 2c**) suggesting that the binding of the wild-type hepta-peptide was

sequence specific. This binding analysis demonstrates that the C-terminal peptide of the

sarbecovirus spike contains sufficient sequence and structural information to interact directly with

αCOPI-WD40. This is consistent with prior COPI binding analyses with peptides corresponding

to host dibasic motifs [27,28]. In contrast, this sarbecovirus hepta-peptide demonstrates a lack of

binding to β'COPI-WD40 (**Figure 2d**). This selectivity for αCOPI-WD40 is consistent with that

reported for a similar spike hepta-peptide (^{1377}Phe-Glu-Lys-Val-His-Val-Gln^{1383}) from porcine

epidemic diarrhea virus (PEDV), an α-coronavirus [28]. Mutation of the K-x-H motif to K-x-K in the sarbecovirus mutant peptide, i.e., [1267]Gly-Val-Lys-Leu-Lys-Tyr-Thr[1273], demonstrates similar selectivity for αCOPI-WD40 over β'COPI-WD40 (**Figure 2e, f**). Cellular studies suggest enhanced interactions of this mutant spike sequence with COPI subunits [15,30]. It is likely that this mutation affects the local conformation of the full-length spike protein leading to modulation of COPI interactions in a cellular environment.

One of the first analyses of COPI involvement in SARS-CoV spike trafficking employed cellular lysate pull-downs to show enhanced spike-COPI interactions under acidification to pH 6.5 [10]. As the sarbecovirus hepta-peptide contains His[1271] in the K-x-H motif, we asked if this acidification would affect hepta-peptide interaction with αCOPI-WD40. A nearly 3-fold enhancement in binding between the wild-type hepta-peptide and αCOPI-WD40 (equilibrium K_D=1.40±0.15 μM) was observed upon acidification to pH 6.5, likely due to partial protonation of the His residue in the hepta-peptide K-x-H motif (**Figure 2g, Table 1**). Relative to pH 7.5, this lower pH accelerated the association rate of the hepta-peptide with αCOPI-WD40 by a factor of 1.7 while concomitantly suppressing complex dissociation by another 1.7-fold (**Table 1**). As such, acidification was inferred to be a key factor in stabilizing the hepta-peptide complex with αCOPI-WD40. All subsequent BLI assays were performed at pH 6.5.

The binding of β'COPI-WD40 to the wild type spike peptide is still substantially weaker than that with αCOPI-WD40 in low pH (**Figure 2h**).

The terminal residues in the spike hepta-peptide are key modulators of αCOPI-WD40 binding and dissociation

Homology modeling was employed to analyze the structural basis of interaction between SARS-CoV-2 spike hepta-peptide and αCOPI-WD40 domain. This modeling was based on a prior co-crystal structure of αCOPI-WD40 domain with a dibasic peptide [28]. Apart from the Lys[1269] and His[1271] residues in the K-x-H motif, the terminal Tyr[1272]-Thr[1273] residues in the hepta-peptide are within interaction distance of αCOPI-WD40 surface residues (**Figure 3a**). This is intriguingly suggestive of a role of these two spike residues in binding αCOPI. Interestingly, the two N-terminal residues in the hepta-peptide, i.e., Gly[1267]-Val[1268], make no contact with the αCOPI-WD40 surface. To evaluate the role of the spike residues in binding αCOPI-WD40, *in silico* alanine scanning mutagenesis of the modeled spike hepta-peptide was performed (**Table 2**). The spike Lys[1269] and His[1271] residues that constitute the K-x-H dibasic motif are predicted to be most crucial for binding αCOPI-WD40 domain. The *in silico* mutations of these residues to Ala yield highly unfavorable free energy changes suggestive of substantially weakened binding to αCOPI-WD40 (**Table 2**).

The Ala mutation of Tyr^{1272} in the spike peptide implies a substantial role of this residue in
stabilization of the spike-αCOPI-WD40 complex. This is likely due to the side chain interaction
between the oxygen atom in Tyr^{1272} side-chain hydroxyl group with the αCOPI-WD40 His^{31} side-
chain NE2 atom, along with main chain interactions of Tyr^{1272}. The terminal residue in the spike,
i.e., Thr^{1273}, is predicted to contribute modestly to the stabilization of the complex with αCOPI-
WD40 (**Table 2**).

Next, we tested this *in silico* model of interactions between the spike hepta-peptide and
αCOPI-WD40 using BLI assays (**Figure 3b-g**). The mutagenesis of Lys^{1269} or His^{1271} in the spike
K-x-H motif to Ala residues abolished binding to αCOPI-WD40 (**Figure 3b, c**). As expected, the
dual Ala mutation of the K-x-H motif does not demonstrate any substantial binding to αCOPI-
WD40 (**Figure 3d**). As such, both basic residues in SARS-CoV-2 spike K-x-H motif are individually
and concomitantly required for αCOPI-WD40 binding. Replacement of either residue is sufficient
to disrupt αCOPI-WD40 binding to the spike hepta-peptide. These data are consistent with the *in*
silico predictions described above as well as with cellular assays on SARS-CoV and SARS-CoV-
2 spike trafficking [10,15,30]. We next tested the contribution of spike hepta-peptide Tyr^{1272} residue to
αCOPI-WD40 binding. A BLI assay of a mutant Tyr^{1272}→Ala spike hepta-peptide (^{1267}Gly-Val-Lys-
Leu-His-<u>Ala</u>-Thr^{1273}) yielded an equilibrium K_D=3.77±0.34 µM, which is 2.7-fold weaker than the
wild-type spike peptide (**Figure 3e, Table 1)**. Although this mutation only reduced the rate of
complex formation by 1.3-fold relative to the wild-type hepta-peptide (**Table 1**), it accelerated
complex dissociation by 1.9-fold (**Table 1**). This suggests weakened interactions of the spike
hepta-peptide with αCOPI-WD40 when the aromatic side chain interactions of Tyr^{1272} are
abrogated. Collectively, this BLI analysis indicates that Tyr^{1272} is important for complex stability.
These experimental results are consistent with the above described *in silico* model (**Table 3**).
Next, we evaluated the C-terminal position Thr^{1273} in the spike. A BLI assay of a mutant
Thr^{1273}→Ala hepta-peptide (^{1267}Gly-Val-Lys-Leu-His-Tyr-<u>Ala</u>^{1273}) yielded an equilibrium
K_D=1.13±0.05 µM, which is similar to the wild type hepta-peptide (K_D=1.40±0.15 µM) (**Figure 3f,**
Table 1). This Thr^{1273}→Ala mutation caused a slowing down of αCOPI-WD40 association-
dissociation kinetics by 2.3 and 1.9-fold respectively (**Table 1**). This suggested that a β-branched
residue at the C-terminus may be an important determinant in complex formation kinetics. To
probe further, we generated a Thr^{1273}→Val mutant hepta-peptide, which maintains a β-branched
residue at the C-terminus. Val has methyl groups at the two side chain γ positions, which replace
a methyl and a hydroxyl group at equivalent γ positions in Thr. A BLI assay of this mutant hepta-
peptide showed a 2.3-fold weakened interaction with αCOPI-WD40 relative to the wild type, with
an equilibrium K_D=3.20±0.04 µM (**Figure 3g, Table 1**). Interestingly, this mutant demonstrated

1.5-fold slower association kinetics and 1.5-fold more rapid dissociation than for the wild type
hepta-peptide (**Table 1**). Compared to the Thr[1273]→Ala hepta-peptide, this Thr[1273]→Val mutant
weakened binding by 2.8-fold while accelerating αCOPI-WD40 complex association and
dissociation kinetics by 1.5 and 2.8-fold respectively (**Table 1**). These data suggest that the spike
C-terminal Thr residue side-chain provides interactions that modulate dissociation of the complex
with αCOPI-WD40. Interestingly, a prior analysis implicated β-branched residues at the
penultimate position in the PEDV spike sequence in modulating interactions with COPI-WD40
domains [28].
215
Electrostatics of spike hepta-peptide C-terminus drive dissociation from αCOPI-WD40
We performed an *in silico* analysis of the human proteome to gain insights into whether
the spike extended dibasic motif demonstrates consistency with host dibasic motifs and their
environment. We identified 119 sequences predicted to be membrane proteins that terminate with
K-x-H-x-x and K-x-K-x-x dibasic motifs (**Supplementary Table T2**). These sequences were
aligned and analyzed for the frequency of 20 amino acids at each of the positions in the dibasic
motif and the two terminal residues following the motif (**Figure 4a, Supplementary Table T3**).
This analysis revealed novel details about the dibasic motif. First, it was inferred that the
predominant dibasic motif is K-x-K-x-x rather than K-x-H-x-x by nearly an order of magnitude.
Second, only a low frequency (0.07) of the sequences has an aromatic residue at the penultimate
position, which corresponds to Tyr[1272] in the SARS-CoV-2 spike. β-Branched residues Leu, Ile,
Val, Ser, and Thr are found at a high frequency of 0.38 at this penultimate position. Third, acidic
residues at the C-terminus are observed in nearly a quarter (frequency=0.24) of the sequences.
Overall, with a frequency of 0.42, the C-terminal position has a strong tendency to be occupied
by charged residues such as Arg, Asp, Glu, His, and Lys. Hydroxyl side chain containing Thr,
which corresponds to Thr[1273] in SARS-CoV-2 spike, is a low frequency residue (0.05). In our *in*
silico model of the hepta-peptide complexed with αCOPI-WD40, the side chain of this Thr[1273]
residue is within interaction distance of a cluster of basic residues in αCOPI-WD40 (Arg[13], Lys[15],
and Arg[300], **Figure 4b**). Hence, we hypothesized that the presence of a charged residue at this
spike position would modulate interactions with αCOPI-WD40. This was supported by our *in silico*
analysis, which predicted stabilization of the complex when an acidic Glu residue replaced Thr[1273]
in the spike hepta-peptide (**Table 2**).
We next tested the role of this spike C-terminal residue in modulating αCOPI-WD40
binding using a BLI assay (**Figure 4c-e**). We employed three distinct mutations of the spike hepta-
peptide at this position, i.e., acidic (Glu), basic (Arg), and neutral (Gln) (**Figure 4c-e**). The

presence of an acidic Glu residue at the C-terminus was found to substantially strengthen binding

of the hepta-peptide to αCOPI-WD40 with an equilibrium $K_D=0.31\pm0.01$ μM, which is 4.5-fold

tighter than the binding of the wild type hepta-peptide sequence (**Figure 4c, Table 2**). This is

consistent with our *in silico* model and is strongly suggestive of an electrostatic interaction

between the Glu[1273] side chain and αCOPI Arg[13], Lys[15], and Arg[299] side chains to stabilize the

complex. Furthermore, the rate of dissociation of αCOPI-WD40 domain from the Glu[1273] hepta-

peptide is 6-fold slower than that of the wild-type spike hepta-peptide (**Table 1**). In fact, during the

time course of our experiment, we did not observe complete dissociation of this complex with the

Glu[1273] containing hepta-peptide. To eliminate the possibility of non-specific interactions, we

employed a hepta-peptide with a scrambled sequence (**Supplementary Figure S3**). Next, we

tested whether modifying side-chain charge at the C-terminal position of the spike hepta-peptide

affects complex formation with αCOPI-WD40. Relative to Glu[1273], the binding between αCOPI-

WD40 domain was weakened when neutral Gln[1273] was substituted into the hepta-peptide

(equilibrium $K_D=0.78\pm0.04$ μM, **Figure 4d**). However, the binding of the hepta-peptide with Gln[1273]

was still 1.8-fold tighter than that of the wild-type hepta-peptide (**Table 1**). In contrast, basic Arg[1273]

in the spike hepta-peptide (equilibrium $K_D=1.73\pm0.24$ μM) yielded an interaction strength similar

to the wild type sequence (equilibrium $K_D=1.40\pm0.15$ μM, **Figure 4e**). The amide carbonyl group

in the Gln[1273] side-chain likely interacts with the basic residue cluster on αCOPI-WD40 through

hydrogen bonding. This stabilizing interaction is disrupted when Gln is replaced by Arg[1273] in the

spike hepta-peptide. Intriguingly, the rate of association with αCOPI-WD40 was slowed down

relative to the wild type hepta-peptide by a factor of 2.6 for Glu[1273] and Gln[1273], whereas it was

similar to that of Arg[1273] containing hepta-peptide (**Table 2**). Overall, these data establish a critical

role of the C-terminal position in the SARS-CoV-2 spike in modulating binding to αCOPI-WD40.

Low frequency of acidic residues at the spike C-terminus

Given the inference that a C-terminal acidic residue strengthens binding to αCOPI-WD40,

which likely interferes with spike release, we asked if a typical coronavirus spike demonstrates a

paucity for such C-terminal acidic residues. A sequence and phylogenetic analysis of the five C-

terminal residues in the spike protein of coronaviruses was performed to determine the frequency

of Asp and Glu at the C-terminus (**Figure 5**). None of the coronavirus spike proteins that

demonstrate the K-x-H-x-x, K-x-K-x-x, or K-x-R-x-x dibasic motif has a C-terminal acidic residue.

It has been suggested that bats are the likely genetic source of human β-coronaviruses [31-

33]. Apart from bats, during the SARS-CoV epidemic and the ongoing SARS-CoV-2 pandemic,

zoonotic reservoirs such as civets and pangolins have been suggested to be involved in

coronavirus transmission [33-38]. Our phylogenetic analysis showed that the extended coatomer

binding motif in the spike, i.e., [1269]Lys-Leu-His-Tyr-Thr[1273], is conserved in coronavirus isolates

from these animals (**Figure 5**). Moreover, this conservation of the extended motif is seen in the

WHO deemed emerging variants of concern for SARS-CoV-2, i.e., α (Pango lineage B.1.1.7), β

(B.1.351), γ (P.1), δ (B.1.617.2), and the latest O (B.1.1.529) [39]. This is indicative of a strong

selection pressure to maintain this COPI-interacting sequence in the spike protein.

A polar αCOPI-WD40 interface for spike hepta-peptide binding

We subsequently focused our attention on the spike binding residues in αCOPI-WD40.

The *in silico* modeling of SARS-CoV-2 spike hepta-peptide shows that interaction with αCOPI-

WD40 domain involves predominantly polar residues (**Table 3**). Amongst these residues, Arg^{57},

Asp^{115}, and Tyr^{139} provide the highest level of stabilization to spike hepta-peptide binding. The

Arg^{57} side-chain interacts with the main chain carbonyl of spike His^{1271}, Tyr^{1272}, and Thr^{1273} (**Figure**

6a). The Asp^{115} side-chain forms a bond with the terminal NZ atom in the spike Lys^{1269} side-chain

(**Figure 6b**). This side-chain of spike Lys^{1269} is further stabilized by an interaction with the hydroxyl

oxygen in Tyr^{139} side-chain (**Figure 6c**). Hence, the side chains of αCOPI-WD40 Arg^{57}, Asp^{115},

and Tyr^{139} residues provide an extensive and polar interaction network for binding of the spike

hepta-peptide. Therefore, mutagenesis of these three αCOPI-WD40 residues to Ala is predicted

to disrupt interactions with the spike hepta-peptide as suggested by our *in silico* analysis (**Table**

3).

Next, the role of αCOPI-WD40 Arg^{57}, Asp^{115}, and Tyr^{139} residues in binding the spike

hepta-peptide was tested. We generated three single-site mutants of αCOPI-WD40 wherein Arg^{57},

Asp^{115}, and Tyr^{139} residues were individually mutated to Ala. These mutants were expressed

heterologously and purified from Expi293 cells. Analysis by size-exclusion chromatography (SEC)

suggested an overall similarity in hydrodynamic radius with the wild-type αCOPI-WD40 domain

(**Supplementary Figure S2**). These three mutants were analyzed for binding to the wild-type

spike hepta-peptide by BLI assays. None of the three mutants demonstrates any binding to the

wild type sequence of the spike hepta-peptide (**Figure 6d-f**). This demonstrated that αCOPI-

WD40 residues Arg^{57}, Asp^{115}, and Tyr^{139} are individually critical for binding the spike hepta-

peptide. Disruption of even one of these interactions is likely sufficient to destabilize the spike-

COPI complex.

Structural basis of conformational changes in an αCOPI-WD40 mutant

The results of αCOPI-WD40 Arg[57], Asp[115], and Tyr[139] mutagenesis led us to ask if the loss
of binding to the spike hepta-peptide was due to disruption of a single critical interaction or due to
larger alterations in the protein structure. This is because the surface of αCOPI-WD40 contains
an intricate network of residues with charged and polar side-chains. Mis-sense mutations that
alter charge balance could modify αCOPI-WD40 conformations. To address this question, we
crystallized αCOPI-WD40 Arg[57]→Ala and Tyr[139]→Ala mutants. The αCOPI-WD40 Asp[115]→Ala
mutant did not yield crystals in the conditions we tested. The crystal structures of αCOPI-WD40
Arg[57]→Ala and Tyr[139]→Ala mutants were determined by X-ray diffraction to a resolution of 1.2Å
and 1.5Å, respectively (**Table T1**).

The αCOPI-WD40 Arg[57]→Ala mutant structure demonstrated novel structural alterations
that had previously not been reported in the crystal structures of wild-type αCOPI-WD40 or the
related β'COPI-WD40 (**Supplementary Figure S4a**). The mutation of Arg[57] to an Ala residue
generated a cavity in the spike hepta-peptide binding site. This change led to a 62° rotation of a
nearby Tyr[97] residue side chain into the newly generated cavity in αCOPI-WD40 (**Supplementary**
Figure S4b). In parallel, the residue Asp[73] underwent a substantial conformational change. This
residue interacts with the side chain of Arg[57] in the wild-type αCOPI-WD40 structure. However,
the loss of stabilizing interactions from the Arg[57] side chain and the reorientation of Tyr[97] caused
the Asp[73] side chain to rotate away by 73° from its initial position (**Supplementary Figure S4b**).
These conformational changes are accompanied by a 1.1Å and 0.8Å movement of Tyr[139] and
His[31] side chains respectively closer towards the spike hepta-peptide as inferred from our *in silico*
model. In contrast, the side chain of Lys[15] moves 1.7Å away from the inferred hepta-peptide
position. As such, the binding site and its vicinity demonstrate a substantially modified interaction
network in Arg[57]→Ala mutant.

Next, we asked if Arg[57]→Ala mutation and the associated rotameric changes caused any
main chain reorganization in αCOPI-WD40. To obtain a global overview of changes in the main
chain geometry, the differences in Ramachandran angles were calculated between
corresponding residues in the wild type and Arg[57]→Ala crystal structures (**Supplementary Figure**
S4c). The top peak in this difference Ramachandran plot, i.e., peak 1, in this analysis corresponds
to a substantial main chain twist at Gly[72], which is near the Arg[57]→Ala mutation site. This
conformational change is associated with the Asp[73] side chain rotation and repositioning of the
main chain atoms from Gly[72] to Val[77], which are pushed away from the domain core consistent
with the reorientation of the Asp[73] side chain (**Supplementary Figure S4d**). Peaks 2 and 4 in this
difference Ramachandran plot correspond to changes in surface loops that are 19Å and 25Å from
the mutation site and are likely due to crystal contacts. Peaks 3 and 6 correspond to main chain

rearrangement in the mutation site, i.e., Arg[57]→Ala. This is likely a combination of the mutation

and modifications to side chain rearrangements in the neighborhood of Ala[57] described above.

Peak 5 is associated with a surface loop 31Å from the mutation site. This loop demonstrates weak

electron density and is only partly ordered. Hence, three of the top six peaks, i.e., 1, 3, and 6, in

this analysis are associated with considerable rearrangement of the αCOPI-WD40 surface upon

the mutation of basic Arg[57] to neutral Ala including the site for spike hepta-peptide binding.

In contrast to the Arg[57]→Ala substitution, the crystal structure of αCOPI-WD40 Tyr[139]→Ala

mutant demonstrated no substantial changes as compared to the wild-type structure

(**Supplementary Figure S4c, e**). No major rearrangements of side chains or main chains were

observed. This lack of conformational rearrangement contrasts with the structural changes in the

αCOPI-WD40 Arg[57] →Ala mutant structure. It is likely that the electroneutral change from Tyr[139]

to Ala does not perturb the local electrostatic surface sufficiently to alter protein conformation.

Hence, disruption of spike hepta-peptide binding in this Tyr[139]→Ala mutant is due to the loss of a

single side chain hydroxyl group. Collectively, the crystal structures of αCOPI-WD40 Arg[57]→Ala

and Tyr[139]→Ala mutants reveal distinct and contrasting structural principles by which spike hepta-

peptide binding is disrupted.

An analysis of crystal packing in the αCOPI-WD40 structures reported here showed that

the peptide binding site residues are in contact with symmetry related chains. We subsequently

asked if distinct crystal packing may have contributed to these different structural consequences

of Arg[57]→Ala and Tyr[139]→Ala mutations. However, similar crystal packing interactions are

provided by residues His[267] and Lys[309] from a symmetry-related αCOPI-WD40 chain to the spike

hepta-peptide binding site. As such, conformational differences in these two αCOPI-WD40

mutants are due to altered interaction chemistry in the hepta-peptide binding site.

Conservation of αCOPI-WD40 residues critical for spike hepta-peptide binding

The mutagenesis, BLI, and crystallographic analyses described here are focused on

αCOPI. Hence, we asked if αCOPI Arg[57], Asp[115], and Tyr[139] residues are conserved in bats,

pangolins, camels, and humans, which have been implicated as zoonotic reservoirs and hosts for

β-, coronaviruses [40-44]. Overall, αCOPI is 96.5-99.8% identical in these multicellular higher

organisms (**Supplementary Table T4**). In contrast, αCOPI conservation is relatively moderate

between these organisms and yeast at 46.8-47.1% sequence identity. An analysis of αCOPI

sequence conservation across 150 species using the CONSURF server [45] demonstrated that

Arg[57] and Asp[115] are completely conserved whereas Tyr[139] is replaced by Phe or Trp in 5.3% and

0.7% of the sequences, respectively (**Supplementary Data**). Importantly, all these three αCOPI

residues are found to be 100% identical in yeast, bat, pangolin, camel, and human αCOPI. This
suggests an evolutionary pressure on these three residues in binding dibasic motifs in host
proteins, which is exploited by the sarbecovirus spike to hijack the host COPI machinery. This
conservation of the three αCOPI residues extends to chicken, which is has been suggested to be
a host for γ- and δ-coronaviruses [31]. αCOPI residues such as Lys[15] and His[31] that are suggested
to be involved in spike hepta-peptide binding by our *in silico* analysis demonstrate complete
conservation. Interestingly, this conservation extends to *S. cerevisiae* β'COPI wherein αCOPI
residues Arg[57] and Asp[115] are replaced by Arg[59] and Asp[117] in β'COPI, respectively. However,
αCOPI Tyr[139] is semi-conserved and is replaced by Phe[142] in β'COPI.

Discussion

Once the sarbecovirus spike is delivered from ER to Golgi, its trafficking to the progeny
virus assembly site consists of three distinct steps, i.e., spike-COPI binding in donor membranes
such as cis-Golgi, inter-organelle trafficking, and dissociation of spike-COPI at the destination,
which is ERGIC [8]. This trafficking pathway can be disrupted by either weakening of spike-COPI
binding leading to premature complex dissociation or enhanced stability of this complex, which
interferes with spike release. This is supported by recent cellular imaging and biochemical
analysis of the SARS-CoV-2 spike protein [15]. Hence, elucidating the determinants of spike-COPI
interactions is fundamental to understanding sarbecovirus assembly.

Employing a spike hepta-peptide and a purified aCOPI-WD40 domain, the present
investigation expounds on the biophysical and structural bases of spike-COPI interactions. We
demonstrate that direct binding of purified αCOPI-WD40 domain to the SARS-CoV-2 spike hepta-
peptide is modulated by an extended coatomer binding motif that stretches beyond the spike K-
x-H residues. Our data show that residues such as acidic Glu in the C-terminal position in the
spike likely interact with complementary charged basic residues in αCOPI-WD40. This interaction
strengthens spike binding to the host αCOPI. This analysis is consistent with a recent preprint
that shows a key role of this SARS-CoV-2 spike C-terminal position in pull-down assays of the
spike cytosolic domain with COPI subunits [30]. A second cellular investigation has recently shown
that the inferred stabilization of the spike-COPI complex by a Lys[1269]-x-His[1271]→Lys[1269]-x-Lys[1271]
spike mutation has dramatic effects on SARS-CoV-2 spike processing and trafficking (Jennings
et al., 2021). This functional analysis suggests a key role of spike-COPI complex dissociation in
modulating spike trafficking and function. Hence, it is likely that residues that strengthen spike-
COPI complex stability beyond that from wild type interactions are avoided in the spike C-
terminus. This includes acidic Glu and unbranched Ala residue that stabilize the αCOPI-WD40

domain as demonstrated in the present investigation. Interestingly, our analysis of the human membrane proteome suggests that the occurrence of a charged residue such as Glu and β-branched residues is a high probability event at the C-terminus of dibasic motifs. This raises an intriguing question of whether such charged residues present a structural and biophysical disadvantage to spike-COPI interactions, and hence, are selected against in sarbecoviruses. Such a highly stabilized complex may not undergo dissociation in ERGIC to release the spike for processing and downstream virion assembly. Interestingly, acidic residues are absent from the C-terminus of coronavirus spike proteins that have a dibasic motif for COPI dependent trafficking. We note that complex formation of Glu^{1273} or Gln^{1283} containing spike hepta-peptide is substantially slower than in the wild type. Even Arg^{1273}, which lacks charge complementarity with αCOPI-WD40 residues shows slower association kinetics. Glu, Gln, and Arg have long side chains unlike Thr^{1273}, which is suggestive of a role of side chain size in modulating interactions with COPI.

On the coronavirus side, the spike dibasic motif and adjacent residues demonstrate complete conservation in β-coronavirus isolates from bats, pangolins, civets, camels, and humans [33-38,46]. However, variations in K-x-H motif as well the absence of a dibasic motif in the spike in animal isolates of coronaviruses suggest that this mechanism of direct interactions between COPI and the spike may not be universal. It has been suggested that dibasic motifs in coronavirus proteins other than the spike may be involved in modulating COPI dependent trafficking by oligomerization with the spike protein [9]. An example is bovine coronavirus, which demonstrates a dibasic motif in the enzyme hemagglutinin esterase but not in the spike protein [47]. The interaction of this enzyme with the spike protein [48] offers a potential route for COPI dependent trafficking of the spike.

our investigation identifies three αCOPI residues Arg^{57}, Asp^{115}, and Tyr^{139}, as essential for spike hepta-peptide binding. These αCOPI residues are completely conserved across organisms associated with β-coronavirus infections such as bats, pangolins, camels, and humans and in chicken, which is infected by γ- and δ-coronaviruses. This suggests a likely conserved mechanism for COPI dependent spike trafficking. Interestingly, the critical αCOPI residues identified in the present analysis are broadly consistent with a prior genetic and biophysical study that implicated αCOPI Arg^{57} and Lys^{15}, and β'COPI Arg^{59} and Asp^{117} (equivalent to αCOPI Asp^{115}) as critical for dibasic motif binding, retrograde trafficking, and growth of yeast cells [27].

Building on this prior investigation, our crystallographic analysis of αCOPI Arg^{57}→Ala and Tyr ^{139}→Ala mutants presents two complementary structural results to substantially advance the understanding of how these residues are critical for COPI architecture. The αCOPI Arg^{57}→Ala

mutant demonstrates a rearrangement of the spike hepta-peptide binding site and of neighboring residues whereas the Tyr[139]→Ala mutant structure is largely similar to the wild type αCOPI structure. Yet, both mutants demonstrate the same functional outcome, i.e., loss of spike hepta-peptide binding. Given this structural sensitivity of αCOPI, and presumably β'COPI, to changes in electrostatics, this raises an interesting question about the structural basis of how mutations in these subunits alter normal retrograde trafficking. It is relevant to note that the *COPA* gene, which encodes the human homolog of αCOPI, has been implicated in a set of clinical disorders collectively known as the COPA syndrome [49,50]. Here, mis-sense mutations including ones that modify side chain charge in the WD40 domains compromise COPA protein function in retrograde trafficking [49]. Based on data presented here, it would be of interest to investigate the structural basis of this dysfunction to gain deeper insights into COPI biology.

In conclusion, our present analysis and supporting prior investigations demonstrate that the extended dibasic motif in the sarbecovirus spike functions as an effective tool to hijack the COPI complex involved in retrograde trafficking. In broader terms, our structural analysis provides a basis to further investigate the structural and functional consequences of αCOPI and β'COPI mutations in disrupting retrograde trafficking.

Methods

Protein and peptide production

The *S. pombe* αCOPI-WD40 domain was synthesized by TOPGENE and cloned in pcDNA3.1(+) with a C-terminal strep-tag for affinity purification. Five mutations (Leu[181]→Lys, Leu[185]→Lys, Ile[192]→Lys, Leu[196]→Lys and Phe[197]→Lys) were incorporated in the gene to improve solubility as suggested previously [28]. Expression was performed in Expi293 mammalian cells using the Thermo Fisher ExpiFectamine expression kit. Protein purification was performed by affinity chromatography of the clarified cellular lysate followed by SEC in a Superdex 75 chromatography column. Arg[57]→Ala, Asp[115]→Ala, and Tyr[139]→Ala mutants of αCOPI-WD40 domain were expressed and purified as described for the wild type protein. The purified αCOPI-WD40 domain in 150 mM NaCl, 5 mM dithiotreitol (DTT), 10% glycerol, and either 20 mM Tris-HCl (pH 7.5) or 50 mM MES-NaOH (pH 6.5) was flash-frozen in liquid nitrogen until further experimentation. All mutations in pcDNA3.1(+)-αCOPI-WD40 were made GenScript.

β'COPI-WD40 (residues 1-301) from *S. cerevisiae* was cloned in pSUMO vector with an N-terminal strep-tag, a Hisx8 tag, and a Ulp1 protease cleavage site, and expressed overnight in

E. coli pLysS cells at 18°C. This fusion protein was purified by affinity chromatography and SEC

in 150 mM NaCl, 5 mM dithiotreitol (DTT), 10% glycerol, and 20 mM Tris-HCl (pH 7.5) followed

by overnight digestion with Ulp1 protease. The digested β'COPI-WD40 domain was subjected to

negative purification by Ni-NTA and SEC and was flash-frozen in liquid nitrogen until further

experimentation.

Peptide synthesis was performed by Biomatik (USA) with an N-terminal biotin tag and a

$(PEG)_4$ linker between the tag and the peptide. No modification was performed at the C-terminus

of the peptides thereby leaving a free terminal carboxylate group.

BLI assay

Biotinylated spike hepta-peptides were tethered to streptavidin (SA) biosensors (FortéBio)

in a 96-well plate format. Purified αCOPI-WD40 domain was provided as the analyte. Kinetics

measurements for determination of binding affinity were performed on an Octet RED96 system

(FortéBio). Data acquisition was carried out using the Data Acquisition 11.1 suite. Briefly, SA

biosensors were hydrated in 200 µL of kinetics buffer (20mM Tris-HCl (pH 7.5) or 50 mM MES-

NaOH (pH 6.5), 150 mM NaCl, 5 mM DTT, 10% glycerol, 0.2 mg/ml bovine serum albumin (BSA),

and 0.002% Tween 20) for 10 minutes prior to binding. The spike hepta-peptide (5 µg/ml) was

loaded on the biosensors for 15 seconds. A baseline was established by rinsing the biosensor

tips in the kinetic buffer for 30 seconds. This was followed by association with αCOP in varying

concentrations over 60 seconds and dissociation in the baseline well for 90 seconds. A

temperature of 25°C and a shake speed of 1000 rpm was maintained during acquisition. All

experiments were carried out in triplicates. A new sensor was used for each replicate. Data

processing and analysis were performed in the FortéBio Data Analysis 11.1 software suite. Raw

data was subtracted from the 0 µM αCOP signal as a reference. The baseline step immediately

before the association step was used for the alignment of the y-axis. An inter-step correction

between the association and dissociation steps was performed. Reference subtracted curves

were processed with the Savitzky-Golay filtering method and subjected to global fitting using a

1:1 binding model. All fits to BLI data had R^2 value (goodness of fit) >0.9.

Crystallization and structure determination

Purified αCOPI-WD40 domain was concentrated to 2 mg/ml in 20 mM Tris-HCl (pH 7.5),

150 mM NaCl, 5 mM DTT and 10% glycerol buffer. Crystal trays were set up with the hanging

drop vapor diffusion method with 0.5 µL of αCOPI-WD40 mixed with an equal volume of reservoir

buffer. Crystals were within 48 hours at 22°C in 20% PEG3350 and 0.25 M sodium citrate tribasic

dihydrate. Crystals were cryo-protected in mother liquor supplemented with 20% ethylene glycol

and flash-frozen in liquid nitrogen. Purified Arg[57]→Ala and Tyr[139]→Ala mutants of αCOP were

concentrated to ~2.2 mg/ml and crystallized as described for the wild type protein. Crystals for

Arg[57]→Ala were obtained in 22% PEG3350 and 0.2 M trisodium citrate and for Tyr[139]→Ala in 18%

PEG3350 and 0.2M potassium-sodium tartrate. The crystals for Arg[57]→Ala and Tyr[139]→Ala

αCOPI-WD40 mutants were cryoprotected in 20% glycerol and 20% ethylene glycol, respectively.

X-ray diffraction data for wild-type αCOPI-WD40 was collected at the beamline GM/CA 23-ID-D

of the Advanced Photon Source at the Argonne National Laboratory and at the National

Synchrotron Light Source II (NSLS II) beamline 17-ID-1 AMX at the Brookhaven National

Laboratory for the mutants. The X-ray diffraction data for the wild type protein crystals was

indexed, integrated and scaled using HKL3000 [51] whereas those for the mutants were processed

using XDS [52] as part of the data acquisition and processing pipeline at the beamline. The data

processing statistics are given in **Supplementary Table T1**. The scaled data were merged using

AIMLESS in CCP4 suite [53]. Molecular replacement was performed in Phenix using a previously

determined αCOPI-WD40 domain structure (PDB ID 4J87) as the search model [54,55]. Iterative

model building and refinement were performed in Phenix.refine [56] and Coot [57]. Figures were

generated in PyMol. Part of the software used here was curated by SBGrid [58].

Analysis of Ramachandran angles

The crystal structures of wild type, Arg[57]→Ala, and Tyr[139]→Ala αCOPI-WD40 were

analyzed in Molprobity [59]. For each structure pair, i.e., wild type with Arg[57]→Ala or wild type with

Tyr[139]→Ala, per residue difference in Ramachandran angles was determined using equation [1]-

$(\delta=\sqrt{((\psi_{WT}-\psi_m)^2+(\phi_{WT}-\phi_m)^2)})$ [1]

Here, Ramachandran angles for wild type and mutant structures are represented as (ψ_{WT}, ϕ_{WT})

and (ψ_m, ϕ_m) respectively. Each structure-pair was superimposed in PyMol and inspected to

ensure consistency with the results of the Ramachandran angle analysis. We identified three

instances of surface exposed residues (Asp[96], Asn[257] in the wild type coordinates, and Ser[11] in

Arg[57]→Ala coordinates) where the structures were highly similar between corresponding main

chain atoms in the wild type and mutant but the sign of a dihedral angle close to 180° had been

flipped. The Cα rmsd of short penta-residue stretches of the polypeptide chain centered at each

of these residues was 0.14Å, 0.06Å, and 0.18Å, respectively. The signs of the Ramachandran

angles for these residues were corrected manually.

***In silico* analysis of sarbecovirus spike hepta-peptide with αCOPI-WD40**

Structural modeling of the SARS-CoV-2 spike C-terminus peptide (sequence: GVKLHYT)

in complex with the αCOPI-WD40 domain was performed using homology modeling in Modeller

[60] and the structure of the αCOP-WD40 complexed with Emp47p peptide (PDB ID 4J8B) as a

template. Prior to computational mutagenesis, models were processed with FastRelax [61] in

Rosetta (v. 3.5), with backbone and side chain atoms constrained to the input coordinates. The

command line parameter settings for FastRelax execution ("relax" executable) used were:

-relax:constrain_relax_to_start_coords

-relax:coord_constrain_sidechains

-relax:ramp_constraints false

-ex1

-ex2

-use_input_sc

-correct

-no_his_his_pairE

-no_optH false

-flip_HNQ

-nstruct 1

Computational mutagenesis simulations to predict effects on binding affinities (ΔΔGs) for

point substitutions were performed using a previously described protocol implemented in Rosetta

(v. 2.3) [62]. Default parameters were used, with the exception of extra rotamers allowed during

packing of modeled side chains, specified by command line parameters:

-extrachi_cutoff 1 -ex1 -ex2 -ex3

Sequence analysis of dibasic motifs in the human membrane proteome

UNIPROT identifiers of secreted and membrane-bound human proteins, as well as
secreted/membrane-bound protein isoforms, were downloaded from the Human Protein Atlas
(http://www.proteinatlas.org) [63]. The corresponding protein sequences were obtained from
UNIPROT, leading to approximately 6800 sequences, which were parsed using an in-house Perl
script to identify C-terminal motif residues. Of these, 119 sequences that demonstrated a C-
terminal dibasic motif were analyzed further for amino acid propensities in the dibasic motif and
neighboring residues.

Accession numbers of protein sequences used in this investigation

Spike protein (bat coronavirus CDPHE15/USA/2006, S5YNL4; bat coronavirus HKU10,
K4JZD2; bat *Rhinolopus* alpha- coronavirus/HuB2013, A0A0U1WJW2; human coronavirus 229E,
P15423; rodent coronavirus, A0A2H4MWY1; mink coronavirus strain WD1127, D9J1Z4; bat
coronavirus HKU8, B1PHK2; bat alpha-coronavirus SAX2011, A0A0U1WHD7; bat alpha-
coronavirus SC2013, A0A0U1UZD0; porcine epidemic diarrhea virus strain CV777, Q91AV1; bat
coronavirus 512/2005, Q0Q466; bat coronavirus HKU2, A8JP08; human coronavirus NL63,
Q6Q1S2; Bat alphacoronavirus, A0A4Y6A7L9; human coronavirus OC43, P36334; beta-
coronavirus HKU24, A0A0A7UZR7; human coronavirus HKU1 isolate N5, Q0ZME7; murine
coronavirus strain A59, P11224; bat coronavirus/Zhejiang2013, A0A088DJY6; hedgehog
coronavirus 1, A0A4D6G1A4; Middle East respiratory syndrome-related coronavirus isolate
United Kingdom/H123990006/2012, K9N5Q8; bat coronavirus HKU5, A3EXD0; bat coronavirus
HKU4, A3EX94; Rousettus bat coronavirus, A0A1B3Q5V4; bat coronavirus HKU9, A3EXG6;
SARS-CoV-2, P0DTC2; SARS-CoV, P59594; avian infectious bronchitis virus strain Beaudette,
P11223; bulbul coronavirus HKU11-934, B6VDW0; common moorhen coronavirus HKU21,
H9BR35; porcine coronavirus HKU15, X2G836; munia coronavirus HKU13-3514, B6VDY7; night
heron coronavirus HKU19, H9BR17; bat coronavirus RaTG13, A0A6B9WHD3; Middle East
respiratory syndrome coronavirus isolate Camel/Qatar_2_2014, KJ650098.1; Civet SARS-CoV
SZ3/2003, AY304486.1; Pangolin coronavirus cDNA31-S, MT799526.1; bat coronavirus
279/2005, Q0Q475; bat coronavirus Rp3/2004, Q3I5J5; bat SARS-like coronavirus WIV1,
AGZ48828.1; porcine transmissible gastroenteritis coronavirus strain Purdue, P07946; bovine
coronavirus strain Mebus, P15777), αCOPI (bat, XP_032949522.1; camel, XP_031291824.1;

chicken, H9L3L2; civet, XP_036778629.1; human, P53621; yeast, NP_595279.1), and β'COPI

(yeast, Q96WV5). The phylogenetic analysis of these sequences was performed in PhyML 3.0 [64].

A list of αCOPI sequences and their multiple sequence alignment from the CONSURF server [45]

is given in **Supplementary Data**.

Statistics and Reproducibility

All BLI experiments were performed in independent triplicates. A statistical correlation

coefficient (CC1/2) was used for crystallographic resolution estimation between half datasets. 5%

of the crystallographic reflections were omitted from refinement to calculate R_{free} and to avoid

over-fitting.

Table 1: Dissociation and rate constants for the interaction of spike hepta-peptides with
COPI-WD40 domains.

Peptide sequence[a]	COPI-WD40 analyte[b]	pH	Equilibrium K_D (µM)	Kinetic K_D (µM)	k_{on} (10^3/Ms)	k_{off} (10^{-1}/s)
GV**KL**H**Y**T	α	7.5	4.17 ± 0.04	2.75 ± 0.09	89.96 ± 2.72	2.50 ± 0.03
GV**KL**H**Y**T	β'	7.5	n.d.	n.d.	n.d.	n.d.
VL**K**TY**HG**	α	7.5	n.d.	n.d.	n.d.	n.d.
GV**KL**<u>**K**</u>YT	α	7.5	4.17 + 0.31	2.71 + 0.05	67.3 + 1.45	1.84 + 0.01
GV**KL**<u>**K**</u>YT	β'	7.5	n.d.	n.d.	n.d.	n.d.
GV**KL**H**Y**T	α	6.5	1.40 ± 0.15	0.95 ± 0.02	152.13 ± 3.45	1.45 ± 0.01
GV**KL**H**Y**T	β'	6.5	n.d.	n.d.	n.d.	n.d.
GV<u>**A**</u>L**HY**T	α	6.5	n.d.	n.d.	n.d.	n.d.
GV**KL**<u>**A**</u>YT	α	6.5	n.d.	n.d.	n.d.	n.d.
GV<u>**A**</u>L<u>**A**</u>YT	α	6.5	n.d.	n.d.	n.d.	n.d.
GV**KL**H<u>**A**</u>T	α	6.5	3.77 ± 0.34	2.32 ± 0.07	120.87± 3.39	2.80 ± 0.03
GV**KL**HY<u>**A**</u>	α	6.5	1.13 ± 0.05	1.14 ± 0.02	66.67 ± 0.80	0.76 ± 0.01
GV**KL**HY<u>**V**</u>	α	6.5	3.20 ± 0.04	2.06 ± 0.05	102.67 ± 2.61	2.12 ± 0.02
GV**KL**HY<u>**E**</u>	α	6.5	0.31 ± 0.01	0.38 ± 0.01	58.96 ± 0.54	0.24 ± 0.003
GV**KL**HY<u>**Q**</u>	α	6.5	0.78 ± 0.04	1.15 ± 0.01	60.05 ± 0.75	0.66 ± 0.01
GV**KL**HY<u>**R**</u>	α	6.5	1.73 ± 0.24	1.29 ± 0.04	57.60 ± 0.61	1.77 ± 0.01
KEVYL**HG**	α	6.5	n.d.	n.d.	n.d.	n.d.
GV**KL**H**Y**T	α-Ala[57]	6.5	n.d.	n.d.	n.d.	n.d.
GV**KL**H**Y**T	α-Ala[115]	6.5	n.d.	n.d.	n.d.	n.d.
GV**KL**H**Y**T	α-Ala[139]	6.5	n.d.	n.d.	n.d.	n.d.

[a]The sequences of immobilized sarbecovirus spike peptides are given in the left column. The
dibasic motif residues are highlighted in bold. Mutations in the peptides relative to the wild-type
sequence are highlighted in bold and are underlined.
[b]α and β' represent wild-type αCOPI-WD40 and β'COPI-WD40, respectively. α-Ala[57], α-Ala[115],
and α-Ala[139], represent the Arg[57]→Ala, Asp[115]→Ala, and Tyr[139]→Ala mutants of αCOPI-WD40,
respectively.
610

611 **Table 2: *In silico* mutagenesis of extended dibasic motif in SARS-CoV-2 spike**

Spike hepta-peptide mutation	Change in interaction energy (ΔΔG, Rosetta units)[1]	Predicted binding change with αCOPI-WD40
Lys^{1269}→Ala	1.9	Weakened
Leu^{1270}→Ala	0.1	Moderate effect
His^{1271}→Ala	1.4	Weakened
Tyr^{1272}→Ala	0.8	Weakened
Thr^{1273}→Ala	0.1	Moderate to no effect
Thr^{1273}→Glu	-1.4	Strengthened

613 [1]Rosetta energy unit ΔΔG values correspond approximately to energies in kcal/mol, as seen in
614 previous studies [62,65], therefore predicted moderate disruptive substitutions have scores > 0.6
615 and highly disruptive substitutions have scores > 1.2, while predicted substantial improved
616 binding corresponds to scores < -1.2.

617   **Table 3: Interaction residues in *in silico* model of spike hepta-peptide with αCOPI-WD40**

Residue in SARS-CoV-2 spike hepta-peptide	αCOPI-WD40 contact residue within 3.5Å
Lys^{1269}	Asp^{96}, Asp^{115}, Tyr^{139}
Leu^{1270}	Arg^{99}
His^{1271}	Arg^{57}, Arg^{99}, Met^{141}, Leu^{157}, Asn^{216}
Tyr^{1272}	His^{31}, Arg^{57}
Thr^{1273}	Arg^{13}, Lys^{15}, His^{31}, Arg^{57}, Arg^{300}, Trp^{302}

619   **Table 4: Effects of *in silico* αCOPI-WD40 mutagenesis on spike hepta-peptide binding**

αCOPI-WD40 mutation	Change in interaction energy (ΔΔG, Rosetta Units)	Predicted binding change with αCOPI-WD40
Lys^{15}→Ala	0.8	Weakened
His^{31}→Ala	0.8	Weakened
Arg^{57}→Ala	1.2	Weakened
Asp^{115}→Ala	0.9	Weakened
Tyr^{139}→Ala	1.1	Weakened

 Figure Legends

 Figure 1: Organization of the coronavirus spike protein. The spike protein is divided into an
ecto-domain (gray), a trans-membrane domain (green), and a cytosolic domain (yellow-white-
cyan). The cytosolic domain includes a cysteine rich tract (yellow) and a dibasic motif for COPI
interactions (cyan). This overall organization and the dibasic motif are conserved in the spike
protein of SARS-CoV, SARS-CoV-2, and MERS-CoV, which have been implicated in wide-spread
human disease. The underlined residues correspond to the peptide sequence utilized in this
manuscript.

Figure 2: Direct binding interaction of sarbecovirus spike hepta-peptide with αCOPI-WD40.
(a) Structural conservation of αCOPI-WD40 domain determined in the present investigation
(yellow) and a previous structure (magenta). Arrow highlights main chain differences between
these two αCOPI-WD40 structures in Gly[168]-Ala[188]. **(b-h)** BLI assay of N-biotinylated spike hepta-
peptide with COPI-WD40 domain. One representative experiment of three is shown in panels **(b-
h)**. Color code for concentrations is given at the bottom of the figure. The equilibrium K_D is
provided with each sensorgram for comparison. **(b)** The spike wild type peptide sequence
demonstrates dose-dependent binding to αCOPI-WD40 domain. **(c)** Scrambling of the hepta-
peptide sequence abolishes binding suggesting sequence-specific interaction. **(d)** β'COPI-Wd40
demonstrates no interaction with the immobilized hepta-peptide. The mutant peptide, Gly-Val-
Lys-Leu-Lys-Tyr-Thr, shows dose-dependent binding to **(e)** αCOPI-WD40 but not **(f)** β'COPI-
WD40. **(g)** Acidification enhances binding between the wild-type spike hepta-peptide and αCOPI-
WD40 domain. **(h)** β'COPI-WD40 shows weakly enhanced binding to the spike hepta-peptide
upon acidification. "n.d." implies not determined for weak interactions.

**Figure 3: Structure-guided mutagenesis of spike hepta-peptide and binding analysis with
αCOPI-WD40 domain. (a)** *In silico* model of the spike hepta-peptide complexed with αCOPI-
WD40 domain (yellow surface). The hepta-peptide is shown as a ribbon in rainbow colors from N
(blue) to C (red) terminus. The Cα-atoms in the hepta-peptide are shown as spheres. The side
chains of residues that interact with αCOPI-WD40 are shown as a stick. **(b-g)** BLI analyses of
αCOPI-WD40 binding to spike hepta-peptide mutants. The color code of BLI traces is given at the
bottom of the figure. One representative experiment of three is shown. Color code for
concentrations is given at the bottom of the figure. The mutation in the spike hepta-peptide
sequence is highlighted in bold and is underlined. The equilibrium K_D is provided with each

sensorgram for comparison. Mutagenesis of, **(b)** Lys[1269], **(c)** His[1271], or **(d)** both abolishes binding

to αCOPI-WD40. **(e)** In contrast, Tyr[1272]→Ala mutation only weakens binding to αCOPI-WD40.

The middle panel shows weak binding of αCOPI-WD40 domain with a hepta-peptide wherein

Lys[1269] has been mutated to Ala. **(f)** Mutagenesis of Thr[1273] to Ala in the spike hepta-peptide leads

to moderately enhanced binding to αCOPI-WD40 whereas mutagenesis to Val[1273] weakens

binding **(g)**.

Figure 4: *In silico* and biophysical analysis of spike C-terminal position in αCOPI-WD40

binding. (a) Sequence logo generated from the alignment of K-x-H(K)-x-x sequence in 119

proteins predicted to be in the human membrane proteome. This shows the abundance of Lys in

the first and third positions, low frequency of aromatic residues in the penultimate position, and

the abundance of Asp and Glu in the C-terminal position. **(b)** An *in silico* model of the spike hepta-

peptide on αCOPI-WD40 (yellow surface) shows an abundance of basic residues in the vicinity

of the terminal Thr[1273] spike residue. Panels **(c-e)** show results of a BLI analysis of binding

between spike hepta-peptide mutants and αCOPI-WD40. The equilibrium K_D is provided with

each sensorgram for comparison. Color code for concentrations is given at the bottom of the

figure. Stabilization of the spike hepta-peptide complexed with αCOPI-WD40 is observed when

the terminal position contains either, **(c)** acidic Glu[1273], or **(d)** neutral Gln[1273] residue. **(e)** In

contrast, basic Arg[1273] in the spike hepta-peptide does not favor enhanced binding. These data

show a role of this terminal hepta-peptide position in modulating tight binding to αCOPI-WD40. In

(c-e), one representative experiment of three is shown.

Figure 5: Phylogenetic analysis of coronavirus spike C-terminal sequence. This phylogram

was generated from the alignment of five C-terminal residues in the spike protein. This penta-

residue spike sequence is shown in italics to the right of each coronavirus. The residues in the

dibasic motif are highlighted in cyan. The C-terminal acidic residues are highlighted in red. An

acidic Asp is seen at the C-terminus of the spike proteins of only those coronaviruses that lack a

dibasic motif.

Figure 6: Structure-guided mutagenesis of αCOPI-WD40 domain and binding analysis with

spike hepta-peptide. Panels **(a-c)** highlight the hepta-peptide interactions (within 4Å) of **(a)** Arg[57],

(b) Asp[115], and **(c)** Tyr[139] residues in αCOPI-WD40 in an *in silico* model. These three interacting

αCOPI-WD40 residues are shown as yellow-red-blue sticks whereas the other residues are

shown as a yellow surface for simplicity. The corresponding interacting residues in the spike

hepta-peptide are labelled and shown as green-red-blue sticks and spheres for Cα atoms. The

BLI analysis of Arg[57]→Ala, Asp[115]→Ala, and Tyr[139]→Ala mutants with the wild type spike hepta-

peptide is shown in panels **(d)**, **(e)**, and **(f)** respectively. All three mutants demonstrate no

substantial binding of the spike hepta-peptide. One representative experiment of three is shown

in panels **(d-f)**.

Acknowledgements: We thank Prof. David Owen (University of Cambridge, Cambridge,

England) for critical comments, Dr. Jonathan Goldberg (Memorial Sloan Kettering Cancer Center,

New York City, USA), Dr. Elena Goldberg (Memorial Sloan Kettering Cancer Center, New York

City, USA), and Dr. Lauren Jackson (Vanderbilt University, Nashville USA) for advice on protein

purification, Dr. Travis Gallagher (National Institute of Standards and Technology, Rockville USA)

for advice on crystallographic data collection, Dr. Andrey Galkin (University of Maryland Baltimore

USA) for advice on BLI assays, and Dr. Jean Jakoncic (NSLSII beamline 17-ID-1 AMX), Dr.

Stephan Corcoran (APS beamline 23-ID-D), and Dr. Darren Sherrell (APS beamline 19-ID) for

advice during crystallographic data collection and processing, and Ms. Corrinne Wilson for

assistance in manuscript preparation. SSH acknowledges support from the University of Maryland

School of Medicine, University of Maryland MPower, and MPower COVID-19 Response Fund

Award. This article was supported by funds through the Maryland Department of Health's

Cigarette Restitution Fund Program, University of Maryland Marlene and Stewart Greenebaum

Comprehensive Cancer Center (National Cancer Institute - Cancer Center Support Grant (CCSG)

- P30CA134274), The Holden Comprehensive Cancer Center at The University of Iowa and its

National Cancer Institute Award P30CA086862. This research used resources of the Advanced

Photon Source, a U.S. Department of Energy (DOE) Office of Science User Facility, operated for

the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-

06CH11357. GM/CA@APS has been funded by the National Cancer Institute (ACB-12002) and

the National Institute of General Medical Sciences (AGM-12006, P30GM138396). Access to

Sector 84 laboratories at the Advanced Protein Characterization Facility (APCF) was possible

through funding provided by NIH grant GM115586 and DOE contract DE-AC02-06CH11357. This

research used resources AMX of the National Synchrotron Light Source II, a U.S. DOE Office of

Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory

under contract no. DE-SC0012704.

Author contributions:

DD: Data acquisition, analysis, interpretation, and manuscript preparation.

SS: Data acquisition, analysis, interpretation, and manuscript preparation.

SK: Data acquisition, analysis, and interpretation.

MM: Data acquisition.

NJS: Data acquisition, analysis, interpretation, and manuscript preparation.

LG: Data acquisition, analysis, interpretation, and manuscript preparation.

BGP: Data acquisition, analysis, interpretation, and manuscript preparation.

SSH: Conception, data acquisition, analysis, interpretation, and manuscript preparation.

Competing interests: The authors declare no competing interests.

Data availability: Coordinates for the crystal structures have been deposited in the Protein Data

Bank with IDs: 7S22 (wild type), 7S16 ($Arg^{57} \rightarrow Ala$), and 7S23 ($Tyr^{139} \rightarrow Ala$). The plasmids for

constructs pSUMO-β′-COPI-WD40, pcDNA3.1(+)-αCOPI-WD40, pcDNA3.1(+)-αCOPI-WD40

mutant $Arg^{57} \rightarrow Ala$, pcDNA3.1(+)-αCOPI-WD40 mutant $Asp^{115} \rightarrow Ala$, and pcDNA3.1(+)-αCOPI-

WD40 mutant $Tyr^{139} \rightarrow Ala$ will be deposited in www.addgene.org.

References:

Stadler, K. *et al.* SARS-beginning to understand a new virus. *Nat Rev Microbiol* **1**, 209-218, doi:10.1038/nrmicro775 (2003).

Ahmed, A. E. The predictors of 3- and 30-day mortality in 660 MERS-CoV patients. *BMC Infect Dis* **17**, 615, doi:10.1186/s12879-017-2712-2 (2017).

Zaki, A. M., van Boheemen, S., Bestebroer, T. M., Osterhaus, A. D. & Fouchier, R. A. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820, doi:10.1056/NEJMoa1211721 (2012).

Polack, F. P. *et al.* Safety and efficacy of the BNT162b2 mRNA COVID-19 vaccine. *N Engl J Med* **383**, 2603-2615, doi:10.1056/NEJMoa2034577 (2020).

USFDA. U.S. Food and Drug Administration. Moderna COVID-19 vaccine [FDA briefing document]. Silver Spring, MD: U.S. Food and Drug Administration, Vaccines and Related Biological Products Advisory Committee; 2020., doi:https://www.fda.gov/media/144434/download (2020).

Beniac, D. R., Andonov, A., Grudeski, E. & Booth, T. F. Architecture of the SARS coronavirus prefusion spike. *Nat Struct Mol Biol* **13**, 751-752, doi:10.1038/nsmb1123 (2006).

Song, H. C. *et al.* Synthesis and characterization of a native, oligomeric form of recombinant severe acute respiratory syndrome coronavirus spike glycoprotein. *J Virol* **78**, 10328-10335, doi:10.1128/JVI.78.19.10328-10335.2004 (2004).

Klumperman, J. *et al.* Coronavirus M proteins accumulate in the Golgi complex beyond the site of virion budding. *J Virol* **68**, 6523-6534, doi:10.1128/JVI.68.10.6523-6534.1994 (1994).

Lontok, E., Corse, E. & Machamer, C. E. Intracellular targeting signals contribute to localization of coronavirus spike proteins near the virus assembly site. *J Virol* **78**, 5913-5922, doi:10.1128/JVI.78.11.5913-5922.2004 (2004).

McBride, C. E., Li, J. & Machamer, C. E. The cytoplasmic tail of the severe acute respiratory syndrome coronavirus spike protein contains a novel endoplasmic reticulum retrieval signal that binds COPI and promotes interaction with membrane protein. *J Virol* **81**, 2418-2428, doi:10.1128/JVI.02146-06 (2007).

Cosson, P. & Letourneur, F. Coatomer interaction with di-lysine endoplasmic reticulum retention motifs. *Science* **263**, 1629-1631, doi:10.1126/science.8128252 (1994).

Gaynor, E. C., te Heesen, S., Graham, T. R., Aebi, M. & Emr, S. D. Signal-mediated retrieval of a membrane protein from the Golgi to the ER in yeast. *J Cell Biol* **127**, 653-665, doi:10.1083/jcb.127.3.653 (1994).

Jackson, M. R., Nilsson, T. & Peterson, P. A. Retrieval of transmembrane proteins to the endoplasmic reticulum. *J Cell Biol* **121**, 317-333, doi:10.1083/jcb.121.2.317 (1993).

Townsley, F. M. & Pelham, H. R. The KKXX signal mediates retrieval of membrane proteins from the Golgi to the ER in yeast. *Eur J Cell Biol* **64**, 211-216 (1994).

Jennings, B. C., Kornfeld, S. & Doray, B. A weak COPI binding motif in the cytoplasmic tail of SARS-CoV-2 spike glycoprotein is necessary for its cleavage, glycosylation, and localization. *FEBS Lett* **595**, 1758-1767, doi:10.1002/1873-3468.14109 (2021).

Letourneur, F. *et al.* Coatomer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell* **79**, 1199-1207, doi:10.1016/0092-8674(94)90011-6 (1994).

Dodonova, S. O. *et al.* A structure of the COPI coat and the role of coat proteins in membrane vesicle assembly. *Science* **349**, 195-198, doi:10.1126/science.aab1121 (2015).

Duden, R., Griffiths, G., Frank, R., Argos, P. & Kreis, T. E. β-COP, a 110 kd protein associated with non-clathrin-coated vesicles and the Golgi complex, shows homology to β-adaptin. *Cell* **64**, 649-665, doi:10.1016/0092-8674(91)90248-w (1991).

19 Hara-Kuge, S. *et al.* En bloc incorporation of coatomer subunits during the assembly of
COP-coated vesicles. *J Cell Biol* **124**, 883-892, doi:10.1083/jcb.124.6.883 (1994).
20 Harrison-Lavoie, K. J. *et al.* A 102 kDa subunit of a Golgi-associated particle has homology
to beta subunits of trimeric G proteins. *EMBO J* **12**, 2847-2853 (1993).
21 Malhotra, V., Serafini, T., Orci, L., Shepherd, J. C. & Rothman, J. E. Purification of a novel
class of coated vesicles mediating biosynthetic protein transport through the Golgi stack.
Cell **58**, 329-336, doi:10.1016/0092-8674(89)90847-7 (1989).
22 Serafini, T. *et al.* A coat subunit of Golgi-derived non-clathrin-coated vesicles with
homology to the clathrin-coated vesicle coat protein β-adaptin. *Nature* **349**, 215-220,
doi:10.1038/349215a0 (1991).
23 Waters, M. G., Serafini, T. & Rothman, J. E. 'Coatomer': A cytosolic protein complex
containing subunits of non-clathrin-coated Golgi transport vesicles. *Nature* **349**, 248-251,
doi:10.1038/349248a0 (1991).
24 Stenbeck, G. *et al.* beta'-COP, a novel subunit of coatomer. *EMBO J* **12**, 2841-2845
(1993).
25 Eugster, A., Frigerio, G., Dale, M. & Duden, R. The α- and β′-COP WD40 domains mediate
cargo-selective interactions with distinct di-lysine motifs. *Mol Biol Cell* **15**, 1011-1023,
doi:10.1091/mbc.e03-10-0724 (2004).
26 Fiedler, K., Veit, M., Stamnes, M. A. & Rothman, J. E. Bimodal interaction of coatomer
with the p24 family of putative cargo receptors. *Science* **273**, 1396-1399,
doi:10.1126/science.273.5280.1396 (1996).
27 Jackson, L. P. *et al.* Molecular basis for recognition of dilysine trafficking motifs by COPI.
Dev Cell **23**, 1255-1262, doi:10.1016/j.devcel.2012.10.017 (2012).
28 Ma, W. & Goldberg, J. Rules for the recognition of dilysine retrieval motifs by coatomer.
EMBO J **32**, 926-937, doi:10.1038/emboj.2013.41 (2013).
29 Schroder-Kohne, S., Letourneur, F. & Riezman, H. α-COP can discriminate between
distinct, functional di-lysine signals in vitro and regulates access into retrograde transport.
J Cell Sci **111 (Pt 23)**, 3459-3470 (1998).
30 Cattin-Ortolá, J. *et al.* Sequences in the cytoplasmic tail of SARS-CoV-2 spike facilitate
expression at the cell surface and syncytia formation. *Nature Communications* **12**, 5333
(2021).
31 Woo, P. C. *et al.* Discovery of seven novel mammalian and avian coronaviruses in the
genus deltacoronavirus supports bat coronaviruses as the gene source of
alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of
gammacoronavirus and deltacoronavirus. *J Virol* **86**, 3995-4008, doi:10.1128/JVI.06540-
11 (2012).
32 Drexler, J. F., Corman, V. M. & Drosten, C. Ecology, evolution and classification of bat
coronaviruses in the aftermath of SARS. *Antiviral Res* **101**, 45-56,
doi:10.1016/j.antiviral.2013.10.013 (2014).
33 Li, W. *et al.* Bats are natural reservoirs of SARS-like coronaviruses. *Science* **310**, 676-
679, doi:10.1126/science.1118391 (2005).
34 Ge, X. Y. *et al.* Isolation and characterization of a bat SARS-like coronavirus that uses the
ACE2 receptor. *Nature* **503**, 535-538, doi:10.1038/nature12711 (2013).
35 Guan, Y. *et al.* Isolation and characterization of viruses related to the SARS coronavirus
from animals in southern China. *Science* **302**, 276-278, doi:10.1126/science.1087139
(2003).
36 Lau, S. K. *et al.* Severe acute respiratory syndrome coronavirus-like virus in Chinese
horseshoe bats. *Proc Natl Acad Sci U S A* **102**, 14040-14045,
doi:10.1073/pnas.0506735102 (2005).
37 Tang, X. C. *et al.* Prevalence and genetic diversity of coronaviruses in bats from China. *J
Virol* **80**, 7481-7490, doi:10.1128/JVI.00697-06 (2006).

38 Xiao, K. *et al.* Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins.
Nature **583**, 286-289, doi:10.1038/s41586-020-2313-x (2020).
39 WHO-2021. https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/, accessed
on 12/03/2021.
40 Bolles, M., Donaldson, E. & Baric, R. SARS-CoV and emergent coronaviruses: Viral
determinants of interspecies transmission. *Curr Opin Virol* **1**, 624-634,
doi:10.1016/j.coviro.2011.10.012 (2011).
41 Han, H. J., Yu, H. & Yu, X. J. Evidence for zoonotic origins of Middle East respiratory
syndrome coronavirus. *J Gen Virol* **97**, 274-280, doi:10.1099/jgv.0.000342 (2016).
42 Latinne, A. *et al.* Origin and cross-species transmission of bat coronaviruses in China. *Nat
Commun* **11**, 4235, doi:10.1038/s41467-020-17687-3 (2020).
43 Roess, A., Carruth, L., Lahm, S. & Salman, M. Camels, MERS-CoV, and other emerging
infections in east Africa. *Lancet Infect Dis* **16**, 14-15, doi:10.1016/S1473-3099(15)00471-
5 (2016).
44 Zhang, T., Wu, Q. & Zhang, Z. Probable pangolin origin of SARS-CoV-2 associated with
the COVID-19 outbreak. *Curr Biol* **30**, 1346-1351 e1342, doi:10.1016/j.cub.2020.03.022
(2020).
45 Ashkenazy, H. *et al.* ConSurf 2016: An improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acids Res* **44**, W344-350,
doi:10.1093/nar/gkw408 (2016).
46 Raj, V. S. *et al.* Isolation of MERS coronavirus from a dromedary camel, Qatar, 2014.
Emerg Infect Dis **20**, 1339-1342, doi:10.3201/eid2008.140663 (2014).
47 Kienzle, T. E., Abraham, S., Hogue, B. G. & Brian, D. A. Structure and orientation of
expressed bovine coronavirus hemagglutinin-esterase protein. *J Virol* **64**, 1834-1838,
doi:10.1128/JVI.64.4.1834-1838.1990 (1990).
48 Nguyen, V. P. & Hogue, B. G. Protein interactions during coronavirus assembly. *J Virol*
71, 9278-9284, doi:10.1128/JVI.71.12.9278-9284.1997 (1997).
49 Watkin, L. B. *et al.* COPA mutations impair ER-Golgi transport and cause hereditary
autoimmune-mediated lung disease and arthritis. *Nat Genet* **47**, 654-660,
doi:10.1038/ng.3279 (2015).
50 Vece, T. J. *et al.* COPA syndrome: A novel autosomal dominant immune dysregulatory
disease. *J Clin Immunol* **36**, 377-387, doi:10.1007/s10875-016-0271-8 (2016).
51 Minor, W., Cymborowski, M., Otwinowski, Z. & Chruszcz, M. HKL-3000: The integration
of data reduction and structure solution-from diffraction images to an initial model in
minutes. *Acta Crystallogr D Biol Crystallogr* **62**, 859-866,
doi:10.1107/S0907444906019949 (2006).
52 Kabsch, W. XDS. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132,
doi:10.1107/S0907444909047337 (2010).
53 Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution? *Acta
Crystallogr D Biol Crystallogr* **69**, 1204-1214, doi:10.1107/S0907444913000061 (2013).
54 McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674,
doi:10.1107/S0021889807021206 (2007).
55 Rossmann, M. G. & Blow, D. M. The detection of sub-units within the crystallographic
asymmetric unit. *Acta Crystallographica* **15**, 24-31, doi:doi:10.1107/S0365110X62000067
(1962).
56 Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with
phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352-367,
doi:10.1107/S0907444912001308 (2012).
57 Emsley, P. & Cowtan, K. Coot: Model-building tools for molecular graphics. *Acta
Crystallogr D Biol Crystallogr* **60**, 2126-2132, doi:10.1107/S0907444904019158 (2004).

58   Morin, A. *et al.* Collaboration gets the most out of software. *Elife* **2**, e01456,
doi:10.7554/eLife.01456 (2013).
59   Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315, doi:10.1002/pro.3330 (2018).
60   Webb, B. & Sali, A. Protein structure modeling with MODELLER. *Methods Mol Biol* **1137**,
1-15, doi:10.1007/978-1-4939-0366-5_1 (2014).
61   Conway, P., Tyka, M. D., DiMaio, F., Konerding, D. E. & Baker, D. Relaxation of backbone
bond geometry improves protein energy landscape modeling. *Protein Sci* **23**, 47-55,
doi:10.1002/pro.2389 (2014).
62   Kortemme, T. & Baker, D. A simple physical model for binding energy hot spots in protein-
protein complexes. *Proc Natl Acad Sci U S A* **99**, 14116-14121,
doi:10.1073/pnas.202485799 (2002).
63   Thul, P. J. *et al.* A subcellular map of the human proteome. *Science* **356**,
doi:10.1126/science.aal3321 (2017).
64   Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321,
doi:10.1093/sysbio/syq010 (2010).
65   Pierce, B. G. *et al.* Computational design of the affinity and specificity of a therapeutic T
cell receptor. *PLoS Comput Biol* **10**, e1003478, doi:10.1371/journal.pcbi.1003478 (2014).
907