Inverse Problem Reveals Conditions for Characteristic Retinal Degeneration Patterns in Retinitis Pigmentosa under the Trophic Factor Hypothesis

Paul A. Roberts [1,*]

[1] *Baden Lab, Centre for Sensory Neuroscience and Computation, School of Life Sciences, University of Sussex, Brighton, UK*

Correspondence*:
Paul A. Roberts
p.a.roberts@univ.oxon.org

ABSTRACT

Retinitis pigmentosa (RP) is the most common inherited retinal dystrophy with a prevalence of about 1 in 4000, affecting approximately 1.5 million people worldwide. Patients with RP experience progressive visual field loss as the retina degenerates, destroying light-sensitive photoreceptor cells (rods and cones), with rods affected earlier and more severely than cones. Spatio-temporal patterns of retinal degeneration in human RP have been well characterised; however, the mechanism(s) giving rise to these patterns have not been conclusively determined. One such mechanism, which has received a wealth of experimental support, is described by the trophic factor hypothesis. This hypothesis suggests that rods produce a trophic factor necessary for cone survival; the loss of rods depletes this factor, leading to cone degeneration. In this paper we formulate a partial differential equation mathematical model of RP in one spatial dimension, spanning the region between the retinal centre (fovea) and the retinal edge (ora serrata). Using this model we derive and solve an inverse problem, revealing for the first time experimentally testable conditions under which the trophic factor mechanism will qualitatively recapitulate the spatio-temporal patterns or retinal regeneration observed in human RP.

Keywords: Partial Differential Equations, Asymptotic Analysis, Retina, Photoreceptors, Rod-derived Cone Viability Factor

1 INTRODUCTION

The group of inherited retinal diseases known as retinitis pigmentosa (RP) causes the progressive loss of visual function. The patterns of visual field loss associated with the human version of this condition have been well characterised; however, the mechanisms underpinning these patterns have yet to be conclusively determined. In this paper, we use mathematical models to predict the conditions under which a trophic factor mechanism could explain these patterns.

The retina is a tissue layer lining the back of the eye containing light-sensitive cells known as photo-receptors, which come in two varieties: rods and cones (Fig. 1A). Rods confer monochromatic vision under low-light (scotopic) conditions, while cones confer colour vision under well-lit (photopic) conditions (Oyster, 1999). In RP, rod function and health are typically affected earlier and more severely than those of cones, with cone loss following rod loss. Rods are lost since they express one of a number of mutant

genes; however, since cones do not express these genes, it is hypothesised that they depend upon rods
either directly or indirectly for their survival (Daiger et al., 2007; Hamel, 2006; Hartong et al., 2006).

A number of mechanisms have been hypothesised to explain secondary cone loss, including trophic factor
(TF) depletion (Aït-Ali et al., 2015; Léveillard et al., 2004; Mei et al., 2016), oxygen toxicity (Stone et al.,
1999; Travis et al., 1991; Valter et al., 1998), metabolic dysregulation (Punzo et al., 2009, 2012), toxic
substances (Ripps, 2002) and microglia (Gupta et al., 2003). While not typically related to spatio-temporal
patterns of retinal degeneration in the literature, it is reasonable to infer that these mechanisms play an
important role in determining spatio-temporal patterns of retinal degeneration.

Grover et al. (1998) have classified the spatio-temporal patterns of visual field loss (and hence photorece-
ptor degeneration) in RP patients into three patterns and six sub-patterns (see Fig. 2). Pattern 1A consists in
a restriction of the peripheral visual field, while Pattern 1B also includes a para-/peri-foveal ring scotoma
(blind spot); Pattern 2 (A, B and C) involves an initial loss of the superior visual field, winding nasally or
temporally into the inferior visual field; lastly, Pattern 3 starts with loss of the mid-peripheral visual field,
before spreading into the superior of inferior visual field and winding around the far-periphery. In all cases
central vision is the best preserved, though it too is eventually lost.

In this paper we explore the conditions under which the TF mechanism (in isolation) can replicate
the patterns of retinal degeneration observed *in vivo*. This mechanism has been studied in detail. Rod
photoreceptors have been shown to produce a TF called rod-derived cone viability factor (RdCVF), which
is necessary for cone survival (Fintz et al., 2003; Léveillard et al., 2004; Mohand-Saïd et al., 1998, 2000,
1997; Yang et al., 2009). Cones do not produce RdCVF, thus, when rods are lost, RdCVF concentration
drops and cone degeneration follows (though it has been suggested that it may ultimately be oxygen toxicity
which kills cones; Léveillard and Sahel, 2017).

Thus far, two groups have developed mathematical models operating under the TF hypothesis. Camacho
et al. have developed a series of (non-spatial) dynamical systems ordinary differential equation models to
describe the role of RdCVF in health and RP (Colón Vélez et al., 2003; Camacho et al., 2010; Camacho and
Wirkus, 2013; Camacho et al., 2014, 2016a,b,c, 2019, 2020, 2021; Wifvat et al., 2021). In Roberts (2021),
we developed the first partial differential equation (PDE) models of the TF mechanism in RP, predicting
the spatial spread of retinal degeneration (see also Roberts et al., 2017, 2018, for our work on the oxygen
toxicity mechanism, together with Roberts et al., 2016b; Burns et al., 2002, for a mathematical model of the
toxic substance mechanism; and Roberts et al., 2016a, for a review on mathematical models of the retina).
It was found that, under the assumption of a spatially uniform mutation-induced rod degeneration rate, ϕ_r,
and TF threshold concentration, f_{crit}, (defined in Section 2.1) the mechanism was unable to replicate *in*
vivo patterns of retinal degeneration.

In this study, we extend our work in Roberts (2021) by formulating and solving an inverse problem
to determine the spatially heterogeneous $\phi_r(\theta)$ and $f_{\mathrm{crit}}(\theta)$ profiles that are required to qualitatively
recapitulate observed patterns of spatio-temporal degeneration in human RP.

2 MATERIAL AND METHODS

2.1 Model Formulation

We begin by formulating a reaction-diffusion PDE mathematical model (a simplified version of the model
presented in Roberts, 2021), posing it on a spherical geometry to replicate that of the human retina. This
geometry is most naturally represented using a spherical polar coordinate system, (r,θ,ϕ), centred in the
middle of the vitreous body, where $r \geq 0$ (m) is the distance from the origin, $0 \leq \theta \leq \pi$ (rad) is the polar

angle and $0 \leq \phi < 2\pi$ (rad) is the azimuthal angle. To create a more mathematically tractable model, we
simplify the geometry by assuming symmetry about the z-axis (directed outward from the origin through
the foveal centre), eliminating variation in the azimuthal direction, and effectively depth-average through
the retina, assuming that it lies at a single fixed distance, $R > 0$ (m), from the origin at all eccentricities, θ
(see Roberts, 2021, for justification). Thus, we have reduced the coordinate system to (R,θ), where R is a
positive constant parameter and $0 \leq \theta \leq \Theta$ is an independent variable, which we bound to range between
the fovea (at $\theta = 0$ rad) and the ora serrata (at $\theta = \Theta = 1.33$ rad; see Fig. 1A). We further simplify the
model by non-dimensionalising; scaling the dependent and independent variables so that they and the
resultant model parameters are dimensionless and hence unitless. This reduces the number of parameters
(including eliminating R) and allows us to identify the dominant terms of the governing equations in the
ensuing asymptotic analysis. For this reason, there are no units to be stated in Figs. 3–10. For the full
dimensional model and non-dimensionalisation see Roberts (2021).

We proceed directly to the dimensionless model, which consists of a system of PDEs in terms of the
dependent variables: TF concentration, $f(\theta, t)$, rod photoreceptor density, $p_r(\theta, t)$, and cone photoreceptor
density, $p_c(\theta, t)$; as functions of the independent variables: polar angle, scaled to lie in the range $0 \leq \theta \leq 1$,
and time, $t > 0$ (see Table 1).

The TF equation is as follows

$$\frac{\partial f}{\partial t} = \underbrace{\frac{D_f}{\sin(\Theta\theta)}\frac{\partial}{\partial \theta}\left(\sin(\Theta\theta)\frac{\partial f}{\partial \theta}\right)}_{\text{diffusion}} + \underbrace{\alpha p_r}_{\text{production}} - \underbrace{\beta f p_c}_{\text{consumption}} - \underbrace{\eta f}_{\text{decay}}, \tag{1}$$

86   where the parameters, D_f, the TF diffusivity, α, the rate of TF production by rods, β, the rate of TF
87   consumption by cones, and η, the rate of TF decay, are positive constants. Trophic factor is free to diffuse
88   across the retina, is produced by rods at a rate proportional to their local density, is consumed by cones at a
89   rate proportional the product of the local TF concentration and the local cone density, and decays at a rate
90   proportional to the local TF concentration. The rod equation takes the following form

$$\frac{\partial p_r}{\partial t} = - \underbrace{\phi_r(\theta)p_r}_{\substack{\text{cell degeneration} \\ \text{(mutation-induced)}}}, \tag{2}$$

where the variable $\phi_r(\theta)$, the rate of mutation-induced rod degeneration, may vary spatially (functional
forms defined in the Results section), or take a constant value, ϕ_r. Rods degenerate due to their expression
of a mutant gene at a rate proportional to their local density and independent of the TF concentration. We
note that Eqn. (2) can be solved to yield $p_r(\theta, t) = p_{r_{\text{init}}}(\theta)e^{-\phi_r(\theta)t}$ (where $p_{r_{\text{init}}}(\theta)$, the initial value of
$p_r(\theta, t)$, is defined below), provided there is no delay in onset or interruption of degeneration. The cone
equation is as follows

$$\frac{\partial p_c}{\partial t} = - \underbrace{p_c\lambda_2(f)}_{\substack{\text{cell degeneration} \\ \text{(TF starvation)}}}. \tag{3}$$

Defining the Heaviside step function, $H(\cdot)$, such that

$$H(x) := \left\{ \begin{array}{ll} 0 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0, \end{array} \right.$$

the function $\lambda_2(f)$ is given by

$$\lambda_2(f) = 1 - H(f - f_{\text{crit}}(\theta)),$$

where the variable $f_{\text{crit}}(\theta)$, the TF threshold concentration, may vary spatially (functional forms defined in
the Results section), or take a constant value, f_{crit}. Cone density remains constant provided the local TF
concentration, $f(\theta, t)$, remains in the healthy range at or above the critical threshold, f_{crit}, while cones
decay (due to TF starvation) at a rate proportional to their local density if $f(\theta, t)$ drops below this threshold.

Having defined the governing equations (Eqs. (1)–(3)), we close the system by imposing boundary and
initial conditions. We apply zero-flux boundary conditions at both ends of the domain,

$$\frac{\partial f}{\partial \theta}(0, t) = 0 = \frac{\partial f}{\partial \theta}(1, t), \tag{4}$$

such that there is no net flow of TF into or out of the domain. This is justified by symmetry at $\theta = 0$, and
is reasonably assumed to hold at $\theta = 1$ in the absence of experimental data. The healthy rod and cone
distributions are given by the following functions

$$\tilde{p}_r(\theta) = B_3 \theta e^{-b_3 \theta},$$
$$\tilde{p}_c(\theta) = B_1 e^{-b_1 \theta} + B_2 e^{-b_2 \theta},$$

103 where the values of the positive constants B_1, B_2, B_3, b_1, b_2 and b_3 are found by fitting to the mean
104 of Curcio et al. (1990)'s measurements of healthy human rod and cone distributions along the temporal
105 horizontal meridian using the Trust-Region Reflective algorithm in Matlab's curve fitting toolbox (see Fig.
106 1B). Lastly, we impose the initial conditions

$$f(\theta, 0) = f_{\text{init}}(\theta), \qquad p_r(\theta, 0) = p_{r_{\text{init}}}(\theta) = \tilde{p}_r(\theta), \qquad p_c(\theta, 0) = p_{c_{\text{init}}}(\theta) = \tilde{p}_c(\theta), \tag{5}$$

where $f_{\text{init}}(\theta)$ is the steady-state solution to Eqs. (1) and (4) with $p_r = p_{r_{\text{init}}}(\theta)$ and $p_c = p_{c_{\text{init}}}(\theta)$ (see
Fig. 3A). Thus, the retina starts in the healthy state in all simulations. See Table 2 for the dimensionless
parameter values (see Roberts, 2021, for dimensional values and justification of parameter values). The
model presented here simplifies that in Roberts (2021) in the following ways: it does not include treatment,
cone OS regeneration, or initial patches of rod or cone loss, while mutation-induced rod loss is active for
all simulations in this study. The present model also adds two new features to the previous model: allowing
the rate of mutation-induced rod degeneration, $\phi_r(\theta)$, and the TF threshold concentration, $f_{\text{crit}}(\theta)$, to vary
spatially, where before they were constant (or piecewise constant in the high f_{crit} subcase).

115
2.2   Numerical Solutions

Numerical (computational) solutions to Eqs. (1)–(5) were obtained using the method of lines (as in
Roberts, 2021), discretising in space and then integrating in time. The time integration was performed
using the Matlab routine `ode15s`, a variable-step, variable-order solver, designed to solve problems
involving multiple timescales such as this (Matlab version R2020a was used here and throughout the paper).
We used a relative error tolerance of 10^{-6} and an absolute error tolerance of 10^{-10}, with the remaining
settings at their default values. The number of spatial mesh points employed varies between simulations,
taking values of 26, 51, 101, 401 or 4001. The upper bound of 4001 mesh points was chosen such that
the distance between mesh points corresponds to the average width of a photoreceptor. In each case the
maximum computationally feasible mesh density was employed, all mesh densities being sufficient to

achieve accurate results. The initial TF profile, $f(\theta, 0) = f_{\text{init}}(\theta)$, was calculated by discretising Eqs. (1)
and (4) at steady-state, using a finite difference scheme, and solving the consequent system of nonlinear
algebraic equations using the Matlab routine fsolve (which employs a Trust–Region–Dogleg algorithm)
with $p_r = p_{r_{\text{init}}}(\theta)$ and $p_c = p_{c_{\text{init}}}(\theta)$.

130
2.3 Inverse Problem

In an inverse problem we seek to determine the model input required to attain a known/desired output.
In this case the known output is the target cone degeneration profile, $t_{\text{degen}}(\theta)$, while the input is either
the rate of mutation-induced rod loss profile, $\phi_r(\theta)$, or the TF threshold concentration profile, $f_{\text{crit}}(\theta)$,
with corresponding inverses denoted as $\phi_r(\theta) = \phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}}(\theta) = f_{\text{crit}_{\text{inv}}}(\theta)$ respectively. When
searching for $\phi_{r_{\text{inv}}}(\theta)$, we hold the TF threshold concentration constant at $f_{\text{crit}}(\theta) = f_{\text{crit}} = 3 \times 10^{-5}$,
while, when searching for $f_{\text{crit}_{\text{inv}}}(\theta)$, we hold the rate of mutation-induced rod loss constant at $\phi_r(\theta) =$
$\phi_r = 7.33 \times 10^{-2}$. We consider a range of target cone degeneration profiles, summarised in Table 3 and
Fig. 5, which qualitatively replicate retinal degeneration Patterns 1A, 1B and 3 seen *in vivo* (taking the
degeneration of the far-peripheral retina to occur in a radially symmetric manner in Pattern 3; see Fig. 2
and Grover et al., 1998). We do not consider patterns of type 2 (to be explored in a future study) since
these cannot be replicated by a 1D model. For each degeneration pattern we consider a set of sub-patterns
to examine how this affects the shape of the inverses. We also consider a uniform target cone degeneration
profile for comparison. For each pattern, we consider the effect of two (biologically realistic) scalings for
the rate of TF production by rods, α, and the rate of TF consumption by cones, β, upon the inverse profiles:
(i) Scaling 1 — for which $\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$ as in Roberts (2021); and (ii) Scaling 2 —
for which $\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$.

We solve the inverse problem both analytically and numerically (computationally), as described in
Sections 2.3.1 and 2.3.2 below. Analytical approximations are computationally inexpensive and provide
deeper insight into model behaviour, while numerical solutions, though computationally intensive, are
more accurate.

152
2.3.1 Analytical Inverse

To derive analytical (algebraic) approximations for the inverses, $\phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}_{\text{inv}}}(\theta)$, we perform an
asymptotic analysis, seeking the leading order behaviour of Eqs. (1)–(5). Proceeding as in Roberts (2021)
(where we considered a steady-state problem), we choose $\epsilon = O(10^{-2})$ and scale the parameters $\eta = \epsilon^{-1}\eta'$
and $b_1 = \epsilon^{-1}b_1'$, introducing the new scaling $\phi_r(\theta) = \epsilon\phi_r'(\theta)$, as we study the time-dependent problem here.
We consider two possible (biologically realistic) scalings on α and β: (i) Scaling 1 — for which $\alpha = \epsilon^{-2}\alpha'$
and $\beta = \epsilon^{-3}\beta'$ as in Roberts (2021) (corresponding to $\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$); and (ii)
Scaling 2 — for which $\alpha = \epsilon^{-1}\alpha'$ and $\beta = \epsilon^{-2}\beta'$ (corresponding to $\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$).
All remaining parameters are assumed to be $O(1)$. We also scale the dependent variable $p_c(\theta) = \epsilon p_c'(\theta)$,
and assume $f(\theta) = O(1)$ and $p_r(\theta) = O(1)$.

Applying the above scalings and dropping the dashes, Eqn. (2) becomes

$$\frac{\partial p_r}{\partial t} = -\epsilon\phi_r(\theta)p_r.$$

Thus, on this (fast) timescale, the rod density is constant. Since we are interested in the timescale upon
which rods degenerate, we scale time as $t' = \epsilon t$ such that the decay term enters the dominant balance. Thus,

on this slow timescale, after dropping the dashes, we have that

$$\frac{\partial p_r}{\partial t} = -\phi_r(\theta)p_r,$$

163 such that, at leading order, $p_{r_0}(\theta, t) = p_{r_{\text{init}_0}}(\theta)e^{-\phi_r(\theta)t} = B_3\theta e^{-b_3\theta}e^{\phi_r(\theta)t}$.

164   We are interested here in the regime in which cones have not yet degenerated, thus we assume the leading
165 order cone density remains constant at $p_{c_0}(\theta) = p_{c_{\text{init}_0}}(\theta) = B_2 e^{-b_2\theta}$.

Applying Scaling 1 and the slow timescale to Eqn. (1) we obtain

$$\epsilon\frac{\partial f}{\partial t} = D_f\frac{\partial^2 f}{\partial \theta^2} + D_f\Theta\cot(\Theta\theta)\frac{\partial f}{\partial \theta} + \epsilon^{-2}\alpha p_r - \epsilon^{-2}\beta p_c f - \epsilon^{-1}\eta f.$$

Since the TF dynamics occur on a faster timescale than mutation-induced rod loss, we make a quasi-steady-state approximation (QSSA), assuming that the TF concentration instantaneously takes its steady-state profile, for any given rod density profile, as the rods degenerate ($\epsilon\partial_t f \sim 0$). Thus, at leading order, we obtain

$$f_{0_{\text{QSSA}}}(\theta) = \frac{\alpha p_{r_0}(\theta, t)}{\beta p_{c_0}(\theta)}.$$

166 Rearranging this expression and assuming that cone degeneration initiates when $f_{0_{\text{QSSA}}}(\theta) = f_{\text{crit}}(\theta)$, we
167 obtain the cone degeneration time profile,

$$t_{\text{degen}}(\theta) = \frac{1}{\phi_r(\theta)}\left(\log\left(\frac{\alpha B_3}{\beta B_2 f_{\text{crit}}(\theta)}\theta\right) - (b_3 - b_2)\theta\right), \tag{6}$$

168 the inverse mutation-induced rod degeneration rate profile,

$$\phi_{r_{\text{inv}}}(\theta) = \frac{1}{t_{\text{degen}}(\theta)}\left(\log\left(\frac{\alpha B_3}{\beta B_2 f_{\text{crit}}}\theta\right) - (b_3 - b_2)\theta\right), \tag{7}$$

169 and the inverse TF threshold concentration profile,

$$f_{\text{crit}_{\text{inv}}}(\theta) = \frac{\alpha B_3}{\beta B_2}\theta e^{-((b_3-b_2)\theta+\phi_r t_{\text{degen}}(\theta))}. \tag{8}$$

Alternatively, if we apply Scaling 2 and the slow timescale to Eqn. (1) we obtain

$$\epsilon\frac{\partial f}{\partial t} = D_f\frac{\partial^2 f}{\partial \theta^2} + D_f\Theta\cot(\Theta\theta)\frac{\partial f}{\partial \theta} + \epsilon^{-1}\alpha p_r - \epsilon^{-1}\beta p_c f - \epsilon^{-1}\eta f.$$

Applying the QSSA and proceeding as above we find

$$f_{0_{\text{QSSA}}}(\theta) = \frac{\alpha p_{r_0}(\theta, t)}{\beta p_{c_0}(\theta) + \eta},$$

170 with cone degeneration time profile,

$$t_{\text{degen}}(\theta) = \frac{1}{\phi_r(\theta)}\left(\log\left(\frac{\alpha B_3}{(\beta B_2 + \eta e^{b_2\theta})f_{\text{crit}}(\theta)}\theta\right) - (b_3 - b_2)\theta\right), \tag{9}$$

171 inverse mutation-induced rod degeneration rate profile,

$$\phi_{r_{\mathrm{inv}}}(\theta) = \frac{1}{t_{\mathrm{degen}}(\theta)} \left(\log \left(\frac{\alpha B_3}{(\beta B_2 + \eta e^{b_2 \theta}) f_{\mathrm{crit}}} \theta \right) - (b_3 - b_2)\theta \right), \tag{10}$$

172 and inverse TF threshold concentration profile,

$$f_{\mathrm{crit}_{\mathrm{inv}}}(\theta) = \frac{\alpha B_3}{(\beta B_2 + \eta e^{b_2 \theta})} \theta e^{-((b_3 - b_2)\theta + \phi_r t_{\mathrm{degen}}(\theta))}. \tag{11}$$

173
174
2.3.2 Numerical Inverse

The numerical inverse is calculated by repeatedly solving the forward problem (Eqs. (1)–(5)) for different
values of the input ($\phi_r(\theta)$ or $f_{\mathrm{crit}}(\theta)$), with the aim of converging upon the inverse ($\phi_{r_{\mathrm{inv}}}(\theta)$ or $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$).
To find $\phi_{r_{\mathrm{inv}}}(\theta)$ we use the Matlab routine fminsearch (which uses a simplex search method), while to
obtain $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ the Matlab routine patternsearch (which uses an adaptive mesh technique) was found to
be more effective. In both cases the objective function was taken as the sum of squares of the difference
between the target cone degeneration profile, $t_{\mathrm{degen}}(\theta)$, and the contour described by $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$
(along which cone degeneration is deemed to have initiated). Eqs. (1)–(5) were solved at each iteration as
described in Section 2.2. Numerical inverses were calculated only at those locations (eccentricities) where
the analytical inverse failed to generate a $t_{\mathrm{degen}}(\theta)$ profile matching the target profile, the analytical inverse
being assumed to hold at all other eccentricities.

3 RESULTS

We begin by calculating the cone degeneration profiles, $t_{\mathrm{degen}}(\theta)$, in the case where both the rate of
mutation induced rod degeneration, ϕ_r, and the TF threshold concentration, f_{crit}, are spatially uniform
(or piecewise constant). We set the standard value for $\phi_r = 7.33 \times 10^{-2}$ and consider the subcases (i)
$f_{\mathrm{crit}} = 3 \times 10^{-5}$ for $0 \leq \theta \leq 1$ (Fig. 4A), and (ii) $f_{\mathrm{crit}} = 0.3$ for $\theta > 0.13$ while $f_{\mathrm{crit}} = 3 \times 10^{-5}$ for
$\theta \leq 0.13$ (Fig. 4B), as were explored in Roberts (2021). These subcases correspond to the situation in
which the TF threshold concentration lies beneath the minimum healthy TF value at all retinal locations
(i), and the situation in which foveal cones are afforded special protection compared to the rest of the
retina, such that they can withstand lower TF concentrations (ii). For notational simplicity, we shall refer
to subcase (ii) simply as $f_{\mathrm{crit}} = 0.3$ in what follows. As with Figs. 6–9, we consider both Scaling 1 and
Scaling 2 (see Section 2.3) on the rate of TF production by rods, α, and the rate of TF consumption by
cones, β, calculating both analytical and numerical solutions.

Cone degeneration initiates at the fovea ($\theta = 0$) in Fig. 4A and at $\theta = 0.13$ in Fig. 4B, spreading
peripherally (rightwards) in both cases, while degeneration also initiates at the ora serrata ($\theta = 1$) under
Scaling 2 in both Fig. 4A and Fig. 4B, spreading centrally. Degeneration occurs earlier in Fig. 4B than
in Fig. 4A and earlier for Scaling 2 than for Scaling 1 (except near the fovea in Fig. 4A). Numerical and
analytical solutions agree well, only diverging close to fovea in Fig. 4A, where the analytical solution
breaks down. None of these patterns of degeneration match those seen *in vivo* (see Fig. 2).

In Figs. 6–9 we calculate the $\phi_r(\theta) = \phi_{r_{\mathrm{inv}}}(\theta)$ and $f_{\mathrm{crit}}(\theta) = f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ profiles required to qualitatively
replicate the cone degeneration profiles, $t_{\mathrm{degen}}(\theta)$, observed *in vivo* (Fig. 5), by solving the associated
inverse problems (see Section 2.3). Analytical inverses are plotted across the domain ($0 \leq \theta \leq 1$), while

numerical inverses are calculated and plotted only at those locations (eccentricities) where the analytical
inverse fails to generate a $t_{\text{degen}}(\theta)$ profile matching the target profile.

In Fig. 6 we calculate inverses for a Uniform degeneration profile. While this pattern is not typically
observed in humans (though present in animal models of RP), we consider this case as a point of comparison
with the non-uniform patterns explored in Figs. 7–9. Both inverses, $\phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}_{\text{inv}}}(\theta)$, are monotone
increasing for Scaling 1, and increase initially for Scaling 2 before reaching a maximum and decreasing
toward the ora serrata (at $\theta = 1$). Consequently, Scaling 1 and 2 inverses, while close near the fovea
($\theta = 0$), diverge toward the ora serrata, this effect being more prominent for $f_{\text{crit}_{\text{inv}}}(\theta)$. The inverse profiles
have a similar shape to the $t_{\text{degen}}(\theta)$ profiles in Fig. 4 (see Discussion). Numerical solutions reveal lower
values of the inverses near the fovea, where the analytical approximations break down.

Inverses for linear (Fig. 7A and B), concave up (quadratic) (Fig. 7C and D) and concave down (quadratic)
(Fig. 7E and F) Pattern 1A degeneration profiles are shown in Fig. 7. Inverses are monotone increasing
functions for both Scalings 1 and 2 in Fig. 7A,B,E and F, and for Scaling 1 in Fig. 7C and D, while the
inverses increase initially for Scaling 2 before reaching a maximum and decreasing toward the ora serrata in
Fig. 7C and D. Numerical solutions reveal lower values of the inverses near the fovea, where the analytical
approximations break down.

Fig. 8 shows inverses for linear (Fig. 8A and B), quadratic (Fig. 8C and D) and exponential (Fig. 8E and
F) Pattern 1B degeneration profiles. Inverses resemble vertically flipped versions of the $t_{\text{degen}}(\theta)$ profiles
in Fig. 5C (see Discussion). Numerical solutions reveal lower values of the inverses near the fovea, where
the analytical approximations break down, and higher values in some regions away from the fovea in Fig.
8A–D.

In Fig. 9 we calculate inverses for linear 1 (Fig. 9A and B), linear 2 (Fig. 9C and D), quadratic (Fig.
9E and F) and cubic (Fig. 9G and H) Pattern 3 degeneration profiles. Inverses resemble vertically flipped
versions of the $t_{\text{degen}}(\theta)$ profiles in Fig. 5D (see Discussion). Numerical solutions reveal lower values of
the inverses near the fovea, where the analytical approximations break down, and higher values in some
regions away from the fovea in Fig. 9C–F and H.

Lastly, in Fig. 10, we show simulation results of proportional cone loss for analytical and numerical
$\phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}_{\text{inv}}}(\theta)$, for Uniform (Scaling 1, Fig. 10A–D), concave up Pattern 1A (Scaling 1, Fig.
10E–H), linear Pattern 1B (Scaling 2, Fig. 10I–L) and quadratic Pattern 3 (Scaling 2, Fig. 10M–P)
target degeneration profiles. Cone degeneration profiles generally show good agreement with the target
$t_{\text{degen}}(\theta)$ profiles. There is some divergence from $t_{\text{degen}}(\theta)$ for the analytical inverses near the fovea and
at discontinuous or nonsmooth portions of $t_{\text{degen}}(\theta)$; this is mostly corrected by the numerical inverses.
Premature degeneration near the fovea might be avoided biologically, but cannot be completely avoided
numerically, even when using the more accurate numerical inverses, since it is necessary to replace the
Heaviside step function in $\lambda_2(f)$ (see Eqn. (3)) with a hyperbolic `tanh` function to satisfy the smoothness
requirements for the numerical solver, with the result that the initiation of cone degeneration is sensitive to
the low TF concentrations ($< 10^{-4}$) in that region.

4 DISCUSSION

The spatio-temporal patterns of retinal degeneration observed in human retinitis pigmentosa (RP) are
well characterised; however, the mechanistic explanation for these patterns has yet to be conclusively
determined. In this paper, we have formulated a one-dimensional (1D) reaction-diffusion partial differential
equation (PDE) model (modified from Roberts, 2021) to predict RP progression under the trophic factor

(TF) hypothesis. Using this model, we solved inverse problems to determine the rate of mutation-induced
rod loss profiles, $\phi_r(\theta) = \phi_{r_{\text{inv}}}(\theta)$, and TF threshold concentration profiles, $f_{\text{crit}}(\theta) = f_{\text{crit}_{\text{inv}}}(\theta)$, that
would be required to generate spatio-temporal patterns of cone degeneration qualitatively resembling
those observed *in vivo*, were the TF mechanism solely responsible for RP progression. In reality, multiple
mechanisms (including oxidative damage and metabolic dysregulation, Punzo et al., 2009, 2012; Stone
et al., 1999; Travis et al., 1991; Valter et al., 1998) likely operate in tandem to drive the initiation and
propagation of retinal degeneration in RP. By using mathematics to isolate the TF mechanism, in a way
that would be impossible to achieve experimentally, we are able to determine the conditions under which
the TF mechanism alone would recapitulate known phenotypes. Having identified these conditions, this
paves the way for future biomedical and experimental studies to test our predictions.

Spatially uniform $\phi_r(\theta)$ and $f_{\text{crit}}(\theta)$ profiles fail to replicate any of the *in vivo* patterns of degeneration
(Fig. 4), showing that heterogenous profiles are required, all else being equal. Throughout this paper we
have considered two scalings on the rate of TF production by rods, α, and the rate of TF consumption by
cones, β (denoted as Scalings 1 and 2, see Section 2.3 for details). Under Scaling 1, the rod:cone ratio (Fig.
3B) dominates the model behaviour (see Eqn. (6)), leading to a monotone, central to peripheral pattern of
degeneration, while under Scaling 2, the trophic factor decay term, ηf, enters the dominant balance (see
Eqn. (9)), such that degeneration initiates at both the fovea and (later) at the ora serrata, the degenerative
fronts meeting in the mid-/far-periphery (Fig. 4).

We solved the inverse functions, $\phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}_{\text{inv}}}(\theta)$, both analytically (algebraically) and numeri-
cally (computationally). Analytical solutions are approximations; however, they have the advantage of
being easier to compute (increasing their utility for biomedical researchers) and provide a more intuitive
understanding of model behaviour, while numerical solutions are more accurate, though computationally
expensive. We calculated the inverses for a range of target cone degeneration profiles, consisting of a
Uniform profile and profiles which qualitatively replicate those found *in vivo*: Pattern 1A, Pattern 1B and
Pattern 3 (Pattern 2 being inaccessible to a 1D model; see Table 3 and Fig. 5).

The shapes of the inverse functions are determined partly by the rod and cone distributions, $\tilde{p}_r(\theta)$ and
$\tilde{p}_c(\theta)$, and partly by the target cone degeneration profile, $t_{\text{degen}}(\theta)$ (see Eqs. (7), (8), (10) and (11)). As
such, in the Uniform case (Fig. 6), the Scaling 1 inverse profiles take a similar shape to the rod:cone ratio
(Fig. 3B), the inverses being lower towards the fovea to compensate for the smaller rod:cone ratio and hence
lower supply of TF to each cone. The Scaling 2 inverse profiles follow a similar trend but decrease toward
the ora serrata after peaking in the mid-/far-periphery due to the greater influence of the trophic factor
decay term under this scaling. Interestingly, the shapes of these inverse profiles bear a striking resemblance
to the cone degeneration profiles for spatially uniform $\phi_r(\theta)$ and $f_{\text{crit}}(\theta)$ (Fig. 4). This is because lower
values of the inverses are required to delay degeneration, in those regions where cones would otherwise
degenerate earlier, to achieve a uniform degeneration profile. The inverse functions resemble vertically
flipped versions of the target cone degeneration profiles for Patterns 1A, 1B and 3 (Figs. 7–9), this being
more apparent for Patterns 1B and 3 due to their more distinctive shapes. This makes sense since lower
inverse values are required for later degeneration times. Scaling 2 inverses typically lie below Scaling 1
inverses, compensating for the fact that degeneration generally occurs earlier under Scaling 2 than under
Scaling 1 for any given $\phi_r(\theta)$ and $f_{\text{crit}}(\theta)$.

Analytical inverses give rise to cone degeneration profiles that accurately match the target cone degenera-
tion profiles, except near the fovea (centred at $\theta = 0$, where the validity of the analytical approximation
breaks down) and where the target $t_{\text{degen}}(\theta)$ profile is nonsmooth or discontinuous (i.e. linear and quadratic
Pattern 1B, and linear 1, linear 2 and quadratic Pattern 3; see Fig. 10 for examples). Numerical inverses

improve accuracy in these regions, consistently taking lower values near the fovea, delaying degeneration
where it occurs prematurely under the analytical approximation.

We have assumed throughout this study that at least one of $\phi_r(\theta)$ and $f_{\text{crit}}(\theta)$ is spatially uniform. It is
possible, however, that both vary spatially. In this case there are no unique inverses; however, if the profile
for one of these functions could be measured experimentally, then the inverse problem for the remaining
function could be solved as in this paper.

This work could be extended both experimentally and theoretically. Experimental and biomedical studies
could measure how the rate of mutation-induced rod loss and TF threshold concentration vary with location
in the retina, noting the spatio-temporal pattern of retinal degeneration and comparing with the inverse
$\phi_{r_{\text{inv}}}(\theta)$ and $f_{\text{crit}_{\text{inv}}}(\theta)$ profiles predicted by our models. Curcio et al. (1993) have previously measured
variation in the rate of rod loss in normal (non-RP) human retinas (where rods degenerated most rapidly in
the central retina); however, to the best of our knowledge, detailed measurements have not been made in
the RP retina. Measuring the TF threshold concentration may prove more challenging, but may be possible
in animal models or inferred from *in vitro/ex vivo* studies. In future work, we will extend our mathematical
model to two spatial dimensions, accounting for variation in the azimuthal dimension, and adapt the model
to consider animal retinas for which the photoreceptor distribution has been well characterised (e.g. rats,
mice and pigs, Chandler et al., 1999; Gaillard et al., 2009; Ortín-Martínez et al., 2014).

In conclusion, we have formulated and solved a mathematical inverse problem to determine the rate of
mutation-induced rod loss and TF threshold concentration profiles required to explain the spatio-temporal
patterns of retinal degeneration observed in human RP. Inverse profiles were calculated for a set of
qualitatively distinct degeneration patterns, achieving a close match with the target cone degeneration
profiles. Predicted inverse profiles await future experimental verification.

CONFLICT OF INTEREST STATEMENT

313 The author declares that the research was conducted in the absence of any commercial or financial
314 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

315 PAR: conceptualisation, methodology, software, validation, formal analysis, investigation, data curation,
316 writing —- original draft, writing —- review and editing, visualisation, and project administration.

FUNDING

317 P.A.R. is funded by BBSRC (BB/R014817/1).

ACKNOWLEDGMENTS

318 P.A.R. thanks Tom Baden for allowing the time to pursue this research.

REFERENCES

Aït-Ali, N., Fridlich, R., Millet-Puel, G., Clérin, E., Delalande, F., Jaillard, C., et al. (2015). Rod-derived
cone viability factor promotes cone survival by stimulating aerobic glycolysis. *Cell* 161, 817–832.
doi:https://doi.org/10.1016/j.cell.2015.03.023
Burns, J., Clarke, G., and Lumsden, C. J. (2002). Photoreceptor death: Spatiotemporal patterns arising
from one-hit death kinetics and a diffusible cell death factor. *Bull. Math. Biol.* 64, 1117–1145.
doi:https://doi.org/10.1006/bulm.2002.0320

Camacho, E. T., Brager, D., Elachouri, G., Korneyeva, T., Millet-Puel, G., Sahel, J. A., et al. (2019). A
mathematical analysis of aerobic glycolysis triggered by glucose uptake in cones. *Sci. Rep.* 9, 4162.
doi:https://doi.org/10.1038/s41598-019-39901-z

Camacho, E. T., Colón Vélez, M. A., Hernández, D. J., Bernier, U. R., van Laarhoven, J., and Wirkus,
S. (2010). A mathematical model for photoreceptor interactions. *J. Theor. Biol.* 267, 638–646.
doi:https://doi.org/10.1016/j.jtbi.2010.09.006

Camacho, E. T., Dobreva, A., Larripa, K., Rădulescu, A., Schmidt, D., and Trejo, I. (2021). *Mathematical*
Modeling of Retinal Degeneration: Aerobic Glycolysis in a Single Cone (Cham: Springer International
Publishing), vol. 22 of *Association for Women in Mathematics Series*. 135–178. doi:https://doi.org/10.
1007/978-3-030-57129-0_7

Camacho, E. T., Lenhart, S., Melara, L. A., Villalobos, M. C., and Wirkus, S. (2020). Optimal control with
MANF treatment of photoreceptor degeneration. *Math. Med. Biol.* 37, 1–21. doi:https://doi.org/10.1093/
imammb/dqz003

Camacho, E. T., Léveillard, T., Sahel, J. A., and Wirkus, S. (2016a). Mathematical model of the role
of RdCVF in the coexistence of rods and cones in a healthy eye. *Bull. Math. Biol.* 78, 1394–1409.
doi:https://doi.org/10.1007/s11538-016-0185-x

Camacho, E. T., Melara, L. A., Villalobos, M. C., and Wirkus, S. (2014). Optimal control in the treatment
of retinitis pigmentosa. *Bull. Math. Biol.* 76, 292–313. doi:https://doi.org/10.1007/s11538-013-9919-1

Camacho, E. T., Punzo, C., and Wirkus, S. A. (2016b). Quantifying the metabolic contribution to
photoreceptor death in retinitis pigmentosa via a mathematical model. *J. Theor. Biol.* 408, 75–87.
doi:https://doi.org/10.1016/j.jtbi.2016.08.001

Camacho, E. T., Radulescu, A., and Wirkus, S. (2016c). Bifurcation analysis of a photoreceptor interaction
model for retinitis pigmentosa. *Commun. Nonlinear Sci. Numer. Simulat.* 38, 267–276. doi:https:
//doi.org/10.1016/j.cnsns.2016.02.030

Camacho, E. T. and Wirkus, S. (2013). Tracing the progression of retinitis pigmentosa via photoreceptor
interactions. *J. Theor. Biol.* 317, 105–118. doi:https://doi.org/10.1016/j.jtbi.2012.09.034

Chandler, M. J., Smith, P. J., Samuelson, D. A., and Mackay, E. O. (1999). Photoreceptor density of the
domestic pig retina. *Vet. Ophthalmol.* 2, 179–184. doi:https://doi.org/10.1046/j.1463-5224.1999.00077.x

Colón Vélez, M. A., Hernández, D. J., Bernier, U. R., van Laarhoven, J., and Camacho, E. T. (2003).
Mathematical Models for Photoreceptor Interactions. Tech. rep., Cornell University, Department of
Biological Statistics and Computational Biology

Curcio, C. A., Millican, C. L., Allen, K. A., and Kalina, R. E. (1993). Aging of the human photoreceptor
mosaic: evidence for selective vulnerability of rods in central retina. *Invest. Ophthalmol. Vis. Sci.* 34,
3278–3296

Curcio, C. A., Sloan, K. R., Kalina, R. E., and Hendrickson, A. E. (1990). Human photoreceptor topography.
J. Comp. Neurol. 292, 497–523. doi:https://doi.org/10.1002/cne.902920402

Daiger, S. P., Bowne, S. J., and Sullivan, L. S. (2007). Perspective on genes and mutations causing retinitis
pigmentosa. *Arch. Ophthalmol.* 125, 151–158. doi:10.1001/archopht.125.2.151

Fintz, A. C., Audo, I., Hicks, D., Mohand-Saïd, S., Léveillard, T., and Sahel, J. (2003). Partial characteriza-
tion of retina-derived cone neuroprotection in two culture models of photoreceptor degeneration. *Invest.*
Ophthalmol. Vis. Sci. 44, 818–825. doi:https://doi.org/10.1167/iovs.01-1144

Gaillard, F., Kuny, S., and Sauvé, Y. (2009). Topographic arrangement of S-cone photoreceptors in the
retina of the diurnal nile grass rat (arvicanthis niloticus). *Invest. Ophthalmol. Vis. Sci.* 50, 5426–5434.
doi:https://doi.org/10.1167/iovs.09-3896

Grover, S., Fishman, G. A., and Brown Jr, J. (1998). Patterns of visual field progression in patients with
retinitis pigmentosa. *Ophthalmology* 105, 1069–1075. doi:https://doi.org/10.1016/S0161-6420(98)
96009-2

Gupta, N., Brown, K. E., and Milam, A. H. (2003). Activated microglia in human retinitis pigmentosa,
late-onset retinal degeneration, and age-related macular degeneration. *Exp. Eye Res.* 76, 463–471.
doi:https://doi.org/10.1016/S0014-4835(02)00332-9

Hamel, C. (2006). Retinitis pigmentosa. *Orphanet. J. Rare Dis.* 1, 40. doi:https://doi.org/10.1186/
1750-1172-1-40

Hartong, D. T., Berson, E. L., and Dryja, T. P. (2006). Retinitis pigmentosa. *Lancet* 368, 1795–1809.
doi:https://doi.org/10.1016/S0140-6736(06)69740-7

Léveillard, T., Mohand-Saïd, S., Lorentz, O., Hicks, D., Fintz, A. C., Clérin, E., et al. (2004). Identification
and characterization of rod-derived cone viability factor. *Nat. Genet.* 36, 755–759. doi:https://doi.org/10.
1038/ng1386

Léveillard, T. and Sahel, J. A. (2017). Metabolic and redox signaling in the retina. *Cell. Mol. Life Sci.* 74,
3649–3665. doi:https://doi.org/10.1007/s00018-016-2318-7

Mei, X., Chaffiol, A., Kole, C., Yang, Y., Millet-Puel, G., Clérin, E., et al. (2016). The thioredoxin
encoded by the rod-derived cone viability factor gene protects cone photoreceptors against oxidative
stress. *Antioxidants & Redox Signaling* 24, 909–923. doi:https://doi.org/10.1089/ars.2015.6509

Mohand-Saïd, S., Deudon-Combe, A., Hicks, D., Simonutti, M., Forster, V., Fintz, A. C., et al. (1998).
Normal retina releases a diffusible factor stimulating cone survival in the retinal degeneration mouse.
Proc. Natl. Acad. Sci. 95, 8357–8362. doi:https://doi.org/10.1073/pnas.95.14.8357

Mohand-Saïd, S., Hicks, D., Dreyfus, H., and Sahel, J. A. (2000). Selective transplantation of rods delays
cone loss in a retinitis pigmentosa model. *Arch. Ophthalmol.* 118, 807–811. doi:10.1001/archopht.118.6.
807

Mohand-Saïd, S., Hicks, D., Simonutti, M., Tran-Minh, D., Deudon-Combe, A., Dreyfus, H., et al. (1997).
Photoreceptor transplants increase host cone survival in the retinal degeneration (rd) mouse. *Ophthalmic
Res.* 29, 290–297. doi:https://doi.org/10.1159/000268027

Ortín-Martínez, A., Nadal-Nicolás, F. M., Jiménez-López, M., Alburquerque-Béjar, J. J., Nieto-López,
L., García-Ayuso, D., et al. (2014). Number and distribution of mouse retinal cone photoreceptors:
Differences between an albino (Swiss) and a pigmented (C57/BL6) strain. *PLoS One* 9, e102392.
doi:https://doi.org/10.1371/journal.pone.0102392

Oyster, C. W. (1999). *The Human Eye: Structure and Function* (Sinauer Associates Inc.)

Punzo, C., Kornacker, K., and Cepko, C. L. (2009). Stimulation of the insulin/mTOR pathway delays cone
death in a mouse model of retinitis pigmentosa. *Nat. Neurosci.* 12, 44–52. doi:https://doi.org/10.1038/
nn.2234

Punzo, C., Xiong, W., and Cepko, C. L. (2012). Loss of daylight vision in retinal degeneration: Are
oxidative stress and metabolic dysregulation to blame? *J. Biol. Chem.* 287, 1642–1648. doi:https:
//doi.org/10.1074/jbc.R111.304428

Ripps, H. (2002). Cell death in retinitis pigmentosa: Gap junctions and the 'bystander' effect. *Exp. Eye
Res.* 74, 327–336. doi:https://doi.org/10.1006/exer.2002.1155

Roberts, P. A. (2021). Mathematical models of retinitis pigmentosa: The trophic factor hypothesis. *bioRxiv*
doi:https://doi.org/10.1101/2021.03.25.436992

Roberts, P. A., Gaffney, E. A., Luthert, P. J., Foss, A. J. E., and Byrne, H. M. (2016a). Mathematical and
computational models of the retina in health, development and disease. *Prog. Retin. Eye. Res.* 53, 48–69.
doi:https://doi.org/10.1016/j.preteyeres.2016.04.001

Roberts, P. A., Gaffney, E. A., Luthert, P. J., Foss, A. J. E., and Byrne, H. M. (2016b). Retinal oxygen distribution and the role of neuroglobin. *J. Math. Biol.* 73, 1–38. doi:https://doi.org/10.1007/s00285-015-0931-y

Roberts, P. A., Gaffney, E. A., Luthert, P. J., Foss, A. J. E., and Byrne, H. M. (2017). Mathematical models of retinitis pigmentosa: The oxygen toxicity hypothesis. *J. Theor. Biol.* 425, 53–71. doi:https://doi.org/10.1016/j.jtbi.2017.05.006

Roberts, P. A., Gaffney, E. A., Whiteley, J. P., Luthert, P. J., Foss, A. J. E., and Byrne, H. M. (2018). Predictive mathematical models for the spread and treatment of hyperoxia-induced photoreceptor degeneration in retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 59, 1238–1249. doi:https://doi.org/10.1167/iovs.17-23177

Stone, J., Maslim, J., Valter-Kocsi, K., Mervin, K., Bowers, F., Chu, Y., et al. (1999). Mechanisms of photoreceptor death and survival in mammalian retina. *Prog. Retin. Eye Res.* 18(6), 689–735. doi:https://doi.org/10.1016/S1350-9462(98)00032-9

Travis, G. H., Sutcliffe, J. G., and Bok, D. (1991). The retinal degeneration slow (rds) gene product is a photoreceptor disc membrane-associated glycoprotein. *Neuron* 6, 61–70. doi:https://doi.org/10.1016/0896-6273(91)90122-G

Valter, K., Maslim, J., Bowers, F., and Stone, J. (1998). Photoreceptor dystrophy in the RCS rat: roles of oxygen, debris, and bFGF. *Invest. Ophthalmol. Vis. Sci.* 39, 2427–2442

Wifvat, K., Camacho, E. T., Wirkus, S., and Léveillard, T. (2021). The role of RdCVFL in a mathematical model of photoreceptor interactions. *J. Theor. Biol.* 520, 110642. doi:https://doi.org/10.1016/j.jtbi.2021.110642

Yang, Y., Mohand-Said, S., Danan, A., Simonutti, M., Fontaine, V., Clérin, E., et al. (2009). Functional cone rescue by RdCVF protein in a dominant model of retinitis pigmentosa. *Mol. Ther.* 17, 787–795. doi:https://doi.org/10.1038/mt.2009.28

Figure 1. Diagrams of the human eye and retinal photoreceptor distribution (reproduced, with permission, from Roberts et al., 2017). **(A)** Diagram of the (right) human eye, viewed in the transverse plane, illustrating the mathematical model geometry. The model is posed on a domain spanning the region between the foveal centre, at $\theta = 0$, and the ora serrata, at $\theta = \Theta$, along the temporal horizontal meridian, where θ measures the eccentricity. Figure originally reproduced, with modifications, from `http://www.nei.nih.gov/health/coloboma/coloboma.asp`, courtesy: National Eye Institute, National Institutes of Health (NEI/NIH). **(B)** Measured and fitted photoreceptor profiles, along the temporal horizontal meridian, in the human retina. Cone profile: $\tilde{p}_c(\theta) = B_1 e^{-b_1\theta} + B_2 e^{-b_2\theta}$, and rod profile: $\tilde{p}_r(\theta) = B_3 \theta e^{-b_3\theta}$ (see Table 2 for dimensionless parameter values). The photoreceptor profile is the sum of the rod and cone profiles ($\tilde{p}_r(\theta) + \tilde{p}_c(\theta)$). Experimental data provided by Curcio and published in Curcio et al. (1990).

FIGURES & TABLES

Figure 2. Characteristic patterns of retinal degeneration in human RP (reproduced, with permission and modifications, from Roberts et al., 2018). Degeneration patterns can be classified into three cases and six subcases (classification system described in Grover et al., 1998). Large grey arrows indicate transitions between stages of degeneration and small red arrows indicate the direction of degeneration propagation. See text for details.

Table 1. Variables employed in the non-dimensional mathematical model (Eqs. (1)–(5)).

Variable	Description
θ	Eccentricity
t	Time
$f(\theta, t)$	Trophic factor concentration
$p_r(\theta, t)$	Rod density
$p_c(\theta, t)$	Cone density

Figure 3. Initial conditions and ratio of rods to cones. **(A)** initial conditions used in all simulations, consisting of healthy rod and cone profiles and the corresponding RdCVF profiles under Scalings 1 and 2 (the legend applies to **(A)** only). **(B)** variation in the healthy rod:cone ratio, $\tilde{p}_r(\theta)/\tilde{p}_c(\theta)$, with eccentricity. To obtain $f_{\text{init}}(\theta)$ in **(A)**, Eqs. (1) and (4) were solved at steady-state using the finite difference method, with 4001 mesh points, where $p_r(\theta) = p_{r_{\text{init}}}(\theta)$ and $p_c(\theta) = p_{c_{\text{init}}}(\theta)$. Under Scaling 1, $\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$, while under Scaling 2, $\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$. Remaining parameter values as in Table 2.

Figure 4. Cone degeneration profiles. Graphs show the time, $t_{\text{degen}}(\theta)$, at which cones degenerate due to RdCVF deprivation, with constant rate of mutation-induced rod degeneration, $\phi_r = 7.33 \times 10^{-2}$, and constant TF threshold concentrations: $f_{\text{crit}} = 3 \times 10^{-5}$ **(A)** and $f_{\text{crit}} = 0.3$ **(B)**. The solid black and dashed green curves correspond to Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$), while the solid blue and dashed red curves correspond to Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$). The black and blue solid curves are analytical approximations, obtained by plotting Eqs. (6) and (9) respectively, while the green and red dashed curves are $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$ contours, obtained by solving Eqs. (1)–(5) using the method of lines with 401 mesh points. **(A)** simulation spans ~17.7 years in dimensional variables; **(B)** simulation spans ~2.8 years in dimensional variables. Insets show magnified portions of each graph. Cone degeneration initiates at the fovea ($\theta = 0$) in **(A)** and at $\theta = 0.13$ in **(B)**, spreading peripherally (rightwards) in both cases. Degeneration occurs earlier in **(B)** than in **(A)** and for Scaling 2 than for Scaling 1 (except near the fovea in **(A)**). Remaining parameter values as in Table 2.

Figure 5. Target cone degeneration profiles. Panels show cone degeneration profiles, $t_{\mathrm{degen}}(\theta)$, qualitatively replicating typical spatio-temporal patterns of retinal degeneration in RP: **(A)** Uniform, **(B)** Pattern 1A, **(C)** Pattern 1B and **(D)** Pattern 3. We seek to replicate these patterns by finding appropriate $\phi_{r_{\mathrm{inv}}}(\theta)$ and $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ profiles in Figs. 6–9. Inset diagrams show the corresponding 2D patterns of retinal degeneration — white regions: healthy retina, black regions: degenerate retina, and red arrows: direction of propagation of degeneration. Parameters: $t_0 = 100$ (~ 11.0 years), $t_1 = 150$ (~ 16.6 years), $t_2 = 200$ (~ 22.1 years, $\theta_{d_1} = 0.1$ (~ 7.6 degrees), $\theta_{d_2} = 0.55$ (~ 41.9 degrees), $\theta_{d_3} = 0.3$ (~ 22.9 degrees), $\theta_{d_4} = 0.4$ (~ 30.5 degrees) and $\theta_{d_5} = 0.6$ (~ 45.7 degrees). Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Figure 6. Inverse mutation-induced rod degeneration rate and TF threshold concentration — Uniform target cone degeneration profile. **(A)** inverse mutation-induced rod degeneration rate, $\phi_{r_{\text{inv}}}(\theta)$ ($f_{\text{crit}} = 3 \times 10^{-5}$); **(B)** inverse TF threshold concentration, $f_{\text{crit}_{\text{inv}}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$). The solid black and dashed green curves correspond to Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$), while the solid blue and dashed red curves correspond to Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$). The black and blue solid curves are analytical approximations to the inverses, obtained by plotting Eqs. (7) and (10) respectively **(A)**, and Eqs. (8) and (11) respectively **(B)**. The green and red dashed curves are numerical inverses, obtained by using the Matlab routines fminsearch **(A)** and patternsearch **(B)** to calculate the ϕ_r and f_{crit} profiles for which the contour described by $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$ matches the target cone degeneration profile, $t_{\text{degen}}(\theta)$. Eqs. (1)–(5) were solved at each iteration using the method of lines, with 101 mesh points. Insets show magnified portions of each graph. Numerical inverses are calculated and plotted only at those locations (eccentricities) where the analytical inverse fails to generate a $t_{\text{degen}}(\theta)$ profile matching the target profile. Inverses are monotone increasing for Scaling 1, and increase initially for Scaling 2 before reaching a maximum and decreasing toward the ora serrata ($\theta = 1$). Numerical solutions reveal lower values of the inverses near the fovea ($\theta = 0$) than the analytical approximations suggest. Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Table 2. Parameters employed in the non-dimensional mathematical model (Eqs. (1)–(5)). Values are given to three significant figures (radians are dimensionless).

Parameter	Description	Value
Θ	Eccentricity of the ora serrata	1.33 rad
D_f	Trophic factor diffusivity	0.237
α	Rate of trophic factor production by rods	7.01×10^2 or 7.01×10^4
β	Rate of trophic factor consumption by cones	1.79×10^4 or 1.79×10^6
η	Rate of trophic factor decay	1.79×10^2
ϕ_r	Rate of mutation-induced rod degeneration	7.33×10^{-2}
f_{crit}	Trophic factor threshold concentration	3×10^{-5} or 0.3
B_1	Cone profile parameter	1.56
B_2	Cone profile parameter	0.158
B_3	Rod profile parameter	10.6
b_1	Cone profile parameter	71.8
b_2	Cone profile parameter	2.67
b_3	Rod profile parameter	3.06

Figure 7. Inverse mutation-induced rod degeneration rate and TF threshold concentration — Pattern 1A target cone degeneration profiles. **(A)**, **(C)** and **(E)** inverse mutation-induced rod degeneration rate, $\phi_{r_{\text{inv}}}(\theta)$ ($f_{\text{crit}} = 3 \times 10^{-5}$); **(B)**, **(D)** and **(F)** inverse TF threshold concentration, $f_{\text{crit}_{\text{inv}}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$). **(A)** and **(B)** linear target cone degeneration profile, $t_{\text{degen}}(\theta)$; **(C)** and **(D)** concave up quadratic $t_{\text{degen}}(\theta)$ profile; **(E)** and **(F)** concave down quadratic $t_{\text{degen}}(\theta)$ profile. The solid black and dashed green curves correspond to Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$), while the solid blue and dashed red curves correspond to Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$). The black and blue solid curves are analytical approximations to the inverses, obtained by plotting Eqs. (7) and (10) respectively **(A)**, **(C)** and **(E)**, and Eqs. (8) and (11) respectively **(B)**, **(D)** and **(F)**. The green and red dashed curves are numerical inverses, obtained by using the Matlab routines fminsearch **(A)**, **(C)** and **(E)**, and patternsearch **(B)**, **(D)** and **(F)** to calculate the ϕ_r and f_{crit} profiles for which the contour described by $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$ matches the target cone degeneration profile, $t_{\text{degen}}(\theta)$. Eqs. (1)–(5) were solved at each iteration using the method of lines, with 26, 51 or 101 mesh points. Insets show magnified portions of each graph. Numerical inverses are calculated and plotted only at those locations (eccentricities) where the analytical inverse fails to generate a $t_{\text{degen}}(\theta)$ profile matching the target profile. Inverses are monotone increasing functions for both scalings in **(A)**, **(B)**, **(E)** and **(F)**, and for Scaling 1 in **(C)** and **(D)**, while the inverses increase initially for Scaling 2 before reaching a maximum and decreasing toward the ora serrata ($\theta = 1$) in **(C)** and **(D)**. Numerical solutions reveal lower values of the inverses near the fovea ($\theta = 0$) than the analytical approximations suggest. Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Figure 8. Inverse mutation-induced rod degeneration rate and TF threshold concentration — Pattern 1B target cone degeneration profiles. **(A)**, **(C)** and **(E)** inverse mutation-induced rod degeneration rate, $\phi_{r_{\text{inv}}}(\theta)$ ($f_{\text{crit}} = 3 \times 10^{-5}$); **(B)**, **(D)** and **(F)** inverse TF threshold concentration, $f_{\text{crit}_{\text{inv}}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$). **(A)** and **(B)** linear target cone degeneration profile, $t_{\text{degen}}(\theta)$; **(C)** and **(D)** quadratic $t_{\text{degen}}(\theta)$ profile; **(E)** and **(F)** exponential $t_{\text{degen}}(\theta)$ profile. The solid black and dashed green curves correspond to Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$), while the solid blue and dashed red curves correspond to Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$). The black and blue solid curves are analytical approximations to the inverses, obtained by plotting Eqs. (7) and (10) respectively **(A)**, **(C)** and **(E)**, and Eqs. (8) and (11) respectively **(B)**, **(D)** and **(F)**. The green and red dashed curves are numerical inverses, obtained by using the Matlab routines fminsearch **(A)**, **(C)** and **(E)**, and patternsearch **(B)**, **(D)** and **(F)** to calculate the ϕ_r and f_{crit} profiles for which the contour described by $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$ matches the target cone degeneration profile, $t_{\text{degen}}(\theta)$. Eqs. (1)–(5) were solved at each iteration using the method of lines, with 51 or 101 mesh points. Insets show magnified portions of each graph. Numerical inverses are calculated and plotted only at those locations (eccentricities) where the analytical inverse fails to generate a $t_{\text{degen}}(\theta)$ profile matching the target profile. Inverses resemble vertically flipped versions of the $t_{\text{degen}}(\theta)$ profiles. Numerical solutions reveal lower values of the inverses near the fovea ($\theta = 0$) than the analytical approximations suggest and higher values in some regions away from the fovea in **(A)**–**(D)**. Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Figure 9. Inverse mutation-induced rod degeneration rate and TF threshold concentration — Pattern 3 target cone degeneration profiles. **(A)**, **(C)**, **(E)** and **(G)** inverse mutation-induced rod degeneration rate, $\phi_{r_{inv}}(\theta)$ ($f_{crit} = 3 \times 10^{-5}$); **(B)**, **(D)**, **(F)** and **(H)** inverse TF threshold concentration, $f_{crit_{inv}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$). **(A)** and **(B)** linear 1 target cone degeneration profile, $t_{degen}(\theta)$; **(C)** and **(D)** linear 2 $t_{degen}(\theta)$ profile; **(E)** and **(F)** quadratic $t_{degen}(\theta)$ profile; **(G)** and **(H)** cubic $t_{degen}(\theta)$ profile. The solid black and dashed green curves correspond to Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$), while the solid blue and dashed red curves correspond to Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$). The black and blue solid curves are analytical approximations to the inverses, obtained by plotting Eqs. (7) and (10) respectively **(A)**, **(C)**, **(E)** and **(G)**, and Eqs. (8) and (11) respectively **(B)**, **(D)**, **(F)** and **(H)**. The green and red dashed curves are numerical inverses, obtained by using the Matlab routines fminsearch **(A)**, **(C)**, **(E)** and **(G)**, and patternsearch **(B)**, **(D)**, **(F)** and **(H)** to calculate the ϕ_r and f_{crit} profiles for which the contour described by $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$ matches the target cone degeneration profile, $t_{degen}(\theta)$. Eqs. (1)–(5) were solved at each iteration using the method of lines, with 26, 51 or 101 mesh points. Insets show magnified portions of each graph. Numerical inverses are calculated and plotted only at those locations (eccentricities) where the analytical inverse fails to generate a $t_{degen}(\theta)$ profile matching the target profile. Inverses resemble vertically flipped versions of the $t_{degen}(\theta)$ profiles. Numerical solutions reveal lower values of the inverses near the fovea ($\theta = 0$) than the analytical approximations suggest and higher values in some regions away from the fovea in **(C)**–**(F)** and **(H)**. Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Figure 10. Simulations of proportional cone loss for a range of inverse mutation-induced rod degeneration rates and TF threshold concentrations. Plots show the proportion of cones remaining compared to local healthy values, $p_c(\theta, t)/\tilde{p}_c(\theta)$, across space and over time. **(A)**, **(E)**, **(I)** and **(M)** analytical inverse mutation-induced rod degeneration rate, $\phi_{r_{\mathrm{inv}}}(\theta)$ ($f_{\mathrm{crit}} = 3 \times 10^{-5}$); **(B)**, **(F)**, **(J)** and **(N)** numerical $\phi_{r_{\mathrm{inv}}}(\theta)$ ($f_{\mathrm{crit}} = 3 \times 10^{-5}$); **(C)**, **(G)**, **(K)** and **(O)** analytical inverse TF threshold concentration, $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$); **(D)**, **(H)**, **(L)** and **(P)** numerical $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ ($\phi_r = 7.33 \times 10^{-2}$). **(A)**–**(D)** Uniform target cone degeneration profile, $t_{\mathrm{degen}}(\theta)$, with Scaling 1 ($\alpha = 7.01 \times 10^4$ and $\beta = 1.79 \times 10^6$); **(E)**–**(H)** Pattern 1A quadratic concave up $t_{\mathrm{degen}}(\theta)$ profile with Scaling 1; **(I)**–**(L)** Pattern 1B linear $t_{\mathrm{degen}}(\theta)$ profile with Scaling 2 ($\alpha = 7.01 \times 10^2$ and $\beta = 1.79 \times 10^4$); **(M)**–**(P)** Pattern 3 quadratic $t_{\mathrm{degen}}(\theta)$ profile with Scaling 2. Eqs. (1)–(5) were solved using the method of lines, with 26, 51 or 101 mesh points. Analytical and numerical $\phi_{r_{\mathrm{inv}}}(\theta)$ and $f_{\mathrm{crit}_{\mathrm{inv}}}(\theta)$ are as plotted in Figs. 6–9. Solid red curves denote the contours along which $p_c(\theta, t)/\tilde{p}_c(\theta) = 0.99$, while dashed green curves show the target $t_{\mathrm{degen}}(\theta)$ profiles. Cone degeneration profiles generally show good agreement with the target $t_{\mathrm{degen}}(\theta)$ profiles. There is some divergence from $t_{\mathrm{degen}}(\theta)$ for the analytical inverses near the fovea ($\theta = 0$) and at discontinuous or nonsmooth portions of $t_{\mathrm{degen}}(\theta)$; this is mostly corrected by the numerical inverses. Cone degeneration profile formulas and parameters are given in Table 3. Remaining parameter values as in Table 2.

Table 3. Target cone degeneration profiles, $t_{\mathrm{degen}}(\theta)$.

Degeneration Pattern	Sub-pattern	Cone Degeneration Time ($t_{\mathrm{degen}}(\theta)$)
Uniform	—	t_1
Pattern 1A	linear	$t_2 - (t_2 - t_0)\theta$
	quadratic (concave up)	$(t_2 - t_0)(\theta - 1)^2 + t_0$
	quadratic (concave down)	$t_2 - (t_2 - t_0)\theta^2$
Pattern 1B	linear	$t_2 - \frac{(t_2-t_1)}{\theta_{d_1}}\theta \quad$ if $\theta \leq \theta_{d_1}$ $t_0 + \frac{(t_1-t_0)}{(\theta_{d_2}-\theta_{d_1})}(\theta - \theta_{d_1}) \quad$ if $\theta_{d_1} < \theta \leq \theta_{d_2}$ $t_1 + \frac{(t_1-t_0)}{(1-\theta_{d_2})}(\theta_{d_2} - \theta) \quad$ if $\theta \geq \theta_{d_2}$
	quadratic	$\frac{(t_2-t_1)}{\theta_{d_1}^2}(\theta - \theta_{d_1})^2 + t_1 \quad$ if $\theta \leq \theta_{d_1}$ $t_1 - \frac{(t_1-t_0)}{(\theta_{d_2}-1)^2}(\theta - \theta_{d_2})^2 \quad$ if $\theta > \theta_{d_1}$
	exponential	$A_1 e^{-a_1\theta} + A_2\theta e^{-a_2\theta} + A_3$
Pattern 3	linear 1	$t_2 - \frac{(t_2-t_0)}{\theta_{d_4}}\theta \quad$ if $\theta \leq \theta_{d_4}$ $t_0 + \frac{(t_1-t_0)}{(1-\theta_{d_4})}(\theta - \theta_{d_4}) \quad$ if $\theta \geq \theta_{d_4}$
	linear 2	$t_2 - \frac{(t_2-t_1)}{\theta_{d_3}}\theta \quad$ if $\theta \leq \theta_{d_3}$ $t_0 \quad$ if $\theta_{d_3} < \theta \leq \theta_{d_5}$ $t_0 + \frac{(t_1-t_0)}{(1-\theta_{d_5})}(\theta - \theta_{d_5}) \quad$ if $\theta \geq \theta_{d_5}$
	quadratic	$\frac{(t_2-t_1)}{\theta_{d_3}^2}(\theta - \theta_{d_3})^2 + t_1 \quad$ if $\theta \leq \theta_{d_3}$ $\frac{(t_1-t_0)}{(1-\theta_{d_4})^2}(\theta - \theta_{d_4})^2 + t_0 \quad$ if $\theta > \theta_{d_3}$
	cubic	$C_3\theta^3 + C_2\theta^2 + C_1\theta + C_0$

Parameter Values*

$$t_0 = 100 \qquad t_1 = 150 \qquad t_2 = 200 \qquad \theta_{d_1} = 0.1 \qquad \theta_{d_2} = 0.55$$
$$\theta_{d_3} = 0.3 \qquad \theta_{d_4} = 0.4 \qquad \theta_{d_5} = 0.6 \qquad A_1 = 125 \qquad A_2 = 600$$
$$A_3 = 75 \qquad a_1 = 71.8 \qquad a_2 = 3.06$$
$$C_0 = t_2 = 200$$
$$C_1 = \frac{-2(t_2-t_0)+3(t_2-t_0)\theta_{d_4}-(t_2-t_1)\theta_{d_4}^3}{\theta_{d_4}(1-\theta_{d_4})^2} = -5.78 \times 10^2$$
$$C_2 = \frac{(t_2-t_0)-3(t_2-t_0)\theta_{d_4}^2+2(t_2-t_1)\theta_{d_4}^3}{\theta_{d_4}^2(1-\theta_{d_4})^2} = 1.01 \times 10^3$$
$$C_3 = \frac{-(t_2-t_0)+2(t_2-t_0)\theta_{d_4}-(t_2-t_1)\theta_{d_4}^2}{\theta_{d_4}^2(1-\theta_{d_4})^2} = -4.86 \times 10^2$$

* We choose θ_{d_1} and θ_{d_2} such that $\theta_{d_2} = (\theta_{d_1} + 1)/2$, so that θ_{d_2} lies half way between $\theta = \theta_{d_1}$ and $\theta = 1$.