Genome size evolution in the beetle genus *Diabrotica*.

Dimpal Lata[1], Brad S Coates[2], Kimberly K O Walden[3], Hugh M Robertson[3] and

Nicholas J Miller[1*]

1. Department of Biological Sciences, Illinois Institute of Technology, Chicago, Illinois,

USA.

2. USDA-ARS, Corn Insects & Crop Genetics Research Unit, Ames, Iowa, USA.

3. Department of Entomology, University of Illinois at Urbana-Champaign, Urbana,

Illinois USA.

*Correspondence: Nicholas J Miller, 298 R A Pritzker Science Center 3101 S Dearborn

St, Chicago, Illinois 60616, nmiller11@iit.edu, +1 312 567 3288.

<u>Abstract</u>

Diabrocite corn rootworms are one of the most economically significant pests of maize

in the United States and Europe and an emerging model for insect-plant interactions.

Genome sizes of several species in the genus *Diabrotica* were estimated using flow

cytometry along with that of *Acalymma vittatum* as an outgroup. Within the *Diabrotica*

subgroups fucata and virgifera, genome sizes ranged between 1.59 - 1.68 gigabase

pairs (Gb) and between 2.31- 2.65 Gb, respectively, and the *Acalymma vittatum*

genome size was around 1.69 Gb. This result indicated that a substantial increase in

genome size occurred in the ancestor of the virgifera group. Further analysis of fucata

group and virgifera group genome sequencing reads indicated that the genome size

difference between the *Diabrotica* subgroups could be attributed to a higher content of

transposable elements, mostly miniature inverted-transposable elements (MITEs) and

gypsy-like long terminal repeat (LTR) retroelements.

<u>Introduction</u>

The family Chrysomelidae is one of the largest families of phytophagous beetles (Order:

Coleoptera), with nearly 40,000 species. A large number of species are important

agricultural and forestry pests causing negative economic impacts (Reid, 1995; Nie et

al., 2020). A subtribe of Chrysomelidae, Diabroticina includes important agricultural

pests from the genera *Acalymma, Cerotoma,* and *Diabrotica* (Toepfer et al., 2009).

Many species of the genus *Acalymma* are specialists on Cucurbitaceae, with *Acalymma*

vittatum, the striped cucumber beetle, being one of the key pests of cucurbits in the

northeastern United States (Lewis et al., 1990). The bean leaf beetle, *Cerotoma*

trifurcata, is an important pest of leguminous crops such as peas and soybeans

throughout the eastern USA (Koch et al., 2004). *Diabrotica*, the most diverse genus

(Eben & Espinosa de los Monteros, 2013), is among the most destructive insect pests

impacting US agriculture. *Diabrotica spp.* are divided into three groups: signifera, fucata,

and virgifera, with the latter two containing recognized pest species (Krysan, 1986). The

species in the fucata group are multivoltine and polyphagous, while species in the

virgifera group are univoltine and oligophagous (Branson & Krysan, 1981; Krysan,

1982). *Diabrotica undecimpunctata* (southern corn rootworm) within the fucata group is

a generalist feeder that feeds on several crops, including cucurbits, peanuts, and maize

in the southern USA (Jackson et al., 2005). *Diabrotica virgifera virgifera* (western corn

rootworm), *Diabrotica barberi* (northern corn rootworm), and *Diabrotica virgifera zeae*

(Mexican corn rootworm) from the virgifera group are specialist feeders and are pests of

maize. *Diabrotica virgifera virgifera* is most abundant in the US Corn Belt but is found

throughout much of the United States as well as parts of Canada and Mexico.

Diabrotica virgifera virgifera and *D.barberi* are sympatric in the northern part of the US

Corn Belt, while *D. v. zeae,* is sympatric with *D. v. virgifera* over part of their range in

Texas, Arizona, and Mexico (Bragard et al., 2019)*.* Diabroticina species show a wide

range of host breadth, from monophagous to polyphagous. It has been found that the

ancestors of Diabroticina were monophagous. The genus *Acalymma* has retained the

ancestral monophagous habit, and the genus *Cerotoma* evolved oligophagy while the

genus *Diabrotica* underwent an increase in diet breadth, shifting from monophagy to

polyphagy. The fucata subgroup has retained the polyphagous characteristic, the

virgifera subgroup, on the other hand, has become secondarily oligophagous (Eben &

Espinosa de los Monteros, 2013).

As the name "corn rootworm" suggests, these insects cause substantial economic

damage to maize. The western corn rootworm, *Diabrotica virgifera virgifera*, is

considered one of the most destructive pests of maize throughout the US Corn Belt.

More than $1 billion is spent on its control and lost yield every year (Sappington et al.,

2006). The species has also been introduced into Europe and has become widespread

because of a combination of transatlantic introductions and intra-continental movement

(Ciosi et al., 2008; Miller et al., 2010; Ciosi et al., 2011). *Diabrotica barberi* is a serious

maize pest but is less widespread than *D. v. virgifera* (Capinera, 2008). Although

D.undecimpunctata is more widely distributed it is unable to survive the winter

temperatures of the US Corn Belt, and it is considered an occasional pest of maize.

Diabroticina pests have repeatedly evolved behavioral and physiological

adaptations to several management strategies like chemical control, crop rotation, and

transgenic maize expressing *Bt* proteins (Levine & Oloumi-Sadeghi, 1991; Toepfer et

al., 2009; Miller et al., 2009; Eben & Espinosa de los Monteros, 2013; Devos et al.,

2013; Gassmann et al., 2014). *Diabrotica virgifera virgifera* evolved resistance to

cyclodiene insecticides during the late 1950s (Ball & Weekman, 1962; Sappington et al.,

2006; Miller et al., 2009), and later to carbamates, the organophosphate methyl-

parathion (Meinke et al., 1998; Miller et al., 2009), and most recently to pyrethroids

(Pereira et al., 2017; Souza et al., 2019; Souza et al., 2020). Both *D. v. virgifera* and *D.*

barberi have overcome crop rotation strategies through oviposition behavior (O'Neal et

al., 2002) and extended diapause (Krysan et al., 1984), respectively. Resistance to

transgenic maize expressing Cry3Bb1 and Cry34/35Ab1 proteins has also evolved in *D.*

v. virgifera and *D. barberi* (Gassmann et al., 2014; Calles-Torrez et al., 2019).

Diabrotica virgifera virgifera has also been shown to be capable of evolving resistance

to RNA interference (RNAi) in the laboratory (Khajuria et al., 2018).

Diabroticites have not only evolved resistance to control tactics but have also

evolved tolerance to plant-derived secondary metabolites used by plants to confer a

degree of protection against several insect herbivores. The benzoxazinoid DIMBOA

(2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one) is one of the major secondary

metabolites produced by maize plants (Sasai et al., 2009). Cucurbitacins are another

group of bitter secondary metabolites found in the plants of the family Cucurbitaceae

(Rehm & Wessels, 1957). Studies have shown that *D. undecimpunctata* and *D. v.*

virgifera from the fucata and virgifera groups of *Diabrotica* can tolerate and sequester

the toxic secondary metabolites of plants such as cucurbitacins (Metcalf, 1986; Tallamy

et al., 1998) and DIMBOA (Robert et al., 2017) respectively. Although it has been found

that both these species can tolerate DIMBOA, they respond distinctly to this metabolite.

It was found *D. v. virgifera* gained significantly more dry weight when fed wild-type

plants for 10 days compared to being fed mutant plants, deficient for benzoxazinoid

biosynthesis. However, *D. undecimpunctata* performed equally well when fed on both

types of plants (Alouw & Miller, 2015). The enhanced performance of specialist *D. v.*

virgifera may be related to its ability to use DIMBOA as a signal to locate nutritious parts

of roots, while the generalist from the fucata group does not (Robert et al., 2012).

Further, RNA-Seq studies showed transcripts encoding for a CYP9-like cytochrome

P450 monooxygenase were expressed in *D. v. virgifera* larvae feeding on wild type

plants but not in larvae feeding on benzoxazinoid-deficient mutant plants (Miller & Zhao,

2015), suggesting a cytochrome P450 mediated adaptation to benzoxazinoids in *D. v.*

virgifera.

As several species in the *Diabrotica* genus can cause damage to crops and

adapt to control measures, there has been considerable interest in obtaining sequences

of their genomes and understanding genetic mechanisms for adaptations. Most

Diabrotica genetics and genomics research so far has been concentrated on *D. v.*

virgifera (Gray et al., 2009; Miller et al., 2010). The genome of *D. v. virgifera* is one of

the larger genomes among beetles and is estimated to be around 2.58 Gb (Coates et

al., 2012) whereas the average genome size for Coleoptera is 0.76Gb (Schoville et al.,

2018; Gregory, 2021). Sequencing the *D. v. virgifera* genome has posed a challenge.

Increased sizes of eukaryotic genomes are generally attributed to corresponding

increased numbers of repetitive DNA elements (Kidwell, 2002), where a large proportion

of repeats are composed of different transposable element (TE) sequences (Kojima,

2019). Eukaryotic transposons are divided into retroelements that propagate by an

RNA intermediate (class I) and DNA elements (class II) that mobilize by a "cut-and-

paste" mechanism (Finnegan, 1989; Wicker et al., 2007).

Studies also suggest that the *D. v. virgifera* genome contains a high proportion of

repetitive elements (Coates et al., 2012, 2014). The cadherin gene of *D. v. virgifera* is

approximately 13.3 fold larger than the *Tribolium castaneum* ortholog due to much

larger introns. The presence of numerous MITE-like elements within the cadherin gene

of *D. v. virgifera* indicates that the difference in the gene size is due to the insertion of

transposable elements in the *D. v. virgifera* introns (Coates et al., 2012). Class I BEL-

like long terminal repeat (LTR) retrotransposons have been also found in the *D. v.*

virgifera genome (Coates et al., 2014).

Although the genome size of *D. v. virgifera* has been reported, no genome size

data have been obtained for the other species in the genus *Diabrotica* and related

genera. Since the genome size of *D. v. virgifera* is relatively large, we hypothesized that

there has been a recent expansion in genome size in the lineage leading to it. We

tested this hypothesis by estimating and comparing the genome sizes of *D. v. virgifera*

with those of several *Diabrotica* species and an outgroup species, *A. vittatum.* As a high

proportion of repetitive elements were found in the cadherin gene of *D. v. virgifera,* we

further hypothesized that genome size expansion in the lineage leading to *D.v. virgifera*

was due to a general increase in repetitive elements*.* To test this hypothesis we looked

at the nature and quantity of repetitive elements in the virgifera group and compared it

with the fucata group.

<u>Materials and Methods</u>

Sample Collection

Specimens of *D. v. virgifera* and *A. vittatum* were collected from a maize field in Illinois

in 2017, while *D. barberi* were collected from Wisconsin by Tracy Schilder, Wisconsin

Department of Agriculture. Specimens of *D. v. zeae* were collected from Texas by

Thomas Sappington, US Department of Agriculture Agricultural Research Service, and

D. balteata were provided by Blair Siegfried and Heather McAuslane, University of

Florida, from a laboratory colony. *Diabrotica undecimpunctata* were obtained from Crop

Characteristics (Farmington, Minnesota, USA). Adult *Periplaneta americana,* which

were used as an external reference (Guo et al., 2015; He et al., 2016) for flow

cytometric measurement, were obtained from Carolina Biological Supply (Burlington,

North Carolina, USA). All samples were flash-frozen in liquid nitrogen and preserved at -

80°C.

Sample Preparation

Genome size estimates were generated for eight individuals from five species of

Diabrotica and one species of *Acalymma*. Preparations of nuclei were based on the

method of Hare and Johnston (2012). The heads of single individuals were

homogenized in 1ml of cold Galbraith buffer (4.26 g of $MgCl_2$, 8.84 g of sodium citrate,

4.2g of MOPS, 1ml of Triton-x, and 1mg of boiled ribonuclease A into 1 liter of ddH_20)

placed in a 7ml Kontes Dounce. The homogenate was filtered through a 20 µm nylon

mesh in a 1.5ml Eppendorf tube. Nuclei were stained with propidium iodide (PI) at 50

µg/ml, in the dark at 4°C for an hour. In addition to the test sample, the brain tissue of *P.*

americana was used as a standard (Hanrahan & Johnston, 2011). The brain tissue of *P.*

americana was dissected out, and the nuclear suspension was prepared and stained as

described above.

Flow Cytometric Analysis

Stained nuclei were analyzed using an Attune NxT Flow Cytometer (Thermo Fisher

Scientific, Waltham, Massachusetts). The propidium iodide-stained nuclei were excited

by exposing them to the 488 nm blue laser. Red fluorescence from the propidium iodide

was collected using a 610/20 emission filter in the YL2 detector channel. The calibration

of the flow cytometer was performed using a standard manufacturer's protocol before

use. The samples were inverted and mixed several times to resuspend and then loaded

onto the cytometer for analysis. During each sample run, the linearity of the

fluorescence measurement was confirmed by checking that the mean channel number

of the 4C nuclei (G2 phase) was double that of 2C nuclei (G1 phase). At least 1000

nuclear events were collected under each unknown and standard 2C peak. The nuclei

peak (PI fluorescence histogram) and coefficient of variation (CV) for each peak of

interest (sample and standard) were obtained using the gating function in the Attune

Software. The coefficient of variation (CV) was evaluated and confirmed to be less than

5 percent which is considered appropriate for accurate genome size estimates (Dolezel

et al., 2007; Tomaszewska et al., 2021). The known genome size of the standard

(3.41Gb, Harahan & Johnson, 2011) and the relative fluorescence obtained from the

sample and standard were then used to estimate the genome size using the following

formula:

Sample 1C DNA content = [(sample 2C mean peak position) / (standards 2C mean

peak position)] * standards 1C.

Genome size variations were analyzed using analysis of variance (ANOVA) and

Tukey's honest significant difference (HSD) post-hoc analyses using R statistical

software (version: 4.10) (*R Foundation for Statistical Computing, Vienna, Austria.*,

2020). Letters were assigned showing significance based on Tukey HSD post-hoc test

using R statistical software (version: 4.10) (R Core Team 2021).

Sequencing Data

Data from Illumina whole-genome shotgun sequencing were used to analyze the

repetitive DNA content of *D. barberi, D.undecimpunctata,* and *D. v. virgifera*. Adult *D.*

barberi (*n* = 71) and *D. undecimpunctata* (*n* = 50) were collected from maize fields near

Ames, Iowa, and Monmouth, Illinois, respectively. Each sample was pooled by species,

flash-frozen in liquid nitrogen and ground in liquid nitrogen, and then DNA extracted

from ~3.0 mg of tissue using the Qiagen DNeasy Blood and Tissue Extraction kit, with

modifications as described (Coates et al., 2014). Two micrograms of extracted DNA

was submitted to the Iowa State University DNA Facility (Ames, IA, USA) from which

~500bp insert indexed sequencing libraries were generated using the Illumina TruSeq

v2 Library Construction Kit (Illumina, San Diego, CA). Single-end 100-bp reads were

generated from *D. barberi* and *D. undecimpunctata* libraries in separate lanes of an

Illumina HiSeq2500. Raw reads were submitted to the Biotechnology Information

(NCBI) Short Read Archive (SRA) under accessions SRR13363759 and SRR13364002

for *D. barberi* and *D. undecimpunctata*, respectively.

Diabrotica virgifera virgifera adult females of inbred line Ped12-6-A-3 were flash-frozen

prior to genomic DNA isolation. Briefly, whole beetles were homogenized in an SDS-

based cell lysis solution followed by overnight incubation with Proteinase K at 55°C.

Cellular debris was pelleted and RNA was digested with RNaseA. The homogenate was

mixed with a high-salt solution and incubated overnight at 4°C. The DNA in the

supernatant was precipitated overnight with ethanol at -20°C. DNA was quantified on an

Invitrogen Qubit. A paired-end short-insert genomic DNA library was prepared at the

Roy J. Carver Biotechnology Center at the University of Illinois at Urbana-Champaign

using an Illumina TruSeq DNAseq Sample Prep kit. Reads were sequenced to 100bp

with the Illumina TruSeq SBS sequencing kit version 3 on an Illumina HiSeq 2000

instrument using Casava 1.8 for basecalling.

Sequencing generated 90 million single-end reads of 100bp for *D. barberi,* 118 million

single-end reads of 100bp for *D.undecimpunctata*, and 116 million 100-bp paired-end

reads for *D. v. virgifera*.

Annotation and Quantification of Repeat Content

Raw reads were quality-filtered using fastp software (version 0.20.1) with a minimum 20

average Phred score. Reads mapping to mitochondrial genome sequences of

Diabrotica species available through the NCBI website (KF658070.1, KF669870.1) were

identified (minimap2 v2.17) and filtered out as implemented in the SSRG workflow

(Pombert, 2021). Repetitive elements in the genomes of *D.undecimpunctata, D. barberi,*

and *D. v. virgifera* were assembled and quantified using dnaPipeTE v1.3 (Goubert et al.,

2015) and annotated using the DeepTE tool (Yan et al., 2020). To quantify the

proportion of TEs, dnaPipeTE uses samples of sequence reads instead of genome

assemblies, making this pipeline (dnaPipeTE) applicable for genomes with lower

sequencing depth. The pipeline performed assembly of repetitive reads into contigs

from low coverage sampling of raw reads using Trinity (Grabherr et al., 2011) and

annotated them using RepeatMasker (Smit et al., 1996-2004) with built-in Repbase

libraries (Bao et al., 2015, version 2017-01-27). Quantification was done by mapping a

random sample of reads onto the assembled repeats. The parameters set as the

benchmark for repeat content analysis for genomes greater than 500Mb (Goubert et al.,

2015), including the coverage parameter, were used to run dnaPipeTE. The pipeline

was run for all three species using 0.1x coverage. Additionally, 0.1x coverage was

chosen based on the high N50 metric and plateauing point of transposable elements i.e,

increasing the coverage beyond 0.1x only marginally increased the proportion of

transposable elements for all three species. The dnaPipeTE pipeline does not annotate

novel repeats that do not match an entry in the included Repbase library. A high

proportion of repeats from each of the three beetle species were not annotated by

dnaPipeTE. DeepTE, a deep learning method based on convolutional neural networks,

was used to classify and annotate the unknown TEs. DeepTE uses eight trained models

to classify TEs into superfamilies and orders. All the TE contigs assembled by

dnaPipeTE were analyzed using DeepTE, whether or not they had been previously

classified by dnaPipeTE. Combining the results of the assembly and quantification by

dnaPipeTE with the classification results from DeepTE allowed the abundance of repeat

families in the genomes of all three species to be determined.

For comparison with the dnaPipeTE *de-novo* assembly of repetitive elements, the

percentage of repetitive elements in the *D. v. virgifera* genome assembly (NCBI RefSeq

accession GCF_003013835.1) was analyzed with RepeatModeler version 2.0.1 (Flynn

et al., 2019).   Repeatmodeler is a *de-novo* transposable element identification package

that uses three repeat finding programs (RECON, RepeatScout, and

LtrHarvest/Ltr_retriever) to discover repetitive DNA sequences in the genome. These

repetitive DNA sequences were annotated by repeatClassifier based on the similarity to

RepBase and Dfam databases. The annotated library produced was used as input to

RepeatMasker to detect and mask repeats in the genome. Default parameters were

used to run RepeatModeler.

<u>Results</u>

Genome size from flow cytometry:

The genome sizes of *D. v. zeae*, *D. v. virgifera,* and *D. barberi* from the virgifera group

were estimated to be 2.65Gb ± 0.01, 2.64 Gb ± 0.02, and 2.31Gb ± 0.04, respectively.

The genomes of *D. balteata* and *D.undecimpunctata* from the fucata group were

estimated at 1.68Gb ± 0.01 and 1.59 Gb ± 0.00. The outgroup species *A. vittatum*

genome size is estimated to be 1.69 Gb ± 0.01. An analysis of variance (ANOVA)

showed a significant difference (p < 2×10^{-16}) in the genome sizes of the species

under study. A subsequent Tukey HSD test showed that there were no significant

differences in genome size between *D. v. virgifera* and *D. v. zeae*, between *D. balteata*

and *D.undecimpunctata,* or between *D. balteata* and *A. vittatum*. The estimated genome

size for each species with their phylogenetic relationships is shown in Figure I.

Figure I: Genome size evolution within the genus *Diabrotica.* Phylogeny of *Diabrotica* and outgroup

Acalymma vittatum is based on Eben & Espinosa de los Monteros, 2013. Letters a - d indicate groups

with no significant difference in mean 1C-value (Tukey HSD; α = 0.05).

Repeat Content analysis:

The repeatomes of *D.v.virgifera, D. barberi,* and *D.undecimpunctata* comprised 72.5%,

70.4%, and 52.7% of their genomes, respectively. The repeat content obtained via

Repeat Modeler for the draft *D. v. virgifera* genome was 57.4%. As the assembly of the

D.v. virgifera genome was based on short-reads, TEs were expected to be under-

represented because of the difficulty of assembling individual copies. TE-rich large

genomes are difficult to assemble and often end up with high levels of fragmentation

around repetitive regions leading to underestimation of TE content (Green, 2002).

To further investigate the classes of repeat families that contributed to the genome size

variation between the two groups of *Diabrotica*, the DeepTE annotations were coupled

with the dnaPipeTE abundance quantification to estimate the abundance of different

repeat elements in the genomes of *D. v. virgifera, D.barberi,* and *D.undecimpunctata*.

The TEs that accounted for most of the difference in the genome size of the two groups

were annotated as class II DNA Tc1-mariner Miniature-repeat Transposable Elements

(MITEs) (ClassII_DNA_TcMar_MITE) and class II DNA hAT MITE

(ClassII_DNA_hAT_MITE) of transposable elements and class I long terminal repeat

(LTR) Gypsy (ClassI_LTR_Gypsy) transposable elements (Figure II). The *D. v. virgifera*

genome contained a large amount of ClassII_DNA_TcMar_MITE(0.62 Gb),

ClassII_DNA_hAT_MITE (0.24 Gb), and ClassI_LTR_Gypsy(0.31 Gb). Similarly, *D.*

barberi also had a high amount of ClassII_DNA_TcMar_MITE(0.53 Gb),

ClassII_DNA_hAT_MITE (0.19 Gb), and ClassI_LTR_Gypsy(0.26 Gb). The *D.*

undecimpunctata had a lower amount of ClassII_DNA_TcMar_MITE(0.16 Gb),

ClassII_DNA_hAT_MITE (0.06 Gb), and ClassI_LTR_Gypsy(0.17 Gb). Transposable

elements from other repeat families such as nLTRS, helitrons, and others from class I

and class II DNA elements were not as prominent as those mentioned above. Single-

low copy sequences representing the non-repetitive portion of the genomes of *D. v.*

virgifera, *D.barberi*, and *D.undecimpunctata* totaled 0.73 Gb, 0.68 Gb, and 0.75 Gb,

respectively.

Figure II: Quantities of transposable elements present in the genomes of three species of *Diabrotica.* The

boxed transposable elements are those with high contributions to the genome size variation between the

groups of fucata and virgifera.

Discussion:

Diabrotica is the most diverse genus within the subtribe Diabroticina and includes 354

species native to America. Only species from the fucata and virgifera groups occur in

the United States (Krysan, 1986), while species from the signifera group are endemic to

South America. The economically significant pest species in this genus either belong to

the virgifera group or to the fucata group, justifying the need to study them

comprehensively. The signifera group species are not of economic importance, and

their biology is also mostly unknown (Clark et al., 2001) and so the group is

understudied. The expansion and adaptability of the pest species generates a sense of

urgency to study them.

Our results demonstrated that the genome size of virgifera group species are

approximately 1Gb larger than that of the fucata group and *Acalymma* species. The

genome size for *D. v. virgifera* obtained in this study and in a previous study (Coates et

al., 2012) are consistent. Our genome size results, when coupled with the species'

phylogenetic relationship, indicated that an expansion in genome size occurred in the

common ancestor of the virgifera group leading to *D. barberi, D. v. virgifera,* and *D. v.*

zeae. There was also a significant difference in the genome sizes of *D. barberi* and *D.*

virgifera subsp. suggesting a possible further expansion of the genome size in the

common ancestor of *D. v. virgifera* and *D. v. zeae*.

The estimated divergence time between *Acalymma* and *Diabrotica* species and

between the *Diabrotica* species group was dated at 45 million years ago (Mya) and 31

Mya respectively. Another radiation within the Diabrotica genus took place around 17

Mya resulting in virgifera group (Eben & Espinosa de los Monteros, 2013). Repeat

elements of species from fucata and virgifera were studied to understand the basis of

genome size expansion in *Diabrotica.* The genomes of the virgifera and fucata group of

Diabrotica species contained many common TEs, but the abundance of three TE

families differed substantially and accounted for much of the difference in genome size

between the two groups and also within the virgifera group between *D.barberi* and *D.*

virgifera subsp. The MITE-like Tc1-mariner and hAT elements and LTR Gypsy

retroelements were more abundant in the virgifera group. Miniature-repeat

Transposable Elements (MITEs) are short AT-rich (<0.5kb) derivatives of DNA elements

whose internal sequence lacks an open reading frame (Lu et al., 2012), contain

conserved terminal inverted repeats, are flanked by target site duplications, and are

closely associated with euchromatic genes (Kuang et al., 2009). Gypsy elements are

one of the most abundant classes of the long terminal repeat (LTR)-retrotransposons

superfamily with large numbers of copies found in almost all the plants, animals, and

fungi tested (Thomas-Bulle et al., 2018). Generally, class I transposable elements have

been reported to be in high abundance in insect genomes, such as *Tribolium*

castaneum (Wang et al., 2008), *Drosophila* (Clark et al., 2007)*, Bombyx mori* (Osanai-

Futahashi et al., 2008) in comparison to class II elements. However, in *Diabrotica*, we

discovered an abundance of class II elements and, to some extent, class I elements.

Genome size varies enormously among eukaryote species (Hidalgo et al., 2017), and

among beetles in the Family Chrysomelidae (Hanrahan & Johnston, 2011; Petitpierre et

al., 1993). There are studies showing a positive correlation between genome size and

phenotypic traits such as body size in groups of invertebrates, including aphids (Finston

et al., 1995) and flies (Ferrari & Rai, 1989). Equally, there are studies showing negative

correlations between phenotypic traits and genome sizes, such as the relation between

body size and genome size for the *Pimelia* and *Phylan* genera in the beetle family

Tenebrionidae (Palmer et al., 2003; Palmer & Petitpierre, 1996). Cases of negative

correlation of genome size with pupal development have been reported in ladybird

beetles and with the overall development rate in *Tribolium* (Carreras et al., 1991;

Gregory et al., 2003). It has been previously hypothesized that genome size negatively

correlates with the host range in insect herbivores (Zhang et al., 2019). However, there

are also studies showing that genome size may be positively correlated with the host

range in insect herbivores (Matsubayashi & Ohshima, 2015; Calatayud et al., 2016).

This contradiction in results, along with our data suggests that the genome size may not

be related directly to the host range.

Comparative genomic studies in insects have revealed that repeat elements can make

large contributions to genome size variation. Variation in TE abundance can be seen

both within and among species (Lynch, 2007). Honeybees, with a genome size of

230Mb, show very few repeat elements, representing a case of TE extinction

(Weinstock et al., 2006). Similarly, the small genome of *Belgica antarctica,* the Antarctic

midge, (99 Mb) is also due to the reduction of repeats in the genome (Kelley et al.,

2014). There are also cases of TE proliferations, such as in *Locusta migratoria,* that led

to its large genome size of 6.5 Gb (Wang et al., 2014). Cases of increase in genome

size as a consequence of transposable elements have also been reported in wood

white (Leptidea) butterflies and North American fireflies (Lampyridae) (Talla et al., 2017;

Lower et al., 2017). This further suggests that genome size in insects is fairly plastic and

largely driven by the loss and gain of transposable elements.

Host genomes have employed several mechanisms to suppress transposable

element expression and mobility (Bourque et al., 2018) which includes epigenetic

silencing either through histone modifications or DNA methylation, targeted

mutagenesis, small RNA interference, as well as sequence-specific repressors such as

the recently profiled KRAB zinc-finger proteins (Fouché et al., 2020; Maupetit-Mehouas

& Vaury, 2020). At the same time, some transposable elements have evolved regulatory

sequences controlling their own copy number to autonomously replicate in the genome

(Lohe & Hartl, 1996; Saha et al., 2015). TE de-repression is triggered by environmental

stimuli, in particular stress (Bundo et al., 2014; Fouché et al., 2020; Voronova et al.,

2014) impacting transcription levels and increasing transpositional activity (Dubin et al.,

2018). Additionally, there are other factors influencing TE mobilization for example

demethylation and the removal of repressive histone marks during epigenetic

reprogramming stages (Russell & LaMarre, 2018)

Transposable elements are a major component of many eukaryotic genomes and

have been associated with several mechanisms shaping eukaryotic genome function,

structure, and evolution (Charlesworth & Charlesworth, 1983; Chuong et al., 2017).

These transposons are capable of changing position within the genome, leading to

mutations by inserting themselves into functional regions and causing change by either

modifying or eliminating gene expression (Feschotte, 2008; Oliver & Greene, 2009). TE

integration and excision can introduce novel variation at the individual and population-

scale (McClintock, 1950; Wendel & Wessler, 2000) introducing phenotypic modifications

through changes in epigenetic mechanisms and providing genetic motifs (Jiang et al.,

2004; Kapitonov & Jurka, 2007; Stuart et al., 2016; Wessler et al., 1995). They may also

lead to genomic rearrangement (Maumus et al., 2015; Mat Razali et al., 2019). In

particular, MITES and LTR retrotransposon have played an essential role in shaping

gene structure and altering gene function (Wessler et al., 1995).

Initially, MITEs were found as key components of plant genomes, as they are

frequently associated with genes with high copy numbers indicating a possible role in

gene expression and genome evolution (Santiago et al., 2002; Oki et al., 2008). They

are also found in animals including mosquitoes, *Drosophila*, fish, and humans (Deprá et

al., 2012). Similar to MITES, LTR retrotransposons were also first discovered in plants.

They are usually located largely in intergenic regions and are often the single largest

component of plant genomes (Feschotte et al., 2002; Kumar & Bennetzen, 1999).

Previous studies have also revealed that both MITES and LTR retrotransposons can

regulate downstream gene expression by inserting into promoter regions (Wang et al.,

2017; Butelli et al., 2012; Liu et al., 2019).

Mostly, TE insertions are presumed to be deleterious or neutral, but some have

been shown to be selectively advantageous. There are several studies showing that TE-

mediated insertions have led to insecticide resistance. In pink bollworm *Pectinophora*

gossypiella, a major pest of cotton (Rostant et al., 2012), several independent TE

insertions in the *PgCad1* gene conferred resistance to *Bt* Cry1Ac toxin (Fabrick et al.,

2011; Wang et al., 2019). Cases of resistance to *Bacillus thuringiensis* (Bt) toxins have

also been reported in *Heliothis virescens* which is caused by disruption of a cadherin-

superfamily gene by TE insertion (Gahan, 2001). TE insertions in xenobiotic metabolism

related genes such as those encoding cytochrome P450 monooxygenases and

glutathione S-transferases in *Helicoverpa armigera* are the cause of resistance to

insecticides (Klai et al., 2020). Another example demonstrating that TEs can produce

adaptive mutations has been reported in *D. melanogaster* (Rostant et al., 2012; Gilbert

et al., 2021). An increased resistance to dichlorodiphenyltrichloroethane (DDT) in *D.*

melanogaster has been reported due to Cyp6g1 upregulation caused by insertion of the

Accord transposon in the 5′ regulatory region of the Cyp6g1 gene (Chung et al., 2007).

Similarly, TE insertion, which truncates the CHKov1 gene in *D. melanogaster*, confers

resistance towards organophosphate (Aminetzach, 2005).

The role played by TEs, if any, in Diabroticina evolution, both long term such as

speciation and host plant shifts, and in recent evolution, such as their adaptation to pest

management, is unknown. However, the ongoing effort of the United States Department

of Agriculture Agricultural Research Service (USDA-ARS) Ag100Pest initiative to

sequence other species of *Diabrotica* may help us understand and answer these

questions.

Acknowledgments:

We would like to thank Dr. Thomas Sappington, Dr. Blair Siegfried, Dr. Heather

McAuslane, and Tracy Schilder for providing insect samples. We also thank Dr. J.

Spencer Johnston and David Leclerc for their advice and guidance on flow cytometry

methods and Dr. Clément Goubert's advice on dnaPipeTE. This work was supported

and funded by USDA NIFA award number 2011-67009-30134 and by the USDA

Agricultural Research Service (ARS) (CRIS Project 5030-22000-019-00D). This article

reports the results of research only, and mention of any commercial product or service

does not constitute a recommendation for its use by USDA. The findings and

conclusions in this publication are those of the authors and should not be construed to

represent any official USDA or United States government determination of policy.

USDA is an equal opportunity employer and provider.

References:

Alouw, J. C., & Miller, N. J. (2015). Effects of benzoxazinoids on specialist and

generalist *Diabrotica* species. *Journal of Applied Entomology*, *139*(6), 424–431.

https://doi.org/10.1111/jen.12194

Aminetzach, Y. T. (2005). Pesticide resistance via transposition-mediated adaptive

gene truncation in *Drosophila*. *Science*, *309*(5735), 764–767.

https://doi.org/10.1126/science.1112699

Ball, H. J., & Weekman, G. T. (1962). Insecticide resistance in the adult western corn

rootworm in Nebraska. *Journal of Economic Entomology*, *55*(4), 439–441.

https://doi.org/10.1093/jee/55.4.439

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive

elements in eukaryotic genomes. *Mobile DNA*, *6*(1), 11.

https://doi.org/10.1186/s13100-015-0041-9

Bourque, G., Burns, K. H., Gehring, M., Gorbunova, V., Seluanov, A., Hammell, M.,

Imbeault, M., Izsvák, Z., Levin, H. L., Macfarlan, T. S., Mager, D. L., & Feschotte,

C. (2018). Ten things you should know about transposable elements. *Genome*

Biology, *19*, 199. https://doi.org/10.1186/s13059-018-1577-z

Bragard, C., Dehnen-Schmutz, K., Di Serio, F., Gonthier, P., Jacques, M., Jaques Miret,

J. A., Justesen, A. F., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A.,

Parnell, S., Potting, R., Reignault, P. L., Thulke, H., Van der Werf, W., Vicent

Civera, A., Yuen, J., Zappalà, L., Czwienczek, E., & MacLeod, A. (2019). Pest

categorisation of *Diabrotica virgifera zeae* . *EFSA Journal*, *17*(11).

https://doi.org/10.2903/j.efsa.2019.5858

Branson, T. F., & Krysan, J. L. (1981). Feeding and oviposition behavior and life cycle

strategies of *Diabrotica* : An evolutionary view with implications for pest

management. *Environmental Entomology*, *10*(6), 826–831.

https://doi.org/10.1093/ee/10.6.826

Bundo, M., Toyoshima, M., Okada, Y., Akamatsu, W., Ueda, J., Nemoto-Miyauchi, T.,

Sunaga, F., Toritsuka, M., Ikawa, D., Kakita, A., Kato, M., Kasai, K., Kishimoto,

T., Nawa, H., Okano, H., Yoshikawa, T., Kato, T., & Iwamoto, K. (2014).

Increased L1 retrotransposition in the neuronal genome in schizophrenia.

Neuron, *81*(2), 306–313. https://doi.org/10.1016/j.neuron.2013.10.053

Butelli, E., Licciardello, C., Zhang, Y., Liu, J., Mackay, S., Bailey, P., Reforgiato-

Recupero, G., & Martin, C. (2012). Retrotransposons control fruit-specific, cold-

dependent accumulation of anthocyanins in blood oranges. *The Plant Cell*, *24*(3),

1242–1255. https://doi.org/10.1105/tpc.111.095232

Calatayud, P.-A., Petit, C., Burlet, N., Dupas, S., Glaser, N., Capdevielle-Dulac, C., Le

Ru, B., Jacquin-Joly, E., Kaiser-Arnauld, L., Harry, M., & Vieira, C. (2016). Is

genome size of Lepidoptera linked to host plant range? *Entomologia*

Experimentalis et Applicata, *159*(3), 354–361. https://doi.org/10.1111/eea.12446

Calles-Torrez, V., Knodel, J. J., Boetel, M. A., French, B. W., Fuller, B. W., & Ransom,

J. K. (2019). Field-evolved resistance of northern and western corn rootworm

(Coleoptera: Chrysomelidae) populations to corn hybrids expressing single and

pyramided Cry3Bb1 and Cry34/35Ab1 Bt proteins in North Dakota. *Journal of*

Economic Entomology, *112*(4), 1875–1886. https://doi.org/10.1093/jee/toz111

Capinera, J. L. (2008). Northern corn rootworm, *Diabrotica barberi* Smith & Lawrence

(Coleoptera: Chrysomelidae). In J. L. Capinera (Ed.), *Encyclopedia of*

Entomology (pp. 2616–2619). Springer Netherlands. https://doi.org/10.1007/978-

1-4020-6359-6_2242

Carreras, I., A., A.-F., Juan, C., & Petitpierre, E. (1991). Tasa de desarrollo en *Tribolium*

brevicornis (Coleoptera: Tenebrionidae) y su relación con el tamaño del genoma.

Charlesworth, B., & Charlesworth, D. (1983). The population dynamics of transposable

elements. *Genetics Research*, *42*(1), 1–27.

https://doi.org/10.1017/S0016672300021455

Chung, H., Bogwitz, M. R., McCart, C., Andrianopoulos, A., Ffrench-Constant, R. H.,

Batterham, P., & Daborn, P. J. (2007). *Cis*-regulatory elements in the *Accord*

retrotransposon result in tissue-specific expression of the *Drosophila*

melanogaster insecticide resistance gene *Cyp6g1*. *Genetics*, *175*(3), 1071–1077.

https://doi.org/10.1534/genetics.106.066597

Chuong, E. B., Elde, N. C., & Feschotte, C. (2017). Regulatory activities of transposable

elements: From conflicts to benefits. *Nature Reviews. Genetics*, *18*(2), 71–86.

https://doi.org/10.1038/nrg.2016.139

Ciosi, M., Miller, N. J., Kim, K. S., Giordano, R., Estoup, A., & Guillemaud, T. (2008).

Invasion of Europe by the western corn rootworm, *Diabrotica virgifera virgifera*:

Multiple transatlantic introductions with various reductions of genetic diversity.

Molecular Ecology, *17*(16), 3614--3627. https://doi.org/10.1111/j.1365-

294X.2008.03866.x

Ciosi, M., Miller, N. J., Toepfer, S., Estoup, A., & Guillemaud, T. (2011). Stratified

dispersal and increasing genetic variation during the invasion of Central Europe

by the western corn rootworm, *Diabrotica virgifera virgifera*. *Evolutionary*

Applications, *4*(1), 54–70. https://doi.org/10.1111/j.1752-4571.2010.00133.x

Clark, A. G., Eisen, M. B., Smith, D. R., Bergman, C. M., Oliver, B., Markow, T. A.,

Kaufman, T. C., Kellis, M., Gelbart, W., Iyer, V. N., Pollard, D. A., Sackton, T. B.,

Larracuente, A. M., Singh, N. D., Abad, J. P., Abt, D. N., Adryan, B., Aguade, M.,

Akashi, H., … *Broad Institute Genome Sequencing Platform. (2007). Evolution

of genes and genomes on the *Drosophila* phylogeny. *Nature*, *450*(7167), 203–

218. https://doi.org/10.1038/nature06341

Clark, T. L., Meinke, L. J., & Foster, J. E. (2001). Molecular phylogeny of *Diabrotica*

beetles (Coleoptera: Chrysomelidae) inferred from analysis of combined

mitochondrial and nuclear DNA sequences. *Insect Molecular Biology*, *10*(4),

303–314. https://doi.org/10.1046/j.0962-1075.2001.00269.x

Coates, B. S., Alves, A. P., Wang, H., Walden, K. K. O., French, B. W., Miller, N. J.,

Abel, C. A., Robertson, H. M., Sappington, T. W., & Siegfried, B. D. (2012).

Distribution of genes and repetitive elements in the *Diabrotica virgifera virgifera*

genome estimated using BAC sequencing. *Journal of Biomedicine and*

Biotechnology, *2012*, 1–9. https://doi.org/10.1155/2012/604076

Coates, B. S., Fraser, L. M., French, B. W., & Sappington, T. W. (2014). Proliferation

and copy number variation of BEL-like long terminal repeat retrotransposons

within the *Diabrotica virgifera virgifera* genome. *Gene*, *534*(2), 362–370.

Deprá, M., Ludwig, A., Valente, V. L., & Loreto, E. L. (2012). Mar, a MITE family of hAT

transposons in *Drosophila. Mobile DNA*, *3*(1), 13. https://doi.org/10.1186/1759-

8753-3-13

Devos, Y., Meihls, L. N., Kiss, J., & Hibbard, B. E. (2013). Resistance evolution to the

first generation of genetically modified *Diabrotica*-active *Bt*-maize events by

western corn rootworm: Management and monitoring considerations. *Transgenic*

Research, *22*(2), 269–299. https://doi.org/10.1007/s11248-012-9657-4

Dolezel, J., Greilhuber, J., & Suda, J. (2007). *Flow Cytometry with Plant Cells: Analysis*

of Genes, Chromosomes and Genomes (p. 454).

https://doi.org/10.1002/9783527610921

Dubin, M. J., Mittelsten Scheid, O., & Becker, C. (2018). Transposons: A blessing curse.

Current Opinion in Plant Biology, 42, 23–29.

https://doi.org/10.1016/j.pbi.2018.01.003

Eben, A., & Espinosa de los Monteros, A. (2013). Tempo and mode of evolutionary

radiation in Diabroticina beetles (genera Acalymma, Cerotoma, and Diabrotica).

ZooKeys, 332, 207–321. https://doi.org/10.3897/zookeys.332.5220

Fabrick, J. A., Mathew, L. G., Tabashnik, B. E., & Li, X. (2011). Insertion of an intact

CR1 retrotransposon in a cadherin gene linked with Bt resistance in the pink

bollworm, Pectinophora gossypiella. Insect Molecular Biology, 20(5), 651–665.

https://doi.org/10.1111/j.1365-2583.2011.01095.x

Ferrari, J. A., & Rai, K. S. (1989). Phenotypic correlates of genome size variation in

Aedes albopictus. Evolution; International Journal of Organic Evolution, 43(4),

895–899. https://doi.org/10.1111/j.1558-5646.1989.tb05187.x

Feschotte, C. (2008). Transposable elements and the evolution of regulatory networks.

Nature Reviews. Genetics, 9(5), 397–405. https://doi.org/10.1038/nrg2337

Feschotte, C., Jiang, N., & Wessler, S. R. (2002). Plant transposable elements: Where

genetics meets genomics. Nature Reviews Genetics, 3(5), 329–341.

https://doi.org/10.1038/nrg793

Finnegan, D. J. (1989). Eukaryotic transposable elements and genome evolution.

Trends in Genetics: TIG, 5(4), 103–107. https://doi.org/10.1016/0168-

9525(89)90039-5

Finston, T. L., Hebert, P. D. N., & Foottit, R. B. (1995). Genome size variation in aphids.

Insect Biochemistry and Molecular Biology, *25*(2), 189–196.

https://doi.org/10.1016/0965-1748(94)00050-R

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A.

F. (2019). *RepeatModeler2: Automated genomic discovery of transposable*

element families [Preprint]. Genomics. https://doi.org/10.1101/856591

Fouché, S., Badet, T., Oggenfuss, U., Plissonneau, C., Francisco, C. S., & Croll, D.

(2020). Stress-Driven Transposable Element De-repression Dynamics and

Virulence Evolution in a Fungal Pathogen. *Molecular Biology and Evolution*,

37(1), 221–239. https://doi.org/10.1093/molbev/msz216

Gahan, L. J. (2001). Identification of a gene associated with Bt resistance in *Heliothis*

virescens. *Science*, *293*(5531), 857–860.

https://doi.org/10.1126/science.1060949

Gassmann, A. J., Petzold-Maxwell, J. L., Clifton, E. H., Dunbar, M. W., Hoffmann, A. M.,

Ingber, D. A., & Keweshan, R. S. (2014). Field-evolved resistance by western

corn rootworm to multiple *Bacillus thuringiensis* toxins in transgenic maize.

Proceedings of the National Academy of Sciences, *111*(14), 5141–5146.

https://doi.org/10.1073/pnas.1317179111

Gilbert, C., Peccoud, J., & Cordaux, R. (2021). Transposable elements and the

evolution of insects. *Annual Review of Entomology*, *66*(1), 355–372.

https://doi.org/10.1146/annurev-ento-070720-074650

Goubert, C., Modolo, L., Vieira, C., ValienteMoro, C., Mavingui, P., & Boulesteix, M.

(2015). De novo assembly and annotation of the Asian tiger mosquito (*Aedes*

albopictus) repeatome with dnaPipeTE from raw genomic reads and comparative

analysis with the yellow fever mosquito (*Aedes aegypti*). *Genome Biology and*

Evolution, *7*(4), 1192–1205. https://doi.org/10.1093/gbe/evv050

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I.,

Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E.,

Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C.,

Lindblad-Toh, K., … Regev, A. (2011). Full-length transcriptome assembly from

RNA-seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–

652. https://doi.org/10.1038/nbt.1883

Gray, M. E., Sappington, T. W., Miller, N. J., Moeser, J., & Bohn, M. O. (2009).

Adaptation and invasiveness of western corn rootworm: Intensifying research on

a worsening pest. *Annual Review of Entomology*, *54*(1), 303–321.

https://doi.org/10.1146/annurev.ento.54.110807.090434

Green, P. (2002). Whole-genome disassembly. *Proceedings of the National Academy of*

Sciences of the United States of America, *99*(7), 4143–4144.

https://doi.org/10.1073/pnas.082095999

Gregory, T. R. (2021). *Animal genome size database*. http://www.genomesize.com/

Gregory, T. R., Nedvěd, O., & Adamowicz, S. (2003). C-value estimates for 31 species

of ladybird beetles (Coleoptera: Coccinellidae). *Hereditas*, *139*, 121–127.

https://doi.org/10.1111/j.1601-5223.2003.01766.x

Guo, L. T., Wang, S. L., Wu, Q. J., Zhou, X. G., Xie, W., & Zhang, Y. J. (2015). Flow

cytometry and K-mer analysis estimates of the genome sizes of *Bemisia tabaci*

B and Q (Hemiptera: Aleyrodidae). *Frontiers in Physiology*, *6*.

https://doi.org/10.3389/fphys.2015.00144

Hanrahan, S. J., & Johnston, J. S. (2011). New genome size estimates of 134 species

of arthropods. *Chromosome Research*, *19*(6), 809–823.

https://doi.org/10.1007/s10577-011-9231-6

Hare, E. E., & Johnston, J. S. (2012). Genome size determination using flow cytometry

of propidium iodide-stained nuclei. In V. Orgogozo & M. V. Rockman (Eds.),

Molecular Methods for Evolutionary Genetics (Vol. 772, pp. 3–12). Humana

Press. https://doi.org/10.1007/978-1-61779-228-1_1

He, K., Lin, K., Wang, G., & Li, F. (2016). Genome sizes of nine insect species

determined by flow cytometry and k-mer analysis. *Frontiers in Physiology*, *7*.

https://doi.org/10.3389/fphys.2016.00569

Hidalgo, O., Pellicer, J., Christenhusz, M., Schneider, H., Leitch, A., & Leitch, I. (2017).

Is there an upper limit to genome size? *Trends in Plant Science*, *22*.

https://doi.org/10.1016/j.tplants.2017.04.005

Jackson, D. M., Sorensen, K. A., Sorenson, C. E., & Story, R. N. (2005). Monitoring

cucumber beetles in sweetpotato and cucurbits with kairomone-baited traps.

Journal of Economic Entomology, *98*(1), 12.

Jiang, N., Bao, Z., Zhang, X., Eddy, S. R., & Wessler, S. R. (2004). Pack-MULE

transposable elements mediate gene evolution in plants. *Nature*, *431*(7008),

569–573. https://doi.org/10.1038/nature02953

Kapitonov, V. V., & Jurka, J. (2007). Helitrons on a roll: Eukaryotic rolling-circle

transposons. *Trends in Genetics: TIG*, *23*(10), 521–529.

https://doi.org/10.1016/j.tig.2007.08.004

Kelley, J. L., Peyton, J. T., Fiston-Lavier, A.-S., Teets, N. M., Yee, M.-C., Johnston, J.

S., Bustamante, C. D., Lee, R. E., & Denlinger, D. L. (2014). Compact genome of

the Antarctic midge is likely an adaptation to an extreme environment. *Nature*

Communications, *5*(1), 4611. https://doi.org/10.1038/ncomms5611

Khajuria, C., Ivashuta, S., Wiggins, E., Flagel, L., Moar, W., Pleau, M., Miller, K., Zhang,

Y., Ramaseshadri, P., Jiang, C., Hodge, T., Jensen, P., Chen, M., Gowda, A.,

McNulty, B., Vazquez, C., Bolognesi, R., Haas, J., Head, G., & Clark, T. (2018).

Development and characterization of the first dsRNA-resistant insect population

from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *PLOS ONE*,

13(5), e0197059. https://doi.org/10.1371/journal.pone.0197059

Kidwell, M. G. (2002). Transposable elements and the evolution of genome size in

eukaryotes. *Genetica*, *115*(1), 49–63. https://doi.org/10.1023/a:1016072014259

Klai, K., Chénais, B., Zidi, M., Djebbi, S., Caruso, A., Denis, F., Confais, J., Badawi, M.,

Casse, N., & Mezghani Khemakhem, M. (2020). Screening of *Helicoverpa*

armigera mobilome revealed transposable element insertions in insecticide

resistance genes. *Insects*, *11*(12), 879. https://doi.org/10.3390/insects11120879

Koch, R. L., Burkness, E. C., & Hutchison, W. D. (2004). Confirmation of bean leaf

beetle, *Cerotoma trifurcata*, feeding on cucurbits. *Journal of Insect Science*, *4*(1).

https://doi.org/10.1093/jis/4.1.5

Kojima, K. K. (2019). Structural and sequence diversity of eukaryotic transposable

elements. *Genes & Genetic Systems*, *94*(6), 233–252.

https://doi.org/10.1266/ggs.18-00024

Krysan, J. L. (1982). Diapause in the nearctic species of the *virgifera* group of

Diabrotica: Evidence for tropical origin and temperate adaptations. *Annals of the*

Entomological Society of America, *75*(2), 136–142.

https://doi.org/10.1093/aesa/75.2.136

Krysan, J. L. (1986). Introduction: Biology, distribution, and identification of pest

Diabrotica. In J. L. Krysan & T. A. Miller (Eds.), *Methods for the Study of Pest*

Diabrotica (pp. 1–23). Springer. https://doi.org/10.1007/978-1-4612-4868-2_1

Krysan, J. L., Jackson, J. J., & Lew, A. C. (1984). Field Termination of Egg Diapause in

Diabrotica with New Evidence of Extended Diapause in *D. barberi* (Coleoptera:

Chrysomelidae). *Environmental Entomology*, *13*(5), 1237–1240.

https://doi.org/10.1093/ee/13.5.1237

Kuang, H., Padmanabhan, C., Li, F., Kamei, A., Bhaskar, P. B., Ouyang, S., Jiang, J.,

Buell, C. R., & Baker, B. (2009). Identification of miniature inverted-repeat

transposable elements (MITEs) and biogenesis of their siRNAs in the

Solanaceae: New functional implications for MITEs. *Genome Research*, *19*(1),

42–56. https://doi.org/10.1101/gr.078196.108

Kumar, A., & Bennetzen, J. L. (1999). Plant retrotransposons. *Annual Review of*

Genetics, *33*, 479–532. https://doi.org/10.1146/annurev.genet.33.1.479

Levine, E., & Oloumi-Sadeghi, H. (1991). Management of Diabroticite Rootworms in

Corn. Annual Review of Entomology, 36(1), 299-255.

https://doi.org/10.1146/annurev.en.36.010191.001305

Lewis, P. A., Lampman, R. L., & Metcalf, R. L. (1990). Kairomonal attractants for

Acalymma vittatum (Coleoptera: Chrysomelidae). *Environmental Entomology*,

19(1), 8–14. https://doi.org/10.1093/ee/19.1.8

Liu, Y., Tahir ul Qamar, M., Feng, J.-W., Ding, Y., Wang, S., Wu, G., Ke, L., Xu, Q., &

Chen, L.-L. (2019). Comparative analysis of miniature inverted–repeat

transposable elements (MITEs) and long terminal repeat (LTR) retrotransposons

in six *Citrus* species. *BMC Plant Biology*, *19*(1), 140.

https://doi.org/10.1186/s12870-019-1757-3

Lohe, A. R., & Hartl, D. L. (1996). Autoregulation of mariner transposase activity by

overproduction and dominant-negative complementation. *Molecular Biology and*

Evolution, *13*(4), 549–555.

https://doi.org/10.1093/oxfordjournals.molbev.a025615

Lower, S. S., Johnston, J. S., Stanger-Hall, K. F., Hjelmen, C. E., Hanrahan, S. J.,

Korunes, K., & Hall, D. (2017). Genome size in North American fireflies:

Substantial variation likely driven by neutral processes. *Genome Biology and*

Evolution, *9*(6), 1499–1512. https://doi.org/10.1093/gbe/evx097

Lu, C., Chen, J., Zhang, Y., Hu, Q., Su, W., & Kuang, H. (2012). Miniature inverted-

repeat transposable elements (MITEs) have been accumulated through

amplification bursts and play important roles in gene expression and species

diversity in *Oryza sativa*. *Molecular Biology and Evolution*, *29*(3), 1005–1017.

https://doi.org/10.1093/molbev/msr282

Lynch, M. (2007). *The origins of genome architecture*. Sinauer.

Mat Razali, N., Cheah, B. H., & Nadarajah, K. (2019). Transposable elements adaptive

role in genome plasticity, pathogenicity and evolution in fungal phytopathogens.

International Journal of Molecular Sciences, *20*(14).

https://doi.org/10.3390/ijms20143597

714 Matsubayashi, K. W., & Ohshima, I. (2015). Genome size increase in the phytophagous

715   ladybird beetle *Henosepilachna vigintioctomaculata* species complex

716   (Coleoptera: Coccinellidae): genome size increase in ladybird beetle.

717   *Entomological Science*, *18*(1), 134–137. https://doi.org/10.1111/ens.12090

718 Maumus, F., Fiston-Lavier, A.-S., & Quesneville, H. (2015). Impact of transposable

719   elements on insect genomes and biology. *Current Opinion in Insect Science*, 7,

720   30–36. https://doi.org/10.1016/j.cois.2015.01.001

721 Maupetit-Mehouas, S., & Vaury, C. (2020). Transposon Reactivation in the Germline

722   May Be Useful for Both Transposons and Their Host Genomes. *Cells*, *9*(5), 1172.

723   https://doi.org/10.3390/cells9051172

724 McClintock, B. (1950). The origin and behavior of mutable loci in maize. *Proceedings of*

725   *the National Academy of Sciences of the United States of America*, *36*(6), 344–

726   355. https://doi.org/10.1073/pnas.36.6.344

727 Meinke, L. J., Siegfried, B. D., Wright, R. J., & Chandler, L. D. (1998). Adult

728   susceptibility of Nebraska western corn rootworm (Coleoptera: Chrysomelidae)

729   populations to selected insecticides. *Journal of Economic Entomology*, *91*(3),

730   594–600. https://doi.org/10.1093/jee/91.3.594

731 Metcalf, R. L. (1986). Coevolutionary adaptations of rootworm beetles (Coleoptera:

732   Chrysomelidae) to cucurbitacins. *Journal of Chemical Ecology*, *12*(5), 1109–

733   1124. https://doi.org/10.1007/BF01638999

734 Miller, N. J., Guillemaud, T., Giordano, R., Siegfried, B. D., Gray, M. E., Meinke, L. J., &

735   Sappington, T. W. (2009). Genes, gene flow and adaptation of *Diabrotica*

736   *virgifera virgifera*. *Agricultural and Forest Entomology*, *11*(1), 47–60.

https://doi.org/10.1111/j.1461-9563.2008.00398.x

Miller, N. J., Richards, S., & Sappington, T. W. (2010). The prospects for sequencing

the western corn rootworm genome. *Journal of Applied Entomology*, *134*(5),

420–428. https://doi.org/10.1111/j.1439-0418.2009.01448.x

Miller, N. J., & Zhao, Z. (2015). Transcriptional responses of *Diabrotica virgifera virgifera*

larvae to benzoxazinoids. *Journal of Applied Entomology*, *139*(6), 416–423.

https://doi.org/10.1111/jen.12193

Nie, R., Andújar, C., Gómez-Rodríguez, C., Bai, M., Xue, H.-J., Tang, M., Yang, C.-T.,

Tang, P., Yang, X.-K., & Vogler, A. P. (2020). The phylogeny of leaf beetles

(Chrysomelidae) inferred from mitochondrial genomes. *Systematic Entomology*,

45(1), 188–204. https://doi.org/10.1111/syen.12387

Oki, N., Yano, K., Okumoto, Y., Tsukiyama, T., Teraishi, M., & Tanisaka, T. (2008). A

genome-wide view of miniature inverted-repeat transposable elements (MITEs)

in rice, *Oryza sativa* ssp.*japonica*. *Genes & Genetic Systems*, *83*(4), 321–329.

https://doi.org/10.1266/ggs.83.321

Oliver, K. R., & Greene, W. K. (2009). Transposable elements: Powerful facilitators of

evolution. *BioEssays: News and Reviews in Molecular, Cellular and*

Developmental Biology, *31*(7), 703–714. https://doi.org/10.1002/bies.200800219

O'Neal, M., Difonzo, C. D., Landis, D., & Meek, D. (2002). Monitoring *Diabrotica*

virgifera virgifera (LeConte) in Michigan soybean fields and subsequent adult

emergence in rotated and continuous cornfields. *Great Lakes Entomologist*, *35*,

173–181.

Osanai-Futahashi, M., Suetsugu, Y., Mita, K., & Fujiwara, H. (2008). Genome-wide

screening and characterization of transposable elements and their distribution

analysis in the silkworm, *Bombyx mori*. *Insect Biochemistry and Molecular*

Biology, *38*(12), 1046–1057. https://doi.org/10.1016/j.ibmb.2008.05.012

Palmer, M., & Petitpierre, E. (1996). Relationship of genome size to body size in *Phylan*

semicostatus(Coleoptera: Tenebrionidae). *Annals of the Entomological Society of*

America, *89*(2), 221–225. https://doi.org/10.1093/aesa/89.2.221

Palmer, M., Petitpierre, E., & Pons, J. (2003). Test of the correlation between body size

and DNA content in *Pimelia* (Coleoptera: Tenebrionidae) from the Canary

islands. *European Journal of Entomology*, *100*, 123–129.

https://doi.org/10.14411/eje.2003.022

Pereira, A. E., Wang, H., Zukoff, S. N., Meinke, L. J., French, B. W., & Siegfried, B. D.

(2015). Evidence of field-evolved resistance to bifenthrin in western corn

rootworm (*Diabrotica virgifera virgifera* LeConte) populations in western

Nebraska and Kansas. *PLOS ONE*, *10*(11), e0142299.

https://doi.org/10.1371/journal.pone.0142299

Petitpierre, E., Segarra, C., & Juan, C. (1993). Genome Size and Chromosomal

Evolution in Leaf Beetles (Coleoptera, Chrysomelidae). *Hereditas*, *119*(1), 1–6.

https://doi.org/10.1111/j.1601-5223.1993.00001.x

Pombert, J.-F. (2021). *PombertLab/SSRG: SSRG-v2.0.1*. Zenodo.

https://doi.org/10.5281/zenodo.5532794

R Core Team (2021). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/

Rehm, S., & Wessels, J. H. (1957). Bitter principles of the cucurbitaceae. VIII.—

Cucurbitacins in seedlings—Occurrence, biochemistry and genetical aspects.

Journal of the Science of Food and Agriculture, *8*(12), 687–691.

https://doi.org/10.1002/jsfa.2740081204

Reid, C. A. M. (1995). A cladistic analysis of subfamilial relationships in the

Chrysomelidae sensu lato (Chrysomeloidea). In: Pakaluk J. & Slipinski S.A.

(eds): *Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating*

the 80th Birthday of Roy A. Crowson. Muzeum i Instytut Zoologii PAN,

Warszawa. 2, 559-631.

Robert, C. A., Veyrat, N., Glauser, G., Marti, G., Doyen, G. R., Villard, N., Gaillard, M.

D. P., Köllner, T. G., Giron, D., Body, M., Babst, B. A., Ferrieri, R. A., Turlings, T.

C. J., & Erb, M. (2012). A specialist root herbivore exploits defensive metabolites

to locate nutritious tissues: A root herbivore exploits plant defences. *Ecology*

Letters, *15*(1), 55–64. https://doi.org/10.1111/j.1461-0248.2011.01708.x

Robert, C. A., Zhang, X., Machado, R. A., Schirmer, S., Lori, M., Mateo, P., Erb, M., &

Gershenzon, J. (2017). Sequestration and activation of plant toxins protect the

western corn rootworm from enemies at multiple trophic levels. *ELife*, *6*, e29307.

https://doi.org/10.7554/eLife.29307

Rostant, W. G., Wedell, N., & Hosken, D. J. (2012). Transposable elements and

insecticide resistance. In *Advances in Genetics* (Vol. 78, pp. 169–201). Elsevier.

https://doi.org/10.1016/B978-0-12-394394-1.00002-X

Russell, S. J., & LaMarre, J. (2018). Transposons and the PIWI pathway: Genome

defense in gametes and embryos. *Reproduction*, *156*(4), R111–R124.

https://doi.org/10.1530/REP-18-0218

Saha, A., Mitchell, J. A., Nishida, Y., Hildreth, J. E., Ariberre, J. A., Gilbert, W. V., &

807   Garfinkel, D. J. (2015). A trans-Dominant Form of Gag Restricts Ty1

808   Retrotransposition and Mediates Copy Number Control. *Journal of Virology*,

809   *89*(7), 3922–3938. https://doi.org/10.1128/JVI.03060-14

810   Santiago, N., Herráiz, C., Goñi, J. R., Messeguer, X., & Casacuberta, J. M. (2002).

811   Genome-wide analysis of the *Emigrant* family of MITEs of *Arabidopsis thaliana*.

812   *Molecular Biology and Evolution*, *19*(12), 2285–2293.

813   https://doi.org/10.1093/oxfordjournals.molbev.a004052

814   Sappington, T. W., Siegfried, B. D., & Guillemaud, T. (2006). Coordinated *Diabrotica*

815   genetics research: Accelerating progress on an urgent insect pest problem.

816   *American Entomologist*, *52*(2), 90–97. https://doi.org/10.1093/ae/52.2.90

817   Sasai, H., Ishida, M., Murakami, K., Tadokoro, N., Ishihara, A., Nishida, R., & Mori, N.

818   (2009). Species-Specific Glucosylation of DIMBOA in Larvae of the Rice

819   Armyworm. *Bioscience, Biotechnology, and Biochemistry*, *73*(6), 1333–1338.

820   https://doi.org/10.1271/bbb.80903

821   Schoville, S. D., Chen, Y. H., Andersson, M. N., Benoit, J. B., Bhandari, A., Bowsher, J.

822   H., Brevik, K., Cappelle, K., Chen, M.-J. M., Childers, A. K., Childers, C.,

823   Christiaens, O., Clements, J., Didion, E. M., Elpidina, E. N., Engsontia, P.,

824   Friedrich, M., García-Robles, I., Gibbs, R. A., … Richards, S. (2018). A model

825   species for agricultural pest genomics: The genome of the Colorado potato

826   beetle, *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Scientific*

827   *Reports*, *8*(1), 1931. https://doi.org/10.1038/s41598-018-20154-1

Smit, A. F. A., Hubley, R., & Green, P. (1996). Repeat-Masker open-3.0.

Http://Www.Repeatmasker.Org. https://ci.nii.ac.jp/naid/10029514778/

Souza, D., Peterson, J. A., Wright, R. J., & Meinke, L. J. (2020). Field efficacy of soil

insecticides on pyrethroid-resistant western corn rootworms (*Diabrotica virgifera*

virgifera LeConte). *Pest Management Science*, *76*(2), 827–833.

https://doi.org/10.1002/ps.5586

Souza, D., Vieira, B. C., Fritz, B. K., Hoffmann, W. C., Peterson, J. A., Kruger, G. R., &

Meinke, L. J. (2019). Western corn rootworm pyrethroid resistance confirmed by

aerial application simulations of commercial insecticides. *Scientific Reports*, *9*(1),

6713. https://doi.org/10.1038/s41598-019-43202-w

Stuart, T., Eichten, S. R., Cahn, J., Karpievitch, Y. V., Borevitz, J. O., & Lister, R.

(2016). Population scale mapping of transposable element diversity reveals links

to gene regulation and epigenomic variation. *ELife*, *5*, e20777.

https://doi.org/10.7554/eLife.20777

Talla, V., Suh, A., Kalsoom, F., Dincă, V., Vila, R., Friberg, M., Wiklund, C., &

Backström, N. (2017). Rapid increase in genome size as a consequence of

transposable element hyperactivity in wood-white (*Leptidea*) butterflies. *Genome*

Biology and Evolution, *9*(10), 2491–2505. https://doi.org/10.1093/gbe/evx163

Tallamy, D. W., Whittington, D. P., Defurio, F., Fontaine, D. A., Gorski, P. M., & Gothro,

P. W. (1998). Sequestered Cucurbitacins and Pathogenicity of *Metarhizium*

anisopliae (Moniliales: Moniliaceae) on Spotted Cucumber Beetle Eggs and

Larvae (Coleoptera: Chrysomelidae). *Environmental Entomology*, *27*(2), 366–

372. https://doi.org/10.1093/ee/27.2.366

Thomas-Bulle, C., Piednoël, M., Donnart, T., Filée, J., Jollivet, D., & Bonnivard, É.

(2018). Mollusc genomes reveal variability in patterns of LTR-retrotransposons

dynamics. *BMC Genomics*, *19*(1), 821. https://doi.org/10.1186/s12864-018-5200-

1

Toepfer, S., Haye, T., Erlandson, M., Goettel, M., Lundgren, J. G., Kleespies, R. G.,

Weber, D. C., Walsh, G. C., Peters, A., Ehlers, R.-U., Strasser, H., Moore, D.,

Keller, S., Vidal, S., & Kuhlmann, U. (2009). A review of the natural enemies of

beetles in the subtribe Diabroticina (Coleoptera: Chrysomelidae): implications for

sustainable pest management. *Biocontrol Science and Technology*, *19*(1), 1–65.

https://doi.org/10.1080/09583150802524727

Tomaszewska, P., Pellny, T. K., Hernández, L. M., Mitchell, R. A. C., Castiblanco, V.,

Vega, J. J. de, Schwarzacher, T., & Heslop-Harrison, P. (J S.). (2021). Flow

cytometry-based determination of ploidy from dried leaf specimens in

genomically complex collections of the tropical forage grass *Urochloa* s. L.

BioRxiv, 2021.03.26.437252. https://doi.org/10.1101/2021.03.26.437252

Voronova, A., Belevich, V., Jansons, A., & Rungis, D. (2014). Stress-induced

transcriptional activation of retrotransposon-like sequences in the Scots pine

(Pinus sylvestris L.) genome. *Tree Genetics & Genomes*, *10*(4), 937–951.

https://doi.org/10.1007/s11295-014-0733-1

Wang, L., Wang, J., Ma, Y., Wan, P., Liu, K., Cong, S., Xiao, Y., Xu, D., Wu, K., Fabrick,

J. A., Li, X., & Tabashnik, B. E. (2019). Transposon insertion causes cadherin

mis-splicing and confers resistance to Bt cotton in pink bollworm from China.

Scientific Reports, *9*(1), 7479. https://doi.org/10.1038/s41598-019-43889-x

Wang, S., Lorenzen, M. D., Beeman, R. W., & Brown, S. J. (2008). Analysis of repetitive

DNA distribution patterns in the *Tribolium castaneum* genome. *Genome Biology*,

9(3), R61. https://doi.org/10.1186/gb-2008-9-3-r61

Wang, X., Fang, X., Yang, P., Jiang, X., Jiang, F., Zhao, D., Li, B., Cui, F., Wei, J., Ma,

C., Wang, Y., He, J., Luo, Y., Wang, Z., Guo, X., Guo, W., Wang, X., Zhang, Y.,

Yang, M., … Kang, L. (2014). The locust genome provides insight into swarm

formation and long-distance flight. *Nature Communications*, *5*(1), 2957.

https://doi.org/10.1038/ncomms3957

Wang, X., Xu, Y., Zhang, S., Cao, L., Huang, Y., Cheng, J., Wu, G., Tian, S., Chen, C.,

Liu, Y., Yu, H., Yang, X., Lan, H., Wang, N., Wang, L., Xu, J., Jiang, X., Xie, Z.,

Tan, M., … Xu, Q. (2017). Genomic analyses of primitive, wild and cultivated

citrus provide insights into asexual reproduction. *Nature Genetics*, *49*(5), 765–

772. https://doi.org/10.1038/ng.3839

Weinstock, G. M., Robinson, G. E., Gibbs, R. A., Weinstock, G. M., Weinstock, G. M.,

Robinson, G. E., Worley, K. C., Evans, J. D., Maleszka, R., Robertson, H. M.,

Weaver, D. B., Beye, M., Bork, P., Elsik, C. G., Evans, J. D., Hartfelder, K., Hunt,

G. J., Robertson, H. M., Robinson, G. E., … Genome assembly: (2006). Insights

into social insects from the genome of the honeybee *Apis mellifera*. *Nature*,

443(7114), 931–949. https://doi.org/10.1038/nature05260

Wendel, J. F., & Wessler, S. R. (2000). Retrotransposon-mediated genome evolution on

a local ecological scale. *Proceedings of the National Academy of Sciences*,

97(12), 6250–6252. https://doi.org/10.1073/pnas.97.12.6250

Wessler, S. R., Bureau, T. E., & White, S. E. (1995). LTR-retrotransposons and MITEs:

Important players in the evolution of plant genomes. *Current Opinion in Genetics*

& Development, *5*(6), 814–821. https://doi.org/10.1016/0959-437x(95)80016-x

Wicker, T., Sabot, F., Hua-Van, A., Bennetzen, J. L., Capy, P., Chalhoub, B., Flavell, A.,

Leroy, P., Morgante, M., Panaud, O., Paux, E., SanMiguel, P., & Schulman, A. H.

(2007). A unified classification system for eukaryotic transposable elements.

Nature Reviews Genetics, *8*(12), 973–982. https://doi.org/10.1038/nrg2165

Yan, H., Bombarely, A., & Li, S. (2020). *DeepTE: A computational method for de novo*

classification of transposons with convolutional neural network [Preprint].

Bioinformatics. https://doi.org/10.1101/2020.01.27.921874

Zhang, S., Gu, S., Ni, X., & Li, X. (2019). Genome size reversely correlates with host

plant range in *Helicoverpa* species. *Frontiers in Physiology*, *10*, 29.

https://doi.org/10.3389/fphys.2019.00029