Hippo pathway and Bonus control developmental cell fate decisions in the *Drosophila* eye

Heya Zhao[1], Kenneth H. Moberg[2], Alexey Veraksa[1, 3, *]

[1] Department of Biology, University of Massachusetts Boston, Boston, MA, 02125, USA

[2] Department of Cell Biology, Emory University School of Medicine, Atlanta, GA, 30322, USA

[3] Lead Contact

* Correspondence: alexey.veraksa@umb.edu

Summary

The canonical function of the Hippo signaling pathway is regulation of organ growth. How this pathway controls cell fate determination is less well understood. Here, we uncover a function of the Hippo pathway in developmental cell fate decisions in the *Drosophila* eye-antennal disc exerted through the interaction of Yorkie (Yki) with the transcriptional regulator Bonus (Bon), an ortholog of mammalian Transcriptional Intermediary Factor 1/tripartite motif (TIF1/TRIM) family proteins. Instead of controlling tissue growth, Yki and Bon promote epidermal and antennal fates at the expense of the eye fate. Proteomic, transcriptomic, and genetic analyses reveal that Yki and Bon control these cell fate decisions by recruiting transcriptional and post-transcriptional co-regulators, and by activating epidermal differentiation genes and repressing Notch target genes. Our work expands the range of functions and regulatory mechanisms under Hippo pathway control.

Keywords: Yki, Bon, cell fate, eye specification, Notch, E(spl)-C, HDAC1, Su(var)2-10, transcriptional regulation, protein interaction

Introduction

Organ growth and cell fate determination are critical processes in organism development. These
processes are controlled and kept in balance by multiple signaling pathways whose dysregulation
leads to developmental abnormalities and diseases. One such pathway is the evolutionarily
conserved Hippo pathway, which is implicated in various physiological functions and pathological
conditions including organ growth, cell differentiation, tissue regeneration, and oncogenesis
(Kulkarni et al., 2020; Zheng and Pan, 2019). The Hippo pathway transduces upstream regulatory
signals to transcriptional outputs that mediate various cellular functions including cell proliferation
and survival (Meng et al., 2016). The core Hippo pathway acts via the Hippo (Hpo)/Warts (Wts)
kinase cascade to negatively regulate the activity of the transcriptional coactivator Yorkie (Yki),
which associates with DNA-binding proteins such as Scalloped (Sd) to control gene expression
(Zheng and Pan, 2019). The canonical transcriptional targets of the Yki-Sd complex in *Drosophila*
include *cyclin E* (*cycE*), *death-associated inhibitor of apoptosis 1* (*diap1*), *bantam* microRNA (*mir-*
ban), and *expanded* (*ex*); these factors promote proliferation, inhibit apoptosis, and enable
negative feedback regulation (Pan, 2010). Although increasing evidence supports an essential
role of the Hippo pathway in cell fate determination in multiple biological contexts (Davis and
Tapon, 2019), the cellular mechanisms remain poorly understood.

The *Drosophila* eye is an excellent model to study gene regulatory networks controlling cell fate
determination (Weasner and Kumar, 2022). Most of the *Drosophila* adult head structures develop
from the larval eye-antennal disc, with the compound eye and ocelli originating from the eye disc
compartment, the antenna and maxillary palp from the antennal compartment, and the head
epidermis from the tissues surrounding the two compartments (Dominguez and Casares, 2005).
Segregation of the mutually antagonistic eye, antennal, and head epidermal fates, which begins
at the second instar larval stage (L2), is regulated by several signaling pathways including Notch,
EGFR, Wingless, and Hedgehog, and retinal determination genes such as *eyeless* (*ey*) and
homothorax (*hth*) (Kumar, 2010; 2018). Alteration of these regulatory inputs can cause a switch
from one fate to another, leading to partial or in some cases complete homeotic transformations
of the affected structures (Kumar and Moses, 2001; Ordway et al., 2021; Wang and Sun, 2012;
Weasner and Kumar, 2013). Key patterning events in the eye are linked to a wave of
differentiation called the morphogenetic furrow (MF) that starts in the early third instar larval stage
(L3) and proceeds from the posterior to the anterior of the eye disc, resulting in the differentiation
of an array of optical units called ommatidia, each consisting of photoreceptor cells, cone cells,

primary pigment cells, interommatidial bristles, and secondary and tertiary pigment cells (also
called interommatidial cells) (Cagan and Ready, 1989a; Charlton-Perkins and Cook, 2010).

Previous studies of the Hippo pathway in *Drosophila* eye differentiation focused on MF
progression, terminal differentiation of photoreceptor cells, and formation of peripodial epithelium
(Jukam et al., 2013; Pojer et al., 2021; Wittkorn et al., 2015; Zhang et al., 2011). Mutant analyses
of the Hippo pathway components *ex*, *Merlin* (*Mer*) and *mob as tumor suppressor* (*mats*) have
suggested an earlier and broader impact of the Hippo pathway in eye specification (Boedigheimer
and Laughon, 1993; Lai et al., 2005; McCartney et al., 2000; Pellock et al., 2007). However, the
involvement of the Hippo pathway in controlling cell fate decisions among the eye, antenna, and
head epidermis remains elusive, and the underlying transcriptional mechanisms are unknown.

We reasoned that the Hippo pathway may have a novel function in controlling the eye-antenna-
epidermis fate determination through yet unknown interactors that regulate the transcriptional
output of the Yki-Sd complex. To identify such interactors, we performed proteomic analyses and
identified a novel Yki-interacting protein, Bonus (Bon). Bon is the only *Drosophila* ortholog of
mammalian TIF1 family proteins TIF1α (TRIM24), TIF1β (TRIM28/KAP1), TIF1γ (TRIM33), and
TIF1δ (TRIM66) (Beckstead et al., 2001; Cammas et al., 2012). TIF1/Bon proteins are chromatin-
associated factors that activate or repress transcription by binding to co-regulators and controlling
chromatin state (Beckstead *et al.*, 2001; Beckstead et al., 2005; Cammas *et al.*, 2012; Ito et al.,
2012). TIF1 proteins play various roles during vertebrate development and are implicated in
cancer (Cambiaghi et al., 2012; Petrera and Meroni, 2012). Invertebrate Bon is essential for
nervous system development, metamorphosis, and cell survival (Allton et al., 2009; Beckstead *et*
al., 2001; Ito *et al.*, 2012; Salzberg et al., 1997).

Here, we present evidence that Bon and the Hippo pathway co-regulate major cell fate decisions
during the development of the *Drosophila* eye. Yki and Bon bind via WW domain-PPxY motif
interactions and cooperate to produce epidermal cells in the eye at the expense of ommatidial
cells, while loss of *bon* induces ectopic eye markers, suggesting that the Hippo pathway and Bon
control the choice between the eye and epidermal fates. The Hippo pathway and Bon also
regulate the eye-antennal specification, with Yki and Bon inhibiting the eye fate and promoting
the antennal fate. Through the analysis of Bon and Yki protein interactors, we have identified
multiple transcriptional and post-transcriptional regulators that are necessary for their control of
cell fate decisions. Transcriptome analysis has revealed that Bon and Yki exert their functions by

jointly activating epidermal differentiation genes and, unexpectedly, repressing Notch target
genes. Overall, we have identified a novel function of the Hippo pathway in the
eye/antenna/epidermis cell fate decisions during *Drosophila* eye development. This function
requires the interaction of Yki with Bon, their recruitment of coregulators, and the joint
transcriptional control of a non-canonical set of target genes.

Results

Bon is a novel Yki interactor
To identify new regulators of the Hippo pathway, we performed affinity purification-mass
spectrometry (AP-MS) using *Drosophila* embryos expressing Yki-EGFP with a ubiquitous driver
da-GAL4 (Wodarz et al., 1995), as well as cultured *Drosophila* S2 cells expressing streptavidin-
binding peptide (SBP)-tagged Yki (see Method Details). We identified the core Hippo pathway
components, accessory regulators, and several putative novel Yki interactors, including Bon
(Figure 1A, Table S1) (Kwon et al., 2013; Oh et al., 2014; Zheng and Pan, 2019). We confirmed
the interaction between Bon and Yki by co-immunoprecipitation (co-IP) of tagged Bon and Yki
proteins in S2 cells (Figures 1C-D). We hypothesized that the Yki-Bon interaction may be
mediated by the two PPxY motifs in Bon (PPLY[507] and PPSY[585]) binding to the WW domains in
Yki (Figure 1B), similar to other known Yki interactors (Huang et al., 2005; Oh *et al.*, 2014).
Mutating the key tyrosine residue to alanine (Zhang et al., 2015) in either single or both WW
domains in Yki strongly reduced its binding to Bon, with the first WW domain (WW1) showing a
stronger reduction in binding upon mutation (Figure 1C). Similarly, substituting tyrosine with
alanine in one or both PPxY motifs in Bon also largely reduced its binding to Yki, with a stronger
reduction by the mutation in the second PPxY motif (PPxY2) (Figure 1D). These results indicate
that Bon interacts with Yki through PPxY motifs in Bon and WW domains in Yki.
89

Activation of Bon or Yki induces epidermal trichomes in adult eyes
We next sought to investigate whether Bon is involved in Yki-mediated growth regulation. RNAi
knockdown of *bon* did not affect the overgrowth or cell proliferation induced by Yki overexpression
in adult eyes, larval eye discs, and wing discs (Figures S1A-D). In addition, RNAi knockdown of
bon in wing discs did not affect the reporters of canonical Yki target genes *diap1* and *ex,* which
mediate Yki's function in growth regulation through anti-apoptosis and feedback regulation (Figure
S1E). Loss-of-function *bon* mutants exhibit ectopic abdominal peripheral neurons in the embryo

(Beckstead *et al.*, 2001; Salzberg *et al.*, 1997). Null mutants of *wts* (*wts^{X1}*) (Xu et al., 1995) and

yki (*yki^{B5}*) (Huang *et al.*, 2005) did not show loss or gain of peripheral neurons and did not affect

the ectopic neuron phenotype in null (*bon^{21B}*) or hypomorphic (*bon^{487}*) *bon* mutant embryos

(Figure S1F). Collectively, these data argue that Yki and Bon have independent functions in

growth regulation and embryonic peripheral nervous system (PNS) development.

Interestingly, overexpression of Bon with *GMR-GAL4*, which drives expression in all cells after

the MF and persists through pupal eye development (Ellis et al., 1993; Hay et al., 1994), resulted

in the formation of trichomes on the surface of adult eyes and disruption of the ommatidial array

(Figures 2A-B and S2A), which is consistent with a previous report (Tardi et al., 2012). Trichomes

are actin-rich, non-neuronal apical extensions, which are present on *Drosophila* epidermal cells

but not retinal cells (Arif et al., 2015; Delon et al., 2003). This trichome induction suggests that

Bon promotes epidermal fate in the eye.

To examine whether the Hippo pathway is involved in Bon trichome induction, we overexpressed

or knocked down key components of the Hippo pathway in the Bon overexpression background

and quantified trichome number. Yki overexpression or *wts* knockdown increased Bon-induced

trichomes (Figures 2C-D and 2H). In contrast, knockdown of *yki* or *sd* strongly suppressed Bon-

induced trichomes, with only mild eye roughness observed in single knockdowns (Figures 2E-F,

2H, S2B, and S2R-T). Thus, the core Hippo pathway and Yki transcriptional activity are essential

for Bon-induced trichomes in adult eyes. Furthermore, overexpression of Bon-PPxA, a Bon

mutant with both PPxY motifs mutated to PPxA that cannot bind to Yki (see Figure 1D), resulted

in significantly fewer trichomes and milder ommatidial defects (Figures 2G-H), indicating that Bon

must bind Yki to efficiently induce eye trichomes.

We then investigated whether Yki activation is sufficient to induce trichomes in adult eyes. Indeed,

Yki overexpression by *GMR-GAL4* resulted in trichome formation in the eye (Figures 2I and S2C-

D). Overexpression of constitutively active Yki-S168A, a Yki mutant insensitive to Wts

phosphorylation and inhibition (Oh and Irvine, 2008), induced even more trichomes (Figures 2J,

2O, and S2G). Knockdown of *wts* also induced more trichomes than Yki overexpression (Figures

2K, 2O, and S2K). Therefore, Yki activation promotes trichome formation, with trichome number

positively correlated with Yki activity. Knockdown of *bon* strongly suppressed trichome formation

induced by wild-type Yki, Yki-S168A, and *wts*-RNAi, with only mild roughness when knocked

down alone (Figures 2L-N, 2P-R, and S2D-Q), suggesting that Bon is necessary for trichome

formation induced by activated Yki. Together, these results show that the Hippo pathway and Bon

jointly regulate the epidermal trichome formation in the eye.

Hippo pathway and Bon control the decision between eye and epidermal cell fates

Next, we investigated cell fate changes underlying ectopic trichome induction in the retina. The

establishment of trichomes occurs at 30-36 hrs after puparium formation (APF) in the wing and

38-39 hrs APF in the notum (Chanut-Delalande et al., 2014; Delon *et al.*, 2003). F-actin and cell

boundary (Dlg) staining of pupal eyes revealed that Bon-induced trichomes are initiated at 40 hrs

APF (Figure S3A-C) and are easily visible at 44 hrs APF (Figures 3A). Bon overexpression

resulted in severe disruption of the ommatidia and produced extra interommatidial-like cells from

which the trichomes derive (Figure 3A). Importantly, the excess interommatidial-like cells were

generated without alteration in DNA synthesis (Figure S1D) or overall apical cell numbers (Figure

S3E), suggesting a failure of differentiation rather than aberrant proliferation or survival.

Interommatidial cells are believed to be the default state in retinal differentiation (Charlton-Perkins

and Cook, 2010; Kumar, 2012). Trichomes on these cells indicate that they may be

reprogrammed into epidermal-like cells. Knockdown of *yki* or *sd* strongly suppressed the initiation

of Bon-induced trichomes and partially restored the ommatidia (Figure 3A). Therefore, Bon

induces epidermal cell fate at the expense of the eye fate in a Yki and Sd-dependent manner.

To determine more precisely which retinal cells were affected, we immunostained pupal eyes at

44 hrs APF for cone cell marker Ct, primary pigment cell marker BarH1, and photoreceptor cell

and bristle group marker Elav (Charlton-Perkins and Cook, 2010). Bon overexpression with *GMR-*

GAL4 strongly inhibited formation of cone cells and primary pigment cells, and affected patterning

of photoreceptor cells and bristle groups (Figures 3A-B and S3D). *yki* or *sd* knockdown in this

background suppressed the loss of cone cells and partially restored the mispatterned

photoreceptor cells and bristle groups (Figures 3A-B and S3D), but did not rescue the loss of

primary pigment cells (Figure S3D), likely because the differentiation of primary pigment cells

requires successful differentiation of cone cells (Nagaraj and Banerjee, 2007). These results

demonstrate that Bon, Yki, and Sd jointly suppress eye fate.

wts knockdown by RNAi also induced trichomes on interommatidial cells at mid-pupal stage,

suggesting that these cells were reprogrammed into the epidermal fate (Figure 3C). Although the

trichomes induced by *wts* RNAi were shorter and thicker than Bon-induced trichomes (Figures 2B

and 2K), they were clearly distinguishable from the bristle shafts of the bristle groups which have

sockets (Figure 3C, Dlg staining) and neuronal input (Figure S3I, Elav staining) (Charlton-Perkins

and Cook, 2010). *bon* knockdown strongly suppressed *wts* RNAi-induced trichome initiation, but

not the excess interommatidial cells and bristles which result from overproliferation (Huang *et al.*,

2005; Udan et al., 2003), suggesting that Bon is specifically necessary for cell fate reprogramming

(Figure 3C). Consistent with previous studies, *wts* RNAi reduced the number of cone cells

(Hamaratoglu et al., 2006; Wittkorn *et al.*, 2015), which was rescued by knockdown of *bon*

(Figures 3C-D). Moreover, *bon* knockdown alone resulted in gain of cone cells, especially at the

periphery of the retina (Figures S3F-G). Loss of *wts* and *bon* had a mild effect on photoreceptor

patterning and showed a mixed outcome for primary pigment cells (either gain or loss per

ommatidium) (Figure S3H). Together, these results show that Bon activation or Hippo pathway

inactivation behind the MF induces epidermal fate at the expense of the eye fate, as represented

by trichome formation and inhibition of cone cells.

We further investigated endogenous control of cell fate determination using loss of function clones

of *wts* and *bon* alleles. Trichomes were apparent in *wts^{X1}* eye clones at 46 hrs APF, and the

trichome-making cells strongly expressed Hth, which normally marks the head epidermis at this

stage (Figure 3E-F) (Lim and Tomlinson, 2006; Wernet et al., 2003). *wts^{X1}* clones also suppressed

eye fate, as evidenced by loss of ommatidia (Dlg in Figure 3F) and photoreceptors (Elav in Figure

3G). In adult eyes, *wts^{X1}* clones produced outgrowths of epidermal tissue covered with trichomes

(Figure 3H). Interestingly, *bon^{21B}* clones led to gain of neuronal eye marker Elav at the basal side

of the larval eye disc, without affecting the normal apical patterns of Elav, suggesting an induction

of ectopic eye tissue (Figure 3I). Altogether, our results using both loss and gain of function

analyses suggest that Bon and Yki (or inactivated core Hippo pathway) suppress eye fate and

promote the epidermal fate.

Hippo pathway and Bon regulate the choice between eye and antennal cell fates

Since eye fate is antagonistic to epidermal and antennal fates (Weasner and Kumar, 2022), we

asked whether Hippo and Bon also regulate the balance of eye-antennal fates. To explore this

hypothesis, we used the early eye driver *ey-GAL4* which is expressed in the entire eye disc

beginning at the L2 stage, and examined L3 eye-antennal discs using Hth and Ct as antennal

markers, and Elav as the differentiated neuron/eye marker (Blochlinger et al., 1993; Kumar and

Moses, 2001; Pai et al., 1998; Wang and Sun, 2012). *wts* knockdown or Yki-S168A

overexpression by *ey-GAL4* significantly suppressed the eye field, with the respective penetrance

of 56% (*wts*-RNAi), 100% (*yki-S168A*-1), and 85% (*yki-S168A*-2), and even occasionally led to a

complete eye-to-antenna transformation, with the respective penetrance of 11%, 5%, and 6%
(Figures 4A-C, 4I, S4A-C, S4E, S4I-K and Table S5). Remarkably, *wts* knockdown or Yki-S168A
overexpression also led to clear eye-to-antenna transformations in some adult flies (Figures 4E-
G). Wildtype Yki overexpression with *ey-GAL4* suppressed the eye field and resulted in extra
bristles (vibrissae) and occasional epidermal outgrowths, but without eye-to-antenna
transformation (Figures S4G-G''').

bon knockdown significantly rescued the smaller eye field resulting from Yki-S168A
overexpression (from 100% to 3%, and 85% to 43%, with two independent lines) and suppressed
formation of double antennae (from 5% to 0%, and 6% to 2.5%, with two lines) (Figures 4D, 4I,
S4D, S4F, and Table S5). In adults, *bon* knockdown also largely rescued Yki-S168A-induced loss
of the eye (Figure 4H). Bon overexpression using *ey-GAL4* suppressed the differentiated eye field
in eye-antennal discs with a penetrance of 20% (Figures 4J, S4O, and Table S5), and led to a
complete or partial loss of the eye with frequent epidermal outgrowths in adults, with a penetrance
of 18% (Figure 4K and Table S5), but without eye-to-antenna transformation. In contrast, Bon-
PPxA overexpression led to largely normal eyes (Figure 4L and Table S5). Altogether, these
results show that early and strong activation of Yki across the eye field can switch the eye fate to
antennal fate in a Bon-dependent manner, while weaker activation of Yki or Bon transforms eye
to epidermis.

To determine whether the eye vs. antennal fate choice is under control of endogenous *yki* and
bon, we tested effects of *yki/bon* loss in the antenna using the *hth-GAL4* driver (Dominguez and
Casares, 2005; Wernet *et al.*, 2003). Since eye and antennal fates are mutually antagonistic and
reciprocally transformable (Wang and Sun, 2012), we hypothesized that loss of *yki* or *bon* in
antenna would promote eye fate. Indeed, *yki* knockdown alone resulted in variable degrees of
antenna loss with 100% penetrance, and moderate antenna-to-eye transformations (13%
penetrance) (Figures 4M-N, 4P, S4L-N, and Table S5). This partial antenna-to-eye transformation
exhibited loss of Ct and gain of Elav in the antennal field, suggesting a suppression of the antennal
fate and differentiation of ectopic eye tissue (Figure 4N). Strikingly, a double knockdown of *yki*
and *bon* resulted in a severe antenna-to-eye transformation with a 26% penetrance, in which the
antennal field marked by Ct was restricted to the center of the eye-antennal disc, and the ectopic
Elav pattern resembled a mirror-image duplication of the endogenous eye field (Figures 4O-P).
These results demonstrate that endogenous Yki and Bon are necessary for the specification of
the antennal fate and the suppression of the eye fate. Altogether, our results support a model of

eye specification regulated by the Hippo pathway and Bon, where Yki and Bon promote epidermal

fate and the antennal fate, while suppressing eye fate (Figure 4Q).

Interactors of Bon and Yki are required for the eye-epidermal fate choice

We sought to further investigate the mechanism through which Yki and Bon regulate the eye-

epidermal fate choice. One potential mechanism is that Bon enhances Yki activity by competing

Yki away from Wts, given that both Bon and Wts bind to Yki through the PPxY-WW interaction.

However, co-IP in S2 cells showed no competition between Bon and Wts for binding to Yki (Figure

S5A). This result suggests that rather than competing with Wts, Bon may employ another

mechanism, such as engaging a special set of interactors to mediate the eye-epidermal cell fate

decision together with Yki.

To identify Bon cofactors, we analyzed the Bon interactome by AP-MS using Bon-SBP expressed

in S2 cells (Figure 5A and Table S2). Yki was identified as one of the top Bon interactors, further

validating their interaction (Figure 5A). In addition, we identified Histone deacetylase 1 (HDAC1),

Suppressor of variegation 2-10 (Su(var)2-10), and Heterogeneous nuclear ribonucleoprotein at

27C (Hrb27C) as top Bon interactors (Figure 5A). HDAC1 is a transcriptional corepressor that is

a key component of several chromatin remodeling complexes, such as NuRD, Sin3, and CoRest

(Seto and Yoshida, 2014). Su(var)2-10 belongs to the PIAS family of SUMO ligases and controls

gene transcription as well as *Drosophila* eye specification (Betz et al., 2001; Hari et al., 2001;

Ninova et al., 2020). Hrb27C is an abundant hnRNP that is involved in various aspects of post-

transcriptional mRNA regulation and is implicated in the modulation of Yki activity (Goodrich et

al., 2004; Mach et al., 2018). We tested whether these three Bon interactors are involved in

trichome formation in adult eyes. Knockdown of *HDAC1* strongly suppressed, whereas its

overexpression enhanced, both Bon- and Yki-S168A-induced trichomes (Figures 5B-C, 5G-H, 5L,

and S5B-C), and knockdown of *Su(var)2-10* or *Hrb27C* suppressed these trichomes (Figures 5D-

E, 5I-J, 5L, and S5B-C). Therefore, HDAC1, Su(var)2-10, and Hrb27C promote Bon- or Yki-

induced epidermal fate in the eye.

The transcription factor Shavenbaby/Ovo (Svb/Ovo) plays a key role in the formation and

patterning of epidermal trichomes (Chanut-Delalande et al., 2006; Delon *et al.*, 2003). Also, Svb

interacts with Yki and regulates apoptosis through *diap1* (Bohere et al., 2018). We tested if

Svb/Ovo was involved in the regulation of Bon- and Yki-induced trichomes in adult eyes.

Overexpression of the constitutive activator OvoB enhanced Bon and Yki-S168A induced

trichomes, while overexpression of the constitutive repressor OvoA, or knockdown of somatic *svb*,
strongly suppressed Bon- and Yki-S168A-induced trichomes (Figures 5F, 5K-L, and S5B-C)
(Andrews et al., 2000). These results suggest that Svb/Ovo is required for ectopic trichome
generation in the eye induced by Bon or Yki.

We then asked if these interactors are involved in Bon and Yki regulated suppression of eye fate.
In pupal eyes, knockdown of *HDAC1* or *svb* with *GMR-GAL4* suppressed Bon-induced trichomes
but did not rescue the loss of cone cells (Figures 5M-N). In contrast, knockdown of *Su(var)2-10*
or *Hrb27C* not only suppressed the trichomes, but also largely rescued the number of cone cells
(Figures 5M-N). Interestingly, *Su(var)2-10* RNAi even partially restored the loss of primary
pigment cells and mispatterned ommatidial array (Figure S5D). These results suggest that
Su(var)2-10 and Hrb27C contribute to both eye fate suppression and epidermal fate promotion in
the pupal eye, while HDAC1 and Svb/Ovo are only involved in the latter function at this stage. In
L3 eye-antennal discs, Bon overexpression with the *bi-GAL4* driver, which drives expression at
the dorsal and ventral margins of the eye disc from L2 stage (Tare et al., 2013), inhibited Elav
expression (Figures 5O-Q), similar to Yki overexpression in a previous report (Wittkorn *et al.*,
2015). Knockdown of *Su(var)2-10* or *HDAC1* suppressed the loss of Elav due to Bon
overexpression (Figures 5R-S and 5O). Furthermore, knockdown of *HDAC1* under *ey-GAL4*
rescued the loss of eye field due to Yki-S168A overexpression (Figures S5E-G). Collectively,
these results suggest that Bon, Yki, Hrb27C, Su(var)2-10, and HDAC1 work together to suppress
eye fate, and promote the epidermal fate with the involvement of Svb/Ovo.

Bon and Yki control cell fate decisions in the eye through transcriptional regulation of joint
target genes
Since Bon, Yki, and interactors described above are all transcriptional or post-transcriptional
regulators, we hypothesized that they mediate cell fate choices in the eye-antennal disc via joint
regulation of a unique set of transcriptional targets. To identify these target genes, we performed
RNA sequencing (RNA-seq) using pupal eyes at 40-41 hrs APF (when the trichomes initiate) that
overexpressed Bon, with or without a simultaneous RNAi knockdown of *yki*. Through differential
gene expression analysis, we identified 216 genes as Bon-activated/Yki-dependent genes, and
119 genes as Bon-repressed/Yki-dependent genes (Figures 6A, S6A-B, and Table S3).
Correlation coefficient analysis revealed the significantly concordant regulation of gene
expression by Bon and Yki (Figure 6B). Also, *ex* and *Rhodopsin 5* (*Rh5)*, which are known targets
of Yki in the pupal eye (Deng et al., 2020; Jukam *et al.*, 2013), were among Yki-dependent Bon-

activated genes (Figures 6A and S6A-B). These transcriptomic data suggest that Bon and Yki

work together to regulate gene expression in pupal eyes, consistent with our genetic results.

We evaluated genes jointly regulated by Bon and Yki by Gene Ontology analysis of Biological

Process (GO_BP, Table S4). Bon-activated/Yki-dependent genes were enriched for GO terms

"epidermal cell differentiation", "cuticle development", and "sensory perception of taste", while

Bon-repressed/Yki-dependent genes were enriched for GO terms "compound eye development"

and "Notch signaling pathway", consistent with the genetic function of Bon and Yki in promoting

epidermal fate and antennal fate as well as suppressing the eye fate (Figure 6C). Bon-

activated/Yki-dependent epidermal genes, *shavenoid* (*sha*), *forked* (*f*), *nyobe* (*nyo*), and *neyo*

(*neo*), are known Svb target genes that are essential for formation and patterning of epidermal

trichomes, suggesting that they may mediate Bon- and Yki-induced trichome formation in the eye

(Figures 6A and 6C) (Chanut-Delalande *et al.*, 2006; Fernandes et al., 2010). Bon-repressed/Yki-

dependent Notch pathway genes, *E(spl)mdelta-HLH*, *E(spl)m3-HLH* and *E(spl)m2-BFM*, are

members of the Enhancer of split gene complex (*E(spl)-C*), which is a major transcriptional target

of Notch (Figures 6A and 6C) (Knust et al., 1992; Lai et al., 2000). Two other Notch target genes,

ct and *Transcription factor AP-2* (*TfAP-2*), were also identified among Bon-repressed/Yki-

dependent genes (Figure 6A) (Kerber et al., 2001; Micchelli et al., 1997). Previous work showed

that Notch signaling promotes eye fate and suppresses antennal fate, and that Notch, E(spl)-C,

Ct, and TfAP-2 are all critical for cell fate establishment in the eye (Cagan and Ready, 1989b;

Kumar and Moses, 2001; Kurata et al., 2000; Lai and Rubin, 2001; Monge et al., 2001; Nagaraj

and Banerjee, 2007; Wang and Sun, 2012). These observations raise the intriguing possibility

that downregulation of Notch targets by Yki and Bon mediates suppression of eye fate and

promotion of antennal fate. Finally, Bon-activated/Yki-dependent genes included sensory

perception genes such as antenna-specific odorant receptor *Or47b*, as well as a group of

gustatory receptors (*Gr64a-f*) that are expressed in multiple sensory organs including the antenna

but excluding the eye, indicating an induction of antennal identity at the transcriptional level

(Figures 6A, 6C, and S6A-B) (Barish et al., 2017; Fujii et al., 2015; Menuz et al., 2014; Vosshall,

2001).

We validated the transcriptional regulation of *sha*, *f*, *E(spl)m3-HLH*, *E(spl)m2-BFM*, and

E(spl)mdelta-HLH by Bon and Yki using target-specific qPCR, which generally confirmed the

RNA-seq findings (Figure 6D). To determine whether the transcriptional targets of Yki and Bon

are under their direct control, we analyzed published ChIP-seq datasets for Yki and Bon, and the

DamID-seq dataset for Sd (Negre et al., 2011; Oh et al., 2013; Vissers et al., 2018). Pairwise overlaps of binding sites were significant, with 75 loci shared among all three proteins, including the *E(spl)-C* region, suggesting a general co-localization of Yki, Bon and Sd on chromosomes and their direct control of *E(spl)-C* (Figure 6E and S6C). Yki and Sd also bound to *ct* and *TfAP-2*, raising the possibility of a direct regulation (Figure S6C). Yki, Sd, and Bon did not associate with the epidermal differentiation genes and antennal sensory receptor genes, except for Yki binding to *neo*, suggesting indirect activation of these genes by Bon and Yki (Figure S6C). To validate the binding of Bon to the *E(spl)mdelta-HLH* genomic locus, we performed ChIP-qPCR using L3 eye-antennal discs expressing Bon-mCherry with *GMR-GAL4*. We chose two DNA segments (P1 and P2) that overlapped the Yki and Sd binding regions (Figure 6F). Bon bound to P2 but did not significantly associate with P1 (Figure 6G). Together, these results suggest that Bon and Yki jointly and directly repress Notch targets and indirectly activate epidermal and antennal markers.

Identification of genes that are directly repressed by Yki and Bon was unexpected and suggests that Bon binding may shift Yki activity towards repression. We further evaluated the repression of *E(spl)-C* by Bon and Yki using a BAC transgene reporter expressing GFP-tagged E(spl)mdelta-HLH (*GFPmδ*) and a transcriptional reporter of *E(spl)m3-HLH* (*m3-GFP*) (Couturier et al., 2019). In the L3 eye disc, both reporters are expressed within and posterior to the MF (Figures 6H and 6J). *wts^{X1}* clones showed strongly reduced expression of *GFPmδ* and *m3-GFP*, and moderately reduced Elav (Figures 6H and S6D). In pupal eyes 25 hrs APF, *GFPmδ* is mainly expressed in primary pigment cells (Figure 6I). *bon^{21B}* clones resulted in a gain of *GFPmδ*-positive cells, suggesting that Bon suppresses *E(spl)-C* and inhibits the primary pigment cell fate (Figure 6I). *wts* knockdown with the *DE-GAL4* driver, which is expressed in the dorsal eye in both columnar cells of the disc proper (DP) and squamous peripodial epithelium (PE) (Figure S6E) (Morrison and Halder, 2010), reduced *GFPmδ*, *m3-GFP*, and Elav (Figures 6J and S6F). Knockdown of *yki* or *sd* expanded *GFPmδ*, *m3-GFP*, and Elav into the PE (Figures 6K and S6G-I), reminiscent of a PE to DP transformation previously reported for loss of *yki* and *sd* (Zhang *et al.*, 2011). *bon* knockdown also led to occasional expansion of *m3-GFP* and Elav into the PE, with a penetrance of 8.3% and 12.5% for two RNAi lines (Figures 6L, S6J, and Table S5). Together, these results show that Bon, Yki and Sd repress, while Wts promotes, the expression of *E(spl)mdelta-HLH* and *E(spl)m3-HLH* in the eye.

We then asked whether the target genes we identified are involved in Bon and Yki control of eye-antenna-epidermal fate determination. Sha overexpression enhanced, whereas *f* knockdown

suppressed, trichomes induced by Bon and Yki-S168A in the eye (Figures 7A-C, 7E-G, 7I, S6K-
L). Therefore, *sha* and *f* promote Bon/Yki-induced epidermal fate in the eye, consistent with their
transcriptional activation by Bon and Yki. In contrast, overexpression of E(spl)mdelta-HLH and
E(spl)m3-HLH suppressed the trichomes induced by Bon and Yki-S168A, suggesting that they
inhibit the epidermal fate in the eye, in agreement with their transcriptional repression by Bon and
Yki (Figures 7D, 7H, 7I, S6M-N). Overexpression of E(spl)m3-HLH significantly rescued the loss
of eye field and completely suppressed the formation of double antennae due to Yki-S168A
overexpression, suggesting that E(spl)-C promotes eye fate and suppresses antennal fate
(Figures 7J-K, S5G, and Table S5). Therefore, Bon and Yki control cell fate decisions in the eye
through transcriptional regulation of a unique set of target genes: activation of *sha* and *f*, and
repression of *E(spl)-C* genes.

Discussion

We have revealed a previously unappreciated and profound role of the Hippo pathway in
controlling cell fate determination in the *Drosophila* eye (Figure 7L). This function depends on the
activity of the novel Yki interactor, Bon. We have uncovered a collaborative regulatory program in
which Yki and Bon interact, and likely function within larger multiprotein complexes that include
other transcriptional regulators. Rather than controlling growth or PNS differentiation, which
represent the previously ascribed independent functions of Yki and Bon, the Yki-Bon module
regulates the proper segregation of the eye, epidermal, and antennal fates in the developing eye.
This function involves the promotion of the epidermal and antennal fates, and the suppression of
the eye fate, via transcriptional regulation of a distinct set of target genes (Figure 7L). Our study
thus provides a molecular mechanism for the biological function of the Hippo pathway and Bon in
cell fate determination during eye development.
391

Hippo pathway and Bon control eye-antenna-epidermis cell fate decisions at two levels
Our results suggest that the Hippo pathway and Bon regulate the developmental cell fate
decisions in the eye at two levels (Figure S7). First, the Yki-Bon complex promotes antennal and
epidermal fates and suppresses the eye fate during the early eye field specification, before the
L3 larval stage. This is supported by the phenotypes observed under various genetic
manipulations of Bon, Wts, Yki, and Sd during the L1 and L2 larval stages, including the reciprocal
transformations of eye and antenna, epidermal outgrowths in the eye, and ectopic eye fate
(Figures 3, 4, 6, S4, and S6). The Yki-Bon module is thus an essential component of the extensive

network of signaling pathways and transcription factors that control these cell fate decisions in early eye development (Kumar and Moses, 2001). Previous studies showed that *ex*, *mer* and *mats* mutants exhibit eye to epidermal transformation and occasionally eye to antenna transformation (in *ex* mutants), suggesting that the upstream Hippo pathway may also regulate the Yki-Bon controlled fate determination at this stage (Boedigheimer and Laughon, 1993; Lai *et al.*, 2005; McCartney *et al.*, 2000; Pellock *et al.*, 2007).

Second, after the segregation of the eye/antenna/epidermis fields and the start of MF in L3, the Yki-Bon complex promotes the epidermal cell fate while suppressing ommatidia, whereas Wts counteracts this activity. This is evidenced by the formation of epidermal trichomes on interommatidial cells as well as suppression of ommatidial cell types, especially cone cells, upon knockdown of *wts* or overexpression of Bon or Yki with the late eye driver *GMR-GAL4* (Figures 2, 3, and S3). Furthermore, our RNA-seq data also revealed cell fate regulation at the gene expression level: the Yki-Bon complex activates epidermal differentiation genes (*sha*, *f*, *nyo* and *neo*), and represses multiple Notch targets (*E(spl)-C*, *ct* and *TfAP-2*) that are required for eye fate establishment and are expressed in ommatidial cells (e.g., *ct* in cone cells and *E(spl)mdelta-HLH* in primary pigment cells) (Figure 6). Although activation of the Yki-Bon complex at this stage did not exhibit an eye-to-antenna transformation phenotype, certain antennal genes (*Gr64s* and *Or47b*) were upregulated, suggesting transformation at the level of gene expression.

The eye-antenna-epidermis fate determination was previously studied during the early stages of eye-antennal disc development (Kumar, 2018). Our work shows that these fates are not completely defined during the early stages, as the retina could still transform into epidermal tissue and express epidermal and even antennal genes when the Hippo pathway and Bon were modulated after MF formation. Interestingly, conditional knockout of *eya* after the MF results in suppression of ommatidia and formation of trichomes in the eye (Jin et al., 2016). This suggests that the retinal determination genes are also involved in eye-epidermal fate decisions during later stages of eye development, and that trichome induction may be a general biological outcome of interference with the eye vs. epidermis specification after the start of MF. Thus, we conclude that the eye is still developmentally plastic at late stages, with a latent epidermal fate that is normally inhibited. Interestingly, this fate is revealed in the insect order Strepsiptera, whose compound eyes are comprised of optical units that are separated by epidermal tissue bearing trichome-like extensions (Buschbeck, 2005).

Yki and Bon control cell fate decisions in the eye by recruiting cofactors and regulating a distinct set of target genes

We have uncovered an unexpected layer of control over eye specification exerted by Yki and Bon at the level of Notch target genes. The Hippo pathway has been reported to control cell fate determination in other biological contexts through regulation of Notch ligands (Reddy et al., 2010; Totaro et al., 2017). Although we have identified several Notch targets that are repressed by Bon and Yki, Notch itself and the key Notch pathway components, Serrate, Delta, and Su(H), were not transcriptionally regulated or found in Yki or Bon protein interactomes (Tables S1-S3). Therefore, we propose that during cell fate determination in the eye, Bon and Yki repress Notch targets such as *E(spl)-C* genes independently from upstream Notch signaling. We note that not all *E(spl)-C* genes are under Bon and Yki control, implying context-dependent regulation and functional divergence of *E(spl)-C* genes, as suggested by previous studies (Table S3) (Couturier *et al.*, 2019; Ligoxygakis et al., 1999). Both Hippo and Notch contribute to cell proliferation and growth of the eye (Dominguez and Casares, 2005; Peng et al., 2009). Our data suggest that Bon is not required for the growth regulation function of the Hippo pathway (Figure S1). Instead, the interaction between Bon and Yki is specifically required for controlling eye specification through regulation of Notch targets. We speculate that Bon may function as a switch that redirects some of Hippo pathway activities from growth regulation to cell fate determination.

So far, *Drosophila* Yki has only been implicated in transcriptional activation (Zheng and Pan, 2019). However, studies in mammalian systems have shown that the YAP/TAZ-TEAD complex (Yki-Sd orthologs) can also function as a transcriptional repressor on non-canonical target genes (Beyer et al., 2013; He et al., 2020; Kim et al., 2015). The direct repression of Notch targets reported here suggests that *Drosophila* Yki can also function in transcriptional repression, likely via recruitment of corepressors mediated by Bon. HDAC1 and its associated corepressor complexes repress gene transcription (including Notch targets) via epigenetic modifications (Borggrefe and Oswald, 2009). We identified HDAC1 and its corepressor, CoRest, in the Bon interactome (Figure 5A), raising the possibility that Bon and Yki repress Notch target genes in part via recruitment of this repressor complex. The involvement of epigenetic regulators is further exemplified by Su(var)2-10, which has a role in chromatin SUMOylation and piRNA target gene silencing (Ninova *et al.*, 2020). Interestingly, Su(var)2-10 can suppress eye fate and even induce antennal fate in a sensitized background (Betz *et al.*, 2001). Due to the strong genetic interaction between Su(var)2-10 and the Bon-Yki complex (Figures 5 and S5), it is possible that chromatin

467 SUMOylation is involved in gene repression by Bon and Yki. Future studies of chromatin status
468 and epigenetic markers may reveal the mechanistic details of gene repression by Bon and Yki.
469
The Hippo pathway and Bon orthologs are conserved and broadly expressed in higher eukaryotes
(Cammas *et al.*, 2012; Zheng and Pan, 2019), raising the possibility that they may also function
together in other species and tissue contexts. The vertebrate Hippo pathway controls
retinogenesis and the cell fate decision between the neuronal retina and non-neuronal retinal
pigment epithelium, which shares similarities with *Drosophila* interommatidial cells (Charlton-
Perkins and Cook, 2010; Lee et al., 2018). It is possible that the TIF1 family proteins are also
involved in this process. The Hippo pathway and TIF1γ have been independently implicated in
controlling hematopoiesis (Lundin et al., 2020; Milton et al., 2014; Rossmann et al., 2021). The
YAP/TAZ-TEAD complex or TIF1β can recruit the transcriptional corepressor NuRD complex for
gene repression in various biological contexts (Beyer *et al.*, 2013; He *et al.*, 2020; Kim *et al.*, 2015;
Schultz et al., 2001). These studies and our data suggest that the biological functions controlled
by the Hippo pathway and Bon and the underlying molecular mechanism we report here may be
evolutionarily conserved.

Acknowledgments
We thank Spyros Artavanis-Tsakonas, Hugo Bellen, Kwang-Wook Choi, Claude Desplan, Maxim
Frolov, Jin Jiang, Eric Lai, Duojia Pan, François Schweisguth, Alexei Tulin, BDSC, VDRC, DGRC
and DSHB for fly stocks and reagents. We thank Georg Halder, Kieran Harvey, Tony Ip, Kenneth
Irvine, Ruth Johnson, Donald Ready, Nicolas Tapon, Barry Thompson, Can Zhang and Lei Zhang
for helpful discussions. We thank Alexander Letizia and Katerina Heath for helping with plasmid
construction. We also thank Jens Rister and Claire Jackan for helpful comments on the
manuscript. Part of the SEM work was supported by Award #S10RR021043 from the National
Center for Research Resources to the UMass Chan EM Core Facility. This work was funded by
NIH grants GM105813 and GM123136 to K.H.M. and A.V.

Author contributions
492 Conceptualization, H.Z., K.H.M., and A.V.; Methodology, H.Z., K.H.M., and A.V.; Formal analysis,
493 H.Z.; Investigation, H.Z.; Resources, K.H.M. and A.V.; Data Curation, H.Z.; Writing – Original
494 Draft, H.Z.; Writing – Review & Editing, H.Z., K.H.M., and A.V.; Visualization, H.Z.; Supervision,
495 A.V.; Funding Acquisition, K.H.M. and A.V.

Declaration of interests

496   The authors declare no competing interests.

Figure Legends

Figure 1. Identification of Bonus as a Yorkie interactor

(A) The Yki protein interaction network. AP-MS results of Yki-SBP from *Drosophila* S2 cells and

Yki-EGFP from embryos were analyzed with Significance Analysis of INTeractome (SAINT) (Choi

et al., 2011). All the interactors (nodes) in this network were identified as significant Yki interactors

(SAINT score ≥ 0.8); these interactions (edges) are not shown to avoid clutter. The shown edges

were incorporated from the STRING database and FlyBase, and include various types of

evidence (Szklarczyk et al., 2019; Thurmond et al., 2019). Also see Method Details.

(B) Schematic diagram showing the major domains and motifs in Yki and Bon. Note two WW

domains in Yki and two PPxY motifs in Bon.

(C) Co-IP of Bon-V5 and wild-type or mutant Yki-HA expressed in S2 cells.

(D) Co-IP of Yki-HA and wild-type or mutant Bon-V5 expressed in S2 cells.

Figure 2. Activation of Bon or Yki induces epidermal trichomes in adult eyes

(A-G, I-N) SEM images of adult *Drosophila* eyes expressing indicated *UAS* transgenes with the

GMR-GAL4 driver. Crosses with Bon were carried out at 25°C, and crosses with Yki, Yki-S168A

and *wts-RNAi* were set up at 25°C and shifted to 29°C after the emergence of first instar larvae.

Arrowheads: interommatidial bristles, arrows: trichomes. Scale bars in left panels: 100 μm;

enlarged views in right panels: 10 μm.

(H, O-R) Quantification of trichome numbers in A-G (H) and I-N (O-R). *GFP* was coexpressed for

transgene dosage compensation. Trichome numbers here and in the other figures were obtained

in an area of 1306 μm^2. Two additional *bon-RNAis* were used that significantly reduced trichome

numbers, with images and quantifications given in Figure S2 and Table S5, respectively. Images

for other quantified genotypes are shown in Figure S2. The number of trichomes and the number

of images quantified (n ≥ 3) for each genotype are provided in Table S5. Statistical analyses are

described in Method Details.

Figure 3. Yki and Bon promote the epidermal fate at the expense of the eye fate

(A, C) Pupal eyes expressing the indicated transgenes with *GMR-GAL4* were stained with phalloidin for F-actin and anti-disc large (Dlg) antibody for cell boundaries, or anti-Cut (Ct) antibody for cone cells. (A) 44 hrs APF grown at 25°C, (C) 40 hrs APF grown at 29°C (equivalent to 48 hrs APF at 25°C). Arrowheads: interommatidial bristles and sockets, arrows: trichomes and the corresponding cells, dashed circles in C: individual cone cell clusters per ommatidium. Scale bars: 10 μm.

(B, D) Quantification of cone cell numbers per area in (A) or per ommatidium in (C), respectively. Cone cell counts in (B) were done in an area of 1×10^4 μm^2 instead of per ommatidium due to a severe phenotype. (D) The *p* values were determined using Fisher's exact test between ommatidia with ≥4 cone cells (CC) and those with <4 CC per ommatidium. The numbers are provided in Table S5.

(E-F) Pupal eyes at 46 hrs APF with mosaic clones of the indicated genotypes were immunostained with anti-Hth antibody for head epidermis, anti-Dlg antibody for cell boundaries, and phalloidin for F-actin. Mutant clones were generated with heat shock at 39 hrs after egg deposition (AED) and marked by loss of GFP. Enlarged views of the boxed regions are shown in the bottom panels. Arrows: trichomes and the corresponding cells in wild type (WT) head cuticle (E) or ectopic epidermis in the eye (*wts*^{X1} clone in F). Scale bars: 50 μm. Exact genotypes of mutants and reporters here and in other figures are given in Method Details.

(G) Pupal eye at 46 hrs APF with mosaic clones of *wts*^{X1} immunostained with anti-GFP antibody to mark the cells outside the clones and anti-Elav antibody for neuronal eye fate. Mutant clones were generated with heat shock at 42 hrs AED. Scale bars: 50 μm.

(H) Representative SEM image of adult eye with mosaic clones of *wts*^{X1}. Mutant clones were generated with heat shock at 39 hrs AED. Enlarged view of the boxed region is shown in the right panel. Arrow: trichome. Scale bars: 20 μm.

(I) L3 eye disc with mosaic clones of the indicated genotypes immunostained with anti-GFP and anti-Elav antibodies. Mutant clones were generated with *eyFLP*. Orthogonal sections (Z) at the dashed lines of basal/apical (2D) views are shown in the left panels. Arrows: the ectopic eye (Elav) and the corresponding clone (loss of GFP) in the basal layer of the *bon*^{21B} clone. The orthogonal sections and their scale bars were scaled 2x along the z axis for easier visualization. Scale bars: 5 μm in Z views and 50 μm in 2D views.

Figure 4. Yki and Bon promote antennal fate and suppress eye fate

(A-D') L3 eye-antennal discs expressing indicated transgenes with *ey-GAL4* were immunostained with anti-Hth antibody for the antennal field. Expression region of the driver is visualized by GFP

(A). (A'-D') Schematic illustrations for the structure of the eye-antennal disc and the pattern of Hth

staining in A-D. Arrows: eye discs, arrowheads: antennal discs. Scale bars: 50 µm.

(E-H) Representative SEM images of adult heads expressing the indicated transgenes with *ey-*

GAL4. (F) *wts-RNAi* had double antennae with triple aristae and fused a1 and a2 segments. (G)

Yki-S168A had double or triple antennae with one arista each and an ectopic maxillary palp. Right

panels: enlarged views of the phenotypes, arrows: eyes, white arrowheads: antennal a3

segments, yellow arrowheads: aristae, white mp and yellow mp: endogenous and ectopic

maxillary palps, respectively. Scale bars: 50 µm.

(I) Quantification of the phenotypes in eye-antennal discs shown in A-D, S4A-S4F and S4I-S4K.

P values were determined using Fisher's exact test between normal and abnormal eye disc

populations within indicated comparisons. The numbers are provided in Table S5.

(J-J') L3 eye-antennal disc expressing *bon-mCherry* with *ey-GAL4-2* immunostained with anti-Ct

antibody for the antennal field and anti-Elav antibody for the neuronal eye fate. mCherry and Ct

signals were colored green and red, respectively. (J') Schematic illustration of J. Control with GFP

expressed under the same driver is shown in S4O-O'. Quantifications are provided in Table S5.

Scale bars: 50 µm.

(K-L) Representative SEM images of adult heads expressing the indicated transgenes with *ey-*

GAL4. Right panels: enlarged views of the phenotypes, arrow: eye, white arrowhead: antennal a3

segment, yellow arrowhead: arista, eo: epidermal outgrowth. Quantifications are provided in Table

S5. Scale bars: 50 µm.

(M-O') L3 eye-antennal discs expressing indicated transgenes with *hth-GAL4* were

immunostained with anti-Ct and anti-Elav antibodies. Expression region of the driver in disc proper

is shown with GFP (M). Representative moderate (N) and severe (O) antenna-to-eye

transformations are shown. (M'-O') Schematic illustrations of M-O. Scale bars: 50 µm.

(P) Quantification of the phenotypes in eye-antennal discs shown in M-O and S4L-S4N. P values

were determined using Fisher's exact test between normal and abnormal antennal disc

populations (*GFP* vs. *yki-RNAi-1*) or between severe transformation and non-severe phenotypes

(*yki-RNAi-1 + bon-RNAi-1* vs. *yki-RNAi-1*). The numbers are provided in Table S5.

(Q) Summary model showing the function of Wts, Yki, Sd and Bon during cell fate decisions in

the eye: the Yki-Bon complex promotes epidermal fate and antennal fate, and suppresses eye

fate.

Figure 5. Interactors of Bon and Yki are required for the eye-epidermis fate decision

(A) Bon protein interactome showing significant interactors (SAINT score ≥ 0.8) identified from AP-MS of Bon-SBP in S2 cells. Interactors highlighted with colors were tested in genetic experiments.

(B-K) SEM images of adult eyes expressing indicated *UAS* transgenes with the *GMR-GAL4* driver. *w-RNAi-1* was coexpressed for transgene dosage compensation. Crosses with Bon were carried out at 25°C, and crosses with Yki-S168A were set up at 25°C and shifted to 29°C after the emergence of first instar larvae. Scales bars in left panels: 100 µm; enlarged views in right panels: 10 µm.

(L) Quantification of the trichome numbers for the indicated genotypes in B-K and S5B-C.

(M) Pupal eyes at 44 hrs APF expressing the indicated transgenes with *GMR-GAL4* were stained with phalloidin for F-actin and anti-Dlg antibody for cell boundaries, or anti-Ct antibody for cone cells. Scale bars: 10 µm.

(N) Quantification of cone cell numbers per 1×10^4 µm^2 area in M. Dashed line indicates the mean cone cell number in WT (*GMR>mCherry*) shown in Figures 3A-B. The numbers are provided in Table S5.

(O) Quantification of the phenotypes in eye-antennal discs shown in P-S. P values were determined using Fisher's exact test between normal and loss-of-Elav populations within indicated comparisons. The numbers are provided in Table S5.

(P-S) L3 eye-antennal discs expressing indicated transgenes with the *bi-GAL4* driver were immunostained with anti-Elav antibody. Enlarged views of the dorsal margins (boxed) of the eye discs are shown in the bottom panels. Arrows: Elav expression at the dorsal margins. Scale bars: 50 µm.

Figure 6. Bon and Yki jointly regulate a set of non-canonical transcriptional target genes

(A) Differential gene expression analysis of RNA-seq data from 40-41 hrs pupal eyes. Output from DEseq2 (Love et al., 2014) was plotted using $\log_2$ fold changes (FC) for *GMR>bon-mCherry + yki-RNAi-1* vs. *GMR>bon-mCherry* (y axis) and that for *GMR>bon-mCherry* vs. *GMR>mCherry* (WT) (x axis). Dashed lines: FC = 1.5 in both directions. Red circles: Bon-activated/Yki-dependent genes. Blue circles: Bon-repressed/Yki-dependent genes. All colored circles fulfill *p* adj. ≤ 0.05. Solid circles with labels: genes of interest. Labels underlined: genes validated with qRT-PCR.

(B) Pearson correlation coefficient analysis of genes jointly regulated by Bon and Yki. Axes are the same as in A. All genes with FC ≥ 1.5 in both directions and *p* adj. ≤ 0.05 were included in the analysis.

(C) Terms of interest from GO_BP analysis of Yki-dependent Bon-activated (red) and -repressed (blue) genes. Complete GO_BP analysis is provided in Table S4.

(D) qRT-PCR validation for the genes of interest using 40-41 hrs APF pupal eyes with the genotypes *GMR>mCherry* (ctrl), *GMR>bon-mCherry*, and *GMR>bon-mCherry + yki-RNAi-1*.

(E) A Venn diagram showing the overlaps of binding sites among Yki, Bon and Sd. All pairwise overlaps have *p* < 0.0001. Previously published datasets used in the analysis were: Yki ChIP-seq from wing discs, Bon ChIP-seq from embryos, and Sd DamID-seq from larval CNS (Negre *et al.*, 2011; Oh *et al.*, 2013; Vissers *et al.*, 2018).

(F) Schematic diagram showing a genomic region around the transcription start site (arrow) of *E(spl)mdelta-HLH* (-2.2 kb to +0.5 kb). Yki and Sd binding regions from previous publications are indicated (Oh *et al.*, 2013; Vissers *et al.*, 2018). P1 and P2: amplicons for ChIP-qPCR.

(G) ChIP-qPCR analysis for the direct binding of Bon to the two genomic loci (P1 and P2) shown in F, using L3 eye-antennal discs from *GMR>bon-mCherry*.

(H) L3 eye-antennal discs with mosaic clones of the indicated genotypes were immunostained with anti-GFP antibody for *GFPmδ* reporter and anti-Elav antibody for neuronal eye fate. Mutant clones were generated with *eyFLP* and marked by loss of mRFP. Enlarged views of the boxed regions in *wts^{X1}* clone are shown in the bottom panels. Scale bars: 50 μm.

(I) Pupal eyes at 25 hrs APF of control (*DE>w-RNAi*) and *bon^{21B}* mosaic clones were immunostained with anti-GFP antibody for *GFPmδ* reporter and anti-Dlg antibody for cell boundaries. Mutant clones were generated with *eyFLP* and marked by loss of mRFP. Dashed circle: individual ommatidium, arrowheads: ectopic expression of *GFPmδ* in *bon^{21B}* clone. Scale bars: 10 μm.

(J) L3 eye-antennal discs expressing indicated transgenes with *DE-GAL4*. The expression region of the driver is shown with GFP in the top panel. Signal from *m3-GFP* reporter was amplified by immunostaining with anti-GFP antibody. Bracket: loss of *m3-GFP* in the dorsal compartment of *DE>wts-RNAi*. Scale bars: 50 μm.

(K-L) L3 eye-antennal discs expressing indicated transgenes with *DE-GAL4* were immunostained with anti-GFP antibody for *m3-GFP* reporter and anti-Elav antibody for neuronal eye fate. Left and middle panels are focused at the level of endogenous peripodial epithelium (PE). Middle panels: enlarged views of the boxed regions, right panels: orthogonal sections at the dashed lines, brackets: gain of *m3-GFP* and Elav in the PE layer of the dorsal compartment. The orthogonal sections and their scale bars were scaled 2x along the z axis for easier visualization. Scale bars: 5 μm in orthogonal views, 50 μm in others.

Figure 7. Bon and Yki control the eye-antenna-epidermis fate determination through their

joint transcriptional targets.

(A-H) SEM images of adult eyes expressing indicated *UAS* transgenes with the *GMR-GAL4* driver.

w-RNAi-2 was coexpressed for transgene dosage compensation. Crosses with Bon were carried

out at 25°C, and crosses with Yki-S168A were set up at 25°C and shifted to 29°C after the

emergence of first instar larvae. Scales bars in left panels: 100 µm; enlarged views in right panels:

10 µm.

(I) Quantification of the trichome numbers for the indicated genotypes in A-H and S6K-N.

(J-K') L3 eye-antennal discs expressing indicated transgenes with *ey-GAL4* were immunostained

with anti-Ct and anti-Elav antibodies. (J'-K') Schematic illustrations of J-K. Scale bars: 50 µm.

Quantification of the phenotypes is shown in S5G, and the detailed numbers are given in Table

S5.

(L) Left: independent functions of Yki and Bon in growth regulation and embryonic PNS

development. Right: Yki and Bon jointly regulate cell fate decisions in the eye through recruitment

of transcriptional and post-transcriptional regulators and transcriptional control of a set of non-

canonical target genes.

Methods

Plasmid construction

pMK33-yki-SBP was generated by cloning Yki-RG isoform from pMT-yki-V5 (Zhang *et al.*, 2015)

into pMK33-SBP-C vector (Yang and Veraksa, 2017). pUASTattB-yki-EGFP was generated by

cloning Yki-RG and EGFP tag into pUASTattB vector (Bischof et al., 2007). pMT-yki-HA was

generated by cloning Yki-RG and HA tag into pMT-V5-His vector (Invitrogen, Cat# V412020). The

Y281A, Y350A mutations of WW domains in Yki were generated by Q5 Site-Directed Mutagenesis

Kit (NEB, Cat# E0554S) using pMT-yki-HA as a template. The numbering of the key tyrosine

residues used in this work corresponds with the first reported yki sequence (yki-RH) which has a

stretch of additional 23 aa at the N-terminus (Huang *et al.*, 2005). All tags were added to the C-

terminus of the genes.

pMT-bon-V5, pMK33-bon-SBP, and pUASTattB-bon-mCherry were generated by cloning Bon-

RB isoform from pFlc-1-bon (DGRC, RE48191). The Y507A, Y585A mutations in PPxY motifs of

Bon were generated by the Q5 Site-Directed Mutagenesis Kit. pUASTattB-bon-mCherry and

pUASTattB-bon-Y507A Y585A-mCherry were constructed using NEBuilder HiFi DNA Assembly

Cloning Kit (NEB, Cat# E5520S). All tags were added to the C-terminus of the genes.

pMT-Myc-wts was generated by cloning the coding sequence of wts from pFlag-wts plasmid

kindly offered by Maxim Frolov Lab. Myc tag was added to the N-terminus.

Stable cell lines

S2 cells were transfected with pMK33-yki-SBP and pMK33-bon-SBP constructs using the

Effectene transfection reagent (Qiagen, Cat# 301427). Stable cell lines were selected with

medium containing 300 µg/ml of Hygromycin B (MilliporeSigma, Cat# H3274). Yki-SBP and Bon-

SBP stable cells were induced overnight with 0.07 mM CuSO$_4$, and protein expression was

verified by western blot using anti-SBP antibody (1:1000, Santa Cruz, Cat# sc-101595).

Transgenic fly lines

pUASTattB-yki-EGFP plasmid was injected into the attP2 site in *D. melanogaster* embryos

(Genetic Services, Inc). pUASTattB-bon-mCherry and pUASTattB-bon-Y507A Y585A-mCherry

plasmids were injected into the attP40 site in *D. melanogaster* embryos (Rainbow Transgenic

Flies, Inc). *yw*, Chr. 2 and Chr. 3 balancer stocks were used in standard crossing schemes to

establish the homozygous transgenic lines.

Genetic crosses

To score the trichome phenotype, *UAS-bon-mCherry*, *UAS-yki-EGFP*, *UAS-yki-S168A-1* (*UAS-*

yki.S168A.GFP.HA, BDSC: 28816) and *UAS-wts*-RNAi (VDRC: 111002) were combined with the

eye driver *GMR-GAL4* to generate *UAS-bon-mCherry/CyO wg-lacZ; GMR-GAL4/TM6B*, *GMR-*

GAL4 UAS-yki-EGFP/TM6B, *GMR-GAL4 UAS-yki-S168A-GFP-1/TM6B* and *UAS-wts*-RNAi*/CyO*

wg-lacZ; GMR-GAL4/TM6B fly lines, respectively. *GFP*, *w*-RNAi-1, or *w*-RNAi-2 were

coexpressed for transgene dosage compensation.

To achieve higher GAL4 activity, all crosses with *GMR-GAL4 UAS-yki-EGFP*, *GMR-GAL4 UAS-*

yki-S168A-GFP-1, and *UAS-wts*-RNAi *GMR-GAL4* were set up at 25°C and shifted to 29°C after

the emergence of first instar larvae. All the crosses with *UAS-bon-mCherry/CyO wg-lacZ; GMR-*

GAL4/TM6B were carried out at 25°C.

All the crosses using *ey-GAL4* and *ey-GAL4-2* were carried out at 29°C except for the cross of

ey-GAL4 and *UAS-wts*-RNAi which was carried out at 25°C. The cross of *da-GAL4* and *UAS-yki-*

EGFP for embryo collection and affinity purification was carried out at RT. All other crosses were

carried out at 25°C.

Mosaic clones with hsFLP/FRT were generated by heat-shocking first instar larvae with indicated

time after egg deposition (AED) at 37°C for 1 hr. Genotypes used for clones are as follow:

3E: *hsFLP;; FRT82B/FRT82B Ubi-GFP*

3F-G: *hsFLP;; FRT82B wts^{X1}/FRT82B Ubi-GFP*

3H: *hsFLP;; FRT82B wts^{X1}/FRT82B Sb63b*

3I ctrl: *eyFLP;; FRT82B/FRT82B Ubi-GFP*

3I *bon^{21B}*: *eyFLP;; FRT82B bon^{21B}/FRT82B Ubi-GFP*

6H ctrl: *eyFLP;; GFPmδ/+; FRT82B/FRT82B Ubi-mRFP*

6H *wts^{X1}*: *eyFLP;; GFPmδ/+; FRT82B wts^{X1}/FRT82B Ubi-mRFP*

6I *bon^{21B}*: *eyFLP;; GFPmδ/+; FRT82B bon^{21B}/FRT82B Ubi-mRFP*

S6D ctrl: *eyFLP;; m3-GFP/+; FRT82B/FRT82B Ubi-mRFP*

S6D *wts^{X1}*: *eyFLP;; m3-GFP/+; FRT82B wts^{X1}/FRT82B Ubi-mRFP*

Affinity Purification

Affinity purification from S2 cells. Blank S2 cells (control) and stable cell lines with 50 ml dense

culture were induced overnight (~16 hrs) with 0.07 mM CuSO$_4$ at 25°C. Cells were collected and

lysed on ice for 20 min using 1 ml of ice-cold Default Lysis Buffer (DLB) (50 mM Tris pH 7.5, 125

mM NaCl, 5% (v/v) glycerol, 0.2% (v/v) IGEPAL, 1.5 mM MgCl$_2$, 25 mM NaF, 1 mM Na$_3$VO$_4$, 1

mM DTT, and 2x cOmplete Protease Inhibitor (MilliporeSigma, Cat# 11836145001, 1 tablet per

25 ml lysis buffer)). The lysate was centrifuged at 16,000 rcf for 20 min at 4°C, and the

supernatant was incubated with 50 μl of packed Streptavidin beads (Thermo Fisher Scientific,

Cat# 53117) for 2 hrs at 4°C with rotation. The beads were then washed 5 times with 1 ml DLB.

Proteins on the beads were eluted with 300 μl DLB containing 2 mM biotin and precipitated by

adding 1/10 volume of 8.74 M Trichloroacetic acid solution (TCA, Fisher Scientific, Cat# BP555-

500). Precipitated proteins were washed once with 500 μl of 0.874 M TCA and four times with

ice-cold acetone. The protein pellet was dried overnight at RT and incubated in 40 μl of 2x SDS

sample buffer at 95°C for 5 min. Samples were assessed by silver staining using NuPAGE 4-12%

Bis-Tris Protein Gel (Thermo Fisher Scientific, Cat# NP0335) and SilverQuest Silver Staining Kit

(Thermo Fisher Scientific, Cat# LC6070). The protein samples for mass spectrometry were
separated on a short SDS-PAGE gel (8% Tris-Glycine) followed by Coomassie blue staining. Two
gel pieces (>75 kDa and < 75 kDa) for each sample were submitted to Taplin Mass Spectrometry
Facility at Harvard Medical School for mass spectrometry analysis, as described (Yang and
Veraksa, 2017).

Affinity purification from embryos. 5 L fly cages with apple juice plates (Yang et al., 2017) were
set up for *yw* (control) and the cross *da-GAL4* x *UAS-yki-EGFP*. Flies were allowed to lay eggs
for 15 hrs at RT, then the apple juice plates containing the embryos were incubated for 3 hrs at
25°C. Approximately 1 g embryos were dechorionated with 50% (v/v) Clorox bleach and washed
with water. Collected embryos were then mixed with 6 ml of ice-cold DLB containing 0.5% (v/v)
IGEPAL and 10 µM MG132 in a glass homogenizer and lysed on ice using 20-30 strokes with a
tight pestle. The lysate was kept on ice for 20 min and centrifuged at 16,000 rcf for 20 min at 4°C.
The supernatant was filtered with pre-chilled 0.45 µm filter and incubated with 50 µl of packed
Pierce Control Agarose Resin (Thermo Fisher Scientific, Cat# 26150) for 30 min at 4°C with
rotation to minimize unspecific binding. The lysate was then incubated with 20 µl of packed GFP-
Trap Agarose (Bulldog Bio, Cat# GTA020) for 2 hrs at 4°C with rotation. After binding, the GFP
beads were washed 5 times with 1 ml DLB containing 0.5% (v/v) IGEPAL and 10 µM MG132,
followed by addition of 40 µl of 4x SDS sample buffer and heating at 95°C for 6 min. The samples
were then assessed by silver staining and the protein samples for mass spectrometry were
prepared and submitted as above.

Mass Spectrometry and data analysis
In-gel trypsin digestion and liquid chromatography/tandem mass spectrometry (nanoLC-MS/MS)
were performed by Taplin Mass Spectrometry Facility at Harvard Medical School. Mass
spectrometry results from two gel pieces for each sample were combined, and the results from
the experimental and control samples were analyzed with Significance Analysis of INTeractome
(SAINT) (v2.5.0) using the total peptides identified for each protein (Choi *et al.*, 2011). Any protein
with the SAINT score (average probability) ≥ 0.8 was considered as a high-confidence interactor
and was included in the interactome. SAINT analysis included the following numbers of samples:
Yki-SBP, four experiments and four controls; Yki-EGFP, three experiments and five controls; Bon-
SBP, two experiments and three controls.

Yki protein interactome. Yki protein interactors with SAINT score ≥ 0.8 either in S2 cells or in

embryos were included in the Yki protein interactome. The Yki protein interactome was further

populated with the interactions incorporated from the STRING database (v11.0) and FlyBase

(vFB2020_04) (Szklarczyk *et al.*, 2019; Thurmond *et al.*, 2019). All proteins in the Yki interactome

from the current study were imported into the STRING database and analyzed with default

settings: full network, medium confidence 0.4 and all active interaction sources (Textmining,

Experiments, Databases, Co-expression, Neighborhood, Gene Fusion and Co-occurrence). The

summary of physical interactions of Yki in FlyBase was selected to show neighbor-neighbor

interactions, and the summary of genetic interactions was selected to show both suppression and

enhancement. The interaction data from FlyBase were exported through esyN (Bean et al., 2014),

while only the interactions between the proteins that were identified as components of the Yki

protein interactome from the current study were incorporated into the Yki protein interactome. The

Yki protein interactome was visualized with Cytoscape (Shannon et al., 2003). Nodes represent

Yki and Yki interactors identified in the current study. Edges represent the interactions

incorporated from STRING and FlyBase. Clustering was done manually based on FlyBase

annotations and publications (Thurmond *et al.*, 2019). Gene symbols were updated to FlyBase

version FB2020_04, released Aug 18, 2020.

Bon protein interactome. Bon protein interactors with SAINT score ≥ 0.8 from S2 cells were

included in the Bon protein interactome which was visualized with Cytoscape. The nodes and

edges represent the interactors and interactions identified in the current study, respectively. Gene

symbols were updated to FlyBase version FB2020_04, released Aug 18, 2020.

Co-Immunoprecipitation

S2 cells were transfected with indicated plasmids or blank pMT-V5-His vector (Invitrogen, Cat#

V412020) using the Effectene transfection reagent (Qiagen, Cat# 301427). 24 hrs after

transfection, cells were induced with 0.35 mM $CuSO_4$ overnight at 25°C, collected and lysed on

ice for 20 min with 600 µl of ice-cold DLB. The lysate was centrifuged at 16,000 rcf for 20 min at

4°C. 40 µl supernatant was mixed with 20 µl of 4x SDS sample buffer and heated at 95°C for 6

min to generate lysate samples. The rest of the lysate was incubated with 20 µl of packed anti-

V5 beads (MilliporeSigma, Cat# A7345) or anti-HA beads (MilliporeSigma, Cat# E6779) for 2 hrs

at 4°C. The beads were then washed 3 times (or 4 times for co-IP with Myc-wts) with 1 ml DLB,

mixed with 40 µl of 4x SDS sample buffer, and heated at 95°C for 6 min to generate IP samples.

The lysate samples and IP samples were subjected to SDS-PAGE followed by western blot using

Odyssey Blocking Buffer (PBS) (LI-COR Biosciences, Cat# 927-40003) as blocking buffer, mouse

anti-V5 antibody (1:1000, MilliporeSigma, Cat# V8012), Rabbit anti-HA antibody (1:1000,

MilliporeSigma, Cat# H6908) and Mouse anti-Myc antibody (1:1000, Cell Signaling, Cat# 2276S)

as primary antibody, and Goat anti-Mouse IgG (1:20,000, LI-COR Biosciences, Cat# 926-68070)

and Donkey anti-Rabbit IgG (1:20,000, LI-COR Biosciences, Cat# 926-32213) as secondary

antibody. Blotting membranes were scanned with LI-COR Odyssey CLx Imaging Systems.

EdU incorporation assay

EdU incorporation assay was performed using Click-iT® EdU Imaging Kit (Thermo Fisher

Scientific, C10086) with adapted procedures from (Gouge and Christensen, 2010). Eye discs

were dissected in unsupplemented Schneider's *Drosophila* Medium and incubated in 100 μM EdU

for 20 min in the dark at RT. Discs were then washed three times with 1x PBS, fixed in 3.7% (v/v)

Formaldehyde (MilliporeSigma, Cat# F8775) in 1x PBS for 15 min in the dark at RT, followed by

three washes with 1x PBS and permeabilization in PB-Triton (1x PBS containing 0.3% (v/v) Triton

X-100) for 20 min at RT. Discs were then washed twice with 3% BSA in 1x PBS and incubated in

Click-iT reaction cocktail containing Alexa Fluor 488 (for *bon-mChery*) or 594 (for *yki-S168A-GFP*)

for 30 min in the dark at RT. After washing once with 3% BSA in 1x PBS and once with 1x PBS,

discs were mounted in ProLong Gold Antifade Mountant with DAPI (Thermo Fisher Scientific,

Cat# P36931). All Incubation and washing were carried out on a platform shaker with mild shaking.

Immunostaining, fluorescence and bright-field microscopy

Antibodies and phalloidin. All antibodies and phalloidin reagents used in this study are listed in

the Key Resources Table. All primary antibodies from DSHB were used at 1:50 (v/v) dilution. All

primary and secondary antibodies and phalloidin from Thermo Fisher Scientific were used at

1:500 (v/v) dilution. Mouse anti-β-Gal antibody (Promega, Cat# Z3783) was used at 1:100 (v/v)

dilution. Guinea Pig anti-Bon antibody was used at 1: 5000 (v/v) dilution (Beckstead *et al.*, 2001).

Guinea Pig anti-Hth antibody was used at 1:1000 (v/v) dilution (Ozel et al., 2021). Rabbit anti-

BarH1 was used at 1:500 (v/v) dilution (Kang et al., 2014). Phalloidin 405 (Biotium, Cat# 00034-

T) was used at 1:100 (v/v) dilution.

Embryo staining. Stage 16 embryos were collected and dechorionated with 50% (v/v) Clorox

bleach, rinsed with water, and fixed in 20 ml of fixative containing 4% (v/v) paraformaldehyde

(Electron Microscopy Sciences, Cat# 15710), 25% (v/v) Heptane and 1x PBS for 20 min at RT. 8

ml of methanol was added to remove the vitelline membrane. Fixed and devitellinized embryos

were washed three times with 1.4 ml methanol, four times with 1.4 ml ethanol and twice with PBT

(1x PBS containing 0.1% (v/v) Tween 20). Embryos were subsequently incubated in 1 ml of

blocking solution (1:1 (v/v) of Western Blocking Reagent (Roche, Cat#11921681001) and PBT)

for 2 hrs at RT and incubated overnight in 500 µl of blocking solution containing 22C10 primary

antibody (DSHB, Cat# 22c10) at 4°C. Embryos were then washed five times with 0.1% BSA (m/v

in PBT) and incubated in 1 ml of blocking solution for 1 hr at RT. The embryos were then incubated

in secondary antibody diluted in blocking solution for 2 hrs in the dark at RT, washed sequentially

in 0.1% BSA, PBT and 1x PBS, and mounted with ProLong Gold Antifade Mountant with DAPI.

All incubation and washing were carried out on a nutator.

L3 imaginal disc staining. 3rd instar larval wing discs and eye-antennal discs were dissected in

ice-cold 1x PBS and were fixed in 3.7% (v/v) formaldehyde for 15-20 min at RT. The discs were

washed three times with 1x PBS, permeabilized with PB-Triton for 20 min at RT and were

incubated in primary antibodies diluted in blocking solution (1:1 (v/v) of Western Blocking Reagent

and PBT) overnight at RT. The discs were then washed four times with PBT, incubated in

secondary antibodies diluted in blocking solution for 2-3 hrs in the dark at RT, washed four times

with PBT, and mounted with ProLong Gold Antifade Mountant with DAPI. All incubation and

washing were carried out on a platform shaker with mild shaking.

Pupal eye staining. Pupal eye staging, dissection and staining procedures were adapted from

(Hsiao et al., 2012; Walther and Pichaud, 2006). Newly formed white pupae were circled on the

fly vial. 32-46 hrs after puparium formation (APF), pupal eyes were dissected with forceps in ice-

cold 1x PBS, transferred to 3.7% (v/v) formaldehyde (MilliporeSigma, Cat# F8775) and fixed for

15-20 min at RT. Fixative was replaced with 1x PBS and washed three times followed by

permeabilizing with PB-Triton for 30 min. The pupal eyes were then incubated in primary

antibodies diluted in PB-Triton overnight at RT, washed four times with PB-Triton, and incubated

in secondary antibody or Phalloidin diluted in PB-Triton for 3 hrs in the dark at RT. The pupal eyes

were then washed four times with PB-Triton and mounted with ProLong Gold Antifade Mountant

(Thermo Fisher Scientific, Cat# P10144). All incubation and washing were carried out on a

platform shaker with mild shaking.

Microscopy. Fluorescent images were acquired using Zeiss LSM 880 Confocal Microscope. For

embryos, 20x objective and 1 Airy Unit (AU) pinhole were used. For L3 wing discs, 20x objective

was used to take a z stack, and maximum intensity projection of the entire disc proper was

performed to generate final images. For L3 eye-antennal discs, 20x or 63x objective was used to
take z stacks with optimal step size, and maximum intensity projection was performed for the
entire disc proper, except for the anti-Ct staining which was focused on the disc proper while
minimizing the inclusion of the cone cell focal plane, or for the orthogonal sections and their
corresponding 2D views which are indicated in the figures and legends. For pupal eyes, 63x
objective and z stack were used with focal plane set to the apical surface, except for the final
images of *GFPmδ* and anti-Elav staining which were generated by maximum intensity projection
of the entire pupal eye unless indicated otherwise. In mispatterned pupal eyes overexpressing
Bon, Ct-positive cone cells and BarH1-positive primary pigment cells were distinguished from
bristle groups which have both Ct and BarH1 expression in the basal nuclei (Beam and Moberg,
2010; Cadigan and Nusse, 1996; Charlton-Perkins and Cook, 2010). Bright-field images of adult
eyes were taken under Zeiss Stemi 2000-C Microscope with an attached-on camera at 50x
magnification. Due to the difference in size, only images from female adult flies were analyzed.
Images were processed with Fiji (Schindelin et al., 2012).
903

Scanning electron microscopy (SEM)
Preparation of adult flies for SEM was adapted from (Pepple et al., 2007; Wolff, 2011). 1-2 days
old adult flies were sequentially dehydrated in 1 ml of 25%, 50%, 75%, 100%, 100% and 100%
ethanol (v/v in water) for 8-16 hrs per step at RT (overnight for the first step). The flies were then
chemically dried in 500 µl of 25%, 50%, 75%, 100%, 100% and 100% Hexamethyldisilazane (v/v
in ethanol) (HMDS, Electron Microscopy Sciences, Cat# 16700) for 30 min per step at RT. Most
of the HMDS was then removed and the remaining HMDS was allowed to be dried under vacuum
in a desiccator containing Indicating Drierite (W.A. Hammond, Cat# 23005) overnight at RT. The
samples were then mounted on Aluminum Mount using Carbon Adhesive Tape, and stored in
Mount Holder (Electron Microscopy Sciences, Cat# 75220, 77816 and 76510). The samples were
subsequently coated with gold at 50 mAmps for 150 secs with Denton Vacuum Desk V sputter
coater or coated with platinum for 8 nm with Denton Vacuum 502-B for samples in Figures S2O-
T. Most images were acquired with JEOL JSM-6010LA IntouchScope Scanning Electron
Microscope using 20 kV voltage, 10 mm working distance, 40-50 spot size, and 150x, 300x and
1000x magnifications for the whole eye, zoomed-in view for ectopic antenna, and zoomed-in view
for trichomes, respectively. Images in Figures S2O-T were acquired with FEI Quanta 200 FEG
MK II Scanning Electron Microscope using 10 kV voltage, 10 mm working distance, 2.5 spot size,
and 150x and 1000x magnifications for the whole eye and zoomed-in view, respectively. Due to
the difference in size, only images from female adult flies were analyzed. Images were processed

with Fiji. The numbers of trichomes were counted with Fiji in an area of 1306 µm^2 (41.73 µm x

31.30 µm) per image for all genotypes.

Total RNA preparation

Total RNA preparation method was adapted from (Zhang *et al.*, 2015). Pupal eyes were dissected

from 40-41 hrs APF pupae (newly formed white pupae were staged for 40 hrs at 25°C and

dissected within 1 hr at RT) in ice cold RNase-free 1x PBS diluted from RNase-free 10x PBS

(Fisher Scientific, Cat# BP3994) with RNase-free water (Thermo Fisher Scientific, Cat#

10977015). Dissected pupal eyes were immediately transferred to 100 µl of TRIzol (Thermo

Fisher Scientific, Cat# 15596026) and kept at RT for 5 min to lyse. Samples in TRIzol were stored

at -80°C until 60 pupal eyes for each genotype were collected. All TRIzol samples for each

genotype were then thawed and combined, and the volume was brought up to 500 µl per sample.

Total RNA was extracted twice by adding 100 µl of chloroform, vigorously shaking for 15 secs,

incubating at RT for 3 min, centrifuging at 11,000 rcf at 4°C for 15 min, and collecting the aqueous

phase. The total RNA was then precipitated by adding 1 volume of 70% ethanol (v/v in RNase-

free water), and purified with RNeasy Mini Kit (QIAGEN, Cat# 74104, 79254).

RNA-sequencing, data analysis, and qPCR validation

The RNA-seq libraries were prepared from 100 µg of total RNA per genotype with NEBNext Ultra

II RNA Library Prep Kit mRNA Isolation Module (NEB, Cat# E7775S, E7490S, E7500S). Quality

control and sequencing of the RNA-seq libraries were performed by GENEWIZ, Inc. using Agilent

TapeStation and Illumina HiSeq 4000 with a 2x150 paired-end configuration.

The adapter sequences were trimmed from raw reads, and reads shorter than 18 nt were removed

by Cutadapt (v2.9) (Martin, 2011). The reads mapped to tRNAs, rRNA, snRNA and snoRNA were

removed by mapping through Bowtie 2 (v2.3.5.1) with very sensitive setting and maximum

fragment size of 1000 in addition to default setting (Langmead and Salzberg, 2012). The tRNA,

rRNA, snRNA and snoRNA reference sequences were obtained from FlyBase (dmel-all-tRNA-

r6.33.fasta and dmel-all-miscRNA-r6.33.fasta) (Thurmond *et al.*, 2019). The remaining reads

were mapped to *D. melanogaster* genome Ensembl_BDGP6 (Flicek et al., 2014) by STAR

(v2.7.0e) (Dobin et al., 2013) with default parameters. Gene counting was achieved by

featureCounts (Subread v1.6.2) with overlapping or polycistronic genes counted as a fraction

(Liao et al., 2014). Differential gene expression analysis was performed with DEseq2 (v1.30.1)

through DEBrowser (v1.16.3) with default settings (Kucukural et al., 2019; Love *et al.*, 2014), and

genes with maximum count fewer than 10 were filtered out. Normalization method used in
DESeq2 was Median Ratio Normalization (MRN) (Maza et al., 2013). The Pearson Correlation
Coefficient analysis was performed by ggpubr R package. The Gene Ontology (GO) analysis was
performed with DAVID Bioinformatics Resources (v6.8) (Huang da et al., 2009a; b).

cDNA was generated from total RNA using SuperScript III First-Strand Synthesis System (Thermo
Fisher Scientific, Cat# 18080051). qPCR was performed with biological and technical triplicates
using PowerUp SYBR Green Master Mix (Thermo Fisher Scientific, Cat# A25742) on QuantStudio
3 Real-Time PCR System. Gene expression was normalized to ribosomal protein *rp49*. P values
were calculated using the values of ΔCq (Cq (gene of interest) - Cq(rp49)) (Yuan et al., 2006).
The sequences of the primers are listed in the Key Resources Table.

DNA binding analysis and ChIP-qPCR
Overlap in the DNA binding locus for Yki, Bon, and Sd was analyzed and visualized by
ChIPpeakAnno using published ChIP-seq datasets of Yki (GSE38594) and Bon (GSE25921) after
converting to dm6 coordinates, and DamID-seq dataset of Sd (GSE120731) (Negre *et al.*, 2011;
Oh *et al.*, 2013; Vissers *et al.*, 2018; Zhu et al., 2010). The p values between every two datasets
were determined by the hypergeometric test using totalTest number for number of potential peaks
(Zhu *et al.*, 2010). The totalTest number here (89144) was calculated by dividing the *Drosophila*
genome (143.7 Mb) (dos Santos et al., 2015) with the largest mean peak width from three datasets
(1612 bp for Sd DamID), assuming that these three proteins can potentially bind anywhere in the
genome. The presence of ChIP peak and DamID peak at Bon-Yki jointly regulated genes was
analyzed using IGV (Robinson et al., 2011).

ChIP assay was performed using Chromatin Immunoprecipitation (ChIP) Assay Kit
(MilliporeSigma, Cat# 17-295) with adaptations based on the application notes on GFP- and RFP-
Trap by ChromoTek. 100 Eye-antennal discs for each replicate were dissected in ice-cold 1xPBS.
Protein and DNA were crosslinked by incubating the discs in 1.8% (v/v) formaldehyde (Thermo
Fisher Scientific, Cat# 28906) in 1x PBS for 15 min at RT and quenched with 225 mM Glycine for
5 min at RT (Oh *et al.*, 2013). Discs were washed three time with 1x PBS, transferred to 200 µl
SDS lysis buffer (per 100 discs) containing 2.5 x EDTA-free mini cOmplete Protease Inhibitor
(MilliporeSigma, Cat# 11836170001, 1 tablet per 4 ml SDS lysis buffer), homogenized with Pellet
Pestle for 30 secs on ice, and incubated for 10 min on ice. The lysate was sonicated using
Bioruptor sonicator for 15 min (30 secs on and 30 secs rest at high power) in 4°C water bath to

shear the DNA. Samples were centrifuged at 16,000 rcf for 10 min at 4°C, and the supernatant
was diluted 10-fold in ChIP Dilution Buffer containing 2.5 x mini cOmplete inhibitor. Samples were
pre-cleared with 75 µl (50% slurry) of Pierce Control Agarose Resin for 1 hr at 4°C with rotation,
and after a brief centrifugation, 5% sample was saved as input sample. The supernatant was then
incubated with 25 µl (50% slurry) of RFP-Trap agarose overnight at 4°C with rotation. The agarose
beads were washed as indicated in the kit. The protein-DNA complex was eluted twice with 250
µl elution buffer for 15 min, first at RT on a nutator and second on a 65°C shaker. Input samples
and elutes were reverse cross-linked with 0.2 M NaCl at 65°C overnight, digested with RNase A
(Thermo Fisher Scientific, Cat# EN0531) for 30 min at 37°C and Proteinase K (Thermo Fisher
Scientific, Cat# 26160) as indicated in the kit. DNA was recovered by phenol/chloroform extraction
and ethanol precipitation.

1002

1003 qPCR was performed with biological triplicates and technical duplicates using PowerUp SYBR
1004 Green Master Mix on QuantStudio 3 Real-Time PCR System. Ratio to the input was presented.
1005 P values were calculated using the values of ΔCq (Cq (ChIPped) - Cq(input)). The sequences of
1006 the primers are listed in the Key Resources Table.

1007

1008

1009 **Quantification And Statistical Analysis**

1010

Statistical significance of categorical variables was determined by Fisher's exact test with two-
sided p value calculation using GraphPad Prism (v9.1.0). Unless indicated otherwise, in all other
statistical analyses, data were presented as mean ± SD of at least three replicates, and the
significance was determined by unpaired t-test with Welch's correction using GraphPad Prism
(v9.1.0). The numbers (n) and values of each sample used per experiment are provided in Table
S5. The p values are presented as follows: ns ($p > 0.05$, not significant), * ($p ≤ 0.05$), ** ($p ≤ 0.01$),
*** ($p ≤ 0.001$), **** ($p ≤ 0.0001$).

1018

1019 **Supplemental Tables**

1020

Table S1. Yki interactome (related to Figure 1).
Table S2. Bon interactome (related to Figure 5).
Table S3. Differential gene expression analysis (related to Figure 6 and S6).
Table S4. GO_BP analysis of joint targets of Bon and Yki (related to Figure 6).

Table S5. Quantifications of phenotypes (related to Figure 2, 3, 4, 5, 6, 7, S1, S2, S3, S4, S5,
and S6).

References

Allton, K., Jain, A.K., Herz, H.M., Tsai, W.W., Jung, S.Y., Qin, J., Bergmann, A., Johnson, R.L., and
Barton, M.C. (2009). Trim24 targets endogenous p53 for degradation. Proc Natl Acad Sci U S A
106, 11612-11616. 10.1073/pnas.0813177106.
Andrews, J., Garcia-Estefania, D., Delon, I., Lu, J., Mevel-Ninio, M., Spierer, A., Payre, F., Pauli,
D., and Oliver, B. (2000). OVO transcription factors function antagonistically in the Drosophila
female germline. Development *127*, 881-892.
Arif, S., Kittelmann, S., and McGregor, A.P. (2015). From shavenbaby to the naked valley:
trichome formation as a model for evolutionary developmental biology. Evol Dev *17*, 120-126.
10.1111/ede.12113.
Barish, S., Li, Q., Pan, J.W., Soeder, C., Jones, C., and Volkan, P.C. (2017). Transcriptional
profiling of olfactory system development identifies distal antenna as a regulator of subset of
neuronal fates. Sci Rep *7*, 40873. 10.1038/srep40873.
Beam, C.K., and Moberg, K. (2010). The gang of four gene regulates growth and patterning of
the developing Drosophila eye. Fly (Austin) *4*, 104-116. 10.4161/fly.4.2.11890.
Bean, D.M., Heimbach, J., Ficorella, L., Micklem, G., Oliver, S.G., and Favrin, G. (2014). esyN:
network building, sharing and publishing. PLoS One *9*, e106035. 10.1371/journal.pone.0106035.
Beckstead, R., Ortiz, J.A., Sanchez, C., Prokopenko, S.N., Chambon, P., Losson, R., and Bellen,
H.J. (2001). Bonus, a Drosophila homolog of TIF1 proteins, interacts with nuclear receptors and
can inhibit betaFTZ-F1-dependent transcription. Mol Cell *7*, 753-765.
Beckstead, R.B., Ner, S.S., Hales, K.G., Grigliatti, T.A., Baker, B.S., and Bellen, H.J. (2005). Bonus,
a Drosophila TIF1 homolog, is a chromatin-associated protein that acts as a modifier of
position-effect variegation. Genetics *169*, 783-794. 10.1534/genetics.104.037085.
Betz, A., Lampen, N., Martinek, S., Young, M.W., and Darnell, J.E., Jr. (2001). A Drosophila PIAS
homologue negatively regulates stat92E. Proc Natl Acad Sci U S A *98*, 9563-9568.
10.1073/pnas.171302098.
Beyer, T.A., Weiss, A., Khomchuk, Y., Huang, K., Ogunjimi, A.A., Varelas, X., and Wrana, J.L.
(2013). Switch enhancers interpret TGF-beta and Hippo signaling to control cell fate in human
embryonic stem cells. Cell Rep *5*, 1611-1624. 10.1016/j.celrep.2013.11.021.
Bischof, J., Maeda, R.K., Hediger, M., Karch, F., and Basler, K. (2007). An optimized transgenesis
system for Drosophila using germ-line-specific phiC31 integrases. Proc Natl Acad Sci U S A *104*,
3312-3317. 10.1073/pnas.0611511104.
Blochlinger, K., Jan, L.Y., and Jan, Y.N. (1993). Postembryonic patterns of expression of cut, a
locus regulating sensory organ identity in Drosophila. Development *117*, 441-450.
Boedigheimer, M., and Laughon, A. (1993). Expanded: a gene involved in the control of cell
proliferation in imaginal discs. Development *118*, 1291-1301.

Bohere, J., Mancheno-Ferris, A., Al Hayek, S., Zanet, J., Valenti, P., Akino, K., Yamabe, Y., Inagaki,
S., Chanut-Delalande, H., Plaza, S., et al. (2018). Shavenbaby and Yorkie mediate Hippo signaling
to protect adult stem cells from apoptosis. Nat Commun *9*, 5123. 10.1038/s41467-018-07569-0.
Borggrefe, T., and Oswald, F. (2009). The Notch signaling pathway: transcriptional regulation at
Notch target genes. Cell Mol Life Sci *66*, 1631-1646. 10.1007/s00018-009-8668-7.
Buschbeck, E.K. (2005). The compound lens eye of Strepsiptera: morphological development of
larvae and pupae. Arthropod Structure & Development *34*, 315-326.
https://doi.org/10.1016/j.asd.2005.04.002.
Cadigan, K.M., and Nusse, R. (1996). wingless signaling in the Drosophila eye and embryonic
epidermis. Development *122*, 2801-2812. 10.1242/dev.122.9.2801.
Cagan, R.L., and Ready, D.F. (1989a). The emergence of order in the Drosophila pupal retina.
Dev Biol *136*, 346-362. 10.1016/0012-1606(89)90261-3.
Cagan, R.L., and Ready, D.F. (1989b). Notch is required for successive cell decisions in the
developing Drosophila retina. Genes Dev *3*, 1099-1112. 10.1101/gad.3.8.1099.
Cambiaghi, V., Giuliani, V., Lombardi, S., Marinelli, C., Toffalorio, F., and Pelicci, P.G. (2012).
TRIM proteins in cancer. Adv Exp Med Biol *770*, 77-91. 10.1007/978-1-4614-5398-7_6.
Cammas, F., Khetchoumian, K., Chambon, P., and Losson, R. (2012). TRIM involvement in
transcriptional regulation. Adv Exp Med Biol *770*, 59-76. 10.1007/978-1-4614-5398-7_5.
Chanut-Delalande, H., Fernandes, I., Roch, F., Payre, F., and Plaza, S. (2006). Shavenbaby
couples patterning to epidermal cell shape control. PLoS Biol *4*, e290.
10.1371/journal.pbio.0040290.
Chanut-Delalande, H., Hashimoto, Y., Pelissier-Monier, A., Spokony, R., Dib, A., Kondo, T.,
Bohere, J., Niimi, K., Latapie, Y., Inagaki, S., et al. (2014). Pri peptides are mediators of ecdysone
for the temporal control of development. Nat Cell Biol *16*, 1035-1044. 10.1038/ncb3052.
Charlton-Perkins, M., and Cook, T.A. (2010). Building a fly eye: terminal differentiation events of
the retina, corneal lens, and pigmented epithelia. Curr Top Dev Biol *93*, 129-173. 10.1016/B978-
0-12-385044-7.00005-9.
Choi, H., Larsen, B., Lin, Z.Y., Breitkreutz, A., Mellacheruvu, D., Fermin, D., Qin, Z.S., Tyers, M.,
Gingras, A.C., and Nesvizhskii, A.I. (2011). SAINT: probabilistic scoring of affinity purification-
mass spectrometry data. Nat Methods *8*, 70-73. 10.1038/nmeth.1541.
Couturier, L., Mazouni, K., Corson, F., and Schweisguth, F. (2019). Regulation of Notch output
dynamics via specific E(spl)-HLH factors during bristle patterning in Drosophila. Nat Commun
10, 3486. 10.1038/s41467-019-11477-2.
Davis, J.R., and Tapon, N. (2019). Hippo signalling during development. Development *146*.
10.1242/dev.167106.
Delon, I., Chanut-Delalande, H., and Payre, F. (2003). The Ovo/Shavenbaby transcription factor
specifies actin remodelling during epidermal differentiation in Drosophila. Mech Dev *120*, 747-
758. 10.1016/s0925-4773(03)00081-9.
Deng, H., Yang, L., Wen, P., Lei, H., Blount, P., and Pan, D. (2020). Spectrin couples cell shape,
cortical tension, and Hippo signaling in retinal epithelial morphogenesis. J Cell Biol *219*.
10.1083/jcb.201907018.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.
10.1093/bioinformatics/bts635.

Dominguez, M., and Casares, F. (2005). Organ specification-growth control connection: new in-
sights from the Drosophila eye-antennal disc. Dev Dyn *232*, 673-684. 10.1002/dvdy.20311.
dos Santos, G., Schroeder, A.J., Goodman, J.L., Strelets, V.B., Crosby, M.A., Thurmond, J.,
Emmert, D.B., Gelbart, W.M., and FlyBase, C. (2015). FlyBase: introduction of the Drosophila
melanogaster Release 6 reference genome assembly and large-scale migration of genome
annotations. Nucleic Acids Res *43*, D690-697. 10.1093/nar/gku1099.
Ellis, M.C., O'Neill, E.M., and Rubin, G.M. (1993). Expression of Drosophila glass protein and
evidence for negative regulation of its activity in non-neuronal cells by another DNA-binding
protein. Development *119*, 855-865.
Fernandes, I., Chanut-Delalande, H., Ferrer, P., Latapie, Y., Waltzer, L., Affolter, M., Payre, F.,
and Plaza, S. (2010). Zona pellucida domain proteins remodel the apical compartment for
localized cell shape changes. Dev Cell *18*, 64-76. 10.1016/j.devcel.2009.11.009.
Flicek, P., Amode, M.R., Barrell, D., Beal, K., Billis, K., Brent, S., Carvalho-Silva, D., Clapham, P.,
Coates, G., Fitzgerald, S., et al. (2014). Ensembl 2014. Nucleic Acids Res *42*, D749-755.
10.1093/nar/gkt1196.
Fujii, S., Yavuz, A., Slone, J., Jagge, C., Song, X., and Amrein, H. (2015). Drosophila sugar
receptors in sweet taste perception, olfaction, and internal nutrient sensing. Curr Biol *25*, 621-
627. 10.1016/j.cub.2014.12.058.
Goodrich, J.S., Clouse, K.N., and Schupbach, T. (2004). Hrb27C, Sqd and Otu cooperatively
regulate gurken RNA localization and mediate nurse cell chromosome dispersion in Drosophila
oogenesis. Development *131*, 1949-1958. 10.1242/dev.01078.
Gouge, C.A., and Christensen, T.W. (2010). Drosophila Sld5 is essential for normal cell cycle
progression and maintenance of genomic integrity. Biochem Biophys Res Commun *400*, 145-
150. 10.1016/j.bbrc.2010.08.033.
Hamaratoglu, F., Willecke, M., Kango-Singh, M., Nolo, R., Hyun, E., Tao, C., Jafar-Nejad, H., and
Halder, G. (2006). The tumour-suppressor genes NF2/Merlin and Expanded act through Hippo
signalling to regulate cell proliferation and apoptosis. Nat Cell Biol *8*, 27-36. 10.1038/ncb1339.
Hari, K.L., Cook, K.R., and Karpen, G.H. (2001). The Drosophila Su(var)2-10 locus regulates
chromosome structure and function and encodes a member of the PIAS protein family. Genes
Dev *15*, 1334-1348. 10.1101/gad.877901.
Hay, B.A., Wolff, T., and Rubin, G.M. (1994). Expression of baculovirus P35 prevents cell death
in Drosophila. Development *120*, 2121-2129.
He, L., Yuan, L., Yu, W., Sun, Y., Jiang, D., Wang, X., Feng, X., Wang, Z., Xu, J., Yang, R., et al.
(2020). A Regulation Loop between YAP and NR4A1 Balances Cell Proliferation and Apoptosis.
Cell Rep *33*, 108284. 10.1016/j.celrep.2020.108284.
Hsiao, H.Y., Johnston, R.J., Jr., Jukam, D., Vasiliauskas, D., Desplan, C., and Rister, J. (2012).
Dissection and immunohistochemistry of larval, pupal and adult Drosophila retinas. J Vis Exp,
4347. 10.3791/4347.
Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009a). Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res *37*, 1-
13. 10.1093/nar/gkn923.
Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009b). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nat Protoc *4*, 44-57.
10.1038/nprot.2008.211.

Huang, J., Wu, S., Barrera, J., Matthews, K., and Pan, D. (2005). The Hippo signaling pathway
coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the Drosophila
Homolog of YAP. Cell *122*, 421-434. 10.1016/j.cell.2005.06.007.
Ito, H., Sato, K., Koganezawa, M., Ote, M., Matsumoto, K., Hama, C., and Yamamoto, D. (2012).
Fruitless recruits two antagonistic chromatin factors to establish single-neuron sexual
dimorphism. Cell *149*, 1327-1338. 10.1016/j.cell.2012.04.025.
Jin, M., Eblimit, A., Pulikkathara, M., Corr, S., Chen, R., and Mardon, G. (2016). Conditional
knockout of retinal determination genes in differentiating cells in Drosophila. FEBS J *283*, 2754-
2766. 10.1111/febs.13772.
Jukam, D., Xie, B., Rister, J., Terrell, D., Charlton-Perkins, M., Pistillo, D., Gebelein, B., Desplan,
C., and Cook, T. (2013). Opposite feedbacks in the Hippo pathway for growth control and neural
fate. Science *342*, 1238016. 10.1126/science.1238016.
Kang, J., Yeom, E., Lim, J., and Choi, K.W. (2014). Bar represses dPax2 and decapentaplegic to
regulate cell fate and morphogenetic cell death in Drosophila eye. PLoS One *9*, e88171.
10.1371/journal.pone.0088171.
Kerber, B., Monge, I., Mueller, M., Mitchell, P.J., and Cohen, S.M. (2001). The AP-2 transcription
factor is required for joint formation and cell survival in Drosophila leg development.
Development *128*, 1231-1238.
Kim, M., Kim, T., Johnson, R.L., and Lim, D.S. (2015). Transcriptional co-repressor function of the
hippo pathway transducers YAP and TAZ. Cell Rep *11*, 270-282. 10.1016/j.celrep.2015.03.015.
Knust, E., Schrons, H., Grawe, F., and Campos-Ortega, J.A. (1992). Seven genes of the Enhancer
of split complex of Drosophila melanogaster encode helix-loop-helix proteins. Genetics *132*,
505-518.
Kucukural, A., Yukselen, O., Ozata, D.M., Moore, M.J., and Garber, M. (2019). DEBrowser:
interactive differential expression analysis and visualization tool for count data. Bmc Genomics
20, 6. 10.1186/s12864-018-5362-x.
Kulkarni, A., Chang, M.T., Vissers, J.H.A., Dey, A., and Harvey, K.F. (2020). The Hippo Pathway as
a Driver of Select Human Cancers. Trends Cancer *6*, 781-796. 10.1016/j.trecan.2020.04.004.
Kumar, J.P. (2010). Retinal determination the beginning of eye development. Curr Top Dev Biol
93, 1-28. 10.1016/B978-0-12-385044-7.00001-1.
Kumar, J.P. (2012). Building an ommatidium one cell at a time. Developmental Dynamics *241*,
136-149. https://doi.org/10.1002/dvdy.23707.
Kumar, J.P. (2018). The fly eye: Through the looking glass. Dev Dyn *247*, 111-123.
10.1002/dvdy.24585.
Kumar, J.P., and Moses, K. (2001). EGF receptor and Notch signaling act upstream of
Eyeless/Pax6 to control eye specification. Cell *104*, 687-697. 10.1016/s0092-8674(01)00265-3.
Kurata, S., Go, M.J., Artavanis-Tsakonas, S., and Gehring, W.J. (2000). Notch signaling and the
determination of appendage identity. Proc Natl Acad Sci U S A *97*, 2117-2122.
10.1073/pnas.040556497.
Kwon, Y., Vinayagam, A., Sun, X., Dephoure, N., Gygi, S.P., Hong, P., and Perrimon, N. (2013).
The Hippo signaling pathway interactome. Science *342*, 737-740. science.1243971 [pii]
10.1126/science.1243971.

Lai, E.C., Bodner, R., and Posakony, J.W. (2000). The enhancer of split complex of Drosophila
includes four Notch-regulated members of the bearded gene family. Development *127*, 3441-
3455.
Lai, E.C., and Rubin, G.M. (2001). Neuralized is essential for a subset of Notch pathway-
dependent cell fate decisions during Drosophila eye development. Proc Natl Acad Sci U S A *98*,
5637-5642. 10.1073/pnas.101135498.
Lai, Z.C., Wei, X., Shimizu, T., Ramos, E., Rohrbaugh, M., Nikolaidis, N., Ho, L.L., and Li, Y. (2005).
Control of cell proliferation and apoptosis by mob as tumor suppressor, mats. Cell *120*, 675-
685. 10.1016/j.cell.2004.12.036.
Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357-359. 10.1038/nmeth.1923.
Lee, M., Goraya, N., Kim, S., and Cho, S.H. (2018). Hippo-yap signaling in ocular development
and disease. Dev Dyn *247*, 794-806. 10.1002/dvdy.24628.
Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.
10.1093/bioinformatics/btt656.
Ligoxygakis, P., Bray, S.J., Apidianakis, Y., and Delidakis, C. (1999). Ectopic expression of
individual E(spl) genes has differential effects on different cell fate decisions and underscores
the biphasic requirement for notch activity in wing margin establishment in Drosophila.
Development *126*, 2205-2214.
Lim, H.Y., and Tomlinson, A. (2006). Organization of the peripheral fly eye: the roles of Snail
family transcription factors in peripheral retinal apoptosis. Development *133*, 3529-3537.
10.1242/dev.02524.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.
Lundin, V., Sugden, W.W., Theodore, L.N., Sousa, P.M., Han, A., Chou, S., Wrighton, P.J., Cox,
A.G., Ingber, D.E., Goessling, W., et al. (2020). YAP Regulates Hematopoietic Stem Cell
Formation in Response to the Biomechanical Forces of Blood Flow. Dev Cell *52*, 446-460 e445.
10.1016/j.devcel.2020.01.006.
Mach, J., Atkins, M., Gajewski, K.M., Mottier-Pavie, V., Sansores-Garcia, L., Xie, J., Mills, R.A.,
Kowalczyk, W., Van Huffel, L., Mills, G.B., and Halder, G. (2018). Modulation of the Hippo
pathway and organ growth by RNA processing proteins. Proc Natl Acad Sci U S A *115*, 10684-
10689. 10.1073/pnas.1807325115.
Martin, M. (2011). Cutadapt Removes Adapter Sequences From High-Throughput Sequencing
Reads. EMBnet.journal *17*, 10-12. 10.14806/ej.17.1.200.
Maza, E., Frasse, P., Senin, P., Bouzayen, M., and Zouine, M. (2013). Comparison of
normalization methods for differential gene expression analysis in RNA-Seq experiments: A
matter of relative size of studied transcriptomes. Commun Integr Biol *6*, e25849.
10.4161/cib.25849.
McCartney, B.M., Kulikauskas, R.M., LaJeunesse, D.R., and Fehon, R.G. (2000). The
neurofibromatosis-2 homologue, Merlin, and the tumor suppressor expanded function
together in Drosophila to regulate cell proliferation and differentiation. Development *127*,
1315-1324.

Meng, Z., Moroishi, T., and Guan, K.L. (2016). Mechanisms of Hippo pathway regulation. Genes Dev *30*, 1-17. 10.1101/gad.274027.115.

Menuz, K., Larter, N.K., Park, J., and Carlson, J.R. (2014). An RNA-seq screen of the Drosophila antenna identifies a transporter necessary for ammonia detection. PLoS Genet *10*, e1004810. 10.1371/journal.pgen.1004810.

Micchelli, C.A., Rulifson, E.J., and Blair, S.S. (1997). The function and regulation of cut expression on the wing margin of Drosophila: Notch, Wingless and a dominant negative role for Delta and Serrate. Development *124*, 1485-1495.

Milton, C.C., Grusche, F.A., Degoutin, J.L., Yu, E., Dai, Q., Lai, E.C., and Harvey, K.F. (2014). The Hippo pathway regulates hematopoiesis in Drosophila melanogaster. Curr Biol *24*, 2673-2680. 10.1016/j.cub.2014.10.031.

Monge, I., Krishnamurthy, R., Sims, D., Hirth, F., Spengler, M., Kammermeier, L., Reichert, H., and Mitchell, P.J. (2001). Drosophila transcription factor AP-2 in proboscis, leg and brain central complex development. Development *128*, 1239-1252.

Morrison, C.M., and Halder, G. (2010). Characterization of a dorsal-eye Gal4 Line in Drosophila. Genesis *48*, 3-7. 10.1002/dvg.20571.

Nagaraj, R., and Banerjee, U. (2007). Combinatorial signaling in the specification of primary pigment cells in the Drosophila eye. Development *134*, 825-831. 10.1242/dev.02788.

Negre, N., Brown, C.D., Ma, L., Bristow, C.A., Miller, S.W., Wagner, U., Kheradpour, P., Eaton, M.L., Loriaux, P., Sealfon, R., et al. (2011). A cis-regulatory map of the Drosophila genome. Nature *471*, 527-531. 10.1038/nature09990.

Ninova, M., Chen, Y.A., Godneeva, B., Rogers, A.K., Luo, Y., Fejes Toth, K., and Aravin, A.A. (2020). Su(var)2-10 and the SUMO Pathway Link piRNA-Guided Target Recognition to Chromatin Silencing. Mol Cell *77*, 556-570 e556. 10.1016/j.molcel.2019.11.012.

Oh, H., and Irvine, K.D. (2008). In vivo regulation of Yorkie phosphorylation and localization. Development *135*, 1081-1088. 10.1242/dev.015255.

Oh, H., Slattery, M., Ma, L., Crofts, A., White, K.P., Mann, R.S., and Irvine, K.D. (2013). Genome-wide association of Yorkie with chromatin and chromatin-remodeling complexes. Cell Rep *3*, 309-318. S2211-1247(13)00015-6 [pii] 10.1016/j.celrep.2013.01.008.

Oh, H., Slattery, M., Ma, L., White, K.P., Mann, R.S., and Irvine, K.D. (2014). Yorkie promotes transcription by recruiting a histone methyltransferase complex. Cell Rep *8*, 449-459. 10.1016/j.celrep.2014.06.017.

Ordway, A.J., Teeters, G.M., Weasner, B.M., Weasner, B.P., Policastro, R., and Kumar, J.P. (2021). A multi-gene knockdown approach reveals a new role for Pax6 in controlling organ number in Drosophila. Development *148*. 10.1242/dev.198796.

Ozel, M.N., Simon, F., Jafari, S., Holguera, I., Chen, Y.C., Benhra, N., El-Danaf, R.N., Kapuralin, K., Malin, J.A., Konstantinides, N., and Desplan, C. (2021). Neuronal diversity and convergence in a visual system developmental atlas. Nature *589*, 88-95. 10.1038/s41586-020-2879-3.

Pai, C.Y., Kuo, T.S., Jaw, T.J., Kurant, E., Chen, C.T., Bessarab, D.A., Salzberg, A., and Sun, Y.H. (1998). The Homothorax homeoprotein activates the nuclear localization of another homeoprotein, extradenticle, and suppresses eye development in Drosophila. Genes Dev *12*, 435-446. 10.1101/gad.12.3.435.

Pan, D. (2010). The hippo signaling pathway in development and cancer. Dev Cell *19*, 491-505.
10.1016/j.devcel.2010.09.011.
Pellock, B.J., Buff, E., White, K., and Hariharan, I.K. (2007). The Drosophila tumor suppressors
Expanded and Merlin differentially regulate cell cycle exit, apoptosis, and Wingless signaling.
Dev Biol *304*, 102-115. 10.1016/j.ydbio.2006.12.021.
Peng, H.W., Slattery, M., and Mann, R.S. (2009). Transcription factor choice in the Hippo
signaling pathway: homothorax and yorkie regulation of the microRNA bantam in the
progenitor domain of the Drosophila eye imaginal disc. Genes Dev *23*, 2307-2319.
10.1101/gad.1820009.
Pepple, K.L., Anderson, A.E., Frankfort, B.J., and Mardon, G. (2007). A genetic screen in
Drosophila for genes interacting with senseless during neuronal development identifies the
importin moleskin. Genetics *175*, 125-141. 10.1534/genetics.106.065680.
Petrera, F., and Meroni, G. (2012). TRIM proteins in development. Adv Exp Med Biol *770*, 131-
141. 10.1007/978-1-4614-5398-7_10.
Pojer, J.M., Manning, S.A., Kroeger, B., Kondo, S., and Harvey, K.F. (2021). The Hippo pathway
uses different machinery to control cell fate and organ size. iScience *24*, 102830.
10.1016/j.isci.2021.102830.
Reddy, B.V., Rauskolb, C., and Irvine, K.D. (2010). Influence of fat-hippo and notch signaling on
the proliferation and differentiation of Drosophila optic neuroepithelia. Development *137*,
2397-2408. 10.1242/dev.050013.
Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26. 10.1038/nbt.1754.
Rossmann, M.P., Hoi, K., Chan, V., Abraham, B.J., Yang, S., Mullahoo, J., Papanastasiou, M.,
Wang, Y., Elia, I., Perlin, J.R., et al. (2021). Cell-specific transcriptional control of mitochondrial
metabolism by TIF1gamma drives erythropoiesis. Science *372*, 716-721.
10.1126/science.aaz2740.
Salzberg, A., Prokopenko, S.N., He, Y., Tsai, P., Pal, M., Maroy, P., Glover, D.M., Deak, P., and
Bellen, H.J. (1997). P-element insertion alleles of essential genes on the third chromosome of
Drosophila melanogaster: mutations affecting embryonic PNS development. Genetics *147*,
1723-1741.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.
Schultz, D.C., Friedman, J.R., and Rauscher, F.J., 3rd (2001). Targeting histone deacetylase
complexes via KRAB-zinc finger proteins: the PHD and bromodomains of KAP-1 form a
cooperative unit that recruits a novel isoform of the Mi-2alpha subunit of NuRD. Genes Dev *15*,
428-443. 10.1101/gad.869501.
Seto, E., and Yoshida, M. (2014). Erasers of histone acetylation: the histone deacetylase
enzymes. Cold Spring Harb Perspect Biol *6*, a018713. 10.1101/cshperspect.a018713.
Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski,
B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of
biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.
Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein-protein association

networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res *47*, D607-D613. 10.1093/nar/gky1131.
Tardi, N.J., Cook, M.E., and Edwards, K.A. (2012). Rapid phenotypic analysis of uncoated
Drosophila samples with low-vacuum scanning electron microscopy. Fly (Austin) *6*, 184-192.
10.4161/fly.20525.
Tare, M., Puli, O.R., Moran, M.T., Kango-Singh, M., and Singh, A. (2013). Domain specific genetic
mosaic system in the Drosophila eye. Genesis *51*, 68-74. 10.1002/dvg.22355.
Thurmond, J., Goodman, J.L., Strelets, V.B., Attrill, H., Gramates, L.S., Marygold, S.J., Matthews,
B.B., Millburn, G., Antonazzo, G., Trovisco, V., et al. (2019). FlyBase 2.0: the next generation.
Nucleic Acids Res *47*, D759-D765. 10.1093/nar/gky1003.
Totaro, A., Castellan, M., Battilana, G., Zanconato, F., Azzolin, L., Giulitti, S., Cordenonsi, M., and
Piccolo, S. (2017). YAP/TAZ link cell mechanics to Notch signalling to control epidermal stem cell
fate. Nat Commun *8*, 15206. 10.1038/ncomms15206.
Udan, R.S., Kango-Singh, M., Nolo, R., Tao, C., and Halder, G. (2003). Hippo promotes
proliferation arrest and apoptosis in the Salvador/Warts pathway. Nat Cell Biol *5*, 914-920.
10.1038/ncb1050.
Vissers, J.H.A., Froldi, F., Schroder, J., Papenfuss, A.T., Cheng, L.Y., and Harvey, K.F. (2018). The
Scalloped and Nerfin-1 Transcription Factors Cooperate to Maintain Neuronal Cell Fate. Cell Rep
25, 1561-1576 e1567. 10.1016/j.celrep.2018.10.038.
Vosshall, L.B. (2001). The molecular logic of olfaction in Drosophila. Chem Senses *26*, 207-213.
10.1093/chemse/26.2.207.
Walther, R.F., and Pichaud, F. (2006). Immunofluorescent staining and imaging of the pupal and
adult Drosophila visual system. Nat Protoc *1*, 2635-2642. 10.1038/nprot.2006.379.
Wang, C.W., and Sun, Y.H. (2012). Segregation of eye and antenna fates maintained by mutual
antagonism in Drosophila. Development *139*, 3413-3421. 10.1242/dev.078857.
Weasner, B.M., and Kumar, J.P. (2013). Competition among gene regulatory networks imposes
order within the eye-antennal disc of Drosophila. Development *140*, 205-215.
10.1242/dev.085423.
Weasner, B.P., and Kumar, J.P. (2022). The early history of the eye-antennal disc of Drosophila
melanogaster. Genetics. 10.1093/genetics/iyac041.
Wernet, M.F., Labhart, T., Baumann, F., Mazzoni, E.O., Pichaud, F., and Desplan, C. (2003).
Homothorax switches function of Drosophila photoreceptors from color to polarized light
sensors. Cell *115*, 267-279. 10.1016/s0092-8674(03)00848-1.
Wittkorn, E., Sarkar, A., Garcia, K., Kango-Singh, M., and Singh, A. (2015). The Hippo pathway
effector Yki downregulates Wg signaling to promote retinal differentiation in the Drosophila
eye. Development *142*, 2002-2013. 10.1242/dev.117358.
Wodarz, A., Hinz, U., Engelbert, M., and Knust, E. (1995). Expression of crumbs confers apical
character on plasma membrane domains of ectodermal epithelia of Drosophila. Cell *82*, 67-76.
10.1016/0092-8674(95)90053-5.
Wolff, T. (2011). Preparation of Drosophila eye specimens for scanning electron microscopy.
Cold Spring Harb Protoc *2011*, 1383-1385. 10.1101/pdb.prot066506.
Xu, T., Wang, W., Zhang, S., Stewart, R.A., and Yu, W. (1995). Identifying tumor suppressors in
genetic mosaics: the Drosophila lats gene encodes a putative protein kinase. Development *121*,
1053-1063.

Yang, L., Paul, S., DuBois-Coyne, S., Kyriakakis, P., and Veraksa, A. (2017). Medium-scale
Preparation of Drosophila Embryo Extracts for Proteomic Experiments. J Vis Exp.
10.3791/55804.
Yang, L., and Veraksa, A. (2017). Single-Step Affinity Purification of ERK Signaling Complexes
Using the Streptavidin-Binding Peptide (SBP) Tag. Methods Mol Biol *1487*, 113-126.
10.1007/978-1-4939-6424-6_8.
Yuan, J.S., Reed, A., Chen, F., and Stewart, C.N., Jr. (2006). Statistical analysis of real-time PCR
data. BMC Bioinformatics *7*, 85. 10.1186/1471-2105-7-85.
Zhang, C., Robinson, B.S., Xu, W., Yang, L., Yao, B., Zhao, H., Byun, P.K., Jin, P., Veraksa, A., and
Moberg, K.H. (2015). The ecdysone receptor coactivator Taiman links Yorkie to transcriptional
control of germline stem cell factors in somatic tissue. Dev Cell *34*, 168-180. S1534-
5807(15)00326-3 [pii]
10.1016/j.devcel.2015.05.010.
Zhang, T., Zhou, Q., and Pignoni, F. (2011). Yki/YAP, Sd/TEAD and Hth/MEIS control tissue
specification in the Drosophila eye disc epithelium. PLoS One *6*, e22278.
10.1371/journal.pone.0022278.
Zheng, Y., and Pan, D. (2019). The Hippo Signaling Pathway in Development and Disease. Dev
Cell *50*, 264-282. 10.1016/j.devcel.2019.06.003.
Zhu, L.J., Gazin, C., Lawson, N.D., Pages, H., Lin, S.M., Lapointe, D.S., and Green, M.R. (2010).
ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. BMC
Bioinformatics *11*, 237. 10.1186/1471-2105-11-237.
1387