Short-term instantaneous prophylaxis and efficient treatment against SARS-CoV-2 in hACE2 mice conferred by an intranasal nanobody (Nb22)

Xilin Wu[1,2]†, Yaxing Wang[3]†, Lin Cheng[4]† , Fengfeng Ni[5,6]†, Linjing Zhu [1,2]†, Sen Ma[3], Bilian Huang[1], Mengmeng Ji[7], Huimin Hu[5,6]，Yuncheng Li[5,6]，Shijie Xu[8], Haixia Shi[8], Doudou Zhang[8], Linshuo Liu[8], Waqas Nawaz[1], Qinxue Hu[5,9]，Sheng Ye[3,10]*, Yalan Liu[5]*, Zhiwei Wu[1,7, 11, 12,13]*

1. Center for Public Health Research, Medical School, Nanjing University, Nanjing, P.R. China.
2. Department of Antibody, Abrev Biotechnology Co., Ltd. Nanjing, P.R. China.
3. Tianjin Key Laboratory of Function and Application of Biological Macromolecular Structures, School of Life Sciences, Tianjin University, Tianjin, P.R. China.
4. Institute for Hepatology, National Clinical Research Center for Infectious Disease, Shenzhen Third People's Hospital, Shenzhen, P.R. China.
5. State Key Laboratory of Virology, Wuhan Institute of Virology, Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan, P.R. China
6. Savaid Medical School, University of Chinese Academy of Sciences, Beijing, P.R. China
7. School of Life Sciences, Ningxia University, Yinchuan, P.R. China.
8. Department of Antibody, Y-clone Medical Science Co. Ltd. Suzhou, P.R. China.
9. Institute for Infection and Immunity, St George's University of London, London, UK
10. Life Sciences Institute, Zhejiang University, Zhejiang, P.R. China.
11. Jiangsu Key Laboratory of Molecular Medicine, Medical School, Nanjing University, Nanjing, P.R. China.
12. State Key Laboratory of Analytical Chemistry for Life Science, Nanjing University, Nanjing, P.R. China.
13. Lead contact

†These authors contributed equally to this work.

*Corresponding author: Z. Wu, E-mail: wzhw@nju.edu.cn, S. Ye, E-mail: sye@tju.edu.cn and Y. Liu, E-mail: liuyl@wh.iov.cn

Mailing address: 22 Hankou Road, Nanjing, Jiangsu 210093. China

Phone: +86 (25) 8368-6092. Fax: +86 (25) 8359-6023.

Declaration of interests: The authors have declared no conflict of interest.

Abstract (234 words)

Current COVID-19 vaccines need to take at least one month to complete inoculation

and then become effective. Around 51% global population are still not fully vaccinated.

Instantaneous protection is an unmet need among those who are not fully vaccinated.

In addition, breakthrough infections caused by SARS-CoV-2 are widely reported. All

these highlight the unmet needing for short-term instantaneous prophylaxis (STIP) in

the communities where SARS-CoV-2 is circulating. Previously, we reported

nanobodies isolated from an alpaca immunized with the spike protein, exhibiting

ultrahigh potency against SARS-CoV-2 and its variants. Herein, we found that Nb22,

among our previously reported nanobodies, exhibited ultrapotent neutralization against

Delta variant with an IC_{50} value of 0.41 ng/ml (5.13 pM). Furthermore, the crystal

structural analysis revealed that the binding of Nb22 to WH01 and Delta RBDs both

effectively blocked the binding of RBD to hACE2. Additionally, intranasal Nb22

exhibited protection against SARS-CoV-2 Delta variant in the post-exposure

prophylaxis (PEP) and pre-exposure prophylaxis (PrEP). Of note, intranasal Nb22 also

demonstrated high efficacy against SARS-CoV-2 Delta variant in STIP for seven days

administered by single dose and exhibited long-lasting retention in the respiratory

system for at least one month administered by four doses, providing a means of

instantaneous short-term prophylaxis against SARS-CoV-2. Thus, ultrahigh potency,

long-lasting retention in the respiratory system as well as stability at room-temperature

make the intranasal or inhaled Nb22 to be a potential therapeutic or STIP agent against

SARS-CoV-2.

Brief summary: Nb22 exhibits ultrahigh potency against Delta variant in vitro and is

exploited by crystal structural analysis; furthermore, animal study demonstrates high

effectiveness in the treatment and short-term instantaneous prophylaxis in hACE2 mice

via intranasal administration.

Keywords: SARS-CoV-2, Delta variant, B.1.617.2, Nanobody, Nb22, STIP, Structure,
Instantaneous prophylaxis, Instantaneous protection

Highlights:

1) Nb22 exhibits ultrapotent neutralization against Delta variant with an IC_{50} value of 0.41 ng/ml (5.13 pM).

2) Structural analysis elucidates the ultrapotent neutralization of Nb22 against Delta variant.

3) Nb22 demonstrates complete protection in the treatment of Delta variant infection in hACE2 transgenic mice.

4) We complete the proof of concept of STIP against SARS-CoV-2 using intranasal Nb22 with ultrahigh potency and long-lasting retention in respiratory system.

Graphic Abstract

Introduction:

SARS-CoV-2 has given rise to the COVID-19 pandemic [1], resulting in massive disruption of social and economic activities. Global vaccination has provided protection against the catastrophic outcome of the pandemic. However, a number of individuals are either not fully vaccinated or cannot mount adequate responses to the vaccine. Additionally, current COVID-19 vaccines require multiple doses to achieve full effectiveness and the immunity wanes within a matter of months, which increases the risk of infection and demands the use of agents for providing instantaneous protection at vulnerable times. Several antibodies were approved for emergency use within 7 days of high-risk exposure in the Post-exposure prophylaxis (PEP) against SARS-CoV-2 infection [2,3]. However, there is no licensed agent in preventing infection before exposure to SARS-CoV-2 (i.e., as pre-exposure prophylaxis, PrEP). A few PrEP studies in animal model demonstrated that antibodies exhibited accelerated clearance of SARS-CoV-2 when administered 1-3 days prior to infection[2,4-6]. The efficacy was not fully explored when antibodies were administered more than three days prior to SARS-CoV-2 infection. To the best of our knowledge, there is no effective intervention to prevent SARS-CoV-2 infection in advance of one week or longer. Therefore, there is a research gap on short-term instantaneous prophylaxis (STIP) that prevention can take effective immediately following antibody infusion and last for one week or longer. As such, STIP is an unmet need for the prevention against SARS-CoV-2.

The Delta variant, also known as B.1.617.2, was first identified in India in December 2020 and has become predominant in many countries, characterized by the spike protein mutations T19R, L452R, T478K, D614G, P681R, D950N and a double deletion at 157-158 [7-11]. It has been designated as a Variant of Concern (VOC) and is believed to be 60% more transmissible than Alpha variant [12]. Delta variant poses a challenge to the available COVID-19 vaccines, such as the protective effectiveness of AstraZeneca and Pfizer vaccines against Delta variant was reduced to 60% and 88%, respectively [11,12]. More recently, a newly emerged variant, Omicron, has spread rapidly in parts of the world and drawn attention for its potential impact on the global public health; however,

in most of the world including China, Delta variant remains the dominant virus and the

focus of the containment efforts. Recent research indicated that the Delta variant

partially but significantly resisted neutralization by mAbs including Bamlanivimab,

SARS-CoV-2 convalescent sera and vaccine-elicited antibodies [13,14]. While B1-182.1

and A23-58.1, recently isolated from convalescent donors, exhibited ultrapotent

neutralization against Delta variant with IC_{50} values of 1.0 and 1.6 ng/ml, respectively

[15].

To date, a growing number of nanobodies, single-domain fragments of camelid heavy-

chain antibodies or VHH, were reported for the prophylaxis or treatment of SARS-CoV-

2 infection. However, nanobodies with potent neutralization against Delta variant were

rarely reported [6,16-22]. As SARS-CoV-2 is transmitted through and replicates in

respiratory tract and lungs, and does not transverse in blood[1,23], we believe that,

nanobodies's exceptional resistance to extreme pH and high temperature[24], makes them

ideal candidates to be administered via intranasal or oral rout, directly to the site of viral

infection. Previously, we identified three ultrapotent nanobodies against the initial

strain of SARS-CoV-2, Wuhan-Hu-01 (WH01); accordingly, one of the intranasally

delivered nanobody was shown to protect hACE2 mice infected by WH01 strain[25].

Here, we compared the neutralizing potency of the aforementioned nanobodies against

various circulating SARS-CoV-2 variants. Nb22-Fc was identified to exhibit increased

neutralization potency against Delta variant compared to WH01 strain, to which the

antibody was originally raised. The binding characterization and crystal structural

analysis were conducted to further elucidate the potential mechanism. Furthermore,

therapeutic studies demonstrate that intranasal Nb22 exhibited complete protection

against SARS-CoV-2 Delta variant in hACE2 mice. Additionally, we comprehensively

evaluated the prophylactic efficacy of Nb22 when intranasally administered at 1, 3, 5,

7 days prior to SARS-CoV-2 infection. Notably, single dose of intranasal Nb22 still

exhibited protective against hACE2 mice even when administered 7 days prior to

infection of Delta variant. Moreover, four doses of intranasal Nb22 could maintain

long-lasting retention in respiratory system for more than one month. All these indicate

that intranasal Nb22 could be applied not only in the PrEP and PEP but also in the STIP,

filling the gap between the long-term lagging prevention and PreP.

Results

Potent neutralization of Delta variant by nanobodies

We previously reported the discovery and characterization of three potent neutralizing

nanobodies against WH01 strain and its variants with IC_{50} values of ~1 ng/ml. These

three nanobodies (Nb15-Fc, Nb22-Fc and Nb31-Fc) were identified to bind to RBD [25].

Neutralization experiments were further conducted to measure their activity against the

circulating variants including variants of concern (VOC) comprising Alpha (B.1.1.7

with N501Y), Beta (B.1.351 with E484K and N501Y), Delta (B.1.617.2 with L452R

and T478K) and Gamma (P.1 with K417T, E484K and N501Y), as well as variants of

interest (VOI) comprising Eta (B.1.525 with E484K), Iota (B.1.526 with E484K),

Epsilon (B.1.429 with L452R), and Kappa (B.1.617.1 with L452R and E484Q) [7-10].

Nb15-Fc exhibited increased potency against Alpha variant, but decreased potency

against Delta variant or Epsilon as compared with WH01, the RBD used to select the

nanobodies. Nb31-Fc exhibited reduced potency against Alpha, Delta and Epsilon

variants relative to WH01 or D614G variant (Fig. 1A-G). Interestingly, Nb22-Fc

exhibited about 2.5-fold increased neutralizing potency against Delta variant with an

IC_{50} value of 0.41 ng/ml (5.13 pM) compared to WH01 with an IC_{50} of 1.01 ng/ml

(12.63 pM). Notably, Nb22-Fc also exhibited around 8.4-fold increased neutralization

potency against Delta variant relative to variant Alpha with an IC_{50} of 3.45 ng/ml (43.13

pM) (Fig. 1A-G). Impressively, Nb22-Fc also exhibited outstanding neutralizing curve

against authentic Delta variant compared to Nb15-Fc and Nb31-Fc (Fig. 1F-G). All

three nanobodies failed to neutralize variants containing E484K/Q mutation, suggesting

that E484K/Q mutation in RBD could lead to the resistance to all three nanobodies.

Altogether, Nb15-Fc presented the most potent neutralization against variant Alpha

with an IC_{50} of 0.18 ng/ml, and Nb15-Fc and Nb31-Fc still retained potent

neutralization of variants containing L452R and T478K mutations in RBD (Fig. 1E-F),

though with reduced potency like most other anti-RBD antibodies [7,12]. Of note, Nb22-
Fc exhibited the most potent neutralization against pseudotyped or authentic Delta
variant virus among three nanobodies (Fig. 1A-G).

Characterization of Nb22-Fc binding to RBD

To explore antibody binding characteristics to the RBD with respect to their
neutralization of Delta variant, the interactions of three nanobodies with variant RBDs
were analyzed using biolayer interferometry (BLI). Nb15-Fc, Nb22-Fc and Nb31-Fc
showed high affinity interactions with RBD of Delta variant at 1.86 nM, 0.31 nM and
0.31 nM, respectively (Fig. 2A-D). However, the ultrahigh affinity of Nb22-Fc and
Nb31-Fc to the RBD of Delta variant did not fully reflect the neutralization potency as
Nb22-Fc neutralized Delta variant with markedly more potency than that of Nb31-Fc,
suggesting that affinity is not the only factor dictating the neutralization activity.
Furthermore, Nb22-Fc exhibited increased affinity with Delta variant RBD relative to
other variant RBDs (Fig. 2A-D), which is in line with the increased potency conferred
by Nb22-Fc against Delta variant as compared with other variants.

Moreover, immunofluorescence analysis revealed that Nb22-Fc specifically interacted
with spike protein from WH01, D614G, Alpha, Epsilon and Delta variants on the
surface of transfected 293T cells, whereas no binding with the spike protein from other
variants containing E484K/Q mutation (Fig. 2E). These results were substantiated by
flow cytometric results (Fig. 2F). Overall, these specific binding characteristics are
consistent with its specific neutralization properties.

Structural analysis of RBD-Nb22 complex

Structural analysis of Nb22 interaction with RBD was performed to address the
ultrahigh potency of Nb22 against WH01 strain and Delta variant. Initially, we
determined the crystal structure of WH01 RBD-Nb22 complex at a resolution of 2.7 Å
(Fig. 3A and Table S1). Nb22 adopts a typical β-barreled structure, and contains three
variable complementarity-determining regions (CDR) binding to RBD. The buried

surface area (BSA) was 800 $Å^2$, mainly constituting of hydrogen bonds and hydrophobic interactions. 14 residues constituting epitope of three CDRs were identified using a distance of <4 Å as the cutoff (Fig. 3B). For CDR1, T30 and S33 formed two hydrogen bonds with S494 of RBD, while the hydrophobic interactions included A32 and F34 of Nb22 and Y449, L452, F490 and Q493 of RBD (Fig. 3C and Table S2). N57 of CDR2 interacted with G485 by hydrogen bond, and the hydrophobic interactions were mediated by I56 and Y489 (Fig. 3D). CDR3 is a relatively longer region with only one hydrogen bond (Y119 and G446). The side chain of P104 inserted into the hydrophobic cavity formed by F101, R107, Y453, F456 and Y495 (Fig. 3E). Apart from the five hydrogen bonds in CDR regions, the interface of Nb22 and RBD was stabilized by two additional hydrogen bonds consisting of G1, S75, N450 and E484 (Fig. 3F). Interactions were also facilitated by the hydrophobic network constituted by P2, Q3, V4, G28, G29, R73 and D74 of Nb22 (Fig. 3G and Table S2).

Superimposition of the structure of WH01 RBD-Nb22 complex and RBD-hACE2 (PDB code: 6MOJ) immediately elucidates the structural basis of neutralization, in which the binding of Nb22 to RBD effectively blocks the binding of RBD to hACE2 during virus infection. Firstly, the binding site of Nb22 on RBD partially overlaps with that of hACE2 (Fig. S1A). Secondly, the loop (V102-Y117) of Nb22 clashes with two α-helices of the N-terminus of hACE2 (Fig. S1B).

To elucidate the increased potency of Nb22 in neutralizing Delta variant, we determined the Delta RBD-Nb22 complex structure at a resolution of 2.9 Å, which revealed that two distinct mutations, T478K and L452R, had differing impact on the binding K478 locates outside the CDR binding regions, and does not disturb the interaction surface (Fig. 4A and Table S1). Therefore, T478K mutation does not affect Nb22 neutralization (Fig. 4A). However, the mutation of hydrophilic leucine to positively charged arginine (R) at position 452 significantly enhances the interactions of RBD with the CDR3 region of Nb22. Two additionally formed hydrogen bonds, T30-R452 and S33-Q493, pull the CDR3 loop of Nb22 closer to R452 region of RBD, as revealed by the

superimposition of the structures of WH01 and Delta RBD-Nb22 (Fig. 4B, 4C and Table S2), explaining enhanced neutralization activity of Nb22 against the Delta variant.

Nb22 exhibits room-temperature stability in vitro and long-lasting retention in vivo

Nanobodies exhibit various advantages including thermostability. We reported that nanobodies could retain 100% activity even after being incubated at 70 °C for one hour and retain 83% activity after 80 °C treatment for one hour[25]. Further evaluation showed that Nb22 could maintain full activity for at least two months at room temperature and did not lose any activity even undergoing five rounds of freeze-thawing (Fig. 5A and 5B), indicating that Nb22-Fc is highly stable and idea for non-cold chain storage and use at room-temperature.

To determine Nb22-Fc distribution in vivo, YF®750 SE-labeled Nb22 (Nb22-YF750) was administered via intranasal (i.n.) in a mouse model. The fluorescence could be readily detected in respiratory system including nasopharynx, trachea and lung 2h post infusion. The fluorescence was detectable for more than seven days after a single dose of 200 ug (average of 10 mg/kg) Nb22-YF750 administration, which is in agreement with our previous reports. As expected, the fluorescence could be detected for more than one month when four doses of 200 μg (average of 10 mg/kg) Nb22-YF750 were administered every week (Fig. 5C-E and Fig. S2). Nb22-Fc could also be detected in the blood, indicating that Nb22-Fc could also exert its activity in the blood after bypassing the respiratory system (Fig. 5F). All these observations of prolonged retention of Nb22-Fc in the respiratory system suggest the potential application of the antibody for STIP against SARS-CoV-2 infection. Taken together, intranasal Nb22-Fc could be developed as an STIP reagent for its long-lasting retention in the respiratory system and a portable therapeutics thanks to its room-temperature stability.

Intranasal Nb22 is highly efficacious in the STIP, PreP and PeP in hACE2 transgenic mice challenged by Delta variant

To evaluate the efficacy of Nb22-Fc in vivo, hACE2 transgenic mice were challenged

with 1×10^5 PFU SARS-CoV-2 Delta variant (CRST：1633.06.IVCAS 6.7593) and

conducted as we previously reported[25]. hACE2 mice were divided into nine groups

(n=3-6) as shown in Figure 6A and 200 μg (average of 10 mg/kg) Nb22-Fc was

administered via i.n. before or after Delta variant challenge to evaluate the antibody's

prophylactic or therapeutic efficacy against Delta variant infection. Viral RNA in the

lungs was detected in virus control group named as SARS-CoV-2 group (Group 2).

Animals in Control Nb group Group 3 were challenged with Delta variant and received

control nanobody treatment one hour post infection. As expected, high copy numbers

of viral RNA were also detected in control Nb mice without significant difference

compared to SARS-CoV-2 group (Fig. 6B).

In order to evaluate the prophylactic duration conferred by Nb22, a single dose of Nb22

was administered via i.n. at days 1, 3, 5, and 7, respectively, prior to Delta variant

challenge. Viral RNA copies increased over the course of Nb22-Fc administration (Fig.

6B). As expected, viral RNA copies in the aforementioned prophylactic groups were

all significantly lower than that in the control Nb group, indicating that even a single

dose of Nb22-Fc could provide protection against Delta variant infection in hACE2

transgenic mice for 7 days. Nb22 administered in -7d and -5d before challenge

significantly reduced viral load though failed to provide complete protection (Fig. 6B).

Notably, Nb22 exhibited significant prevention against SARS-CoV-2 infection in the -

3d Nb22 (Group 6) and -1d Nb22 group (Group 7) (Fig. 6B). All these indicate that

Nb22-Fc provides better protection at the earlier time points upon infection.

In the therapeutic group, viral RNA copies in the animals treated with Nb22 at 1 hour

or 1 day postinfection were undetectable in 5/5 mice in the groups 8 and 9, suggesting

that Nb22 had complete protection of hACE2 mice against Delta variant infection (Fig.

6B). The viral RNA results were also validated by immunofluorescence (IF) staining

and HE staining in the lungs (Fig. 6C and 7E). We noted that mice challenged by Delta

variant did not show obvious weight loss even in 4 days post infection (Fig. S3). In

summary, Nb22 exhibited high efficacy in both the prevention and therapy of hACE2

transgenic mice challenged with Delta variant. Nb22 provided complete protection in

PEP (in 1h Nb22 group and 1d Nb22 group) and exhibited high efficacy in PrEP (in -

1d Nb22 and -3d Nb22 group). Impressively, a single dose of Nb22 could maintain

effectiveness in the prevention against Delta variant infection for at least seven days (in

-7d Nb22 group), indicating the potential application for STIP against SARS-CoV-2.

Discussion

To date, a small number of nanobodies with ultrahigh potency against SARS-CoV-2

and its variants have been reported [15,26,27], whereas nanobodies with potent

neutralization against the currently dominant Delta variant were rarely reported. Our

results revealed that three previously reported nanobodies [25], retained ultrahigh potency

in neutralization against the Delta variant. Among them, Nb22-Fc with an IC_{50} value of

0.41ng/ml (5.13 pM) is outstanding with increased neutralization of Delta variant

relative to Alpha variant. The Nb22 binding to RBD provides mechanistic insight into

the enhanced neutralization against Delta variant, suggesting that the increased binding

affinity enhanced the neutralizing potency against Delta variant relative to Alpha

variant (Fig. 1). Given that most anti-RBD, anti-NTD antibodies or convalescent sera

or vaccine-elicited antibodies showed reduced neutralization of Delta variant relative

to that of Alpha variant [7,12], the increased neutralization activity of Nb22-Fc against

Delta variant is particularly striking and the structural basis of the phenomenon is of

interest for understanding the neutralization mechanisms.

The structural analysis further illustrated the characteristics of Nb22 binding to WH01

and Delta RBD and the mechanisms of viral inhibition. Nb22 binding to RBD

effectively blocks the binding of RBD to hACE2 during virus infection. The binding

site of Nb22 on RBD overlaps with that of hACE2 (Fig. S1A), and the loop (V102-

Y117) of Nb22 clashes with two α-helixes of the N-terminus of hACE2 (Fig. S1B). In

addition, crystal structural analysis showed that T478K mutation of Delta variant are

located outside 800 $Å^2$ BSA of Nb22 interacting with RBD and do not perturb the

interaction between Nb22 and the RBD of Delta variant. Of note, the guanidine moiety

in the L452R mutation forms an additional hydrogen bond with the hydroxyl group of T30 of Nb22, pulling the CDR3 loop of Nb22 closer, and leading to an extra hydrogen bond between S33 of Nb22 and Q493 of RBD (Fig. 4). Consequently, the BSA extends from 800 $Å^2$ to 835 $Å^2$, in comparison with that of Nb22-WH01 RBD. All these contribute to the enhanced binding and neutralizing potency of Nb22 against Delta variant.

Compared to conventional antibodies for passive immunization, nanobodies are efficiently produced in prokaryotic expression systems at low cost and possess favorable biophysical properties including high thermostability[28]. We reported that nanobodies could remain 100% activity even incubated at 70 ℃ for one hour and are amenable to engineering of multimeric nanobody constructs[25]. Such nanobodies exhibit high effectiveness against virus infection via intranasal administration[25]. The results reveal that Nb22 could maintain full activity for more than two months at room temperature and does not lose any activity even after undergoing five rounds of freeze-thawing.

Our results demonstrate that a single dose of intranasal Nb22 could exhibit efficacy in the STIP, PrEP and PEP against SARS-CoV-2 infection in hACE2 mice. Of note, a single dose of intranasal Nb22 could maintain efficacy against SARS-CoV-2 infection for at least one week in hACE2 mice, which would readily serve as STIP. Our antibody distribution results also revealed that Nb22 could retain in respiratory tracts for at least one month when weekly administered via i.n.. As such, we anticipate that Nb22 could provide one month prevention against SARS-CoV-2 infection when administered intranasally every week.

VLong–term lagging prevention against SARS-CoV-2 conferred by approved vaccine usually takes more than one month to be effective and then lasts for months or years[29]. Vaccine efficacy has been shown to wan within months after vaccination [7,11]. Instantaneous prevention of SARS-CoV-2 is also needed for individuals who do not take vaccines when SARS-CoV-2 is circulating. A few studies in animal model just demonstrated that antibodies exhibited accelerated clearance of SARS-CoV-2 in PrEP when administered 1-3 days prior to infection[2,4-6]. Whereas, to the best of our

knowledge, few studies have fully investigated the STIP that prevention could be

readily effective immediately following inoculation and last for more than one week to

one month for people at high risk of SARS-CoV-2 infection, which can also serve to

reduce transmission during the asymptomatic stage of the infection As such, our results

demonstrate that intranasal Nb22 with ultrahigh potency and long-lasting retention in

the lung could satisfy the need of STIP against SARS-CoV-2.

In summary, structural analysis provides a mechanistic explanation to the enhanced

sensitivity of Delta variant and the increased neutralization potency of this antibody.

The structural analysis may further guide the rational design of pan-coronavirus

vaccines and therapeutics. Nb22 exhibited one of the ultrahigh neutralization potencies

among the reported antibodies or nanobodies against Delta variant infection [7,12,26,30,31].

We presented proof of concept of STIP against SARS-CoV-2 using our Nb22 and

suggest STIP as a new prophylactic strategy for long-lasting antibodies to prevent virus

infection. Although the newly emerged Omicron variant is spreading globally, the Delta

variant remains the dominant variant and likely the deadliest in most of the world;

therefore, the ultrahigh potent, and thermal stable Nb22 is an excellent candidate for

intranasal or inhalable anti-SARS-Cov-2 agent for both therapy or prophylaxis,

especially including STIP.

Materials and Methods

Expression of nanobodies

The Fc1 gene (CH2-CH3) of the human monoclonal antibody was fused with the VHH

gene of nanobodies (named as Nb-Fc or Nbs) to assist the purification and prolong the

half-life of the Nb antibody, following our previously published protocol [32]. The Nbs

were finally cloned into the pcDNA3.4 eukaryotic expression vector (Invitrogen),

which were transfected into 293F cells (cat.# R79007, Thermo Scientific) to produce

Nb-Fcs. Nb fused with Fc was purified using Protein G (cat.# 20399, Thermo

Scientific).

Enzyme linked immunosorbent assay (ELISA) analysis

Antibody quantification and antibody characterization were tested by ELISA as our

previously reported method[33], with modifications. In brief, the protein was coated to

ELISA plates (Corning) at a concentration of 0.5 μg/ml. After washing 2-4 times, 5%

non-fat milk in PBS was added and incubated for blocking at 37 °C for 1 h. After

washing, 100 μl serially diluted sera or purified antibody was added and incubated at

37 °C for 1 h. Following washing, secondary antibody of goat anti-llama IgG (H+L)

with HRP (Novus, cat.# NB7242, 1:10000 dilution) or goat anti-human IgG with HRP

was added and incubated at 37 °C for 1 h. Accordingly, 3,3′,5,5′-Tetramethylbenzidine

(TMB, Sigma) substrate was added at 37 °C for 10 minutes (min); and 10 μl 0.2 M

H_2SO_4 was added to stop the reaction. The optical densities at 450 nm (OD450) were

measured using the Infinite 200 (Tecan, Ramsey, MN, USA). Antibody quantification

in the sera was calculated according to the standard curve generated by purified

antibody.

Neutralization activity of nanobodies against pseudovirus

Pseudovirus neutralization assay was carried out following our previously published

protocol[25], with the follow modifications. Briefly, pseudovirus of SARS-CoV-2

variants was produced by co-transfection of pNL4-3.Luc.R-E-, an HIV-1 NL4-3

luciferase reporter vector that comprises defective Nef, Env and Vpr (HIV AIDS

Reagent Program), and pVAX1 (Invitrogen) expression vectors encoding the spike

proteins of respective variants into 293T cells (ATCC). Supernatants containing

pseudovirus were collected after 48 hours (h), and viral titers were determined by

luciferase assay in relative light units (Bright-Glo Luciferase Assay Vector System,

Promega Biosciences). Human codon optimized S genes of SARS-CoV-2 variants were

synthesized, and the corresponding pseudoviruses were produced following the above

protocol. For neutralization assay, SNB02, an Nb-Fc specific against SFTSV [32], served

as a negative control. Neutralization assays were conducted by incubating pseudovirus

with serial dilutions of purified nanobodies at 37 °C for 1 h. HEK293T-ACE2 cells

(cat.# 41107ES03, Yeasen Biotech Co., Ltd. China) (approximately 2.5×10^4 per well)

were added in duplicate to the virus-antibody mixture. Half-maximal inhibitory

concentrations (IC_{50}) of the evaluated nanobodies were determined by luciferase

activity 48 h following exposure to virus-antibody mixture, and analyzed by GraphPad

Prism 8.01 (GraphPad Software Inc.).

Immunofluorescence and flow cytometric analysis

Immunofluorescence and flow cytometric analysis were conducted following our

previously published protocol [34], with minor modifications. Briefly, S gene sequences

for SARS-CoV-2 spike protein of various SARS-CoV-2 variants were obtained from

the GISAID website (https://gisaid.org). S genes were synthesized and constructed as

expression plasmids by GenScript. The plasmids were transfected into 293T cells

(ATCC) cultured in 12-well plates. Next, 48 hours post transfection, the cells were

washed by PBS and fixed with 4% paraformaldehyde for 20 minutes at room

temperature. The purified Nb-Fc was used to stain the 293T cells, followed by Alexa

Fluor 488 AffiniPure goat Anti-human IgG (H+L) (1:500 dilution) (109-545-003,

Jackson ImmunoResearch). For immunofluorescence analysis, the cells on the plate

were examined and the images were acquired using an OLYMPUS IX73. For flow

cytometric analysis, the cells were resuspended in 500 µl PBSF buffer (PBS+2% FBS)

and analyzed using ACEA NovoCyte TM (Agilent Biosciences); non-transfected 293T

cells served as a negative control.

Affinity determination by Bio-Layer Interferometry (BLI)

We measured antibody affinity using a ForteBio OctetRED 96 BLI (Molecular Devices

ForteBio LLC, Fremont, CA) with shaking at 1,000 rpm at 25 °C [25]. To determine the

affinity of Nbs with human Fc tag, Nb-Fcs were loaded to anti-human Fc (AHC)

biosensors (cat.# 18-5060, Fortebio) in a kinetic buffer (PBS, 0.02% (v/v) Tween-20,

pH 7.0) for 200 sec prior to baseline equilibration for 180 sec in a kinetic buffer.

Association of SARS-CoV-2 RBD in a three-fold dilution series from 33.3 nM to 1.2

nM was performed prior to dissociation for 180 sec. After each cycle, the biosensors

were regenerated through 3 brief pulses of 5 sec each with 100 mM pH 2.7 glycine-

HCL followed by a running buffer. The data were baseline subtracted before fitting

using a 1:1 binding model and the ForteBio data analysis software. K_D, Ka and Kd

values were determined by applying a global fit to all data.

Expression and purification of WH01 and Delta RBD protein for crystal

structural analysis

The WH01 and Delta RBD were expressed using the Bac-to-Bac baculovirus system.

The two pAcgp67-RBD (residues 333–530) plasmid with a C-terminal 8×His tag were

transfected into Sf9 cells using Cellfectin II Reagent (Invitrogen) to produce the

recombinant baculoviruses. After 3 rounds of amplification, Hi5 cells were infected

with baculoviruses at an MOI of 4 at a density of 2×10^6 cells/ml. The supernatants of

cell culture containing the secreted RBD were harvested at 60 h after infection. The

RBD was purified by Ni-NTA resin (GE Healthcare). Nonspecific contaminants were

removed by washing the resin with 20 mM Tris-HCl, 150 mM NaCl, pH 7.5, and the

target proteins were eluted with elution buffer containing 20 mM Tris-HCl, 150 mM

NaCl, 500 mM imidazole, pH 7.5. The eluted proteins were further purified by

Superdex 75 (GE Healthcare, USA) and stored in 20 mM Tris-HCl, 150 mM NaCl, pH

7.5.

Expression and purification of Nb22 for crystal structural analysis

The VHH gene for Nb22 was amplified by PCR and cloned into a pET21a vector with

*Bam*H I and *Xho* I restriction sites. The recombinant plasmids were transformed into

Escherichia coli. BL21 (DE3). The cells were cultured in LB medium and grown to

$OD_{600} = 0.8$ at 37°C. Isopropyl-D-1-thiogalactopyranoside (IPTG) was added to a final

concentration of 1.0 mM to induce the protein expression, and the cultures were grown

at 16 °C overnight. Cells were harvested by centrifugation at 4,500 rpm for 15 min, re-

suspended and homogenized in the lysis buffer containing 20 mM Tris-HCl, 150 mM

NaCl, pH 7.5 using ultrasonic. Cell debris was removed by centrifugation at 18,000

rpm for 30 min. The supernatants were added to Ni- NTA resin (GE Healthcare, USA).

The nonspecific contaminants were eluted by washing the resin with the lysis buffer

containing 10 mM imidazole. The target protein with 6 x His tag, named as Nb22, was

subsequently eluted with the lysis buffer containing 500 mM imidazole. Nb22 was

eluted and purified by Superdex 75 (GE Healthcare, USA).

Crystallization, structural determination and data acquisition

The complexes were prepared by mixing WH01 or Delta RBD and Nb22 at a 1:1.2

molar ratio and incubating at 4 °C overnight. The complexes were further purified by

Superdex 75 (GE Healthcare, USA) to remove the excess nanobody. The crystals were

screened by vapor-diffusion sitting-drop method at 16˚C. The crystals appeared and

reached their final size within 3 days in a well solution comprising 0.1 M HEPES (pH

7.0), 5% *v/v* (+/-)-2-Methyl-2,4-pentanediol (MPD), 10% polyethylene glycol (PEG)

10000 (WH01 RBD-Nb22) and 0.1 M Tris (pH 7.0), 37.5% Jeffamine (Delta RBD-

Nb22), respectively.

To collect data, a single crystal was mounted on a nylon loop and was flash-cooled with

a nitrogen gas stream at 100 K. Diffraction data of WH01 RBD-Nb22 was collected on

BL18U1 at Shanghai Synchrotron Radiation Facility (SSRF) at a wavelength of

0.97915 Å. While, the Delta RBD-Nb22 was collected on BL02U1 at a wavelength of

0.97918 Å. Data were processed and scaled using the HKL3000 package and

autoPROC [35]. The structures were elucidated using the molecular replacement (MR)

method in PHASER program [36] with the structure of SARS-CoV-2 RBD (PDB code:

7CJF) [37] as the initial searching model. The model was built into the modified

experimental electron density using COOT [38] and further refined in PHENIX [39]. The

final refinement statistics are summarized in Table S1. Structural figures were prepared

by PyMOL. Epitope and paratope residues, as well as their interactions, were identified

by PISA (http://www.ebi.ac.uk/pdbe/prot_int/pistart.html) at the European

Bioinformatics Institute.

Pharmacokinetics (PK) of Nb22-Fc *in vivo*.

Purified Nb22-Fc were injected via intranasal (*i.n.*) into BALB/c mice (Qing Long Shan Animal Center, Nanjing, China) at a dose of 10 mg/kg. The concentration of Nb22-Fc in serum was measured by ELISA. The $T_{1/2}$ of Nb22-Fc was caculated as ln (2)/k, where k is a rate constant expressed reciprocally of the x axis time units by the plateau followed one phase decay or one phase decay equation in the GraphPad software.

Spatial distribution of Nb22-Fc *in vivo*

Spatial distribution of Nb22-Fc were conducted following our previously published protocol[25], with minor modifications. Nb22-Fc labeled with far infrared dye YF®750 SE (US EVERBRIGHT INC, YS0056) were named as Nb22-YF750. 10 mg/kg Nbs-YF750 were administered via intranasal into nude mice (18-22g, Qing Long Shan Animal Center, Nanjing, China). Images were recorded at Ex:740 nm/Em:780 nm by NightOWL LB 983 (Berthold, Germany) at the indicated time point. Images were analyzed using Indigo imaging software Ver. A 01.19.01.

Evaluating the efficacy of Nb22-Fc in SARS-CoV-2 infected hACE2 mice.

The efficacy of Nb22-Fc against SARS-CoV-2 were evaluated according to our previously published protocol[25], with minor modifications. In brief, A total of 43 8-week-old male transgenic hACE2 mice (C57BL/6J) (cat.# T037630, GemPharmatech Co., Ltd., Nanjing, China) were challenged with $1x\ 10^5$ PFU SARS-CoV-2 Delta variant (CRST：1633.06.IVCAS 6.7593) per mouse. The mice were randomly divided into nine groups (n=3-6) for either prophylactic or therapeutic evaluation, as described in Figure 5A. Mice without any treatment and challenge were taken as blank control (No SARS-CoV-2, n=3). Mice challenged with SARS-CoV-2 were taken as infection control (SARS-CoV-2, n=5). 250 μg/mouse (average of 10 mg/kg) SNB02 (Y-Clone, China), an anti-SFTSV antibody constructed by Nb fused with human Fc1 (Nb-Fc) [32], was intranasally injected 1 hour (h) after infection and was taken as an isotype antibody treated control (Control Nb, n=3). For the prophylactic group, mice were intranasally injected with Nb22-Fc at a dose of 250 μg/mouse (average of 10 mg/kg) at 7 days (d),

5d, 3d, 1d before infection (named as -7d Nb22, -5d Nb22, -3d Nb22, -1d Nb22,

respectively, n=5-6). For the therapeutic group, mice were intranasally injected with

Nb22-Fc at a dose of 250 μg/mouse (average of 10 mg/kg) 1 h or 24 h after infection

(named as 1h Nb22 and 1d Nb22, n=5, respectively). Body weight of mouse was

recorded daily. Given that hACE2 transgenic mice typically clear virus within 5-7 days

after SARS-CoV-2 infection[40], the mice were sacrificed at 4 days post infection (dpi).

Subsequently, lung tissues were harvested for viral load determination and tissue

sections for immunofluorescence (IF) and hematoxylin and eosin (H&E) staining. All

experiments were conducted in a Biosafety Level 3 (BSL-3) facility.

Viral load measurement by quantitative RT-PCR.

Viral load was measured by quantitative real-time PCR (qRT-PCR) on RNA extracted

from the supernatant of lung homogenates as reported previously [41]. Briefly, lung

homogenates were prepared by homogenizing perfused lung using an electric

homogenizer. The inactivated samples were transferred from the BSL-3 to BSL-2

laboratory and total RNA was extracted from the collected supernatant. Each RNA

sample was reverse transcribed to 50 μl cDNA with HiScript II Q RT SuperMix for

qPCR (+gDNA wiper) (R223-01). 5 μl cDNA was added into a 25 μl qRT-PCR reaction

containing the ChamQ SYBR qPCR Master Mix (High ROX Premixed) (Q341-02,

Vazyme Biotech, China) and primers designed to target the nucleocapsid protein of

SARS-CoV-2 (5′- GGGGAACTTCTCCTGCTAGAAT -3′ and 5′-

CAGACATTTGCTCTCAAGCTG -3′). The samples were run in triplicate on an

ABI 7900 Real-Time System (Applied Biosystems, Thermo Fisher Scientific). The

following cycling conditions were performed: 1 cycle of 50 °C for 2 min, 1 cycle of

95 °C for 10 min, and 40 cycles of 95 °C for 15 sec and 58 °C for 1 min.

Immunofluorescence staining of SARS-CoV-2-infected cells and H&E staining in

tissues.

The lung tissues were immersed in 10% neutral buffered formalin (cat.# Z2902, Sigma)

for 24 h. After the formalin fixation, the tissues were placed in 70% ethanol (Merck)

and subsequently embedded with paraffin. Tissue sections (5-µm thick) were prepared

for H&E staining and immunofluorescence staining for SARS-CoV-2 detection using

the Coronavirus nucleocapsid protein (NP) antibody (cat. 40143-MM05, Sino

Biological). Images were collected under a Pannoramic MIDI system (3DHISTECH,

Thermo) using Pannoramic scanner software and analyzed by ImageJ (NIH).

Quantification and statistical analysis

All statistical analyses were carried out using GraphPad Prism 8.01 software (GraphPad)

or OriginPro 8.5 software (OriginLab). ANOVA or Mann-Whitney test was performed

for group comparisons. $P < 0.05$ was considered as statistically significant with mean

±SEM or mean ±SD.

Study approval

The study and the protocol for this research were approved by the Center for Public

Health Research, Medical School, Nanjing University. All animal experimental

procedures without infection were approved by the Committee on the Use of Live

Animals by the Ethics Committee of Nanjing University. All animals infected by

SARS-CoV-2 were performed in Biosafety Level 3 animal facilities in accordance with

the recommendations for care and use of the Institutional Review Board of Wuhan

Institute of Virology of the Chinese Academy of Sciences (Ethics Number:

WIVA11202111). All the authors declare their compliance with publishing ethics.

Author contributions

XW conducted most experiments, analyzed the data and wrote the draft manuscript. LC

conducted all the neutralization experiments. LZ, BH, MJ, SX, HS, DZ, LL, WN

provided technical assistance. YW, SM and SY conducted the structural analysis. FN,

YL, HH, QH and YL, evaluated the efficacy of Nb22 in SARS-CoV-2 infected

transgenic hACE2 mice. ZW designed the study, directed and revised the manuscript.

All authors critically reviewed the draft manuscript and approved the final version.

Acknowledgements

We thank Prof. Guo. for providing the plasmid of RBD. The X-ray data were collected

using Shanghai Synchrotron Radiation Facility on BL18U1 and BL02U1.   The

efficacy of Nb22 in SARS-CoV-2 infected transgenic hACE2 mice was evaluated in

Wuhan National Biosafety Laboratory, Chinese Academy of Sciences. This work was

supported by National Science Foundation of China (NSFC) (No. 81803414 to X.W.,

31970149 to Z.W.), the Major Research and Development Project (2018ZX10301406

to Z.W.), Ministry of Science and Technology (2020YFA0908500 to S.Y.), the National

Natural Science Foundation of China (31971127 to S.Y. and 81801998 to Y.W.), Tianjin

Natural Science Foundation (20JCQNJC01570 to Y.W.), Nanjing University-Ningxia

University Collaborative Project (Grant# 2017BN04 to Z.W.), Jiangsu Province Natural

Science Foundation for Young Scholar (Grant# BK20170653 to X.W.), Key Natural

Science Foundation of Jiangsu Province (Grant# ZDA2020014 to X.W.), the

Fundamental Research Funds for the Central Universities（Grant# 0214-14380523 to

X.W.）Jiangsu province "Innovative and Entrepreneurial talent" and Six Talent Peaks

Project of Jiangsu Province, the Emergency Prevention and Control Capacity Program

for New Severe Infectious diseases of National Institute for Viral Disease Control and

Prevention, and the 135 Strategic Program of Chinese Academy of Sciences, the

Science and Technology Innovation Committee of Shenzhen Municipality

(JCYJ20180228162229889 to L.C.), and the National Science Foundation of China (No.

31970172 to Y.L.).

Reference

1.   Zhou, P., *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin.
Nature (2020).

2.   Baum, A., *et al.* REGN-COV2 antibodies prevent and treat SARS-CoV-2 infection in rhesus
macaques and hamsters. *Science (New York, N.Y.)* **370**, 1110-1115 (2020).

3.   Chen, P., *et al.* SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19.
N. Engl. J. Med. (2020).

4.   Zost, S.J., *et al.* Potently neutralizing and protective human antibodies against SARS-CoV-2.
Nature **584**, 443-+ (2020).

5.   Rogers, T.F., *et al.* Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from
disease in a small animal model. *Science* (2020).

6.   Pymm, P., *et al.* Nanobody cocktails potently neutralize SARS-CoV-2 D614G N501Y variant
and protect mice. *Proc Natl Acad Sci U S A* **118**(2021).

7.   Liu, C., *et al.* Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and convalescent
serum. *Cell* (2021).

8.   Garcia-Beltran, W.F., *et al.* Multiple SARS-CoV-2 variants escape neutralization by vaccine-
induced humoral immunity. *Cell* (2021).

9.   Davies, N.G., *et al.* Increased mortality in community-tested cases of SARS-CoV-2 lineage
B.1.1.7. *Nature* (2021).

10.   Plante, J.A., *et al.* The variant gambit: COVID-19's next move. *Cell Host & Microbe* **29**, 508-
515 (2021).

11.   Bernal, J.L., *et al.* Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *N.*
Engl. J. Med. (2021).

12.   Planas, D., *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization.
Nature (2021).

13.   Wall, E.C., *et al.* AZD1222-induced neutralising antibody activity against SARS-CoV-2 Delta
VOC. *Lancet* **398**, 207-209 (2021).

14.   Yadav, P.D., *et al.* Neutralization of variant under investigation B.1.617 with sera of BBV152
vaccinees. *Clinical infectious diseases : an official publication of the Infectious Diseases*
Society of America (2021).

15.   Wang, L., *et al.* Ultrapotent antibodies against diverse and highly transmissible SARS-CoV-2
variants. *Science (New York, N.Y.)* **373**(2021).

16.   Xiang, Y., *et al.* Versatile and multivalent nanobodies efficiently neutralize SARS-CoV-2.
Science (New York, N.Y.) **370**, 1479-1484 (2020).

17.   Schoof, M., *et al.* An ultrapotent synthetic nanobody neutralizes SARS-CoV-2 by stabilizing
inactive Spike. *Science (New York, N.Y.)* **370**, 1473-1479 (2020).

18.   Huo, J.D., *et al.* Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block interaction
with ACE2. *Nature Structural & Molecular Biology* **27**, 846-+ (2020).

19.   Hanke, L., *et al.* An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor interaction.
Nature Communications **11**(2020).

20.   Wu, Y., *et al.* Identification of Human Single-Domain Antibodies against SARS-CoV-2. *Cell*

658    *Host & Microbe* **27**, 891-+ (2020).

659 21. Dong, J.*, et al.* Development of multi-specific humanized llama antibodies blocking SARS-
660    CoV-2/ACE2 interaction with high affinity and avidity. *Emerging Microbes & Infections* **9**,
661    1034-1036 (2020).

662 22. Nambulli, S.*, et al.* Inhalable Nanobody (PiN-21) prevents and treats SARS-CoV-2 infections
663    in Syrian hamsters at ultra-low doses. *Science advances* **7**(2021).

664 23. Wang, W.L.*, et al.* Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *Jama-
665    J Am Med Assoc* **323**, 1843-1844 (2020).

666 24. Wu, Y., Jiang, S. & Ying, T. Single-Domain Antibodies As Therapeutics against Human viral
667    Diseases. *Frontiers in immunology* **8**(2017).

668 25. Wu, X.*, et al.* A potent bispecific nanobody protects hACE2 mice against SARS-CoV-2
669    infection via intranasal administration. *Cell reports*, 109869-109869 (2021).

670 26. Guttler, T.*, et al.* Neutralization of SARS-CoV-2 by highly potent, hyperthermostable, and
671    mutation-tolerant nanobodies. *Embo Journal*.

672 27. Wrapp, D.*, et al.* Structural Basis for Potent Neutralization of Betacoronaviruses by Single-
673    Domain Camelid Antibodies. *Cell* **181**, 1436-1441 (2020).

674 28. Steeland, S., Vandenbroucke, R.E. & Libert, C. Nanobodies as therapeutics: big opportunities
675    for small antibodies. *Drug Discov Today* **21**, 1076-1113 (2016).

676 29. Dai, L. & Gao, G.F. Viral targets for vaccines against COVID-19. *Nature Reviews Immunology*
677    (2020).

678 30. Koenig, P.A.*, et al.* Structure-guided multivalent nanobodies block SARS-CoV-2 infection and
679    suppress mutational escape. *Science* **371**, 691-+ (2021).

680 31. Taylor, P.C.*, et al.* Neutralizing monoclonal antibodies for treatment of COVID-19. *Nature
681    Reviews Immunology* **21**, 382-393 (2021).

682 32. Wu, X.*, et al.* A single-domain antibody inhibits SFTSV and mitigates virus-induced
683    pathogenesis in vivo. *JCI insight* **5**(2020).

684 33. Wu, X.*, et al.* Induction of neutralizing antibodies by human papillomavirus vaccine generated
685    in mammalian cells. *Antibody Therapeutics* (2019).

686 34. Huang, B.*, et al.* Potent Neutralizing Humanized Antibody With Topical Therapeutic Potential
687    Against HPV18-Related Cervical Cancer. *Frontiers in immunology* **12**(2021).

688 35. Minor, W., Cymborowski, M., Otwinowski, Z. & Chruszcz, M. HKL-3000: the integration of
689    data reduction and structure solution--from diffraction images to an initial model in minutes.
690    *Acta Crystallogr D Biol Crystallogr* **62**, 859-866 (2006).

691 36. McCoy, A.J.*, et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674 (2007).

692 37. Jin, Z.*, et al.* Structure of M(pro) from SARS-CoV-2 and discovery of its inhibitors. *Nature* **582**,
693    289-293 (2020).

694 38. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta
695    Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

696 39. Adams, P.D.*, et al.* PHENIX: a comprehensive Python-based system for macromolecular
697    structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).

698 40. Bao, L.N.*, et al.* The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature* **583**,
699    830-+ (2020).

700 41. Cao, Y.L.*, et al.* Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-
701    Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-+ (2020).

Figure Legends

Figure 1 Characterizing nanobodies neutralizing circulating variants of SARS-CoV-2. The neutralization curve of Nb15-Fc (**A**), Nb22-Fc (**B**), Nb31-Fc(**C**) and SNB02 (**D**) inhibiting SARS-CoV-2 pseudovirus of circulating variants. Nb-Fcs and SNB02 were all constructed as the format of VHH fused with human Fc1. SNB02 was taken as an antibody control specific for SFTS virus. (**E**) The summary curve of IC_{50} of Nb-Fcs exhibiting potent neutralization against SARS-CoV-2 variants. (**F**) The neutralization potency of Nb-Fcs was evaluated based on authentic SARS-CoV-2 Delta variant plaque reduction neutralization test. (**G**) The summary table of IC_{50} and IC_{80} of Nb-Fcs in A-C and F, displaying potent neutralization. Data are represented as mean ± SD. All experiments were repeated at least twice.

Figure 2. Characterizing the binding of Nbs. Kinetic binding curve of Nb15-Fc (**A**), Nb22-Fc (**B**) and Nb31-Fc (**C**) at the concentration 33.3 nM,11.1nM, 3.7nM and 1.2 nM with RBD of Delta variant, respectively, detected by BLI. Binding curves are colored black, and fit of the data to a 1:1 binding model is colored red. (**D**) Representative binding curve of various RBD as indicated to Nb22-Fc tested by BLI. Nb22-Fc binding with RBD from representative SARS-CoV-2 variants detected by immunofluorescence assay (**E**) and flow cytometric analysis (**F**), respectively. Mock served as a cell control without plasmid transfection. Images were visualized under the ×10 objective. All experiments were repeated at least twice.

Figure 3. Structural analysis of Nb22 and WH01 RBD complex. (**A**) The overall complex structure of Nb22 and WH01 RBD. The CDR1 (red), CDR2 (blue), CDR3 (green) of Nb22 (pink) and WH01 RBD (orange) are displayed in cartoon representation. (**B**) The epitope of Nb22 shown in surface representation. The CDR regions are colored in red, blue and green, respectively. The interaction between CDR1 (**C**), CDR2 (**D**), CDR3 (**E**) and WH01 RBD. (**F**) The hydrogen bonds of the interface between Nb22 and WH01 RBD. The hydrogen bonds are shown in cyan dash line. (**G**) The hydrophobic network between Nb22 and WH01 RBD. All the residues are shown in sticks.

Figure 4. Structural analysis of Nb22 and Delta RBD complex. (**A**) The two mutation sites of Delta RBD. K478 is located outside the CDR binding regions, R452 is on the CDR2 recognized epitope. R452 and K478 are colored in cyan, and the epitope of CDRs is colored identical to Fig. 3. (**B**) The superimposition of WH01 RBD-Nb22 (orange and pink) and Delta RBD-Nb22 (light blue and yellow). (**C**) The hydrogen

743 bonds on the interaction interface of Delta RBD-Nb22. R452 and Q493 form two
744 additional hydrogen bonds with T30 and S33 of Nb22 The residues identified are shown
745 in sticks.
746
Figure 5. Characterizing Nb22-Fc stability in vitro and pharmaceutics in vivo. (**A**)
Binding curve of RBD with Nb22-Fc detected by ELISA after storage at room
temperature in the indicated time points including 0 d, 5 d, 10 d, 30 d, 40 d, 50 d and
60 d. (**B**) Binding curve of RBD with Nb22 detected by ELISA after the indicated
rounds of freeze-thawing including 0, 1, 3, 5 rounds, respectively. (**C**) Pharmacokinetic
of Nb22-Fc labeled with dye YF®750 SE via intranasal administration was detected.
200 µg Nb22-YF750 was infused at day 0, 7, 14 and 21 respectively. The optical
imaging of mice upper half body was measured by NightOwl LB 983 0.08 d (2h) post
Nb22-Fc infusion or at indicated time point labeled at the top of panel. The mice in the
red dash line figure were sacrificed at the indicated time point in the left of panel for
analysis of the fluorescence intensity of lung. (**D**)The fluorescence intensity of upper
half body of mice in (**C**) was summarized. (**E**)The fluorescence intensity of lung in lung
column of (C) was summarized. (**F**) Bioavailability and t1/2 of Nb22 in BALB/c
mice.Nb22 was intranasally (i.n.) administered into mice (n=3, Female) at 200 µg
(average of 10 mg/kg mice. Serum concentrations of the Nbs were determined at
various time points by ELISA. $T_{1/2}$, time of half-life. Data represent mean ± SEM.
763
764
Figure 6. The efficacy of Nb22s evaluated in hACE2 transgenic mice challenged
by SARS-CoV-2. (**A**) Experimental schedule of Nb22s in the prevention and treatment
of SARS-CoV-2 infection. Bottom, table summary of groups (n = 3–6 mice) with
different treatments. (**B**) Viral loads in lungs among 9 groups were measured by qRT-
PCR. The name of each group in the x axis was indicated as in the table in (A). Each
dot represents one mouse. The limit of detection was 1000 copies/mg referenced to
blank control (No-SARS-CoV-2 group). Data are represented as mean ± SEM; Mann-
Whitney test was performed to compare treatment group with the SARS-CoV-2 control
group. (**C**) Sections of lung were analyzed by immunofluorescence staining using
antibodies specific to SARS-CoV-2 NP in red and DAPI for nuclei in blue, respectively.
The fluorescence signal intensity of red was taken as a quantitative indicator for viral
infection, which was calculated by ImageJ software. ns, no significance; *p < 0.05, **p
< 0.01, ***p < 0.001. All experiments of (B) and (C) was repeated twice.
778
Figure 7. Representative sections of lung from hACE2 mice were analyzed by
H&E staining and immunofluorescence staining. Representative lung tissue sections
from the mice indicated in figure 6C including No-SARS-CoV-2, SARS-CoV-2,
Control Nb, -7d-Nb22, -5d-Nb22, -3d-Nb22, 1h-Nb22 and 1d-Nb22 group were
analyzed by immunofluorescence staining using antibodies specific to SARS-CoV-2
NP in red and DAPI for nuclei in blue, respectively. The corresponding representative
lung tissue sections were also analyzed by H&E staining. Immunofluorescence and HE
Images were visualized under the indicated bar.

Supplemental Materials

Supplemental Figure 1. Nb22 blocks the binding of hACE2 to WH01 RBD. (**A**) Overlap of Nb22 and hACE2 binding sites on WH01 RBD. hACE2 binding site on WH01 RBD is shown in cyan line. Nb22 binding site is shown in pink line. The overlap region is represented by ellipses with dashed lines. (**B**) The loop (V102-Y117) of Nb22 is clashed with the two helixes on N-terminal of hACE2. The loop is colored in red and helixes are colored in green.

Supplemental Figure 2. Spational distribution of Nb22 labeled with dye YF®750 SE. Mice were dissected and detected by NightOwl LB 983 after 200 μg Nb22-YF®750 SE infusion into mice as indicated in figure 5C. The fluorescence intensity was measured at 2 hours (0d-Nb22), 7d (7d-Nb22), 14d (14d-Nb22), 21d（21d-Nb22）and 28d（8d-Nb22） after infusion of Nb22 via i.n., respectively. Blank, blank mice without infusion of any antibody, was taken as blank control. The fluorescence intensity of various organs including trachea (Tr), lung (Lu), heart (H) , stomach(St), intestine (In), liver(Li), spleen (Sp), kidney (Ki), bladder (B), were analyzed

Supplemental Figure 3. Body weight of mice. Body weight of mice in figure 6 was recorded at the indicated time point.

Supplemental Table 1. Data collection and refinement statistics

Supplemental Table 2. Residues contributed to interaction between Nb22 and RBD were identified by PISA at the European Bioinformatics Institute.

Figures

Virus Name		IC$_{50}$ (ng/ml)			IC$_{80}$ (ng/ml)		
		Nb$_{15}$-Fc	Nb$_{22}$-Fc	Nb$_{31}$-Fc	Nb$_{15}$-Fc	Nb$_{22}$-Fc	Nb$_{31}$-Fc
Pseudovirus	WH01	0.57	1.01	0.64	1.38	1.76	1.7
	D614G	0.51	0.45	0.64	0.82	1.06	1.53
	Alpha	0.18	3.45	8.5	0.46	5.76	15.53
	Delta	13.9	0.41	1.96	42.42	1.44	11.55
	Epsilon	7.78	1.07	3.41	15.36	4.46	13.77
Live	Delta	195.6	12.3	34.97	403	21.26	56.71

Figure 1 Characterizing nanobodies neutralizing circulating variants of SARS-CoV-2. The neutralization curve of Nb15-Fc (**A**), Nb22-Fc (**B**), Nb31-Fc(**C**) and SNB02 (**D**) inhibiting SARS-CoV-2 pseudovirus of circulating variants. Nb-Fcs and SNB02 were all constructed as the format of VHH fused with human Fc1. SNB02 was taken as an antibody control specific for SFTS virus. (**E**) The summary curve of IC$_{50}$ of Nb-Fcs exhibiting potent neutralization against SARS-CoV-2 variants. (**F**) The neutralization potency of Nb-Fcs was evaluated based on authentic SARS-CoV-2 Delta variant plaque reduction neutralization test. (**G**) The summary table of IC$_{50}$ and IC$_{80}$ of Nb-Fcs in A-C and F, displaying potent neutralization. Data are represented as mean ± SD. All experiments were repeated at least twice.

Figure 2. Characterizing the binding of Nbs. Kinetic binding curve of Nb15-Fc (**A**),
Nb22-Fc (**B**) and Nb31-Fc (**C**) at the concentration 33.3 nM,11.1nM, 3.7nM and 1.2
nM with RBD of Delta variant, respectively, detected by BLI. Binding curves are
colored black, and fit of the data to a 1:1 binding model is colored red. (**D**)
Representative binding curve of various RBD as indicated to Nb22-Fc tested by BLI.
Nb22-Fc binding with RBD from representative SARS-CoV-2 variants detected by
immunofluorescence assay (**E**) and flow cytometric analysis (**F**), respectively. Mock
served as a cell control without plasmid transfection. Images were visualized under the
×10 objective. All experiments were repeated at least twice.

Figure 3. Structural analysis of Nb22 and WH01 RBD complex. (**A**) The overall
complex structure of Nb22 and WH01 RBD. The CDR1 (red), CDR2 (blue), CDR3
(green) of Nb22 (pink) and WH01 RBD (orange) are displayed in cartoon
representation. (**B**) The epitope of Nb22 shown in surface representation. The CDR
regions are colored in red, blue and green, respectively. The interaction between CDR1
(**C**), CDR2 (**D**), CDR3 (**E**) and WH01 RBD. (**F**) The hydrogen bonds of the interface
between Nb22 and WH01 RBD. The hydrogen bonds are shown in cyan dash line. (**G**)
The hydrophobic network between Nb22 and WH01 RBD. All the residues are shown
in sticks.

Figure 4. Structural analysis of Nb22 and Delta RBD complex. (**A**) The two
mutation sites of Delta RBD. K478 is located outside the CDR binding regions, R452
is on the CDR2 recognized epitope. R452 and K478 are colored in cyan, and the epitope
of CDRs is colored identical to Fig. 3. (**B**) The superimposition of WH01 RBD-Nb22
(orange and pink) and Delta RBD-Nb22 (light blue and yellow). (**C**) The hydrogen
bonds on the interaction interface of Delta RBD-Nb22. R452 and Q493 form two
additional hydrogen bonds with T30 and S33 of Nb22 The residues identified are shown
in sticks.

Figure 5. Characterizing Nb22-Fc stability in vitro and pharmaceutics in vivo. (**A**)
Binding curve of RBD with Nb22-Fc detected by ELISA after storage at room
temperature in the indicated time points including 0 d, 5 d, 10 d, 30 d, 40 d, 50 d and
60 d. (**B**) Binding curve of RBD with Nb22 detected by ELISA after the indicated
rounds of freeze-thawing including 0, 1, 3, 5 rounds, respectively. (**C**) Pharmacokinetic
of Nb22-Fc labeled with dye YF®750 SE via intranasal administration was detected.
200 μg Nb22-YF750 was infused at day 0, 7, 14 and 21 respectively. The optical

imaging of mice upper half body was measured by NightOwl LB 983 0.08 d (2h) post
Nb22-Fc infusion or at indicated time point labeled at the top of panel. The mice in the
red dash line figure were sacrificed at the indicated time point in the left of panel for
analysis of the fluorescence intensity of lung. (**D**)The fluorescence intensity of upper
half body of mice in (**C**) was summarized. (**E**)The fluorescence intensity of lung in lung
column of (**C**) was summarized. (**F**) Bioavailability and t1/2 of Nb22 in BALB/c
mice.Nb22 was intranasally (i.n.) administered into mice (n=3, Female) at 200 μg
(average of 10 mg/kg mice. Serum concentrations of the Nbs were determined at
various time points by ELISA. $T_{1/2}$, time of half-life. Data represent mean ± SEM.

Group	Name	Reagent	Numbers (n)	Antibody infusion i.n. Time	SARS-CoV-2 Challenge i.n. 0h
Group 1	No SARS-CoV-2	Blank	3	/	/
Group 2	SARS-CoV-2	PBS	5	/	√
Group 3	Control Nb	SNB02	3	1h	√
Group 4	Nb22 -7d	Nb22	6	-7d	√
Group 5	Nb22 -5d		6	-5d	√
Group 6	Nb22 -3d		5	-3d	√
Group 7	Nb22 -1d		5	-1d	√
Group 8	Nb22 1h		5	1h	√
Group 9	Nb22 1d		5	1d	√

**Figure 6. The efficacy of Nb22s evaluated in hACE2 transgenic mice challenged
by SARS-CoV-2.** (**A**) Experimental schedule of Nb22s in the prevention and treatment
of SARS-CoV-2 infection. Bottom, table summary of groups (n = 3–6 mice) with
different treatments. (**B**) Viral loads in lungs among 9 groups were measured by qRT-
PCR. The name of each group in the x axis was indicated as in the table in (A). Each
dot represents one mouse. The limit of detection was 1000 copies/mg referenced to
blank control (No-SARS-CoV-2 group). Data are represented as mean ± SEM; Mann-

Whitney test was performed to compare treatment group with the SARS-CoV-2 control
group. (**C**) Sections of lung were analyzed by immunofluorescence staining using
antibodies specific to SARS-CoV-2 NP in red and DAPI for nuclei in blue, respectively.
The fluorescence signal intensity of red was taken as a quantitative indicator for viral
infection, which was calculated by ImageJ software. ns, no significance; *$p < 0.05$, **p
< 0.01, ***$p < 0.001$. All experiments of (B) and (C) was repeated twice.

885
Figure 7. Representative sections of lung from hACE2 mice were analyzed by
H&E staining and immunofluorescence staining. Representative lung tissue sections
from the mice indicated in figure 6C including No-SARS-CoV-2, SARS-CoV-2,

Control Nb, -7d-Nb22, -5d-Nb22, -3d-Nb22, 1h-Nb22 and 1d-Nb22 group were
analyzed by immunofluorescence staining using antibodies specific to SARS-CoV-2
NP in red and DAPI for nuclei in blue, respectively. The corresponding representative
lung tissue sections were also analyzed by H&E staining. Immunofluorescence and HE
Images were visualized under the indicated bar.

894