Title:

Genetic epidemiology of blood type, disease and trait variants, and genome-wide

genetic diversity in over 11,000 domestic cats

Short title: Genetic epidemiology of over 11,000 cats

Authors:

Heidi Anderson[1], Stephen Davison[1], Katherine M. Lytle[1], Leena Honkanen[1], Jamie Freyer[1], Julia Mathlin[1], Kaisa

Kyöstilä[2,3,4], Laura Inman[1], Annette Louviere[1], Rebecca Chodroff Foran[1], Oliver P. Forman[1], Hannes Lohi[2,3,4],

Jonas Donner[1].

[1] Wisdom Panel Research Team, Wisdom Panel, Kinship, Portland, OR, USA

[2] Department of Medical and Clinical Genetics, University of Helsinki, Helsinki, Finland

[3] Department of Veterinary Biosciences, University of Helsinki, Helsinki, Finland

[4] Folkhälsan Research Center, Helsinki, Finland

17
18
19 # Abstract
20

In the largest DNA-based study of domestic cat to date, 11,036 individuals (10,419 pedigreed cats from 91

breeds and breed types and 617 non-pedigreed cats) were genotyped via commercial panel testing,

elucidating the distribution and frequency of known genetic variants associated with blood type, disease and

physical traits across cat breeds. Blood group determining variants, which are relevant clinically and in cat

breeding, were genotyped to assess the across breed distribution of blood types A, B and AB. Extensive panel

testing identified 13 disease-associated variants in 48 breeds or breed types for which the variant had not

previously been observed, strengthening the argument for panel testing across populations. The study also

indicates that multiple breed clubs have effectively used DNA testing to reduce disease-associated genetic

variants within certain pedigreed cat populations. Appearance-associated genetic variation in all cats is also

discussed. Additionally, we combined genotypic data with phenotype information and clinical documentation,

actively conducted owner and veterinarian interviews, and recruited cats for clinical examination to investigate

the causality of a number of tested variants across different breed backgrounds. Lastly, genome-wide

informative SNP heterozygosity levels were calculated to obtain a comparable measure of the genetic diversity

in different cat breeds.

This study represents the first comprehensive exploration of informative Mendelian variants in felines by

screening over 10,000 domestic cats. The results qualitatively contribute to the understanding of feline variant

heritage and genetic diversity and demonstrate the clinical utility and importance of such information in

supporting breeding programs and the research community. The work also highlights the crucial commitment

of pedigreed cat breeders and registries in supporting the establishment of large genomic databases that when

combined with phenotype information can advance scientific understanding and provide insights that can be

applied to improve the health and welfare of cats.

INTRODUCTION

Precision or genomic medicine, encompassing understanding of individual genetic variability in disease risk

and the customization of healthcare by patient subgroups, is now feasible and likely to become a future

standard-of-care in veterinary medicine and care of companion animals, including cats (1). Fueling this

development for felines specifically are improved genomic resources and technologies, combined with a

steady increase in the discovery of Mendelian disease and trait variants (2,3). Genetic testing is a form of

precision/genomic medicine when used for diagnosing diseases or traits of clinical relevance, as the results

are patient-specific and can potentially be used to tailor treatment to the disease and the patient (4). Direct-to-

consumer genetic testing is now readily available, and further empowers owners to proactively invest in

precision medicine for their pet.

There is also an opportunity for the insight gained from feline population-based genomic data to have a

transformational impact on felines and humans (1). Domestic cats are valued companions to humans, with

owners seeking high quality veterinary care to support long-term wellness and positive outcomes for disease

treatment. Cat genomes are structured in a similar way to humans, with 90% of their genes having a human

homologue; and cats and humans also suffer from many similar diseases (1). Domestic cats, like dogs, can

serve as naturally occurring models for many human diseases, in which the development of therapeutic treatments may be helpful for both veterinary and human patients (1,5–9).

While domestication of cats approximately 10,000 years ago did not introduce drastic genomic changes compared with undomesticated Felidae (10), pedigreed cats emerged about 150 years ago and do represent small genetically unique subpopulations of domestic cats. The genetic research of isolated populations in humans, dogs, and more recently cats has vastly improved identification of genetic disease and trait variants for use in genetic diagnostics and precision medicine. As an example, researchers have identified specific genetic variants of the *CMAH* gene that determine blood type in different cat breeds (11–15). Research on genetically isolated populations has further highlighted that one of the consequences of isolation -reduced genetic diversity- can manifest as an increase in the number of health conditions that are identified (16–19). For more than a decade, it has been common practice to eradicate disease-associated variants from pedigreed cat breeding populations using DNA testing. However, the focus on eradicating single DNA variants from a breed could contribute to severe loss of genetic diversity, especially if implemented strictly instead of thoughtfully (20).

Our previous work has elucidated that comprehensive screening of genetic variants in dogs is convenient and justified as it provides information to support breeding programs, veterinary care and health research (21). Further large-scale multiplex screening approaches were taken to characterize canine disease heritage and the relative frequency and distribution of disease-associated variants across breeds (22), and to explore the frequency of known canine appearance-associated variants among dog breeds (23). As we will demonstrate, such efforts are now equally feasible in cats and hold comparable promise for gaining insight into the genetic epidemiology of feline diseases and traits to better inform feline breeding decisions and establish the foundation for precision medicine of individuals, populations, and breeds.

This study represents the first comprehensive genetic evaluation of known feline disease and trait variants through the examination of 80 variants in 10,419 pedigreed cats and 617 non-pedigreed cats. These results provide a first glance into feline mutation heritage across nearly 100 breeds and breed types, and underscore the importance of large-scale population screening studies in improving veterinary diagnostics, breeding programs, and health recommendations for all cats.

Materials and Methods

Study Sample Population

The cat study population consisted of 10,419 pedigreed cat samples and 617 non-pedigreed cats, which was obtained during the development and provision of the commercially available MyCatDNA™ and Optimal Selection™ Feline tests (Wisdom Panel, Helsinki, Finland and Wisdom Panel, Vancouver, WA, USA, respectively) between 2016 and 2021. The 10,419 pedigreed cat samples represented 91 breeds and breed varieties, with 84 (89.3%) breeds and varieties represented by 5 or more individuals (S1 Table).

The breed of a cat was reported by its owner with accompanying registration information confirming the cat was registered with The International Cat Association (TICA) Fédération Internationale Féline (FiFe), The Cat Fanciers' Association (CFA), or World Cat Federation (WCF) standards. A few additional breeds not yet recognized by any major breed registry but with an established community of breed hobbyists were also considered breeds for the purposes of this study. The non-pedigreed cat sample set consisted of mixed breeds; breed crosses or random-bred cats. The tested cats were most often from the United States of America (54.9%), while cats from Finland (17.4%), Canada (5.3%), United Kingdom (3.5%), Norway (3.5%) and Sweden (3.3%), Russia (2.5%) and France (1%) represented other notable subgroups (>1% of the sample).

Ethics Statement

Feline DNA was obtained by Wisdom Panel as owner submitted, non-invasive cheek swab samples, and as cheek swab and blood samples collected at certified veterinary clinics in accordance with international standards for animal care and research. All cat owners provided consent for the use of their cat's DNA sample in scientific research. University's biobank samples were collected under the permit ESAVI/6054/04.10.03/2012 by the Animal Ethics Committee of the State Provincial Office of Southern Finland, Hämeenlinna, Finland.

Microarray development

A custom genotyping microarray for selected feline disease and trait associated variants (S2 Table) was developed based on the robust and widely utilized Illumina Infinium XT platform (Illumina, Inc., San Diego, CA,

USA), commercially available as the Complete for Cats™ / MyCatDNA™ / Optimal Selection™ Feline tests. The

microarray was designed and validated for use following the same protocol and principles as previously

described for canines (21). Firstly, public databases (3) and searches of the scientific literature were used to

identify likely causal variants for feline Mendelian disorders and traits. Secondly, genotyping assays for the

identified variants were designed according to the manufacturer's guidelines (Illumina, Inc.). At least three

technical replicates of each target sequence were included in the array design. Thirdly, the validation of each

specific disease and trait assay was completed with the use of control samples of known genotype, or synthetic

oligonucleotides in the case of rare conditions for which no control samples were available. In addition, owner-

provided photographs contributed to phenotypic validation of trait variant tests.

Genotyping

Microarray genotyping analyses were carried out following manufacturer-recommended standard protocols for

the Illumina XT platform (Illumina, Inc.). All genotype data from samples with call rates below 98% of the

analyzed markers were discarded in order to ensure high quality data, and all disease-associated variant calls

were confirmed by manual review. Selected disease-associated variant findings were genotyped using

standard capillary sequencing on an ABI3730xl DNA Analyzer platform (Thermo Fisher Scientific, Waltham,

MA, USA) as a secondary technology to provide further validation of results that were unexpected for the

breed. Sequencing was performed at the Sanger Sequencing Unit of the Finnish Institute of Molecular

Medicine (FIMM). The DNA extractions and PCR-product preparation and purification were carried out as

previously described in detail (21) using ~20 ng of genomic template DNA and an Amplitaq Gold Master Mix-

based protocol according to the manufacturer's instructions (Applied Biosystems, Waltham, MA, USA). The

heterozygosity (Hz) values for each individual and breed were calculated from the genotypes of 7,815

informative SNPs that differentiate between individuals and evenly spaced informative SNPs with a median

intermarker distance of ~312 kilobases.

Clinical validations

Medical history information for genetically affected cats was collected through interviews with cat owners and

veterinary clinicians. Clinical examinations were performed for confirmation of a Factor XII deficiency diagnosis

by collecting a blood sample for an activated partial thromboplastin time screening test through IDEXX

Laboratories (IDEXX Europe B.V., Hoofddorp, The Netherlands). Progressive Retinal Atrophy diagnosis was

confirmed via ophthalmic examination performed by an ECVO (European College of Veterinary

Ophthalmologists) board-certified veterinary specialist. A set of blood type determination data was available

through the records of the former Genlab (Niini, Helsinki, Finland). All additional phenotype information (clinical

or trait) and documentation was obtained through voluntary owner submissions.

Results

Overview of genotyping results

A total of 11,036 domestic cats, mainly pedigreed cats (N=10,419) representing 91 breeds and breed types

and an additional 617 non-pedigreed cats, were genotyped using a custom Illumina XT microarray. Genetic

screening included genotyping of 7,815 informative SNP markers across the genome and 82 variants

associated with blood type, diseases, and/or physical appearance; 78 of which were evaluated in the entire

11,036 cat study sample and four more recently included variants were screened for in a subset of 2,186

samples (19.8% of the study sample; S1 Table, and S2-S3 Tables). We observed 51 (62.1%) of the 82 tested

variants at least once in this study cohort, including 22 (44.9%) disease-associated variants and 29 (56.9%)

variants associated with an appearance-associated trait or blood type (Table 1, S3 Table). The maximum

number of disease-associated variants observed in any one individual was four. We observed that 2,480

(22.5%) of the tested cats had at least one disease-associated variant present, and 452 (4.1%) of the tested

cats were potentially at risk for at least one health condition, in accordance with the disorder modes of

inheritance. Of the disease-associated variants, 19 were observed in pedigreed cats, and 12 were also found

in non-pedigreed cats. Three disease-associated variants were solely observed in non-pedigreed cat samples

(Fig 1).

While we observed several disease-associated variants in the breeds with documented occurrence, we also

detected 17 disease-associated variants in additional breeds or breed types with no previously reported

frequency. These disease-associated variant discoveries and frequencies in additional breeds are listed in

Table 2.

In this study, we also investigated the tested variants' significance in additional breeds by combining genotype

data with information obtained from cat owners and/or veterinary clinicians through a variety of means,

including voluntarily submitted photographs, submission of clinical documentation, or interviews with a Wisdom

Panel veterinarian or geneticist. In addition, genetically affected cats were recruited for clinicopathological

investigations where possible. Finally, the variant frequencies were examined in the light of genome-wide

genetic diversity measured as a percentage of heterozygous SNPs.

Fig 1. Distribution of disease-associated variants within pedigreed and non-pedigreed cat populations.

Table 1. Tested disease-associated variants and their frequencies in all cats.

Tested Disease-associated Variant	MOI	Gene	Variant Details	Rank	Variant Allele Frequency (%)
Factor XII Deficiency	AR	*F12*	c.1631G>C	1.	7.01%*
Pyruvate Kinase Deficiency (PK-def)	AR	*PKLR*	c.693+304G>A	2.	2.94 %
Factor XII Deficiency	AR	*F12*	c.1321delC	1.	1.34 %
Progressive Retinal Atrophy (rdAc-PRA)	AR	*CEP290*	IVS50 + 9T>G	4.	1.13 %
Progressive Retinal Atrophy	AR	*KIF3B*	c.1000G>A	5.	1.11 %
Hypertrophic Cardiomyopathy; HCM (A31P)	AD	*MYBPC3*	c.91G>C	6.	0.85 %
MDR1 Medication Sensitivity	AR	*ABCB1*	11930_1931del TC	7.	0.58 %
Osteochondrodysplasia and Earfold	AD	*TRPV4*	c.1024G>T	8.	0.42 %
Hypertrophic Cardiomyopathy; HCM (R820W)	AD	*MYBPC3*	c.2460C>T	9.	0.20 %
Spinal Muscular Atrophy	AR	*LIX1 -LNPEP*	a ~140 kb partial gene deletion	10.	0.06 %
Polycystic Kidney Disease (PKD)	AD	*PKD1*	c.10063C>A	11.	0.05 %
Congenital Myasthenic Syndrome	AR	*COLQ*	c.1190G>A	12.	0.05 %
Cystinuria Type B	AR	*SLC7A9*	c.881T>A	13.	0.05 %
Hyperoxaluria Type II	AR	*GRHPR*	point mutation, G to A, at the 3# splice acceptor site of intron 4	14.	0.04 %

Disease	MOI	Gene	Variant		Frequency
Familial Episodic Hypokalemic Polymyopathy	AR	WNK4	c.2899C>T	15.	0.02 %
Lipoprotein Lipase Deficiency	AR	LPL	c.1234G>A	16.	0.02 %
Burmese Head Defect	AR	ALX1	c.496delCTCTCAGGACTG	16.	0.02 %
GM2 Gangliosidosis Type II	AR	HEXB	c.1244-8_1250del15	18.	0.01 %
Glycogen Storage Disease	AR	GBE1	g. IVS11+1552_IVS12-1339 del6	18.	0.01 %
Myotonia Congenita	AR	CLCN1	c.1930+1G>T	20.	0.005 %
Acute Intermittent Porphyria (5 variants)	AD	HMBS	c.842_844delGAG, c.189dupT, c.250G>A, c.107_110delACAG, c.826-1G>A		NA
Autoimmune Lymphoproliferative Syndrome	AR	FASL	c.413_414insA		NA
Congenital Adrenal Hyperplasia	AR	CYP11B1	Exon 7 G>A		NA
Congenital Erythropoietic Porphyria	AR	UROS	c.331G>A + c.140C>T		NA
Cystinuria Type 1A	AR	SCL3A1	c.1342C>T		NA
Cystinuria Type B (2 variants)	AR	SCL7A9	c.706G>A, c.1175C>T		NA
Dihydropyrimidinase Deficiency	AR	DPYS	c.1303G>A		NA
Glutaric Aciduria Type II	AR	ETFDH	c.692T>G		NA
GM1 Gangliosidosis	AR	GLB1	c.1448G>C		NA
GM2 Gangliosidosis	AR	GM2A	c.390_393GGTC		NA
GM2 Gangliosidosis Type II (2 variants)	AR	HEXB	c.1467_1491inv, c.667C>T		NA
Hemophilia B (2 variants)	X-linked	F9	c.247G>A, c.1014C>T		NA
Hypotrichosis	AR	FOXN1	c.1030_1033delCTGT		NA
Mucopolysaccharidosis Type I	AR	IDUA	c.1107_1109delCGA / c.1108_1110GAC		NA
Mucopolysaccharidosis Type VI	AR	ARSB	c.1558G>A		NA
Mucopolysaccharidosis Type VII (2 variants)	AR	GUSB	c.1052A>G, c.1421T>G + c.1424C>T		NA
Progressive Retinal Atrophy	AR	AIPL1	c.577C>T		NA
Sphingomyelinosis; Niemann-Pick C1	AR	NPC1	c.2864G>C		NA
Sphingomyelinosis; Niemann-Pick C2	AR	NPC2	c.82+5G>A		NA
Vitamin D-Dependent Rickets	AR	CYP27B1	c.637G>T		NA
Potential Non-causal Variant[#]					
Mucopolysaccharidosis Type VI Modifier		ARSB	c.1427T>C		2.51 %
Congenital Erythropoietic Porphyria		UROS	c.140C>T		0.74 %

*The frequency value is based on a subset of 2,186 samples (19.8% of the full study sample) screened for this variant. # Genetic screening for these variants is available, but the predictive value of disease is low. Abbreviations: MOI; mode of inheritance, AR; autosomal recessive, AD; autosomal dominant, NA; not applicable.

178 Table 2. Summary of disease-associated variant findings in additional breeds.

Tested Disease-associated Variant	Breed(s) variant was previously characterized in	Additional breed(s) with variant allele frequency (No. of cats tested)
Cystinuria Type B	non-pedigree cat Maine Coon Sphynx	Maine Coon Polydactyl 1% (150) Siberian 0.4% (559)
Factor XII Deficiency; F12 c.1631+G>C*	non-pedigree cat Bengal Maine Coon Siamese	Balinese 16.67% (76) Bombay 50.00% (5) British Shorthair 2.08% (347) Cashmere 33.33% (3) Devon Rex 8.76% (447) Donskoy 75.00% (4) Elf 6.25% (8) Exotic Shorthair 22.22% (9) Highlander 2.12% (94) Himalayan 50.00% (2) Lykoi 7.89% (19) Maine Coon Polydactyl 1.16% (43) Minuet 12.50% (4) Minuet Longhair 13.64% (11) Munchkin 3.33% (15) Munchkin Longhair 20.00% (11) Neva Masquerade 4.17% (12) Oriental Longhair 25.00% (2) Oriental Shorthair 8.33% (12) Persian 10.00% (5) Peterbald 33.33% (6) Ragdoll 6.04% (298) Savannah 25.00% (12) Scottish Fold Shorthair 4.84% (31) Scottish Straight 2.38% (21) Scottish Straight Longhair 10.00% (5) Selkirk Rex 23.53% (17) Selkirk Rex Longhair 18.75% (16) Siberian 3.09% (97) Sphynx 4.55% (11) Tennessee Rex 42.31% (13) Turkish Angora 19.23% (13)

Disease	Breed	Other breeds / populations
Factor XII Deficiency; F12 c.1321delC	non-pedigree cat Bengal Maine Coon Siamese	American Shorthair 1.04% (49) Balinese 0.66% (76) Cymric 6.25% (16) Highlander 0.48% (201) Himalayan 3.13% (16) Maine Coon Polydactyl 3.02% (150) Manx 3.45% (30) Minuet 4.55% (12) Munchkin 2.63% (40) Munchkin Longhair 5.00% (11) Ragdoll 0.05% (1115) Savannah 3.21% (80) Tennessee Rex 43.75% (35)
GM2 Gangliosidosis Type II	Burmese	non-pedigree cat 0.08% (617)
Hypertrophic Cardiomyopathy; HCM (A31P)	Maine Coon	Maine Coon Polydactyl 3.67% (150) non-pedigree cat 0.16% (617) Pixiebob Longhair 4.17% (12)
Hypertrophic Cardiomyopathy; HCM (R820W)	Ragdoll	American Bobtail Longhair 5.71% (35) American Bobtail Shorthair 5.56% (9) Highlander 2.12% (201) Munchkin 1.25% (40) RagaMuffin 0.85% (118) non-pedigree cat 0.08% (617)
MDR1 Medication Sensitivity	non-pedigree cat Ragdoll Russian Blue Siamese	Balinese 1.32% (76) Maine Coon 2.79% (1971) Maine Coon Polydactyl 3.33% (150) Turkish Angora 0.45% (110)
Osteochondrodysplasia and Earfold	Scottish Fold	non-pedigree cat 0.6% (617)
Polycystic Kidney Disease (PKD)	>10 breeds	Maine Coon 0.13% (1971) Scottish Straight 0.82% (61)
Progressive Retinal Atrophy	Bengal	Highlander 3.48% (201) Highlander Shorthair 3.33% (30) Savannah 0.63% (80) non-pedigree cat (Bengal Mix) 0.08% (617)

Progressive Retinal Atrophy (rdAc-PRA)	>10 breeds	American Shorthair 1.02% (49) Devon Rex 1.68% (447) Donskoy 2.94% (17) European Shorthair 0.55% (91) Havana Brown 1.67% (30) Highlander 0.48% (201) Manx 5% (30) non-pedigree cat 1.13% (617) Oriental Longhair 19.61% (51) Pixiebob Longhair 4.17% (12) Ragdoll 0.49% (1115) Savannah 6.25% (80) Scottish Fold Shorthair 1.32% (76)
Pyruvate Kinase Deficiency (PK-def)	>10 breeds	American Bobtail Shorthair 5.56% (9) Caracat 50.00% (1) Chausie 8.33% (6) European Shorthair 0.55% (91) Highlander 1.99% (201) Highlander Shorthair 3.33% (30) Lykoi 3.37% (104) Maine Coon Polydactyl 10.67% (150) Minuet Longhair 4.17% (24) Munchkin 2.56% (40) Neva Masquerade 2.17% (23) Pixiebob 13.16% (21) Pixiebob Longhair 18.18% (12) Ragdoll 0.09% (1115) Sphynx 0.09% (547) Toyger 3.85% (13) Turkish Angora 0.48% (110)
Spinal Muscular Atrophy	Maine Coon	Highlander 0.48% (201) Maine Coon Polydactyl 0.33% (150)

*The frequency value is based on a subset of 2,186 samples (19.8% of the full study sample) screened for this variant.

Genetic epidemiology of the common AB blood group system across breeds and breed

types

The major feline AB blood groups including blood type A, blood type B and the rare blood type AB are caused

by functional differences in the cytidine monophospho-N-acetylneuraminic acid hydroxylase enzyme encoded

by the *CMAH* gene (11–15). According to current convention (2019 typing panel) (14,15), genetic testing for

blood types A, B and AB should be based on panel testing of four likely causal variants of *CMAH* to assist

veterinary clinicians and breeders in recognizing, confirming, and avoiding blood incompatibilities. In this study,

a subset of 2,179 cats (19.7%) was genotyped for all four *CMAH* variants of the proposed genotyping scheme:

common b variant 1, c.268T>A; b variant 2, c.179G>T (discovered in Turkish breeds); the c variant, c.364C>T

resulting in blood type AB; and the b variant 3, c.1322delT (discovered in the Ragdoll), which was more recently

added to the WISDOM PANEL genotyping platform. All 11,036 cats were genotyped for the first three variants.

We observed b variant 1, b variant 2; and the c variant to be widely distributed across breeds with frequencies

of 12.6%, 1.6% and 1.5% in all cats, respectively (S4 Table). No more than two variants in total were found in

any individual cat. Across the 2,179 genotyped cats representing 69 breeds and varieties, the b variant 3 was

exclusively found in Ragdolls (16.9% allele frequency) and in a Ragdoll mix (S4 Table). Since the b variant 3

was found only in cats with Ragdoll ancestry, applying the assumption that the entire dataset was consistent

with this finding allowed genotypic interpretations for blood type determination to be based on three variants

(the b variant 1 and 2 and the c variant) in all breeds except for Ragdoll, in which a four variant-based

interpretation of genotype (including the b variants 1, 2, and 3, and the c variant) was used.

Based on genetic blood type determination, blood type B was most common in the following five breeds or

breed types: American Curl (40.4%), British Shorthair breed types (20.3%), Cornish Rex (33%), Devon Rex

(30.3%) and Havana Brown (20%). In addition, the breeds in which the rare blood type AB was present with a

frequency of >1% were European Shorthair (2.2%), Lykoi (1%), Scottish Fold (3.3%), RagaMuffin (3.4%),

Ragdoll (2.2%) and Russian Blue breed types (1.5%). The proportions of type A, type B and type AB blood for

each breed or breed group with >15 individuals tested are shown in Table 3 (breed groups listed in S5 Table).

In this study, blood typing results from serological tests were available for 220 cats. We found that blood types

assigned according to the cat's genotype and serological tests were 99.5% concordant. One Ragdoll with c/c

genotype shown to result in blood type AB (11,14,15), had been determined serologically to have blood type

A according to the cat's owner, although retesting to confirm this result or further clinical investigation was not

performed.

Table 3. Genetically determined proportions of type A, type B and type AB blood for each breed or breed type

with >15 individuals tested.

Breeds and breed types	No. of tested cats	Type B (b/b)	Type AB (c/b, c/c)	Type A (A/A, A/b, A/c)
American Curl	47	40.4 %	0.0 %	59.6 %
Cornish Rex	106	33.0 %	0.9 %	66.0 %
Devon Rex	446	30.3 %	0.0 %	69.7 %
British Shorthair types	395	20.3 %	0.0 %	79.7 %
Havana Brown	30	20.0 %	0.0 %	80.0 %
American Bobtail	44	15.9 %	0.0 %	84.1 %
Toybob	56	12.5 %	0.0 %	87.5 %
Turkish Van	40	12.5 %	0.0 %	87.5 %
Scottish Fold	90	8.9 %	3.3 %	87.8 %
Scottish Straight	75	12.0 %	0.0 %	88.0 %
Turkish Angora	110	11.8 %	0.0 %	88.2 %
Birman	174	10.9 %	0.0 %	89.1 %
Chartreux	84	8.3 %	0.0 %	91.7 %
Sphynx	547	7.9 %	0.0 %	92.1 %
Lykoi	104	5.8 %	1.0 %	93.3 %
Pixiebob	33	6.1 %	0.0 %	93.9 %
European Shorthair	91	3.3 %	2.2 %	94.5 %
Selkirk Rex	121	5.0 %	0.0 %	95.0 %
Exotic Shorthair	68	4.4 %	0.0 %	95.6 %
RagaMuffin	118	0.8 %	3.4 %	95.8 %
non-pedigree cat	616	3.6 %	0.6 %	95.8 %
Ragdoll	1115	1.7 %	2.2 %	96.1 %
Munchkin breed types	109	2.8 %	0.0 %	97.2 %
Manx breed types	46	2.2 %	0.0 %	97.8 %
American Shorthair	49	2.0 %	0.0 %	98.0 %
Siberian breed types	582	1.9 %	0.0 %	98.1 %
Russian Blue breed types	65	0.0 %	1.5 %	98.5 %
Persian breed types	136	1.5 %	0.0 %	98.5 %
Burmese breed types	161	1.2 %	0.0 %	98.8 %
Maine Coon	2121	0.8 %	0.0 %	99.2 %
Highlander	231	0.4 %	0.0 %	99.6 %
Siamese breed types	233	0.4 %	0.0 %	99.6 %
Bengal	1706	0.0 %	0.1 %	99.9 %
Abyssinian breed types	214	0.0 %	0.0 %	100.0 %
Donskoy	17	0.0 %	0.0 %	100.0 %
Egyptian Mau	55	0.0 %	0.0 %	100.0 %
Khaomanee	21	0.0 %	0.0 %	100.0 %
Korat	51	0.0 %	0.0 %	100.0 %
LaPerm	35	0.0 %	0.0 %	100.0 %
Norwegian Forest Cat	121	0.0 %	0.0 %	100.0 %
Ocicat	76	0.0 %	0.0 %	100.0 %

Oriental breed types	230	0.0 %	0.0 %	100.0 %
Peterbald	17	0.0 %	0.0 %	100.0 %
Savannah	80	0.0 %	0.0 %	100.0 %
Singapura	39	0.0 %	0.0 %	100.0 %
Tennessee Rex	35	0.0 %	0.0 %	100.0 %
Tonkinese	41	0.0 %	0.0 %	100.0 %

Factor XII Deficiency and Pyruvate Kinase Deficiency are widespread blood disorders in the cat population

Factor XII Deficiency is a widely distributed heritable disorder in the domestic cat population (24). Factor XII Deficiency is a clinical hemostatic defect that manifests as a prolonged activated partial thromboplastin time (aPTT) which would be observed in a presurgical coagulation assay, but does not require transfusions (25). Two variants of the *F12* gene (c.1321delC and c.1631G>C) have been identified in a colony of inbred cats from the United States and in a litter of cats from Japan, respectively. Both variants segregate consistently with cases of Factor XII Deficiency. The variants also co-segregate, although variant c.1631G>C is likely to have originated first, as it is observed without the c.1321delC variant. These variants are both considered common in the domestic cat (25–27). The most severe aPTT prolongation is observed in cats homozygous for both variants (25). In accordance with the previous observations (25), we noted that the c.1321delC variant always co-segregates with c.1631G>C, while one or two copies of the latter can be inherited in the absence of the c.1321delC variant. The observed variant frequencies for the c.1321delC and c.1631G>C variants were 1.3% and 7% in all cats (S4 Table). The frequency of the c.1631C variant was based on a subset of 2186 genotyped cats, as the variant represented a more recent discovery added to the WISDOM PANEL genotyping platform. Notably, the c.1631G>C variant present alone without the second variant is observed at high frequency in the Donskoy (75%), Bombay (50%) and Himalayan (50%) breeds, while Tennessee Rex represents a breed with high frequency for both variants (>40%) observed together (S4 Table). In all, we discovered the two variants of *F12* gene present in 40 additional breeds. Our data confirm the presence of the identified *F12* variants in the following breeds in which cases of Factor XII Deficiency have been documented: Himalayan, Maine Coon, Manx, Munchkin, Oriental Shorthair, Persian, Ragdoll, Siberian and Siamese. The c.1321delC variant was not discovered in any of the 121 genotyped Norwegian Forest Cats, nor was the c.1631G>C variant (examined in 16 Norwegian Forest Cats) despite documented cases of Factor XII Deficiency in the breed (25). Consequently, it may be possible that these cats are affected by other candidate

variants of *F12* such as the c.1549C>T variant, which was recently identified (25), but not screened in this study. Furthermore, we could not confirm the tested variants' presence in the Turkish Van. We, therefore, could not corroborate the documented association with Factor XII Deficiency (25), though we did discover the presence of the c.1631G>C variant in the related Turkish Angora breed. Finally, we obtained laboratory results from a 10-month-old intact female Maine Coon homozygous for the c.1321delC and c.1631G>C variants of *F12* gene. This individual showed prolonged aPTT of >180 seconds (laboratory reference <13.4 seconds), confirming Factor XII Deficiency in the Maine Coon and the association of the tested variants with clinical signs.

Pyruvate Kinase Deficiency (PK-def) is an inherited anemia characterized by low levels of the pyruvate kinase enzyme. This insufficient presence of the PK enzyme causes red blood cells to break easily, resulting in hemolytic anemia. A single nucleotide substitution variant (c.693+304G>A) of intron 5 in the genomic DNA splice site of the *PKLR* gene has been associated with the manifestation of PK-def in the Abyssinian and Somali breeds (28), but has also been previously identified in at least 15 additional cat breeds. PK-def associated with the identified variant has marked clinical variability, including variation in the age of disease onset and disease severity (28,29). Here we report the presence of the *PKLR* variant in an additional 35 breeds and breed types (Table 1). Where the breed was represented by at least ten individuals in our study set, the breeds with the highest allele frequencies were the Pixiebob (16.6%), Pixiebob Longhair (13.5%), Egyptian Mau (12.7%) and Maine Coon Polydactyl (10.5%). The updated *PKLR* variant frequencies for the Abyssinian and Somali breeds were 3.1% and 2.2%, respectively (S4 Table). Several young cats from Bengal, Maine Coon and Maine Coon Polydactyl breeds had two copies of the PK-def associated variant and are therefore clinically at risk. In the scope of this study, we were able to interview the owner of a 3-year-9-month-old female Bengal cat. The owner reported a veterinarian had obtained a complete blood count (CBC) for this cat, however, the test results at that time did not reveal the presence of anemia. Additionally, we reached out to several breed clubs to ask whether clinical PK-def has been documented in the Maine Coon, but discussions were inconclusive. Further clinical data are required to confirm clinical signs manifesting in the additional breeds identified with the *PKLR* variant.

Panel screening reveals autosomal dominant disease-associated variants affecting cats in additional breeds

We screened for four feline disease-associated variants that most closely follow an autosomal dominant mode

of inheritance in clinical settings: Polycystic Kidney Disease (PKD) (30), two Hypertrophic Cardiomyopathy

(HCM) variants (31,32), and Osteochondrodysplasia and Earfold (33).

Polycystic Kidney Disease (PKD) is a severe autosomal dominant (homozygous lethal) condition in which

clusters of cysts present at birth develop in the kidney and other organs, causing chronic kidney disease which

can lead to kidney failure (34,35). PKD is caused by a stop codon in exon 29 of *PKD1* (c.10063C>A), resulting

in a truncated form of the gene, which was discovered in Persian cats with ~40% frequency in the Persian cat

population worldwide (36–39). Genetic testing was introduced into the breeding programs of Persians and

some Persian-related cats. We report that the overall frequency of the *PKD1* variant in the screened breeds

has reduced notably from what was previously reported (S4 Table). The *PKD1* variant was identified in the

Maine Coon, a breed in which it had not been previously documented in the peer-reviewed literature. Clinical

manifestation of PKD in a genetically affected female Maine Coon, diagnosed at the age of 3 months, was

confirmed after interviewing the cat's owner and assessing associated diagnostic documentation including an

ultrasound of the kidneys in which numerous, round, well-defined cysts were observed bilaterally throughout

the renal cortex and medulla. This finding provides confirmation that genetic screening for the *PKD1* variant in

the Maine Coon is a clinically relevant marker for PKD.

Hypertrophic cardiomyopathy (HCM) is the most common heart disease in domestic cats. Two independent

condition predisposing variants of the *MYBPC3* gene c.91G>C p.(A31P) and c.1024G>T p.(R820W) have

been associated with HCM in Maine Coon and Ragdoll breeds, respectively (31,32,40). In the heterozygous

state, the likelihood of developing clinical HCM early in life is very low. However, supporting the autosomal

dominant mode of inheritance, regional diastolic and systolic dysfunction has been observed in heterozygous

asymptomatic cats (41,42). In the homozygous state, the development of HCM (A31P) is highly likely in the

Maine Coon with risk increasing with age (43). Similarly, in Ragdoll cats the (R820W) heterozygous cats have

a normal life expectancy, while homozygous cats are likely to have a shortened life span (44). In the present

study, we found higher frequencies of HCM (A31P) and HCM (R820W) variants present in additional breeds

(Table 1). The observed cats were young, and all were carrying one copy of either of the variant alleles; no

further clinical validations were pursued in the scope of this study. Yet, given the association of these likely

causal variants with HCM, these variants suggest potential molecular explanations for cases of HCM in these

additional breeds.

Osteochondrodysplasia and Earfold is a highly penetrant autosomal dominant condition caused by a missense

variant (c.1024G>T) in the *TRPV4* gene resulting in congenital degenerative osteochondrodysplasia or

"Scottish Fold Syndrome", which results in skeletal deformities such as a short, thick, inflexible tail and

malformation of the distal fore- and hindlimbs, which can lead to a stilted gait (33). We observed at least one

copy of the *TRPV4* variant in all 90 tested Scottish Fold cats, and the *TRPV4* variant was absent in all 75

Scottish Straight cats tested. We also discovered one copy of the *TRPV4* variant in a crossbred cat resulting

from the mating of a Scottish Fold and a Highlander. As the ear phenotype of the kitten was curled-back as

presented in the Highlander breed, rather than folded forward as seen in the Scottish Fold, observation of the

TRPV4 variant was not entirely expected. However, the kitten did present a stiff and inflexible shortened tail,

characteristic of *TRPV4* variant carriers. It therefore would appear that the yet unknown variant that is causing

the Highlander ear type masks the Scottish Fold ear phenotype caused by the *TRPV4* variant, when the two

variants are inherited together. In another recent study, a cat registered as an American Curl with curled ears

was diagnosed with osteochondrodysplasia and genotypically showed one copy of *TRPV4* variant (45). Thus,

we report a second case of Osteochondrodysplasia in which the cat's ear phenotype belied the presence of

the causal variant.

Molecular heterogeneity of feline hereditary retinal dystrophies

The disease-associated variants for retinal dystrophies screened in this study include *CEP290*, *KIF3B* and

AIPL1. The *CEP290* variant associated with late-onset Progressive Retinal Atrophy (rdAc-PRA) is present in

many pedigreed breeds (46,47); here we document a frequency of 1.1% in all cats. We have identified the

presence of the *CEP290* variant in 20 additional breeds. Some of the highest *CEP290* variant frequencies

were observed in the Peterbald (26.5%) and in one of the additionally identified breeds, the Oriental Longhair

(19.6%). The variant of the *KIF3B* gene, recently associated with an early-onset Progressive Retinal Atrophy

in the Bengal (48), was present in 6.9% of the Bengal breed (resulting in a frequency of 1.1% in all cats) (Table

1-2, and S4 Table). This variant was additionally discovered in the Highlander breed types and the Savannah.

The *AIPL1* variant associated with Progressive Retinal Atrophy (discovered in the Persian) was the rarest

variant associated with retinal dystrophies (49), that was screened in a subset of 2,186 samples and not

observed at all (Table 1, S4 Table). To pursue clinical validation of our findings, we recruited a 10-year-3-

month-old female Oriental Longhair, homozygous for *CEP290*. Clinical validation was initiated with an owner

interview, in which the owner reported no apparent changes in the cat's behavior that were suggestive of vision

loss. Yet, during an ophthalmic examination, a marked discoloration of pigmentation of the tapetal fundus with

a slight vascular attenuation was noted, confirming the presence of retinal degeneration. We demonstrate that

rdAc-PRA does manifest clinically in the Oriental Longhair. Some vision may be retained longer than the

previously reported 3-7 years (5), at least for this particular breed example.

Feline MDR1 Medication Sensitivity associated with adverse medication reactions in the

Maine Coon

Feline MDR1 Medication Sensitivity is a disorder associated with severe adverse reactions after exposure to

medications that use the p-glycoprotein drug transporter. This genetic condition is caused by a two base pair

deletion within exon 15 of the *ABCB1* gene resulting in abnormal p-glycoprotein (50). While functional p-

glycoprotein plays a significant part in the blood-brain barrier that prevents various drugs and chemicals in the

bloodstream from entering the brain, a defective p-glycoprotein allows more drugs to cross this barrier, thus

increasing the neurological effects of some medications. Severe macrocyclic lactone-induced neurologic

toxicosis has previously been reported in cats homozygous for the MDR1 variant receiving either a

subcutaneously administered dose of ivermectin or a topically administered eprinomectin-containing

antiparasitic product labeled for cats (50,51). We report the frequency of the *ABCB1* variant as 0.6% of all

genotyped cats, in addition to the discovery of the variant in the Maine Coon, Ragdoll, Siamese and Turkish

Van breeds (Tables 1-2; S4 Table). Our veterinarians interviewed three owners of cats identified as

homozygous for the MDR1 variant to assess their cat's medical history. The cats consisted of a 1-year-4-

month-old intact female Maine Coon, a 2-year-3-month-old intact female Ragdoll, and a 3-year-2-month-old

male Maine Coon. Only one of the cats had undergone anesthesia, and reportedly showed a delayed recovery

with mild lethargy documented on the following day, but fully recovered. All three cats had been administered

topical flea medications (of varying brands) with no discernable side effects, however, none of the medications

applied contained eprinomectin.

Genetic diagnosis plays a crucial role in the diagnosis of uncommon inherited disease

We identified a cat genetically affected with Myotonia Congenita, an uncommon recessively inherited disorder

manifesting as an inability of the muscles to relax after contraction, which is caused by a variant in the *CLCN1*

gene (52). This is a sporadic condition that was discovered in a rescue domestic cat population in Winnipeg,

Canada. While the variant was not identified in any pedigreed cats (which make up a large proportion of the

study sample), we discovered two copies of the variant in a single non-pedigreed domestic cat in Oregon,

United States. In the owner interview, we learned that the genetic diagnosis was crucial in assisting with clinical

diagnosis. While this is an incurable condition, having the correct diagnosis helps ensure that the cat is getting

appropriate supportive care. The owner confirmed that this cat, initially misdiagnosed with flea bite anemia at

the age of 13 weeks, has a disease manifestation that includes fainting spells when startled, prolonged

prolapse of the nictitating membrane, hypertrophic musculature, flattened ears, motor dysfunction, shortened

gait, and limited range of motion in the jaw. The cat also shows a characteristic "smile" after a yawn or a meow

due to delayed relaxation of the muscle in the upper lip as well as commonly has the paws protracted (Fig 2A).

However, the owner mentioned that this cat, currently 4-year-2-month-old, is not drooling or showing any dental

defects, which differs from the original disease description (52).

Panel screening enables dissociation of variants with clinical disease

In this study we determined the frequency of the c.140C>T p.(S47F) variant of the *UROS* gene to be 0.7%

across the entire study sample (Table 1, S4 Table). This variant was previously discovered along with

c.331G>A p.(G111S) in a cat manifesting Congenital Erythropoietic Porphyria (CEP) (53). However, we did

not observe the c.331G>A variant in any of the tested cats. While the original research revealed no cats

carrying solely the c.140C>T variant, functional studies have shown that c.140C>T alone does not significantly

alter the protein function (53). The manifestation of CEP includes distinctively stained brownish-yellow teeth

that turn fluorescent pink under UV light. Due to the high frequency of c.140C>T in some cat breeds and

because some DNA testing laboratories offer tests for the two variants separately, our veterinarians reached

out to the owners of cats homozygous for only the c.140C>T variant. Owners were contacted in the cases of

five c.140C>T homozygous cats: a 1-year-4-month-old intact female RagaMuffin, a 1-year-5-month-old intact

male RagaMuffin, a 1-year-7-month-old intact female Siberian, a 2-year-10-month-old intact female Singapura

and a 3-year-7-month-old intact Toybob; none of these cats had clinical signs suggestive of CEP. Thus, the

existing evidence strongly suggests that c.140C>T is a benign variant when it is not inherited along with

c.331G>A. We would advise breeders and DNA laboratories not to consider c.140C>T alone a precursor for

CEP.

Mucopolysaccharidosis Type VI (MPS VI), a lysosomal storage disease caused by a deficiency of *N*-

acetylgalactosamine-4-sulfatase (4S), is another disease in which the roles of two variants, independently

inherited in this case, have been under discussion (54). While the MPS VI variant c.1427T>C p.(L476P) of the

ARSB gene is associated with severe disease (55), the c.1558G>A p. (D520N) variant of the *ARSB* gene is

sometimes referred to as the "mild" type (56). However, it is necessary for the (D520N) variant to be inherited

as a compound heterozygote with one copy of the L476P variant for the disease to manifest in its mild form.

We have elected to call the D520N variant MPS VI Modifier. The D520N variant is present in large numbers

of cats, with a frequency of 2.5% (Table 1, S4 Table). It is also the major allele in the Havana Brown, with an

allele frequency of 76.7%. Additionally, we confirm that MPS VI (L476P) is a scarce variant, and this study

cohort did not identify any cats with MPS VI variant in concordance with previously described observations

(54). The high variant frequency of the benign MPS VI Modifier and very low frequency of the MPS VI variant

(absent in many breeds) further confirm that breeding to avoid MPS VI should concentrate solely on managing

the MPS VI variant.

Appearance-associated variants distributed across breeds and breed types

In this study sample, the ancestral form of the appearance-associated (trait) variant was the major allele,

except for the *ASIP* gene in which the derived allele a (non-agouti/solid color) showed a 56.2% variant

frequency in all cats (S4 Table). The rarest derived allele was the e allele of the *MC1R* gene associated with

the coat color Amber (discovered in the Norwegian Forest Cat), only observed in one random-bred cat from

Finland. The observed cat was homozygous for the derived allele with pictures confirming the expression of

the Amber coat color (Fig 2B).

Expectedly, the screened trait variants were present in a high number of breeds and breed types (S4 Table).

Genotyping results showed overall concordance with observed phenotypes, strongly supporting the variants'

causality, with one exception where our findings are supportive of the potential additional candidate locus may

play a role in the regulation of gloves phenotype (59). In Birman cats, the two adjacent missense variants of

the *KIT* gene associated with breed-defining white gloving phenotype (58) were observed with a high variant

frequency of 95.6% in the breed. These variants were also observed as the minor allele in a large number of

breeds and breed types, and in homozygote state in some individuals of Chartreux, Highlander, Maine Coon,

Ragdoll and Siberian breeds, in which they were not associated with gloving based on photographic evidence

and owner reports. Of the phenotyped individuals some white areas of hair were seen in the belly or toes in

(3/17) Maine Coons and (1/2) Siberian cats.

The polydactyly variant *Hw* was the most typical *LIMBR1* gene variant observed in polydactylous cats,

including the Maine Coon, Pixiebob and Highlander breed types, and some non-pedigreed cats from North

America. The *Hw* variant (variant 1) of *LIMBR1* was observed to be incompletely penetrant, with cats

presenting with four to seven toes per paw. Higher penetrance without a more extreme phenotypic

manifestation was seen in homozygous cats, which is in line with previous observations (59). However, extra

toes did not manifest solely in the front feet as previously reported; photographic evidence and owner-provided

details revealed the presence of extra toes on all four feet, which was formerly considered to be characteristic

of the *UK1* (variant 2) and *UK2* (variant 3) variants only (60) (Fig 2C). The only cat with two copies of the *UK1*

variant had two extra digits on each paw, per the owner's description. Various owner-reported cases of

polydactyl cats testing negative for the screened variants were also noted. Such cats are likely to be carrying

additional variants of the *LIMBR1* gene not yet identified.

Additional variants are also likely to be associated with shortened tail phenotype. The c.5T>C variant in the

gene *HES7* (discovered in the Japanese Bobtail) associated with a shortened and kinked tail (61) was

observed in 100% of the Japanese Bobtail cats studied, and was also highly prevalent in the Kurilian Bobtail

(96.4%) and Mekong Bobtail (66.7%), with some minor allele frequencies observed in additional bobtail breeds

(S4 Table). All studied cats with one or more copies of *HES7* variant and available phenotypic information

presented with a shortened tail.

The three tested variants (c.998delT, c.1169delC and c.1199delC) of the *T-box* gene (discovered in the Manx)

were found in the American Bobtail, Cymric, Highland Lynx, Highlander, Manx, Pixiebob and Toybob.

However, photographic evidence in these breeds revealed several short-tailed cats that were not carrying any

known bobtail variant in addition to the cats that were the long-tailed versions of these breeds. The absence

of known variants was especially profound in Highlander breed types and Toybob, in which cats with no known

bobtail variant had a prevalence of 50-60% and 78.6%, respectively (S4 Table).

We also discovered that the c.816+1G>A variant in the *KRT71* gene which is known to result in the hairless

phenotype of the Sphynx cat had a variant frequency of 74.7% in the Sphynx breed (S4 Table). Compound

heterozygotes for the Sphynx c.816+1G>A variant with the Devon Rex associated curly coat variant are also

hairless. A subset of 2,186 samples were genotyped for the Devon Rex curly coat in this study, however, we

also identified 22 hairless Sphynx cats without any copies of the Sphynx variant suggesting that an additional

unknown variant in the same or another gene entirely causes the hairless phenotype in some cats of this

breed. We also confirmed that the Sphynx variant was not present in the Donskoy and Peterbald breeds, which

also represent hairless phenotypes. It was recently suggested that a novel 4 base pair variant of the *LPAR6*

gene identified in a Peterbald cat potentially results in a hairless coat phenotype as a homozygote or as a

compound heterozygote with the c.250_253_delTTTG variant in the *LPAR6* gene which causes the curly coat

of the Cornish Rex (57). We show a variant frequency of 25% for the Cornish Rex coat variant in the Donskoy,

but did not identify any Cornish Rex coat variant carriers among the 17 tested Peterbald cats of this study.

Fig 2. A) The signature "smile" of a cat with Myotonia Congenita; B) The rare coat color phenotype Amber in

a random-bred cat from Finland; C) Polydactyly (variant 1) also associated with extra toes in all four feet.

Genome-wide analysis of genetic diversity demonstrates differences between and within cat breeds

The entire data set of 11,036 samples was genotyped for 7,815 informative SNP markers distributed across

the genome. In the pedigreed cat population, the median heterozygosity was 34.0% and the typical range

(defined as the 10th and 90th percentile) was 27.2%-38.3%, in the non-pedigreed population, the median

heterozygosity was 38.8%; and the typical range was 29.8%-41.3% (Fig 3). Of the 91 breeds tested, the

median heterozygosity was calculated for 56 breeds and breed types were represented by at least 15

individuals in the dataset (S6 Table). The most diverse breeds include three of the newer cat breeds; the short-

legged Munchkin, produced from a sibling mating followed by regular non-pedigree cat outcrosses (2,62); the

Highlander, a crossbreed of two recent experimental hybrid cat breeds the Desert Lynx and Jungle Curl; and

the Lykoi breed founded by unrelated cats expressing hypotrichosis, whose unique sparse and roaned coat

phenotype may be caused by any of six different variants of the *HR* gene from six independent lineages found

in four different states of the United States, Canada and France (63). The heterozygosity levels of European

Shorthair, Norwegian Forest Cat, Siberian and Manx, developed from the local domestic populations with likely

a larger diversity in the founder population, were above average compared with the entire pedigreed cat

population. The lowest median heterozygosity measures in any pedigreed cat population were observed in the

Burmese, Birman, Havana Brown, Korat, Singapura and cat breeds of the Siamese group (such as Balinese,

Siamese and Oriental Shorthair), in line with previous observations (20). A full breakdown of the diversity levels

per breed can be found in Supplemental Table 5.

Fig 3. The median genetic diversity in pedigreed and non-pedigreed cat populations with typical range (the

10th and 90th percentile).

Discussion

In the largest DNA-based feline study cohort to date, a custom genetic panel screening test was utilized to

determine blood type, disease and phenotypic trait heritage, as well as the relative genome-wide genetic

diversity in 11,036 domestic cats representing 91 breeds and breed types.

One specific area of focus was blood type determination, which is particularly important in cats due to its link

with Neonatal Isoerythrolysis, a significant cause of fading kitten syndrome and neonatal death if the blood

types of breeding pairs are not appropriately identified. We identified 17 breeds in which at least one out of ten

cats in the population were blood type B. We also observed the rare blood type AB, previously described with

notable prevalence only in the Ragdoll, also present in the European Shorthair, Scottish Fold breed types, and

the RagaMuffin. We confirm a high concordance between blood type determination based on the proposed

DNA genotyping scheme for purpose-bred cats, and serological blood type based on the 220 results available

for analysis in this study. This provides further justification for the use of DNA-based blood type determination

approaches in felines, provided that the appropriate carefully validated genetic variants are assayed.

Panel screening of disease-associated variants provides clinically relevant information. In addition to blood

type determination, we show that the genomic data available today can assist in disease diagnosis, treatment,

and preventative care. Through the comprehensive investigation of variant allele frequencies in this study

cohort, we re-evaluate and provide updated variant frequency information compared to estimates provided in

conjunction with the original variant discoveries. Our data suggest that disease-associated variant frequencies

are now lower for many conditions (GM2 in Burmese, Hypokalemia in Burmese, HCM-A31P in Maine Coon,

HCM-R820W in Ragdoll, PKD in Persian) compared to the frequencies at the time of their discovery, perhaps

reflecting change over time within the breed, presumably due to genetic testing combined with informed

breeding selections. We note in particular that some variants, such as rdAC-PRA, PKD, HCM-A31P and HCM-

R820W appear more common in additional breeds than in the original breed in which they were discovered,

likely due to lack of awareness and inadvertent selection. For example, the *PKD1* variant, initially discovered

to affect nearly 40% of Persian cats, was found in this study at higher frequency in breeds with Persian

background or non-related breeds than it was in the Persian breed. In fact, none of the Persian cat samples in

this study were identified as carriers of the *PKD1* variant. Additional recent studies indicate that the prevalence

of PKD in Persian cats of Iranian origin continues to be high (64), but also that *PKD1* is common in pedigreed

and non-pedigreed cats in Japan and Turkey (65,66). Taken together, the *PKD1* variant should be seen as a

potential genetic cause of PKD in any breed, such as the previously documented Neva Masquerade (67),

Chartreux (54,68) or the Maine Coon as reported in this study. In this study we mainly focused on genotyping

pedigreed cats and had a relatively small sample size of non-pedigreed cats. We nevertheless discovered 13

likely causal disease-associated variants in non-pedigreed cats.

This study sample provides an extensive investigation of disease-associated variant heritage across nearly

100 cat breeds. Large scale screening studies of isolated subpopulations of a species such as pedigreed cats

hold great value as a secondary independent tool for validating original discoveries in as often there are made

by focusing on a limited number of individuals from a single breed. Investigations that extend beyond the

original discovery breed enable researchers to conclusively to understand the causal relationships between

variants and diseases. For each disease-associated variant discovered in additional breeds in this study, a

total of 13 different variants across tens of breeds, our follow up investigations applied a similar validation

protocol as previously recommended for dogs by combining genotype information with clinical information

collected and evaluated by veterinarians to assess variant manifestation in different breed backgrounds (22).

These clinical phenotype evaluation studies are crucial to ensure genetic counseling information that truly

offers solutions to improve the health of cats, and they are fueled by the cat community and individual breeders'

willingness to provide phenotype information and clinical documentation and participate in veterinary

examinations. Here we confirm a strong relationship between several disease-associated variants (*F12*, *PKD1*,

TRPV4, *CEP290*, *ABCB1* and *CLCN1*) and their clinical manifestations. We further suggest investigating the

role of the two *MYBPC3* variants in contributing to HCM in additional breeds, and raising awareness that PK-

def is a genetically common condition that may result in highly variable clinical signs and perhaps be similarly

underdiagnosed in cats as it is in humans (69). Finally, after evaluation, we report two associated disease

variants that have little value as markers for genetic disease. Both the c.140C>T variant of the *UROS* gene

previously co-segregating with a second variant associated with CEP and the c.1558G>A variant of *ARSB*

associated with mild MPS VI disease as a compound heterozygote with the c.1427T>C variant (severe

disease-associated variant) were found to be the major variants in some breeds without any health impact.

We advise DNA testing laboratories to discontinue offering a test for CEP based solely on the use of the

c.140C>T variant. Moreover, similar to previous investigations (54), we further emphasize that prevention of

MPS VI should focus entirely on managing the c.1427T>C severe variant in the cat population. The MPS VI

Modifier is an asymptomatic variant that is contributing to a mild phenotypic expression of disease in compound

heterozygotes (54–56), suggesting selecting against the c.1558G>A variant is not justified or recommended,

as it would also reduce the genetic variation in the breed.

The appearance of the cat is influenced by various genes which are often monitored by genetic testing to

inform breeding pair selection. In this study, all cats were tested for 26 appearance-associated variants.

Information was provided on the frequency at which the trait variants are encountered across breeds, to explain

the observed phenotypes. The same trait variants influencing coat color/type and morphology are highly

frequent in cats of various breed backgrounds, providing evidence of likely causality. However, we note that

the *KIT* gene variant associated with breed-characteristic gloves (white feet) phenotype in Birman cats (58), is

observed in various individuals of other breeds without gloves phenotype. Moreover, while most of the trait

phenotypes were explained by the known variants, the previously discovered variants associated with

shortened tail, extra digits and hairlessness could only explain the presence of some of these phenotypes.

Our analysis of genetic diversity in cat breed populations shows a wide range of diversity levels within and

between breeds. We found evidence that, as expected, more recently formed breeds with a more significant

number of founding individuals and breeds allowing continued outcrossing tend to have the greatest diversity

levels. Maintaining diversity in closed populations is challenging, and the use of outcrossing may help maintain

and potentially increase diversity levels if widely adopted. The importance of preserving diversity for health

and vigor has been widely documented (70–72). Additionally, the disease-associated variant findings in the

non-pedigreed cat samples set demonstrate the significance of genetic screening for known disease-

associated variants as well.

In conclusion, we demonstrate that several feline disease-associated variants are more widespread across cat

breeds and varieties than previously reported, with both dominant and recessive Mendelian disease-

associated variants observed in additional breeds and often at higher allele frequency than the breeds in which

they were originally discovered. This, in part, demonstrates the effectiveness of proactive genetic testing, which

has reduced disease-associated variant frequencies in notably affected breeds. We have also shown that

some disease-associated variants are very rare and limited to specific breeds. We report the prevalence at

which the three clinically relevant feline blood types occur within breeds and breed types and provide trait

variant frequencies across the feline population. We have combined genotype information with phenotypic

information to investigate and re-evaluate causality in different breed backgrounds, confirming causal

relationships for some variants and weak evidence of penetrance for other variants. In summary, genetic

testing can be used to inform breeding decisions aiming to prevent genetic disease, while a concurrent goal

should be to maintain genetic diversity in a breed's population, helping to sustain the breed. As more cats are

genotyped, we will learn more about the feline variant heritage in the broader domestic cat population, leading

to improved advice to all cat owners. Direct-to-consumer tests help to further raise awareness of various

inherited conditions in cats, provide information that owners can share with their veterinarians, and in time, as

more genotypic and phenotypic data are collected, will enable the genetics of common complex feline disease

to be deciphered, paving the way for personalized precision healthcare with the potential to ultimately improve

welfare for all cats.

Acknowledgements

We extend our warmest thanks to all the cat owners and breeders who enabled the present study by voluntarily

submitting photos and clinical documentation through their interest in advancing feline genetics research; we

issue special thanks to the participants in the Wisdom Panel-The International Cat Association (TICA) State

of the Cat Study for their early support and enthusiasm for this work. We thank Professor Leslie Lyons from

the University of Missouri and various anonymous cat owners for providing validation samples. Excellent

advice and support were provided by TICA, Finnish Cat Association, Anthony Hutcherson, Adriana Kajon, Suvi

Ruotanen, Dr. Casey A. Brookhart-Knox, Dr. Jason Huff and Dr. Susan Puckett; all of whom are deeply

thanked.

Supporting Material

S1 Table. The summary of 11,036 tested pedigreed and non-pedigreed cats.

S2 Table. Tested disease and trait associated variants.

S3 Table. All tested disease and trait genotype data for 11,036 tested cats.

S4 Table. All tested disease and trait variant frequencies for 11,036 tested cats.

S5 Table. Clustered breed information for representing the proportions of the different blood types with > tested

individuals.

S6 Table. Genetic diversity for all breeds with >15 individuals tested.

References

1.   Lyons LA. Cats – telomere to telomere and nose to tail. Trends Genet. 2021;June 28:1–3. Available
from: https://doi.org/10.1016/j.tig.2021.06.001

2.   Buckley RM, Davis BW, Brashear WA, Farias FHG, Kuroki K, Graves T, et al. A new domestic cat genome
assembly based on long sequence reads empowers feline genomic medicine and identifies a novel
gene for dwarfism. PLoS Genet. 2020;16(10):1–28. Available from:
http://dx.doi.org/10.1371/journal.pgen.1008926

3.   Online Mendelian Inheritance in Animals, OMIA. Sydney School of Veterinary Science. Available from:
https://omia.org/

4.   Buckley RM, Lyons LA. Precision/Genomic Medicine for Domestic Cats. Vet Clin North Am - Small Anim
Pract. 2020;50(5):983–90. Available from: https://doi.org/10.1016/j.cvsm.2020.05.005

5.   Narfström K, Holland Deckman K, Menotti-Raymond M. The Domestic Cat as a Large Animal Model
for Characterization of Disease and Therapeutic Intervention in Hereditary Retinal Blindness. J
Ophthalmol. 2011;2011:1–8.

6.   Supsavhad, Wachiraphan; Dirksen, Wessel P.; Martin, Chelsea K.; Rosol TJ. Animal models of head and
neck squamous cell carcinoma. Vet J. 2016;210:7–16.

7.   Packialakshmi B, Stewart IJ, Burmeister DM, Chung KK, Zhou X. Large animal models for translational
research in acute kidney injury. Ren Fail. 2020;42(1):1042–58. Available from:
https://doi.org/10.1080/0886022X.2020.1830108

8.   Freeman, GH; Halton J. Note on exact treatment of contingency, goodness of fit and other problems
of significance. Biometrika. 1951;38:141–9.

9.   Winkler PA, Occelli LM, Petersen-Jones SM. Large Animal Models of Inherited Retinal Degenerations:
A Review. Cells. 2020;9(4).

10.   Ottoni, Claudio; Van Neer, Wim; De Cupere, Bea; Daligault, Julien; Guimaraes, Silvia; Peters, Joris;
Spassov, Nikolai; Prendergast, Mary E.; Boivin, Nicole; Morales-Muniz, Arturo; Balasescu, Adrian;
Becker, Cornelia; Benecke, Norbert; Boroneath, Adrina; B E-M. The paleogenetics of cat dispersal in
the ancient world. Nat Ecol Evol. 2017;1(0139):1–7.

11.   Gandolfi B, Grahn RA, Gustafson NA, Proverbio D, Spada E, Adhikari B, et al. A novel variant in CMAH
is associated with blood type AB in Ragdoll cats. PLoS One. 2016 May 1;11(5).

12.   Bighignoli B, Niini T, Grahn RA, Pedersen NC, Millon L V., Polli M, et al. Cytidine monophospho-N-
acetylneuraminic acid hydroxylase (CMAH) mutations associated with the domestic cat AB blood
group. BMC Genet. 2007 Jun 6;8.

13.   Omi T, Nakazawa S, Udagawa C, Tada N, Ochiai K, Chong YH, et al. Molecular characterization of the
cytidine monophosphate-n-acetylneuraminic acid hydroxylase (CMAH) gene associated with the
feline AB blood group system. PLoS One. 2016 Oct 1;11(10).

14. Kehl, Alexandra; Mueller, E; Giger U. CMAH genotyping survey for blood types A, B and C (AB) in
purpose-bred cats. Anim Genet. 2019;doi:10.1111/age.12778.

15. Kehl A, Heimberger K, Langbein-Detsch I, Boehmer S, Raj K, Mueller E, et al. Molecular
characterization of blood type A, B, and C (AB) in domestic cats and a CMAH genotyping scheme. PLoS
One. 2018 Sep 1;13(9).

16. Gandolfi B, Gruffydd-Jones TJ, Malik R, Cortes A, Jones BR, Helps CR, et al. First WNK4-Hypokalemia
Animal Model Identified by Genome-Wide Association in Burmese Cats. PLoS One. 2012 Dec 28;7(12).

17. Bradbury AM, Morrison NE, Hwang M, Cox NR, Baker HJ, Martin DR. Neurodegenerative lysosomal
storage disease in European Burmese cats with hexosaminidase β-subunit deficiency. Mol Genet
Metab. 2009 May;97(1):53–9.

18. Balmer L, O'leary CA, Menotti-Raymond M, David V, O'brien S, Penglis B, et al. Mapping of diabetes
susceptibility LOCI in a domestic cat breed with an unusually high incidence of diabetes mellitus.
Genes (Basel). 2020;11(11):1–11.

19. Lyons LA, Erdman CA, Grahn RA, Hamilton MJ, Carter MJ, Helps CR, et al. Aristaless-Like Homeobox
protein 1 (ALX1) variant associated with craniofacial structure and frontonasal dysplasia in Burmese
cats. Dev Biol. 2016;409(2):451–8.

20. Gandolfi B, Alhaddad H, Abdi M, Bach LH, Creighton EK, Davis BW, et al. Applications and efficiencies
of the first cat 63K DNA array. Sci Rep. 2018;8(1):1–15.

21. Donner J, Kaukonen M, Anderson H, Möller F, Kyöstilä K, Sankari S, et al. Genetic panel screening of
nearly 100 mutations reveals new insights into the breed distribution of risk variants for canine
hereditary disorders. PLoS One. 2016;11(8):1–18.

22. Donner J, Anderson H, Davison S, Hughes AM, Bouirmane J, Lindqvist J, et al. Frequency and
distribution of 152 genetic disease variants in over 100,000 mixed breed and purebred dogs. PLoS
Genet. 2018;14(4):1–20.

23. Dreger DL, Hooser BN, Hughes AM, Ganesan B, Donner J, Anderson H, et al. True Colors: Commercially-
acquired morphological genotypes reveal hidden allele variation among dog breeds, informing both
trait ancestry and breed potential. PLoS One. 2019;14(10):1–24.

24. Brooks, M; DeWilde L. Feline factor XII deficiency. Compend Contin Educ Pr Vet. 2006;28:148–56.

25. Maruyama H, Brooks MB, Stablein A, Frye A. Factor XII de fi ciency is common in domestic cats and
associated with two high frequency F12 mutations. Gene. 2019;706(December 2018):6–12. Available
from: https://doi.org/10.1016/j.gene.2019.04.053

26. Maruyama H, Hosoe H, Nagamatsu K, Kano R. A novel missense mutation in the factor XII gene in a
litter of cats with factor XII deficiency. 2017;822–6.

27. Bender DE, Kloos MT, Pontius JU, Hinsdale ME, Bellinger DA. Molecular Characterization of Cat Factor
XII Gene and Identification of a Mutation Causing Factor XII Deficiency in a Domestic Shorthair Cat
Colony. Vet Pathol. 2015 Mar 16;52(2):312–20.

28. Grahn RA, Grahn JC, Ct Penedo M, Helps CR, Lyons LA. Erythrocyte Pyruvate Kinase Deficiency
mutation identified in multiple breeds of domestic cats. 2012. Available from: http://www.

29. Kohn, Barbara; Fumi C. Clinical course of pyruvate kinase deficiency in Abyssinian and Somali cats. J
Feline Med Surg. 2008;10(2):145–53.

30. Lyons LA, Biller DS, Erdman CA, Lipinski MJ, Young AE, Roe BA, et al. Feline polycystic kidney disease
mutation identified in PKD1. J Am Soc Nephrol. 2004 Oct;15(10):2548–55.

31.   Meurs KM, Sanchez X, David RM, Bowles NE, Towbin JA, Reiser PJ, et al. A cardiac myosin binding
663   protein C mutation in the Maine Coon cat with familial hypertrophic cardiomyopathy. Hum Mol
664   Genet. 2005 Dec 1;14(23):3587–93.

665   32.   Meurs KM, Norgard MM, Ederer MM, Hendrix KP, Kittleson MD. A substitution mutation in the myosin
666   binding protein C gene in ragdoll hypertrophic cardiomyopathy. Genomics. 2007 Aug;90(2):261–4.

667   33.   Gandolfi B, Alamri S, Darby WG, Adhikari B, Lattimer JC, Malik R, et al. A dominant TRPV4 variant
668   underlies osteochondrodysplasia in Scottish fold cats. Osteoarthr Cartil. 2016 Aug;24(8).

669   34.   Biller DS, DiBartola SP, Eaton KA, Pflueger S, Wellman ML., Radin MJ. Inheritance of polycystic kidney
670   disease in Persian cats. J Hered. 1996;87(1):1–5.

671   35.   Biller DS; Chew DJ. No Polycystic kidney disease in a family of Persian catsTitle. J Am Vet Med Assoc.
672   1990;196(8):1288–90.

673   36.   Barrs VR, Gunew M, Foster SF, Beatty JA, Malik R. Prevalence of autosomal dominant polycystic kidney
674   disease in Persian cats and related-breeds in Sydney and Brisbane. Aust Vet J. 2001;79(4):257–9.

675   37.   Cannon MJ, Barr FJ, Rudorf H, Bradley KJ, Gruffydd-Jones TJ, MacKay A. Prevalence of polycystic kidney
676   disease in Persian cats in the United Kingdom. Vet Rec. 2001;149(14):409–11.

677   38.   Beck C, Lavelle R. Feline polycystic kidney disease in Persian and other cats: a prospective study using
678   ultrasonography. Aust Vet J. 2001;79(3):181–4.

679   39.   Barthez PY, Rivier P, Begon D. Prevalence of polycystic kidney disease in Persian and Persian related
680   cats in France. J Feline Med Surg. 2003;5(6):345–7.

681   40.   Longeri M, Ferrari P, Knafelz P, Mezzelani A, Marabotti A, Milanesi L, et al. Myosin-Binding Protein C
682   DNA Variants in Domestic Cats (A31P, A74T, R820W) and their Association with Hypertrophic
683   Cardiomyopathy. J Vet Intern Med. 2013 Mar;27(2):275–85.

684   41.   Sampedrano CC, Chetboul V, Mary J, Tissier R, Abitbol M, Serres F, Gouni V, Thomas A, Pouchelon J-L.
685   Prospective Echocardiographic and Tissue Doppler Imaging Screening of a Population of Maine Coon
686   Cats Tested for the A31P Mutation in the Myosin-Binding Protein C Gene: A Specific Analysis of the
687   Heterozygous Status. J Vet Intern Med. 2009;23:91–9.

688   42.   Pellegrino A, Daniel AGT, Pereira GG, Itikawa PH, Larsson MHMA. Assessment of regional left
689   ventricular systolic function by strain imaging echocardiography in phenotypically normal and
690   abnormal Maine coon cats tested for the A31p mutation in the MYBPC3 gene. Can J Vet Res.
691   2017;81(2):137–46.

692   43.   Godiksen MTN, granstrøm S, Koch J, Christiansen M. Hypertrophic cardiomyopathy in young Maine
693   Coon cats caused by the p.A31P cMyBP-C mutation - the clinical significance of having the mutation.
694   Acta Vet Scand. 2011;53(1):7. Available from: http://www.actavetscand.com/content/53/1/7

695   44.   Borgeat, Kieran; Casamian-Sorrosal, Domingo; Helps, Chris; Luis Fuentes, Virginia; Connolly DJ.
696   Association of the myosin binding protein C3 mutation (MYBPC3 R820W) with cardiac death in a
697   survey of 236 Ragdoll cats. J Vet Cardiol. 2014;16(2):73–80.

698   45.   Takanosu M, Hattori Y. Osteochondrodysplasia in scottish fold cross-breed cats. J Vet Med Sci.
699   2020;82(12):1769–72.

700   46.   Menotti-Raymond M, David VA, Schäffer AA, Stephens R, Wells D, Kumar-Singh R, et al. Mutation in
701   CEP290 discovered for cat model of human retinal degeneration. J Hered. 2007 May;98(3):211–20.

702   47.   Menotti-Raymond M, David VA, Pflueger S, Roelke ME, Kehler J, O'Brien SJ, et al. Widespread retinal
703   degenerative disease mutation (rdAc) discovered among a large number of popular cat breeds. Vet J.

2010;186(1):32–8. Available from: http://dx.doi.org/10.1016/j.tvjl.2009.08.010

48.   Cogné B, Latypova X, Senaratne LDS, Martin L, Koboldt DC, Kellaris G, et al. Mutations in the Kinesin-
2 Motor KIF3B Cause an Autosomal-Dominant Ciliopathy. Am J Hum Genet. 2020 Jun 4;106(6):893–
904.

49.   Alhaddad H, Gandolfi B, Grahn RA, Rah HC, Peterson CB, Maggs DJ, et al. Genome-wide association
and linkage analyses localize a progressive retinal atrophy locus in Persian cats. Mamm Genome.
2014;25(7–8):354–62.

50.   Mealey KL, Burke NS. Identification of a nonsense mutation in feline ABCB1. J Vet Pharmacol Ther.
2015 Oct 1;38(5):429–33.

51.   Mealey KL, Burke NS, Connors RL. Role of an ABCB11930_1931del TC gene mutation in a temporal
cluster of macrocyclic lactone–induced neurologic toxicosis in cats associated with products labeled
for companion animal use. JAVMA. 2021;259(1):72–6.

52.   Gandolfi B, Daniel RJ, O'brien DP, Guo LT, Youngs MD, Leach SB, et al. A novel mutation in CLCN1
associated with feline myotonia congenita. PLoS One. 2014 Oct 30;9(10).

53.   Clavero S, Bishop DF, Giger U, Haskins ME, Desnick RJ. Feline congenital erythropoietic porphyria: Two
homozygous UROS missense mutations cause the enzyme deficiency and porphyrin accumulation.
Mol Med. 2010 Sep;16(9–10):381–8.

54.   Lyons LA, Grahn RA, Genova F, Beccaglia M, Hopwood JJ, Longeri M. Mucopolysaccharidosis VI in cats
- clarification regarding genetic testing. BMC Vet Res. 2016 Jul 2;12(1).

55.   Yogalingam G, Litjens T, Bielicki J, Crawley AC, Muller V, Anson DS, et al. Feline mucopolysaccharidosis
type VI: Characterization of recombinant N- acetylgalactosamine 4-sulfatase and identification of a
mutation causing the disease. J Biol Chem. 1996;271(44):27259–65. Available from:
http://dx.doi.org/10.1074/jbc.271.44.27259

56.   Crawley AC, Yogalingam G, Muller VJ, Hopwood JJ. Two Feline Mucopolysaccharidosis Type VI
Mutations with Three Phenotypes Two Mutations within a Feline Mucopolysaccharidosis Type VI
Colony Cause Three Different Clinical Phenotypes. Vol. 101, J. Clin. Invest. 1998. Available from:
http://www.jci.org

57.   Lettice LA, Hill AE, Devenney PS, Hill RE. Point mutations in a distant sonic hedgehog cis-regulator
generate a variable regulatory output responsible for preaxial polydactyly. Hum Mol Genet. 2008 Apr
1;17(7).

58.   Montague MJ, Li G, Golfi B, Khan R, Aken BL, Searle SMJ, et al. Comparative analysis of the domestic
cat genome reveals genetic signatures underlying feline biology and domestication. Proc Natl Acad
Sci U S A. 2014 Dec 2;111(48):17230–5.

59.   Alhaddad H, Abdi M, Lyons LA. Patterns of allele frequency differences among domestic cat breeds
assessed by a 63K SNP array. PLoS One. 2021;16(2 February):1–19. Available from:
http://dx.doi.org/10.1371/journal.pone.0247092

60.   Lyons LA, Creighton EK, Alhaddad H, Beale HC, Grahn RA, Rah HC, et al. Whole genome sequencing in
cats, identifies new models for blindness in AIPL1 and somite segmentation in HES7. BMC Genomics.
2016;17(1).

61.   Rodney AR, Buckley RM, Fulton RS, Fronick C, Richmond T, Helps CR, et al. A domestic cat whole exome
sequencing resource for trait discovery. Sci Rep. 2021;11(1):1–13. Available from:
https://doi.org/10.1038/s41598-021-86200-7

60.   Struck AK, Braun M, Detering KA, Dziallas P, Neßler J, Fehr M, et al. A structural UGDH variant

associated with standard Munchkin cats. BMC Genet. 2020;21(1):1–10.

62.   Buckley RM, Gandolfi B, Creighton EK, Pyne CA, Bouhan DM, Leroy ML, et al. Werewolf, there wolf:
749   Variants in hairless associated with hypotrichia and roaning in the lykoi cat breed. Genes (Basel). 2020
750   Jun 1;11(6):1–15.

751   63.   Noori Z, Moosavian HR, Esmaeilzadeh H, Vali Y, Fazli M. Prevalence of polycystic kidney disease in
752   Persian and Persian related-cats referred to Small Animal Hospital, University of Tehran, Iran. Iran J
753   Vet Res. 2019;20(2):151–4.

754   64.   Sato R, Uchida N, Kawana Y, Tozuka M, Kobayashi S, Hanyu N, et al. Epidemiological evaluation of cats
755   associated with feline polycystic kidney disease caused by the feline PKD1 genetic mutation in Japan.
756   J Vet Med Sci. 2019;81(7):1006–11.

757   65.   Bilgen N, Bişkin Türkmen M, Çınar Kul B, Isparta S, Şen Y, Akkurt MY, et al. Prevalence of PKD1 gene
758   mutation in cats in Turkey and pathogenesis of feline polycystic kidney disease. J Vet Diagnostic
759   Investig. 2020;32(4):549–55.

760   66.   Jasik A, Kulesza M. Polycystic kidney disease in a Neva Masquerade cat. J Small Anim Pract.
761   2014;55(7):387.

762   67.   Helps C,Tasker S, Harley R. Correlation of the feline PKD1 genetic mutation with cases of PKD
763   diagnosed by pathological examination. Exp Mol Pathol. 2007;83(2):264–8.

764   68.   Secrest MH, Storm M, Carrington C, Casso D, Gilroy K, Pladson L, et al. Prevalence of pyruvate kinase
765   deficiency: A systematic literature review. Eur J Haematol. 2020;105(2):173–84.

766   69.   Montague MJ, Li G, Golfi B, Khan R, Aken BL, Searle SMJ, et al. Comparative analysis of the domestic
767   cat genome reveals genetic signatures underlying feline biology and domestication. Proc Natl Acad Sci
768   U S A. 2014 Dec 2;111(48):17230–5.

769   70.   Benson JF, Hostetler JA, Onorato DP, Johnson WE, Roelke ME, Brien SJO, et al. HHS Public Access.
770   2020;80(5):958–67.

771   71.   Johnson W, Onorato D, Roelke M, Land E, Cunningham M, Belden R, et al. Genetic restoration of the
772   Florida panther Final environmental assessment. Science (80-). 2010;329(5999):1641–5. Available
773   from:
774   http://science.sciencemag.org/content/329/5999/1641.short%0AJordan_1994_Genetic_restoration
775   _of_the_Florida_panther.pdf

776   72.   Akeberegn D, Getabalew M, Getahun D, Alemneh T, Zewdie D. Importance of hybrid vigor or heterosis
777   for   animal   breeding.   Biochem   Biotechnol   Res.   2019;7(July):1–4.   Available   from:
778   https://www.researchgate.net/publication/334416265