Phase-amplitude coupling and phase synchronization between medial temporal, frontal and posterior brain regions support episodic autobiographical memory recall

Nicolas Roehri[1,*], Lucie Bréchet[2,3,4,*], Martin Seeber[1], Alvaro Pascual-Leone[3,4,5], Christoph M. Michel[1,2]

[1]Functional Brain Mapping Laboratory, Department of Fundamental Neurosciences, Campus Biotech, University of Geneva, 1201 Geneva, Switzerland.

[2]Center for Biomedical Imaging (CIBM), Lausanne and Geneva, 1015 Lausanne, Switzerland

[3]Hinda and Arthur Marcus Institute for Aging Research and Deanna and Sidney Wolk Center for Memory Health, Hebrew SeniorLife, Boston, MA, USA

[4]Department of Neurology, Harvard Medical School, Boston, MA, USA

[5]Guttmann Brain Health Institute, Institut Guttman de Neurorehabilitació, Barcelona, Spain

*The first two authors contributed equally.

Corresponding author:

Christoph M. Michel
Campus Biotech
Neuroscience Department
University of Geneva
9 chemin des Mines
1211 Genève 20, Switzerland
email:christoph.michel@unige.ch

Abstract

Episodic autobiographical memory (EAM) is a complex cognitive function that emerges from the coordination of specific and distant brain regions. Specific brain rhythms, namely theta and gamma oscillations and their synchronization, are thought of as putative mechanisms enabling EAM. Yet, the mechanisms of inter-regional interaction in the EAM network remain unclear in humans at the whole brain level. To investigate this, we analyzed EEG recordings of participants instructed to retrieve autobiographical episodes. EEG recordings were projected in the source space, and time-courses of atlas-based brain regions-of-interest (ROIs) were derived. Directed phase synchrony in high theta (7-10 Hz) and gamma (30-80 Hz) bands and high theta-gamma phase-amplitude coupling were computed between each pair of ROIs. Using network-based statistics, a graph-theory method, we found statistically significant networks for each investigated mechanism. In the gamma band, two sub-networks were found, one between the posterior cingulate cortex (PCC) and the medial temporal lobe (MTL) and another within the medial frontal areas. In the high theta band, we found a PCC to ventromedial prefrontal cortex (vmPFC) network. In phase-amplitude coupling, we found the high theta phase of the left MTL biasing the gamma amplitude of posterior regions and the vmPFC. Other regions of the temporal lobe and the insula were also phase biasing the vmPFC. These findings suggest that EAM, rather than emerging from a single mechanism at a single frequency, involves precise spatio-temporal signatures mapping on distinct memory processes. We propose that the MTL orchestrates activity in vmPFC and PCC via precise phase-amplitude coupling, with vmPFC and PCC interaction via high theta phase synchrony and gamma synchronization contributing to bind information within the PCC-MTL sub-network or valuate the candidate memory within the medial frontal sub-network.

Keywords: episodic autobiographical memory, phase synchronization, phase-amplitude coupling, EEG, network.

Abbreviations: EAM: episodic autobiographical memory, PAC: phase-amplitude coupling, ROI: region of interest, dPLI: directed Phase Lag Index, nMI: normalized Modulation Index, NBS: network-based statistic, dFPN: dorsal Frontoparietal Network, MTL: medial temporal lobe, ACC: anterior cingulate cortex, MCC: median cingulate cortex, PCC: posterior cingulate cortex, SFG: superior frontal gyrus, SFGmedOrb: medial orbital part of the SFG, pSFG: posterior superior frontal gyrus, vmPFC: ventromedial prefrontal cortex, SCG: subcallosal gyrus, Rec: gyrus rectus, STG: superior temporal gyrus, MTG: middle temporal gyrus, ITG: inferior temporal gyrus, PrCu: precuneus, SPG: superior parietal gyrus, IPG: inferior parietal gyrus, SMG: supramarginal gyrus, PHG: parahippocampal gyrus, HIPP: hippocampus, AMY: amygdala, Th: Thalamus.

Introduction

Episodic autobiographical memory (EAM) is a complex cognitive function that involves a specific large-scale network of interacting brain regions to store and later retrieve our lives' experiences. However, the precise mechanisms that enable the specialized brain areas to encode or retrieve EAM remain unclear. Brain oscillations and synchronization within and across large-scale networks facilitate long-distance communication and promote cognitive functions in the healthy human brain (Buzsáki 2006). Particularly, the slower theta (3-10Hz) and faster gamma (>25Hz) oscillations have been related to memory processes but likely play distinct mechanistic roles. Gamma oscillations have been suggested related to memory-related synaptic changes (Nakazono et al. 2018; Sederberg et al. 2007). Gamma phase synchrony in the neocortex is well-known to facilitate neuronal communication and functional integration thanks to the precise temporal coincidence of spiking activity (Fries 2009), and in turn binds distributed memory representations that are needed to retrieve episodic memories (Düzel, Penny, and Burgess 2010; Fell and Axmacher 2011). Theta oscillations are the dominant rhythms recorded in the hippocampus (Buzsaki 2002) and appear to serve as a mechanism of EAM memories constituted of shorter sub-episodes, whose temporal dynamics during recall are supported by the same mechanism as during encoding (Buzsaki 2002; Buzsáki and Moser 2013). Theta oscillations have been associated with both encoding and retrieval of episodic memories (Friese et al. 2013; Osipova et al. 2006; Sederberg et al. 2003).

Generally, theta phase synchrony is supposed to coordinate the coactivation of distant regions whose local neuronal assembly would be synchronized in the gamma range (Fell and Axmacher 2011). This hypothesis can also be seen as cross-frequency phase-amplitude coupling (PAC); the gamma activity would occur at a specific phase of the low-frequency oscillation (Canolty and Knight 2010; Fell and Axmacher 2011; Florin and Baillet 2015; Sirota et al. 2008; Tort et al. 2008). The low-frequency oscillation here acts as a metronome that coordinates neural activity in the same region or a remote

region, making a putative mechanism for long-range communication. Using simultaneous recordings

from both the hippocampus and neocortex in behaving rats and mice, Sirota et al. (Sirota et al. 2008)

showed that the theta oscillations in the hippocampus modulate gamma oscillations in the neocortex

through PAC. Thus, the coupling between hippocampal theta and neocortical gamma oscillations

seems to be a critical mechanism supporting the large-scale memory network's interactions (Axmacher

et al. 2010; Nyhus and Curran 2010).

Very few human studies to date have explored the complex communication within and between

regions involved in EAM. A few recent magnetoencephalography (MEG) studies of personal memories

have started to investigate the mechanisms for the organized communication within and between

broad networks at different spatiotemporal scales (Fuentemilla et al. 2014; Hebscher, Meltzer, and

Gilboa 2019). For example, Fuentemilla et al. (Fuentemilla et al. 2014) found selective phase-

synchronization in theta frequency between the medial temporal lobe (MTL), medial prefrontal cortex,

and precuneus that was higher during EAM than general semantic retrieval. In a follow-up study,

Fuentemilla et al. (2018) sought to identify larger-scale patterns in higher frequencies and found neural

synchrony, specifically in the gamma frequency range underlying episodic autobiographical

recollection. Hebscher et al. (2019) used theta-burst transcranial magnetic stimulation to examine

whether interactions between theta and gamma frequencies facilitate the recall of personal memories

and found that memory retrieval is supported by MTL-cortical communication mediated by theta

phase coupling and theta-gamma PAC. Although such research emphasizes the importance of brain

oscillations in memory processing and integration, these studies relied either on seed-based methods

calculating the phase-synchrony between a selected region and the rest of the brain, thus missing

potential interactions between regions not interacting with the seed, or scalp level connectivity

analysis without any consideration of the source-level network reconstruction. Therefore, there is a

need to analyze the interactions both at the source level and between all brain regions without

selecting only some brain areas based on their coactivation during the EAM processes or a priori

knowledge to obtain a complete picture of the interaction between brain regions.

In the present study, we investigated the dynamic interactions between neuronal networks during

EAM retrieval, particularly in the gamma and high theta frequencies. We focused on the upper range

of the theta band (7-10 Hz; high theta) because, while the frequency of the human hippocampal theta

is rather controversial (Jacobs 2014; de la Prida 2019; Miller et al. 2018), Fuentemilla et al. (2014) found

that when participants were asked to retrieve personal episodes and elaborate them during 30 s during

MEG recording, the peak theta frequency linked to the MTL-Precuneus-mPFC phase synchronization

was above 7 Hz. We assessed the directed phase connectivity and phase-amplitude coupling at play

during EAM recollection, leveraging the high-temporal resolution of EEG and the 64 channels dataset

reported in (Bréchet et al. 2019). We first computed the average power in high theta and gamma bands

in the source space to investigate which parts of the brain were more active during the EAM retrieval

than during a control task. We then applied two types of directed connectivity measures, the directed

phase lag index (dPLI) (Stam and van Straaten 2012) and the normalized modulation index (nMI)

(Özkurt and Schnitzler 2011; Tort et al. 2008) to quantify the coupling at the whole-brain level between

84 regions of interest (ROIs). The dPLI is a measure of phase-synchronization, which is robust to source

leakage and analyzes the direction of the information flow. The nMI allows quantifying the PAC

between the phase of low-frequency time-series and the amplitude of the high-frequency course.

Materials and Methods

Participants

15 healthy participants (30.5 ± 5.5 years, 10 female) were included in the study. The dataset was acquired and described in Bréchet et al. 2019. The local ethics committee's institutional review board approved this study, and all participants provided written informed consent prior to the experiment.

Experimental paradigm

Participants were instructed to either retrieve episodic, self-related memories ("memory condition") or perform arithmetic calculations ("math condition") while undergoing hd-EEG recording. There were 40 trials in total, each trial lasting 30 s. On each trial, a personal photograph (e.g., photo of a participant with a birthday cake) or a calculation (e.g., 447-7=) was presented for 3 s, followed by 27 s period of closed eyes, during which participants either retrieved the EAM or subtracted/added numbers. Participants were asked to retrieve the personal events in as much detail as possible or fully focus on the calculations. At the end of each trial, participants were asked to rate how much they were able to either relive the past self-related events or pay attention to the self-unrelated calculation. The ratings were similar between the memory and math conditions. For more detail on the experimental paradigm, see (Bréchet et al. 2019).

EEG recording and preprocessing

EEG was recorded with a 64-channel BrainAmp EEG system (Brain Products, Munich, Germany) at 5000Hz. The EEG recording was down-sampled to 500Hz and filtered using a zero-phase digital 4th-order Butterworth bandpass filter between 1Hz and 80Hz. Infomax-based Independent Component Analysis (ICA) was applied to remove oculomotor, cardiac, and muscle artifacts based on the channels

with maximal amplitude, the topography, and time course of the component (as recommended in Jung

et al. 2000). Particular attention was devoted to removing long-standing muscle artifacts, which may

overlap and contaminate the genuine gamma activity produced by the brain in the same frequency

band (30-80Hz). In other words, we removed independent components whose topographies were

limited to one or two channels and whose time-courses showed strong activity in the gamma band

(Jung et al. 2000). Bad electrodes were interpolated using a 3-D spherical spline (Perrin et al. 1989) and

the data were re-referenced to the common average reference. Finally, the remaining artifacts were

marked and excluded from further analysis. The pre-processing stages were performed using the

Fieldtrip Toolbox (Oostenveld et al. 2011).

EEG source imaging

The forward model for source localization, based on realistic head geometry and conductivity data

considering skull thickness (Locally Spherical Model with Anatomical Constraints [LSMAC]) (Brunet,

Murray, and Michel 2011), was calculated for a standard set of 64-electrode positions co-registered to

the Montreal Neurological Institute (MNI) atlas. The inverse solution space consisted of about 5000

point sources equally distributed in gray matter volume. We computed the inverse solution matrix

using the linear distributed source localization procedure LAURA (Grave De Peralta Menendez et al.

2004). The lead-field matrix and the inverse matrix were computed using the freely available software

Cartool (Brunet et al. 2011; Michel and Brunet 2019);

https://sites.google.com/site/cartoolcommunity/.

Frequency-band power analyses

We computed the analytic signal of the EEG traces using the Hilbert transform and projected the real

and imaginary parts in the inverse space using the inverse operator to obtain the average power in

high theta and gamma bands at each point source for the memory and math conditions. Before the

Hilbert transform, the signals were bandpass filtered in either high theta (7-10Hz) or gamma (30-80Hz)

bands. The square of the norm of the resulting three-dimensional complex signal was calculated to

obtain a single real power time-course for each point source. A standardization across time was applied

for each point source to eliminate activation biases (Bréchet et al. 2019; Michel and Brunet 2019).

These values were then averaged across time and trials, resulting in an average power value at each

participant's source location.

Whole-brain connectivity analysis

We used 84 regions of interest (ROIs) to build the connectivity matrices. In Cartool, each source was

assigned to one of the ROIs defined by the second version of the AAL atlas (Rolls, Joliot, and Tzourio-

Mazoyer 2015; Tzourio-Mazoyer et al. 2002). S1 Figure shows the position of the centroid of each ROI

in a 3D brain mesh and S1 Table lists the abbreviations and anatomical descriptions of the AAL ROIs.

Notably, the AAL atlas is an anatomical parcellation of the brain based on gyri and sulci rather than

functional specificity. Therefore, we used several ROIs to describe functional brain areas. For example,

the vmPFC was described as in Gilboa and Marlatte 2017 and McCormick et al. 2018 with the three

following ROIs: the medial orbital part of the superior frontal gyrus, gyrus rectus and the subcallosal

gyrus. We projected the filtered data in either high theta or gamma bands into the source space to

obtain the different ROIs' time-courses. We took the first component of the singular value

decomposition computed over all the time-series of every source (independently of their orientation)

belonging to a given ROI. This method was described in Rubega et al. 2019 and used in several other

studies (Carboni et al. 2019, 2020; Damborská et al. 2020). The singular value decomposition has a sign

ambiguity, which can bias directed connectivity measures. To resolve this indeterminacy, we used the

method proposed by Bro, Acar, and Kolda 2008, which allocates the sign based on the majority of

vectors represented by the component. The non-artifacts sections of the trials were concatenated, and

the following connectivity measures were calculated.

We used two types of directed connectivity measures to quantify the coupling between ROIs: the

directed Phase Lag Index [dPLI] (Stam and van Straaten 2012) and the normalized Modulation Index

[nMI] (Canolty et al. 2006; Özkurt and Schnitzler 2011).

Directed Phase Lag Index: The dPLI measures the probability that the difference of the instantaneous

phases of two signals is positive, i.e., the probability that one signal has its instantaneous phase leading

the instantaneous phase of another signal. Formally:

$$224 \qquad dPLI_{xy} := \frac{1}{N} \sum_{t=1}^{N} H\big(\varphi_x(t) - \varphi_y(t)\big)$$

where H is the Heaviside step function, N is the number of samples, φ_x and φ_y are respectively the

instantaneous phase of the time series x and y obtained after applying the Hilbert transform. $dPLI_{xy}$

is bounded between 0 and 1. By definition, x is phase leading compared to y when $dPLI_{xy} > 0.5$ and

phase lagging when $dPLI_{xy} < 0.5$. This measure was applied to the high theta and gamma filtered

time-series separately.

Normalized Modulation Index: The nMI quantifies the phase-amplitude coupling (PAC) between the

phase of low-frequency time-series and the amplitude of the high-frequency course. Mathematically:

$$233 \qquad nMI_{xy} := \frac{1}{\sqrt{N}} \frac{\left| \sum_{t=1}^{N} a_\gamma(t) e^{i\phi_{h\theta}(t)} \right|}{\sqrt{\sum_{t=1}^{N} a_\gamma(t)^2}}$$

a_γ is the amplitude of the analytic signal filtered in gamma, $\phi_{h\theta}$ is the instantaneous phase of the

analytic signal filtered in high theta, and N is the number of samples. Note that the direction of the

connectivity does not carry the usual causality meaning but rather that the phase of region A is coupled

to the amplitude of region B (i.e., A → B means that $\phi_{h\theta}^{A}$ is coupled to a_{γ}^{B}).

Statistical analysis

Activation maps: We applied a two-sided paired t-test to each point source's average power across

participants to investigate which parts of the brain were more active during the memory than the math

condition at the group level. A cluster-based permutation procedure implemented in FieldTrip (Maris

and Oostenveld 2007; Oostenveld et al. 2011) was used to correct multiple comparisons. Clusters are

formed by grouping together neighboring point sources whose t-values were higher than the initial

threshold. The initial threshold for cluster definition was set to p < 0.0005, and the final threshold for

significance of the summed t value within clusters was set to p < 0.05 with 5000 permutations.

Connectivity Networks: Networks are defined by nodes and links. In this study, the nodes are the ROIs,

and the links are the values of the connectivity of each pair of nodes. We used the network-based

statistic (NBS) (Zalesky, Fornito, and Bullmore 2010) to compare the connectivity matrices. First, the

connectivity matrices were subjected to a t-test and then corrected for multiple comparisons using a

connected-component-based permutation procedure. In graph theory, a connected component is a

set of nodes where any two nodes (e.g., A and B) are connected either directly (A-B) or indirectly (i.e.,

via other nodes: A-C-D-B). As for the cluster-based statistics, one can derive a measure for each

component present after the first level threshold, presently the number of links, and test this measure

against values obtained after permutation. Importantly, NBS only works for undirected networks as it

uses a breadth-first search (Hopcroft and Tarjan 1973). We modified this part to work with a directed

network by searching for weakly connected components, i.e., seeking connected components while

ignoring link direction as implemented in Matlab's *conncomp*. We chose weakly over strongly

connected components because strongly connected components have a stricter definition, which may

not be realistic in terms of brain networks. A component is defined as strongly connected if and only

if for every pair of nodes, there is a directed path between any two nodes. For instance, A→B→C is

weakly connected because there is no directed path to go from B or C to A, while A→B→C→A is

strongly connected, as there is a directed path that connects B to A (B→C→A) and C to A (C→A).

As proposed by Zalesky and colleagues (Zalesky et al. 2010), the data can be transformed prior to the

t-test to ensure a meaningful first level threshold. Both the nMI and dPLI are bounded by 0 and 1.

Therefore we applied a logit transform. The logit transform is a monotonically increasing function and

transforms values from $(0,1)$ to $(-\infty, +\infty)$, specifically logit(0.5) = 0. The above inequalities relative to

the dPLI are conserved except that the critical value became 0 instead of 0.5.

We directly contrasted the memory and math conditions through a paired t-test as the nMI matrices

are non-symmetric ($nMI_{xy} \neq nMI_{yx}$). The permutations were obtained by randomly exchanging the

conditions within subjects. However, the dPLI matrices are anti-symmetric ($dPLI_{xy} = -dPLI_{yx}$) and

a direct comparison of the conditions is thus not possible. Therefore, we first applied a one-sample t-

test against 0 to determine the presence of a preferred direction between nodes across participants

and then removed the links present in both memory and math conditions. Here the permutations were

obtained by randomly transposing the matrices, i.e., changing the direction of the links. The initial

threshold for connected-component definition was set to p < 0.005. The final threshold for significance

of the number of connections within connected components was set to p < 0.05 with 5000

permutations. We also defined three different types of nodes: a.) source nodes are nodes with only

outgoing links, b.) sink nodes with only incoming links, and c.) intermediate nodes have incoming and

outgoing links. When describing the significant networks, we will give their number of links and nodes

and, for the sake of readability, only detail the connectivity of their strongest nodes, i.e., nodes with

more than three links. The complete networks are shown in the respective figures and supplementary

tables.

Comparison of the distance between connected regions: Finally, we calculated the distance between

the centroids of pairs of connected ROIs for each connectivity type. We performed a one-way ANOVA

to test whether there exists a difference in these profiles of distance. Post hoc t-tests were used to

explore differences between all the pairs of connectivity types and were controlled for multiple

comparisons by the Dunn-Sidák approach (Šidák 1967). All reported p-values were corrected for

multiple comparisons by one of the above approaches.

Results

Power changes in source space

We found several statistically significant clusters of point sources in gamma and high theta bands when

comparing the power in the memory condition to the control/math condition **(Fig. 1, Table 1)**. Specific

to the gamma frequency, we found 5 significant clusters (Fig. 1a, Table 1a). The first cluster (sum of t:

982.03, p < 0.01) had its maximum t-value located in the right superior temporal gyrus (STG), and the

right temporal pole and supramarginal gyrus (SMG) were its anterior and posterior limits, respectively.

It also included part of the right middle temporal gyrus (MTG), insula, rolandic operculum, and

postcentral gyrus. The second cluster (sum of t: 624.44, p < 0.01) showed a maximum t-value in the

left precuneus (PrCu) and included the PrCu bilaterally and the left superior parietal (SPG) and inferior

parietal gyri (IPG). The third cluster (sum of t: 258.80, p < 0.01) bilaterally included the median cingulate

cortex (MCC) and posterior superior frontal gyrus (pSFG), and its peak was in the left MCC. The peak

of the fourth cluster was found in the left SMG, and this cluster (sum of t: 89.88, p<0.01) was composed

of the left SMG, IPG, and postcentral gyrus. The fifth cluster (sum of t: 32.29, p<0.05) included the right

caudate nucleus and thalamus with a peak in the right caudate nucleus.

We identified 4 significant clusters in the high theta frequency (Fig. 1b, Table 1b). The first cluster (sum

of t: 204.54, p < 0.01) had its maximum t-value located in the left SPG and included the left postcentral

gyrus, IPG, precentral gyrus, and the left PrCu. The second cluster (sum of t: 60.62, p < 0.01) showed a

peak in the right superior frontal gyrus (SFG) and included the right pSFG. The third cluster (sum of t:

56.78, p < 0.01) bilaterally included part of the pSFG and the left SFG with a peak in the left pSFG. The

fourth cluster (sum of t: 18.87, p < 0.05) showed a maximum t-value in the right SPG and partially

included the postcentral gyrus.

Fig. 1 Significant clusters in a. gamma and b. high theta band when contrasting the memory vs. math

conditions. *Only slices containing significant clusters are shown. Numbers identify the different clusters*

and are detailed in Table 1

322 ***Table 1 Anatomical description of the statistically significant clusters obtained when contrasting***

323 ***memory and math condition in the a. gamma band and b. high theta band.***

a. gamma band								
# Cluster	size	p	AAL regions	peak region	peak t	X	Y	Z
1	193	0.0014	R STG R MTG R insula R rolandic operculum R PostCG R Putamen R SMG R STGpole R MTGpole R heschl gyrus R ITG R Pallidum	R STG	6.51	55	-3	-3
2	128	0.0032	L precuneus R precuneus L SPG L IPG L SOG L MCC L cuneus R MCC L MOG	L precuneus	5.68	-3	-55	42
3	54	0.0056	L MCC R MCC L pSFG R pSFG	L MCC	5.29	-3	-10	42
4	19	0.0090	L SMG L IPG L PostCG	L SMG	5.27	-62	-29	36
5	7	0.0130	R Caudate R Thalamus	R Caudate	4.70	16	-3	23
b. high theta band								
# Cluster	size	p	AAL regions	peak region	peak t	X	Y	Z
1	37	0.0030	L SPG L PostCG L IPG L precuneus L PreCG	L SPG	7.10	-36	-62	62
2	13	0.0078	R SFG R pSFG	R SFG	4.89	29	-3	68
3	12	0.0080	L pSFG R pSFG L SFG	L pSFG	5.05	-10	-3	74
4	4	0.0120	R SPG R PostCG	R SPG	5.12	29	-55	68

The lists of AAL regions included in a cluster are sorted in a descending order according to their number

of point sources within the cluster. Sum of t-values correspond to the summation of the t-values within

a cluster. The size corresponds to the number of point sources making up the cluster. The region where

the peak t-value is found is given under the peak region column. X, Y and Z are MNI coordinates in mm

of the peak. Only regions with more than 1% of their size included in the clusters are listed. Annotations:

L: left, R: Right, STG: superior temporal gyrus, MTG: middle temporal gyrus, ITG: inferior temporal

gyrus, PreCG: precentral gyrus, PostCG: postcentral gyrus, SMG: supramarginal gyrus, SPG: superior

parietal gyrus, IPG: inferior parietal gyrus, SOG: superior occipital gyrus, MOG: middle occipital gyrus,

MCC: median cingulate cortex, SFG: superior frontal gyrus, pSFG: posterior superior frontal gyrus.

Directed functional connectivity

As shown in **Figure 2 and S2 Table**, the gamma band directed connectivity analysis revealed a network

($p < 0.01$) composed of 16 links and 16 nodes (including 9 source, 1 intermediate, and 6 sink nodes).

No statistically significant network was found in the math condition, and therefore no link was

removed in the network found in the memory condition as described in the Method section. Most of

the links connected nodes in the frontoparietal midline and left MTL. Particularly, the amygdala (AMY)

was the sink node with the most incoming links (4 links) originating from the hippocampus (HIPP),

parahippocampal gyrus (PHG), thalamus (Th), and left PCC. The right subcallosal gyrus (SCG) was the

second strongest sink node with 3 links connected to the right PCC, pSFG, and left Th. With the same

number of incoming links, the left anterior cingulate cortex (ACC) was connected to the left and right

pSFG and the medial orbital part of the right SFG (SFGmedOrb). The pSFG was the source node with

the most outgoing links, connecting the ACC and right SCG.

Fig. 2 Significant directed network in gamma in the memory condition. *The statistically significant network (16 links and 16 nodes, p<0.01) is represented in 3D along the a.) coronal, b.) sagittal and c.) axial axes. The connectivity matrix is shown in d. Triangles and squares encode nodes with only outgoing or incoming links, i.e., source or sink nodes, respectively. Circles correspond to intermediate nodes with both incoming and outgoing links. The normalized grand average of the connectivity is color-coded. One corresponds to the maximum value. The size of the node encodes its number of (outgoing or incoming) links. Annotations: L: left, R: Right, Th: thalamus, SCG: subcallosal gyrus, AMY: amygdala, HIPP: hippocampus, PCC: posterior cingulate cortex, ITG: inferior temporal gyrus, PHG: parahippocampal gyrus, SFG: superior frontal gyrus, pSFG: posterior SFG, ACC: anterior cingulate cortex, STGpole: polar part of the superior temporal gyrus, SFGmedOrb: medial orbital part of the SFG, ITG: inferior temporal gyrus, MOG: middle occipital gyrus.*

As shown in **Figure 3 and S3 Table**, the high theta directed connectivity analysis revealed the presence

of a medial posterior-to-anterior network (p < 0.01) composed of 21 links and 18 nodes (including 4

source and 14 sink nodes). Only one link was present in both memory and math conditions and was

removed only to keep links specific to the memory condition as described in the Methods section. The

left PCC was the source region with the most outgoing links (9 links). This region drove the right ACC

and bilaterally the gyrus rectus (Rec), SCG, Th, and MCC. The right PCC was driving 5 regions: the right

Rec and bilaterally the Th and SCG. The right inferior temporal gyrus (ITG) led the right Rec,

SFGmedOrb, opercular part of the IFG, and the orbital part of both SFG.   Interestingly, the two

strongest source nodes (left and right PCCs) were mostly connected to the nodes of the vmPFC (defined

by the SFGmedOrb, Rec, and SCG nodes).

 Fig. 3 Significant directed network in high theta in the memory condition. *The statistically significant*

network (21 links and 18 nodes, p<0.01) is represented in 3D along the a.) coronal, b.) sagittal and c.)

axial axes. The connectivity matrix is shown in d. Triangles and squares encode nodes with only

outgoing or incoming links, i.e., source or sink nodes, respectively. The normalized grand average of the

connectivity is color-coded; one corresponds to the maximum value. The size of the node encodes its

number of (outgoing or incoming) links. Annotations: L: left, R: Right, PCC: posterior cingulate cortex,

SCG: subcallosal gyrus, Rec: gyrus rectus, ACC: anterior cingulate cortex, Th: thalamus, MCC: median

cingulate cortex, PrCu: precuneus, SFG: superior frontal gyrus, pSFG: posterior SFG, ITG: inferior

temporal gyrus, SFGorb: orbital part of the SFG, ITG: inferior temporal gyrus, SFGmedOrb: medial

orbital part of the SFG.

Phase-amplitude coupled network

Compared to the gamma and high theta directed networks, the direction of the PAC-directed network

links carries information about the phase to amplitude relationship. For a given pair of connected

nodes, the high theta phase of the node with the outgoing link is coupled to the gamma amplitude of

the node with the incoming link. We identified a statistically significant PAC network ($p < 0.05$) when

comparing the memory to the math conditions **(Fig. 4 and S4 Table)**. This network was composed of

20 links and 20 nodes (including 12 source, 3 intermediate, and 5 sink nodes). The left SFGmedOrb

(part of the vmPFC) was the sink region with the most incoming links (5 links). Its gamma amplitude

was coupled to the high theta phase of the left PHG, HIPP, STG, ITG, and lingual gyrus. The right SPG

was the second sink node with the most incoming links (4 links). It was coupled to the left HIPP and

MCC, and right AMY and superior occipital gyrus. The right Rec (also a node of the vmPFC) had 3 links

coming from the right MTG pole, right insula, and left MTG. The left fusiform gyrus received 3

connections from the right STG, precentral gyrus, and angular gyrus. Interestingly, we found that the

left HIPP had its high theta phase coupled to gamma amplitude of the two sink nodes with the highest

number of links, namely the SFGmedOrb (part of the vmPFC) and right SPG.

Fig. 4 Significant phase-amplitude coupling directed network when contrasting the memory and

math conditions. *The statistically significant network (20 links and 20 nodes, p<0.05) is represented in*

3D along the a.) coronal, b.) sagittal and c.) axial axes. The connectivity matrix is shown in d. Triangles

and squares encode nodes with only outgoing or incoming links, i.e., source or sink nodes, respectively.

Circles correspond to intermediate nodes with both incoming and outgoing links. The normalized grand

average of the connectivity is color-coded. One corresponds to the maximum value. The size of the node

encodes its number of (outgoing or incoming) links. Annotations: L: left, R: Right, INS: insula, MTG:

middle temporal gyrus, ANG: angular gyrus, MTGpole: polar part of the MTG, PHG: parahippocampal

gyrus, STG: superior temporal gyrus, FUSI: fusiform gyrus, Rec: gyrus rectus, SFG: superior frontal gyrus,

SFGorb: orbital part of the SFG, HIPP: hippocampus, SPG: superior parietal gyrus, SFGmed: medial part

of the superior frontal gyrus, MCC: median cingulate cortex, ITG: inferior temporal gyrus, SFGmedOrb:

medial orbital part of the SFG, AMY: amygdala, LING: lingual gyrus, PreCG: precentral cortex, SOG:

superior occipital gyrus.

ROI to ROI distance profile in different connectivity type

We applied one-way ANOVA to test whether the distributions of the distances between two linked

ROIs were different across connectivity types and found a significant result (F = 11.32, p < 0.0001,

Fig. 5). Furthermore, the post-hoc t-test revealed that the distances were higher when the ROIs were

connected by PAC than by high theta (p < 0.05) or gamma phase synchrony (p < 0.001).

Fig. 5 Boxplot of the distances between ROIs connected either by high theta, gamma phase

synchrony, or phase-amplitude coupling. *The one-way ANOVA showed that these distributions were*

not originating from the same distribution (F = 11.32, p < 0.0001). Post-hoc t-tests indicated that the

distances between ROIs connected via phase-amplitude coupling (nMI) were longer than via phase

synchrony. (: p<0.05, ***: p<0.001, the reported p-value were adjusted after the Dunn-Sidák*

procedure correcting for multiple comparisons). dPLI: directed phase lag index, nMI: normalized

Modulation Index, hθ: high theta, γ: gamma.

Discussion

A brief summary of main findings

Our results showed frequency-specific activation of PrCu, SMG, MCC, and STG in the gamma band and

SPG, SFG, and pSFG in the high theta band. We also found a frequency-specific directed phase-

synchrony network that highlighted the importance of frontoparietal midline and MTL interactions.

Specifically, in the gamma band, the left AMY ROI was driven by other MTL regions (HIPP, PHG) and

left PCC, and the pSFG, ACC, and vmPFC were interconnected. In the high theta band, a posterior-to-

anterior network was found with the PCC driving the vmPFC. Finally, the high theta-gamma PAC

network was dense and showed that the MTL phase organized the left vmPFC and right SPG. Overall,

the results reveal distinct spatio-temporal signatures in the high theta and gamma bands, with specific

interactions that we hypothesize reflect distinct cognitive processes of self-referential

autobiographical memory retrieval.

Increased gamma activity in regions linked to self-relevant processing

The brain regions with increased gamma activity were primarily found in the left PrCu, left SMG, right

STG, and MCC. Generally, these gamma activity clusters seem to occur in regions whose role is related

to the first-person perspective and self-orientation.

The PrCu and the SMG have been implicated in self-relevant processing. The PrCu is supposed to shift

the frame of reference from an allocentric to an egocentric one (Hebscher, Levine, and Gilboa 2018),

and stimulation of this region has been shown to alter the autobiographical memory retrieval

(Hebscher, Ibrahim, and Gilboa 2020; Hebscher et al. 2019). In a case report, it was reported that

epileptic seizures involving the left SMG provoked autonomic hallucinations (i.e., seeing the double of

oneself from an internal point of view) in a patient investigated with intracerebral EEG (Fonti et al.

2020). In addition, these two regions (PrCu and SMG) were also found in the fMRI analysis in the

original paper of the analyzed dataset (Bréchet et al. 2019), strengthening the implication of these

regions in EAM.

The MCC is known to be involved in the head and body's reflexive orientation in space (Vogt 2016).

This region could work in concert with the PrCu and SMG areas to create an internal representation

and sense of self-orientation during EAM recall. The right STG is often listed as activated during

recollection of EAM (Boccia, Teghil, and Guariglia 2019; Bréchet et al. 2018) but does not belong to the

core regions, unlike the anterior temporal area (Svoboda, McKinnon, and Levine 2006). Of note, it

could be possible that, because of the lower spatial resolution of the EEG, the activation of the right

temporal junction and the right anterior temporal area merged into a single cluster.

Activation of the dorsal Frontoparietal Network in high theta band

Increased activation of the SFG, pSFG, and SPG was found in a high theta band in the memory condition

compared to the math condition. Because the clusters found in the SFG do not entirely delineate this

gyrus, it can be further interpreted as belonging to the superior frontal sulcus. These regions are known

to be part of the dorsal Frontoparietal Network (dFPN) (Petersen and Posner 2012; Ptak, Schnider, and

Fellrath 2017). Its functions are diverse and include motor planning and imagery, mental rotation,

spatial attention, and working memory (Ptak et al. 2017). At first, it could seem unexpected to find

such a visual attentional network more active in the EAM condition than in the math condition.

However, during the memory condition, the participants were asked to relive past events depicted on

self-related photographs for more than twenty seconds. This requires manipulating and maintaining

the scene internally, which is in line with functions attributed to the dFPN. Although the math condition

is a working memory task, the regions of the dFPN were less involved in that condition. Solving simple

arithmetic operations rely more on the frontal network (Ishii et al. 2014). Our previous study showed

that one specific microstate occurred more often during the math than memory condition (Bréchet et

al. 2019). This microstate's source localization revealed activity in the dorsal ACC, IFG, and IPG,

attributed to the executive control system (Ferris and Peterson 2012).

MTL organizes local gamma synchronization in prefrontal and parietal cortices

Our PAC result showed that the left hippocampus' high theta phase orchestrated the gamma activity

in the medial orbital part of the superior frontal gyrus (belonging to vmPFC) and right superior parietal

gyrus. According to the "reader-initiated" mechanism hypothesis (Buzśaki and Wang 2012; Sirota et al.

2008), the hippocampus through theta oscillations can initiate long-range reciprocal communication

with neocortical gamma oscillations. In that case, the hippocampus would act as a metronome that

coordinates neocortical gamma oscillators. Those gamma oscillations then locally, i.e., within the same

region, synchronize to combine information sent back to the hippocampus. We determine that the left

AMY node received information from the left PCC, Th, PHG, and HIPP via gamma phase synchrony.

Because of the lower spatial resolution of EEG, it might be possible that the AMY node activity globally

represents the activity of the anterior MTL rather than activity solely of the AMY. Similarly, we interpret

the PHG and HIPP nodes as representing the MTL activity and only refer to their name to describe their

connections within the network. This gamma phase synchrony could be the reciprocal connection

initiated via PAC by the left MTL with the right superior parietal gyrus, which sends information back

via the PCC to the anterior MTL.

A possible role of this reciprocal connection is to perform the high-resolution contextual binding to

generate a coherent scene (Yonelinas 2013). Indeed the nodes of this gamma sub-network (AMY, HIPP,

PHG, PCC) form the scene-reconstruction network (Andrews-Hanna et al. 2010; Axelrod, Rees, and Bar

2017; Hassabis and Maguire 2007), which was also found in the original paper's BOLD connectivity

analysis and EEG microstate analysis (Bréchet et al. 2019). Furthermore, as defined in AAL, the PCC ROI

contains part of the retrosplenial cortex (Rolls et al. 2015). This region works in concert with the Th,

which sends head direction information, to switch between allocentric and egocentric perspective

(Vann, Aggleton, and Maguire 2009). The anterior hippocampus, which is more engaged during scene

and events reconstruction than the posterior hippocampus (Zeidman and Maguire 2016), reconstructs

the scene from an allocentric viewpoint. This gamma connectivity from the PCC/retrosplenial cortex,

Th, and PHG to anterior MTL could help orient EAM recollection to a personal view.

The second node entrained by the hippocampal phase was in the vmPFC. This node was part of a more

extensive gamma phase synchrony sub-network constituted by the ACC, pSFG, and other nodes of the

vmPFC. Fuentemilla et al. (2018) showed that large-scale gamma synchrony supports autobiographical

recollection and that this synchrony also occurred in the midline frontal MEG sensor, in line with our

findings. The putative role of the vmPFC is to monitor memory veracity and correctness (Barry and

Maguire 2019; Eichenbaum 2017; Gilboa and Moscovitch 2017; Hebscher and Gilboa 2016; Ritchey

and Cooper 2020). This frontal gamma phase synchronization could be the mechanism by which the

vmPFC combines information from the prefrontal cortex to verify the retrieved memory's validity.

Finally, when analyzing the distance between the pairs of coupled ROIs, we showed that the distance

between ROIs connected via PAC was higher than that via high theta or gamma phase synchronization.

PAC thus seems to be the preferred mechanism for long-range coupling, while high theta or gamma

phase synchronization appears to be the preferred mechanism for local coupling. Our results suggest

that the MTL by entraining with its high theta phase local gamma synchronization in remote and

functionally diverse regions, namely the medial prefrontal and the posterior associative cortex, permits

to reconstruct a coherent and valid self-relevant episode.

 PCC leads and vmPFC follows the network dynamics

 On the one hand, we found nodes within the vmPFC as sink nodes in the high theta phase synchrony

 network and the PAC network. This result suggests that the vmPFC is driven via the high theta

 oscillations in the former network and that gamma activity is coupled to other regions' high theta

 phase in the latter network. On the other hand, we found the PCC to exclusively contain source nodes

 in high theta and gamma phase synchrony network, i.e., the PCC was phase leading other regions.

 Specifically, the network analysis revealed a PCC to vmPFC connection in the high theta band. This

 result is supported by anatomical connections, namely the cingulum bundle, between these two

 regions (Bubb, Metzler-Baddeley, and Aggleton 2018) and a recent paper showing the same link in the

 alpha band within the Default Mode Network (Wang et al. 2019). The PCC is supposed to play a critical

 role in integrating relevant episodic information emerging from posterior cortices (Andrews-Hanna,

 Saxe, and Yarkoni 2014; Bird et al. 2015). This directed connectivity suggests that once episodic

 elements are integrated within the PCC, they are sectioned within high theta cycles and sequentially

 made available via high theta phase synchrony to the vmPFC to either maintain or withhold a memory

 candidate.

 Furthermore, the vmPFC's gamma activity was coupled to regions other than the left MTL (as discussed

 above), including the left STG, ITG, the right MTG pole, and the right insula. These regions are linked

 to either semantic memory processing or emotion processing and are known to contribute to EAM

 recollection by adding semantic and emotional features (Svoboda et al. 2006). This PAC relationship

 suggests that the vmPFC is phase-biased by these regions to constrain the memory validation

 procedure by the semantic features supporting the EAM.

 The fact that the vmPFC was mainly a sink node seems to contradict the results of McCormick et al.

 (2020). While the memory and baseline conditions are relatively similar, our studies differ in several

 ways. First, they focused on the five first seconds of the trial while we analyzed the whole trial.

 Therefore, their finding could describe the construction phase of the EAM while ours encompasses

both the construction and elaboration of EAM. Moreover, the hippocampal and vmPFC ROIs in

McCormick et al. (2020) were selected by contrasting the memory and math/baseline condition and

corresponded to areas with attenuation or decrease of broadband (1-30 Hz) activity. Our ROIs were

based on an anatomical atlas. Finally, in McCormick et al. (2020), dynamic causal modelling was used

to determine which of the hippocampus or the vmPFC was driving the other regions, which, as far as

we know, does not model phase-amplitude coupling or phase synchronization (i.e., the two

mechanisms considered in our investigation). Rather than contradicting each other, our study expands

the findings of McCormick et al. (2020) and conveys different, complementary information about the

dynamic procedure of EAM recall.

Potential limitations and the curse of double-dipping

A recent paper from the OHBM COBIDAS committee (Pernet et al. 2020) emphasized the problem of

double-dipping (Kriegeskorte et al. 2009, 2010). This problem could arise when secondary statistical

analysis (in our case, the connectivity analysis) is applied to ROIs selected with the same statistical

criterion using the same dataset. To avoid this issue, we could have selected ROIs with large power in

both conditions or with larger power in the memory vs. math condition but using an independent

dataset. Rather than selecting ROIs, we avoided double-dipping by applying whole-brain analyses.

Interestingly, the regions with high theta or gamma power increase were almost not present in the

three obtained networks. The PrCu, SMG, and MCC showed a greater gamma power in the memory

than in the math condition, but they were absent in the gamma phase synchrony and PAC network.

Likewise, the SPG and SFG were absent in the high theta phase network while showing a high theta

power increase. The results would have been strikingly different if only the activated ROIs had been

selected for the network analysis. In fact, ROIs with higher activity are not necessarily connected to

each other as their activity could evolve independently of each other, especially in our case with trials

lasting several tens of seconds. Moreover, brain areas that do not exhibit a difference in power

between the conditions could be differently connected in the two conditions. These interactions would

be missed if connectivity analysis is applied only to ROIs with larger activity in the condition of interest.

We thus believed that applying whole-brain connectivity analysis, as proposed here, is more

appropriate to capture networks completely.

PAC integrates the gamma band's amplitude and thus could be biased by changes in amplitude.

However, the regions whose gamma activity is synchronized to the high theta phase of another region

are not present in the gamma's activation map, especially the vmPFC. Therefore, the PAC network

cannot be uniquely attributed to differences in gamma power between the conditions.

Finally, gamma activity in scalp EEG can be highly influenced by noise or muscle artifacts. If such

artifacts were still present after ICA denoising, they would generate zero-lag connectivity between

regions affected by the artefact. As the dPLI is insensitive to zero-lag connectivity and no self-loop, i.e.,

a region is connected to itself, is present in the PAC network, it is unlikely that artifact bias the obtained

networks.

Conclusion

In sum, our findings suggest that the left medial temporal lobe (Hipp, AMG and PHG), the ventromedial

prefrontal cortex (vmPFC), and parietal regions (PCC and SPG) work in tandem during recall of self-

relevant episodic memories. Importantly, this synergy occurs thanks to complex coordination via

frequency-specific and cross-frequency coupling. The medial temporal lobe orchestrates the gamma

activity in the ventromedial prefrontal cortex and parietal regions with its high theta phase. These two

areas synchronize in phase their gamma activity with neighboring regions to, possibly, valuate and

construct the memory. Finally, a high theta synchronization occurs between the posterior cingulate

cortex and medial prefrontal regions, which could show the transfer of sub-episodes of memory from

posterior to frontal regions. Future studies should investigate these relationships utilizing a higher

number of electrodes and a paradigm allowing the separation of the activity of regions related to

semantic memory to refine these episodic autobiographical memory networks further.

Data and Code availability statement

The participants informed consent form did not include permission to allow data to be uploaded to an online repository. Data and materials are available upon direct request and are subject to anonymization to protect the privacy of participants. Requestors must sign a formal data sharing agreement.

Data were analyzed using open source and open access tools. EEG recording were analyzed using Fieldtrip (Oostenveld et al. 2011), available at: https://github.com/fieldtrip/fieldtrip. The lead-field matrix and the inverse matrix were computed using the freely available software Cartool (Brunet et al. 2011; Michel and Brunet 2019); available at https://sites.google.com/site/cartoolcommunity/. The Network Based Statistic toolbox(Zalesky et al. 2010) is available at https://www.nitrc.org/projects/nbs.

Acknowledgements

Funding: This work was supported by the Swiss National Science Foundation to LB (grant No. 187949) and CM (grant No. 320030_184677), and by the National Center of Competence in Research (NCCR) 'SYNAPSY - The Synaptic Bases of Mental Diseases' financed by the Swiss National Science Foundation (SNF, grant n° 51AU40_125759).

References

Andrews-Hanna, Jessica R., Jay S. Reidler, Jorge Sepulcre, Renee Poulin, and Randy L. Buckner. 2010. "Functional-Anatomic Fractionation of the Brain's Default Network." *Neuron* 65(4):550–62.

Andrews-Hanna, Jessica R., Rebecca Saxe, and Tal Yarkoni. 2014. "Contributions of Episodic Retrieval and Mentalizing to Autobiographical Thought: Evidence from Functional Neuroimaging, Resting-State Connectivity, and FMRI Meta-Analyses." *NeuroImage* 91:324–35.

Axelrod, Vadim, Geraint Rees, and Moshe Bar. 2017. "The Default Network and the Combination of

Cognitive Processes That Mediate Self-Generated Thought." *Nature Human Behaviour*

1(12):896–910.

Axmacher, Nikolai, Melanie M. Henseler, Ole Jensen, Ilona Weinreich, Christian E. Elger, and Juergen

Fell. 2010. "Cross-Frequency Coupling Supports Multi-Item Working Memory in the Human

Hippocampus." *Proceedings of the National Academy of Sciences of the United States of*

America 107(7):3228–33.

Barry, Daniel N. and Eleanor A. Maguire. 2019. "Remote Memory and the Hippocampus: A

Constructive Critique." *Trends in Cognitive Sciences* 23(2):128–42.

Bird, Chris M., James L. Keidel, Leslie P. Ing, Aidan J. Horner, and Neil Burgess. 2015. "Consolidation

of Complex Events via Reinstatement in Posterior Cingulate Cortex." *Journal of Neuroscience*

35(43):14426–34.

Boccia, Maddalena, Alice Teghil, and Cecilia Guariglia. 2019. "Looking into Recent and Remote Past:

Meta-Analytic Evidence for Cortical Re-Organization of Episodic Autobiographical Memories."

Neuroscience and Biobehavioral Reviews 107(March):84–95.

Bréchet, Lucie, Denis Brunet, Gwénaël Birot, Rolf Gruetter, Christoph M. Michel, and João Jorge.

2019. "Capturing the Spatiotemporal Dynamics of Self-Generated, Task-Initiated Thoughts with

EEG and FMRI." *NeuroImage* 194(February):82–92.

Bréchet, Lucie, Petr Grivaz, Baptiste Gauthier, and Olaf Blanke. 2018. "Common Recruitment of

Angular Gyrus in Episodic Autobiographical Memory and Bodily Self-Consciousness." *Frontiers in*

Behavioral Neuroscience 12(November):1–10.

Bro, R., E. Acar, and Tamara G. Kolda. 2008. "Resolving the Sign Ambiguity in the Singular Value

Decomposition." *Journal of Chemometrics* 22(2):135–40.

Brunet, Denis, Micah M. Murray, and Christoph M. Michel. 2011. "Spatiotemporal Analysis of

Multichannel EEG: CARTOOL." *Computational Intelligence and Neuroscience* 2011.

Bubb, Emma J., Claudia Metzler-Baddeley, and John P. Aggleton. 2018. "The Cingulum Bundle:

Anatomy, Function, and Dysfunction." *Neuroscience and Biobehavioral Reviews*

92(January):104–27.

Buzsaki, Gyorgy. 2002. "Theta Oscillations in the Hippocampus." *Neuron* 33:325–40.

Buzsáki, György. 2006. *Rhythms of the Brain*.

Buzsáki, György and Edvard I. Moser. 2013. "Memory, Navigation and Theta Rhythm in the

Hippocampal-Entorhinal System." *Nature Neuroscience* 16(2):130–38.

Buzśaki, Gÿorgy and Xiao Jing Wang. 2012. "Mechanisms of Gamma Oscillations." *Annual Review of*

Neuroscience 35:203–25.

Canolty, R. T., N. M. Barbaro, E. Edwards, H. E. Kirsch, S. S. Dalal, S. S. Nagarajan, R. T. Knight, M. S.

Berger, and M. Soltani. 2006. "High Gamma Power Is Phase-Locked to Theta Oscillations in

Human Neocortex." *Science* 313(5793):1626–28.

Canolty, Ryan T. and Robert T. Knight. 2010. "The Functional Role of Cross-Frequency Coupling."

Trends in Cognitive Sciences 14(11):506–15.

Carboni, M., M. Rubega, G. R. Iannotti, P. De Stefano, G. Toscano, S. Tourbier, F. Pittau, P. Hagmann,

S. Momjian, K. Schaller, M. Seeck, C. M. Michel, P. van Mierlo, and S. Vulliemoz. 2019. "The

Network Integration of Epileptic Activity in Relation to Surgical Outcome." *Clinical*

Neurophysiology 130(12):2193–2202.

Carboni, Margherita, Pia De Stefano, Bernd J. Vorderwülbecke, Sebastien Tourbier, Emeline Mullier,

Maria Rubega, Shahan Momjian, Karl Schaller, Patric Hagmann, Margitta Seeck, Christoph M.

Michel, Pieter van Mierlo, and Serge Vulliemoz. 2020. "Abnormal Directed Connectivity of

Resting State Networks in Focal Epilepsy." *NeuroImage: Clinical* 27(March):102336.

Damborská, Alena, Eliška Honzírková, Richard Barteček, Jana Hořínková, Sylvie Fedorová, Šimon

Ondruš, Christoph M. Michel, and Maria Rubega. 2020. "Altered Directed Functional

Connectivity of the Right Amygdala in Depression: High-Density EEG Study." *Scientific Reports*

10(1):1–14.

Düzel, Emrah, Will D. Penny, and Neil Burgess. 2010. "Brain Oscillations and Memory." *Current*

Opinion in Neurobiology 20(2):143–49.

Eichenbaum, Howard. 2017. "Prefrontal-Hippocampal Interactions in Episodic Memory." *Nature*

Reviews Neuroscience 18(9):547–58.

Fell, Juergen and Nikolai Axmacher. 2011. "The Role of Phase Synchronization in Memory Processes."

Nature Reviews Neuroscience 12(2):105–18.

Ferris, Daniel P. and Steven M. Peterson. 2012. "Combined Head Phantom and Neural Mass Model

Validation of Effective Connectivity Measures." *Journal of Acquired Immune Deficiency*

Syndromes 59(2):593–600.

Florin, Esther and Sylvain Baillet. 2015. "The Brain's Resting-State Activity Is Shaped by Synchronized

Cross-Frequency Coupling of Neural Oscillations." *NeuroImage* 111:26–35.

Fonti, Davide, Stanislas Lagarde, Julia Scholly, Anne Lépine, Didier Scavarada, Monica Puligheddu,

Aileen McGonigal, and Fabrice Bartolomei. 2020. "Anatomical Electroclinical Correlations during

an SEEG-Recorded Seizure with Autoscopic Hallucination." *Epileptic Disorders : International*

Epilepsy Journal with Videotape 22(6):817–22.

Fries, Pascal. 2009. "Neuronal Gamma-Band Synchronization as a Fundamental Process in Cortical

Computation." *Annual Review of Neuroscience* 32:209–24.

Friese, Uwe, Moritz Köster, Uwe Hassler, Ulla Martens, Nelson Trujillo-Barreto, and Thomas Gruber.

2013. "Successful Memory Encoding Is Associated with Increased Cross-Frequency Coupling

between Frontal Theta and Posterior Gamma Oscillations in Human Scalp-Recorded EEG."

NeuroImage 66:642–47.

Fuentemilla, L., G. R. Barnes, E. Düzel, and B. Levine. 2014. "Theta Oscillations Orchestrate Medial

Temporal Lobe and Neocortex in Remembering Autobiographical Memories." *NeuroImage*

85:730–37.

Fuentemilla, Lluís, Daniela J. Palombo, and Brian Levine. 2018. "Gamma Phase-Synchrony in

Autobiographical Memory: Evidence from Magnetoencephalography and Severely Deficient

Autobiographical Memory." *Neuropsychologia* 110(August 2017):7–13.

Gilboa, Asaf and Hannah Marlatte. 2017. "Neurobiology of Schemas and Schema-Mediated

Memory." *Trends in Cognitive Sciences* 21(8):618–31.

Gilboa, Asaf and Morris Moscovitch. 2017. "Ventromedial Prefrontal Cortex Generates Pre-Stimulus

Theta Coherence Desynchronization: A Schema Instantiation Hypothesis." *Cortex* 87:16–30.

Grave De Peralta Menendez, Rolando, Micah M. Murray, Christoph M. Michel, Roberto Martuzzi, and

Sara L. Gonzalez Andino. 2004. "Electrical Neuroimaging Based on Biophysical Constraints."

NeuroImage 21(2):527–39.

Hassabis, Demis and Eleanor A. Maguire. 2007. "Deconstructing Episodic Memory with

Construction." *Trends in Cognitive Sciences* 11(7):299–306.

Hebscher, Melissa and Asaf Gilboa. 2016. "A Boost of Confidence: The Role of the Ventromedial

Prefrontal Cortex in Memory, Decision-Making, and Schemas." *Neuropsychologia* 90:46–58.

Hebscher, Melissa, Christine Ibrahim, and Asaf Gilboa. 2020. "Precuneus Stimulation Alters the

Neural Dynamics of Autobiographical Memory Retrieval." *NeuroImage* 210(January):116575.

Hebscher, Melissa, Brian Levine, and Asaf Gilboa. 2018. "The Precuneus and Hippocampus Contribute

to Individual Differences in the Unfolding of Spatial Representations during Episodic

Autobiographical Memory." *Neuropsychologia* 110(March 2017):123–33.

Hebscher, Melissa, Jed A. Meltzer, and Asaf Gilboa. 2019. "A Causal Role for the Precuneus in

Network-Wide Theta and Gamma Oscillatory Activity during Complex Memory Retrieval." *ELife*

8:1–20.

Hopcroft, J. E. and Robert Tarjan. 1973. "Efficient Algorithms for Graph Manipulations."

Communication of the ACM 16(6):372–78.

Ishii, Ryouhei, Leonides Canuet, Tsutomu Ishihara, Yasunori Aoki, Shunichiro Ikeda, Masahiro Hata,

Themistoklis Katsimichas, Atsuko Gunji, Hidetoshi Takahashi, Takayuki Nakahachi, Masao Iwase,

and Masatoshi Takeda. 2014. "Frontal Midline Theta Rhythm and Gamma Power Changes

during Focused Attention on Mental Calculation: An MEG Beamformer Analysis." *Frontiers in*

Human Neuroscience 8(JUNE):1–10.

Jacobs, Joshua. 2014. "Hippocampal Theta Oscillations Are Slower in Humans than in Rodents:

Implications for Models of Spatial Navigation and Memory." *Philosophical Transactions of the*

Royal Society B: Biological Sciences 369(1635).

Jung, Tzyy Ping, Scott Makeig, Colin Humphries, Te Won Lee, Martin J. Mckeown, Vicente Iragui, and

Terrence J. Sejnowski. 2000. "Removing Electroencephalographic Artifacts by Blind Source

Separation." *Psychophysiology* 37(2):163–78.

Kriegeskorte, Nikolaus, Martin A. Lindquist, Thomas E. Nichols, Russell A. Poldrack, and Edward Vul.

2010. "Everything You Never Wanted to Know about Circular Analysis, but Were Afraid to Ask."

Journal of Cerebral Blood Flow and Metabolism 30(9):1551–57.

Kriegeskorte, Nikolaus, W. Kyle Simmons, Patrick S. Bellgowan, and Chris I. Baker. 2009. "Circular

Analysis in Systems Neuroscience: The Dangers of Double Dipping." *Nature Neuroscience*

12(5):535–40.

de la Prida, Liset Menendez. 2019. "The Theta Stone: 3 to 10 Hz Oscillations to Decipher the Human

Brain Code." *Epilepsy Currents* 19(2):126–28.

Maris, Eric and Robert Oostenveld. 2007. "Nonparametric Statistical Testing of EEG- and MEG-Data."

Journal of Neuroscience Methods 164(1):177–90.

McCormick, Cornelia, Daniel N. Barry, Amirhossein Jafarian, Gareth R. Barnes, and Eleanor A.

Maguire. 2020. "VmPFC Drives Hippocampal Processing during Autobiographical Memory Recall

Regardless of Remoteness." *Cerebral Cortex* 1–16.

McCormick, Cornelia, Elisa Ciaramelli, Flavia De Luca, and Eleanor A. Maguire. 2018. "Comparing and

Contrasting the Cognitive Effects of Hippocampal and Ventromedial Prefrontal Cortex Damage:

A Review of Human Lesion Studies." *Neuroscience* 374:295–318.

Michel, Christoph M. and Denis Brunet. 2019. "EEG Source Imaging: A Practical Review of the

Analysis Steps." *Frontiers in Neurology* 10(APR).

Miller, Jonathan, Andrew J. Watrous, Melina Tsitsiklis, Sang Ah Lee, Sameer A. Sheth, Catherine A.

Schevon, Elliot H. Smith, Michael R. Sperling, Ashwini Sharan, Ali Akbar Asadi-Pooya, Gregory A.

Worrell, Stephen Meisenhelter, Cory S. Inman, Kathryn A. Davis, Bradley Lega, Paul A. Wanda,

Sandhitsu R. Das, Joel M. Stein, Richard Gorniak, and Joshua Jacobs. 2018. "Lateralized

Hippocampal Oscillations Underlie Distinct Aspects of Human Spatial Memory and Navigation."

Nature Communications 9(1).

Nakazono, Tomoaki, Heechul Jun, Mathew Blurton-Jones, Kim N. Green, and Kei M. Igarashi. 2018.

"Gamma Oscillations in the Entorhinal-Hippocampal Circuit Underlying Memory and Dementia."

Neuroscience Research 129:40–46.

Nyhus, Erika and Tim Curran. 2010. "Functional Role of Gamma and Theta Oscillations in Episodic

Memory." *Neuroscience & Biobehavioral Reviews* 34(7):1023–35.

Oostenveld, Robert, Pascal Fries, Eric Maris, and Jan Mathijs Schoffelen. 2011. "FieldTrip: Open

Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data."

Computational Intelligence and Neuroscience 2011.

Osipova, Daria, Atsuko Takashima, Robert Oostenveld, Guillén Fernández, Eric Maris, and Ole Jensen.

2006. "Theta and Gamma Oscillations Predict Encoding and Retrieval of Declarative Memory."

Journal of Neuroscience 26(28):7523–31.

Özkurt, Tolga Esat and Alfons Schnitzler. 2011. "A Critical Note on the Definition of Phase-Amplitude

Cross-Frequency Coupling." *Journal of Neuroscience Methods* 201(2):438–43.

Pernet, Cyril, Marta I. Garrido, Alexandre Gramfort, Natasha Maurits, Christoph M. Michel, Elizabeth

Pang, Riitta Salmelin, Jan Mathijs Schoffelen, Pedro A. Valdes-Sosa, and Aina Puce. 2020. "Issues

and Recommendations from the OHBM COBIDAS MEEG Committee for Reproducible EEG and

MEG Research." *Nature Neuroscience*.

Perrin, F., J. Pernier, O. Bertrand, and J. F. Echallier. 1989. "Spherical Splines for Scalp Potential and

Current Density Mapping." *Electroencephalography and Clinical Neurophysiology* 72(2):184–87.

Petersen, Steven E. and Michael I. Posner. 2012. "The Attention System of the Human Brain: 20 Years

After." *Annual Review of Neuroscience* 35:73–89.

Ptak, Radek, Armin Schnider, and Julia Fellrath. 2017. "The Dorsal Frontoparietal Network: A Core

System for Emulated Action." *Trends in Cognitive Sciences* 21(8):589–99.

Ritchey, Maureen and Rose A. Cooper. 2020. "Deconstructing the Posterior Medial Episodic

Network." *Trends in Cognitive Sciences* 24(6):451–65.

Rolls, Edmund T., Marc Joliot, and Nathalie Tzourio-Mazoyer. 2015. "Implementation of a New

Parcellation of the Orbitofrontal Cortex in the Automated Anatomical Labeling Atlas."

NeuroImage 122:1–5.

Rubega, M., M. Carboni, M. Seeber, D. Pascucci, S. Tourbier, G. Toscano, P. Van Mierlo, P. Hagmann,

G. Plomp, S. Vulliemoz, and C. M. Michel. 2019. "Estimating EEG Source Dipole Orientation

Based on Singular-Value Decomposition for Connectivity Analysis." *Brain Topography*

32(4):704–19.

Sederberg, Per B., Michael J. Kahana, Marc W. Howard, Elizabeth J. Donner, and Joseph R. Madsen.

2003. "Theta and Gamma Oscillations during Encoding Predict Subsequent Recall." *Journal of*

Neuroscience 23(34):10809–14.

Sederberg, Per B., Andreas Schulze-Bonhage, Joseph R. Madsen, Edward B. Bromfield, David C.

McCarthy, Armin Brandt, Michele S. Tully, and Michael J. Kahana. 2007. "Hippocampal and

Neocortical Gamma Oscillations Predict Memory Formation in Humans." *Cerebral Cortex*

17(5):1190–96.

Šidák, Zbyněk. 1967. "Rectangular Confidence Regions for the Means of Multivariate Normal

Distributions." *Journal of the American Statistical Association* 62(318):626–33.

Sirota, Anton, Sean Montgomery, Shigeyoshi Fujisawa, Yoshikazu Isomura, Michael Zugaro, and

György Buzsáki. 2008. "Entrainment of Neocortical Neurons and Gamma Oscillations by the

Hippocampal Theta Rhythm." *Neuron* 60(4):683–97.

Stam, C. J. and E. C. W. van Straaten. 2012. "Go with the Flow: Use of a Directed Phase Lag Index

(DPLI) to Characterize Patterns of Phase Relations in a Large-Scale Model of Brain Dynamics."

NeuroImage 62(3):1415–28.

Svoboda, Eva, Margaret C. McKinnon, and Brian Levine. 2006. "The Functional Neuroanatomy of

Autobiographical Memory: A Meta-Analysis." *Neuropsychologia* 44(12):2189–2208.

Tort, Adriano B. L., Mark A. Kramer, Catherine Thorn, Daniel J. Gibson, Yasuo Kubota, Ann M.

Graybiel, and Nancy J. Kopell. 2008. "Dynamic Cross-Frequency Couplings of Local Field

Potential Oscillations in Rat Striatum and Hippocampus during Performance of a T-Maze Task."

Proceedings of the National Academy of Sciences of the United States of America

105(51):20517–22.

Tzourio-Mazoyer, N., B. Landeau, D. Papathanassiou, F. Crivello, O. Etard, N. Delcroix, B. Mazoyer,

and M. Joliot. 2002. "Automated Anatomical Labeling of Activations in SPM Using a

Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain." *NeuroImage*

15(1):273–89.

Vann, Seralynne D., John P. Aggleton, and Eleanor A. Maguire. 2009. "What Does the Retrosplenial

Cortex Do?" *Nature Reviews Neuroscience* 10(11):792–802.

Vogt, Brent A. 2016. "Midcingulate Cortex: Structure, Connections, Homologies, Functions and

Diseases." *Journal of Chemical Neuroanatomy* 74:28–46.

Wang, Regina W. Y., Wei Li Chang, Shang Wen Chuang, and I. Ning Liu. 2019. "Posterior Cingulate

Cortex Can Be a Regulatory Modulator of the Default Mode Network in Task-Negative State."

Scientific Reports 9(1):1–12.

Yonelinas, Andrew P. 2013. "The Hippocampus Supports High-Resolution Binding in the Service of

Perception, Working Memory and Long-Term Memory." *Behavioural Brain Research* 254:34–44.

Zalesky, Andrew, Alex Fornito, and Edward T. Bullmore. 2010. "Network-Based Statistic: Identifying

Differences in Brain Networks." *NeuroImage* 53(4):1197–1207.

Zeidman, Peter and Eleanor A. Maguire. 2016. "Anterior Hippocampus: The Anatomy of Perception,

Imagination and Episodic Memory." *Nature Reviews Neuroscience* 17(3):173–82.

Right	Left	Abbreviations		Right	Left	Abbreviations
1	43	AMY		22	64	IOG
2	44	ANG		23	65	MOG
3	45	CAL		24	66	SOG
4	46	ACC		25	67	SCG
5	47	MCC		26	68	PHG
6	48	PCC		27	69	PCL
7	49	CUN		28	70	IPG
8	50	IFGoper		29	71	SPG
9	51	IFGorb		30	72	PostCG
10	52	IFGtri		31	73	PreCG
11	53	SFGmedOrb		32	74	PrCu
12	54	MFGorb		33	75	Rec
13	55	MFG		34	76	ROL
14	56	SFGmed		35	77	pSFG
15	57	SFGorb		36	78	SMG
16	58	SFG		37	79	ITG
17	59	FUSI		38	80	MTG
18	60	HES		39	81	MTGpole
19	61	HIPP		40	82	STGpole
20	62	INS		41	83	STG
21	63	LING		42	84	Th

S1 Figure. Positions of the 84 nodes in 3D and their abbreviation.
The nodes are positioned at the centroid of the ROIs of the AAL atlas and
represented in 3D along the a.) coronal, b.) sagittal and c.) axial axes. The
tables give the abbreviations relative to the different node numbers.

S1 Table. Anatomical description and abbreviations of the 84 regions used in this study and positions of their centroid in MNI space in mm.

Numbers	Abbreviations	Regions	X	Y	Z	Numbers	Abbreviations	Regions	X	Y	Z
1	R AMY	Right amygdala	26	0	-22	43	L AMY	Left amygdala	-27	-2	-22
2	R ANG	Right angular gyrus	44	-62	38	44	L ANG	Left angular gyrus	-46	-64	35
3	R CAL	Right calcarine fissure and surrounding cortex	13	-76	8	45	L CAL	Left calcarine fissure and surrounding cortex	-8	-82	5
4	R ACC	Right anterior cingulate cortex	6	35	18	46	L ACC	Left anterior cingulate cortex	-6	35	13
5	R MCC	Right median cingulate cortex	7	-12	40	47	L MCC	Left median cingulate cortex	-8	-20	42
6	R PCC	Right posterior cingulate cortex	5	-44	22	48	L PCC	Left posterior cingulate cortex	-5	-46	25
7	R CUN	Right cuneus	11	-80	28	49	L CUN	Left cuneus	-8	-82	25
8	R IFGoper	Right inferior frontal gyrus, opercular part	48	13	18	50	L IFGoper	Left inferior frontal gyrus, opercular part	-50	11	18
9	R IFGorb	Right Inferior frontal gyrus, orbital part	37	30	-16	51	L IFGorb	Left Inferior frontal gyrus, orbital part	-37	29	-15
10	R IFGtri	Right inferior frontal gyrus, triangular part	47	28	12	52	L IFGtri	Left inferior frontal gyrus, triangular part	-47	29	10
11	R SFGmedOrb	Right superior frontal gyrus, medial orbital	7	53	-8	53	L SFGmedOrb	Left superior frontal gyrus, medial orbital	-7	55	-7
12	R MFGorb	Right middle frontal gyrus, orbital part	32	51	-12	54	L MFGorb	Left middle frontal gyrus, orbital part	-32	49	-12
13	R MFG	Right middle frontal gyrus	36	31	33	55	L MFG	Left middle frontal gyrus	-35	31	34
14	R SFGmed	Right superior frontal gyrus, medial	7	49	29	56	L SFGmed	Left superior frontal gyrus, medial	-7	47	30
15	R SFGorb	Right superior frontal gyrus, orbital part	17	47	-17	57	L SFGorb	Left superior frontal gyrus, orbital part	-18	47	-15
16	R SFG	Right superior frontal gyrus, dorsolateral	22	31	41	58	L SFG	Left superior frontal gyrus, dorsolateral	-21	32	41
17	R FUSI	Right fusiform gyrus	33	-42	-22	59	L FUSI	Left fusiform gyrus	-33	-46	-21
18	R HES	Right Heschl's gyrus	42	-23	12	60	L HES	Left Heschl's gyrus	-47	-19	10
19	R HIPP	Right hippocampus	28	-22	-12	61	L HIPP	Left hippocampus	-27	-22	-12
20	R INS	Right insula	38	2	1	62	L INS	Left insula	-37	5	3
21	R LING	Right lingual gyrus	15	-68	-5	63	L LING	Left lingual gyrus	-16	-68	-5
22	R IOG	Right inferior occipital gyrus	38	-82	-10	64	L IOG	Left inferior occipital gyrus	-37	-81	-11
23	R MOG	Right middle occipital gyrus	35	-81	20	65	L MOG	Left middle occipital gyrus	-37	-82	15
24	R SOG	Right superior occipital gyrus	23	-84	31	66	L SOG	Left superior occipital gyrus	-19	-85	29
25	R SCG	Right subcallosal gyrus	8	14	-11	67	L SCG	Left subcallosal gyrus	-10	13	-13
26	R PHG	Right parahipocampal gyrus	25	-17	-23	68	L PHG	Left parahipocampal gyrus	-24	-22	-22
27	R PCL	Right paracentral lobule	7	-35	66	69	L PCL	Left paracentral lobule	-8	-28	67
28	R IPG	Right inferior parietal gyrus, excluding supramarginal and angular gyri	45	-47	50	70	L IPG	Left inferior parietal gyrus, excluding supramarginal and angular gyri	-45	-48	45
29	R SPG	Right superior parietal gyrus	24	-62	59	71	L SPG	Left superior parietal gyrus	-26	-63	56
30	R PostCG	Right postcentral gyrus	42	-26	48	72	L PostCG	Left postcentral gyrus	-47	-23	45
31	R PreCG	Right precentral gyrus	41	-10	49	73	L PreCG	Left precentral gyrus	-41	-7	49
32	R PrCu	Right precuneus	9	-59	41	74	L PrCu	Left precuneus	-9	-58	46
33	R Rec	Right gyrus rectus	6	34	-19	75	L Rec	Left gyrus rectus	-7	36	-19
34	R ROL	Right rolandic operculum	50	-10	14	76	L ROL	Left rolandic operculum	-48	-12	13
35	R pSFG	Right posterior superior frontal gyrus	7	0	60	77	L pSFG	Left posterior superior frontal gyrus	-7	3	58
36	R SMG	Right supramarginal gyrus	56	-35	35	78	L SMG	Left supramarginal gyrus	-59	-33	29
37	R ITG	Right inferior temporal gyrus	52	-31	-25	79	L ITG	Left inferior temporal gyrus	-50	-28	-26
38	R MTG	Right middle temporal gyrus	55	-41	-2	80	L MTG	Left middle temporal gyrus	-58	-37	-3
39	R MTGpole	Right temporal pole: middle temporal gyrus	38	13	-35	81	L MTGpole	Left temporal pole: middle temporal gyrus	-37	13	-36
40	R STGpole	Right temporal pole: superior temporal gyrus	44	12	-20	82	L STGpole	Left temporal pole: superior temporal gyrus	-42	12	-21
41	R STG	Right superior temporal gyrus	57	-25	7	83	L STG	Left superior temporal gyrus	-56	-22	6
42	R Th	Right thalamus	11	-20	6	84	L Th	Left thalamus	-11	-20	7

S2 Table. Table listing the links of the significant directed network in gamma in the memory condition.

Leading	Lagging	Normalized grand average	t value
L Th	R SCG	1.00	4.00
L Th	L AMY	0.88	3.77
L HIPP	L AMY	0.86	4.48
L PCC	L AMY	0.78	5.32
R PCC	R SCG	0.74	3.35
L PCC	L ITG	0.70	3.58
L PHG	L AMY	0.69	3.56
R pSFG	L ACC	0.61	4.98
L pSFG	L ACC	0.54	3.94
R STGpole	L SFGmedOrb	0.51	3.54
R SFGmedOrb	L SFGmedOrb	0.51	4.80
R pSFG	R ACC	0.47	3.60
R SFGmedOrb	L ACC	0.46	3.81
L pSFG	R ACC	0.43	3.33
R pSFG	R SCG	0.36	3.39
L ITG	L MOG	0.34	3.56

Annotations: L: left, R: Right, Th: thalamus, SCG: subcallosal gyrus, AMY: amygdala, HIPP: hippocampus, PCC: posterior cingulate cortex, ITG: inferior temporal gyrus, PHG: parahippocampal gyrus, SFG: superior frontal gyrus, pSFG: posterior SFG, ACC: anterior cingulate cortex, STGpole: polar part of the superior temporal gyrus, SFGmedOrb: medial orbital part of the SFG, ITG: inferior temporal gyrus, MOG: middle occipital gyrus.

S3 Table. Table listing the links of the significant directed network in high theta in the memory condition.

Leading	Lagging	Normalized grand average	t value
R PCC	R SCG	1.00	3.74
R PCC	L SCG	1.00	3.90
R PCC	R Rec	1.00	3.67
L PCC	L SCG	0.94	3.75
L PCC	R SCG	0.93	3.61
L PCC	L Rec	0.91	3.47
L PCC	R Rec	0.91	3.42
L PCC	R ACC	0.84	3.29
R PCC	L Th	0.83	3.50
R PCC	R Th	0.82	3.47
L PCC	L Th	0.79	3.42
L PCC	R Th	0.74	3.29
L PCC	R MCC	0.67	3.61
L PCC	L MCC	0.66	3.54
L PrCu	R pSFG	0.57	3.55
R ITG	R SFGorb	0.52	3.99
R ITG	R SCG	0.51	3.47
L PrCu	R MCC	0.50	3.80
R ITG	R SFGmedOrb	0.44	3.30
R ITG	L SFGorb	0.40	3.38
R ITG	R IFGoper	0.37	3.90

Annotations: L: left, R: Right, PCC: posterior cingulate corteax, SCG: subcallosal gyrus, Rec: gyrus rectus, ACC: anterior cingulate cortex, Th: thalamus, MCC: median cingulate cortex, PrCu: precuneus, SFG: superior frontal gyrus, pSFG: posterior SFG, ITG: inferior temporal gyrus, SFGorb: orbital part of the SFG, ITG: inferior temporal gyrus, SFGmedOrb: medial orbital part of the SFG.

S4 Table. Table listing the links of the significant phase-amplitude coupling directed network when contrasting the memory vs. math conditions.

Phase	Amplitude	Normalized grand average	t value
R INS	L MTG	1.00	3.93
R ANG	L MTG	0.95	3.51
R MTGpole	L PHG	0.95	3.31
R STG	L FUSI	0.89	3.55
R MTGpole	L Rec	0.85	3.55
R MTGpole	L SFGorb	0.84	3.40
R ANG	L FUSI	0.82	3.43
L HIPP	R SPG	0.82	3.37
L SFGmed	L STG	0.81	4.31
L MCC	R SPG	0.79	3.39
L ITG	L SFGmedOrb	0.78	3.32
L PHG	L SFGmedOrb	0.77	3.36
L HIPP	L SFGmedOrb	0.77	3.35
L STG	L SFGmedOrb	0.76	3.50
R AMY	R SPG	0.76	3.41
L LING	L SFGmedOrb	0.76	3.42
R INS	L Rec	0.74	3.52
R PreCG	L FUSI	0.71	3.83
L MTG	L Rec	0.64	3.42
R SOG	R SPG	0.61	3.52

Annotations: L: left, R: Right, INS: insula, MTG: middle temporal gyrus, ANG: angular gyrus, MTGpole: polar part of the MTG, PHG: parahippocampal gyrus, STG: superior temporal gyrus, FUSI: fusiform gyrus, Rec: gyrus rectus, SFG: superior frontal gyrus, SFGorb: orbital part of the SFG, HIPP: hippocampus, SPG: superior parietal gyrus, SFGmed: medial part of the superior frontal gyrus, MCC: median cingulate cortex, ITG: inferior temporal gyrus, SFGmedOrb: medial orbital part of the SFG, AMY: amygdala, LING: lingual gyrus, PreCG: precentral cortex, SOG: superior occipital gyrus.