EXPERIMENTAL EVOLUTION OF A PHEROMONE SIGNAL

Short title: Experimental sex pheromone evolution

Thomas Blankers[1,2][†], Elise Fruitet[1][†], Emily Burdfield-Steel[1][†], Astrid T. Groot[1]

[1] *Evolutionary and Population Biology, Institute for Biodiversity and Ecosystem Dynamics,*

University of Amsterdam

[2] Author for correspondence: thomasblankers<at>gmail.com

[†] These authors contributed equally

SUMMARY STATEMENT

This study reveals a mechanism that can facilitate evolution of sex pheromone components under

selection independent of other components of the sex pheromone that are not under selection.

ABSTRACT

Sexual signals are important in speciation, but understanding their evolution is complex as these

signals are often composed of multiple, genetically interdependent components. To understand

how signals evolve, we thus need to consider selection responses in multiple components and

account for the genetic correlations among components. One intriguing possibility is that

selection changes genetic correlations between components under selection and those not under

selection, reducing evolutionary constraints. However, this hypothesis remains largely untested

empirically. In this study, we investigate the evolutionary response of the multi-component

female sex pheromone blend of the moth *Heliothis subflexa* to 10 generations of artificial

selection. We observed a selection response of about 3/4s of a phenotypic standard deviation in

the components under selection. Interestingly, other pheromone components that are

biochemically and genetically linked to the components under selection did not change. We

found that this component-specific selection response was likely facilitated due to reduced

genetic coupling between the component under selection and the components not under

selection, resulting from changes in genetic covariances within the 10 generations of selection.

Our findings provide rare empirical support for an intriguing mechanism by which a sexual

signal can respond to selection without possible constraints from indirect selection responses.

Key words: Sexual communication, Genetic variance-covariance, Pheromone, Artificial

selection, Lepidoptera

INTRODUCTION

Animal taxa that engage in sexual communication typically show high among-species diversity

in sexual signals (Andersson, 1994; Coyne and Orr, 2004; Ritchie, 2007; Schaefer and Ruxton,

2015; Wiens and Tuschhoff, 2020). Sexual signals are generally hypothesized to diversify due to

directional or disruptive selection (Ritchie, 2007; Schaefer and Ruxton, 2015; West-Eberhard,

2014; Wilkins et al., 2013), although their evolution is still a mystery in many species. Since

sexual signals play an important role in the origin and maintenance of species and contribute to

biodiversity (Coyne and Orr, 2004), it is important to assess whether there are constraints to their

selection response and identify the mechanisms that can mitigate those constraints.

Understanding selection responses in sexual signals is challenging, because signals are often

composed of multiple components (Candolin, 2003; Higham and Hebets, 2013; Rowe, 1999).

For example, mating songs can vary both in pitch and in rhythm (Blankers et al., 2015; Tanner et

al., 2017; Wilkins et al., 2015), color signals can be composed of multiple, functionally distinct

patches (Cole and Endler, 2015; Grether et al., 2004), and sex pheromones are often blends of

multiple chemical compounds (Ferveur, 2005; Linn et al., 1987). To understand how

multicomponent signals evolve, we thus need to consider the selection response in multiple

dimensions simultaneously. Signal components can have a shared genetic or developmental

basis, or can be subject to correlated selection pressures (Armbruster et al., 2014; Cheverud,

1996). The resulting genetic correlations among signals components can influence how selection

on the phenotype translates to changes in the underlying genotypes (Chenoweth and Blows,

2006). To understand how the genotype-phenotype map of sexual signals influences the selection

response, we thus need to determine the genetic correlations between the different signal

components.

Statistical frameworks in quantitative genetics, in particular the (multivariate) breeder's equation,

allow us to predict and quantitatively understand selection responses in correlated traits. In this

framework, the response to selection is a function of the genetic (G) and phenotypic (P)

variance-covariance matrix and the selection gradient: selection acts on the P matrix and the

resulting response is constrained by the G matrix (Lande, 1979; Lande and Arnold, 1983; Lynch

and Walsh, 1998). The difficulty in predicting selection responses of multivariate traits is that

selection acting on multiple components may be counterbalancing, e.g. directional selection on

one trait, but stabilizing selection on correlated traits, resulting in evolutionary constraints

(Barton and Turelli, 1989). Counterbalancing selection is likely prevalent in the evolution of

sexual signals, as choosing individuals may favor higher or lower values of some component of

the signal, while changes in correlated components may result in reduced mate attraction.

These evolutionary constraints can be overcome if genetic variances and covariances themselves

respond to selection, thus reshaping the G matrix (Arnold et al., 2008; Barton and Turelli, 1989;

Eroukhmanoff, 2009; Jones et al., 2003; Melo and Marroig, 2014; Revell, 2007; Roff and

Fairbairn, 2012). Theory predicts that the G matrix will vary through time, because on one hand

selection erodes genetic variance (Barton and Turelli, 1989; Estes and Arnold, 2007), while on

the other hand mutation and introgression add new variation, albeit more slowly. Moreover,

genetic correlations can respond to selection directly, especially if they result from interactions

among unlinked genetic loci affecting the co-expression of multiple traits (Wolf et al., 2005), or

from selection acting on correlations directly (Armbruster et al., 2014; Roff and Fairbairn, 2012;

Svensson et al., 2021). Interestingly, directional selection can change genetic covariances and

increase modularity, meaning that groups (modules) of traits become genetically independent

from other modules, in a way that facilitates the phenotypic response to selection (Melo and

Marroig, 2014). Empirical work has shown that patterns of covariation can indeed evolve both

across populations and species in nature (Bégin and Roff, 2003; Berner et al., 2010; Blankers et

al., 2017; Garant et al., 2008; Gosden and Chenoweth, 2014) and during artificial selection

experiments (Careau et al., 2015; Hine et al., 2011; Uesugi et al., 2017). However, it is still

unclear how changes in the phenotypic selection response are related to changes in genetic
(co)variance through time.

In this study, we explored the response in phenotypic means and genetic (co)variances to
artificial selection on the female sex pheromone of the moth *Heliothis subflexa* (Lepidoptera,
Noctuidae). Like many other moths, *H. subflexa* females secrete a sex pheromone blend to which
conspecific males are attracted. These sex pheromone blends are species-specific and vary
among species in both the presence/absence of components as well as in relative amounts (or
ratios) of the components (Cardé and Haynes, 2004; Schneider, 1992).

The sex pheromone blend of *H. subflexa* females consists of 11 compounds, with the following
components that are critically important for conspecific male attraction: (Z)-11-hexadecenal
(Z11-16:Ald) as the major sex pheromone component, (Z)-9-hexadecenal (Z9-16:Ald) and (Z)-
11-hexadecenol (Z11-16:OH) as the two secondary sex pheromone components, without which
H. subflexa males are not attracted (Groot et al., 2007; Vickers, 2002). Interestingly, the acetate
esters (Z)-7-hexadecenyl acetate (Z7-16:OAc), (Z)-9-hexadecenyl acetate (Z9-16:OAc), and (Z)-
11-hexadecenyl acetate (Z11-16:OAc), from here on referred to as "acetates", have a dual role:
these acetates attract conspecific males, while simultaneously repelling males of *H. virescens*
(Groot et al., 2006; Vickers and Baker, 1997). In geographic regions where *H. virescens* is
present, acetates are more abundant in the *H. subflexa* pheromone compared to where this
species is absent (Groot et al., 2009a). This suggests that the acetates are subject to divergent
selection across a geographic cline. The relative amounts of the other components are
hypothesized to be under stabilizing selection across the range, as in general in moth pheromone
communication the mean blend is preferred over deviations from the mean (Allison and Cardé,
2008; Groot et al., 2010; Kárpáti et al., 2013; Linn et al., 1997; Löfstedt, 1990; Zhu et al., 1997).

To explore the selection response of the acetates, we selected for higher and lower amounts of
acetates during 10 generations of truncation selection. Since the acetates vary geographically
(Groot et al., 2009a) and have a genetic basis that is partially independent of other components
(Groot et al., 2009b), we hypothesized that the relative amount of acetates can evolve in response
to univariate selection for higher/lower acetates but that genetic variance will be reduced after
selection. However, since acetates also partially share their genetic basis with other components
and since all sex pheromone compounds are produced through the same biosynthetic pathway,

we also hypothesized that there will be indirect selection responses in other pheromone

components, specifically in the unsaturated aldehydes, Z9-16:Ald and Z11-16:Ald, and alcohol,

Z11-16:OH (Fig 1). Since some of the correlations among components may reduce the selection

response, we also expected genetic covariances to change during selection, specifically in a way

that facilitates the phenotypic selection response.

Figure 1. Simplified biosynthetic pathway of the *Heliothis subflexa* sex pheromone, modified from (Groot et al.,

2009b) with permission. Desaturation and β-oxidation produce mono-unsaturated acyl-CoA precursors from 18 or

16 carbon acyl CoA derivatives, which are then modified to form acetate esters, alcohols, and aldehydes through

specific enzymatic conversions. The Δ9 and Δ11 desaturases create a double bond between the 9th and 10th or 11th

and 12th carbon of the 18- or 16-carbon acyl-CoA derivatives, respectively. β-oxidation shortens the chain length

from 18 to 16 carbons. Note that (Z)9-16 acyl-CoA can be formed through two alternative routes. The compounds

present in the pheromone glands of *H. subflexa* are in grey boxes.

METHODS

Insect rearing

The laboratory population of *H. subflexa* was founded from animals collected in the field in 2006 and has been reared at the University of Amsterdam since 2011 using single pair matings to maintain genetic diversity. There has been occasional exchange among *H. subflexa* populations at North Carolina State University, Amsterdam, and the Max Planck Institute for Chemical Ecology in Jena. Eggs collected from single pair matings were kept in Petri dishes (Ø 85 mm) with artificial wheat germ/soy flour-based diet (BioServ Inc., Newark, DE, USA) at room temperature for approximately 10 days, after which larvae were reared in separate individual 37-mL cups filled with the same artificial diet and kept at 25°C and 60% relative humidity with 14h:10h light-dark cycle. Upon emergence, adults were provided sugar water. Pairs of males and females were housed in 375 mL paper cups covered with gauze and kept under the same conditions as the late-instar larvae and virgin adults. The mating pairs were provided with sugar water. To stimulate oviposition, a freshly cut gooseberry fruit was placed on top of the gauze. Once the eggs began to hatch, the gauze and the eggs and larvae on it, were transferred to Petri dishes, which were then placed at room temperature until larvae were transferred to individual cups. The females that produced fertile offspring were collected and phenotyped, if still alive. All matings were assigned unique numbers. This way, we obtained a full pedigree of all individuals in the selection and control lines.

Phenotyping

Female sex pheromone can be extracted from mated and old females by injecting them with Pheromone Biosynthesis Activating Neuropeptide, i.e. PBAN (Groot et al., 2005). Females were injected with a 7.5 pmol (2 µL of a 0.0146 µg/µL) PBAN solution to activate pheromone production in the mated females. After a 90-minute incubation time, female glands were extruded by squeezing the abdomen then fixed by firmly holding the abdomen with forceps just anterior of the gland. The gland was excised with microdissection scissors. Excess abdominal tissue and eggs that remained in the ovipositor were removed, after which the glands were

submerged in 50 μL hexane containing 200 ng pentadecane as an internal standard. After 30

minutes, the glands were removed and the extracts stored at -20°C until analysis.

Pheromone extracts were analyzed by injecting the concentrated samples into a splitless inlet of a

7890A GC (Agilent Technologies, Santa Clara, CA, USA). The area under the pheromone peaks

was calculated using integration software implemented in Agilent ChemStation (version

B.04.03). Pheromone peak areas were obtained for the 11 pheromone components: two 14-C

aldehydes (14:Ald, Z7-14:Ald), four 16-C aldehydes (16:Ald, Z7-16:Ald, Z9-16:Ald, Z11-

16:Ald), the three acetates (Z7-16:OAc, Z9-16:OAc, Z11-16:OAc), and two 16-C alcohols (Z9-

16:OH, Z11-16:OH). Z7-16:Ald and Z9-16:Ald were difficult to separate by GC and were

therefore integrated as one peak (referred to Z9-16:Ald). Absolute amounts (in ng) of each

compound were calculated relative to a 200 ng pentadecane internal standard. All downstream

analyses were done in R 3.6.1 (R Core Team, 2019). Samples containing < 20 ng were excluded,

because the ratios of the components in such low titers cannot be reliably measured in the

chromatogram. Relative amounts were calculated by dividing the absolute amounts by the total

amount across all 11 components.

Data transformation

Selection was performed based on relative amounts of acetates. However, describing

relationships among relative amounts is problematic, because they sum to 100%, thereby

mathematically constraining the (co)variation in the pheromone and biasing the analysis. We

therefore transformed pheromone measurements to log-contrasts for all down-stream analyses in

this study. This approach breaks the interdependency and normalizes the data. Since the divisor

used in the contrast of the variable can no longer be part of any downstream analyses, we chose

14:Ald as the divisor, because this component has a small but clearly detectable peak in the

chromatogram, while it is irrelevant for male response behavior (Heath et al., 1990). Prior to

downstream analyses, samples with a χ^2-distributed Mahalanobis distance score (calculated using

the 'mahalanobis' function in the 'stats' package) that exceeded a threshold value corresponding

to a Bonferroni-corrected P-value < 0.05 were removed, which resulted in removing 64 out of a

total of 2,861 samples. These samples showed abnormal pheromone ratios and were present

across all selection and control lines and generations. We are therefore confident that these

samples represent outliers, e.g. due to extraction or measurement errors, and are not

representative of any relevant biological phenomena.

Selection

Since phenotyping consisted of extracting the sex pheromone gland invasively, selection was

performed post-mating. Each generation, we continued with those families that had a maternal

phenotype satisfying an increasingly stringent threshold. Initially, a so-called "high" and "low"

line were started with offspring from females with a relative amount of acetates above 22% or

below 16%, respectively. These values were based on the distribution of the relative amounts of

acetates in the starting population, representing the 1st and 3rd quantile. These thresholds were

kept for the first 3 generations. In subsequent generations, we increased these thresholds to >24%

or <14% (generation $4 - 5$), and >26% or <12% (generation $6 - 9$). The high line consisted of a

total of 2,236 breeding females across nine generations, ranging from $189 - 295$ matings per

generation. After outlier removal, a total of 1,234 high line females were phenotyped, or between

89 and 171 per generation during the selection phase. The low line consisted of 2,250 breeding

females ($189 - 296$ per generation) and, after outlier removal, phenotypes were measured for a

total of 1,180 females ($57 - 169$ per generation). We stopped selecting in generation 10, but

continued to phenotype ~25 females per line and per generation until generation 13. Throughout

the selection experiment, we also maintained a control line from which we phenotyped between

6 and 111 females (median 20 females) per generation. For detailed sample sizes see Table S1.

Phenotypic selection response

To test the hypothesis that acetates can respond to selection and that other pheromone

components change due to indirect selection, the selection responses as measured by the log-

contrasts were visualized using the R-package 'ggplot'. In addition, we also calculated the

average difference between selected and control lines in units of standard deviation of the

starting population.

Genetic (co)variance selection response

To test whether genetic variances decreased with selection and whether the structure of the

genetic variance-covariance matrix was affected by selection, we first ran multi-response animal

models implemented in the R-package 'MCMCglmm'. We limited our analyses to the four

components that have been show to affect male response: Z9-16:Ald, Z11-16:Ald, Z11-16:OAc,

and Z11-16:OH (in all cases, the log-contrast to 14:Ald was used). Model formulation roughly

followed the MCMCglmm manual (Hadfield, 2010) and Jarrod Hadfield's course notes

(Hadfield, 2012), as well as the animal model tutorial by Pierre de Villemereuil (Villemereuil,

2012). High and low line individuals had separate pedigrees, but shared ancestors for which we

had pedigree data up to three generations prior to the onset of the experiment (in total 1,065

breeding females). To minimize the influence from priors on the covariances, for which we had

no prior expectations with high degrees of belief, we formulated flat, uninformative priors for the

variance-covariance structure of the random effect ('animal'). To evaluate the models, we

checked chain convergence and assessed effective sample sizes and levels of autocorrelation. To

check whether the prior did not contribute unduly to posterior estimates, we also compared

genetic (co)variance estimates among univariate, bi-variate, and full (tetra-variate) models and

using different, more informative priors. We found models with flat priors to perform best.

To observe changes in genetic (co)variances at higher sample sizes, we partitioned the data in the

starting generation (generation 0) and generation blocks of three generations each ($1 - 3, 4 - 6, 7$

$- 9$) for both the high and low line. For each generation block, we obtained posterior estimates of

the additive genetic variance, V_A for each of the four log-contrasts from the multi-response

animal model. For each posterior sample of V_A, we calculated the coefficient of additive genetic

variance, CV_A (where $CV_A = \frac{V_A}{phenotypic\ mean}$), which provides a standardized measure of the

evolvability of a trait that is independent of other variance components. We also extracted the

posterior genetic correlations and visually inspected pairwise genetic correlations between Z11-

16:OAc and each of the other three components. Lastly, we examined changes in the G matrix by

inspecting the trait loadings on genetic principle components. The four genetic principle

components were obtained for every posterior sample using the 'eigen' function. We focused on

the first two axes, gPC1 (also known as g_{max}, the direction in phenotypic space containing the

largest fraction of the genetic variance) and gPC2, because these axes jointly described $> 90\%$ of

the genetic variance (see Results).

RESULTS

Selection response in sex pheromone

All three acetates responded to selection for higher/lower relative total amounts of acetates in the

high and low line, respectively (Fig 2). The response was more pronounced in the more abundant

Z9 and Z11 isomers compared to the less abundant Z7-16:OAc (Fig 2). The selection response

was characterized by an immediate response in the direction of selection (from generation 0 to

generation 1), followed by a gradual progression towards increasing/decreasing relative amounts

in the respective selection lines. In the low line, the selection response flattened after cessation of

selection, while in the high line, the average ratio of Z11:16:OAc to 14:Ald was higher in

generations 11-13 compared to generations 7-10 (Fig 2D).

Fig 2. Selection response in the acetates. Values shown are mean ± SEM. The control line is indicated in black, the

high line in red (dotted) and the low line in blue (dashed). The dashed vertical line indicates the generation with

adult females that developed from the last offspring that went through selection. A – C: Selection responses of the

three acetates log-contrasted to 14:Ald. For the control line, 111 females were phenotyped in generation zero, 71 in

generation one, and between 6 and 36 for the remaining generations. For the low and high line, respectively 189 –

296 and 189 - 295 females per generation were phenotyped. D: Difference in phenotypic means between selection

lines and control in units of starting population standard deviation. The points and error bars with brighter hues show

the average ± standard deviation for generation 1 – 3, 4 – 6, 7 – 10, and 11 – 13, respectively. E: Selection response

of the relative total amount of acetates.

The other pheromone components showed no differentiation among selection lines or between

selected and control lines (Fig 3A-C). The total amount of pheromone measured across the 11

biologically active components remained constant through time, indicating that changing ratios

were independent of the pheromone titer (Fig 3D).

Fig 3. Indirect selection response. Values shown are mean ± SEM for log-contrasts to 14:Ald of the three pheromone

components that make up the minimal blend for male attraction and for the absolute amount of pheromone across all

11 biologically active components. The control line is indicated in black, the high line in red and the low line in

blue. Dashed vertical lines indicate the generation with adult females that developed from the last offspring that

went through truncation selection. Sample sizes as in Fig 2.

Selection response in genetic (co)variances

In testing whether the selection response was associated with (i) a reduction in genetic variance

in Z11-16:OAc (a change in the diagonal elements of the G matrix) and (ii) re-orientation of

genetic covariances among pheromone components (changes in the off-diagonal elements of the

G matrix), we found no evidence for decreasing genetic variance in Z11-16:OAc (Fig 4A) or in

the other components (Fig S1). In contrast to genetic variances, we did find variation in the

genetic covariance structure across generations. First, the posterior mode of the pairwise genetic

correlations differed between the starting and selected generations. Although the change in the

covariance between Z11-16:OAc and the two aldehydes was limited (Fig 4B,C), genetic

covariance between Z11-16:OAc and Z11-16:OH switched direction (from negative to positive)

and became more pronounced, especially in the low line (Fig 4D).

Fig 4. Genetic variation and correlations. The posterior mode and 95% HPD interval of the genetic correlation
between the log-contrast of Z11-16:OAc on 14:Ald and each of the other three log-contrasts is shown for the starting
generation (S, in black), the low (blue) and high (red) lines. 111 females were phenotyped for the starting
generation. For the low line, 443, 284, and 330 phenotyped females were in generation blocks 1-3, 4-6, and 7-9,
respectively. For the high line 465, 324, and 344 phenotyped females were in generation block 1-3, 4-6, and 7-9,
respectively. Pedigree data were obtained for 1,065 mating pairs prior to the first generation of selection and for an
additional 2,250 and 2,236 pairs for the low and high line, respectively.

Second, the partitioning of genetic variance among gPC1 and gPC2 evolved. The first two
genetic principle components changed in magnitude of their eigenvalue (i.e. the amount of
genetic variance they describe) and in their loadings (i.e. in their correlation with the pheromone
components; Fig 5). More of the total genetic variance across the four components was in the
direction of gPC1 in the high and low lines (~80%) compared to the starting generation (59.1%).
This means that more of the genetic variance of the sex pheromone was biased towards a single
dimensions in phenotypic space. Moreover, variation in the component under selection, Z11-
16:OAc, was more strongly associated with gPC2 and less with gPC1 in generations after
compared to before selection, while the other traits showed a trend in the opposite direction (Fig
5). These changes in the G matrix resulted in a more modular pheromone blend, because Z11-
16:OAc, which was under selection here, became uncoupled from the components that were not
under selection.

Fig 5. Eigen-analysis of genetic principle components. For the starting generation and generation 1-3, 4-6, and 7-9
of the low (top) and high (bottom) lines, the posterior mode of the correlation between each of the pheromone ratios

and the first and second principle components is shown by the direction and size of the arrows. The genetic variance
explained by PC1 and PC2 is indicated on the axes.

DISCUSSION

In this study, we investigated how the selection response depends on and shapes the genetic
architecture of a multicomponent sex pheromone signal. Through truncation selection, we
gradually increased and decreased the relative amount of acetates across 10 generations. As we
hypothesized, we found that the acetates responded readily to selection, diverging three-quarters
of a standard deviation away from the control line in 10 generations in both the high and low
lines. However, in contrast to our hypothesis that selection would also result in indirect responses
in the other traits, we found that the response was limited to the acetates only. In addition, we
found that levels of genetic variance were constant through time, opposite to the expected
erosion of genetic variance in response to selection. Lastly, in line with our expectations, the
genetic covariance structure changed in a way that likely facilitated a selection response in the
pheromone. We discuss each of these results further below.

Univariate responses in multicomponent signals

In nature, correlational selection may be quite common (Svensson et al., 2021) and sexual
signals are often found to be subject to multivariate stabilizing and directional selection (Bentsen
et al., 2006; Blankers et al., 2015; Brooks et al., 2005; Devigili et al., 2015; Fisher et al., 2009;
Gerhardt and Brooks, 2009; Hine et al., 2011; Oh and Shaw, 2013; Ryan and Rand, 2003;
Tanner et al., 2017). In our experiment, we applied a univariate selection gradient on the relative
amount of Z11-16:OAc to consider the evolvability of multivariate traits to univariate selection
gradients. We justified this univariate selection based on the found geographic variation in the
relative amount of Z11:16OAc in *H. subflexa*. Females are likely specifically selected for higher
rates of acetates to avoid heterospecific mate attraction in regions where the congener *H.*
virescens occurs (Groot et al., 2009a).

Our observed selection response shows that there are no phenotypic or genetic constraints to the
evolution of Z11-16:OAc (or the other acetates), as we reached similar or even higher levels of
divergence compared to those observed across the range of *H. subflexa*. After 10 generations in

our selection experiment, the high line individuals had on average almost 20% more Z11-
16:OAc compared to the low line (Fig 2E), while *H. subflexa* populations in the eastern US were
found to have blends with up to 10% more Z11-16:OAc compared to populations in the south-
western US and Mexico (Groot et al., 2009a). The selection response that we found was also
surprisingly univariate, as only the three acetate esters and none of the other components showed
divergence between the high and low lines. This indicates that the acetates can evolve
independently from the other components. This is surprising, both in the context of the observed
genetic correlations in the starting population and in later generations (Fig 4) and in the context
of what is known about the shared biochemical pathways across the pheromone components
(Groot et al. 2009a; Fig 1). However, it is not an uncommon result of artificial selection
experiments (Hill and Caballero, 1992; Saltz et al., 2017), including for sexual signals (Ritchie
and Kyriacou, 1996). Thus, genetic correlations among sexual signal components may quickly
dissolve in the absence of correlational selection, which likely facilitates their evolution on short
time scales in response to directional selection.

We acknowledge that our study does not include replicate lines in the experiment. This is
because each selection line consisted of hundreds of single pair matings per generation, as *H.*
subflexa is highly sensitive to inbreeding. We thus needed to maintain large populations of
individuals that are mated in pairs and also individually reared to avoid cannibalism, which
makes these experiments very laborious. Therefore, we chose to rear two lines in separate
directions rather than two replicates of only the high or low line. The fact that we found parallel
results between the lines in the selection response both in phenotypic means and genetic
covariances lends confidence to the patterns observed. However, we do avoid drawing
conclusions based on differences between the lines, because these can both reflect differential
selection effects or sampling variance.

No change in genetic variance

The effect of selection on genetic variation is a key topic in evolutionary biology. Strong
stabilizing or directional selection is expected to erode genetic variance (Barton and Turelli,
1989; Estes and Arnold, 2007). However, when many loci are contributing to variation in the
trait under selection, selection might leave no detectable signature on the genetic variance
associated with that trait (Bulmer, 1971; Johnson and Barton, 2005). Although our truncated

selection increased and decreased the phenotypic mean by three-quarters of a standard deviation

in the high and low line, respectively, we observed no significant decrease in the coefficient of

additive genetic variance. There are several explanations for this observation. First, several

genetic factors may be involved. Previously, genetic mapping studies revealed several QTL

underlying difference in sex pheromone composition in *H. subflexa* (Groot et al., 2013). Since

variation in pheromone composition in *H. subflexa* is also explained by fitness variation

(Blankers et al., 2021), many different genetic factors likely contribute small additional fractions

to the total variance. Another explanation comes from population genetics: re-orientation of

genetic covariances in response to bottlenecks may free up additive genetic variance, thereby

paradoxically increasing levels of genetic variance (Carson, 1990; Templeton, 2008). Since

truncation selection is effectively a non-random bottleneck of the population, this may also

explain the maintenance of genetic variance that we observed. Lastly, to avoid inbreeding

depression, we maintained large populations and avoided mating first and second-degree

relatives. Such a mating scheme likely counteracts some loss in genetic variability due to

selection (Du et al., 2021).

Changing genetic covariance

In contrast to genetic variance, we observed a clear trend in changing genetic covariances.

Specifically, the correlation between Z11-16:OH and Z11-16:OAc changed in both sign and

magnitude. Since both Z11-16:OH and Z11-16:OAc reduce heterospecific mate attraction

(Vetter and Baker, 1983; Vickers and Baker, 1997), and both traits covary with fitness and are

condition-dependent (Blankers et al., 2021), the negative genetic correlation in the starting

population could be the result of a trade-off that is selected against when artificial selection

favors higher rates of acetates.

Tracking changes in G matrices through time, we observed another interesting phenomenon:

after selection, the genetic variance in Z11-16:OAc was more strongly associated with genetic

(g) PC2 compared to before selection, while genetic variance described by gPC1 increased

relative to the other gPCs. Thus, more of the genetic variance was allocated in fewer dimensions

of genetic variation, while the component under selection was uncoupled from the major

dimension of genetic variance. We acknowledge that the observed changes are subtle and that

they cannot be unequivocally attributed to selection. However, we observed similar patterns of

G matrix evolution in the high and low line and these patterns fit the prediction that genetic

variances and covariances are reoriented to align the phenotype with the dominant direction of

selection (Melo and Marroig, 2014). These changes in the G matrix can likely facilitate a more

specific response where only Z11-16:OAc evolves in the absence of indirect selection responses:

relative to the selection regime, the modularity (i.e. the genetic independence of trait modules) of

the pheromone increased.

Conclusion

From our results, we conclude that: 1) pheromone components in *H. subflexa* can evolve in

response to selection independent of other components of the sex pheromone; 2) selection need

not erode the genetic variance in order to drive this phenotypic change; 3) selection alters the

genetic correlations among pheromone components. Our study thus shows that univariate

selection responses in multicomponent sexual signals are possible, even though pairwise genetic

correlations were high both prior to and during selection. Our study also shows that sexual signal

components can respond to selection without reductions in genetic variance, but with rapid

changes in the genetic covariance structure. These results correspond to geographic variation in

the sexual signal found in this species and help to explain how sexual signals in general can

respond to selection without running into constraints from indirect selection responses or

depleted genetic variation. This may be one reason why sexual signals have evolved rapidly and

repeatedly throughout the evolutionary history of animal taxa, thereby contributing to the

speciation process.

ACKNOWLEDGEMENTS

418 We thank Dennis van Veldhuizen for his invaluable contribution to the rearing of our moths. We

419 also thank Laura Caton, Frederica Lotito, and Lilian Seip for their help with the selection lines.

420 We further thank Jet ten Berge for digitizing the pedigree, Michiel van Wijk for commenting on

421 a previous version of the manuscript, and all members of the Groot lab for their helpful and

422 insightful discussions.

COMPETING INTERESTS

424 The authors declare no conflict of interest

AUTHOR CONTRIBUTIONS

FUNDING

This study was funded by the following grants: the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 794254 awarded to TB, NWO grant award ALWOP.2015.075 awarded to ATG, an IMPRS grant of the Max Planck Society awarded to EF, and an ASAB Research grant awarded to EBS.

DATA ACCESSIBILITY

Data and analysis scripts will be archived in Dryad.

FIGURE LEGENDS

Figure 1. Simplified biosynthetic pathway of the *Heliothis subflexa* sex pheromone, modified from (Groot et al., 2009b) with permission. Desaturation and β-oxidation produce mono-unsaturated acyl-CoA precursors from 18 or 16 carbon acyl CoA derivatives, which are then modified to form acetate esters, alcohols, and aldehydes through specific enzymatic conversions. The Δ9 and Δ11 desaturases create a double bond between the 9th and 10th or 11th and 12th carbon of the 18- or 16-carbon acyl-CoA derivatives, respectively. β-oxidation shortens the chain length from 18 to 16 carbons. Note that (Z)9-16 acyl-CoA can be formed through two alternative routes. The compounds present in the pheromone glands of *H. subflexa* are in grey boxes.

Fig 2. Selection response in the acetates. Values shown are mean ± SEM. The control line is indicated in black, the high line in red (dotted) and the low line in blue (dashed). The dashed vertical line indicates the generation with adult females that developed from the last offspring that went through selection. A – C: Selection responses of the three acetates log-contrasted to 14:Ald. For the control line, 111 females were phenotyped in generation zero, 71 in generation one, and between 6 and 36 for the remaining generations. For the low and high line, respectively 189 – 296 and 189 - 295 females per generation were phenotyped. D: Difference in phenotypic means between selection lines and control in units of starting population standard deviation. The

points and error bars with brighter hues show the average ± standard deviation for generation 1 –

3, 4 – 6, 7 – 10, and 11 – 13, respectively. E: Selection response of the relative total amount of

acetates.

Fig 3. Indirect selection response. Values shown are mean ± SEM for log-contrasts to 14:Ald of

the three pheromone components that make up the minimal blend for male attraction and for the

absolute amount of pheromone across all 11 biologically active components. The control line is

indicated in black, the high line in red and the low line in blue. Dashed vertical lines indicate the

generation with adult females that developed from the last offspring that went through truncation

selection. Sample sizes as in Fig 2.

Fig 4. Genetic variation and correlations. The posterior mode and 95% HPD interval of the

genetic correlation between the log-contrast of Z11-16:OAc on 14:Ald and each of the other

three log-contrasts is shown for the starting generation (S, in black), the low (blue) and high (red)

lines. 111 females were phenotyped for the starting generation. For the low line, 443, 284, and

330 phenotyped females were in generation blocks 1-3, 4-6, and 7-9, respectively. For the high

line 465, 324, and 344 phenotyped females were in generation block 1-3, 4-6, and 7-9,

respectively. Pedigree data were obtained for 1,065 mating pairs prior to the first generation of

selection and for an additional 2,250 and 2,236 pairs for the low and high line, respectively.

Fig 5. Eigen-analysis of genetic principle components. For the starting generation and generation

1-3, 4-6, and 7-9 of the low (top) and high (bottom) lines, the posterior mode of the correlation

between each of the pheromone ratios and the first and second principle components is shown by

the direction and size of the arrows. The genetic variance explained by PC1 and PC2 is indicated

on the axes.

REFERENCES

Allison, J. D. and Cardé, R. T. (2008). Male pheromone blend preference function measured in

choice and no-choice wind tunnel trials with almond moths, Cadra cautella. *Anim. Behav.*

75, 259–266.

Andersson, M. (1994). *Sexual Selection*. Princeton University Press.

Armbruster, W. S., Pélabon, C., Bolstad, G. H. and Hansen, T. F. (2014). Integrated

phenotypes: understanding trait covariation in plants and animals. *Philos. Trans. R. Soc. B*

Biol. Sci. **369**, 20130245.

Arnold, S. J., Bürger, R., Hohenlohe, P. A., Ajie, B. C. and Jones, A. G. (2008).

Understanding the evolution and stability of the G-matrix. *Evolution* **62**, 2451–2461.

Barton, N. and Turelli, M. (1989). Evolutionary Quantitative Genetics: How Little Do We

Know. *Annu. Rev. Genet.* **23**, 337–370.

Bégin, M. and Roff, D. A. (2003). The constancy of the G matrix through species divergence

and the effects of quantitative genetic constraints on phenotypic evolution: a case study in

crickets. *Evolution* **57**, 1107–1120.

Bentsen, C. L., Hunt, J., Jennions, M. D. and Brooks, R. (2006). Complex multivariate sexual

selection on male acoustic signaling in a wild population of Teleogryllus commodus. *Am.*

Nat. **167**, E102–E116.

Berner, D., Stutz, W. E. and Bolnick, D. I. (2010). Foraging trait (co)variances in stickleback

evolve deterministically and do not predict trajectories of adaptive diversification. *Evolution*

(N. Y). **64**, 2265–2277.

Blankers, T., Hennig, R. M. and Gray, D. A. (2015). Conservation of multivariate female

preference functions and preference mechanisms in three species of trilling field crickets. *J.*

Evol. Biol. **28**, 630–641.

Blankers, T., Gray, D. A. and Hennig, R. M. (2017). Multivariate Phenotypic Evolution:

Divergent Acoustic Signals and Sexual Selection in Gryllus Field Crickets. *Evol. Biol.* **44**,

43–55.

Blankers, T., Lievers, R., Plata, C., van Wijk, M., van Veldhuizen, D. and Groot, A. T.

(2021). Sex pheromone signal and stability covary with fitness. *R. Soc. Open Sci.* **8**,

210180.

Brooks, R., Hunt, J., Blows, M. W., Smith, M. J., Bussière, L. F. and Jennions, M. D.
(2005). Experimental Evidence for Multivariate Stabilizing Selection. *Evolution (N. Y).* **59**,
871–880.

Bulmer, M. G. (1971). The Effect of Selection on Genetic Variability. *Am. Nat.* **105**, 201–211.

Candolin, U. (2003). The use of multiple cues in mate choice. *Biol. Rev.* **78**, 575–595.

Cardé, R. T. and Haynes, K. F. (2004). Structure of the pheromone communication channel in
moths. In *Advances in insect chemical ecology* (ed. Cardé, R. T.) and Millar, J. G.), pp.
283–332. Cambridge University Press Cambridge.

Careau, V., Wolak, M. E., Carter, P. A. and Garland, T. (2015). Evolution of the additive
genetic variance–covariance matrix under continuous directional selection on a complex
behavioural phenotype. *Proc. R. Soc. B Biol. Sci.* **282**,.

Carson, H. L. (1990). Increased genetic variance after a population bottleneck. *Trends Ecol.*
Evol. **5**, 228–230.

Chenoweth, S. F. and Blows, M. W. (2006). Dissecting the complex genetic basis of mate
choice. *Nat. Rev. Genet.* **7**, 681–692.

Cheverud, J. M. (1996). Developmental Integration and the Evolution of Pleiotropy. *Integr.*
Comp. Biol. **36**, 44–50.

Cole, G. L. and Endler, J. A. (2015). Variable environmental effects on a multicomponent
sexually selected trait. *Am. Nat.* **185**, 452–468.

Coyne, J. A. and Orr, A. H. (2004). *Speciation*. Sinauer Associates, Sunderland, MA.

Devigili, A., Evans, J. P., Di Nisio, A. and Pilastro, A. (2015). Multivariate selection drives
concordant patterns of pre- and postcopulatory sexual selection in a livebearing fish. *Nat.*
Commun. **6**, 8291.

Du, M., Bernstein, R., Hoppe, A. and Bienefeld, K. (2021). Short-term effects of controlled
mating and selection on the genetic variance of honeybee populations. *Heredity (Edinb).*
733–747.

Eroukhmanoff, F. (2009). Just how much is the G-matrix actually constraining adaptation?
Evol. Biol. **36**, 323–326.

Estes, S. and Arnold, S. J. (2007). Resolving the paradox of stasis: models with stabilizing
selection explain evolutionary divergence on all timescales. *Am. Nat.* **169**, 227–244.

Ferveur, J. F. (2005). Cuticular hydrocarbons: Their evolution and roles in Drosophila
pheromonal communication. *Behav. Genet.* **35**, 279–295.

Fisher, H. S., Mascuch, S. J. and Rosenthal, G. G. (2009). Multivariate male traits misalign
with multivariate female preferences in the swordtail fish , Xiphophorus birchmanni. *Anim.*
Behav. **78**, 265–269.

Garant, D., Hadfield, J. D., Kruuk, L. E. B. and Sheldon, B. C. (2008). Stability of genetic
variance and covariance for reproductive characters in the face of climate change in a wild
bird population. *Mol. Ecol.* **17**, 179–188.

Gerhardt, H. C. and Brooks, R. (2009). Experimental analysis of multivariate female choice in
gray treefrogs (Hyla versicolor): Evidence for directional and stabilizing selection.
Evolution (N. Y). **63**, 2504–2512.

Gosden, T. P. and Chenoweth, S. F. (2014). The evolutionary stability of cross-sex, cross-trait
genetic covariances. *Evolution (N. Y).* **68**, 1687–1697.

Grether, G. F., Kolluru, G. R. and Nersissian, K. (2004). Individual colour patches as
multicomponent signals. *Biol. Rev. Camb. Philos. Soc.* **79**, 583–610.

Groot, A. T., Fan, Y., Brownie, C., Jurenka, R. A., Gould, F. and Schal, C. (2005). Effect of
PBAN on pheromone production by mated Heliothis virescens and Heliothis subflexa
females. *J. Chem. Ecol.* **31**, 15–28.

Groot, A. T., Horovitz, J. L., Hamilton, J., Santangelo, R. G., Schal, C. and Gould, F.
(2006). Experimental evidence for interspecific directional selection on moth pheromone
communication. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 5858–5863.

Groot, A. T., Santangelo, R. G., Ricci, E., Brownie, C., Gould, F. and Schal, C. (2007).
Differential Attraction of Heliothis subflexa Males to Synthetic Pheromone Lures in Eastern

US and Western Mexico. *J. Chem. Ecol.* **33**, 353–368.

Groot, A. T., Inglis, O., Bowdridge, S., Santangelo, R. G., Blanco, C., López, J. D., Vargas, A. T., Gould, F. and Schal, C. (2009a). Geographic and temporal variation in moth chemical communication. *Evolution (N. Y).* **63**, 1987–2003.

Groot, A. T., Estock, M. L., Horovitz, J. L., Hamilton, J., Santangelo, R. G., Schal, C. and Gould, F. (2009b). QTL analysis of sex pheromone blend differences between two closely related moths: insights into divergence in biosynthetic pathways. *Insect Biochem. Mol. Biol.* **39**, 568–577.

Groot, A. T., Blanco, C. A., Claβen, A., Inglis, O., Santangelo, R. G., Lopez, J., Heckel, D. G. and Schal, C. (2010). Variation in Sexual Communication of the Tobacco Budworm, *Heliothis virescens*. *Southwest. Entomol.* **35**, 367–372.

Groot, A. T., Staudacher, H., Barthel, A., Inglis, O., Schöfl, G., Santangelo, R. G., Gebauer-Jung, S., Vogel, H., Emerson, J., Schal, C., et al. (2013). One quantitative trait locus for intra- and interspecific variation in a sex pheromone. *Mol. Ecol.* **22**, 1065–1080.

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. *J. Stat. Softw.* **33**, 1–22.

Hadfield, J. D. (2012). MCMCglmm Course Notes.

Heath, R. R., Mitchell, E. R. and Cibrian Tovar, J. (1990). Effect of release rate and ratio of (Z)-11-hexadecen-1-ol from synthetic pheromone blends on trap capture ofHeliothis subflexa (Lepidoptera: Noctuidae). *J. Chem. Ecol.* **16**, 1259–1268.

Higham, J. P. and Hebets, E. A. (2013). An introduction to multimodal communication. *Behav. Ecol. Sociobiol.* **67**, 1381–1388.

Hill, W. G. and Caballero, A. (1992). Artificial selection experiments. *Annu. Rev. Ecol. Syst.* **23**, 287–310.

Hine, E., McGuigan, K. and Blows, M. W. (2011). Natural selection stops the evolution of male attractiveness. *Proc. Natl. Acad. Sci.* **108**, 3659–64.

Johnson, T. and Barton, N. (2005). Theoretical models of selection and mutation on

589  quantitative traits. *Philos. Trans. R. Soc. B Biol. Sci.* **360**, 1411–1425.

590 **Jones, A. G., Arnold, S. J. and Bürger, R.** (2003). Stability of the g -matrix in a population

591  experiencing pleiotropic mutation, stabilizing selection, and genetic drift. *Evolution (N. Y).*

592  **57**, 1747–1760.

593 **Kárpáti, Z., Tasin, M., Cardé, R. T. and Dekker, T.** (2013). Early quality assessment lessens

594  pheromone specificity in a moth. *Proc. Natl. Acad. Sci.* **110**, 7377 LP – 7382.

595 **Lande, R.** (1979). Quantitative genetic analysis of multivariate evolution, applied to brain: body

596  size allometry. *Evolution* **33**, 402–416.

597 **Lande, R. and Arnold, S. J.** (1983). The measurement of selection on correlated characters.

598  *Evolution (N. Y).* **37**, 1210–1226.

599 **Linn, C. E. J., Campbell, M. G. and Roelofs, W. L.** (1987). Pheromone components and active

600  spaces: what do moths smell and where do they smell it? *Science* **237**, 650–652.

601 **Linn, C. E., Young, M. S., Gendle, M., Glover, T. J. and Roelofs, W. L.** (1997). Sex

602  pheromone blend discrimination in two races and hybrids of the European corn borer moth,

603  Ostrinia nubilalis. *Physiol. Entomol.* **22**, 212–223.

604 **Löfstedt, C.** (1990). Population variation and genetic control of pheromone communication

605  systems in moths. *Entomol. Exp. Appl.* **54**, 199–218.

606 **Lynch, M. and Walsh, B.** (1998). *Genetics and analysis of quantitative Traits*. Sinauer,

607  Sunderland, MA.

608 **Melo, D. and Marroig, G.** (2014). Directional selection can drive the evolution of modularity in

609  complex traits. *Proc. Natl. Acad. Sci.* **112**,.

610 **Oh, K. P. and Shaw, K. L.** (2013). Multivariate sexual selection in a rapidly evolving speciation

611  phenotype. *Proc. R. Soc. B-Biological Sci.* **280**, 20130482.

612 **R Core Team** (2019). R: A Language and Environment for Statistical Computing.

613 **Revell, L. J.** (2007). The G matrix under fluctuating correlational mutation and selection.

614  *Evolution (N. Y).* **61**, 1857–1872.

Ritchie, M. G. (2007). Sexual Selection and Speciation. *Annu. Rev. Ecol. Evol. Syst.* **38**, 79–102.

Ritchie, M. G. and Kyriacou, C. P. (1996). Artificial selection for a courtship signal in
Drosophila melanogaster. *Anim. Behav.* **52**, 603–611.

Roff, D. A. and Fairbairn, D. J. (2012). The evolution of trade-offs under directional and
correlational selection. *Evolution (N. Y).* **66**, 2461–2474.

Rowe, C. (1999). Receiver psychology and the evolution of multicomponent signals. *Anim.*
Behav. **58**, 921–931.

Ryan, M. J. and Rand, a S. (2003). Sexual selection in female perceptual space: how female
túngara frogs perceive and respond to complex population variation in acoustic mating
signals. *Evolution* **57**, 2608–2618.

Saltz, J. B., Hessel, F. C. and Kelly, M. W. (2017). Trait Correlations in the Genomics Era.
Trends Ecol. Evol. **32**, 279–290.

Schaefer, H. M. and Ruxton, G. D. (2015). Signal Diversity, Sexual Selection, and Speciation.
Annu. Rev. Ecol. Evol. Syst. **46**, 573–592.

Schneider, D. (1992). 100 years of pheromone research - An essay on lepidoptera.
Naturwissenschaften **79**, 241–250.

Svensson, E. I., Arnold, S. J., Bürger, R., Csilléry, K., Draghi, J., Henshaw, J. M., Jones, A.
G., De Lisle, S., Marques, D. A., McGuigan, K., et al. (2021). Correlational selection in
the age of genomics. *Nat. Ecol. Evol.* **5**,.

Tanner, J. C., Ward, J. L., Shaw, R. G. and Bee, M. A. (2017). Multivariate phenotypic
selection on a complex sexual signal. *Evolution (N. Y).* **71**, 1742–1754.

Templeton, A. R. (2008). The reality and importance of founder speciation in evolution.
BioEssays **30**, 470–479.

Uesugi, A., Connallon, T., Kessler, A. and Monro, K. (2017). Relaxation of herbivore-
mediated selection drives the evolution of genetic covariances between plant competitive
and defense traits. *Evolution (N. Y).* **71**, 1700–1709.

Vetter, R. S. and Baker, T. C. (1983). Behavioral responses of maleHeliothis virescens in a

sustained-flight tunnel to combinations of seven compounds identified from female sex
pheromone glands. *J. Chem. Ecol.* **9**, 747–759.

Vickers, N. J. (2002). Defining a synthetic pheromone blend attractive to male Heliothis
subflexa under wind tunnel conditions. *J. Chem. Ecol.* **28**, 1255–1267.

Vickers, N. J. and Baker, T. C. (1997). Flight of Heliothis virescens males in the field in
response to sex pheromone. *Physiol. Entomol.* **22**, 277–285.

Villemereuil, P. de (2012). Animal Model Tutorial.

West-Eberhard, M. J. (2014). Darwin's forgotten idea: The social essence of sexual selection.
Neurosci. Biobehav. Rev. **46**, 501–508.

Wiens, J. J. and Tuschhoff, E. (2020). Songs versus colours versus horns: what explains the
diversity of sexually selected traits? *Biol. Rev.* **95**, 847–864.

Wilkins, M. R., Seddon, N. and Safran, R. J. (2013). Evolutionary divergence in acoustic
signals : causes and consequences. *Trends Ecol. Evol.* **28**, 156–166.

Wilkins, M. R., Shizuka, D., Joseph, M. B., Hubbard, J. K. and Safran, R. J. (2015).
Multimodal signalling in the North American barn swallow: A phenotype network
approach. *Proc. R. Soc. B Biol. Sci.* **282**,.

Wolf, J. B., Leamy, L. J., Routman, E. J. and Cheverud, J. M. (2005). Epistatic pleiotropy
and the genetic architecture of covariation within early and late-developing skull trait
complexes in mice. *Genetics* **171**, 683–694.

Zhu, J., Chastain, B. B., Spohn, B. G. and Haynes, K. F. (1997). Assortative mating in two
pheromone strains of the cabbage looper moth, Trichoplusia ni. *J. Insect Behav.* **10**, 805–
817.

664