High-throughput small molecule screen identifies inhibitors of microsporidia invasion and

proliferation in *C. elegans*.

Brandon M. Murareanu[1], Noelle V. Antao[2], Winnie Zhao[1], Aurore Dubuffet[4], Hicham El Alaoui[4], Jessica

Knox[1,3], Damian C. Ekiert[2,5], Gira Bhabha[2], Peter J. Roy[1,3], & Aaron W. Reinke[1#].

[1]Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

[2]Department of Cell Biology, New York University School of Medicine, New York, New York, United States

of America

[3]The Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto

[4]Université Clermont Auvergne, CNRS, Laboratoire Microorganismes: Génome et Environnement, F-

63000 Clermont-Ferrand, France

[5]Department of Microbiology, New York University School of Medicine, New York, New York, United

States of America

#Corresponding author

aaron.reinke@utoronto.ca

Abstract

Microsporidia are a diverse group of fungal-related obligate intracellular parasites that infect most animal

phyla. Despite the emerging threat that microsporidia have become to humans and agricultural animals,

few reliable treatment options exist. To identify novel chemical inhibitors of microsporidia infection, we

developed a high-throughput screening method using *Caenorhabditis elegans* and the microsporidia

species *Nematocida parisii*. We screened the Spectrum Collection of 2,560 FDA-approved compounds

and natural products to identify compounds that prevent *C. elegans* progeny inhibition caused by *N.*

parisii infection. We developed a semi-automated method for quantifying *C. elegans* progeny number in

liquid culture, confirming 11 candidate microsporidia inhibitors. We show that five compounds prevent

microsporidia infection by inhibiting spore firing, and demonstrate that one compound, dexrazoxane,

slows infection progression. We also show that these compounds have activity against several other microsporidia species, including those which infect humans. Together, our results demonstrate the effectiveness of *C. elegans* as a model host for drug discovery against intracellular pathogens and provide a scalable high-throughput system for the identification and characterization of additional microsporidia inhibitors.

Introduction

Microsporidia are a diverse group of parasites, comprising over 1400 species that can infect most major animal phyla (1–3). Many microsporidia species infect agriculturally important animals. This includes invertebrate-infecting species such as *Nosema ceranae* and *Nosema apis* which infect honey bees, *Enterocytozoon hepatopenaei* which infects shrimp, *Hepatospora eriocheir* which infects crabs, and *Nosema bombycis* which infects silkworms (4–7). Additionally, there are several species that infect farmed fish, including *Loma salmonae* which infects salmon and trout, as well as *Enterospora nucleophila* which infects gilthead sea bream (8, 9). Microsporidia infection in these animals can result in reduced body size, fewer offspring, and increased mortality (4–9). The economic impact of microsporidia infections is high. The infection of silkworms has triggered historical collapses of the sericulture industry, and microsporidia infections are estimated to cost over $200 million USD annually to Thailand's shrimp industry (7, 10). Livestock such as pigs, cattle, and sheep, as well as pets such as dogs, cats and rabbits are infected by *Encephalitozoon* species and *Enterocytozoon bieneusi*. Humans are also infected by these species, with infections in the immunocompromised being more prevalent (11). Microsporidia commonly infect animals, with over half of honeybee hives and approximately 40% of pigs reported to be infected (11, 12). Microsporidia are also emerging pathogens, with the threat posed by many of these species only being recognized in the last several decades (13).

Despite the threat that these parasites pose to human and animal health, few therapeutic options exist. Fumagillin is a compound from the fungus *Aspergillus fumigatus* which inhibits methionine aminopeptidase type 2 (MetAP2), and has been used since the 1950s to treat microsporidia infections in

honeybees (14). However, recent reports suggest that fumagillin may be ineffective against *N. ceranae* in honeybees and *E. hepatopenaei* in shrimp (15, 16). In addition, fumagillin causes toxicity in humans and its use in beekeeping is banned in some countries (17). The other most used therapeutic agent against microsporidia is albendazole, which disrupts β-tubulin function. Several microsporidia species, including *E. bieneusi,* have β-tubulin variants associated with albendazole resistance, and as a result, these species are not susceptible to the drug (18). Several other approaches for the drug treatment of microsporidia infections have been described, including inhibition of chitin synthase, as well as inhibition of spore firing by blocking calcium channels (18, 19). Microsporidia only grow inside of host cells, making screening for inhibitors challenging. Several screens to identify microsporidia inhibitors have been performed, but these have been limited to less than 100 compounds at a time due to a lack of applicable high-throughput screening assays (20, 21).

The model organism *Caenorhabditis elegans* is a powerful system to study infectious diseases and is widely amenable to high-throughput drug screens. *C. elegans* is commonly infected in nature by the microsporidian *Nematocida parisii,* which has been used as model to study microsporidia spore exit, host immunity, and proteins used by microsporidia to interact with its host (22–26). The infection of *C. elegans* begins when *N. parisii* spores are ingested. Spores then germinate in the intestinal lumen, where their unique invasion apparatus known as a polar tube is fired, enabling sporoplasm deposition into intestinal epithelial cells. Intracellularly, the sporoplasm initiates a proliferative process of multiplication by binary or multiple fission, known as merogony, producing meronts in direct contact with the host cytosol (27, 28). Following proliferation, meronts undergo sporogony to form spores which then exit the worm, with over 100,000 spores being produced by each infected animal (29). Infection of *C. elegans* by *N. parisii* results in impaired growth, reduced progeny production, and lethality (23, 27). The ease of culturing and growing large numbers of animals, along with easily discernible phenotypes, has made *C. elegans* a powerful platform to identify novel anthelmintic, antibiotic, and antifungal agents (30–37).

To discover additional microsporidia inhibitors, we used the *C. elegans / N. parisii* model system to

develop a high-throughput, liquid-based drug screening assay in which compounds were scored for their

ability to prevent infection-induced progeny inhibition. Using this assay, we screened the Spectrum

Collection of 2,560 FDA-approved compounds and natural products, and identified 11 chemical inhibitors

of microsporidia infection, which we confirmed using a semi-automated method for quantifying *C. elegans*

progeny number in liquid culture. We report that five of these compounds, including the known serine

protease inhibitor ZPCK and four compounds that share a quinone structure, prevent microsporidia

invasion in *C. elegans* by inhibiting spore firing. Additionally, the iron chelator and topoisomerase II

inhibitor dexrazoxane prevents infection progression. Together, this work describes methods to screen

thousands of putative microsporidia inhibitors and identifies novel microsporidia inhibitors that either

block microsporidia invasion or proliferation.

Results

High-throughput screen of 2,560 Spectrum Collection compounds reveals 11 microsporidia

inhibitors

To identify inhibitors of microsporidia infection, we developed a novel screening assay for chemical

inhibitors based on the observation that *C. elegans* progeny production is greatly inhibited when infected

with *N. parisii* (23, 38). In our screening assay, L1 animals in 96-well plates were grown in liquid and

infected with *N. parisii* spores for six days at 21°C (Fig. 1A, see methods). In the absence of *N. parisii*

spores, *C. elegans* larvae develop into adults and produce progeny. In the presence of spores, the

animals are smaller and produce fewer progeny, providing a convenient visual indication of infection. This

inhibition of progeny production is prevented by the known microsporidia inhibitor fumagillin (Fig. 1B).

Using this assay, we tested 2,560 FDA-approved compounds and natural products from the Spectrum

Collection for their ability to prevent infection-induced progeny inhibition. After incubation with compounds

for six days, each well was visually assessed for *C. elegans* progeny production. We identified 25 initial

compounds that when added to wells, resulted in the production of more progeny than the vehicle, DMSO,

controls. Upon retesting, 11 compounds were confirmed to reproducibly restore *C. elegans* progeny

production in the presence of spores (Table S1).

To quantify the inhibitory effect of each compound, we developed a semi-automated approach to quantify

progeny number in liquid culture, (Fig. 1A, see methods). Animals were grown in wells as described

above, stained with the dye rose bengal, and imaged using a flatbed scanner. Images were processed

with consistent parameters to highlight stained animals and analyzed with WorMachine to detect and

count the number of animals in each well (Fig. S1A-C) (39). Counts of animals detected using

WorMachine correlated well with those counted manually (Fig. S1D). Additionally, there was good

correlation (average R^2 of 0.71) between technical replicates (Fig. S1E). This approach is similar to a

recently published method, but with the advantage of using a relatively cheap flatbed scanner for imaging

(40–42). We observed that each of the 11 compounds was able to significantly increase the number of

progeny produced by animals under infection conditions (Fig. 1C). This effect is even more pronounced

when considering that for six of the compounds, there was a significant reduction in progeny production

in uninfected animals, indicating moderate host toxicity (Fig. 1D).

Our initial screen identified compounds that could rescue the ability of *C. elegans* to produce progeny in

the presence of *N. parisii* spores. To determine whether the compounds have a direct effect on

microsporidia infection, we performed continuous infection assays. L1 animals were infected continuously

with *N. parisii* spores in the presence of compounds for 4 days at 21°C in 24-well plates in liquid (Fig.

2A). After incubation, animals were fixed and stained with direct yellow 96 (DY96), a fluorescent dye that

binds chitin, a crucial component of both the microsporidia spore wall and *C. elegans* embryos (Fig. 2B)

(43, 44). First, we observed that every compound significantly increased the proportion of adult animals

containing embryos in the presence of *N. parisii* spores (Fig. 2C). These results are consistent with our

data from the initial screen showing that all compounds increased progeny production in the presence of

spores (Fig. 1C). Second, we determined that the control inhibitor fumagillin and 9 of the 11 compounds

inhibited microsporidia infection, as the proportion of animals with newly formed spores was significantly

lower upon treatment with these compounds (Fig. 2D).

N. parisii proliferation is inhibited by dexrazoxane

Microsporidia infection can be inhibited either by blocking invasion, or by preventing proliferation after

infection is established. To distinguish between these possibilities, we performed pulse-chase infection

assays where L1 animals were infected with _N. parisii_ spores for 3 hours, washed to remove excess

spores, and incubated with compounds for 4 days (Fig. 3A). As expected, given its described mode of

action as a MetAP2 inhibitor (17), fumagillin restricted _N. parisii_ proliferation (Fig 3B). Surprisingly, of our

11 compounds, only one, dexrazoxane, inhibited _N. parisii_ proliferation (Fig. 3E). Both fumagillin and

dexrazoxane restored the ability of animals to make embryos when infected, whereas none of the other

compounds were able to do so (Fig. 3D). The effect of dexrazoxane, at a concentration of 60 µM, is

especially striking with a ~1200-fold reduction in the proportion of animals with newly-formed spores,

compared to just ~2-fold for fumagillin, at a concentration of 350 µM (Fig. 3E). At this concentration of

dexrazoxane, we observed no negative effect on uninfected animals (Fig. 1D). Together, these results

suggest that of the compounds identified from our screen, only dexrazoxane inhibits _N. parisii_ after

invasion of _C. elegans_ has occurred.

To determine whether dexrazoxane inhibits microsporidia by slowing proliferation or enhancing parasite

clearance, we used FISH staining of pulse-chase infected animals with probes specific for _N. parisii_ 18S

RNA to visualize the sporoplasms and meronts of the earlier stages of infection prior to new spore

formation (27). Although fumagillin and dexrazoxane both resulted in many fewer animals with newly

formed spores (Fig. 3E), there was no significant difference in the proportion of animals that had at least

some FISH signal (Fig. 3F). When _C. elegans_ is pulse-chase infected with _N. parisii_ for 48 h, many large,

multinucleated meronts are observed. Worms treated with fumagillin contain both large meronts with

many nuclei as well as some parasite cells that fail to progress beyond having 2 nuclei. In contrast,

dexrazoxane treatment leads to almost all parasite cells containing 1 or 2 nuclei, and many parasite cells

displaying irregular morphology (Fig. 3C). Using images of FISH-stained *N. parisii*, we quantified the area

of the infected animal that is covered by sporoplasms and meronts. We observed that fumagillin and

dexrazoxane treatment resulted in a significantly reduced pathogen load, with dexrazoxane having the

strongest effect (Fig. 3G). Together, these results indicate that dexrazoxane can greatly inhibit *N. parisii*

proliferation but does not cause the infection to be cleared.

Dexrazoxane is an iron chelator, and iron levels have been shown to impact the growth of other

microsporidia species (45, 46). To determine if iron levels are important for *N. parisii* growth in *C. elegans*,

we supplemented our liquid cultures with ferric ammonium citrate (FAC) as a water-soluble iron source

(47), and continuously infected L1 animals with a low dose of spores for 4 days. While the proportion of

animals with newly formed spores was slightly higher in the supplemented condition, this effect was not

statistically significant (Fig. S2A). In contrast, addition of FAC resulted in a small, but significant increase

in the proportion of animals containing embryos (Fig. S2B). To determine whether dexrazoxane is likely

acting as an iron chelator in our system, we tested another iron chelator, 2,2'-bipyridyl (BP), for anti-

microsporidial activity in our continuous infection assay, but did not observe an effect (Fig. 3H and S2C).

In addition, FAC supplementation was unable to counteract the anti-microsporidial effect of dexrazoxane

(Fig. 3H and S2C). We also tested the *smf-3(ok1035)* mutant strain RB1074 for sensitivity to

dexrazoxane. RB1074 has ~50% less iron levels compared to the wild-type N2 strain and displays a

striking growth defect upon treatment with the iron chelator BP (47). However, RB1074 did not display

any such growth impairment upon treatment with dexrazoxane (Fig. S3). Taken together, these results

demonstrate that the inhibitory effect of dexrazoxane on *N. parisii* infection is likely independent of its

established function as an iron chelator.

Protease inhibitors and quinone derivatives prevent invasion by inhibiting spore firing

Compounds that displayed strong activity against microsporidia infection in the continuous infection

assays, but not the pulse-chase infections, may be preventing invasion by inhibiting spore firing. In *C.*

elegans, *N. parisii* spores are ingested, and spore firing is triggered in the intestinal lumen (27, 48). To

test if any compounds interfere with this process, we performed spore firing assays on the 7 compounds

with the strongest effect in the continuous infection assays (Fig. 2D). L1 stage *C. elegans* were infected

with *N. parisii* for 3 hours, and then stained with FISH and DY96 (Fig. 4A). This dual staining enables us

to determine whether a given spore in the intestine has fired, based on whether it has released its

sporoplasm (Fig. 4C). We found that treatment with five compounds (ZPCK, menadione, plumbagin,

thymoquinone and dalbergione) resulted in a significantly reduced proportion of fired spores in the

intestine (Fig. 4D). Treatment with these compounds, as well as chloranil, also lead to a notable reduction

in the number of sporoplasms per animal (Fig. 4E). The two compounds that inhibit microsporidia

proliferation, fumagillin and dexrazoxane, had no significant effect on spore firing, although fumagillin

treatment did result in a significant decrease in sporoplasm numbers (Fig. 4D and E).

ZPCK is an irreversible inhibitor of serine proteases, and a microsporidia serine protease has previously

been suggested to be involved in spore firing (49, 50). To determine if other serine protease inhibitors

can prevent *N. parisii* invasion, we tested three additional small molecules (TPCK, TLCK and PMSF) and

three peptides (Antipain, Chymostatin and Leupeptin) known to inhibit serine proteases in a modified

version of our spore firing assay. For these experiments, we incubated spores with serine protease

inhibitors for 24 hours to accentuate their effects, and then used the spores to infect L1 stage *C. elegans*

for 3 hours in the presence of inhibitors (Fig. 4B). Analysis of the proportion of spores fired and the

number of sporoplasms per animal revealed that both ZPCK and the structurally related inhibitor TPCK

displayed strong inhibition of *N. parisii* spore firing and invasion (Fig. 4F and G).

Inhibitors of spore firing could function either by inhibiting an *N. parisii* spore protein or a *C. elegans*

protein necessary for firing and invasion. To distinguish between these possibilities, we incubated spores

for 24 hours with each of the compounds that inhibited spore firing, washed away excess inhibitor, and

then used the treated spores to infect L1 stage *C. elegans* for 3 hours in the absence of inhibitors (Fig.

4B). Treatment with all 6 inhibitors (ZPCK, TPCK, menadione, plumbagin, thymoquinone and

dalbergione) significantly inhibited spore firing and sporoplasm invasion (Fig. 4H and I). These results

demonstrate that all of these inhibitors act directly on *N. parisii* spores to prevent firing and invasion, and

in the case of ZPCK and TPCK, this likely occurs through the inhibition of a serine protease.

Identified compounds inhibit multiple microsporidia species

We next determined if the compounds we identified could inhibit infection by other microsporidia species.

In addition to being infected by *N. parisii*, *C. elegans* is also infected by *Pancytospora epiphaga*. This

species infects the epidermis of *C. elegans* and belongs to the Enterocytozoonida clade which includes

the human infecting species *Vittaformae cornea* and *Enterocytozoon bieneusi* (22, 51). To determine if

dexrazoxane could inhibit *P. epiphaga* proliferation, we infected *C. elegans* with this species and

monitored parasite growth using FISH staining (Fig. 5A) When animals were treated with 350 μM

fumagillin or 60 μM dexrazoxane we observed significantly less parasite than in untreated control

animals, with the least parasite growth occurring in the animals treated with dexrazoxane (Fig. 5B).

Anncaliia algerae infects both mosquitos and humans and belongs to the Neopereziida clade of

microsporidia (51, 52). To test whether dexrazoxane could also prevent inhibition of this species, we

infected human fibroblast cells with *A. algerae* and visualized infection using FISH (Fig. 5C). We observed

a significant dose dependent response of inhibition of parasite division (Fig. 5D). To test whether

dexrazoxane was toxic to cells at the concentrations that prevented *A. algerae* proliferation, we monitored

cell viability and observed no significant toxicity in the concentrations we tested (Fig. 5E).

Encephalitozoon intestinalis infects humans and other mammals and belongs to the Nosematida clade

of microsporidia (51, 53). To determine if the compounds we identified could block spore firing in this

species we pre-treated *E. intestinalis* spores with menadione, plumbagin, thymoquinone, or ZPCK, and

then carried out an in vitro spore germination assay, in which we can directly observe polar tube firing.

We observed a significant decrease in the frequency of germination when spores were pre-treated with

each of these compounds (Fig. 5F). To test whether these compounds impact host cell infection, we

incubated Vero cells with spores that were either untreated or pre-treated with each compound, and

monitored infection with FISH (Fig. 5G). We observed that cells remained largely uninfected when

incubated with spores that had been pre-treated with the compounds, suggesting that all these inhibitors

also prevent invasion (Fig. 5H). In the presence of ZPCK, we identified a small number of single invasion

events (Fig. 5G), but no replication of parasites. Together our results indicate that a diverse set of

microsporidia species can be inhibited by the compounds we identified.

Discussion

To identify inhibitors of microsporidia, we screened 2,560 compounds for their ability to counteract

infection-induced *C. elegans* progeny inhibition. We identified 11 compounds with reproducible inhibitory

activity (Table S1). We found that dexrazoxane inhibits *N. parisii* proliferation, whereas the protease

inhibitors ZPCK and TLCK, along with 4 quinone derivatives, prevent spore firing and invasion (Fig. 6).

Three of the identified inhibitors, curcumin, plumbagin, and thymoquinone, have been previously shown

to inhibit microsporidia infection, although in the case of plumbagin and thymoquinone, the compounds

were effective after microsporidia had infected cells (20, 21). This contrasts with our results, where these

compounds only prevent microsporidia invasion. To our knowledge none of the other compounds we

identified have been previously reported to prevent microsporidia infection, demonstrating the value in

this unbiased approach to identify novel microsporidia inhibitors.

Dexrazoxane is an FDA-approved drug for the prevention of cardiomyopathies caused by

chemotherapeutic drugs in cancer patients (54). Dexrazoxane can be hydrolyzed into a structure that is

similar to the metal ion chelator ethylenediaminetetraacetic acid and is thought to work by reducing the

amount of iron complexed with anthracycline chemotherapeutics, thus reducing the number of superoxide

radicals formed by such interactions (55, 56). Iron is an important metabolite for microsporidia

proliferation. Honey bees infected with *N. ceranae* have reduced levels of iron and the growth of *E.*

cuniculi in macrophages was promoted with the addition of iron and inhibited upon addition of an iron

chelator (45, 46). We found that dexrazoxane could prevent microsporidia proliferation, at a concentration

that has no significant effect on host fitness. Given the established role of dexrazoxane as an iron

chelator, it seemed likely that its mechanism as an anti-microsporidial might involve sequestering iron away from sporoplasms and meronts, thus depriving them of a crucial resource required for growth and infection progression. However, our results do not support this. Neither treatment with the iron chelator BP, nor iron supplementation using FAC, had any significant effect on *N. parisii* infection. Additionally, dexrazoxane still exhibited strong anti-microsporidial effects when *C. elegans* were exposed to *N. parisii* in a high iron environment. These results suggest that the anti-microsporidial properties of dexrazoxane are unlikely to be the result of drug-induced alterations in iron homeostasis. This is contrast to lethality of *C. elegans* caused by the pathogenic bacteria *Pseudomonas aeruginosa*, which can be reversed by treatment with the iron chelator ciclopirox olamine, which was included in our screen, but was not observed to have an effect against *N. parisii* (57). The iron chelator deferoxamine, which inhibits *E. cuniculi* in macrophages, was also included in our screen, but not observed to have activity against *N. parisii* (46). A second proposed mechanism for dexrazoxane is the inhibition of topoisomerase II (58). However, several topoisomerase II inhibitors that have previously been used with *C. elegans* (etoposide, teniposide, dactinomycin, and amsacrine), were included in our initial screen, but were not observed to have activity against *N. parisii* (59). The mechanism of action of dexrazoxane in blocking *N. parisii* proliferation thus remains elusive.

The proteins that control spore firing in microsporidia are unknown (60, 61). Although the *in vivo* signal for firing is undetermined, several different pH and salt conditions can trigger firing *in vitro* (60). Once microsporidia spores are exposed to a host signal, the osmotic pressure of the spore increases forcing the polar tube to be extruded. One protein hypothesized to control spore firing is the subtilisin serine protease which is conserved throughout microsporidia (61). This protein localizes to the poles of *N. bombycis* spores, and the activated version of the protease localizes to the end of the spore from which the polar tube fires (62). Here we show that two irreversible serine protease inhibitors (ZPCK and TPCK) can block firing and prevent invasion. Additionally, we identified four quinone derivatives (menadione, plumbagin, thymoquinone and dalbergione) that also block firing and invasion. For all six of these spore firing inhibitors, we observed that firing is still blocked even when the compounds are washed away

following their use in the pre-treatment of spores. These results demonstrate that the inhibitors likely act directly on the spores to irreversibly block firing, even when the trigger for firing occurs inside of the host animal. As spore firing is the initial step in invasion, blocking spore firing is a promising potential strategy to prevent microsporidia infections in agricultural animals.

The *C. elegans* / *N. parisii* system provides a powerful tool to both uncover and characterize novel microsporidia inhibitors. Our approach is high-throughput, relatively cheap and scalable, allowing the screening of even more diverse collections of small-molecule libraries (63). Our screen is advantageous as it is performed with wild-type animals and pathogens, and by screening based on host fitness, this approach only identifies compounds with minimal host toxicity (35). *C. elegans* is also infected by *P. epiphaga*, providing the opportunity to test the specificity of compound inhibition on multiple species of microsporidia in the same host (22). Several of the compounds we identified also inhibited other microsporidia species, demonstrating the power of using *N. parisii* to discovery novel microsporidia inhibitors. As evidenced by our results, *C. elegans* is a very useful host for determining at what stages of microsporidia infection the identified inhibitors are acting (23). Several strategies to identify inhibitor targets have been successfully used with other eukaryotic parasites, including generating and sequencing parasite variants that are resistant to inhibition or through thermal proteome profiling (64– 66). *C. elegans* is also likely to be useful for future work to identify inhibitor targets in *N. parisii* using these approaches.

Acknowledgements

We thank Hala Tamim El Jarkass and Alexandra R. Willis for providing helpful comments on the manuscript and Hala Tamim El Jarkass for the illustration in figure 6. This work was supported by the Natural Sciences and Engineering Research Council of Canada Undergraduate Student Research Awards (to B.M.), Canadian Institutes of Health Research grant no. 400784 (to A.R.), 173448 (to P.R) , and 313296 (to P.R.), Centre for Collaborative Drug Research pilot project fund (to A. R.), Connaught New Researcher Award (to A.R.), an Alfred P. Sloan Research Fellowship FG2019-12040 (to A.R.), a

Canada Research Chair award (to P.R.), Pew Biomedical Scholars, PEW-00033055 (to G.B.), Searle Scholars Program, SSP-2018-2737 (to G.B.), NIAID R01AI147131 (to G.B.), NIGMS R35GM128777 (to D.C.E.). Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) and we thank WormBase.

Author contributions: B.M., J. K., A.R., and P.R. designed the inhibitor screen. B.M. performed all the experiments with *N. parisii*. W. Z. and A. R. designed the *P. epiphaga* experiments with W. Z. carrying out the experiments. A. D. and H. E. A. designed and carried out the *A. algerae* experiments. N. A., D. E, and G.B. designed the *E. intestinalis* experiments with N. A. carrying out the experiments. B.M. and A.R. analyzed the data and co-wrote the paper with edits from all of the authors.

Competing Interests: The authors declare that they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Methods

Chemical sources

The Spectrum Collection of FDA-approved compounds and natural products was obtained from MicroSource. For retesting and post-screen analyses, individual compounds were obtained from Sigma-Aldrich and MicroSource.

***C. elegans* strains, maintenance and bleach-synchronization**

Animals were cultured as previously described (67). All experiments were performed with the wild-type N2 strain of *Caenorhabditis elegans* unless otherwise indicated. The *smf-3(ok1035)* mutant strain RB1074 was obtained from the CGC. To generate bleach-synchronized L1 stage animals, L4 animals were first picked onto 10 cm NGM plates and grown for ~96 hours at 21°C. Worms were then washed off

plates with M9, and treated with ~4% NaClO / 1 M NaOH solution for 1 – 3 minutes to extract embryos,

then washed with M9. Embryos were incubated in M9 for 18 – 24 hours at 21°C to allow hatching.

***N. parisii* strains and spore preparation**

Nematocida parisii spores were prepared as previously described (23). All experiments were performed

with *N. parisii* strain ERTm1.

96-well plate-based screen for microsporidia inhibitors

C. elegans liquid culture methods and 96-well plate-based screening methods were partially adapted

from established protocols (32, 68). *N. parisii* spores were prepared as described above, mixed in K-

medium (51 mM NaCl, 32 mM KCl, 3 mM CaCl2, 3 mM MgSO4, 3.25 µM cholesterol) with 5x saturated

OP50-1 *E. coli*, and added to 96-well culture plate. 96-well Spectrum Collection plates were thawed from

–80°C. Using a V&P Scientific 96-well pinning tool, 300 nL of DMSO-dissolved compounds were pinned

into screening plate columns 2 – 11, and 300 nL of DMSO was pinned into screening plate columns 1

and 12 to be used for DMSO infected, DMSO uninfected, and fumagillin controls. Bleach-synchronized

L1s were prepared as described above, mixed in K-medium with 5x saturated OP50-1 *E. coli*, and added

to the screening plates. The final volume in each well was 50 µL, with ~100 L1s, *N. parisii* spores at a

final concentration of 6,250 spores/µL, DMSO at a final concentration of <1%, and compounds at a final

concentration of 60 µM except for fumagillin (used at a concentration of 350 µM in all experiments).

Screening plates were covered with adhesive porous film, placed in parafilm wrapped humidity boxes,

and incubated at 21°C, shaking at 180 rpm for 6 days. After incubation, progeny production was scored

manually by visual inspection of the screening plates. Initial hits were rescreened for reproducibility at a

final concentration of 60 µM in the same manner using individual compounds resuspended in DMSO.

The inhibitory effect of compounds that passed rescreening were quantified in separate experiments

using semi-automated methods described below.

Semi-automated quantification of progeny production

Bleach-synchronized L1s were treated with compounds +/- *N. parisii* spores in 96-well plates prepared and incubated as described above. Three wells were assayed for each condition for each biological replicate. After incubation, animals were stained by adding 10 µL of 0.3125 mg/mL Rose Bengal solution to each well using an Integra VIAFLO 96 Electronic pipette. Plates were wrapped in parafilm and incubated at 37°C for 16 – 24 hours. To each well, 240 µL of M9+0.1% Tween-20 was added. Plates were then centrifuged for 1 minute at 2200 x g. Next, 200 µL was removed and 150 µL of M9+0.1% Tween-20 added to each well. 25 µL from each well was then transferred to a white 96-well plate containing 300 µL M9 per well. Plates were imaged using an Epson Perfection V850 Pro flat-bed scanner with the following parameters (dpi = 4800, colour = 24-bit, .jpg compression = 1). Images were edited in GIMP version 2.10 or later to highlight stained animals by removing HTML colour codes #000000 and #FFC9AF, applying unsharp masking with the following parameters (radius = 10, effect = 10, threshold = 10), editing hue saturation with the following parameters (For yellow, green, blue and cyan: lightness = 100, saturation = -100. For red and magenta: lightness = -100, saturation = 100), and exporting each well as a single .tiff image with LZW compression (69). The number of animals in each well was counted using the MATLAB-based phenotypic analysis tool WorMachine with the following parameters (pixel neighbouring threshold = 1, pixel binarization threshold = 30, max object area to remove = 0.003%) (39).

Continuous infection assays

Continuous infection assays were performed in 24-well assay plates with each well containing a final volume 400 µL, ~800 L1s, 5,000 *N. parisii* spores/µL, 60 µM of each compound except for fumagillin (see above), and DMSO at a final concentration of 1%. Three wells were assayed for each compound for each of three biological replicates. Assay plates were covered with adhesive porous film, placed in parafilm wrapped humidity boxes, incubated at 21°C, 180 rpm for 4 days, and stored at 4°C, 20 rpm for 1 – 2 days. After incubation, samples were acetone-fixed, DY96-stained, and subject to fluorescence microscopy as described below.

Pulse infection assays

~8,000 bleach-synchronized L1s and 10 million *N. parisii* spores were added to 6 cm NGM plates with

10 µL 10x OP50-1. Plates were dried in a clean cabinet and incubated for a total of 3 hours at room

temperature. After pulse infection, animals were washed 1 – 2x with 1 mL M9+0.1% Tween-20 to remove

excess spores, then added to transparent 24-well assay plates prepared as described above for the

continuous infection assay, except without adding any spores. Three wells were assayed for each

compound for each of three biological replicates. Assay plates were treated as described for continuous

infection assays, except the incubation period was either 2 or 4 days. Samples incubated for 4 days were

acetone-fixed, FISH-stained and DY96-stained, while samples incubated for 2 days were fixed in 4%

PFA and FISH-stained. Fluorescence microscopy was performed as described below.

Spore firing assays

24-well assay plates were prepared exactly as in the continuous infection assay with *N. parisii* spores at

a final concentration of 5,000 spores/µL, compounds at a final concentration of 60 µM except for

fumagillin (see above), and DMSO at a final concentration of 1%. For all spore firing assays containing

TPCK, spores at a concentration of 10,000 spores/µL were incubated for 24 hours at 21°C in K-medium

with compounds at a concentration of 120 µM, and DMSO at a concentration of 2%. These spores were

then used to prepare assay plates exactly as in the continuous infection assay with final concentrations

as stated above. For all spore firing assays where excess compound was removed prior to infection,

spores were washed 3x with 1 mL K-medium before being used in assay plate preparation. Three wells

were assayed for each compound for each of three biological replicates. Assay plates were covered with

adhesive porous film, placed in parafilm wrapped humidity boxes, and incubated at 21°C, shaking at 180

rpm for 3 hours. After incubation, samples were acetone-fixed, FISH-stained, DY96-stained, and subject

to fluorescence microscopy as described below.

Iron chelation and supplementation

24-well assay plates were prepared exactly as in the continuous infection assay with *N. parisii* spores at

a final concentration of 5,000 spores/µL (normal dose) or 78 spores/µL (low dose), 2,2'-bipyridyl (BP) or

dexrazoxane at a final concentration of 60 µM, water-dissolved ferric ammonium citrate (FAC) at a final concentration of 6.6 mg/mL, and DMSO at a final concentration of 1%. Assay plates were treated as described for continuous infection assays. After incubation, samples were acetone-fixed, DY96-stained, and subject to fluorescence microscopy as described below.

RB1074 drug sensitivity tests

~1000 RB1074 or N2 bleach-synchronized L1s were added to 6 cm NGM plates top plated with 120 µL 10x OP50-1, 2,2'-bipyridyl (BP) or dexrazoxane at a final concentration of 60 µM, and DMSO at a final concentration of 1%. Plates were dried in a clean cabinet for ~1 hour at room temperature, and then incubated at 21°C for 3 days. After incubation, live animals were imaged using a Leica Microsystems dissecting scope.

DY96 staining, fluorescence in situ hybridization (FISH), and microscopy

Post-incubation, samples were washed 1 – 2x with 1 mL M9+0.1% Tween-20 to remove excess OP50, fixed in 700 µL acetone and stored at –20°C, or fixed in 500 µL PFA solution (4% PFA, 1x PBS, 0.1% Tween-20) and stored at 4°C until ready for subsequent steps. Samples were then washed 1 – 2x with 1 mL 1xPBS+0.1% Tween-20 and 1x with 1 mL hybridization buffer (900 mM NaCl, 20 mM Tris HCl, 0.01% SDS). 100 µL FISH staining solution (5 ng/µL FISH probe in hybridization buffer) was added, and samples were incubated for 18 – 24 hours at 46°C. The *N. parisii* 18S rRNA-specific microB FISH probe (ctctcggcactccttcctg) conjugated to Cal Fluor Red 610 (LGC Biosearch Technologies) was used (27). After FISH incubation, samples were washed 1x with 1 mL wash buffer (50 mL hybridization buffer + 5 mM EDTA) to remove excess FISH probe. 500 µL DY96 staining solution (20 µg/µL DY96, 0.1% SDS in 1xPBS+0.1% Tween-20) was added, and samples were incubated for 1 hour at 21°C. After DY96 incubation, samples were suspended in EverBrite™ Mounting Medium with DAPI, and subject to fluorescence microscopy using a ZEISS Axio Imager 2 at 5x – 63x magnification. For continuous infection assays, FISH staining steps were omitted, and just stained with DY96.

Quantification of FISH fluorescence

FISH fluorescence was quantified using FIJI version 2.1.0 (70). Animal area was determined by outlining

boundaries based on DAPI staining. Minimum fluorescence threshold for FISH signal was set to 4000 to

exclude auto-fluorescence and background staining of embryos while maximizing inclusion of FISH-

stained *N. parisii* sporoplasms and meronts. The percentage area of animal covered by FISH signal was

calculated.

P. epiphaga **infection assays**

Spores of *P. epiphaga* strain JUm1396 were prepared the same as *N. parisii*. For infections, ~8,000

bleach-synchronized L1s and 80 million *P. epiphaga* spores were added to 6 cm NGM plates with 4 µL

10X OP50-1. Plates were dried and incubated for a total of 3 hours at room temperature. After the

pulse infection, the animals were washed 2X with M9+0.1% Tween-20 to remove excess spores, then

added to transparent 24-well assay plates. Wells contained a total volume of 400 µL of K-medium + 5x

OP50-1 mixed with ~800 worms. DMSO was added to wells for a final concentration of 1%, fumagillin

was added for a final concentration of 350 µM, and dexrazoxane was added for a final concentration of

60 µM. Assay plates were covered with adhesive porous film, placed in parafilm wrapped humidity

boxes, and incubated at 21°C, 180 rpm for 4 days. Samples were fixed in 4% PFA and FISH-stained

with 5ng/µL of FISH probe specific to *P. epiphaga* 18S RNA (CAL Fluor Red 610-

CTCTATACTGTGCGCACGG). Fluorescence microscopy and quantification of FISH fluorescence was

performed as described for *N. parisii* infections.

Determining the effect of dexrazoxane on *A. algerae* cell division

A. algerae spores were purified as previously described (71). Human Fibroblasts (HFF) cells were seeded

on coverslips at a density of $5x10^4$ cells/well in MEM medium supplemented with 10% inactivated FBS,

2 mM glutamine, 2.5 µg.mL^{-1} amphotericin B, 100 µg mL^{-1} streptomycin, 100 U.mL^{-1} penicillin, 25 µg.mL^{-1}

^{1} gentamicin, at 37°C in a humidified 5% CO_2 atmosphere. Once confluence was reached, cells were

infected with 1×10^6 spores of *A. algerae* for 1 h. After 3 washes with culture medium, infected cells were

incubated for 30-35h at 30°C in culture medium containing either 0, 15, 30, 60 or 120 μM of Dexrazoxane.

Two biological replicates were performed, with either 2 or 3 coverslips tested per replicate. After overnight

fixation with methanol at -20°C, parasites were FISH-stained using a Cy3-labeled probe specific to *A.*

algerae, following the protocol described (71). Samples were also stained with DAPI and DY96. Effects

of Dexrazoxane on parasite divisions was evaluated by counting the number of meronts in each infected

host cell using a ZEISS Axio Imager 2 microscope. The number of divisions for each infected cell was

obtained by the following formula: Number of meronts currently dividing $\times$ 2 + number of meronts not

currently dividing -1.

Determining cytotoxicity of dexrazoxane

HFF cells were seeded in 96-well-microplates at a density of *10^4* cells/well in the culture medium

described above at 37°C in a humidified 5% CO_2 atmosphere for approximately 48 h to reach confluence.

The medium was then replaced by fresh culture medium containing different concentrations of

Dexrazoxane (15, 30, 60 and 120 μM). Negative control (cells with culture medium only) and positive

control (cells with 20% DMSO diluted in medium) were also included. Each condition was tested in 3

separate experiments, in 6 wells per experiment. Dexrazoxane cytotoxicity was evaluated using the

tetrazolium dye MTT as previously described (71).

***E. intestinalis* spore propagation and preparation**

E. intestinalis (ATCC 50506) were grown in Vero cells (ATCC CCL-81) using Dulbecco's Modified Eagle's

Medium with high glucose (DMEM) supplemented with Nonessential amino acids (1X) and 10% heat-

inactivated fetal bovine serum (FBS) at 37°C and with 5% CO_2. At 70%-80% confluence, parasites were

added into a 75 cm^2 tissue culture flask and the media was switched to DMEM supplemented with 3%

FBS. Cells were allowed to grow for ten days and medium was changed every two days. To purify spores,

the infected cells were detached from tissue culture flasks using a cell scraper and placed into a 15 ml

conical tube, followed by centrifugation at 1,300 *xg* for 10 min at 25 °C. Cells were resuspended in sterile

DPBS and mechanically disrupted using a G-27 needle. The released spores were purified using a

Percoll gradient. Equal volumes (5 mL) of spore suspension and 100% Percoll were added to a 15 mL

conical tube, vortexed and then centrifuged at 1,800 *xg* for 30 min at room temperature. The purified

spore pellets were washed three times with sterile DPBS and further purified in a discontinuous Percoll

gradient. Briefly, spore pellets were resuspended in 2 mls of sterile DPBS and layered onto a 10 ml four-

layered percoll gradient (2.5 mls 100% Percoll, 2.5 mls 75% Percoll, 2.5 mls 50% Percoll, 2.5 mls 25%

Percoll), centrifuged at 8600 *xg* for 30 min at RT. Spores that separated into the fourth layer (100%

percoll) were carefully collected and washed twice in 10 mls of sterile DPBS at 3000 *xg* for 5 min at RT.

Purified spore pellets were stored in sterile DPBS at 4°C for further analyses.

***E. intestinalis germination and* infection assays**

Purified *E. intestinalis* spores (2.3 x 10^7 spores) were treated with compounds at a final concentration of

60 μM or DMSO at a final concentration of 0.6%. For both spore firing and infectivity assays, spores were

incubated with compounds for 24h at room temperature.

For polar tube germination assays, 0.3 μl of purified *E. intestinalis* spores was placed on poly-L-lysine

treat slides and allowed to air dry briefly. Next, 3 μl of pre-warmed germination buffer (140 mM NaCl, 5

mM KCl, 1 mM $CaCl_2$, 1 mM $MgCl_2$, and 5% (v/v) H_2O_2 at pH 9.5) was added to the slide and sealed with

a #1.5 18 x 18 mm coverslip. PT firing occurs within ~30 s to 1 min of adding germination buffer to the

spores. PT firing was imaged on a ZEISS Elyra 7 microscope with a Zeiss C-Apochromat 40x/1.2 water

objective with a Dual PCO.Edge 4.2 sCMOS camera. All imaging was performed at 37 C. Z stacks were

collected at 0.12 μm spacing. Germinated spores were defined as those in which the polar tube was

released. At least 100 spores were counted per condition.

For measuring infectivity rates, Vero cells were grown on 12mm diameter, #1.5 coverslips and infected

with spores for 24 h. Cells were fixed in 4% PFA in PBS-T (0.1% Tween 20) for 45 min at room

temperature and then processed for FISH. Prior to mounting, cells were stained with NucBlue to visualize

host and parasite DNA. Coverslips were mounted onto slides using ProLong Diamond Antifade (ThermoScientific) and sealed. All samples were imaged on a Nikon W1 spinning disc confocal microscope with a Nikon Apo 60x 1.40 Oil objective and dual Andor 888 Live EMCCD cameras. Z stacks were collected at 0.3 μm spacing. At least 100 cells were counted per condition.

Statistical analysis

All statistical analyses were conducting using Microsoft Excel and R version 3.6.1 or later accessed via RStudio version 1.2.5019 or later (72–74).

References

1. Murareanu BM, Sukhdeo R, Qu R, Jiang J, Reinke AW. Generation of a Microsporidia Species Attribute Database and Analysis of the Extensive Ecological and Phenotypic Diversity of Microsporidia. mBio 12:e01490-21.

2. Wadi L, Reinke AW. 2020. Evolution of microsporidia: An extremely successful group of eukaryotic intracellular parasites. PLoS Pathog 16:e1008276.

3. Stentiford GD, Feist SW, Stone DM, Bateman KS, Dunn AM. 2013. Microsporidia: diverse, dynamic, and emergent pathogens in aquatic systems. Trends Parasitol 29:567–578.

4. Burnham AJ. 2019. Scientific Advances in Controlling Nosema ceranae (Microsporidia) Infections in Honey Bees (Apis mellifera). Front Vet Sci 6.

5. Chaijarasphong T, Munkongwongsiri N, Stentiford GD, Aldama-Cano DJ, Thansa K, Flegel TW, Sritunyalucksana K, Itsathitphaisarn O. 2020. The shrimp microsporidian Enterocytozoon hepatopenaei (EHP): Biology, pathology, diagnostics and control. Journal of Invertebrate Pathology 107458.

6.   Ding Z, Meng Q, Liu H, Yuan S, Zhang F, Sun M, Zhao Y, Shen M, Zhou G, Pan J, Xue H, Wang W. 2016. First

case of hepatopancreatic necrosis disease in pond-reared Chinese mitten crab, Eriocheir sinensis,

associated with microsporidian. Journal of Fish Diseases 39:1043–1051.

7.   Bhat IA, Buhroo Z, Bhat MA. 2017. Microsporidiosis in silkworms with particular reference to mulberry

silkworm (Bombyx Mori L.). International Journal of Entomology Research 2:01–09.

8.   Picard-Sánchez A, Piazzon MC, Ahmed NH, Del Pozo R, Sitjà-Bobadilla A, Palenzuela O. 2020. Enterospora

nucleophila (Microsporidia) in Gilthead Sea Bream (Sparus aurata): Pathological Effects and Cellular

Immune Response in Natural Infections. Vet Pathol 57:565–576.

9.   Kent ML, Elliott DG, Groff JM, Hedrick RP. 1989. Loma salmonae (Protozoa: Microspora) infections in

seawater reared coho salmon Oncorhynchus kisutch. Aquaculture 80:211–222.

10.   Shinn A, Pratoomyot J, Griffiths D, Trọng T, Thanh Vu N, Jiravanichpaisal P, Briggs M. 2018. Asian Shrimp

Production and the Economic Costs of Disease. Asian Fisheries Science 31S.

11.   Ruan Y, Xu X, He Q, Li L, Guo J, Bao J, Pan G, Li T, Zhou Z. 2021. The largest meta-analysis on the global

prevalence of microsporidia in mammals, avian and water provides insights into the epidemic features of

these ubiquitous pathogens. Parasites & Vectors 14:186.

12.   Ostroverkhova NV, Konusova OL, Kucher AN, Kireeva TN, Rosseykina SA. 2020. Prevalence of the

Microsporidian Nosema spp. in Honey Bee Populations (Apis mellifera) in Some Ecological Regions of North

Asia. 3. Veterinary Sciences 7:111.

13.   Stentiford GD, Becnel JJ, Weiss LM, Keeling PJ, Didier ES, Williams BAP, Bjornson S, Kent ML, Freeman MA,

Brown MJF, Troemel ER, Roesel K, Sokolova Y, Snowden KF, Solter L. 2016. Microsporidia – Emergent

Pathogens in the Global Food Chain. Trends Parasitol 32:336–348.

14.   Katznelson H, Jamieson CA. 1952. Control of Nosema Disease of Honeybees with Fumagillin. Science

115:70–71.

15.   Huang W-F, Solter LF, Yau PM, Imai BS. 2013. Nosema ceranae Escapes Fumagillin Control in Honey Bees.

PLOS Pathogens 9:e1003185.

16.   Tang KFJ, Han JE, Aranguren LF, White-Noble B, Schmidt MM, Piamsomboon P, Risdiana E, Hanggono B.

2016. Dense populations of the microsporidian Enterocytozoon hepatopenaei (EHP) in feces of Penaeus

vannamei exhibiting white feces syndrome and pathways of their transmission to healthy shrimp. Journal

of Invertebrate Pathology 140:1–7.

17.   van den Heever JP, Thompson TS, Curtis JM, Ibrahim A, Pernal SF. 2014. Fumagillin: an overview of recent

scientific advances and their significance for apiculture. J Agric Food Chem 62:2728–2737.

18.   Han B, Weiss LM. 2018. Therapeutic targets for the treatment of microsporidiosis in humans. Expert Opin

Ther Targets 22:903–915.

19.   He Q, Leitch GJ, Visvesvara GS, Wallace S. 1996. Effects of nifedipine, metronidazole, and nitric oxide

donors on spore germination and cell culture infection of the microsporidia Encephalitozoon hellem and

Encephalitozoon intestinalis. Antimicrob Agents Chemother 40:179–185.

20.   Didier ES, Maddry JA, Kwong CD, Green LC, Snowden KF, Shadduck JA. 1998. Screening of compounds for

antimicrosporidial activity in vitro. Folia Parasitologica 45:129–139.

21.   Borges D, Guzman-Novoa E, Goodwin PH. 2020. Control of the microsporidian parasite Nosema ceranae in

honey bees (Apis mellifera) using nutraceutical and immuno-stimulatory compounds. PLOS ONE

15:e0227484.

22.   Zhang G, Sachse M, Prevost M-C, Luallen RJ, Troemel ER, Felix M-A. 2016. A Large Collection of Novel

Nematode-Infecting Microsporidia and Their Diverse Interactions with Caenorhabditis elegans and Other

Related Nematodes. PLoS Pathog 12:e1006093.

23.   Willis AR, Zhao W, Sukhdeo R, Wadi L, Jarkass HTE, Claycomb JM, Reinke AW. 2021. A parental

transcriptional response to microsporidia infection induces inherited immunity in offspring. Science

Advances 7:eabf3114.

24.   Bakowski MA, Desjardins CA, Smelkinson MG, Dunbar TL, Lopez-Moyado IF, Rifkin SA, Cuomo CA, Troemel

ER. 2014. Ubiquitin-mediated response to microsporidia and virus infection in C. elegans. PLoS Pathog

10:e1004200.

25.   Reinke AW, Balla KM, Bennett EJ, Troemel ER. 2017. Identification of microsporidia host-exposed proteins

reveals a repertoire of rapidly evolving proteins. Nat Commun 8:14023.

26.   Estes KA, Szumowski SC, Troemel ER. 2011. Non-lytic, actin-based exit of intracellular parasites from C.

elegans intestinal cells. PLoS Pathog 7:e1002227.

27.   Troemel ER, Félix M-A, Whiteman NK, Barrière A, Ausubel FM. 2008. Microsporidia Are Natural

Intracellular Parasites of the Nematode Caenorhabditis elegans. PLOS Biology 6:e309.

28.   Balla KM, Luallen RJ, Bakowski MA, Troemel ER. 2016. Cell-to-cell spread of microsporidia causes

Caenorhabditis elegans organs to form syncytia. 11. Nature Microbiology 1:1–6.

29.   Luallen RJ, Reinke AW, Tong L, Botts MR, Felix M-A, Troemel ER. 2016. Discovery of a Natural

Microsporidian Pathogen with a Broad Tissue Tropism in Caenorhabditis elegans. PLoS Pathog

12:e1005724.

30.   Ewbank JJ, Zugasti O. 2011. C. elegans: model host and tool for antimicrobial drug discovery. Disease

Models & Mechanisms 4:300–304.

31. Peterson ND, Pukkila-Worley R. 2018. C. elegans in high-throughput screens for anti-infective compounds.

Curr Opin Immunol 54:59–65.

32. Burns AR, Luciani GM, Musso G, Bagg R, Yeo M, Zhang Y, Rajendran L, Glavin J, Hunter R, Redman E, Stasiuk

S, Schertzberg M, Angus McQuibban G, Caffrey CR, Cutler SR, Tyers M, Giaever G, Nislow C, Fraser AG,

MacRae CA, Gilleard J, Roy PJ. 2015. Caenorhabditis elegans is a useful model for anthelmintic discovery. 1.

Nature Communications 6:7485.

33. Anastassopoulou CG, Fuchs BB, Mylonakis E. 2011. Caenorhabditis elegans-based Model Systems for

Antifungal Drug Discovery. Curr Pharm Des 17:1225–1233.

34. Mylonakis E, Ausubel FM, Perfect JR, Heitman J, Calderwood SB. 2002. Killing of Caenorhabditis elegans by

Cryptococcus neoformans as a model of yeast pathogenesis. Proc Natl Acad Sci U S A 99:15675–15680.

35. Moy TI, Ball AR, Anklesaria Z, Casadei G, Lewis K, Ausubel FM. 2006. Identification of novel antimicrobials

using a live-animal infection model. Proc Natl Acad Sci U S A 103:10414–10419.

36. Moy TI, Conery AL, Larkins-Ford J, Wu G, Mazitschek R, Casadei G, Lewis K, Carpenter AE, Ausubel FM.

2009. High-throughput screen for novel antimicrobials using a whole animal infection model. ACS Chem

Biol 4:527–533.

37. O'Reilly LP, Luke CJ, Perlmutter DH, Silverman GA, Pak SC. 2014. C. elegans in high-throughput drug

discovery. Adv Drug Deliv Rev 0:247–253.

38. Luallen RJ, Bakowski MA, Troemel ER. 2015. Characterization of Microsporidia-Induced Developmental

Arrest and a Transmembrane Leucine-Rich Repeat Protein in Caenorhabditis elegans. PLOS ONE

10:e0124065.

39. Hakim A, Mor Y, Toker IA, Levine A, Neuhof M, Markovitz Y, Rechavi O. 2018. WorMachine: machine

learning-based phenotypic analysis tool for worms. BMC Biology 16:8.

40.   Wittkowski P, Violet N, Oelgeschläger M, Schönfelder G, Vogl S. 2020. A quantitative medium-throughput

assay to measure Caenorhabditis elegans development and reproduction. STAR Protocols 1:100224.

41.   Wittkowski P, Marx-Stoelting P, Violet N, Fetz V, Schwarz F, Oelgeschläger M, Schönfelder G, Vogl S. 2019.

Caenorhabditis elegans As a Promising Alternative Model for Environmental Chemical Mixture Effect

Assessment—A Comparative Study. Environ Sci Technol 53:12725–12733.

42.   Mathew MD, Mathew ND, Ebert PR. 2012. WormScan: A Technique for High-Throughput Phenotypic

Analysis of Caenorhabditis elegans. PLOS ONE 7:e33483.

43.   Vávra J, Larsson JIR. 2014. Structure of Microsporidia, p. 1–70. *In* Microsporidia. John Wiley & Sons, Ltd.

44.   Zhang G, Sachse M, Prevost M-C, Luallen RJ, Troemel ER, Félix M-A. 2016. A Large Collection of Novel

Nematode-Infecting Microsporidia and Their Diverse Interactions with Caenorhabditis elegans and Other

Related Nematodes. PLOS Pathogens 12:e1006093.

45.   Rodríguez-García C, Heerman MC, Cook SC, Evans JD, DeGrandi-Hoffman G, Banmeke O, Zhang Y, Huang S,

Hamilton M, Chen YP. 2021. Transferrin-mediated iron sequestration suggests a novel therapeutic strategy

for controlling Nosema disease in the honey bee, Apis mellifera. PLoS Pathogens 17.

46.   Didier ES, Bowers LC, Martin AD, Kuroda MJ, Khan IA, Didier PJ. 2010. Reactive nitrogen and oxygen

species, and iron sequestration contribute to macrophage-mediated control of Encephalitozoon cuniculi

(Phylum Microsporidia) infection in vitro and in vivo. Microbes and Infection 12:1244–1251.

47.   Rajan M, Anderson CP, Rindler PM, Romney SJ, Ferreira dos Santos MC, Gertz J, Leibold EA. 2019. NHR-14

loss of function couples intestinal iron uptake with innate immunity in C. elegans through PQM-1 signaling.

eLife 8:e44674.

48.   Tamim El Jarkass H, Mok C, Schertzberg MR, Fraser AG, Troemel ER, Reinke AW. 2021. An intestinally

secreted host factor limits bacterial colonization but promotes microsporidia invasion of C. elegans. bioRxiv

2021.07.12.452088.

49.   Dang X, Pan G, Li T, Lin L, Ma Q, Geng L, He Y, Zhou Z. 2013. Characterization of a subtilisin-like protease

with apical localization from microsporidian Nosema bombycis. J Invertebr Pathol 112:166–174.

50.   Yavelow J, Caggana M, Beck KA. 1987. Proteases occurring in the cell membrane: a possible cell receptor

for the Bowman-Birk type of protease inhibitors. Cancer Res 47:1598–1601.

51.   Bojko J, Reinke AW, Stentiford GD, Williams B, Rogers MSJ, Bass D. 2022. Microsporidia: a new taxonomic,

evolutionary, and ecological synthesis. Trends in Parasitology 0.

52.   Weiss LM, Takvorian PM. 2021. Anncaliia algerae. Trends in Parasitology 37:762–763.

53.   Hinney B, Sak B, Joachim A, Kváč M. 2016. More than a rabbit's tale – Encephalitozoon spp. in wild

mammals and birds. International Journal for Parasitology: Parasites and Wildlife 5:76–87.

54.   Reichardt P, Tabone M-D, Mora J, Morland B, Jones RL. 2018. Risk–benefit of dexrazoxane for preventing

anthracycline-related cardiotoxicity: re-evaluating the European labeling. Future Oncology 14:2663–2676.

55.   Jones RL. 2008. Utility of dexrazoxane for the reduction of anthracycline-induced cardiotoxicity. Expert

Review of Cardiovascular Therapy 6:1311–1317.

56.   Hasinoff BB. 1998. Chemistry of dexrazoxane and analogues. Semin Oncol 25:3–9.

57.   Kirienko NV, Kirienko DR, Larkins-Ford J, Wählby C, Ruvkun G, Ausubel FM. 2013. Pseudomonas aeruginosa

Disrupts Caenorhabditis elegans Iron Homeostasis, Causing a Hypoxic Response and Death. Cell Host &

Microbe 13:406–416.

58. Deng S, Yan T, Nikolova T, Fuhrmann D, Nemecek A, Gödtel-Armbrust U, Kaina B, Wojnowski L. 2015. The

catalytic topoisomerase II inhibitor dexrazoxane induces DNA breaks, ATF3 and the DNA damage response

in cancer cells. British Journal of Pharmacology 172:2246–2257.

59. Zdraljevic S, Strand C, Seidel HS, Cook DE, Doench JG, Andersen EC. 2017. Natural variation in a single

amino acid substitution underlies physiological responses to topoisomerase II poisons. PLOS Genetics

13:e1006891.

60. Han B, Takvorian PM, Weiss LM. 2020. Invasion of Host Cells by Microsporidia. Front Microbiol 11.

61. Tamim El Jarkass H, Reinke AW. 2020. The ins and outs of host-microsporidia interactions during invasion,

proliferation and exit. Cellular Microbiology 22:e13247.

62. Dang X, Pan G, Li T, Lin L, Ma Q, Geng L, He Y, Zhou Z. 2013. Characterization of a subtilisin-like protease

with apical localization from microsporidian Nosema bombycis. J Invertebr Pathol 112:166–174.

63. P. Wilson BA, C. Thornburg C, J. Henrich C, Grkovic T, R. O'Keefe B. 2020. Creating and screening natural

product libraries. Natural Product Reports 37:893–918.

64. Rosenberg A, Luth MR, Winzeler EA, Behnke M, Sibley LD. 2019. Evolution of resistance in vitro reveals

mechanisms of artemisinin activity in Toxoplasma gondii. PNAS 116:26881–26891.

65. Cowell AN, Istvan ES, Lukens AK, Gomez-Lorenzo MG, Vanaerschot M, Sakata-Kato T, Flannery EL,

Magistrado P, Owen E, Abraham M, LaMonte G, Painter HJ, Williams RM, Franco V, Linares M, Arriaga I,

Bopp S, Corey VC, Gnädig NF, Coburn-Flynn O, Reimer C, Gupta P, Murithi JM, Moura PA, Fuchs O, Sasaki E,

Kim SW, Teng CH, Wang LT, Akidil A, Adjalley S, Willis PA, Siegel D, Tanaseichuk O, Zhong Y, Zhou Y, Llinás

M, Ottilie S, Gamo F-J, Lee MCS, Goldberg DE, Fidock DA, Wirth DF, Winzeler EA. 2018. Mapping the

malaria parasite druggable genome by using in vitro evolution and chemogenomics. Science 359:191–199.

66.   Herneisen AL, Sidik SM, Markus BM, Drewry DH, Zuercher WJ, Lourido S. 2020. Identifying the Target of an

Antiparasitic Compound in Toxoplasma Using Thermal Proteome Profiling. ACS Chem Biol 15:1801–1807.

67.   Lewis JA, Fleming JT. 1995. Basic culture methods. Methods Cell Biol 48:3–29.

68.   Lehner B, Tischler J, Fraser AG. 2006. RNAi screens in Caenorhabditis elegans in a 96-well liquid format and

their application to the systematic identification of genetic interactions. 3. Nature Protocols 1:1617–1620.

69.   The GIMP Development Team. 2018. GIMP. https://www.gimp.org. https://www.gimp.org/. Retrieved 3

September 2021.

70.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S,

Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-

source platform for biological-image analysis. Nat Methods 9:676–682.

71.   Prybylski N, Fayet M, Dubuffet A, Delbac F, Kocer A, Gardarin C, Michaud P, El Alaoui H, Dubessay P. 2022.

Ricin B lectin-like proteins of the microsporidian Encephalitozoon cuniculi and Anncaliia algerae are

involved in host-cell invasion. Parasitology International 87:102518.

72.   Microsoft Corporation. 2021. Microsoft Excel. https://office.microsoft.com/excel.

https://www.microsoft.com/en-ca/microsoft-365/excel. Retrieved 3 September 2021.

73.   R Core Tam. 2021. R: A language and environment for statistical computing. https://wwwR-project.org.

https://www.r-project.org/. Retrieved 3 September 2021.

74.   RStudio Team. 2020. RStudio: Integrated Development for R. https://rstudio.com. https://rstudio.com/.

Retrieved 3 September 2021.

745   **Figures and Tables**

Figure 1. High-throughput drug screen of the Spectrum Collection identifies compounds that restore progeny production in *C. elegans* infected with *N. parisii*. (A) Schematic of small molecule inhibitor screen (see methods). Bleach-synchronized L1 animals were incubated with 60 µM of each compound from the Spectrum Collection for 6 days at 21 °C in the presence of *N. parisii* spores. 2,560 total compounds were screened once, yielding 25 initial hits. The initial hits were retested once, yielding 11 confirmed hits. The effectiveness of these 11 compounds was then quantified across multiple replicates of the screening assay using WorMachine. **(B)** Representative images of wells containing worms. **(B Far Left)** Normal worm growth in the absence of spores. **(B Middle Left)** Microsporidia

infection leads to inhibition of progeny production. **(B Middle & Far Right)** Treatment with anti-microsporidial compounds restores progeny production. **(C & D)** Effect of compounds on progeny production in **(C)** infected and **(D)** uninfected animals. Progeny levels expressed as a percentage of the DMSO uninfected control. Statistical significance evaluated in relation to DMSO controls using two-sided t-tests: *** = *p < 0.001*, ** = *p < 0.01*, * = *p < 0.05, ns = not significant (p > 0.05)*. Data for each condition includes 3 – 6 biological replicates.

Figure 2. Identified compounds inhibit microsporidia infection. (A) Schematic of continuous infection assay (see methods). Bleach-synchronized L1 animals were incubated with compounds for 4 days at 21 °C in the presence of *N. parisii* spores in liquid. Animals were subsequently fixed with acetone, and stained with DAPI and DY96. **(B)** Representative images taken at 5x magnification; scale bars are 500 µm. **(B Far Left)** Normal worm growth in the absence of spores. **(B Middle Left)** Microsporidia infection results in production of new spores highlighted in bright green by DY96, and slows growth thereby preventing development of gravid adults. **(B Middle Right & Far Right)** Treatment with anti-

microsporidial compounds reduces formation of new spores, and restores the development of gravid

adults. **(C)** Percentage of animals that contain embryos (N = ≥200 animals counted per biological

replicate). **(D)** Percentage of animals that contain newly formed spores (N = ≥200 animals counted per

biological replicate). Significance evaluated in relation to DMSO infected controls using one-way ANOVA

with Dunnett's post-hoc test: *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant ($p > 0.05$).

Data for each condition includes 3 biological replicates.

Figure 3. Dexrazoxane prevents microsporidia proliferation. (A) Schematic of pulse-chase infection

assay (see methods). Bleach-synchronized L1 animals were pulse infected with *N. parisii* spores for 3

hours at 21 °C on NGM plates. Excess spores were washed away, and infected animals were incubated

with compounds for 2 or 4 days at 21 °C in liquid. Animals were fixed in acetone or PFA, FISH stained,

and then stained with DY96 and DAPI. **(B)** Representative images of acetone-fixed animals 4 days post

infection taken at 5x magnification; scale bars are 500 μm. **(B Far Left)** Normal growth in uninfected

worms. **(B Middle Left)** Pulse infection results in sporoplasms and meronts highlighted in red by microB

FISH probes and new spores highlighted in bright green by DY96. Pulse infection also slows growth

thereby preventing development of gravid adults. **(B Middle Right & Far Right)** Treatment of pulse-

infected animals with fumagillin or dexrazoxane reduces sporoplasms and meronts, new spores, and

restores development of gravid adults. **(C)** Representative images (z-stack maximum intensity

projections; 7 slices, 0.25 μm spacing) of PFA-fixed animals 2 days post infection taken at 63x

magnification; scale bars are 20 μm. **(C Left)** In the absence of drug treatment, pulse-infected animals

develop large meronts with many nuclei. **(C Middle)** Fumagillin treatment restricts proliferation; both large

and small meronts are observed. **(C Right)** Dexrazoxane treatment restricts proliferation; only small

meronts with one or two nuclei are observed. **(D)** Percentage of animals with embryos (N = ≥170 animals

counted per biological replicate). **(E)** Percentage of animals with newly formed spores (N = ≥170 animals

counted per biological replicate). **(F)** Percentage of animals with FISH signal (N = ≥170 animals counted

per biological replicate). ANOVA not significant ($p = 0.111$). **(G)** Quantitation of FISH fluorescence per

worm (N = 15 animals quantified per biological replicate). **(H)** Effects of iron chelation with BP on infection,

and effects of iron supplementation with FAC on dexrazoxane activity. Percentage of animals with newly

formed spores (N = ≥120 animals counted per biological replicate) after 4 days of continuous infection is

shown. Significance evaluated in relation to DMSO infected controls using one-way ANOVA with

Dunnett's post-hoc test: *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant ($p > 0.05$). Data

for each condition includes 3 biological replicates.

Figure 4. Several compounds prevent microsporidia infection by inhibiting spore firing. (A) Schematic of spore firing assay (see methods). Bleach-synchronized L1 animals were incubated with compounds for 3 hours at 21 °C in the presence of *N. parisii* spores in liquid. Spore prep control was generated by incubating spores in liquid in the absence of *C. elegans*. Animals and spores were fixed in acetone, and stained with microB FISH probes and DY96. **(B)** Schematic of modified spore firing assay (see methods). Spores were incubated with compounds for 24 hours at 21 °C**,** and then used to infect beach-synchronized L1 animals either with or without prior washing to remove excess compounds. **(C)** Representative images of unfired and fired spores at 63x magnification; scale bars are 5 µm. Unfired spore is indicated by white arrowhead. **(D & E)** Effects of microsporidia inhibitors on **(D)** the percentage of fired spores in the *C. elegans* intestinal lumen (N = ≥60 spores counted per biological replicate) and **(E)** the average number of sporoplasms per animal (N = ≥40 animals counted per biological replicate) in a spore firing assay. **(F & G)** Effects of serine protease inhibitors on **(F)** spore firing and **(G)** sporoplasm invasion in a modified spore firing assay without washing away compounds prior to infection (N = ≥39 animals and ≥60 spores counted per biological replicate, except for one chymostatin replicate where only 12 animals and 19 spores were counted). **(H & I)** Effects of spore firing inhibitors on **(H)** spore firing and

(I) sporoplasm invasion in a modified spore firing assay with compounds washed away prior to infection

(N = ≥40 animals and ≥70 spores counted per biological replicate). Significance evaluated in relation to

DMSO controls using one-way ANOVA with Dunnett's post-hoc test: *** = $p < 0.001$, ** = $p < 0.01$, * = p

< 0.05, ns = not significant ($p > 0.05$). Data for each condition includes 3 biological replicates.

Figure 5. Identified inhibitors display activity against multiple diverse microsporidia species. (A-

B) Bleach-synchronized L1 animals were pulse infected with *P. epiphaga* spores for 3 hours on NGM

plates. Excess spores were washed away, and infected animals were incubated with 350 μM fumagillin

or 60 μM dexrazoxane for 4 days at 21 °C in liquid. Animals were fixed and stained with DAPI and a FISH

probe. **(A)** Representative images of animals 4 days post infection; scale bars are 25 μm. **(B)** Quantitation

of FISH fluorescence per worm (N = 15 animals quantified in each condition for three biological

replicates). **(C-D)** Cells infected with *A. algerae* spores for 1 hour with various concentrations of

dexrazoxane for 30-35 hours. **(C)** Representative images of an infected cell; scale bars are 5 µm. **(D)**

Average number of *A. algerae* divisions per cell (Between 30-243 cells were analyzed per each two

biological replicates). **(E)** Host cell viability (experiment consisted of three biological replicates). **(F-H)** *E.*

intestinalis spores were treated with 60 µM inhibitors for 24 hours. Spores were either induced to fire (F)

or used to infect cells for 24 h and then stained with a FISH probe and DAPI (G-H). **(F)** Percentage of

spores that have undergone germination (N = ≥100 spores counted per each of five biological replicates).

(G) Representative images of cells infected with *E. intestinalis* that were either untreated or treated with

ZPCK; scale bars are 10 µm. Merged images are shown on the left and individual channels to the right.

DNA (grey), *E. intestinalis* FISH (yellow). **(H)** Percentage of cells infected (N = ≥100 cells counted per

biological replicate for each of four biological replicates). Significance evaluated in relation to DMSO

infected controls using one-way ANOVA with Dunnett's post-hoc test: *** = $p < 0.001$, ** = $p < 0.01$, * =

$p < 0.05$, ns = not significant ($p > 0.05$).

Figure 6. Compound structures and mechanisms. (A) Structures of compounds containing a quinone

moiety (circled in red) that were identified in the initial screen as inhibitors of microsporidia infection. **(B)**

Microsporidia infection model depicting the stages at which various microsporidia inhibitors act. Two

protease inhibitors and four quinone-derivatives were shown to act directly on *N. parisii* spores to prevent

spore firing and subsequent invasion of sporoplasms. Fumagillin and dexrazoxane act after invasion has

occurred, inhibiting proliferation of sporoplasms and meronts, ultimately reducing parasite burden and

preventing the production of microsporidia spores. Figure made using Biorender.com.

Supplemental material

Figure S1. Quantitation of progeny number using semi-automated image processing. (A) Flatbed

scanner image of a plate following staining and dilution steps. **(B)** Sample images of wells prior to

processing. **(C)** Sample images of wells after processing. **(D)** Correlation between manual counts and

WorMachine counts (N = 24). **(E)** Correlations between pairs of technical replicates as a measure of

technical variability (N = 140 per correlation).

Figure S2. *N. parisii* infection in *C. elegans* is unaffected by altering iron levels. (A & B) Effects of

iron supplementation with FAC on *N. parisii* infection in *C. elegans*. **(A)** Percentage of animals with newly

formed spores (N = ≥140 animals counted per biological replicate; significance evaluated using a two-

sample t-test) and **(B)** percentage of animals with embryos (N = ≥140 animals counted per biological

replicate; significance evaluated using a one-way ANOVA with Tukey's post-hoc test) after 4 days of

continuous infection with a low dose of spores are shown. **(C)** Effects of iron chelation with BP on

infection, and effects of iron supplementation with FAC on dexrazoxane activity. Percentage of animals

with embryos (N = ≥120 animals counted per biological replicate; significance evaluated using a one-way

ANOVA with Tukey's post-hoc test) after 4 days of continuous infection with a normal dose of spores is

shown. $*** = p < 0.001$, $** = p < 0.01$, $* = p < 0.05$, ns = not significant ($p > 0.05$). Data for each condition

includes 3 biological replicates.

| | DMSO | 2,2'-bipyridyl (BP) (60 µM) | Dexrazoxane (60 µM) |

Figure S3. *C. elegans* mutant strain with reduced iron is not sensitive to dexrazoxane. RB1074

animals have 50% less iron compared to N2 and display severe growth defects during treatment with the

iron chelator BP on NGM plates for 3 days from the L1 stage (bottom middle), but not after treatment with

dexrazoxane (bottom right) (47).

Table S1: Compound names and available PubChem CIDs.

Compound	PubChem CID
Ethinyl Estradiol	5991
Menadione	4055
Parthenolide	108068
Plumbagin	10205
Curcumin	969516
Chloranil	8371
4,4'-Dimethoxydalbergione (Dalbergione)	364106
4,6-Dimethoxytoluquinone (Toluquinone)	NA
Thymoquinone	10281
1-Benzyloxycarbonylaminophenethyl chloromethyl ketone (ZPCK)	99625
Dexrazoxane	71384