Impact of pH and removed filtrate on *E. coli* regrowth and microbial community during storage of electro-dewatered biosolids

Tala Navab-Daneshmand [a,b], Bing Guo [b,c], Ronald Gehr [b], Dominic Frigon [b,*]

[a] School of Chemical, Biological, and Environmental Engineering, Oregon State University, 105 SW 26th St., Corvallis, Oregon 97331, USA
[b] Department of Civil Engineering and Applied Mechanics, McGill University, 817 Sherbrooke St. West, Montreal, Quebec H3A 0C3, Canada
[c] Centre for Environmental Health and Engineering (CEHE), Department of Civil and Environmental Engineering, University of Surrey, Guildford, Surrey GU2 7XH United Kingdom

ABSTRACT

Residual biosolids can be land applied if they meet microbiological requirements at the time of application. Electro-dewatering technology is shown to reduce biosolids bacterial counts to detection limits with little potential for bacterial regrowth during incubations. Here, we investigated the impacts on *Escherichia coli* regrowth and microbial communities of biosolids pH, removed nutrients via the filtrate, and produced inhibitory compounds in electro-dewatered biosolids. Findings suggest pH as the primary mechanism impacting *E. coli* regrowth in electro-dewatered biosolids. Propidium monoazide treatments were effective to remove DNA from dead cells based on the removal of obligate anaerobes observed after anaerobic incubation. Analyses of high throughput sequenced data showed lower alpha-diversities associated with electro-dewatering treatment and incubation time. Moreover, biosolids pH and incubation period were the main factors contributing to the variations in microbial community compositions after incubation. Results highlight the role of biosolids pH on regrowing culturable bacteria and overall microbial communities.

Keywords: *E. coli* regrowth, biosolids, electro-dewatering, pH, microbial communities

1. Introduction

Biosolids are the residues produced during the secondary treatment of wastewater and can be land applied if they meet certain standards. The US-EPA classifies biosolids as either Class A or B – in regards to pathogen indicators – for land application (US-EPA, 2003). Class A requirements stipulate a reduction of fecal coliforms as bacterial pathogen indicators to <1,000 most probable number per gram of total solids (MPN/g-TS), whereas Class B requires $<2 \times 10^6$ MPN/g-TS of fecal coliforms. Because of the less stringent standards, land application of Class B biosolids is accompanied by a number of regulatory restrictions with respect to grazing animals and public access. An important caveat is that the above-mentioned US-EPA requirements are to be met at the time of land application and not immediately after biosolids treatment. Accordingly, Class A pathogen requirement must be achieved after transportation and storage periods, or after any mixing with other materials (US-EPA, 2003). Low bacterial pathogen counts immediately after treatment do not guarantee that the inactivation is irreversible (Dentel et al., 2008). For example, centrifuge-dewatered biosolids from mesophilic and thermophilic anaerobic digesters showed 0.8-4.1 logs increase in fecal coliforms and *E. coli* after 24 h storage (Chen et al.,

[*] Corresponding author at: Department of Civil Engineering and Applied Mechanics, McGill University, 817 Sherbrooke Street West, Montréal, Québec, H3A 0C3, Canada. Email address: dominic.frigon@mcgill.ca

2011). Similarly, Higgins and colleagues reported 3.9 logs increase of *E. coli* counts to 10^8-10^9 MPN/g-TS after 1-2 days' incubation in anaerobically digested centrifuged biosolids (Higgins et al., 2007). These counts then decreased back to the detection limits by 20 and 60 days in mesophilic and thermophilic biosolids, respectively.

The electro-dewatering process reduces the water content in raw biosolids from 67-87% w/w to as low as 30% w/w in dewatered solids (Saveyn et al., 2006; Tuan et al., 2008). The process uses an electrical field at the same time as pressure to remove water by electro-osmosis. In addition to dewatering, the 10-min electro-dewatering treatment has been shown to reduce total coliforms and *Escherichia coli* concentrations to below the detection limits (100-600 MPN/g-TS) to meet Class A requirements for bacterial pathogen indicators (Navab-Daneshmand et al., 2012). Our previous work examined the regrowth potential of electro-dewatered biosolids after 10-min dewatering cycles (Navab-Daneshmand et al., 2014a). Regrowth potentials of total coliforms and *E. coli* were compared in electro-dewatered biosolids and heat-treated controls over 7 days under aerobic and anaerobic incubations; no regrowth was observed in either solids (Navab-Daneshmand et al., 2014a). When electro-dewatered and heat-treated biosolids were inoculated with untreated solids to mimic potential on-site contamination, counts in electro-dewatered biosolids returned to the detection limits after 4 d, whereas 4-5 logs increase were observed in heat-treated controls. Regrowth of bacterial pathogens in electro-dewatered biosolids during the storage phase can be affected by several environmental conditions. The most important parameters are the availability of oxygen and other electron acceptors, temperature, water activity, pH, and the availability of nutrients. Our previous work demonstrated that the availability of oxygen as an electron acceptor has no significant impact on the regrowth of total coliforms and *E. coli* in electro-dewatered biosolids during 7-day incubations (Navab-Daneshmand et al., 2014a, 2014b). Storage temperature is another important factor, but because there is rarely much variation in storage temperatures, it was not examined here. The remaining parameters and specific observations in the literature are reviewed below.

Water activity: Water activity is a measure of water availability to microorganisms (i.e., the unbound water supporting microbial activities). It is defined as the ratio of the vapor pressure of water in equilibrium with biosolids to the saturated vapor pressure of pure water. For *E. coli* regrowth to be inhibited, the water activity must be below 0.95 (Tapia et al., 2008). Our previous work reported water activities exceeding 0.95 in electro-dewatered biosolids with total solids as high as ~60% w/w (Navab-Daneshmand et al., 2015). Therefore, water activity is not a controlling parameter for *E. coli* regrowth in electro-dewatered biosolids with the current process limitations for dewatering.

pH: Biosolids pH affects nutrient availability and the metabolic well-being of growing bacteria. Extreme pH values cause enzymes to denature, which impairs their functions. Thus, there are pH minima, maxima, and optima for the growth of each bacterial strain. For *E. coli*, *Salmonella typhimurium*, and *Shigella* spp., the pH minima are approximately 3.90 (Koutsoumanis et al., 2004; Presser et al., 1997), and the pH optima are between 5 to 9 (Small et al., 1994). Slight changes in pH can profoundly affect regrowth dynamics; for example, *E. coli*'s growth rate increases 4.5 times from pH 4 to 5 (Presser et

al., 1997). During biosolids electro-dewatering, electrolysis produces protons at the anode, and cations are electrophoretically removed with the water through the cathode, leading to a decrease in the cake's overall pH (Navab-Daneshmand et al., 2012). It has been suggested that the lower pH of electro-dewatered biosolids (4.6 in the electro-dewatered cake compared to 7.1 in untreated biosolids) could be a factor contributing to the decrease of total coliforms regrowth (Navab-Daneshmand et al., 2014a).

Removal of essential nutrients and precursors: The removal or addition of certain substrates or inhibitors in the biosolids matrix could impact the regrowth potential of bacterial pathogens. For example, the addition of the removed centrate from centrifuge dewatering to the dewatered cake reduced regrowth of fecal coliforms in mesophilically digested biosolids by more than 1.5 logs (Gardner and Oermeci, 2010). The removal or addition of certain autoinducers could resuscitate viable but non-culturable in to a culturable state (Reissbrodt et al., 2002). In another study, the addition of substrate to mesophilically digested biosolids increased fecal coliform counts by 3.4 logs after 24 h (Chen et al., 2011). Justification for this phenomenon is that the filtrate removed during electro-dewatering could contain nutrients that are essential for bacterial regrowth, thereby reducing the regrowth of bacteria in electro-dewatered biosolids, or that the production of compounds during electro-dewatering are inhibitory to bacterial regrowth.

The synergistic effect of the above-mentioned environmental conditions could also impact the regrowth potential of bacterial pathogens. This work investigated the impacts of biosolids pH, removed nutrients via the filtrate, and/or produced inhibitory compounds on *E. coli* regrowth and taxonomic compositions of the microbial communities during incubation of biosolids after electro-dewatering.

2. Materials and methods

2.1. Biosolids
The biosolids were obtained from an activated sludge wastewater treatment plant near Montréal. The plant receives a flow of ~60,000 m^3/d consisting essentially of 50% municipal and 50% industrial sources (mainly pulp and paper, and food industries). It produces over 70 ton/d of residual biosolids. Two cationic polymers were added to the dewatering stream: Flo-CA475 at 1-4 kg/ton-TS before the flotation units, and Flo-CA4800 at 12-21 kg/ton-TS before the centrifuge dewatering units. Solids leaving the centrifuge units had totals solids of 17.1-19.8% w/w and pH of 6.4-7.3 with *E. coli* counts of 3.9-6.1 log MPN/g-TS. Solid samples were collected in plastic bags at the end of the solids handling unit, brought to the McGill laboratory on ice and stored at 4 ºC for up to 2 d before further testing.

2.2. Biosolids treatment and incubation
The laboratory electro-dewatering unit used was a CINETIK® CK-lab model (Ovivo, Boucherville, Québec, Canada). The unit worked with direct current and the maximum voltage and current were set to 60 V and 5.5 A, respectively. Similarly to our previous study (Navab-Daneshmand et al., 2012), ~165 g of wet untreated biosolids were placed on

the filter medium (PPS Ryton, woven) over the perforated stainless steel cathode. The ceramic-coated titanium anode applied 140 kPa constant pressure over the cake. Filtrate produced through the perforated cathode was collected in a filtrate-receiving container during the tests. The 10-min electro-dewatering cycles used in this study were sufficient to reduce *E. coli* counts to below the detection limits.

Heat treatment was used as control treatment, because our previous results demonstrated that high temperature due to Joule heating was the main inactivation mechanism during electro-dewatering (Navab-Daneshmand et al., 2012). As previously described, 30-40 wet g of biosolids were placed in 50 mL glass tubes and incubated in a water bath at 80 ℃ for 16 min to achieve similar heat treatment to the electro-dewatering tests (Navab-Daneshmand et al., 2014a).

After treatment, to mimic biosolids contamination during transport or storage, some microcosms were inoculated with untreated biosolids. For this, 2 wet g of untreated biosolids were dispersed in 40 mL distilled water with a homogenizer (ULTRA-TURRAX® S10N-10G, IKA Works Inc., Wilmington, NC) for 30 s. From the slurry, 1 mL was added to ~50 wet g treated sample and mixed thoroughly by hand. To control for the impact of the nutrients removed with the filtrate during electro-dewatering, in some microcosms the filtrate was added back to the dewatered cake and mixed thoroughly by hand until a uniform texture was achieved. To observe the impact of pH, in some microcosms HCl or NaOH solutions were added to the treated biosolids before incubation. To adjust the pH of the electro-dewatered samples to a level close to that of heat-treated biosolids, 12 mL of 0.5 M NaOH solution were added to ~50 wet g electro-dewatered biosolids and mixed thoroughly by hand. To lower the pH of heat-treated biosolids, 12 mL of 0.5 M HCl solution were added to ~50 wet g heat-treated samples. Similarly, to lower the pH of electro-dewatered biosolids with filtrate additive, 12 mL of 0.5 M HCl solution were added to ~50 wet g electro-dewatered samples that were mixed previously with filtrate.

For each microcosm, approximately 50 wet g of prepared biosolids were placed in 1 L media bottles that had silicon septa on their screw caps. For uniformity, bottles were rotated horizontally on a roller apparatus during incubation (Wheaton Industries Inc., Millville, NJ). The roller operated at room temperature (22 ± 0.5 °C) and was covered to avoid light exposure. To maintain anaerobic conditions during biosolids incubation, bottles were flushed daily with pure nitrogen for 10 min. Anaerobic indicator strips (BD GasPak, Franklin Lakes, NJ) were placed in the bottles to confirm the absence of oxygen. Two sets of tests lasting 7 days each were performed 2 weeks apart.

2.3. Total solids, pH, and *E. coli* counts
Standard Method No 2540-B (APHA et al., 2012) was used to measure total solids. To measure biosolids pH, ~1 wet g of sample was dispersed in 9 mL distilled water with the ULTRA-TURAX® homogenizer (Navab-Daneshmand et al., 2012), and the pH was measured using a gel-filled AgCl combination electrode (Fisher Scientific, Canada).

As described in (Navab-Daneshmand et al., 2012), a modified MPN method with Colilert reagent (IDEXX Laboratories, Westbrook, ME) was used to enumerate *E. coli* using Standard Method No. 9223 (APHA et al., 2012). Samples were serially diluted in clear 96-well microplates and positive well responses were measured by a SpectraMax® microplate reader (Molecular Devices, Sunnyvale, CA). An online MPN calculator was used to calculate MPN/g-TS (Curiale, 2004). The detection limit was determined to be 3 positive wells in the first row, corresponding to 60-170 MPN/g-TS.

2.4. PMA treatment of biosolids

Prior to DNA extraction, some of the samples were treated with propidium monoazide (PMA) to inhibit the amplification of dead or membrane compromised DNA. PMA dye penetrates cell walls and bind to the DNA bases (van Frankenhuyzen et al., 2011). When exposed to light, photolysis of the PMA dye forms a covalent bind that inhibits amplification by polymerase chain reaction (PCR) assays (Nocker and Camper, 2009). In brief, 10 mM PMA solution in 20% dimethyl sulfoxide was added to samples to achieve a final concentration of 100 µL. Samples with PMA solution were incubated in dark for 10 min to allow sufficient binding of the PMA dye to the DNA bases. Following incubation, the samples were exposed to a 600-W halogen light at a distance of 15 cm for 5 min (Taskin et al., 2011).

2.5. DNA extraction and PCR

Genomic DNA was extracted from the PMA-treated and non-PMA-treated biosolids obtained on days 0 and 7 of the experiments using PowerSoil® DNA isolation kits following manufacturer's instructions (Mo Bio Laboratories, Inc., Carlsbad, CA, USA). DNA quality and quantity were assessed using 1% agarose gel electrophoresis and spectrophotometry (260 nm/280 nm ratio). A nested PCR protocol was used to amplify 16S rRNA gene. First, DNA was amplified using universal primer pair 27F and 1492R (Weisburg et al., 1991). Next, amplicons were diluted and used as the template for a second PCR targeting the V6~V8 regions of 16S rRNA gene using primer pairs 926F and 1392R with adaptors (Engelbrektson et al., 2010). Finally, sample-specific barcodes were added to the amplicons by using primer pairs Uniprimer1 and Uniprimer2. The primers and PCR conditions are listed in Table S1. All reactions were carried out with primer concentration at 0.3 µM, Mg^{2+} 1.5 mM, dNTP 200 µM, Taq DNA polymerase 1.25 Unit/50 µl reaction (all reagents from NewEngland Biolab). The amplicons were cleaned up using QIAquick PCR purification kit (QIAGEN), then quantified using Quant-iT™ PicoGreen™ dsDNA Assay Kit (ThermoFisher Scientific).

2.6. Illumina high-throughput sequencing

The amplicon library was sequenced on Illumina MiSeq PE300 (pair-end 300 cycles) platform at McGill University and Génome Québec Innovation Centre (Montréal, QC, Canada). In total, 3,825,375 sequences were generated as de-multiplexed forward and reverse reads. The raw sequences were paired, primer-stripped, quality-filtered, chimera-removed using USEARCH algorithm (Edgar, 2010), resulting in 2,220,746 good-quality sequences. Operational Taxonomic Units (I) were clustered at 97% similarity level using open-reference protocol with the UCLUST algorithm (Edgar, 2010), assigned with 97%

similarity taxonomy from Greengenes database (McDonald et al., 2012; Werner et al., 2012), resulting in 4,026 OTUs in total.

2.7. Statistical analysis

Statistical analysis were performed using R statistical analysis software with "vegan" package (R Core Team, 2019) with a significance level of 0.05. To examine the impact of different treatment factors on the regrowth of *E. coli*, an ANOVA model was applied with pH as a linear regression factor (Sokal and Rohlf, 1995). In addition to pH, biosolids treatment (heat treatment, electro-dewatering, or electro-dewatering with added filtrate), and inoculation with untreated biosolids (with or without) were also considered. The regrowth response variable was defined as the maximum log of *E. coli* counts observed during the 7-d incubation. The pH value for statistical testing was the value observed on the day of maximum *E. coli* counts; this definition was adopted because the pH increased after a few days of incubation in most of the bottles where acid was added to biosolids. Similarly, an ANOVA was applied to analyze the impact of biosolids treatment (untreated, heat treated, electro-dewatered, or electro-dewatered with added filtrate), pH, with or without inoculation with untreated biosolids, with or without PMA treatment, and before or after 7 days of incubation on the changes in microbial community (alpha diversity). The alpha diversity indices were calculated using the number of OTUs, Hill's number derived from exponential of Shannon diversity index, and Pielou's Evenness index. The effect size estimates were obtained by omega squared (ω^2) measure using R "effectsize" package. The canonical correspondence analysis (CCA) was performed using the I table and five variables: pH, incubation time (days 0 and 7), inoculation with untreated biosolids, PMA treatment, and filtrate addition. The last three variables were binary. A subset of samples for CCA including the electro-dewatered with and without added filtrate samples were analyzed using pH, incubation time, inoculation with untreated biosolids, and filtrate addition. Heatmap of the assemblage of the top 10 genera from each sample was constructed using R "gplots" package.

3. Results and discussion

3.1. Biosolids electro-dewatering

A typical electro-dewatering cycle started with a lag in water removal of approximately 1-min. After the onset of water removal, energy consumption and filtrate removal proceeded at constant rates (Figure S1). Similar to energy consumption, the temperature of the dewatering cake also increased at an almost linear rate with time to ~100 °C. By the end of the electro-dewatering cycle, total solids increased from 18.5 ± 1.4% w/w (± indicates the range of two replicates unless otherwise specified) in untreated biosolids to 35.8 ± 0.5% w/w in electro-dewatered cakes (Figure 1a), and the pH decreased from 6.9 ± 0.5 to 4.8 ± 0.1 (Figure 1b). Furthermore, the 10-min Electro-dewatering process reduced *E. coli* counts by at least 2.8 ± 1.1 logs, from 5.0 ± 1.1 in untreated solids to the detection limit in electro-dewatered cake (Figure 1c). These results support the findings from our previous work which showed that high temperatures from Joule heating was the principal mechanism for the inactivation of bacterial pathogen indicators (Navab-Daneshmand et al., 2012).

Figure 1. (a) Total solids, (b) pH, and (c) *E. coli* concentration in untreated (UT), heat treated (HT), electro-dewatered (ED), and electro-dewatered with filtrate addition (EDF) biosolids, with or without inoculation with untreated biosolids, and with or without acid (0.5 M HCl) or base (0.5 M NaOH) addition. Measurements were obtained during 7 d incubation under anaerobic conditions. Bars represent the range of two tests performed two weeks apart. The asterisks (*) are the detection limits (60-170 MPN/g-TS) as these counts were below detection.

3.2. Total solids and pH of biosolids during incubation

Electro-dewatering process increased the total solids content of biosolids to $35.8 \pm 0.5\%$ w/w (Figure 1a). Addition of NaOH to electro-dewatered and inoculated-electro-dewatered samples reduced their solid content to $28.3 \pm 1.0\%$ w/w. Furthermore, addition of the removed filtrate to the electro-dewatered cake reduced total solids to $22.0 \pm 0.3\%$ w/w (Figure 1a). The difference in total solids between untreated and electro-dewatered solids with added filtrate is due to the loss of some of the removed water by evaporation and capillarity retention on electrodes and filtration membrane. In heat-treated controls, biosolids total solids content and pH did not change from those of the untreated solids (Figure 1a and b). In all 13 microcosms, total solids did not vary significantly during the 7-day incubation period.

Biosolids pH decreased from 6.7 ± 0.2 in untreated and heat-treated samples to 4.8 ± 0.1 in electro-dewatered biosolids (Figure 1b). Removed filtrate had a pH of 11.8 ± 0.6, and the addition of filtrate to the electro-dewatered biosolids increased the cake pH to 6.0 ± 0.1 (Figure 1b). During incubation, biosolids pH in samples without any acid or base additive varied between 0.2 and 1.1 pH units. In biosolids with acid or base additive, however, the pH was less stable during the 7-day incubation period and varied between 0.5 and 1.4 pH units (Figure 1b). There was a general increase in pH throughout the incubation period when acid was added, while after base addition 0.6 to 0.8 pH units reduction was observed on day 2 followed by a re-increase of 0.3 to 0.6 pH units.

3.3. Regrowth of *E. coli* during incubation

E. coli levels were reduced by 3-6 logs MPN/g-TS to the detection limits after both electro-dewatering and heat treatment (Figure 1c). Addition of HCl, NaOH, or removed filtrate, or the inoculation of electro-dewatered or heat-treated samples with untreated biosolids, did not significantly increase the *E. coli* levels in electro-dewatered or heat-treated solids before incubation (day 0, Figure 1c). For baseline conditions in which no filtrate, inoculum or acid/base were added, no regrowth of *E. coli* was observed during the 7-day anaerobic incubation in electro-dewatered cake. However, when electro-dewatered samples were inoculated with untreated biosolids, *E. coli* counts increased by 1.2 ± 0.1 logs (Figure 1c). While there was only 0.8 ± 0.4 logs regrowth in heat-treated samples, the inoculated heat-treated biosolids showed 4.0 ± 0.6 logs *E. coli* regrowth. In electro-dewatered biosolids with added filtrate, the inoculation also increased the regrowth from 2.3 ± 1.8 to 3.0 ± 0.2 logs. This is in line with our previous study where lower regrowth was observed in inoculated electro-dewatered biosolids compared to the inoculated heat-treated controls (Navab-Daneshmand et al., 2014a).

To test the impact of pH on *E. coli* regrowth, pH was manipulated different treatments to compare treatments and evaluate different factors independently. For this, NaOH was added to electro-dewatered biosolids, and HCl was added to heat-treated and electro-dewatered with filtrate controls, with similar pH manipulations in inoculated samples. Because of the instabilities observed in biosolids pH during the incubation period, pH was used as a continuous regression variable in ANOVA analysis rather than a measured explanatory variable. Table 1 shows details of the statistical analyses results. Low pH was shown to correlate with lower *E. coli* regrowth ($p < 0.01$). Regression analysis showed an average decrease of 0.9 ± 0.4 and 1.3 ± 0.4 ($\pm$ being the standard error of 12

samples) logs *E. coli* regrowth per unit reduction of pH in non-inoculated and inoculated biosolids, respectively; however, this difference was not significant. Lowering the pH with the addition of HCl in inoculated-heat treated, electro-dewatered with filtrate, and inoculated electro-dewatered with filtrate reduced *E. coli* regrowth by 1.0 to 2.1 logs (Figure 1c). Conversely, increasing the pH (i.e., addition of NaOH) in electro-dewatered and inoculated electro-dewatered biosolids increased regrowth by 3.3 to 3.6 logs (Figure 1c).

Table 1. Results of statistical analysis on the impact of different treatment factors on *E. coli* counts in biosolids after treatment (heat treated, electro-dewatered, or electro-dewatered with added filtrate), with or without inoculation with untreated biosolids after 7 days of incubation. Two sets of experiments (replicates) were performed two weeks apart.

Experimental Factors	Degrees of freedom	p-value	Effect size estimate (ω^2)
Treatment	2	0.85	-0.07
Inoculation	1	0.07	0.10
pH	1	**<0.01**	0.53
Treatment * inoculation	2	0.23	0.05
Treatment * pH	2	0.70	-0.06
Inoculation * pH	1	0.17	0.04

The effect of biosolids treatment on *E. coli* regrowth was not found to be statistically significant (Table 1), suggesting that both removal of nutrients with the filtrate and the possible production of inhibitory compounds did not play a major role in inhibiting regrowth. The impact of the removal of essential nutrients through filtrate can be visualized at both low and high pH levels. For high pH biosolids, inoculated electro-dewatered biosolids with NaOH addition (i.e., filtrate removed) can be compared to inoculated heat-treated controls (i.e., no filtrate removed) where similar *E. coli* regrowth occurred (6.1-6.3 logs, Figure 1c). Likewise, for low pH solids, inoculated electro-dewatered biosolids (i.e., filtrate removed) can be compared to inoculated electro-dewatered with added filtrate and acid, and inoculated heat-treated with added acid controls (i.e., no filtrate removed); there was no significant difference between the regrowth observed in these samples (3.4-3.8 logs regrowth). Therefore, the removal of filtrate does not seem to have a meaningful impact on the regrowth of *E. coli*. The impact of inhibitory compounds that were possibly produced during electro-dewatering can be visualized by comparing inoculated electro-dewatered solids with added filtrate to inoculated heat-treated controls. Although 1-log lower *E. coli* regrowth was observed in inoculated electro-dewatered with filtrate addition (Figure 1c), the pH was also 1 unit lower compared to inoculated heat-treated controls (Figure 1b). Moreover, the lower regrowth in electro-dewatered biosolids with added filtrate seems to be controlled by lower pH in these solids, and therefore, could not be linked to a production of inhibitory compounds during electro-dewatering.

Results suggest that low pH of electro-dewatered biosolids is the principal parameter controlling the regrowth of *E. coli* under the discussed experimental conditions. Electro-dewatering reduces biosolids pH to 4.8 ± 0.1, which is close to the pH minima observed for *E. coli* growth (i.e., pH = 3.9; Presser et al., 1998) (Figure 1b). The higher *E.*

coli regrowth observed in samples with higher pH values is in line with reported increase in *E. coli* growth rates above pH 4 (Presser et al., 1997). Altogether, the impact of low pH on bacterial regrowth was indicated and can be applied to other biosolids treatments. Lowering biosolids pH after treatment could be beneficial for reducing regrowth of bacterial pathogens and providing optimum growth pH for certain crops and plant types. It is known that certain crops such as berries, eggplants, and potatoes require soil pH to be between 4 and 8 (Havlin et al., 2013). If electro-dewatering can be certified as Class A, the technology could be a promising method for the land application of biosolids in terms of fecal indicator bacterial content and pH levels in certain agricultural soils.

Figure 2. Alpha diversity indices (a) number of OTUs, (b) Hill's number derived from Shannon diversity, and (c) Pielou's Evenness in untreated, heat treated, electro-dewatered (ED), and electro-dewatered with added filtrate biosolids, with and without inoculation with untreated biosolids, with or without acid (0.5 M HCl) or base (0.5 M NaOH) addition, and with and without PMA treatment. Samples were collected on days 0 or after 7 d incubation under anaerobic conditions. Associated pH values (circles) are plotted on the

secondary y-axis in plot (a). Bars represent the range of two tests performed two weeks apart.

Table 2. Results of statistical analyses – degrees of freedom (Df), p-values, and effect size estimates (ω^2) – on changes in microbial community (alpha diversity) in biosolids after treatment (untreated, heat treated, electro-dewatered, or electro-dewatered with added filtrate), with or without inoculation with untreated biosolids, with or without PMA treatment, and before or after 7 days of incubation. Two sets of experiments (replicates) were performed two weeks apart.

Experimental factors	Df	Number of OTUs		Hill's number [1]		Pielou's Evenness	
		p	ω^2	p	ω^2	p	ω^2
Treatment	3	**<0.01**	0.64	**<0.01**	0.64	**<0.01**	0.41
Inoculation	1	0.27	<0.01	0.64	–0.02	0.69	–0.02
pH	1	0.37	<0.01	0.73	–0.02	0.66	–0.02
PMA	1	**<0.01**	0.56	**<0.01**	0.26	0.10	0.04
Incubation	1	**<0.01**	0.26	**<0.01**	0.21	**0.03**	0.10
Treatment * inoculation	2	0.39	<0.01	0.15	0.05	0.63	–0.03
Treatment * pH	3	0.05	0.13	0.06	0.12	0.12	0.08
Treatment * PMA	2	**0.03**	0.13	0.10	0.07	0.57	–0.02
Treatment * incubation	2	0.39	<0.01	0.11	0.06	0.35	<0.01
Inoculation * pH	1	0.82	–0.02	0.49	–0.01	0.47	–0.01
pH * PMA	1	0.45	<0.01	0.54	–0.01	0.45	<0.01
pH * incubation	1	0.32	<0.01	0.91	–0.02	0.82	–0.02
PMA * incubation	1	0.51	–0.01	0.90	–0.02	0.88	–0.02

[1] Hill's number = exponential of Shannon diversity index

3.4. Changes in microbial community during incubation

Alpha diversity (within samples) of sequenced 16S rRNA gene amplicon data were calculated using the number of operational taxonomic units (OTUs, defined at 97% identity), Hill's number, and Pielou's Evenness index. Alpha diversity was significantly affected by the treatment of biosolids (i.e., untreated, heat treated, electro-dewatered, or electro-dewatered with added filtrate), PMA treatment, and incubation period ($p < 0.05$, Table 2), but not significantly affected by pH, inoculation or filtrate addition ($p > 0.05$). The interaction of biosolids treatment with pH and PMA treatment also showed significant effect on the number of OTUs.

PMA was used to inhibit extraction and amplification of dead or compromised DNA. Comparing microbial communities' alpha diversity between samples with and without PMA treatment, the number of OTUs and the Hill number were reduced by 41% and 53%, respectively (Figure 2). The distribution of the relative abundances of OTUs shows that the number of least-abundant OTUs were reduced in PMA-treated samples (Figure S2). Previous findings showed that majority of the diversity observed in biosolids without PMA treatment was likely due to dead cells (van Frankenhuyzen et al., 2011), suggesting that the microbial community compositions determined after PMA treatment comprise the truly regrowing species.

Variations in microbial community compositions with experimental factors were analyzed using Canonical correspondence analysis (CCA) to evaluate their correlations; this first CCA model explained 23.4% of the variations (Figure 3). Samples on day 7 treated with PMA tended to appear closer to the day 0 samples than the non-treated samples (Figure 3a), but the distance between the PMA-treated and non-treated sample was relatively small. This is consistent with the observations that PMA treatment removed populations that were present at low abundance and were likely inactive. This suggest that dead cell DNA is being detected without PMA treatment. Therefore, for DNA-based detection methods (e.g., sequencing, qPCR) when both live and dead cells are present at significant levels, PMA treatment is recommended to exclude dead cell DNA.

Figure 3. Canonical correspondence analysis (CCA) of microbial OTUs and treatment variables, pH, inoculation with untreated biosolids (Inoc.), PMA treatment, incubation time (Day), and filtrate addition (F.) in a) untreated (UT), heat-treated (HT), electro-dewatered (ED), and electro-dewatered with filtrate addition (EDF) biosolids and b) electro-dewatered with and without added filtrate. Sample replicates were from two experimental sets.

The incubation period clearly segregated community compositions between samples on day 0 and day 7, and contributed the highest explanation power (6.6%) in the CCA model (Figure 3a). This shows that specific OTUs grew in the community during incubation. The second factor explaining 6.1% of the microbial community variations was the final pH of the biosolids, which oriented in parallel to the CCA2 axis (Figure 3a). Therefore, the composition of the community after 7 days of incubation was modulated by the final pH, which led to 6.1% of the variance. The remaining important experimental factors (filtrate removal, and inoculation with untreated solids) showed similar correlations to the microbial community compositions (i.e., their vector representation point in the same direction, Figure 3a), and together accounted for 8.3% of the variations.

In order to better visualize the impact of inoculation and the addition of electro-dewatering filtrate on the community composition, a second CCA analysis was performed with only the PMA-treated ED samples. Similar to the first CCA, the incubation period and the final pH were the main explanatory factors and they explained together 20.0% of the

variations. Also, the vector projection of pH was at approximately 90° from the other explanatory factors (Figure 3b), further supporting a dominant impact of the pH on the final community profiles. The addition of ED filtrate and inoculation showed similar effects with vector projections pointed in the same direction, and they explained to 7.8% and 7.0% of the variations, respectively. This observation suggests a few possible mechanisms at play. First, it is possible that the addition of ED filtrate re-introduces live bacteria, which could be similar to the ones introduced by inoculation with untreated biosolids. Alternatively, it is possible that the filtrate and the inoculate introduce necessary nutrients that favor certain genera.

Figure 4. Heatmap of the top 10 genera from microbial communities and Pearson correlation coefficient with pH (* indicates $p < 0.05$). Samples are untreated (UT), heat-treated (HT), electro-dewatered (ED), and electro-dewatered with filtrate addition (EDF) biosolids, with or without PMA treatment. I, A and B stands for inoculation, acid and base,

respectively. Sample replicates from two experimental sets are numbered (1) and (2). Bacteria names are shown at genus level, or higher level (family: f_; order: o_; class: c_; phylum: p_; kingdom: k_) if not identified at genus level. Color key indicates relative abundance of genera in each sample. Hierarchical clusters indicate similarities among genera based on Euclidean distance method.

The interpretations of the CCA analyses were refined by performing a cluster analysis on the ensemble of the 10 most abundant genera in each microbial community. This criterion selected 42 genera that accounted for between 72.3% and 98.9% of the reads in each sample. The cluster analysis identified 4 clusters of genera (Figure 4). The last two clusters (Clusters C and D) comprised generally low abundance members of the communities in PMA treated samples on day 7. Cluster D grouped genera that were always only sporadically present in the communities. Cluster C encompassed abundant activated sludge genera and families that are either strict aerobes or respiratory facultative anaerobes with limited fermentative metabolisms (e.g., *Flavobacterium*, *Acinetobacter*, *Rhodobacter*, *Dechloromonas*, and genera in the family *Chitinophagaceae and Methylophylaceae*) (Hunter et al., 2009). These OTUs were either alive immediately after ED and heat treatments, or the cell damage was not extensive enough to allow the PMA to reach DNA. Nonetheless, they decreased in relative abundance (and likely decayed) during the 7-day anaerobic incubations for both the PMA treated and untreated samples.

Cluster A genera were the dominant populations in all biosolids sample after a 7-day anaerobic incubation irrespective of PMA treatment, but they were typically in low relative abundance on day 0, indicating their growth under anaerobic conditions. These genera included *Lactobacillus*, *Enterococcus* and other unidentified genera in the family *Enterobacteriaceae*, *Bacteroides*, and *Clostridium*. All these genera and families have highly competitive fermentative or anaerobic respiratory capabilities (Tao et al., 2020). Furthermore, they are often reported in anaerobic bioreactors (Zhang et al., 2019). Some genera in Cluster A were also abundant before the incubation (e.g., *Lawsonia* and *Oscillospira*), but these are also anaerobic bacteria found in human or animal intestines (Gophna et al., 2017). They are potentially carried by wastewater and remained in the activated sludge (Cai et al., 2014; Guo et al., 2019), indicating their ability to survive in various conditions.

Cluster B genera were generally at low abundances on day 0, and increase in abundance by day 7 of anaerobic incubation except in the sample with only ED treatment (i.e., not ED plus filtrate addition). Similar to Cluster A, the genera in Cluster B have extensive anaerobic respiration and highly competitive fermentative capabilities (Tao et al., 2020). In fact, several OTUs in Cluster B are closely related phylogenetically to OTUs in cluster A. The reduced ability of these OTUs to grow in ED treated biosolids suggest that electro-dewatering removed essential nutrient for the growth of these organisms or that inhibitory compounds that are neutralized by the return of the ED filtrate are produced.

Since pH was shown to largely affect the microbial community variance (Figure 4), the most abundant genera were tested for correlation between their relative abundances and pH values (Figure 4). A few genera were significantly negatively correlated with pH

(Pearson correlation coefficient, $p < 0.05$). *Lactobacillus* and *Acetobacter* were reported in anaerobic digestion systems and related with production of volatile fatty acids (Xu et al., 2014), suggesting their favored growth under low-pH conditions. Unclassified genera in the family *Methylobacteriaceae* and *Planococcaceae* were at very low abundances. *Methylobacteriaceae* may contribute to organic substrate degradation and trace oxygen consumption (Rui et al., 2015). The family *Planococcaceae* was reported as a dominant family in biosolids treated soils and the pH of the soil samples ranged from 5.62 to 5.83, indicating its growth capability under acidic environment (Schlatter et al., 2019).

Bacteroides, *Clostridium*, o_OPB72, f_*Pirellulaceae*, *Epulopiscium*, f_*Peptostreptococcaceae*, *Coprococcus* showed significant positive correlation with pH, most of which were reported in basic conditions. *Bacteroides* and *Clostridium* are core members found in many anaerobic digestion systems (Tao et al., 2020), which are typically maintained at neutral or slightly basic pH conditions. O_OPB72 was isolated from Yellowstone hot spring sediments with a pH of 7.6 at 20 ℃ (Hugenholtz et al., 1998). Some *Pirellulaceae* strains were isolated from natural environment at pH > 7 (Dedysh et al., 2020; Kumar et al., 2021). *Peptostreptococcaceae* are fermentative anaerobes with optimum pH 7.5-8.3, except for the genus *Tepidibacter* (6.0-7.3) (Rosenberg et al., 2014). These results indicate that pH can affect the relative abundance of dominant OTUs due to their different optimum pH conditions for growth.

4. Conclusions

Mechanisms of *E. coli* regrowth and changes in microbial communities in electro-dewatered biosolids were investigated in a microcosm study. The impacts of pH, removed filtrate, and/or produced inhibitory compounds on *E. coli* regrowth and microbial communities were investigated. Results indicate pH as the primary parameter that controls *E. coli* regrowth in electro-dewatered biosolids. Analyses of high throughout sequencing showed lower alpha-diversities associated with electro-dewatering treatment and incubation time. Biosolids pH and incubation period were the main factors contributing to variations in microbial community compositions. Findings highlight the role of biosolids pH on both regrowing culturable bacteria and overall microbial communities.

Acknowledgements
This research was funded by the Natural Sciences and Engineering Research Council of Canada's Collaborative Research and Development program and Ovivo. We thank Frederic Biton (Ovivo), Bruno Desmarais (Ovivo), Alain Silverwood (Ovivo), Céline Gagnon (Aquatech), and Gilbert Samson (Régie d'Assainissement desEaux du Bassin LaPrairie) for their technical support and useful comments.

References
APHA, AWWA, WEF, 2012. Standard Methods for Examination of Water and
      Wastewater. 22nd ed., American Public Health Association, Washington, DC, USA.

Cai, L., Ju, F., Zhang, T., 2014. Tracking human sewage microbiome in a municipal wastewater treatment plant. Appl. Microbiol. Biotechnol. 98, 3317–3326. https://doi.org/10.1007/s00253-013-5402-z

Chen, Y.-C., Murthy, S.N., Hendrickson, D., Araujo, G., Higgins, M.J., 2011. The effect of digestion and dewatering on sudden increases and regrowth of indicator bacteria after dewatering. Water Environ. Res. 83, 773–783. https://doi.org/10.2175/106143010X12851009156727

Curiale, M., 2004. MPN Calculator, Build 24.

Dedysh, S.N., Kulichevskaya, I.S., Beletsky, A.V., Ivanova, A.A., Rijpstra, W.I.C., Damsté, J.S.S., Mardanov, A.V., Ravin, N.V., 2020. Lacipirellula parvula gen. nov., sp. nov., representing a lineage of planctomycetes widespread in low-oxygen habitats, description of the family Lacipirellulaceae fam. nov. and proposal of the orders Pirellulales ord. nov., Gemmatales ord. nov. and Isosphaerales ord. nov. Syst. Appl. Microbiol. 43, 126050. https://doi.org/10.1016/j.syapm.2019.126050

Dentel, S.K., Qi, Y., Herson, D.S., 2008. Improving the assessment of risk from pathogens in biosolids: fecal coliform regrowth, survival, enumeration, and assessment. WATER Sci. Technol. 57, 189–193. https://doi.org/10.2166/wst.2008.005

Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26, 2460–2461. https://doi.org/10.1093/bioinformatics/btq461

Engelbrektson, A., Kunin, V., Wrighton, K.C., Zvenigorodsky, N., Chen, F., Ochman, H., Hugenholtz, P., 2010. Experimental factors affecting PCR-based estimates of microbial species richness and evenness. ISME J. 4, 642–647. https://doi.org/10.1038/ismej.2009.153

Gardner, J., Oermeci, B., 2010. Effect of aging, time, and temperature on fecal coliform counts during centrifugal dewatering and role of centrate in growth inhibition. WATER Environ. Res. 82, 51–61. https://doi.org/10.2175/106143009X442943

Gophna, U., Konikoff, T., Nielsen, H.B., 2017. Oscillospira and related bacteria – From metagenomic species to metabolic features. Environ. Microbiol. 19, 835–841. https://doi.org/10.1111/1462-2920.13658

Guo, B., Liu, C., Gibson, C., Frigon, D., 2019. Wastewater microbial community structure and functional traits change over short timescales. Sci. Total Environ. 662, 779–785. https://doi.org/10.1016/j.scitotenv.2019.01.207

Guo, B., Manchester, M., Luby, T., Frigon, D., 2018. Composition of heterotrophic specialized sub-guilds defined by a positive RNA and polyhydroxyalkanoate correlation in activated sludge. Water Res. 144, 561–571. https://doi.org/10.1016/j.watres.2018.07.059

Havlin, J.L., Beaton, J.D., Tisdale, S.L., Nelson, W.L., 2013. Soil Fertility and Fertilizers: An Introduction to Nutrient Management. Pearson Education, New Jersey, NY.

Higgins, M.J., Chen, Y.-C., Murthy, S.N., Hendrickson, D., Farrel, J., Schafer, P., 2007. Reactivation and growth of non-culturable indicator bacteria in anaerobically digested biosolids after centrifuge dewatering. Water Res. 41, 665–673. https://doi.org/10.1016/j.watres.2006.09.017

Hugenholtz, P., Pitulle, C., Hershberger, K.L., Pace, N.R., 1998. Novel division level bacterial diversity in a Yellowstone hot spring. J. Bacteriol. 180, 366. https://doi.org/10.1128/JB.180.2.366-376.1998

Hunter, C.N., Daldal, F., Thurnauer, M.C., Beatty, J.T., 2009. The Purple Phototrophic Bacteria, Part of the Advances in Photosynthesis and Respiration book series, volume 28. AIPH, Springer, Dordrecht.

Koutsoumanis, K., Kendall, P., Sofos, J., 2004. Modeling the boundaries of growth of *Salmonella typhimurium* in broth as a function of temperature, water activity, and pH. J. Food Prot. 67, 53–59. https://doi.org/10.4315/0362-028X-67.1.53

Kumar, D., Kumar, G., Jagadeeshwari, U., Sasikala, C., Ramana, C.V., 2021. "Candidatus Laterigemmans baculatus" gen. nov. sp. nov., the first representative of rod shaped planctomycetes with lateral budding in the family Pirellulaceae. Syst. Appl. Microbiol. 44, 126188. https://doi.org/10.1016/j.syapm.2021.126188

McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., DeSantis, T.Z., Probst, A., Andersen, G.L., Knight, R., Hugenholtz, P., 2012. An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. ISME J. 6, 610–618. https://doi.org/10.1038/ismej.2011.139

Navab-Daneshmand, T., Beton, R., Hill, R.J., Frigon, D., 2015. Impact of Joule heating and pH on biosolids electro-dewatering. Environ. Sci. Technol. 49, 5417–5424.

Navab-Daneshmand, T., Beton, R., Hill, R.J., Gehr, R., Frigon, D., 2012. Inactivation mechanisms of bacterial pathogen indicators during electro-dewatering of activated sludge biosolids. Water Res. 46, 3999–4008.

Navab-Daneshmand, T., Enayet, S., Gehr, R., Frigon, D., 2014a. Bacterial pathogen indicators regrowth and reduced sulphur compounds' emissions during storage of electro-dewatered biosolids. Chemosphere 113, 109–115. https://doi.org/10.1016/j.chemosphere.2014.04.012

Navab-Daneshmand, T., Gul-E-Hina, Biyela, P., Frigon, D., 2014b. Regrowth of bacterial pathogen indicators after advanced electro-osmotic dewatering of biosolids. Int. J. Environ. Waste Manag. 14, 134–148. https://doi.org/10.1504/IJEWM.2014.064083

Nocker, A., Camper, A.K., 2009. Novel approaches toward preferential detection of viable cells using nucleic acid amplification techniques. Fems Microbiol. Lett. 291, 137–142. https://doi.org/10.1111/j.1574-6968.2008.01429.x

Presser, K., Ratkowsky, D., Ross, T., 1997. Modelling the growth rate of *Escherichia coli* as a function of pH and lactic acid concentration. Appl. Environ. Microbiol. 63, 2355–2360.

Presser, K.A., Ross, T., Ratkowsky, D.A., 1998. Modelling the growth limits (growth/no growth interface) of *Escherichia coli* as a function of temperature, pH, lactic acid concentration, and water activity. Appl. Environ. Microbiol. 64, 1773. https://doi.org/10.1128/AEM.64.5.1773-1779.1998

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Reissbrodt, R., Rienaecker, I., Romanova, J.M., Freestone, P.P.E., Haigh, R.D., Lyte, M., Tschäpe, H., Williams, P.H., 2002. Resuscitation of *Salmonella enterica* serovar typhimurium and enterohemorrhagic *Escherichia coli* from the viable but nonculturable state by heat-stable enterobacterial autoinducer. Appl. Environ. Microbiol. 68, 4788–4794. https://doi.org/10.1128/aem.68.10.4788-4794.2002

Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F., 2014. The Prokaryotes, Firmicutes and Tenericutes. Springer-Verlag Berlin Heidelberg.

Rui, J., Li, J., Zhang, S., Yan, X., Wang, Y., Li, X., 2015. The core populations and co-occurrence patterns of prokaryotic communities in household biogas digesters. Biotechnol. Biofuels 8, 158. https://doi.org/10.1186/s13068-015-0339-3

Saveyn, H., Van der Meeren, P., Pauwels, G., Timmerman, R., 2006. Bench- and pilot-scale sludge electrodewatering in a diaphragm filter press. Water Sci. Technol. 54, 53–60. https://doi.org/10.2166/wst.2006.725

Schlatter, D.C., Paul, N.C., Shah, D.H., Schillinger, W.F., Bary, A.I., Sharratt, B., Paulitz, T.C., 2019. Biosolids and tillage practices influence soil bacterial communities in dryland wheat. Microb. Ecol. 78, 737–752. https://doi.org/10.1007/s00248-019-01339-1

Small, P., Blankemhorn, D., Welty, D., Zinser, E., Slonczewski, J., 1994. Acid and base resistance in *Escherichia coli* and *Shigella flexenri* - role of rpos and growth pH. J. Bacteriol. 176, 1729–1737. https://doi.org/10.1128/jb.176.6.1729-1737.1994

Sokal, R.R., Rohlf, F.J., 1995. Biometry: The Principles and Practice of Statistics in Biological Research. 3rd ed., W. H. Freeman and Co., New York, NY, USA.

Tao, Y., Ersahin, M.E., Ghasimi, D.S.M., Ozgun, H., Wang, H., Zhang, X., Guo, M., Yang, Y., Stuckey, D.C., van Lier, J.B., 2020. Biogas productivity of anaerobic digestion process is governed by a core bacterial microbiota. Chem. Eng. J. 380, 122425. https://doi.org/10.1016/j.cej.2019.122425

Tapia, M.S., Alzamora, S.M., Chirife, J., 2008. Effects of Water Activity (aw) on Microbial Stability: As a Hurdle in Food Preservation, in: Water Activity in Foods: Fundamentals and Applications. Blackwell Publishing, Oxford, UK.

Taskin, B., Gozen, A.G., Duran, M., 2011. Selective quantification of viable *Escherichia coli* bacteria in biosolids by quantitative PCR with propidium monoazide modification. Appl. Environ. Microbiol. 77, 4329–4335. https://doi.org/10.1128/AEM.02895-10

Tuan, P.-A., Jurate, V., Mika, S., 2008. Electrodewatering of sludge under pressure and non-pressure conditions. Environ. Technol. 29, 1075–1084. https://doi.org/10.1080/09593330802180294

US-EPA, 2003. Environmental regulations and technology: Control of pathogens and vector attraction in sewage sludge. U.S. Environmental Protection Agency, Office of Research and Development, Cincinnati, OH.

van Frankenhuyzen, J.K., Trevors, J.T., Lee, H., Flemming, C.A., Habash, M.B., 2011. Molecular pathogen detection in biosolids with a focus on quantitative PCR using propidium monoazide for viable cell enumeration. J. Microbiol. Methods 87, 263–272. https://doi.org/10.1016/j.mimet.2011.09.007

Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA amplification for phylogenetic study. J. Bacteriol. 173, 697. https://doi.org/10.1128/jb.173.2.697-703.1991

Werner, J.J., Koren, O., Hugenholtz, P., DeSantis, T.Z., Walters, W.A., Caporaso, J.G., Angenent, L.T., Knight, R., Ley, R.E., 2012. Impact of training sets on classification of high-throughput bacterial 16s rRNA gene surveys. ISME J. 6, 94–103. https://doi.org/10.1038/ismej.2011.82

Xu, S., Selvam, A., Karthikeyan, Obuli.P., Wong, J.W.C., 2014. Responses of microbial community and acidogenic intermediates to different water regimes in a hybrid solid anaerobic digestion system treating food waste. Spec. Issue Adv. Biol. Treat.

Technol. Sustain. Waste Manag. ICSWHK2013 168, 49–58.
https://doi.org/10.1016/j.biortech.2014.04.090

Zhang, L., Guo, B., Zhang, Q., Florentino, A., Xu, R., Zhang, Y., Liu, Y., 2019. Co-digestion of blackwater with kitchen organic waste: Effects of mixing ratios and insights into microbial community. J. Clean. Prod. 236, 117703.
https://doi.org/10.1016/j.jclepro.2019.117703