Microbial habitability of early Mars lacustrine environments sustained by iron redox cycling

Rachel A. Moore[1]* and Christopher E. Carr[1,2]

Affiliations:

[1] School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, 30332, USA.

[2] Daniel Guggenheim School of Aerospace Engineering, Georgia Institute of Technology, Atlanta, GA, 30332, USA.

*Correspondence to: rmoore305@gatech.edu. Address: Weber Space Science and Technology Building, Room 210D, 275 Ferst Dr NW, Atlanta, GA 30313, USA

One Sentence Summary: This work provides an existence proof of habitability on early Mars and demonstrates modeling processes by which the habitability of extraterrestrial environments can be explored quantitatively.

Preprint: A preprint is available at bioRxiv.

Word Counts: Abstract (117), Main Text (3524)

Abstract

Several studies have reported new data on the estimated compositions of chemical components at Gale crater; however, there is still a lack of information regarding potential past support of biomass and detectable biomarkers of ancient life. In this study we evaluate microbial habitability of early Mars constrained by the recently reconstructed water chemistry at Gale. The modeled community is based on Fe-metabolizing bacteria with the ability to utilize solid-phase iron oxides (e.g., magnetite) as an electron source or sink. Our results illustrate the plausibility of a sustained community in Gale Lake and provides suggestions for future modelled and laboratory-based studies to further evaluate the past habitability of Mars, biosignatures and their preservation potential, and hidden metabolic potential.

1. Introduction

The modern surface of Mars is analogous to Earth's stratosphere: dry, cold, and inundated with DNA-damaging radiation. However, Mars possesses all the critical elements required for life including the essential elements (i.e., C, H, N, O, P, and S), and plausible sources of energy. It is possible that life could be present today in the Mars subsurface, beneath the cryosphere (Jones, Lineweaver and Clarke, 2011). Ancient Mars, with its higher abundance of liquid water, was arguably more hospitable to life as we know it, even while surface fluvial activity was likely transient, and temperatures generally cold (Wordsworth *et al.*, 2021). For instance, Gale crater, a late Noachian to early Hesperian-aged crater (3.5-3.8 Gya), once harbored a liquid water lake (Grotzinger *et al.*, 2015; Figure 1).

Figure 1. A. Aerial view of Gale Crater (137.4°E, −4.6°N) with star representing *Curiosity*'s Bradbury landing site. **B.** Profiles of Gale Crater: ancient (solid line) and modern (dashed line). The water level is not to scale. Modified from Borlina, Ehlmann and Kite, 2015. **C.** Geologic timeline of Gale Crater starting with its formation (LN: Late Noachian, EH: Early Hesperian, LH: Late Hesperian, A: Amazonian). Modified from Deit *et al.*, 2013.

Recently, samples collected by the Mars Science Laboratory rover *Curiosity* were used

to estimate Gale Lake water chemistry (Fukushi et al., 2019). Major ions (Na$^+$, K$^+$, Mg^{2+},

Ca^{2+}, Cl$^-$, SO$_4^{2-}$, HCO^{3-}) and trace (e.g., Zn, Mg, Mn, Cu, Fe) components were

constrained, in addition to pH and dissolved inorganic carbon (CO$_2$). Taken together,

this information provides rigorous constraints for modeling microbial habitability of Gale

Lake.

Compatible microbial redox metabolisms have been suggested, however, they have yet

to be experimentally assessed in an ancient Martian context (Cockell 2020). For

example, several chemo-autotrophic and -heterotrophic redox couples would have been

available for metabolism on the surface of Mars. Iron and sulfate ions have both been

detected (Fukushi *et al.*, 2019) and can be biologically reduced in conjunction with

hydrogen. In addition, there is enough light energy and inorganic carbon to support

anoxic photoautotrophy (Cockell, 2020). Phototrophy could have been productive in the

"photic zone" of Gale Lake where solar energy was abundant and damaging radiation

was attenuated (Cockell 2020; Figure 1 B).

This study approaches the possibility of maintaining a microbial community in Gale Lake

established upon energy from light and iron redox cycling. The potential habitability of

Mars and growing body of data assembled from *Curiosity* is the main reason for the

choice of Gale crater as our study site. Our work proposes the use of *Geobacter*

sulfurreducens and *Rhodopseudomonas palustris* as a simple biological community

model to better understand habitability and production of detectable biomarkers in an

inferred ancient lacustrine environments of Mars. Our work does not preclude other

metabolisms, but provides an existence proof of habitability on early Mars and a

process by which habitability of extraterrestrial environments can be explored

quantitatively.

75 2. Materials and Methods

2.1. Development of genome-scale metabolic models

Previously reconstructed and validated genome-scale metabolic models of *R. palustris*

(*iRpa940*; Alsiyabi, Immethun and Saha, 2019) and *G. sulfurreducens* (*iRM588*;

Mahadevan *et al.*, 2006) were edited before their inclusion in community models.

Several reactions for phototrophic growth in *R. palustris* were de-simplified based on the

PNSB2011 model from Hädicke, Grammel and Klamt (2011) in order to more accurately

track reaction fluxes. In addition, rxn01257, which produces precursors for folate and

aromatic amino acids, was added to *iRpa940* after gapfilling in KBase to allow for

growth in Gale Lake minimal medium (Section 2.3). The enzyme associated with this

reaction (i.e., aminodeoxychorismate synthase) has been identified in other

Rhodopseudomonas spp. (e.g., #A0A1H8TD11).

The gapfilling process itself did not result in any additional added reactions to the model

of *R. palustris*, however, several reaction directions were changed to be unidirectional

(Supplemental file 1). Three reactions were added to *G. sulfurreducens* (i.e., rxn05291,

rxn00197, rxn00257) during gapfilling. Reactions rxn00197, rxn00257, and rxn05291

are part of the Krebs cycle which is known to be present in *G. sulfurreducens* (Zhang *et*

al., 2020). In both models, reactions were expanded upon to model iron redox cycling

(Supplemental file 1), which has been previously demonstrated in this specific set of two

organisms (Byrne *et al.*, 2015). Community models were created using COBRApy's

merge function with objective set to sum (Supplemental file 1). Separate cytoplasmic

compartments were maintained by annotation of the reactions and metabolites for each

organism. The community model is represented in Figure 2.

Figure 2. Model of Gale Lake emphasizing iron redox cycling phenomena.

2.2 Construction of the diurnal community model

A diurnal model was constructed, based on the diuFBA model from Knies et al. 2019,

using the community model of *R. palustris* and *G. sulfurreducens* in section 2.1 as a

scaffold. Briefly, the individual metabolic models were duplicated, and a separate

extracellular space was defined ("night", e2). Photon flux was only allowed in the "day"

model (e0). An ATP maintenance reaction (rxn00062) was added to both organisms so

that they would have to realistically expend energy even when not prioritizing growth.

The reaction rates were based upon how much energy each organism spends on non-

growth-associated maintenance (Klamt, Schuster and Gilles, 2002; Mahadevan *et al.*,

2006).

The day and night models were connected via transfer reactions that allow the "storage"

and use of any metabolite, apart from photons and protons, in a different period other

than when it was produced (Equation 1).

113   [1]

114   The transfer reactions are bidirectional, as a microorganism can theoretically store

115   metabolites from night and use them during the day, as well as store from day to use at

116   night. This is illustrated in Figure 3.

117
118   **Figure 3.** Schematic of the diurnal community model.

A single biomass reaction occurs for *R. palustris* in "Day", and in "Night" for *G.*

sulfurreducens to account for organism growth based upon when it would be prioritized

(Byrne *et al.*, 2015). Both reactions are summed to serve as the overall objective

function (Fig. 3). Flux values were integrated over 12 hours for both light and dark

periods to estimate the total amount of each metabolite during one diurnal cycle (24-

hour period) as in equation 2:

125   [2]

where v is the flux of a given metabolite (mmol gDW^{-1} h^{-1}), and Δt is the duration of

either the light or dark period (12 h). The result, Δc, corresponds to the molar amount of

a given metabolite produced in the light or dark period (mmol gDW^{-1}; Knies *et al.*, 2015).

2.3 Formulation of a Gale Lake minimal medium

An *in-silico* medium (Table 1) was simulated based on the methods of Marinos *et al.*

(2020) and the composition of reconstructed water chemistry and evidence of trace

minerals (Fukushi *et al.*, 2019; Payré *et al.*, 2019). Copper and zinc deposits within Gale

crater were likely adsorbed onto Mn-oxides and thus insoluble (Payré *et al.*, 2019). For

this reason, the bounds for copper and zinc were based on the optimal rate of import

from the external environment for the community model. The remaining metabolites

(Table 1) were treated as soluble within the lake system. The proton flux (H+) reflects

the neutral pH of Gale Lake proposed by Fukushi *et al.* (2019). Concentrations of cobalt

and magnesium were calculated from the *in situ Spirit* and *Opportunity* rovers'

measurements reported in Payré *et al.* (2019) as follows:

$$\frac{PPM(\sim \frac{mg}{kg} \, sediment) \, \times total \, kg \, sediment}{total \, kg \, water} = mg \, kg^{-1} \, water$$

The total mass (6.67×10^6 kg) of sediment in Gale was derived from the estimated lake

area (2×10^{10}; Fukushi *et al.*, 2019) and the average bulk density of Martian sediment

(3×10^3 kg m^{-3}). Total water mass (3×10^{15} kg) was the lower bound estimated in Fukushi

et al. (2019). The upper limit concentration of N$_2$ was calculated using the ideal gas law,

a maximum partial pressure of 0.3 bar (Von Paris *et al.,* 2013; Navarro-Gonzalez et al

2018), a temperature of 273.15 K (Fukushi *et al.*, 2019) and a Martian atmospheric

mass of 2.5×10^{16} kg (*Mars Fact Sheet*, 2021). The lower bound for N$_2$ was based on the

optimal rate of import from the external environment. Water was allowed to freely enter

the extracellular environment at 1000 mmol gDW^{-1} h^{-1}.

Table 1. Estimated chemical composition of Gale Lake and corresponding bounds
utilized in the modelled growth medium.

Metabolite / SEED compound ID	Concentration $(mol\ L^{-1})$	Flux upper bound $(mmol\ gDW^{-1}\ h^{-1})$
Ca^{2+} / cpd29674	$2.4\text{-}4.5\times10^{-2}$ *	24 - 45
Cl^- / cpd00099	$1.1\text{-}2.5\times10^{-1}$ *	110 - 250
Co^{2+} / cpd00149	$1.0\text{-}5.9\times10^{-6}$ ****	$1.0\text{-}5.9\times10^{-3}$
CO_2 / cpd00011	$2.3\text{-}41\times10^{-3}$ *	2.3 - 41
Cu^{2+} / cpd00058	1.0×10^{-7} ***	1.0×10^{-4}
Fe^{2+} / cpd10515	$1.2\text{-}58\times10^{-4}$ *	$1.2\text{-}58\times10^{-1}$
H^+ / cpd00067	1.0×10^{-6} **	1.0×10^{-3}
H_2 / cpd11640	1.2×10^{-2} ***	12
H_2O / cpd00001	1.0	1000
K+ / cpd00205	$1.4\text{-}4.4\times10^{-3}$ *	1.4 – 4.4
Mg / cpd00254	$3.5\text{-}6.0\times10^{-2}$ *	35 - 60
Mn^{2+} / cpd00030	$5.3\text{-}100\times10^{-7}$ ****	$5.3\text{-}100\times10^{-4}$
N_2 / cpd00528	$130\text{-}220\times10^{-5}$ *****	$1300\text{-}2200\times10^{-2}$
Na+ / cpd00971	$9.4\text{-}12\times10^{-2}$ *	94-120
PO_4^{3-} / cpd00009	$5.2\text{-}18000\times10^{-7}$ ***	$5.2\text{-}18000\times10^{-4}$
SO^{2-}_4 / cpd00048	$4.4\text{-}7.2\times10^{-2}$ *	44-72
Zn^{2+} / cpd00034	1.0×10^{-7} ***	1.0×10^{-4}

* Data from Fukushi *et al.* (2019) total dissolved components of calcium, chloride,
carbon dioxide, iron, hydrogen, potassium, magnesium, sodium, and sulfate,
respectively.
** Calculation based on data from Fukushi *et al.* (2019)
*** Concentration and flux based on model requirements.
**** Calculation based on data from Lanza *et al.* (2014) and Berger *et al.* (2019)
***** Calculation based on Von Paris *et al.,* (2013) and Navarro-Gonzalez et al (2018)
158

For models with photoautotrophic growth, photons were allowed to enter the external

environment at a rate of 100 mmol gDW^{-1} hr^{-1} (9.7 µE m^{-2} s^{-1}).

2.5 Flux balance analysis

The metabolic flux distribution of the individual, community, and diurnal models was

calculated using flux balance analysis or dynamic flux balance analysis. The specific

growth rate was used as the objective function (function to be maximized) in all

simulations except for those where maximum yields of hopanoids were estimated as in

Orth, Thiele and Palsson (2010). All simulations were performed using the Python

programming language (version 3.8) in conjunction with open-source GLPK software

(https://www.gnu.org/software/glpk/), and COBRApy (Ebrahim *et al.*, 2013), pandas

(Reback *et al.*, 2020), and NumPy (Harris *et al.*, 2020) libraries.

3. Results

3.1 Genome-scale metabolic models

Table 2 shows the specific growth rate for each model on the different types of

substrates with or without light. Although *R. palustris* is capable of chemotrophic growth

in complete media (Alsiyabi, Immethun and Saha, 2019), it fails to grow in the modelled

medium without the presence of light (Table 2) or a source of organic carbon. Further,

all other models grow slowly at low levels of metabolite flux in the medium (0.001 h^{-1};

Table 2). At higher levels of metabolite flux, community growth is roughly two orders of

magnitude faster than all other models (Table 2). For example, growth of the community

in the presence of light (1.031 h^{-1}) increased ~2000% as compared to growth of the

individual models combined (i.e., 0.005 and 0.032 h^{-1} or 0.037 h^{-1} total).

Table 2. Specific growth rates of the genome-based models under the specified
conditions and constrained to either the low or high bounds of medium metabolite flux.

Electron donor/acceptors	Input carbon source	Light?	Model	Specific growth Rate (h^{-1})
H_2 / Fe	CO_2	NA	*G. sulfurreducens*	0.001–0.005
Fe / CO_2	CO_2	-	*R. palustris*	0.00–0.00
Fe / CO_2	CO_2	+	*R. palustris*	0.001–0.032
H_2, Fe / Fe, CO_2	CO_2	+	Community	0.001–1.031
H_2, Fe / Fe, CO_2	CO_2	-	Community	0.001–1.012

NA: Not applicable

Syntrophic interactions were investigated and are presented in Table 3. Both *R.*

palustris and *G. sulfurreducens* produce metabolic products consumed by the other

member. For example, carbon sources (acetate and citrate) were produced and

exported to the extracellular environment where "cross-feeding" can occur. Acetate

produced by *G. sulfurreducens* was consumed by *R. palustris* at a rate of 142.5 mmol

gDW^{-1} h^{-1} when grown in the higher range of medium fluxes. Likewise, citrate was

produced at a rate of 63.29 mmol gDW^{-1} h^{-1} by *R. palustris* and consumed by *G.*

sulfurreducens (Table 3). Iron (Fe^{2+} and Fe^{3+}) was consumed and produced at the same

rate (4.38 mmol gDW^{-1} h^{-1}) between the microbes and the extracellular environment

(Table 3).

Table 3. Modelled interactions between community members[*].

Metabolite	*R. palustris* (reaction ID)	*G. sulfurreducens* (reaction ID)
Acetate	-0.0053 / -142.5 (rxn05488)	0.0053/ 142.5 (rxn05488)
Fe^{2+}	-0.022 / -4.38 (rxn37614_2)	0.022 / 4.38 (FERCYT)
Fe^{3+}	0.022 / 4.38 (rxn37614_2)	0.022 / -4.38 (FERCYT)
Citrate	-0.0037 / 63.29 (rxn05211)	0.0037 / -63.29 (rxn05211)
CO_2	0.0072 / -94.55 (rxn05467)	-0.22 / 55.41 (rxn05467)

[*] Fluxes are shown in mmol gDW^{-1} h^{-1} for low and high range of medium inputs with light. Negative values
indicate metabolite consumption, whereas positive values indicate production.

3.1.1 Chemolithoautotrophic metabolism of G. sulfurreducens

Flux balance analysis revealed an unexpected hidden autotrophy, namely that *G.*

sulfurreducens was capable of carbon dioxide fixation through the reverse Krebs cycle

(Fig. 4). Here citrate (Table 3) was used (rxn00257) to start the cycle. CO_2 was fixed at

two steps, from succinyl-CoA to 2-oxoglutarate (Low-high = 0.27-43.31 mmol gDW^{-1} h^{-1};

rxn00197) and from 2-oxoglutarate to isocitrate (Low-high = 0.075-30.3 mmol gDW^{-1} h^{-1};

rxn00198).

Figure 4. The reverse Krebs cycle of *G. sulfurreducens* uncovered by flux balance
analysis. Numbers in parentheses refer to reaction fluxes (mmol gDW^{-1} h^{-1}, rounded to
nearest whole number). The arrows refer to the direction in which each reaction is
taking place.

3.2 Dynamic flux balance analysis

Dynamic growth of the community model based on nitrogen (Fig. 5), the limiting

metabolite, was used to calculate the total biomass supported per mL of Gale Lake in

the photic zone. Nitrogen was considered limiting for growth in this context as all other

necessary metabolites were capable of being recycled or produced following initial

input. Independent of initial media metabolite flux (i.e., high or low), 0.23 grams of

biomass were produced per gram of N_2 (Fig. 5). Depending on the amount of nitrogen

present (1.3 – 2.2 mmol N_2 L^{-1}), Gale Lake could have maintained 10^3-10^7 cells per mL

of brine, assuming cells are 10^{-12} grams in weight on average (Bratbak and Dundas,

1984; see supplemental materials).

220
Figure 5. Production of community biomass (solid lines) and consumption of N$_2$
(dashed lines) with low (black) or high (red) flux of metabolites in the medium.

3.3 Hopanoid production

To quantify hopanoid production, the production of precursor farnesyl diphosphate was

used as the objective function. Nitrogen was limited to 1 mmol gDW^{-1} h^{-1} to predict

hopanoid maximum yields per mmol of limiting substrate (N$_2$). Farnesyl diphosphate is

produced at 0.001 mmol per mmol of N$_2$. If, as estimated, Gale Lake contained 13 mmol

N$_2$ L^{-1}, we predict ~0.01 mmol of hopanoid was produced per liter in the photic zone.

This translates to a range of 2×10^{-1} - 2×10^{-2} grams of hopanoid kg^{-1} of Gale Crater

sediment assuming a photic zone depth of 100 m and a sediment thickness of 1-10 m

(see supplemental materials), assuming a preservation event (such as settling after a

mass death event) of 100% of the cells and uniform mixing of settled material within the

sediment.

3.4 Diurnal community model

238
Figure 6. Metabolite concentrations summed by category for 12-hour periods of light
and dark. Circles and triangles represent *G. sulfurreducens* and *R. palustris*,
respectively.

The overall growth rate of the diurnal community model was twofold greater than the

traditional community model (with light, 2.062 h^{-1} vs 1.031 h^{-1}). In addition, *G.*

sulfurreducens produced more biomass (1.998 h^{-1}) as compared to *R. palustris* (0.0645

h^{-1}). Overall, concentrations of metabolites (i.e., consumed and secreted) did not

change significantly between periods of light and dark except for nucleic acids

($7.61{\times}10^3$ vs. $1.27{\times}10^4$) and vitamins and cofactors ($9.74{\times}10^3$ vs. $1.26{\times}10^4$) for *R.*

palustris (Fig. 6, Supplemental File 2). Transfer reaction fluxes indicated that both

organisms prioritized the period in which they were growing. For instance, 5 out of the 8

metabolite categories were associated with a negative transfer flux for *R. palustris*

indicating transfer from night to day (Fig. 7). Likewise, 6 out of the 8 categories were

associated with positive flux values for *G. sulfurreducens* (Fig. 7).

3.4.1 R. palustris transfer reaction fluxes and ATP consumption

254
Figure 7. Metabolites transfer fluxes between light and dark summed by category.
Negative values indicate transfer from night to day, and positive values indicate transfer
from day to night.

The nucleic acid and energy metabolite flux profiles for *R. palustris* (Fig. 7) were

governed mainly by thymidine triphosphate (TTP) and guanosine triphosphate (GTP),

respectively. TTP and GTP were transferred from light to dark where they served as

phosphate donors to produce ATP from ADP (Fig. 7). Carbohydrates (i.e., 1,3-

Bisphospho-D-glycerate and glycerone) were primarily transferred from night to day in

R. palustris (Fig. 7). These were ultimately used to generate NADP and ATP through

the production of phosphate. Inorganic ions like thiosulfate were transferred to daytime

where they were used to create acetate and homocysteine, and to reduce cytochrome

c3. Out of the organic acids, oxaloacetate was transferred between night and day (Fig.

7). Oxaloacetate was primarily used to power the daytime Krebs cycle.

3.4.2 G. sulfurreducens transfer reaction fluxes and ATP consumption

Carbohydrates, inorganic ions, and organic acids were transferred at nonzero fluxes

between night and day for *G. sulfurreducens* (Fig. 7). Of the carbohydrates, acetyl

phosphate and citrate are transferred to night where they are used to make ATP.

Specifically, citrate is used to power the reverse Krebs cycle (Fig. 4). The inorganic ion

flux is dominated primarily by phospshate. Phosphate is transferred from night to day,

where it is used to produce acetylphosphate. Acetylphosphate is used to generate ATP

in both day and night. Any acetate (organic acids) produced is secreted to the

extracellular space.

4. Discussion

In this work we evaluate the microbial habitability of early Mars' lacustrine environment

constrained by reconstructed Gale Crater water chemistry and sustained by the redox

cycling of iron. In this context, we used *G. sulfurreducens* and *R. palustris* as model

organisms because they have been shown to effectively and anoxically use solid-phase

iron oxides (Byrne *et al.*, 2015). Although these specific organisms have been

successfully co-cultured, this is not a known natural association. However, similar

microbial communities have been isolated *in natura*. For example, an Fe(II)-oxidizing

photoferrotroph, *Candidatus Chlorobium masyuteum*, and a putative Fe(III)-reducing

bacterium, *Candidatus Pseudopelobacter ferreus*, were recently co-enriched from the

ferruginous Brownie Lake in Minnesota, USA (e.g., Lambrecht *et al.*, 2021). Acetate and

formate detected in the chemocline of Brownie Lake suggests that the redox cycling of

Fe occurs between these organisms, but it has not yet been confirmed. Until recently, it

was thought that no single organism was capable of both Fe(II)-oxidation and Fe(III)-

reduction. It is now understood that the bacterium *Rhodoferax* strain MIZ03 can utilize

iron as an electron donor and acceptor (Kato and Ohkuma, 2021).

In an evolutionary context, Alpha- and Delta-proteobacteria like *R. palustris* and *G.*

sulfurreducens, respectively, probably did not arise on Earth until well after (<2-3 GYA;

Marin *et al.*, 2017; Wang and Luo, 2021) Gale crater was formed in the Late Noachian

(Fig. 1C). If we assume an origin of life took place ~4 GYA (i.e., either on Earth or Mars;

Cantine and Fournier, 2017; Carr, 2021) and that the hypothesis of common ancestry

between Earth and any possible Mars life is true (Velasco, 2018), these specific

metabolisms would likely have emerged after Gale hosted a lake (~3.8 to 3.1 GYA;

Hurowitz *et al.*, 2017). The modeled organisms are reasonable for testing early Mars

habitability if we assume an earlier second genesis occurred. If instead we assume life

evolved independently (i.e., if the hypothesis of common ancestry is false) on Earth,

these metabolisms also could have emerged in time to inhabit Gale Lake.

It is important to note that a potential caveat of constraint-based modeling is that it does

not account for temperature. Growth rates would be affected by changes in temperature

but not growth yield. In addition, our models assumed a constant light intensity. Daily

light fluctuations have been accounted for in previous models (e.g., Sarkar *et al.*, 2019),

however, this granular approach becomes exponentially complex for multi-species

models and was not necessary for our assessment. Further, as *R. palustris* is a

chemoautotroph and primary producer, we did not supply the models with additional

carbon sources other than CO_2. Organic carbon is present on Mars (Freissinet *et al.*,

2015; Eigenbrode *et al.*, 2018) but its distribution is not known.

In the present study, the individual models were capable of slow growth (Table 2) on the

modelled medium but were predicted to grow faster together as a community, indicating

syntrophy (Table 3). For instance, iron was effectively cycled by being reduced and

oxidized at the same rate (Table 3). Like the findings of Byrne et al. (2015), these

results indicate that iron does not become biologically unavailable to either organism

unless perhaps something were to physically limit access (e.g., biofilm formation). We

also found that the citrate secreted by *R. palustris* (Table 3) was consumed and used by

G. sulfurreducens in the reverse Krebs cycle (Fig. 4), illustrating genome-scale

metabolic modelling as a discovery tool. It has recently been confirmed that *G.*

sulfurreducens can utilize this cycle *in vitro* to fix CO_2 (Zhang *et al.*, 2020). Such hidden

autotrophy could also have implications for planetary protection if microbial metabolism

is used to guide planetary protection requirements.

A diurnal community model was constructed to further assess cross-feeding interactions

and metabolite storage during periods of light and dark (Fig. 3). Previous models have

been used to illustrate the circadian growth of other microbes (e.g., cyanobacteria), but

they have not been modelled in a community setting (e.g., Sarkar *et al.*, 2019). We

18

found that both organisms transferred metabolites to periods in which they prioritized

growth (Fig. 7). For example, *R. palustris* prioritized carbohydrate use during the day to

release phosphate, which is necessary to produce ATP. Additional ATP was produced

at night via nucleoside-diphosphate kinases to feed the non-growth maintenance

reaction. Conversely, *G. sulfurreducens* primarily transferred carbohydrates like citrate

and acetyl phosphate to produce ATP for growth at night.

In the context of early Mars, life, if it existed there, would have likely been present in the

form of a community. Microorganisms in nature are rarely found in isolation, with only

one known example to date (i.e., *Candidatus Desulforudis audaxviator*, Chivian *et al.*,

2008). This is because communities are better able to cope with changes in

temperature, humidity, and nutrients (Palková, 2004). The syntrophic interactions

identified here highlight the benefits of a community, which could allow life to adapt to

varying environmental conditions just as shifts occur in our simple diurnal model.

Beneficial community interactions might also have enabled life to achieve higher

biomass and biosignature production than might otherwise have been possible.

Our calculations show that Gale Lake could have maintained a concentration of cells

($\sim 10^3$-10^7 cells mL^{-1}) similar to modern lakes and oceans. For example, modern

terrestrial ocean waters and saltern brines have been shown to support 10^4-10^6 cells

per mL (Bar-On, Phillips and Milo, 2018); 10^4-10^7 Klempay *et al.*, 2021) depending on

depth and other variables. Terrestrial ocean and lake sediments have also been shown

to harbor 10^4-10^9 cells per cm^3 (Kallmeyer *et al.*, 2012; Wurzbacher *et al.*, 2017).

In addition to predicting the habitability and biomass density of early Mars, it is of

particular interest to calculate the potential densities of biomarkers (i.e., molecular

fossils) that could be used to detect ancient life. For example, certain membrane lipids

like bacteriohopanepolyols or hopanoids can persist for billions of years in the terrestrial

rock record (Brocks *et al.*, 2005). The hopanoids synthesized by many bacteria are

largely resistant to degradation because of their pentacyclic carbon skeleton, and can

be detected through several analytical techniques (e.g., high-performance liquid

chromatography, liquid/gas chromatography–mass spectrometry; Belin *et al.*, 2018).

Here, both *R. palustris* (Welander *et al.*, 2009) and *G. sulfurreducens* (Härtner, Straub

and Kannenberg, 2005) are known to produce biomarker hopanoids under anaerobic

conditions in culture. Our model predicts that hopanoids could have been produced at a

concentration ($\sim 10^{-1}$-10^{-2} g kg^{-1} Gale sediment; see supplemental materials) well above

the limit of detection for miniaturized GCMS systems (~1 PPM, Coy et al., 2011) in the

case where all cells in the photic zone at some steady state timepoint settled and were

evenly distributed into Gale Lake sediments and were 100% preserved. While this

represents an upper bound for a single such event, such events could have occurred

repeatedly, due to dust storms or impact event-driven dust events blocking the sunlight

required for photosynthesis. Future work could link steady state hopanoid (or other

biosignature) production to deposition, preservation, and diagenesis, to yield refined

estimates in support of *in situ* life detection.

370 **5. Conclusions**

Here we have demonstrated that Gale Lake could have provided the energy necessary

to sustain a simple and syntrophic microbial community. It must be noted that our

models and calculations contain several assumptions. For instance, Gale Lake would

not have had an even distribution of light, or of any 'metabolite' for that matter,

throughout its depths. It is also pertinent to note that the densities of all lake

components would have changed greatly over time with evaporation (e.g., Rapin *et al.*,

2019). This would also inevitably lead to reduction in hopanoid concentration potentially

rendering it more difficult to detect. Nevertheless, these data provide guidance for Mars

habitability studies. Future research should further develop and confirm candidate

biosignatures *in vitro* and using other scenarios of habitability on early Mars and on

other worlds.

Acknowledgements

Author Disclosure Statement

The authors declare no conflict of interest.

References

Alsiyabi, A., Immethun, C. M. and Saha, R. (2019) 'Modeling the Interplay between
Photosynthesis, CO2 Fixation, and the Quinone Pool in a Purple Non-Sulfur Bacterium',
Scientific Reports, 9(1), pp. 1–9. doi: 10.1038/s41598-019-49079-z.

Bar-On, Y. M., Phillips, R. and Milo, R. (2018) 'The biomass distribution on Earth',
Proceedings of the National Academy of Sciences of the United States of America,
115(25), pp. 6506–6511. doi: 10.1073/pnas.1711842115.

Belin, B. J. *et al.* (2018) 'Hopanoid lipids: From membranes to plant-bacteria
interactions', *Nature Reviews Microbiology*. Nature Publishing Group, pp. 304–315. doi:
10.1038/nrmicro.2017.173.

Berger, J. A. *et al.* (2019) 'Manganese enrichment pathways relevant to Gale Crater,
Mars: Evaporative concentration and chlorine-induced precipitation', in *50th Lunar and
Planetary Science Conference*.

Borlina, C. S., Ehlmann, B. L. and Kite, E. S. (2015) 'Modeling the thermal and physical
evolution of Mount Sharp's sedimentary rocks, Gale Crater, Mars: Implications for
diagenesis on the MSL Curiosity rover traverse', *Journal of Geophysical Research:*

Planets, 120(8), pp. 1396–1414. doi: 10.1002/2015JE004799.

Bratbak, G. and Dundas, I. (1984) 'Bacterial dry matter content and biomass estimations.', *Applied and environmental microbiology*, 48(4), pp. 755–7. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6508285 (Accessed: 3 October 2018).

Brocks, J. J. *et al.* (2005) 'Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea', *Nature*, 437(7060), pp. 866–870. doi: 10.1038/nature04068.

Byrne, J. M. *et al.* (2015) 'Redox cycling of Fe(II) and Fe(III) in magnetite by Fe-metabolizing bacteria', *Science*, 347(6229), pp. 1473–1476. doi: 10.1126/science.aaa4834.

Cantine, M. D. and Fournier, G. P. (2017) 'Environmental Adaptation from the Origin of Life to the Last Universal Common Ancestor', *Origins of Life and Evolution of Biospheres 2017 48:1*, 48(1), pp. 35–54. doi: 10.1007/S11084-017-9542-5.

Carr, C. E. (2021) *Resolving the History of Life on Earth by Seeking Life As We Know It on Mars*. Atlanta.

Chivian, D. *et al.* (2008) 'Environmental Genomics Reveals a Single-Species Ecosystem Deep Within Earth', *Science*, 322(5899), pp. 275–278. doi: 10.1126/SCIENCE.1155495.

Cockell, C. (2020) *Astrobiology : understanding life in the universe*. 2nd edn.

Deit, L. Le *et al.* (2013) 'Sequence of infilling events in Gale Crater, Mars: Results from morphology, stratigraphy, and mineralogy', *Journal of Geophysical Research: Planets*, 118(12), pp. 2439–2473. doi: 10.1002/2012JE004322.

Ebrahim, A. *et al.* (2013) 'COBRApy: COnstraints-Based Reconstruction and Analysis for Python', *BMC Systems Biology*, 7(1), p. 74. doi: 10.1186/1752-0509-7-74.

Eigenbrode, J. L. *et al.* (2018) 'Organic matter preserved in 3-billion-year-old mudstones at Gale crater, Mars', *Science*, 360(6393), pp. 1096–1101. doi: 10.1126/SCIENCE.AAS9185.

Freissinet, C. *et al.* (2015) 'Organic molecules in the Sheepbed Mudstone, Gale Crater, Mars', *Journal of Geophysical Research: Planets*, 120(3), pp. 495–514. doi: 10.1002/2014JE004737.

Fukushi, K. *et al.* (2019) 'Semiarid climate and hyposaline lake on early Mars inferred from reconstructed water chemistry at Gale', *Nature Communications*, 10(1), pp. 1–11. doi: 10.1038/s41467-019-12871-6.

Grotzinger, J. P. *et al.* (2015) 'Deposition, exhumation, and paleoclimate of an ancient lake deposit, Gale crater, Mars', *Science*, 350(6257). doi: 10.1126/science.aac7575.

Hädicke, O., Grammel, H. and Klamt, S. (2011) 'Metabolic network modeling of redox balancing and biohydrogen production in purple nonsulfur bacteria', *BMC Systems Biology*, 5(1), pp. 1–18. doi: 10.1186/1752-0509-5-150.

Harris, C. R. *et al.* (2020) 'Array programming with NumPy', *Nature*. Nature Research, pp. 357–362. doi: 10.1038/s41586-020-2649-2.

Härtner, T., Straub, K. L. and Kannenberg, E. (2005) 'Occurrence of hopanoid lipids in anaerobic Geobacter species', *FEMS Microbiology Letters*, 243(1), pp. 59–64. doi: 10.1016/j.femsle.2004.11.039.

Hurowitz, J. A. *et al.* (2017) 'Redox stratification of an ancient lake in Gale crater, Mars', *Science*, 356(6341). doi: 10.1126/SCIENCE.AAH6849.

Jones, E. G., Lineweaver, C. H. and Clarke, J. D. (2011) 'An Extensive Phase Space for the Potential Martian Biosphere', *Astrobiology*, 11(10). doi: 10.1089/ast.2011.0660.

Kallmeyer, J. *et al.* (2012) 'Global distribution of microbial abundance and biomass in subseafloor sediment', *Proceedings of the National Academy of Sciences of the United States of America*, 109(40), pp. 16213–16216. doi: 10.1073/pnas.1203849109.

Kato, S. and Ohkuma, M. (2021) 'A Single Bacterium Capable of Oxidation and Reduction of Iron at Circumneutral pH', *Microbiology Spectrum*. Edited by J. A. Gralnick. doi: 10.1128/SPECTRUM.00161-21.

Klamt, S., Schuster, S. and Gilles, E. D. (2002) 'Calculability analysis in underdetermined metabolic networks illustrated by a model of the central metabolism in purple nonsulfur bacteria', *Biotechnology and Bioengineering*, 77(7), pp. 734–751. doi: 10.1002/BIT.10153.

Klempay, B. *et al.* (2021) 'Microbial diversity and activity in Southern California salterns and bitterns: analogues for remnant ocean worlds', *Environmental Microbiology*. doi: 10.1111/1462-2920.15440.

Knies, D. *et al.* (2015) 'Modeling and simulation of optimal resource management during the diurnal cycle in Emiliania huxleyi by genome-scale reconstruction and an extended flux balance analysis approach', *Metabolites*, 5(4), pp. 659–676. doi: 10.3390/metabo5040659.

Lambrecht, N. *et al.* (2021) '"Candidatus Chlorobium masyuteum," a Novel Photoferrotrophic Green Sulfur Bacterium Enriched From a Ferruginous Meromictic Lake', *Frontiers in Microbiology*, 12(July), p. 1768. doi: 10.3389/FMICB.2021.695260.

Lanza, N. L. *et al.* (2014) 'High manganese concentrations in rocks at Gale crater, Mars', *Geophysical Research Letters*, 41(16), pp. 5755–5763. doi: 10.1002/2014GL060329.

Mahadevan, R. *et al.* (2006) 'Characterization of metabolism in the Fe(III)-reducing organism Geobacter sulfurreducens by constraint-based modeling', *Applied and Environmental Microbiology*, 72(2), pp. 1558–1568. doi: 10.1128/AEM.72.2.1558-1568.2006.

Marin, J. *et al.* (2017) 'The Timetree of Prokaryotes: New Insights into Their Evolution and Speciation', *Molecular Biology and Evolution*, 34(2), pp. 437–446. doi: 10.1093/MOLBEV/MSW245.

Marinos, G., Kaleta, C. and Waschina, S. (2020) 'Defining the nutritional input for
genome-scale metabolic models: A roadmap', *PLOS ONE*. Edited by A. Pagnani, 15(8),
p. e0236890. doi: 10.1371/journal.pone.0236890.

Mars Fact Sheet (2021) *NASA*. Available at:
https://nssdc.gsfc.nasa.gov/planetary/factsheet/marsfact.html (Accessed: 19 August
2021).

Orth, J. D., Thiele, I. and Palsson, B. O. (2010) 'What is flux balance analysis?', *Nature*
Biotechnology. NIH Public Access, pp. 245–248. doi: 10.1038/nbt.1614.

Palková, Z. (2004) 'Multicellular microorganisms: laboratory versus nature', *EMBO*
Reports, 5(5), p. 470. doi: 10.1038/SJ.EMBOR.7400145.

Von Paris, P. *et al.* (2013) 'N2-associated surface warming on early Mars', *Planetary*
and Space Science, 82–83, pp. 149–154. doi: 10.1016/j.pss.2013.04.009.

Payré, V. *et al.* (2019) 'Copper enrichments in the Kimberley formation in Gale crater,
Mars: Evidence for a Cu deposit at the source', *Icarus*, 321, pp. 736–751. doi:
10.1016/j.icarus.2018.12.015.

Rapin, W. *et al.* (2019) 'An interval of high salinity in ancient Gale crater lake on Mars',
Nature Geoscience, pp. 1–7. doi: 10.1038/s41561-019-0458-8.

Reback, J. *et al.* (2020) 'pandas'. doi: 10.5281/ZENODO.3715232.

Sarkar, D. *et al.* (2019) 'A diurnal flux balance model of Synechocystis sp. PCC 6803
metabolism', *PLoS Computational Biology*, 15(1), p. e1006692. doi:
10.1371/journal.pcbi.1006692.

Velasco, J. (2018) 'Universal common ancestry, LUCA, and the Tree of Life: three
distinct hypotheses about the evolution of life', *Biology & Philosophy 2018 33:5*, 33(5),
pp. 1–18. doi: 10.1007/S10539-018-9641-3.

Wang, S. and Luo, H. (2021) 'Dating Alphaproteobacteria evolution with eukaryotic
fossils', *Nature Communications 2021 12:1*, 12(1), pp. 1–9. doi: 10.1038/s41467-021-
23645-4.

Welander, P. V. *et al.* (2009) 'Hopanoids play a role in membrane integrity and pH
homeostasis in Rhodopseudomonas palustris TIE-1', *Journal of Bacteriology*, 191(19),
pp. 6145–6156. doi: 10.1128/JB.00460-09.

Wordsworth, R. *et al.* (2021) 'A coupled model of episodic warming, oxidation and
geochemical transitions on early Mars', *Nature Geoscience*, 14(3), pp. 127–132. doi:
10.1038/s41561-021-00701-8.

Wurzbacher, C. *et al.* (2017) 'Shifts among eukaryota, bacteria, and archaea define the
vertical organization of a lake sediment', *Microbiome*, 5(1), p. 41. doi: 10.1186/S40168-
017-0255-9.

Zhang, T. *et al.* (2020) 'The hidden chemolithoautotrophic metabolism of Geobacter
sulfurreducens uncovered by adaptation to formate', *The ISME Journal*, 14, pp. 2078–

2089. doi: 10.1038/s41396-020-0673-8.