Title: Electric shock causes a fear-like persistent behavioral response in the nematode

Caenorhabditis elegans

Authors: Ling Fei Tee[1], Jared J. Young[2], Ryoga Suzuki[1], Keisuke Maruyama[1], Yuto Endo[1,3],

Koutarou D. Kimura[1,3,4]*

Affiliations: 1. Graduate School of Science, Nagoya City University, Nagoya, 467-8501,

Japan; 2: Biology Department, Mills College, Oakland, CA, 94613, USA; 3: Department of

Biological Sciences, Graduate School of Science, Osaka University, Toyonaka, Osaka, 560-

0043, Japan; 4. RIKEN center for Advanced Intelligence Project, Tokyo, 103-0027, Japan

Correspondence: Koutarou D. Kimura, email: kokimura@nsc.nagoya-cu.ac.jp

ABSTRACT

Behavioral persistency reflects internal brain states, which are the foundations of multiple brain functions. However, experimental paradigms that enable genetic analyses of behavioral persistency and its associated brain functions have been limited. Here we report novel persistent behavioral responses caused by electric stimuli in the nematode *Caenorhabditis elegans*. When the animals on bacterial food are stimulated by alternating current, their movement speed suddenly increases more than 2-fold, which persists for minutes even after the electric stimulation is terminated. Genetic analyses reveal that multiple types of voltage-gated channels are required for the response, possibly as the sensors, and neuropeptide signaling regulates the duration of the persistent response. Additional behavioral analyses indicate that the animal's response to electric shock is scalable and has a negative valence. These properties, along with persistence, have been recently regarded as essential features of emotion, suggesting that the animal's response to electric shock may express a form of emotion, such as fear.

INTRODUCTION

Animal behaviors, such as feeding, mating, aggression, and sleeping, are strongly related to internal states in the brain, namely motivation, arousal, drive, and emotion (Anderson, 2016; Berridge, 2004; Kennedy et al., 2014). Because animals can produce different behavioral responses to the same stimulus depending on their brain state, these states are considered to be the foundation from which a variety of behavioral responses emerge (Chen & Hong, 2018; Maimon, 2011). The brain states persist for a certain period of time and transit to a different state based on internal and/or external triggers, which can be observed as transitions among different persistent behavioral states. The neural mechanisms of brain/behavioral states are starting to be revealed. For example, the behavioral states of mating and aggressiveness have been shown to be controlled by relatively small circuits in mice and flies (Anderson, 2016; Hoopfer et al., 2015; Lee & Dan, 2012). However, the mechanisms of persistent brain/behavioral states have been revealed in only limited studies, and, moreover, the molecular basis that generates persistent states is still unclear.

The nematode *Caenorhabditis elegans* has been widely used in neurobiological research because of the feasibility of molecular, physiological, and behavioral analyses of neural functions (Bargmann, 2006; de Bono & Maricq, 2005; Sasakura & Mori, 2013). Recently, persistent behavioral states have also been studied in the animals, especially roaming/dwelling and sleep/arousal. Roaming/dwelling are states of locomotion on bacterial food, either moving over long distances at a constant speed or moving back and forth over short distances (Ben Arous et al., 2009; Fujiwara et al., 2002). Sleep in *C. elegans* is a phenomenon observed just before molt, and meets the definition of sleep in higher animals such as humans, rodents, fishes and flies (Raizen et al., 2008). Both the neural circuits and

genes that control these phenomena are being revealed (Flavell et al., 2020). However, it is

not clear whether *C. elegans* has any other persistent states.

In this study, we report that *C. elegans* exhibits a novel type of persistent behavioral response

to electric stimulus. The animals respond to alternating current (AC) stimulus by immediately

increasing their speed, and the speed increase persists even for minutes after five seconds of

electric stimulus: This result suggests that the response is caused not by direct stimulation of

the motor system for rapid movement but by persistent activity of a specific set of neurons to

generate the behavioral response. Further behavioral analyses indicate that the speed increase

to AC stimulus is scalable and has negative valence. Because persistent behavioral response

is one of the most prominent characteristics of emotions of animals (Abbott, 2020; Anderson

& Adolphs, 2014; Nettle & Bateson, 2012; Paul & Mendl, 2018; Perry & Baciadonna, 2017),

and persistency, scalability and valence are three of the four key features of animal emotions

proposed by Anderson and Adolphs (Anderson & Adolphs, 2014), the speed increase to the

electric shock may express a form of emotion. A series of candidate genetic analyses reveal

that the response is not mediated by well-known chemo- or mechano-sensory mechanisms.

Instead, it requires voltage-gated calcium and potassium channel genes, which are required

for electro-sensation in cartilaginous fishes (Bellono et al., 2017, 2018), suggesting an

evolutionarily conserved mechanism for electro-sensation. Furthermore, we find that

neuropeptide signaling regulates the duration of persistence. These results indicate that the

animals' response to electric shock can be a suitable paradigm to reveal molecular and

physiological mechanisms of electro-sensation as well as persistent brain/behavioral states.

RESULTS

C. elegans' **speed is increased by AC stimulation**

Initially, we started this project by studying *C. elegans* behavioral responses to AC stimuli.

The animals are known to respond to direct current (DC), migrating along the electric field

from the positive end to the negative end (Sukul & Croll, 1978), and a few classes of

chemosensory neurons (ASH and ASJ) were found to be required for their ability to align

themselves according to the DC field (Gabel et al., 2007). However, the animal's migratory

response to AC stimulus has not been reported yet. In our original setup (Fig. 1), several adult

wild-type animals were placed onto 9 cm agar plates seeded with a small bacterial food patch

and subjected to AC stimulation. The complete trajectories produced by the animals were

video-recorded, and their speed was calculated.

Fig. 1. Experimental setup of electric shock experiment. This setup consists of an arbitrary waveform generator, amplifier, multimeter, oscilloscope, camera, and desktop computer.

We first studied the response to AC stimulation covering a range between 15 - 105 V at 60

Hz (the commercial power frequency in Japan), and found that the animals increased their

average speed during electric stimulation by varying amounts (Supplementary Fig. 1). We

then conducted a series of systematic analyses with different voltages and frequencies at 30–

75 V and 0.25–256 Hz (Supplementary Fig. 2). After the analysis, we noticed that an

interesting characteristic of this behavioral phenotype is most apparent when using 4 Hz

stimuli: When animals were stimulated with 30 V, their average speed of movement suddenly

increased more than 2-fold, and this persisted during the electric admission. We named this

behavior the "ON response" (Fig. 2A and C). During this running behavior, the animals

engage in rapid body bends as well as rapid head movements (Supplementary Videos 1 and

2). In the ON response, we did not detect a statistical bias in direction (Supplementary Fig.

3). Unexpectedly, when a stronger electric stimulus of 75 V was applied, it caused a

significant increase in average speed not during but immediately after the stimulus, which we

named the "OFF response" (Fig. 2B). A fraction of the animals responded during the

stimulus in the OFF response condition, while, in the majority of the animals, the speed was

suppressed during the stimulus and then increased immediately after its removal

(Supplementary Fig. 4 and Videos 3 and 4). With other frequencies, ON and OFF responses

were also observed, but were less clear compared to those with 4 Hz (Supplementary Fig. 2).

The range of voltage per length (30–75 V/6 cm = 5–12.5 V/cm) is similar to the range

previously shown to elicit responses to DC (3–12 V/cm) (Gabel et al., 2007), suggesting that

these electric stimuli are physiologically meaningful for the animals. The fact that ON and

OFF responses at 4 Hz were substantially different with only a 2.5-fold difference in the

voltage at the same frequency is interesting because different behavioral responses generally

require much larger differences in stimulus intensity with other stimuli, such as odor

(Bargmann et al., 1993).

Fig. 2. Animals' speed is increased by AC stimulation. **A,** (Left) Speed-time graph with 30 V stimulation at 4 Hz. Thin and thick lines are for individual and average values, respectively. Gray indicates the duration of electric stimulation (0-30 s). (Right) Scatter plot showing average speed of individual animals before, during and after electric stimulation. Each period is 30 s. n = 35. **B,** Speed-time graph (left) and scatter plot (right) with 75 V stimulation at 4 Hz. n = 36. **C,** Cartoons of worm's response to the electric shock. (Left) Before electric stimulation, the animals stay on food patch and maintain their speed at around 0.1 mm/s. (Right) During electric stimulation is delivered, the animals increase speed to around 0.2 - 0.3 mm/s and leave the food patch which persists even after the stimulus is terminated. Statistical values were calculated using Kruskal-Wallis test with Bonferroni correction. ** $p < 0.001$.

We then analyzed whether this response depends on voltage or current by manipulating the

salt concentration in the assay plate (Fig. 3): When 30V is applied to the high salt plate the

current should be similar to the current produced when 75V is applied to a plate with our

standard (control) salt concentration. Conversely, when 75V is applied to the low salt plate,

the current should be similar to the current produced when 30V is applied to the control plate

(Fig. 3A). As shown in Fig. 3, 30 V and 75 V stimuli caused ON and OFF responses,

respectively, regardless of the current value, indicating that the behavioral response depends

on voltage.

Speed increase lasts for several minutes

Next, we examined how long the increased speed persists during and after the stimulus.

When the duration of 30V applied electric shock was 1-2 minutes, significant speed increases

were maintained during the stimulus, lasted for ~1 min after the stimulus, and went back to

the baseline level (Fig. 4A). Interestingly, when the animals were stimulated only for 5 sec,

the speed increase still lasted for 1.5 min. When 4 min stimulus was applied, the increase was

maintained during the stimulus but went back to the baseline level 30 sec after the stimulus.

During 10 min stimulation, the significant speed increase was observed only for 5.5 min.

Thus, we concluded that the ON response caused by 30 V stimulation persists ~5 min at

most.

This result suggested the possibility that the speed increase may decline after several minutes

because of fatigue in motor systems. However, animals stimulated intermittently 5 times for

30 seconds per stimulation maintained a speed increase for a much longer time than those

under the continuous stimulus (Fig. 4B versus "10 minutes" in A). This result supports the

idea that the decrease in speed during the long ON stimulation period is not caused by fatigue

Fig. 3. Speed increase is dependent on voltage, not on current. **A,** Voltage-current graph with different salt concentrations (indicated by different symbols). Each dot represents the measured value on the day of the experiment. The final total osmolarity for sodium chloride (Na$^+$ and Cl$^-$) and sucrose for all the plates was 400 mOsm. **B–E,** Behavioral responses of animals assayed on high-salt plate with 30 V (**B**; n = 32), on control plate with 75 V (**C**; n = 35) or 30 V (**D**; n = 36), or on low-salt plate with 75 V (**E**; n = 34). Stimulation period is indicated by a shaded grey box. **F–I,** Scatter plot showing average speed of individual animals before, during and after electric stimulation, corresponding to the panels **B–E**, respectively. Statistical values were calculated using Kruskal-Wallis test with Bonferroni correction. ** $p < 0.001$.

in the motor system, but possibly by sensory adaptation, which is widely known to adjust the

animal's sensory response to new environments (Wark et al., 2007).

We then tested the persistence of speed increase in the OFF response with 75 V. Five and 30

sec stimuli caused similar or longer persistent responses after the stimulus than 30 V did (Fig.

4C). Remarkably, 45 sec stimulus caused >2 min persistent response, which is the longest

among the responses to 30 and 75 V stimuli after the stimulus. When animals were

stimulated for 1 min, no ON or OFF responses were observed. The fact that the larger

stimulus (75 V) caused longer persistent responses than the smaller one (30 V) suggests that

the response to electric shock is "scalable" (i.e., different strength of stimulus causes different

strength of behavioral response), one of the critical "emotion primitives" together with

persistence (Anderson & Adolphs, 2014).

We then tested the effect of food presence on the speed increase. *C. elegans* move slowly on

the bacterial food lawn and faster out of the lawn (Sawin et al., 2000). As we used a small

food lawn to localize the animal's initial positions to the center of the plate (Fig. 1 and 2C), it

was possible that the electric stimulus caused the animals to move away from the food lawn,

which then caused increased speed due to the absence of food. If this is the case, the animal's

speed would be considerably lower with the electric stimulus when the plates were fully

covered with a bacterial lawn. To test this hypothesis, we compared the time-course of speed

changes on plates with a small patch of food lawn and with a full food lawn. As shown in

Fig. 4D and E (compare Fig. 4A "4 minutes" and C " 30 seconds", respectively), there was no

substantial difference in the time course of speed change between the small food and the full

food plates in ON as well as OFF responses, demonstrating that the speed increase is not

caused by the food absence but by the electric stimulation itself.

Fig. 4. Speed increase persisted for minutes even after the stimulation. **A,** Speed-time graphs of ON response with 30 V stimulation of different time periods, ranging from 5 seconds to 10 minutes. **B,** Speed-time graph for intermittent electric stimulation of 30 seconds, 5 times with 90 s–intervals. **C,** Speed-time graphs of OFF response with 75 V stimulation of different time periods, ranging from 5 seconds to 1 minute. **D** and **E,** Speed-time graphs for electric stimulation of 30 V for 4 minutes (**D**) or 75 V for 30 s (**E**) with animals placed on full food lawn. Shaded regions around the lines represent standard deviation. Statistical values were calculated using Kruskal-Wallis test with Bonferroni correction for the differences from the average speed before the stimulation. * $p < 0.01$, ** $p < 0.001$. Sample numbers were 32–46 per condition, and the details are described in Supplementary Table.

To further confirm that result, we analyzed the animals' speed on a stripe-like food pattern

(Supplementary Fig. 5A). We did not observe a significant difference in speeds when the

animals moved into or out of the food area (Supplementary Fig. 5B). This result suggests that

the electric stimulus may have negative valence that is more influential to the animal's

behavior than the food signal, even though food is critical for their survival. It further

suggests that animals prioritize moving away from a harmful condition, such as the electric

shock, to protect themselves.

Voltage-gated ion channel genes and neuropeptide signaling are required for the AC

response

The molecules required for responses to electric signals have only been revealed in

cartilaginous fishes: Bellono et al. reported that electrosensory cells in little skate and chain

catshark use L-type voltage-gated calcium channels (VGCC) and voltage-gated big-

conductance potassium (BK) channels (Bellono et al., 2017, 2018). To identify gene(s)

required for the response to electric shock in *C. elegans*, we analyzed a series of mutant

strains of candidate genes. Specifically, we tested mutants of genes involved in the animals'

chemo- and mechano-sensation, and the homologues of genes involved in electroreception in

the cartilaginous fishes.

C. elegans' chemo-sensation is largely mediated by the 12 pairs of amphid sensory neurons in

the head, which are classified into the ones using TAX-2 and TAX-4 cyclic nucleotide-gated

channel (CNGC) subunits or the others using OSM-9 and OCR-2 transient receptor potential

(TRP) channel subunits for depolarization (Coburn & Bargmann, 1996; Colbert et al., 1997;

Komatsu et al., 1996; Tobin et al., 2002). In addition to loss-of-function mutants for the

above-mentioned genes, we tested mutants for *che-2*, a gene required for the proper

formation and function of the sensory cilia (Fujiwara et al., 1999). For mechano-sensation,

we analyzed loss- or reduction-of-function alleles of *mec-4, mec-10*, and *trp-4*. *mec-4* and

mec-10 genes encode DEG/ENaC proteins that form a mechanosensory ion channel complex

for transduction of gentle touch (Driscoll & Chalfie, 1991; Huang & Chalfie, 1994), while

trp-4 encodes TRPN (NOMPC) for harsh touch response (Kang et al., 2010). All these

mutant strains exhibited wild-type-like ON and/or OFF responses (panel A in Fig. 5 and 6 for

ON and OFF responses, respectively). Some mutants (*osm-9;ocr-2*, *che-2*, *mec-4*, *mec-10*,

and *tph-1*) exhibited statistical differences in either the ON or OFF response, but not both

(Fig. 5F and 6F), suggesting the partial involvement of these genes, although the defects in

speed increase (i.e. ΔSpeed) were not as severe as the ones of VGCC mutants (see below).

The non-involvement of *tax-4* also indicates that temperature increase caused by the electric

stimulus is not responsible for the speed increase (see Discussion for details).

Fig. 5. Genetic analysis of ON response. A–D, Speed-time graphs of ON response with 30 V stimulation of 4 min on mutants of sensory signaling (**A**), voltage-gated channels (**B**), biogenic amine biosynthesis (**C**), and neuropeptide biosynthesis (**D**). **E**, Scatter plot showing Δspeed of individual wild-type and *egl-3* mutant animals during t = 330-360 s in **D**. **F**, In a series of daily experiments, wild-type N2 and three to five mutant strains were analyzed in parallel. All the N2 data are combined, and the mutant strains are arranged in ascending order of median values in **F**. Statistical values were calculated using Kruskal-Wallis test with Bonferroni correction. ** $p < 0.001$. Sample numbers were 30–36 per mutant strain, and the details are described in the Supplementary Table.

206 We then tested *egl-19*, the orthologue of the L-type VGCC alpha subunit (Lee et al., 1997),

207 which functions in the sensory organ for environmental electric signals for cartilaginous

208 fishes (Bellono et al., 2017, 2018). We found that two reduction-of-function alleles of *egl-19*

209 mutants exhibited strong defects in ON and OFF responses (Fig. 5 and 6, panels B and F).

210 While *egl-19* is expressed widely (Lee et al., 1997), at least one allele of *egl-19* mutants

211 exhibited movement speed comparable to wild-type animals before stimulation

212 (Supplementary Fig. 6), suggesting that the defect in the response is not caused by a problem

213 in the basal locomotory system. These results suggest that the VGCC may be an

214 evolutionarily conserved sensor for environmental electricity.

216 This finding further motivated us to test two other types of voltage-gated calcium channels,

217 namely, N-type (UNC-2) and T-type (CCA-1) VGCCs (Schafer & Kenyon, 1995; Steger et

218 al., 2005), although only L-type VGCC had been found to be involved in electrical responses

219 in the cartilaginous fishes. Unexpectedly, mutants for two alleles of *unc-2* were defective in

220 both ON and OFF responses, while *cca-1* mutants behaved similar to the wild-type controls

221 (Fig. 5 and 6, panels B and F, and Supplementary Fig. 7).

Fig. 6. Genetic analysis of OFF response. A–D, Speed-time graph of OFF response with 75 V stimulation of 30 s on mutants of sensory signaling (**A**), voltage-dependent channels (**B**), biogenic amine biosynthesis (**C**), and neuropeptide biosynthesis (**D**). **E**, Scatter plot showing Δspeed of individual wild-type and *egl-3* mutant animals during $t = 180\text{-}210$ s in **D**. **F**, Scatter plot showing Δspeed of individual animals during 3 min after the stimulation (*i.e., t* = 30-210 s). In a set of daily experiments, wild-type and three to five mutant strains were analyzed in parallel. All the wild-type data are combined, and the mutant strains are arranged in ascending order of median values in **F**. **G**, Scatter plot showing average speed of individual wild-type and *slo-1* mutant animals during 30 s after the stimulation (*i.e., t* = 30-60 s). Statistical values were calculated using Kruskal-Wallis test with Bonferroni correction. ** $p < 0.001$. Sample numbers were 30–36 per mutant strain, and the details are described in the Supplementary Table.

We then investigated the involvement of the BK channel, a voltage-gated potassium channel,

also known to be involved in electro-sensation in cartilaginous fish (Bellono et al., 2017,

2018). Interestingly, two alleles of *slo-1*, the sole ortholog of BK channels in *C. elegans*

(Davies et al., 2003; Wang et al., 2001), also exhibited statistical differences in the ON as

well as the OFF response, at least in some aspects (Fig. 5B and F for ON response, and 6B, F

and G for OFF response). The possible involvement of BK channels in addition to the VGCC

in the *C. elegans'* electrical response suggest that the molecular mechanisms of electro-

sensation may be evolutionarily conserved.

Lastly, we attempted to identify genes required for the behavioral persistency, and considered

the genes involved in the biosynthesis of neuromodulators. We tested *cat-2* (dopamine), *tph-1*

(serotonin), *bas-1* (dopamine and serotonin) and *tdc-1* (tyramine and octopamine) mutant

animals (Alkema et al., 2005; Lints & Emmons, 1999; Loer & Kenyon, 1993; Sze et al.,

2000), and all of these mutants exhibited wild-type-like responses, indicating that these

neuromodulators are not involved (panel C in Fig. 5 and 6). Because dopamine and serotonin

signaling are known to be required for the feeding status-dependent modulation of migratory

speed, these results are also consistent with the fact that feeding status is not the causal reason

for the speed increase (Fig. 4D and E, and Supplementary Fig. 5).

We then further tested the involvement of neuropeptides by using loss- or reduction-of-

function mutations of *egl-3*, a gene required for maturation of pro-neuropeptides (Kass et al.,

2001). Unexpectedly, mutations in both alleles of *egl-3*, *n589* and *ok979*, caused weaker 30 V

ON response and, moreover, much longer persistence of the speed increase after the electric

shock in ON and OFF responses (Fig. 5 and 6, panels D-F), indicating that the persistent

activity in the neural circuit for speed increase is down-regulated by neuropeptide signaling

in wild-type animals.

DISCUSSION

In the present study, we revealed that *C. elegans* exhibits a persistent speed increase in

response to AC stimuli. This behavioral response is characterized by persistence, scalability,

and valence, suggesting that it reflects an emotional state of *C. elegans*, which has never been

reported. In addition, genetic analysis revealed that genes involved in electro-sensation in

cartilaginous fishes and a neuropeptide biosynthesis gene are required for the response,

demonstrating that the AC response of the animals is an ideal paradigm to genetically dissect

the mechanism of electro-sensation as well as that of persistent behavioral states.

Response to electric stimulus and its mechanisms in *C. elegans* and other animal species

In neuroscience research, electricity is used as an unconditioned stimulus with negative

valence to cause associative learning in rodents and in flies (Quinn et al., 1974; Rescorla,

1968). In nature, however, multiple animal species are known to respond to electricity for

survival purposes, such as communication, navigation and/or prey detection (Crampton,

2019; Pettigrew, 1999). For example, weakly electric African fish (*Gnathonemus petersii*) utilize their epidermal electroreceptors to receive self-produced electric signals, allowing the fish to identify, locate, and examine nearby objects (von der Emde et al., 2008). In addition, platypus (*Ornithorhynchus anatinus*) detects electric signals via their duck-like bills to locate and avoid objects when navigating in the water (Scheich et al., 1986). Blind cave salamander (*Proteus anguinus*) perceives a moving back-and-forth direct-current field and its polarity via ampullary organs to survive and navigate in their environment, which is in complete darkness as their eyes are undeveloped (Istenič & Bulog, 1984; Roth & Schlegel, 1988). In invertebrates, bumblebees (*Bombus terrestris*) sense environmental electric fields via sensory hairs to make foraging decisions (Clarke et al., 2013; Sutton et al., 2016). Such wide use of electric signals in the animal kingdom suggests physiological importance of the sensation and behavioral responses to environmental electric signals although the molecular mechanisms for electric sensation have been poorly revealed.

In this study, we established an original experimental paradigm and found that *C. elegans* responds to AC electric stimulus: The animals significantly increase their movement speed during and after the stimulus for minutes. Although the animals have also been reported to respond to DC (Gabel et al., 2007), we consider that the responses to AC and DC are substantially different for the following reasons. (1) In the DC field, the animals moved at a certain angle (~4° per 1 V/cm), which was not observed in our AC stimulus (Supplementary Fig. 3). (2) Movement speed did not change with the DC stimulus (Gabel et al., 2007).

In addition, five pairs of amphid sensory neurons were involved in the DC response (Gabel et al., 2007), while mutations in genes required for sensory signaling in the amphid sensory neurons (*tax-4*, *osm-9*, *ocr-2*, and *che-2*) did not affect the AC response significantly in our

study (Fig. 5 and 6), indicating that DC and AC responses utilize different sensory

mechanisms. Our result also rules out the possibility that the animals respond to increased

agar temperature due to the AC stimulus, because the mutation in *tax-4*, the gene essential for

temperature sensation (Komatsu et al., 1996) did not affect the response. In addition, the

genes required for mechano-sensation (*mec-4*, *mec-10*, and *trp-4*) are not required for the AC

response either.

We found that VGCC and possibly BK channel as well, the voltage-gated calcium and

potassium channels for electro-sensation in the cartilaginous fishes, are involved in the AC

response of *C. elegans*. The involvement of different types of voltage-gated channels in the

sensation of electricity suggests that this mechanism is evolutionarily conserved. It also

suggests that EGL-19 and SLO-1 may function coordinately in a subset of neurons that sense

electricity. Since *egl-19* and *slo-1* are widely expressed in most of the neurons as well as in

muscles (Davies et al., Lee et al., 1997; Wang et al., 2001), it would be interesting to identify

the neurons where these genes function to sense the electric signals.

Electric stimulus causes persistent behavioral response

Persistent neural activity, a sustained neural activity caused by a short term stimulus, plays

critical roles in brain function, such as controlling motivation, arousal, and emotion as well as

working memory and decision-making, although its detailed mechanisms have not been

sufficiently elucidated (Anderson, 2016; Berridge, 2004; Curtis & Lee, 2010; Major and

Tank, 2004). Persistent behavioral state is caused by persistent neural activity, suggesting that

genetic analysis of persistent behavioral state can reveal molecular mechanisms(s) of

persistent neural activity underlying brain functions.

We unexpectedly found that *C. elegans*' high speed response persists after electric shock. In

C. elegans, two other types of persistent behavioral responses related to speed change have

been reported. The first is that the animal's movement speed is elevated at high O_2

concentration in *npr-1(lf)* and in the Hawaiian wild isolate CB4856, which has the same

amino acid variation in *npr-1* (Cheung et al., 2005). In this behavioral response, (1) the

elevated speed returns rapidly to the basal speed when the high O_2 is terminated, (2) the

animals still recognize and aggregate at the edge of a food lawn, and (3) a mutation in the

tax-4 CNGC homolog for sensory depolarization abolishes the response (Coates & de Bono,

2002). Another type of persistent behavioral response is roaming (Flavell et al., 2020;

Fujiwara et al., 2002). Roaming is a behavioral state with high movement speed, although it

is only exhibited when the animals are on food and requires serotonin signaling. Because the

behavioral response to electric shock persists more than 2 min after 30-45 sec stimulus with

75 V and more than 1.5 min after only 5 sec stimulus, is not affected by food stimulus, and

does not require CNGC activity or serotonin signaling, electric shock response is likely

different from the above-mentioned two behavioral responses, and its analysis may provide a

unique opportunity for genetic dissection of a persistent behavioral state and neural activity.

The 30 or 75 V of voltage used in this study may appear artificial. However, we consider that

the responses of *C. elegans* to these stimuli reflect physiologically meaningful biological

mechanisms for the following reasons: (1) The range of voltage per length (30–75 V/6 cm =

5–12.5 V/cm) is similar to the one used to study the animal's DC response (3–12 V/cm)

(Gabel et al., 2007). (2) The electric current flowing inside the worm's body could be

weak because it depends on the resistance of its body and cuticle. (3) Only a 5 second

stimulus causes a persistent response that lasts more than a minute, meaning that the electric

shock itself is just a trigger and what we observe is a physiological response to that trigger.

(4) Fear conditioning in rodents is also triggered by electrical shock. The speed increase

behavior we observed may resemble fleeing, one of the most common responses caused by

fear in higher animals and humans (Adolphs, 2013; Bliss-Moreau, 2017; Mobbs & Kim,

2015).

Response to the electric stimulus may reflect a form of emotion

Emotions are internal brain states triggered by certain types of environmental stimuli, which

are associated with cognitive, behavioral, and physiological responses (Abbott, 2020;

Anderson & Adolphs, 2014; Nettle & Bateson, 2012; Perry & Baciadonna, 2017). Recently,

multiple species of invertebrates are considered to possess internal brain states that resemble

what we consider to be emotions (Bacqué-Cazenave et al., 2017; Cwyn et al., 2016; Gibson

et al., 2015; Hamilton et al., 2016; Mohammad et al., 2016; Pascal et al., 2014). One of the

most prominent characteristics of emotion across animal species is its persistence: For

example, even a transient environmental stimulus can cause a persistent behavioral response,

such as courtship, aggressive, and defensive behavior (Abbott, 2020; Anderson & Adolphs,

2014; Nettle & Bateson, 2012; Paul & Mendl, 2018; Perry & Baciadonna, 2017).

Anderson and Adolphs proposed a new framework to study emotions across animal species,

wherein hallmarks of an emotional state are persistence, scalability, valence, and

generalization. In addition to persistence (Fig. 4A–C), we also consider that the electric

response has negative valence. This is because the animals ignore food during the electric

shock response (Fig. 4D and E, and Sup. Fig. 5), despite the fact that food is one of the most

influential signals for *C. elegans*, affecting many aspects of their behavior. For example,

during the high speed state caused by high O_2, animals still recognize and stay at the edge of

a food lawn (Cheung et al., 2005; Coates & de Bono, 2002), suggesting that the electric

shock signal has a strong negative valence that overrides the strong positive valence of food.

The third point is the scalability—stronger stimulus causes stronger behavioral response.

Compared to the 30 V stimulus, the 75 V stimulus results in a larger number of immobile

animals during the stimulus period (right panels in Fig. 2A and B) as well as a longer-lasting

high speed response after the stimulus (Fig. 4A and C). The fourth point is generalization –

the same emotional state can be triggered by different stimuli and, in turn, the emotional state

triggered by one stimulus can then affect responses to other stimuli. The lack of response to

food during and following our electric stimulus might supports this point as well, as the

emotional state induced by electricity influences the response to food, an entirely different

stimulus.

Taken together, these results suggest that the animal's response to electric shock represents a

form of emotion, probably fear. As we revealed that the persistent aspect of the behavioral

response is regulated by neuropeptide signaling (Fig. 5 and 6), which may resemble the

neuropeptide regulation of fear in mammals including humans (Bowers et al., 2012; Comeras

et al., 2019; van den Burg & Stoop, 2019), the fear-like brain state may be regulated by

evolutionarily conserved molecular mechanisms.

In summary, we found that *C. elegans* persistently responds to electric shock, which is

regulated by voltage-gated ion channels and neuropeptide signaling. Our findings may

suggest the following model (Fig. 7). When the animals sense 30 or 75V AC stimulus at 4

Hz, the stimulus is sensed with the VGCC and BK channel and their internal state transits

from basal speed state to persistent high speed state. The persistent high speed state

eventually returns to the basal speed state, which requires neuropeptide signaling. By taking

advantage of connectome information and the methods for imaging whole brain activity of

identified neurons (Randi & Leifer, 2020; Wen et al., 2021; White et al., 1986; Yemini et al.,

2021), *C. elegans* may become one of the ideal models for revealing the dynamic information

processing involved in the entire neural circuit that regulates emotion.

Fig. 7. Model for mechanism of speed increase caused by electric shock.
394

MATERIALS AND METHODS

C. elegans **strains**

C. elegans strains were maintained with standard procedures (Brenner, 1974). In brief, for

regular cultivation, animals were grown on standard 6 cm nematode growth medium (NGM)

agar plates which had been spread with *E. coli* strain OP50 and incubated at 19.0-19.5 °C.

Strains used were the wild-type strain Bristol N2 and mutant strains PR678 *tax-4(p678)*,

CX4652 *osm-9(ky10);ocr-2(ak47)*, CB1033 *che-2(e1033)*, TU253 *mec-4(u253)*, ZB2551

mec-10(tm1552), TQ296 *trp-4(sy695)*, MT1212 *egl-19(n582)*, DA995 *egl-19(ad995)*, JD21

cca-1(ad1650), CB55 *unc-2(e55)*, VC854 *unc-2(gk366)*, NM1968 *slo-1(js379)*, BZ142 *slo-*

1(eg142), KDK11 *cat-2(tm2261)*, MT7988 *bas-1(ad446)*, GR1321 *tph-1(mg280),* RB993

tdc-1(ok914), VC671 *egl-3(ok979)*, and MT1219 *egl-3(n589)*.

C. elegans **cultivation for electric shock behavioral assay**

Before the behavioral assay, animals were cultivated as described previously (Kimura et al.,

2010). In brief, four adult wild-type animals were placed onto NGM agar plates with OP50

and kept at 19.5°C for 7.5 hours before being removed. After removal, these plates were

incubated at 19.0–19.5 °C for 3 days until the assay day. On the assay day, about 100

synchronized young adult animals were grown on each plate. As some mutant animals had

slower growth or laid fewer eggs than wild-type animals did, the incubation temperature and

number of these mutant animals were adjusted and increased accordingly in order to obtain a

comparable developmental stage (i.e. young adult) and worm number with the wild-type

animals. All behavioral assays were carried out with young adult hermaphrodites.

Experimental instruments for electric shock behavioral assay

The following electric instruments (Fig. 1) were utilized for the electric shock behavioral

assay. A 50 MHz Arbitrary Waveform Generator (FGX-295, Texio Technology Corporation)

was used to generate different types of electric waveforms over a wide range of frequencies.

This waveform generator has an output limit of 10 V. Thus, an AC Power Supply

(PCR500MA, Kikusui Electronics Corp.) was used to amplify the voltage supply. We also

used a Digital Storage Oscilloscope (DCS-1054B, Texio Technology Corporation) in parallel

to measure the voltage and observe the electric waveforms produced as well as a Digital

Multimeter (PC720M, Sanwa Electric Instrument Co., Ltd.) to measure current. A USB

camera (DMK72AUC02, The Imaging Source Co., Ltd.) with a lens (LM16JC5M2, Kowa)

was used to record trajectories produced by the animals.

Electric shock behavioral assay with small OP50 bacterial food patch

Most of the behavioral assays were conducted on 9 cm NGM agar plates seeded with a small

food patch unless indicated otherwise. For the food patch, the bacteria OP50 was grown in

100 mL of LB culture overnight at 37°C, spun down and resuspended in 10 volumes of NGM

buffer, and 5 µL of the suspension was applied at the center of the plate to create a food patch

3 × 10 mm in size on the assay day. This process was used to minimize the thickness of the

food patch as it prevents clear images of worms in the patch. Four animals per plate were

placed in the food patch one hour before the assay to accustom the animals to the

environment and to reduce their movement speed to the basal level. The assay plates were

then inverted and placed onto a custom-made copper plate bridge, whose distance is 6 cm

(Fig. 1). The images were acquired 2 frames per s, and electric shock was delivered with the

conditions described in each figure. Move-tr/2D software (Library Inc., Japan) was used to

calculate the *x-y* coordinates of the animal centroids in each image frame, which were then

analyzed using Excel (Microsoft) or R (The R Project) to calculate the animal's speed. In

general, the moving median for ±1 frame was calculated to remove noise for each animal and

then ensemble averaged for each condition. Baseline speed was calculated from the average

speed over 30 s before the stimulation, and ΔSpeed was calculated by subtracting the baseline

value from each animal's speed during or after the stimulus.

Electric shock behavioral assay with full or strip-like OP50 bacterial food lawn

For the assays conducted with full food lawn, the area of assay plates between the copper

plates were fully seeded or seeded in a stripe-shape with OP50 and kept on the bench

overnight until the assay began. Animals grown in regular cultivation plates were washed in

two droplets of NGM buffer and then transferred to the center of the assay plate and left for 5

minutes. The rest of the procedures were the same as for assays conducted with small food

patch.

To detect the outward and inward movement on the food stripes (Supplementary Fig. 5), the

food positions were first indicated on each image series by the experimenter and the animal's

centroid across the boundary was automatically calculated by a custom-made program.

Investigation of relationship among speed increase, current and voltage

Three different types of NGM agar plates were prepared with varying salt concentrations and

similar osmolarities: High-salt plates had 200 mM sodium chloride; low-salt plates had 10

mM sodium chloride and 380 mM sucrose; control plates had 50 mM sodium chloride and

300 mM sucrose. The purpose of adding sucrose into the plates was to adjust and balance the

osmolarity. The final total osmolarity for sodium chloride (Na^+ and Cl^-) and sucrose for all

the plates was 400 mOsm. The rest of the procedures were the same as for assays conducted

with small food patch.

Data analysis and statistics

All the statistical analyses were performed in R (The R Project). Generally, data of 20 – 50

animals in total from 9 plates from 3 days of experiments for each condition were pooled and

analyzed together. We chose this sample number based on a large scale behavioral analysis of

C. elegans (Yemini et al., 2013). Data is presented as means ± SD unless otherwise specified.

Experimental conditions, such as the electric stimulation or different strains were randomized

on a daily basis.

ACKNOWLEDGEMENT:

We thank Liting Chen for having provided the idea of "worm's emotion" for K.D.K., Yuki

Tanimoto and Yuka Tsuda for the initial phase of the electric shock paradigm, Shinobu

Aoyagi for setting up the system, Chentao Wen for developing the outward/inward scoring

program, and Young-Jai You, Aki Takahashi, Atsuko Nakagawa and the Kimura lab

members for their valuable advice, comments and technical assistance for the study.

Nematode strains were provided by the Caenorhabditis Genetics Center (funded by the NIH

Office of Research Infrastructure Programs P40 OD010440).

FUNDING:

This study was supported by Japan Society for the Promotion of Science (KAKENHI

JP16H06545, 20H05700, 21H00448 and 21K19274 to K.D.K.), Grant-in-Aid for Research in

Nagoya City University (48, 1912011, 1921102 and 2121101), the Joint Research by

National Institutes of Natural Sciences (01112002), Toyoaki Scholarship Foundation, and

RIKEN Center for Advanced Intelligence Project (to K.D.K). L.F.T. was supported by

Japanese Government (MEXT) Scholarship.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES:

Abbott, A. (2020). Inside the mind of an animal. *Nature*, *584*, 182–185.

Adolphs, R. (2013). The Biology of Fear. *Curr Biol.*, *23*(2), 79–93.

https://doi.org/10.1016/j.cub.2012.11.055.The

Alkema, M. J., Hunter-Ensor, M., Ringstad, N., & Horvitz, H. R. (2005). Tyramine Functions

Independently of Octopamine in the Caenorhabditis elegans Nervous System. *Neuron*,

46(2), 247–260. https://doi.org/https://doi.org/10.1016/j.neuron.2005.02.024

Anderson, D. J. (2016). Circuit modules linking internal states and social behaviour in flies

and mice. *Nature Reviews Neuroscience*, *17*(11), 692–704.

https://doi.org/10.1038/nrn.2016.125

Anderson, D. J., & Adolphs, R. (2014). A Framework for Studying Emotions across Species.

Cell, *157*(1), 187–200. https://doi.org/10.1016/j.cell.2014.03.003

Bacqué-Cazenave, J., Cattaert, D., Delbecque, J.-P., & Fossat, P. (2017). Social harassment

induces anxiety-like behaviour in crayfish. *Scientific Reports*, *7*(1), 39935.

https://doi.org/10.1038/srep39935

Bargmann, C. I. (2006). Chemosensation in C. elegans. *WormBook : The Online Review of C.

Elegans Biology*, 1–29. https://doi.org/https://doi.org/10.1895/wormbook.1.123.1

Bargmann, C. I., Hartwieg, E., & Horvitz, H. R. (1993). Odorant-selective genes and neurons

mediate olfaction in C. elegans. *Cell*, *74*(3), 515–527.

https://doi.org/https://doi.org/10.1016/0092-8674(93)80053-H

Bellono, N. W., Leitch, D. B., & Julius, D. (2017). Molecular basis of ancestral vertebrate

electroreception. *Nature*, *543*(7645), 391–396. https://doi.org/10.1038/nature21401

Bellono, N. W., Leitch, D. B., & Julius, D. (2018). Molecular tuning of electroreception in

sharks and skates. *Nature*, *558*(7708), 122–126. https://doi.org/10.1038/s41586-018-

0160-9

Ben Arous, J., Laffont, S., & Chatenay, D. (2009). Molecular and Sensory Basis of a Food

Related Two-State Behavior in C. elegans. *PLOS ONE*, *4*(10), e7584.

https://doi.org/10.1371/journal.pone.0007584

Berridge, K. C. (2004). Motivation concepts in behavioral neuroscience. *Physiology &

Behavior*, *81*(2), 179–209. https://doi.org/https://doi.org/10.1016/j.physbeh.2004.02.004

Bliss-Moreau, E. (2017). Constructing nonhuman animal emotion. *Current Opinion in

Psychology*, *17*, 184–188. https://doi.org/10.1016/j.copsyc.2017.07.011

Bowers, M. E., Choi, D. C., & Ressler, K. J. (2012). Neuropeptide regulation of fear and

anxiety: Implications of cholecystokinin, endogenous opioids, and neuropeptide Y.

Physiology & Behavior, *107*(5), 699–710.

https://doi.org/10.1016/j.physbeh.2012.03.004

Brenner, S. (1974). The Genetics of *Caenorhabditis elegans*. *Genetics*, *77*(1), 71 LP – 94.

http://www.genetics.org/content/77/1/71.abstract

Chen, P., & Hong, W. (2018). Neural Circuit Mechanisms of Social Behavior. *Neuron*, *98*(1),

16–30. https://doi.org/10.1016/j.neuron.2018.02.026

Cheung, B. H. H., Cohen, M., Rogers, C., Albayram, O., & de Bono, M. (2005). Experience-

Dependent Modulation of C. elegans Behavior by Ambient Oxygen. *Current Biology*,

15(10), 905–917. https://doi.org/https://doi.org/10.1016/j.cub.2005.04.017

Clarke, D., Whitney, H., Sutton, G., & Robert, D. (2013). Detection and Learning of Floral

Electric Fields by Bumblebees. *Science*, *340*(6128), 66 LP – 69.

https://doi.org/10.1126/science.1230883

Coates, J. C., & de Bono, M. (2002). Antagonistic pathways in neurons exposed to body fluid

regulate social feeding in Caenorhabditis elegans. *Nature*, *419*(6910), 925–929.

https://doi.org/10.1038/nature01170

Coburn, C. M., & Bargmann, C. I. (1996). A Putative Cyclic Nucleotide–Gated Channel Is

Required for Sensory Development and Function in C. elegans. *Neuron*, *17*(4), 695–706.

https://doi.org/https://doi.org/10.1016/S0896-6273(00)80201-9

Colbert, H. A., Smith, T. L., & Bargmann, C. I. (1997). OSM-9, A Novel Protein with

Structural Similarity to Channels, Is Required for Olfaction, Mechanosensation, and

Olfactory Adaptation in Caenorhabditis elegans. *The Journal of Neuroscience*, *17*(21),

8259 LP – 8269. https://doi.org/10.1523/JNEUROSCI.17-21-08259.1997

Comeras, L. B., Herzog, H., & Tasan, R. O. (2019). Neuropeptides at the crossroad of fear

and hunger: a special focus on neuropeptide Y. *Annals of the New York Academy of*

Sciences, *1455*(1), 59–80. https://doi.org/10.1111/nyas.14179

Crampton, W. G. R. (2019). Electroreception, electrogenesis and electric signal evolution.

Journal of Fish Biology, *95*(1), 92–134. https://doi.org/10.1111/jfb.13922

Cwyn, S., Luigi, B., & Lars, C. (2016). Unexpected rewards induce dopamine-dependent

positive emotion–like state changes in bumblebees. *Science*, *353*(6307), 1529–1531.

https://doi.org/10.1126/science.aaf4454

Davies, A. G., Pierce-Shimomura, J. T., Kim, H., VanHoven, M. K., Thiele, T. R., Bonci, A.,

Bargmann, C. I., & McIntire, S. L. (2003). A Central Role of the BK Potassium Channel

in Behavioral Responses to Ethanol in C. elegans. *Cell*, *115*(6), 655–666.

https://doi.org/10.1016/S0092-8674(03)00979-6

de Bono, M., & Maricq, A.V. (2005). NEURONAL SUBSTRATES OF COMPLEX

BEHAVIORS IN C. ELEGANS. *Annual Review of Neuroscience*, *28*(1), 451–501.

https://doi.org/10.1146/annurev.neuro.27.070203.144259

Driscoll, M., & Chalfie, M. (1991). The mec-4 gene is a member of a family of

Caenorhabditis elegans genes that can mutate to induce neuronal degeneration. *Nature*,

349(6310), 588–593. https://doi.org/10.1038/349588a0

Flavell, S. W., Raizen, D. M., & You, Y.-J. (2020). Behavioral States. *Genetics*, *216*(2), 315–

332. https://doi.org/10.1534/genetics.120.303539

Fujiwara, M., Ishihara, T., & Katsura, I. (1999). A novel WD40 protein, CHE-2, acts cell-

autonomously in the formation of C. elegans sensory cilia. *Development*, *126*(21), 4839–

4848. https://doi.org/10.1242/dev.126.21.4839

Fujiwara, M., Sengupta, P., & McIntire, S. L. (2002). Regulation of body size and behavioral

state of C. elegans by sensory perception and the egl-4 cGMP-dependent protein kinase.

Neuron, *36*(6), 1091–1102. https://doi.org/10.1016/S0896-6273(02)01093-0

Gabel, C. V., Gabel, H., Pavlichin, D., Kao, A., Clark, D. A., & Samuel, A. D. T. (2007).

Neural Circuits Mediate Electrosensory Behavior in Caenorhabditis elegans. *Journal of Neuroscience*, 27(28), 7586–7596. https://doi.org/10.1523/JNEUROSCI.0775-07.2007

Gibson, W. T., Gonzalez, C. R., Fernandez, C., Ramasamy, L., Tabachnik, T., Du, R. R., Felsen, P. D., Maire, M. R., Perona, P., & Anderson, D. J. (2015). Behavioral Responses to a Repetitive Visual Threat Stimulus Express a Persistent State of Defensive Arousal in Drosophila. *Current Biology*, 25(11), 1401–1415. https://doi.org/https://doi.org/10.1016/j.cub.2015.03.058

Hamilton, T. J., Kwan, G. T., Gallup, J., & Tresguerres, M. (2016). Acute fluoxetine exposure alters crab anxiety-like behaviour, but not aggressiveness. *Scientific Reports*, 6(1), 19850. https://doi.org/10.1038/srep19850

Hoopfer, E. D., Jung, Y., Inagaki, H. K., Rubin, G. M., & Anderson, D. J. (2015). P1 interneurons promote a persistent internal state that enhances inter-male aggression in Drosophila. *ELife*, 4, e11346. https://doi.org/10.7554/eLife.11346

Huang, M., & Chalfie, M. (1994). Gene interactions affecting mechanosensory transduction in Caenorhabditis elegans. *Nature*, 367(6462), 467–470. https://doi.org/10.1038/367467a0

Istenič, L., & Bulog, B. (1984). Some evidence for the ampullary organs in the European cave salamander Proteus anguinus (Urodela, Amphibia). *Cell and Tissue Research*, 235(2), 393–402. https://doi.org/10.1007/BF00217865

Kang, L., Gao, J., Schafer, W. R., Xie, Z., & Xu, X. Z. S. (2010). C. elegans TRP Family Protein TRP-4 Is a Pore-Forming Subunit of a Native Mechanotransduction Channel. *Neuron*, 67(3), 381–391. https://doi.org/https://doi.org/10.1016/j.neuron.2010.06.032

Kass, J., Jacob, T. C., Kim, P., & Kaplan, J. M. (2001). The EGL-3 Proprotein Convertase Regulates Mechanosensory Responses of Caenorhabditis elegans. *The Journal of Neuroscience*, 21(23), 9265 LP – 9272. https://doi.org/10.1523/JNEUROSCI.21-23-

09265.2001

Kennedy, A., Asahina, K., Hoopfer, E., Inagaki, H., Jung, Y., Lee, H., Remedios, R., &
Anderson, D. J. (2014). Internal States and Behavioral Decision-Making: Toward an
Integration of Emotion and Cognition. *Cold Spring Harbor Symposia on Quantitative*
Biology, *79*, 199–210. https://doi.org/10.1101/sqb.2014.79.024984

Kimura, K. D., Fujita, K., & Katsura, I. (2010). Enhancement of Odor Avoidance Regulated
by Dopamine Signaling in Caenorhabditis elegans. *The Journal of Neuroscience*, *30*(48),
16365 LP – 16375. https://doi.org/10.1523/JNEUROSCI.6023-09.2010

Komatsu, H., Mori, I., Rhee, J.-S., Akaike, N., & Ohshima, Y. (1996). Mutations in a Cyclic
Nucleotide–Gated Channel Lead to Abnormal Thermosensation and Chemosensation in
C. elegans. *Neuron*, *17*(4), 707–718. https://doi.org/https://doi.org/10.1016/S0896-
6273(00)80202-0

Lee, R. Y. N., Lobel, L., Hengartner, M., Horvitz, H. R., & Avery, L. (1997). Mutations in
the α1 subunit of an L-type voltage-activated Ca2+ channel cause myotonia in
Caenorhabditis elegans. *The EMBO Journal*, *16*(20), 6066–6076.
https://doi.org/https://doi.org/10.1093/emboj/16.20.6066

Lee, S.-H., & Dan, Y. (2012). Neuromodulation of Brain States. *Neuron*, *76*(1), 209–222.
https://doi.org/https://doi.org/10.1016/j.neuron.2012.09.012

Lints, R., & Emmons, S. W. (1999). Patterning of dopaminergic neurotransmitter identity
among Caenorhabditis elegans ray sensory neurons by a TGFbeta family signaling
pathway and a Hox gene. *Development*, *126*(24), 5819–5831.
https://doi.org/10.1242/dev.126.24.5819

Loer, C. M., & Kenyon, C. J. (1993). Serotonin-deficient mutants and male mating behavior
in the nematode Caenorhabditis elegans. *The Journal of Neuroscience*, *13*(12), 5407 LP
– 5417. https://doi.org/10.1523/JNEUROSCI.13-12-05407.1993

Maimon, G. (2011). Modulation of visual physiology by behavioral state in monkeys, mice,

and flies. *Current Opinion in Neurobiology*, *21*(4), 559–564.

https://doi.org/https://doi.org/10.1016/j.conb.2011.05.001

Mobbs, D., & Kim, J. J. (2015). Neuroethological studies of fear, anxiety, and risky decision-

making in rodents and humans. *Current Opinion in Behavioral Sciences*, *5*, 8–15.

https://doi.org/10.1016/j.cobeha.2015.06.005

Mohammad, F., Aryal, S., Ho, J., Stewart, J. C., Norman, N. A., Tan, T. L., Eisaka, A., &

Claridge-Chang, A. (2016). Ancient Anxiety Pathways Influence Drosophila Defense

Behaviors. *Current Biology : CB*, *26*(7), 981–986.

https://doi.org/10.1016/j.cub.2016.02.031

Nettle, D., & Bateson, M. (2012). The Evolutionary Origins of Mood and Its Disorders.

Current Biology, *22*(17), R712–R721.

https://doi.org/https://doi.org/10.1016/j.cub.2012.06.020

Pascal, F., Julien, B.-C., Philippe, D. D., Jean-Paul, D., & Daniel, C. (2014). Anxiety-like

behavior in crayfish is controlled by serotonin. *Science*, *344*(6189), 1293–1297.

https://doi.org/10.1126/science.1248811

Paul, E. S., & Mendl, M. T. (2018). Animal emotion: Descriptive and prescriptive definitions

and their implications for a comparative perspective. *Applied Animal Behaviour Science*,

205, 202–209. https://doi.org/https://doi.org/10.1016/j.applanim.2018.01.008

Perry, C. J., & Baciadonna, L. (2017). Studying emotion in invertebrates: what has been

done, what can be measured and what they can provide. *Journal of Experimental*

Biology, *220*(21), 3856–3868. https://doi.org/10.1242/jeb.151308

Pettigrew, J. D. (1999). Electroreception in monotremes. *Journal of Experimental Biology*,

202(10), 1447–1454. https://doi.org/10.1242/jeb.202.10.1447

Quinn, W. G., Harris, W. A., & Benzer, S. (1974). Conditioned behavior in Drosophila

melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *71*(3), 708–712. https://doi.org/10.1073/pnas.71.3.708

Raizen, D. M., Zimmerman, J. E., Maycock, M. H., Ta, U. D., You, Y. J., Sundaram, M. V., & Pack, A. I. (2008). Lethargus is a Caenorhabditis elegans sleep-like state. *Nature*, *451*(7178), 569–572. https://doi.org/10.1038/nature06535

Randi, F., & Leifer, A. M. (2020). Measuring and modeling whole-brain neural dynamics in Caenorhabditis elegans. *Current Opinion in Neurobiology*, *65*, 167–175. https://doi.org/https://doi.org/10.1016/j.conb.2020.11.001

Rescorla, R. (1968). Probability of shock in the presence and absence of CS in fear conditioning. *Journal of Comparative and Physiological Psychology*, *66 1*, 1–5.

Roth, A., & Schlegel, P. (1988). Behavioral Evidence and Supporting Electrophysiological Observations for Electroreception in the Blind Cave Salamander, *Proteus anguinus* (Urodela). *Brain, Behavior and Evolution*, *32*(5), 277–280. https://doi.org/10.1159/000116556

Sasakura, H., & Mori, I. (2013). Behavioral plasticity, learning, and memory in C. elegans. *Current Opinion in Neurobiology*, *23*(1), 92–99. https://doi.org/https://doi.org/10.1016/j.conb.2012.09.005

Sawin, E. R., Ranganathan, R., & Horvitz, H. R. (2000). C. elegans Locomotory Rate Is Modulated by the Environment through a Dopaminergic Pathway and by Experience through a Serotonergic Pathway. *Neuron*, *26*(3), 619–631. https://doi.org/https://doi.org/10.1016/S0896-6273(00)81199-X

Schafer, W. R., & Kenyon, C. J. (1995). A calcium-channel homologue required for adaptation to dopamine and serotonin in Caenorhabditis elegans. *Nature*, *375*(6526), 73–78. https://doi.org/10.1038/375073a0

Scheich, H., Langner, G., Tidemann, C., Coles, R. B., & Guppy, A. (1986). Electroreception

and electrolocation in platypus. *Nature*, *319*(6052), 401–402.

https://doi.org/10.1038/319401a0

Steger, K. A., Shtonda, B. B., Thacker, C., Snutch, T. P., & Avery, L. (2005). The C. elegans

T-type calcium channel CCA-1 boosts neuromuscular transmission. *Journal of*

Experimental Biology, *208*(11), 2191–2203. https://doi.org/10.1242/jeb.01616

Sukul, N. C., & Croll, N. A. (1978). Influence of potential difference and current on the

electrotaxis of Caenorhaditis elegans. *Journal of Nematology*, *10*(4), 314–317.

http://www.ncbi.nlm.nih.gov/pubmed/19305860%0Ahttp://www.pubmedcentral.nih.gov

/articlerender.fcgi?artid=PMC2617915

Sutton, G. P., Clarke, D., Morley, E. L., & Robert, D. (2016). Mechanosensory hairs in

bumblebees (Bombus terrestris) detect weak electric fields. *Proceedings of the National*

Academy of Sciences, *113*(26), 7261 LP – 7265.

https://doi.org/10.1073/pnas.1601624113

Sze, J. Y., Victor, M., Loer, C., Shi, Y., & Ruvkun, G. (2000). Food and metabolic signalling

defects in a Caenorhabditis elegans serotonin-synthesis mutant. *Nature*, *403*(6769), 560–

564. https://doi.org/10.1038/35000609

Tobin, D. M., Madsen, D. M., Kahn-Kirby, A., Peckol, E. L., Moulder, G., Barstead, R.,

Maricq, A. V, & Bargmann, C. I. (2002). Combinatorial Expression of TRPV Channel

Proteins Defines Their Sensory Functions and Subcellular Localization in C. elegans

Neurons. *Neuron*, *35*(2), 307–318. https://doi.org/https://doi.org/10.1016/S0896-

6273(02)00757-2

van den Burg, E. H., & Stoop, R. (2019). Neuropeptide signalling in the central nucleus of

the amygdala. *Cell and Tissue Research*, *375*(1), 93–101.

https://doi.org/10.1007/s00441-018-2862-6

von der Emde, G., Amey, M., Engelmann, J., Fetz, S., Folde, C., Hollmann, M., Metzen, M.,

& Pusch, R. (2008). Active electrolocation in Gnathonemus petersii: Behaviour, sensory

performance, and receptor systems. *Journal of Physiology Paris*, *102*(4–6), 279–290.

https://doi.org/10.1016/j.jphysparis.2008.10.017

Wang, Z. W., Saifee, O., Nonet, M. L., & Salkoff, L. (2001). SLO-1 potassium channels

control quantal content of neurotransmitter release at the C. elegans neuromuscular

junction. *Neuron*, *32*(5), 867–881. https://doi.org/10.1016/S0896-6273(01)00522-0

Wark, B., Lundstrom, B. N., & Fairhall, A. (2007). Sensory adaptation. *Current Opinion in*

Neurobiology, *17*(4), 423–429. https://doi.org/10.1016/j.conb.2007.07.001

Wen, C., Miura, T., Voleti, V., Yamaguchi, K., Tsutsumi, M., Yamamoto, K., Otomo, K.,

Fujie, Y., Teramoto, T., Ishihara, T., Aoki, K., Nemoto, T., Hillman, E. M. C., &

Kimura, K. D. (2021). 3DeeCellTracker, a deep learning-based pipeline for segmenting

and tracking cells in 3D time lapse images. *ELife*, *10*, e59187.

https://doi.org/10.7554/eLife.59187

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the

nervous system of the nematode Caenorhabditis elegans. In *Philosophical Transactions*

of the Royal Society of London. B, Biological Sciences (Vol. 314, Issue 1165). Royal

Society. https://doi.org/10.1098/rstb.1986.0056

Yemini, E., Jucikas, T., Grundy, L. J., Brown, A. E. X., & Schafer, W. R. (2013). A database

of Caenorhabditis elegans behavioral phenotypes. *Nature Methods*, *10*(9), 877–879.

https://doi.org/10.1038/nmeth.2560

Yemini, E., Lin, A., Nejatbakhsh, A., Varol, E., Sun, R., Mena, G. E., Samuel, A. D. T.,

Paninski, L., Venkatachalam, V., & Hobert, O. (2021). NeuroPAL: A Multicolor Atlas

for Whole-Brain Neuronal Identification in C. elegans. *Cell*, *184*(1), 272-288.e11.

https://doi.org/https://doi.org/10.1016/j.cell.2020.12.012