Common and distinct neurofunctional representations of core and social disgust in the brain: An ALE meta-analysis and MACM characterization

Authors

Xianyang Gan[1], Xinqi Zhou[1], Jialin Li[1,2], Guojuan Jiao[1], Xi Jiang[1], Bharat Biswal[1,3], Shuxia Yao[1], Benjamin Klugah-Brown[1]*, Benjamin Becker[1]*

Affiliations

[1]The Clinical Hospital of Chengdu Brain Science Institute, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, Sichuan 610054, China

[2]Max Planck School of Cognition, Leipzig 04103, Germany

[3]Department of Biomedical Engineering, New Jersey Institute of Technology, New Jersey 7102, United States of America

*Correspondence to

Benjamin Klugah-Brown (bklugah@gmail.com) or Benjamin Becker (ben_becker@gmx.de)

University of Electronic Science and Technology

School of Life Science and Technology

No.2006, Xiyuan Ave

Chengdu, Sichuan

611731, China

ABSTRACT

Disgust represents a multifaceted defensive-avoidance response. On the behavioral level, the response includes withdrawal and a disgust-specific facial expression. While both serve the avoidance of pathogens the latter additionally transmits social-communicative information. Given that common and distinct brain representation of the primary defensive-avoidance response (core disgust) and encoding of the social-communicative signal (social disgust) remain debated we employed neuroimaging meta-analyses to (1) determine brain systems generally engaged in disgust processing, and (2) segregate common and distinct brain systems for core and social disgust. Disgust processing, in general, engaged a bilateral network encompassing the insula, amygdala, occipital and prefrontal regions. Core disgust evoked stronger reactivity in left-lateralized threat detection and defensive response network including amygdala, occipital and frontal regions while social disgust engaged a right-lateralized superior temporal-frontal network engaged in social cognition. Anterior insula, inferior frontal and fusiform regions were commonly engaged during core and social disgust suggesting a common neural basis. We demonstrate a common and separable neural basis of primary disgust responses and encoding of associated social-communicative signals.

Keywords: disgust, fMRI, meta-analysis, activation likelihood estimation (ALE), Meta-analytic connectivity modeling (MACM), defensive-avoidance response, social cognition, face, insula, amygdala

1. INTRODUCTION

Disgust represents a basic emotion and has been conceptualized as a multifaceted and evolutionary adaptive defensive-avoidance response. The disgust response may have initially evolved in the context of rejection of potential poisonous food (Darwin, 1965; Rozin & Fallon, 1987; Rozin, Haidt, & Fincher, 2009) but the same underlying mechanism may have subsequently been adapted to facilitate avoidance of contaminations by pathogens and may even serve the avoidance of harmful social interactions (Curtis, de Barra, & Aunger, 2011; Davey, 2011; Haidt, McCauley, & Rozin, 1994; Kupfer & Fessler, 2018; Rozin & Fallon, 1987; Rozin, Haidt, & McCauley, 2016; Vicario, Rafal, Martino, & Avenanti, 2017). In line with its evolutionary function, disgust is primarily elicited by exposure to potentially poisonous or infectious stimuli such as rotten food, cockroaches or individuals with poor personal hygiene as well as injury- and mutilation-related stimuli (Haidt et al., 1994; Olatunji & Sawchuk, 2005; Rozin & Fallon, 1987; Vicario et al., 2017). Exposure to these stimuli induces a distinct disgust reaction on different levels encompassing subjective experience of revulsion, as well as physiological and behavioral responses. The behavioral response includes rapid withdrawal from the eliciting stimulus as well as a distinct facial expression characterized by a wrinkled nose (Rozin, Lowery, & Ebert, 1994) and raised upper lip (Rozin et al., 1994) reflecting the primary purpose of the disgust reaction in terms of avoiding or rejecting pathogens (Chapman, Kim, Susskind, & Anderson, 2009; Oaten, Stevenson, & Case, 2009; Olatunji & Sawchuk, 2005; Rozin & Fallon, 1987; Rozin et al., 2009; Rozin et al., 1994; Schienle & Wabnegger, 2021) (in line with Vicario et al., (2017) henceforward referred to as *core disgust* processing). The highly distinct facial expression for disgust additionally serves social-communicative purposes (Chapman & Anderson, 2012; Chapman et al., 2009; Vicario et al., 2017; von dem Hagen et al., 2009) such that a disgusted facial expression signals to conspecifics a danger of being exposed to potential pathogens (Burklund, Eisenberger, & Lieberman, 2007; Darwin, 1965; Rozin et al., 1994; Surguladze et al., 2003; von dem Hagen et al., 2009) or an attitude of social rejection (Burklund et al., 2007; Surguladze et al., 2010; von dem Hagen et al., 2009). Accurate decoding of these signals is essential for survival and adaptive social interactions (Darwin, 1965; Ekman, 1993; Fusar-Poli et al., 2009; Malhi et al., 2007; Phillips et al., 1999) (in line with Vicario et al., (2017) perception of a disgusted face as a social-communicative signal is henceforward referred to as *social disgust* processing). Although both, core and social disgust, share the essential function of a defensive-avoidance reaction towards potential pathogen exposure the facial expression of disgust as social signal represents a more indirect and context depended signal which requires further interpretation by the observer (for similar reasoning with respect to other facial expression domains see e.g., Hadjistavropoulos et al., 2011; Zhou, Li, et al., 2020).

Initial evidence for common and separable neurobiological systems underlying core and social disgust responses additionally comes from human lesion models reporting that a patient with focal anterior insula and putamen lesions exhibited intact core disgust reactivity but exhibited selective impairments in recognizing disgust from facial signals (Calder, Keane, Manes, Antoun, & Young, 2000) while a patient with selective lesions of the anterior insula exhibited both, dysregulated core disgust reactivity as well as altered production of disgust expressions (Cantone et al., 2019). A recent study in a larger sample of lesion patients confirmed core as well as social disgust processing impairments in patients with insula-basal ganglia lesions and additionally reported that patients with left hemispheric lesions exhibited pronounced

impairments suggesting a lateralization of disgust processes (Holtmann et al., 2020).

Indirect evidence for common and separable neurobiological systems comes from numerous functional imaging, particularly functional MRI, studies that examined the neural substrates of disgust processing in healthy subjects (the first functional MRI study on disgust was published in 1997) (Phillips et al., 1997). These studies demonstrated that visual, gustatory (Corradi-Dell'Acqua, Tusche, Vuilleumier, & Singer, 2016; Jabbi, Bastiaansen, & Keysers, 2008), olfactory (Berlin et al., 2017; Heining et al., 2004; Meier et al., 2015; Wicker et al., 2003), auditory (Phillips et al., 1998; Zimmer, Höfler, Koschutnig, & Ischebeck, 2016) stimuli can induce the subjective experience of disgust and neural reactivity in the domains of core and social disgust reactivity.

Visual stimuli, such as pictures and film clips depicting core disgust elicitors (e.g., rotten food) or social disgust stimuli (i.e., disgusted faces) in combination with corresponding non-disgust related stimuli have been widely used in separate studies on core or social disgust. However, despite the large number of original studies the key regions involved in disgust in general, as well as core and social disgust in particular, remain strongly debated. For instance, while the insula has been considered and often reported as a key region for general disgust processing (Calder, Lawrence, & Young, 2001; Kirby & Robinson, 2017; Murphy, Nimmo-Smith, & Lawrence, 2003; Phan, Wager, Taylor, & Liberzon, 2002; Vytal & Hamann, 2010) and insula activity has been reported during core (Schäfer, Leutgeb, Reishofer, Ebner, & Schienle, 2009; Schäfer, Schienle, & Vaitl, 2005; Schienle, Höfler, Keck, & Wabnegger, 2020; Schienle et al., 2002; Shapira et al., 2003; Stark et al., 2007) as well as social disgust processing paradigms (Ashworth et al., 2011; Phillips et al., 1997; Wicker et al., 2003), several studies examining disgust processes did not observe insula activity (Deeley et al., 2008; Gorno-Tempini et al., 2001; Schienle et al., 2006; Schienle, Schäfer, Stark, Walter, & Vaitl, 2005b; Stark et al., 2005; Stark et al., 2004; Stark et al., 2003; Trautmann, Fehr, & Herrmann, 2009) which may reflect a general involvement of the insula in interoceptive processes rather than a disgust specific engagement (Li et al., 2019; Stark et al., 2005; Yao, Becker, et al., 2018).

Moreover, several studies suggest a role of the amygdala in disgust processing, probably related to its role in early detection of threat and initiation of defensive responses (e.g., Becker et al., 2012; Zhou, Zhao, et al., 2020). Amygdala activity in response to core disgust stimuli (Pujol et al., 2018; Schienle, Schäfer, Hermann, & Vaitl, 2009; Schienle et al., 2006; Schienle et al., 2002; Schienle, Übel, Schöngaßner, Ille, & Scharmüller, 2014; Stark et al., 2005; Stark et al., 2003; Stark et al., 2007) as well as social disgust stimuli (Burklund et al., 2007; Fitzgerald, Angstadt, Jelsone, Nathan, & Phan, 2006; Phillips et al., 1998; Winston, O'Doherty, & Dolan, 2003) has been reported in several studies, yet a considerable number of studies on the neural basis of disgust failed to find an engagement of the amygdala (Caseras et al., 2007; Deeley et al., 2007; Phillips et al., 2000; Phillips et al., 2004; Phillips et al., 1999; Phillips et al., 1997; Schienle & Scharmüller, 2013; Schroeder et al., 2005; Sprengelmeyer, Rausch, Eysel, & Przuntek, 1998; Surguladze et al., 2010; Wicker et al., 2003). Finally, two studies included different visual disgust stimuli (i.e., core disgust and social disgust stimuli), yet surprisingly did not find robust social disgust related activity or common neural activity during social and core disgust processing, respectively (Schäfer et al., 2005; Schienle et al., 2020).

The ongoing debates about the brain systems generally engaged in disgust processing as well as during core or social disgust processing separately might be partly rooted in the limitations of single fMRI studies and particularly factors such as low and underpowered sample

sizes, variations in processing depths (i.e., implicit vs explicit processing), stimuli sets, experimental designs (e.g., block design vs event-related design), study-specific samples with variations in cultural background or disgust propensity, field strength (e.g., 1T vs 3T) and methods of data analysis may have considerably contributed to the lack of robustly identified neural systems for disgust. The development and application of neuroimaging based meta-analyses have allowed to overcome these limitations by capitalizing on the results from previously published studies (Eickhoff, Bzdok, Laird, Kurth, & Fox, 2012; Laird et al., 2005; Wager, Lindquist, & Kaplan, 2007) thus allowing to determine the convergent and robust neurofunctional basis of emotional and cognitive processes as well as to determine common and distinct neurofunctional underpinnings between processes (e.g., Chen et al., 2018; Feng et al., 2018). Previous studies have capitalized on neuroimaging meta-analyses to successfully determine common and separable brain representations of core emotions or emotional face expressions (Fusar-Poli et al., 2009; Kirby & Robinson, 2017; Murphy et al., 2003; Phan et al., 2002; Vytal & Hamann, 2010).

By capitalizing on neuroimaging meta-analyses, we aimed at determining brain systems that are robustly engaged in general disgust processing, core and social disgust processing as well as to further segregate systems that are common versus specifically involved in core or social disgust processing respectively. Based on overarching disgust frameworks we hypothesized that disgust-related processing will generally recruit regions engaged in motivational aspects, defensive responses and interoception such as the insula, amygdala, prefrontal and occipital regions. Based on the previous literature we further expected that common and separable (domain-specific) networks would underly core and social disgust processes, such that the insula and orbitofrontal regions would be recruited across domains while regions involved in social and self-referential processing would be differentially engaged.

2. METHODS

2.1 Literature search

The present meta-analytic review was conducted in line with the PRISMA guidelines for systematic reviews (Page et al., 2021) and the meta-analytic protocols were preregistered on the Open Science Framework (https://osf.io/pu59t).

Articles published prior to May 2021 were identified using the following keywords: ("functional magnetic resonance imaging" OR "fMRI") AND ("disgust facial expression" OR "disgusted facial expression" OR "disgust face" OR "disgusted face" OR "facial motion, disgust" OR "disgust elicitors" OR "disgusting elicitors" OR "disgust stimuli" OR "disgusting stimuli" OR "disgust visual scenes" OR "disgusting visual scenes" OR "disgust") on the PubMed and Web of Science databases. The search was complemented by a corresponding search on the BrainMap functional database (May 2021; containing 3406 papers and 16901 experiments) using the Sleuth software (version 3.0.4, https://www.brainmap.org/sleuth/) with the following search terms: "Experiments: Context, Normal Mapping", "Experiments: Activations, Activations only", "Experiments: Imaging Modality, fMRI" and "Experiments: Behavioral Domain, Emotion, Negative-Disgust". Finally, reference lists of included articles were searched for relevant articles not identified during the initial database search. Following removal of duplicates 419 neuroimaging papers on disgust processing were retained.

2.2 Inclusion and exclusion criteria

To identify suitable publications according to our criteria titles and abstracts of identified papers were initially examined by two independent reviewers (X. G. and X. Z.). Full texts of eligible papers were downloaded for detailed examination. Any disagreement between these two reviewers was resolved by discussion and a final decision by a third independent reviewer (J. L.). Papers that passed the screening process underwent a thorough and detailed evaluation of eligibility (B. B. and X. G.), studies with ambiguous experimental contrasts were removed at this stage and disagreements were further resolved by another reviewer (B. K-B.). A flowchart illustrating the detailed article screening and exclusion process is shown in **Figure 1**. The following inclusion criteria were applied: (1) studies reporting neuroimaging experiments on disgust processing which presented static pictures or film clips displaying clearly disgust-related contents (e.g., disgusted facial expressions and disgusting visual scenes, to limit variability related to induction modality studies inducing disgust processing via other modalities, such as olfactory, gustatory and auditory stimuli or script-induced imagination were not considered); (2) results were published as original papers in peer-reviewed English-language journals; (3) contrasts for healthy subjects were reported (i.e., healthy volunteers only). Studies reporting only spatial coordinates across or between healthy controls and clinical groups were excluded, but included if data was separately reported for the healthy control group; (4) task-related fMRI was employed; (5) activation contrasts generated by the subtraction approach represent main effects of disgust relative to a baseline or control condition (e.g., disgust faces > neutral faces, disgust elicitors > fixation); (6) activation in terms of higher activity for the disgust-related stimuli reported (deactivations were not included in the analysis because the underlying functional process remains difficult to interpret); (7) results reported in standard stereotactic coordinates (either Talairach or Montreal Neurological Institute [MNI] space); (8) results describing functional, psychophysiological or psychopathological correlations were excluded; (9) other types of emotional faces including schematic faces were excluded; (10) whole-brain results were reported (we excluded studies reporting only region-of-interest, ROI, analyses); (11) for studies reporting multiple but not independent contrasts, only the contrast that substantially reflected the process of interest was included. For instance, if two studies employed disgust stimuli with different intensity levels, only the results corresponding to highest disgust intensity were included (similar approach see Tao, He, Lin, Liu, & Tao, 2021); (12) for studies that included multiple relevant and independent contrasts, all such contrasts were included (similar approach see Tao et al., 2021).

To reduce the possibility of biased study sampling we contacted the corresponding authors of 11 studies for which Talairach or MNI coordinates for the appropriate contrast were not reported (e.g., for healthy controls in a case-control neuroimaging study).

After the in-depth screening procedure, a total of 69 papers (**Supplementary Table 1**) were included in the meta-analyses, and according to the aims of the analyses organized as follows: (1) general disgust processing: 69 papers, 71 experiments, 717 foci, 78 contrasts, 1409 subjects; (2) core disgust processing: 39 papers, 41experiments, 402 foci, 46 contrasts, 897 subjects; (3) social disgust processing: 30 papers, 30 experiments, 315 foci, 32 contrasts, 512 subjects. Based on recent recommendations for activation likelihood estimation (ALE) meta-analyses suggesting that at least 20 experiments are sufficient to determine moderate effects with sufficient statistical power (80%) (Eickhoff et al., 2016) this database allowed us to draw robust inferences.

2.3 Activation likelihood estimation

We performed a series of coordinate-based meta-analyses using the current version of GingerALE 3.0.2 (Eickhoff et al., 2012; Eickhoff et al., 2011; Eickhoff et al., 2009) available via http://www.brainmap.org/ale/. All coordinates (either in Talairach or MNI space) and sample size were manually extracted from the original articles. The coordinates in Talairach space were converted to MNI space using the Lancaster transform algorithm, while Talairach space coordinates transformed from MNI space using the Brett transformation were converted back to MNI space using the Brett transformation (Laird et al., 2010; Lancaster et al., 2007). Briefly, the ALE method estimates the likelihood of each voxel being activated under multiple independent experiments with consideration of participants in each experiment. In other words, ALE aims to find the most robust convergent foci between a number of included experiments. To address spatial uncertainty, ALE treats activation foci from single studies as 3D Gaussian probability distributions. Modeled activation (MA) maps for each study were generated by combining the probabilities of all activation foci for each voxel (Eickhoff et al., 2009; Turkeltaub et al., 2012). The union of these generated MA maps was computed while taking into account the disparities between true activation and noise to determine a voxel-wise ALE score (Klugah-Brown et al., 2020). These ALE scores use the convergence of whole-brain coordinates from experiments under a non-random spatial distribution to create clusters formed by equal probabilities of voxels (Eickhoff et al., 2012; Müller et al., 2018). The ALE map reflects clusters corresponding to the location of convergence. And the resulting non-parametric p values account for the proportion of the random spatial relation between the various experiments under the null distribution (Klugah-Brown et al., 2021; Kohn et al., 2014).

For single study analyses (within-group analyses, i.e., general disgust processing, core disgust processing, and social disgust, respectively), the ALE statistics were calculated at each voxel, and cluster-level family-wise error correction (cluster-FWE) (Eickhoff et al., 2016) was applied to the ALE results to control for multiple comparisons. The voxel-level threshold was set at p < 0.001 (uncorrected cluster forming), and the cluster-level threshold was set at p < 0.05 (Eickhoff, Laird, Fox, Lancaster, & Fox, 2017; Eickhoff et al., 2016; Müller et al., 2018), p values in our analyses were generated by 5,000 permutations (Laird et al., 2010; Tao et al., 2021).

Figure 1. Flowchart illustrating the article screening, exclusion and inclusion process according to PRISMA procedure.

2.4 Contrast and conjunction analyses

The present meta-analysis aimed at determining regions robustly involved in disgust processing across domains as well as to differentiate regions that are convergent and divergent between core and social disgust processing in healthy subjects. To this end we employed a series of contrast and conjunction analyses (between study group analyses, comparing the meta-analytic maps for core and social disgust processing) according to the following steps: (1) single study analyses were performed separately for core disgust processing and social disgust processing; (2) the two single studies to be compared were pooled together using the "Merge & Save Foci" function; (3) a single study analysis was conducted on the pooled foci file employing identical statistical criteria as for the preceding single study analyses; (4) via the separate single study analysis three corrected ALE images were obtained for general disgust processing, core

disgust processing, and social disgust processing, respectively. Then, we further determined common and distinct brain regions by conducting contrast and conjunction analyses by subjecting these images into GingerALE toolbox to obtain the similarities as well as differences in the neurobiological basis of core and social disgust processing. Conjunction analyses were performed to determine common regions across core and social disgust processing which were considered as brain systems involved in disgust across processing domains (domain-general). While contrast analyses were implemented to determine brain regions differentially activated for core and social disgust processing (domain-specific) (See **Supplementary Materials** for detailed methodology of conjunction and contrast analyses).

In accordance with previous studies employing contrast and conjunction analyses (Alain, Du, Bernstein, Barten, & Banai, 2018; Papitto, Friederici, & Zaccarella, 2020), the ALE images were thresholded at p < 0.05 (uncorrected), with minimum cluster size of 100 mm^3 and 10,000 permutations, thereby resulting in a larger number of permutations as compared to single study analyses (Alain et al., 2018; Papitto et al., 2020; Wertheim & Ragni, 2018; Yu, Barron, Tantiwongkosi, Fox, & Fox, 2018).

2.5 Meta-analytic connectivity modeling (MACM) analyses

To further functionally characterize the domain-general disgust regions on the network level MACM was employed. Briefly, MACM examines whole-brain functional connectivity patterns by identifying the above-chance covariance between two or more brain regions (Laird et al., 2009; Robinson et al., 2012; Robinson, Laird, Glahn, Lovallo, & Fox, 2010). This approach is similar to the functional connectivity patterns observed in resting-state fMRI (Kotkowski, Price, Mickle Fox, Vanasse, & Fox, 2018; Reid et al., 2017; Smith et al., 2009), yet further complements the resting-state functional connectivity approach as it provides a measure of functional connectivity during a range of task-associated mental states (Langner, Rottschy, Laird, Fox, & Eickhoff, 2014). Coordinates of the peak activation clusters (i.e., right inferior frontal gyrus and left fusiform gyrus) from the conjunction analyses were selected, with the right inferior frontal gyrus located at (44, 26, -8) and the left fusiform gyrus located at (-42, -50, -14). For the fusiform gyrus clusters, we focused on the peak with the higher ALE value. For the MACM analyses, the regions were defined in MNI space using a 10 mm sphere centered at the peak coordinates using the SPM toolbox MarsBaR (http://marsbar.sourceforge.net). A standard MNI brain template (Colin27_T1_seg_MNI.nii; https://www.brainmap.org/ale/Colin27_T1_seg_MNI.nii.gz) was used to visualize both ROI masks. Next region-specific functional connectivity was employed using the functional database of BrainMap's Sleuth 3.0.4 software. To increase the robustness, we employed two convergent approaches (labeled as MACM-A and MACM-B, respectively) which enable us to determine the functional network characterization by determining co-activated regions in previous experiments. See **Supplementary Materials** for a detailed description of conducting these two separate MACM analyses in Sleuth database. The composition of the two datasets for MACM-A is as follows: (1) the ROI in right inferior frontal gyrus (73 experiments, 891

subjects, 719 foci), (2) the ROI in left fusiform gyrus (28 experiments, 414 subjects, 163 foci). And the composition of the two datasets for MACM-B is the following: (1) the ROI in right inferior frontal gyrus (473 experiments, 7,768 subjects, 7,195 foci), (2) the ROI in left fusiform gyrus (366 experiments, 5,439 subjects, 5,801 foci). Finally, these new datasets were analyzed separately using GingerALE 3.0.2 with the same statistical criteria previously applied in the single study analyses (voxel-level of $p < 0.001$ (uncorrected), cluster-FWE of $p < 0.05$, 5,000 permutations).

2.6 Functional characterization of ALE activations on the behavioral level

To facilitate a robust characterization of the identified regions on the behavioral level a meta-analytic decoding approach was employed. To this end, we capitalized on the Brain Annotation Toolbox (BAT, available at https://istbi.fudan.edu.cn/lnen/info/1173/1788.htm) (Liu et al., 2019), which is based on the activation maps in MNI space for the 217 functional terms from the Neurosynth database (http://neurosynth.org/) (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011). The corresponding functional annotation analysis hypothesizes that voxels within a given cluster/region are more likely to be co-activated by the same terms that are functionally connected with the cluster/region, compared with voxels selected randomly, this way BAT enables the functional characterization of a given region based on the meta-analytically associated terms (Liu et al., 2019). In the present study, we employed this approach to characterize the domain-general disgust (core ∩ social; where ∩ represents intersection) as well as core disgust and social disgust regions. $P < 0.05$ was considered statistically significant based on 10,000 permutations. For further details, please turn to the BAT website (https://istbi.fudan.edu.cn/lnen/softwares/BAT_User_Manual_V1.1.docx).

2.7 Report and visualization of results

For the results presented in tabular overview, the following indices were reported: volume of clusters, lateralization of brain regions, labels of brain regions, peak coordinates from voxels, ALE/Z value (Tao et al., 2021). For the conjunction and contrast results, information of percentage of overlap was further reported. With respect to the statistical indices ALE and Z values are provided for single study analyses; while ALE values are provided for conjunction and Z values are provided for contrast analyses (see Papitto et al., (2020) for details regarding the underlying inferential statistic approach). For visualization, the ALE results were displayed using the current version of Mango 4.1 (https://rii.uthscsa.edu/mango/), overlaid onto a standard MNI brain template (Colin27_T1_seg_MNI.nii).

3. RESULTS

3.1 General disgust processing

The ALE meta-analysis encompassing all disgust processing studies revealed twelve significant clusters (**Table 1, Figure 2a**) that were robustly engaged in general disgust processing. The clusters were located in the bilateral insula, amygdala spreading into the right globus pallidus, fusiform gyrus spreading into the declive, middle occipital gyrus, right inferior occipital gyrus, right lingual gyrus, left inferior temporal gyrus, left inferior parietal lobule, right inferior frontal gyrus, and left superior frontal gyrus. See **Supplementary Materials** for a detailed presentation of percentage of overlap with atlas-based brain regions.

3.2 Core disgust processing

Core disgust robustly engaged ten clusters of significant activation, primarily encompassing the bilateral amygdala (spreading into the right globus pallidus) and insula, left inferior frontal gyrus, left declive, left fusiform gyrus, left inferior parietal lobule, left middle occipital gyrus, left inferior temporal gyrus, left superior frontal gyrus, bilateral inferior occipital gyrus spreading into the adjacent lingual gyrus (**Table 2, Figure 2b**). See **Supplementary Materials** for a detailed presentation of percentage of overlap with atlas-based brain regions.

3.3 Social disgust processing

Examining social disgust processing studies revealed six significant clusters of robust activation located in the right insula and adjacent inferior frontal gyrus, bilateral fusiform gyrus spreading into the right declive, left superior frontal gyrus, right cingulate gyrus, right middle occipital gyrus, and right superior temporal gyrus (**Table 3, Figure 2c**). See **Supplementary Materials** for a detailed presentation of percentage of overlap with atlas-based brain regions.

Figure 2. Overview of significant clusters from the single study analyses. (a) General network of disgust processing in healthy subjects. (b) Network of core disgust processing in healthy subjects. (c) Network of social disgust processing in healthy subjects. Coordinates are in the MNI space. Abbreviations: Amy = Amygdala, CG = Cingulate Gyrus, Dec = Declive, FG = Fusiform Gyrus, GP = Globus Pallidus, IFG = Inferior Frontal Gyrus, Ins = Insula, IOG = Inferior Occipital Gyrus, IPL = Inferior Parietal Lobule, ITG = Inferior Temporal Gyrus, LG = Lingual Gyrus, MFG = Middle Frontal Gyrus, MOG = Middle Occipital Gyrus, SFG = Superior Frontal Gyrus, STG = Superior Temporal Gyrus.

3.4 Common and differential brain regions for core and social disgust processing

Examining the common brain systems by means of conjunction analyses revealed that the right inferior frontal gyrus (extending into the anterior insula) and left fusiform gyrus were robustly activated during core and social disgust processing (domain-general disgust processing) (**Table 4, Figure 3a**).

The contrast analyses revealed stronger recruitment of the left inferior parietal lobule, left postcentral gyrus, left middle occipital gyrus, left inferior temporal gyrus, left limbic-striatal cluster (encompassing the amygdala, hippocampus and globus pallidus), right superior frontal gyrus, left medial frontal gyrus, left inferior frontal gyrus, left declive, left fusiform gyrus and left lingual gyrus for core disgust compared with social disgust (domain-specific core disgust processing), suggesting a left hemisphere predominance (**Table 5, Figure 3b**). In contrast social disgust processing was characterized by a stronger engagement of the right inferior frontal gyrus (extending to the insula), left medial frontal gyrus, right cingulate gyrus, right middle occipital gyrus, right declive, right superior temporal gyrus, right middle temporal gyrus, and bilateral fusiform gyrus as compared to core disgust processing (domain-specific social disgust

processing), reflecting a right hemispheric predominance (**Table 5, Figure 3c**).

Figure 3. Overview of significant clusters from the conjunction and contrast analyses. (a) Common activations for core and social disgust processing in healthy subjects (domain-general disgust processing). (b) Greater activations for Core disgust processing (Core > Social) in healthy subjects (domain-specific regions for core disgust). (c) Greater activations for social disgust processing (Social > Core) in healthy subjects (domain-specific regions for social disgust). Coordinates are in the MNI space. Abbreviations: Amy = Amygdala, CG = Cingulate Gyrus, Cla = Claustrum, Dec = Declive, FG = Fusiform Gyrus, GP = Globus Pallidus, IFG = Inferior Frontal Gyrus, Ins = Insula, IOG = Inferior Occipital Gyrus, IPL = Inferior Parietal Lobule, ITG = Inferior Temporal Gyrus, LG = Lingual Gyrus, MFG = Medial Frontal Gyrus, MOG = Middle Occipital Gyrus, MTG = Middle Temporal Gyrus, PostG = Postcentral Gyrus, PreG = Precentral Cyrus, SFG = Superior Frontal Gyrus, STG = Superior Temporal Gyrus.

3.5 Network level functional characterization via MACM analyses

We adopted two different approaches (labeled as MACM A as well as MACM B) aiming to unveil the functional network of right inferior frontal gyrus and left fusiform gyrus from conjunction analyses across core and social disgust processing.

From MACM-A, co-activation maps for the right inferior frontal gyrus included bilateral insula, bilateral inferior frontal gyrus, right anterior cingulate, bilateral medial frontal gyrus and

left cingulate gyrus (**Table 6, Figure 4a**). Co-activation maps for the left fusiform gyrus were significant for bilateral fusiform gyrus, bilateral parahippocampal gyrus, bilateral amygdala, right putamen and right inferior occipital gyrus (**Table 7, Figure 4a**).

The additional functional network analyses (MACM-B) of right inferior frontal gyrus showed converging results as with MACM-A analyses, including bilateral inferior frontal gyrus, bilateral insula, bilateral medial frontal gyrus, left cingulate gyrus, and right anterior cingulate. MACM-B analyses of left fusiform gyrus also showed highly overlapping regions with MACM-A analyses, including bilateral fusiform gyrus, bilateral amygdala, right putamen and right inferior occipital gyrus (**Figure 4b**, see **Supplementary Table 2** and **Supplementary Table 3** for detailed activated regions). In other words, both functional network characterization analyses confirmed each other.

Figure 4. Overview of the functional networks of right inferior frontal gyrus and left fusiform gyrus by MACM analyses. (a) Network of right inferior frontal gyrus and left fusiform gyrus by MACM-A analyses. (b) Network of right inferior frontal gyrus and left fusiform gyrus by MACM-B analyses. Coordinates are in the MNI space.

3.6 Functional characterization of the ALE maps on the behavioral level

We conducted three separate quantitative functional annotation analyses by testing the associations of functional terms provided by the Neurosynth database with the brain regions robustly engaged during domain-general disgust processing generated from conjunction analyses as well as core disgust processing and social disgust processing obtained from single study

analyses. The top 20 terms were visualized with a larger font size reflecting a smaller p-value (see **Figure 5**). Whereas the domain-general network was mainly characterized by general attention and emotional processes (**Figure 5a**), core disgust processing regions primarily associated with threat- and avoidance-related processes (**Figure 5b**), while social disgust processing revealed social- and adaptation-related terms (**Figure 5c**).

Figure 5. Functional characterization of the neural correlates from three categories. (a) Neural substrates for domain-general disgust processing (Core ∩ Social). (b) Neural substrates for core disgust processing. (c) Neural substrates for social disgust processing. Only the top 20 terms are displayed. The font size of words is proportional to the permutation p value for association between user-provided cluster of regions and each functional term provided by Neurosynth, with smaller p value indicating bigger term size.

4. DISCUSSION

　　Previous studies on the brain systems that mediate disgust-related processes and initial studies that examined common and separable neural representations of direct exposure to disgust elicitors (core disgust) and social-communicative signals of disgust (social disgust) revealed inconsistent results. Against this background we employed neuroimaging meta-analyses that capitalized on a large number of previous fMRI studies to determine the neural substrates robustly engaged in general, core as well as social disgust processing, respectively. Next, by employing meta-analytic conjunction and contrast analyses we further segregated common (domain-general) and distinct (domain-specific) neural correlates of core and social disgust processing and utilized meta-analytic behavioral and network level decoding to functionally characterize the identified brain regions.

4.1. Brain systems mediating general, core and social disgust processing

　　The overarching meta-analysis pooling the data from the 69 qualified studies, encompassing 78 eligible contrasts (general disgust processing) yielded robust engagement of a bilateral network encompassing primarily the bilateral anterior insular cortex, bilateral amygdala and adjacent globus pallidus, inferior and superior frontal, visual occipital (fusiform gyrus, inferior and middle occipital gyri, lingual gyrus) and cerebellar regions. Separate analyses of the subdomains of core and social disgust processing revealed that the core disgust network largely overlapped with the general disgust network while the social disgust processing network additionally engaged superior temporal regions involved in social processing and face processing regions such as fusiform-extrastriate and frontal cortices, mainly in the right hemisphere.

　　The identified network for general disgust processing largely resembles regions that have been consistently engaged in disgust-related processes (Fusar-Poli et al., 2009; Kirby & Robinson, 2017; Vytal & Hamann, 2010; Wager et al., 2015) and may reflect the multifaceted nature of the disgust response. The identified regions may support the engagement of different simultaneous processes which allow a concerted disgust reaction, ranging from initial detection of biologically salient stimuli in occipital visual regions (Tao et al., 2021), early threat detection and implementation of the defensive motor response by the amygdala, globus pallidus, fusiform and cerebellar regions (e.g., Phelps & LeDoux, 2005; Surguladze et al., 2003; Yao, Zhao, et al., 2018), as well as salience allocation and interoceptive awareness of the emotional state in the insula cortex (Craig, 2009; Li et al., 2019; Uddin, 2015), and the subjective experience and interpretation of the emotional state in prefrontal regions (LeDoux & Pine, 2016).

　　Examining the core disgust network, i.e., regions robustly activated by disgust eliciting visual stimuli, revealed a similar network primarily encompassing the bilateral insula and adjacent inferior frontal regions, the bilateral amygdala and adjacent globus pallidus as well as temporo-occipital regions. The overlap may reflect that the direct exposure to potential infectious and harmful stimuli triggers the evolutionary defensive-avoidance circuits engaged in early salience and threat detection and the initiation of the physiological and behavioral defensive response thus facilitating rapid withdrawal. Previous studies have for instance extensively documented reactivity of the amygdala towards aversive stimuli (Zald, 2003) with meta-analytic data demonstrating a higher probability of amygdala activation for stimuli signaling a potential danger, including fear and disgust inducing stimuli, as compared to non-threatening stimuli (Costafreda, Brammer, David, & Fu, 2008), indicating a critical role in threat detection and initiation of

protective responses (Mihov et al., 2013; Schienle, Übel, & Wabnegger, 2016). The amygdala receives dense projections from the visual cortex and via projections with cerebellar regions and the globus pallidus initiates motor and physiological defensive responses towards aversive stimuli (Giovanniello et al., 2020; Lindquist, 2020; Salih et al., 2009; Sambataro et al., 2006), while bilateral projections to the orbitofrontal cortex further mediate withdrawal or avoidance behavior (Rolls, 1999, 2000b). In support of this interpretation the meta-analytic functional characterization of the identified core disgust network revealed primarily terms associated with negative affective responses including e.g., anxiety, arousal as well as disgust.

Separately examining the social disgust studies revealed a right lateralized network encompassing the right anterior insula and inferior frontal gyrus, middle occipital cortex, fusiform gyrus, superior temporal gyrus and cingulate regions. The corresponding meta-analytic functional decoding of the regions revealed primarily terms associated with salience and social communication including attention as well as face and communication. In line with the functional decoding the identified regions have been previously associated with emotional face processing (Fusar-Poli et al., 2009), with the middle occipital cortex involved in early face processing stages (Tu, Qiu, Martens, & Zhang, 2013) and the fusiform gyrus being involved in visual attention and higher level face perception processing (Weiner & Zilles, 2016) while superior temporal regions have been strongly involved in social cognitive processes including theory of mind and mentalizing (Frith, 2007; Zilbovicius et al., 2006).

In contrast to several previous original studies, the present meta-analysis did not reveal a robust engagement of the amygdala or main basal ganglia regions (i.e., caudate and putamen) during social or general disgust processing respectively. However, despite a number of previous studies reporting amygdala engagement during disgusted faces a previous meta-analysis failed to corroborate amygdala engagement during disgust faces processing (Fusar-Poli et al., 2009) and previous studies found either strongest amygdala responses to fear (Mattavelli et al., 2014) or comparable responses to emotional and neutral faces (Derntl et al., 2009; Fitzgerald et al., 2006; Winston et al., 2003) while amygdala lesion patients exhibited intact disgusted face decoding (Becker et al., 2012). With respect to the role of the basal ganglia in disgust previous imaging (Ahn et al., 2014; Benuzzi, Lui, Duzzi, Nichelli, & Porro, 2008; Lassalle et al., 2019; Phillips et al., 2000; Phillips et al., 2004; Phillips et al., 1998; Phillips et al., 1997; Pujol et al., 2018; Shapira et al., 2003; Sprengelmeyer et al., 1998; Tettamanti et al., 2012; Ziegler et al., 2018) and lesion studies (Calder et al., 2000; Sprengelmeyer et al., 1996; Straube et al., 2010) revealed inconsistent results and the present meta-analysis did not confirm a robust engagement.

4.2 Common and distinct neural correlates of core and social disgust processing

Notably the right anterior insula was consistently and specifically identified across both, the separate analyses for core disgust and social disgust. Although the consistent involvement of the insula in disgust processing has long been debated in the original disgust literature (Deeley et al., 2008; Gorno-Tempini et al., 2001; Schienle et al., 2006; Schienle et al., 2005b; Stark et al., 2005; Stark et al., 2004; Stark et al., 2003; Trautmann et al., 2009), several previous studies reported engagement of this region during core disgust as well as social disgust processing (Chapman & Anderson, 2012; Fusar-Poli et al., 2009; Vytal & Hamann, 2010). Further support for a role of the insula in disgust processing comes from lesion (Adolphs, Tranel, & Damasio, 2003; Calder et al., 2000; Kipps, Duggins, McCusker, & Calder, 2007) and intracerebral recording studies (Krolak-

Salmon et al., 2003), and studies in healthy subjects reporting a convergent engagement of the insula across disgust paradigms employing not only visual but also olfactory (Heining et al., 2004; Wicker et al., 2003), gustatory (Corradi-Dell'Acqua et al., 2016; Jabbi et al., 2008) and imagination (Jabbi et al., 2008) procedures. The insula is a functionally heterogenous region and has been involved in general salience and interoceptive processes but also in the awareness and experience of emotional states (e.g., Craig, 2009; Uddin, 2015; Zhou, Li, et al., 2020). Previous studies have demonstrated that negative affective experience induced by visual stimuli (Viinikainen et al., 2010), subjective experienced unpleasantness (Benuzzi et al., 2008), as well as aversive electrical stimulations and unfair offers in social interactions (Corradi-Dell'Acqua et al., 2016) consistently engage the anterior insula confirming its potential role in the representation of negative emotional experiences.

Using a conjunction and contrast meta-analytic approach we aimed to further segregate common and domain-specific neural correlates for core and social disgust processing. In line with the observations from the separate analyses, the right anterior insula extending into the adjacent inferior frontal gyrus and the left fusiform gyrus specifically exhibited convergent engagement during core and social disgust processing. Together with the anterior insula the inferior frontal cortex plays an important role in general emotional processes such as evaluation of biological significant stimuli (Rolls, 2000a, 2004), interoception (Critchley, Wiens, Rotshtein, Öhman, & Dolan, 2004), multimodal representation emotion salient stimuli (Yamasaki, LaBar, & McCarthy, 2002), and emotion recognition (Sprengelmeyer et al., 1998) while the fusiform gyrus plays a role in higher order visual processing (Fusar-Poli et al., 2009) and emotional reactivity towards negative and threatening stimuli (Kirby & Robinson, 2017; Tao et al., 2021). Further meta-analytic network decoding via MACM demonstrated that the identified anterior insula cortex region coupled with a bilateral network encompassing the insula and ventrolateral prefrontal regions as well as anterior cingulate and medial frontal regions. The lateral prefrontal cortex, insula and anterior cingulate exhibit strong anatomical and functional connections (Craig, 2009; Mesulam & Mufson, 1982) and represent key hubs within the salience network, the fronto-parietal executive control network and the sensorimotor networks (Belyk, Brown, Lim, & Kotz, 2017; Cloutman, Binney, Drakesmith, Parker, & Lambon Ralph, 2012; Cox et al., 2010). Functional interactions between these regions have been involved in attentional processes (Wang et al., 2019), emotion regulation (Morawetz, Bode, Baudewig, & Heekeren, 2017) and goal-directed behavior (Dosenbach et al., 2006) which may support disgust processing across core and social domains. The identified left fusiform gyrus network comprised a limbic-occipital network encompassing the amygdala, parahippocampal gyrus, putamen and inferior occipital regions. The functional interplay between these regions has been demonstrated in several previous studies (Amting, Greening, & Mitchell, 2010; Fairhall & Ishai, 2007; Herrington, Taylor, Grupe, Curby, & Schultz, 2011; Molapour, Golkar, Navarrete, Haaker, & Olsson, 2015; Turk-Browne, Norman-Haignere, & McCarthy, 2010) and plays an important role in early threat detection (Tao et al., 2021), suggesting that this network may facilitate detection of potential threats across core and social disgust processes.

Examining regions specifically engaged during the two disgust processing domains revealed stronger activity of several regions of the left hemispheric disgust network during core disgust processing including e.g., occipital visual, limbic, inferior frontal and parietal regions. These regions mainly mirror the results of the core disgust processing network, suggesting that

exposure to potentially dangerous stimuli engages core nodes of the defensive-avoidance circuits, including the amygdala engaged in threat detection (Whalen, 2007), interoceptive and emotional awareness (Critchley et al., 2004) as well as regions engaged in attentional and contextual processes such as inferior parietal and parahippocampal regions (Behrmann, Geng, & Shomstein, 2004; Caspers et al., 2012; Tao et al., 2021). In contrast, for the social disgust processing a right lateralized network encompassing inferior frontal gyrus, superior temporal gyrus as well as anterior cingulate regions was stronger activity. These regions have been strongly engaged in face processing, such that a previous study demonstrated a stronger engagement during emotional faces as compared to emotional scenes in this network (Britton, Taylor, Sudheimer, & Liberzon, 2006), as well as social cognitive processes such as mentalizing and theory of mind, reflecting that the interpretation of social signals of disgust may require more complex social cognitive decoding of the stimuli.

Notably, a strong lateralization effect was observed such that core disgust processing primarily engaged a left-lateralized, while social disgust processing primarily engaged a right-lateralized network. Although we did not have a priori lateralization hypotheses the findings resonate with previous studies suggesting a left lateralization of withdrawal-related activity (Wager, Phan, Liberzon, & Taylor, 2003) while conscious emotional perception is represented in a right lateralized network (Adolphs, Damasio, Tranel, & Damasio, 1996; Davidson & Irwin, 1999; Oliveri et al., 2003).

4.3 Limitations

Results of the present study need to be considered in the context of several limitations. First, core disgust in the present study included contamination and envelope violations (Vicario et al., 2017) while some previous disgust conceptualizations differentiate these stimuli (Haidt et al., 1994; Olatunji & Sawchuk, 2005; Olatunji et al., 2007; Rozin & Fallon, 1987; van Overveld, de Jong, & Peters, 2009). Initial neuroimaging studies revealed inconsistent support of this notion (Borg, de Jong, Renken, & Georgiadis, 2012; Harrison, Gray, Gianaros, & Critchley, 2010; Schienle et al., 2006; Wright, He, Shapira, Goodman, & Liu, 2004) and an increasing number of neuroimaging studies on disgust may allow a reasonable powered examination of further disgust subprocesses in the future. Second, the present database of original studies did not allow to examine sex-differences in the neural basis of disgust processing. Initial studies reported e.g., stronger disgust-related inferior frontal reactivity in women (Caseras et al., 2007), while other studies did not find sex differences (Baumann & Mattingley, 2012; Schienle, Schäfer, Stark, Walter, & Vaitl, 2005a; Stark et al., 2003). Third, the present study included task-based fMRI studies focusing on regional analyses, while recent advanced approaches and findings suggest a distributed neural representation (Zhou, Zhao, et al., 2020) or a functional division between gyri and sulci (Jiang et al., 2015; Jiang et al., 2018; Jiang, Zhang, Zhang, Kendrick, & Liu, 2021) of cognitive and emotional processes. Future studies may capitalize on these developments to further dissect common and differential neural representation of social and non-social processes or different defensive responses such as disgust and fear.

4.4 Conclusions

The present study employed a coordinate-based neuroimaging meta-analytic approach to

determine the neurobiological basis of disgust processing in humans. Exposure to disgust eliciting stimuli induced robust activity in a network encompassing regions engaged in early visual threat detection, emotional experience and defensive motor responses, while social stimuli of disgust primarily engaged regions engaged in facial and social cognitive processes. Across both disgust domains the anterior insula/inferior frontal gyrus and the fusiform gyrus were robustly engaged suggesting an involvement of these regions across disgust processing modalities.

Funding

This work was supported by the National Key Research and Development Program of China (2018YFA0701400, and the National Natural Science Foundation of China (NSFC, 61871420).

Acknowledgements

We would like to thank Zonglin He and Qian Tao from Jinan University (China) for their kind help with resolving some data analyzing issues.

Table 1

Brain regions generally activated by all studies of disgust processing in healthy subjects.

Volume (mm³)	Side	Brain region	BA	MNI coordinates (x, y, z)			ALE value (10⁻²)	Z
2568	right	Amygdala		22	-6	-16	3.34	5.30
	right	Amygdala		24	-2	-26	3.26	5.21
2560	right	Inferior Frontal Gyrus	47	42	26	-6	3.22	5.16
	right	Middle Frontal Gyrus	46	48	18	14	2.28	3.97
	right	Insula	13	44	20	4	1.89	3.42
2272	left	Fusiform Gyrus	37	-42	-48	-16	2.97	4.86
	left	Fusiform Gyrus	37	-44	-58	-8	2.93	4.81
2176	right	Fusiform Gyrus	19	42	-72	-12	2.48	4.24
	right	Fusiform Gyrus	37	42	-52	-18	2.41	4.14
	right	Fusiform Gyrus	37	42	-46	-20	2.24	3.92
	right	Declive		38	-58	-14	2.23	3.89
	right	Inferior Occipital Gyrus	19	48	-72	-2	1.98	3.55
2104	left	Amygdala		-22	-2	-22	4.48	6.57
2024	right	Middle Occipital Gyrus	18	36	-88	6	3.18	5.12
	right	Inferior Occipital Gyrus	18	36	-88	-10	2.82	4.68
1400	right	Lingual Gyrus	17	14	-92	0	3.89	5.93
1304	left	Middle Occipital Gyrus	19	-44	-78	8	2.77	4.61
	left	Inferior Temporal Gyrus	19	-50	-76	0	2.01	3.60
1256	left	Insula		-36	26	-4	2.29	3.99
	left	Insula	13	-34	24	0	2.28	3.97
	left	Insula	13	-44	14	-2	2.12	3.75
1224	left	Inferior Parietal Lobule	40	-62	-24	34	4.07	6.14
960	right	Inferior Frontal Gyrus	9	48	10	24	3.27	5.22

888	left	Superior Frontal Gyrus	9	-6	64	26	2.76	4.60
	left	Superior Frontal Gyrus	9	2	62	18	2.14	3.78

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 2

Brain regions activated during core disgust processing in healthy subjects.

Volume (mm³)	Side	Brain region	BA	MNI coordinates (x, y, z)			ALE value (10⁻²)	Z
2216	left	Amygdala		-22	-2	-22	3.80	6.49
1736	left	Declive		-34	-70	-14	2.53	4.89
	left	Fusiform Gyrus	37	-44	-60	-8	2.09	4.27
	left	Fusiform Gyrus	37	-42	-54	-18	1.69	3.67
1696	right	Amygdala		22	-6	-16	2.85	5.31
	right	Amygdala		26	-2	-24	2.33	4.62
1512	left	Inferior Parietal Lobule	40	-62	-24	34	3.99	6.71
1472	left	Middle Occipital Gyrus	19	-44	-76	8	2.47	4.81
	left	Inferior Temporal Gyrus	19	-50	-76	0	2.01	4.16
1240	left	Superior Frontal Gyrus	9	-6	64	26	2.21	4.44
	left	Superior Frontal Gyrus	9	2	62	18	2.14	4.34
1120	right	Inferior Occipital Gyrus	17	14	-92	-2	2.64	5.04
976	left	Inferior Frontal Gyrus	47	-30	32	-14	2.68	5.09
	left	Insula	13	-34	24	-2	1.73	3.72
896	right	Insula		40	26	-6	2.16	4.37
808	left	Inferior Occipital Gyrus	17	-10	-98	-8	2.14	4.34

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 3

Brain regions activated during social disgust processing in healthy subjects.

Volume (mm³)	Side	Brain region	BA	MNI coordinates (x, y, z)			ALE value (10⁻²)	Z
1848	right	insula	13	48	18	12	2.11	4.66
	right	insula	13	44	20	4	1.81	4.19
	right	Inferior Frontal Gyrus	47	46	28	-8	1.50	3.67
1416	left	Superior Frontal Gyrus	6	2	10	54	1.92	4.37
	left	Superior Frontal Gyrus	6	-2	12	56	1.90	4.33
	right	Cingulate Gyrus	24	6	10	42	1.40	3.51
1392	right	Middle Occipital Gyrus	18	36	-88	8	2.93	5.82
1256	right	Fusiform Gyrus	19	42	-72	-14	2.42	5.12
888	left	Fusiform Gyrus	37	-40	-46	-18	2.31	4.97
728	right	Superior Temporal Gyrus	22	52	-34	4	2.31	4.96

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 4

Brain regions activated by conjunction analysis between core and social disgust processing in healthy subjects.

Volume (mm^3)	Side	Brain region	Percentage of overlap	BA	MNI coordinates (x, y, z)			ALE value (10^{-2})
136	right	Inferior Frontal Gyrus	80% Inferior Frontal Gyrus, 20% Insula.	47	44	26	-8	1.49
16	left	Fusiform Gyrus	100% Fusiform Gyrus.	37	-40	-52	-16	1.21
16	left	Fusiform Gyrus	100% Fusiform Gyrus.	37	-42	-50	-14	1.37

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 5

Brain regions activated by contrast analysis between core and social disgust processing in healthy subjects.

Volume (mm³)	Side	Brain region	Percentage of overlap	BA	MNI coordinates (x, y, z)			Z
Hyper-activation (core > social)								
1256	left	Postcentral Gyrus	67.9% Inferior Parietal Lobule, 32.1% Postcentral Gyrus.	2	-66	-20	34	2.64
	left	Postcentral Gyrus		2	-61.3	-21.3	36.7	2.56
	left	Inferior Parietal Lobule		40	-66	-24	32	2.53
	left	Inferior Parietal Lobule		40	-64.7	-27.3	35	2.47
	left	Inferior Parietal Lobule		40	-64	-28	28	2.34
992	left	Inferior Temporal Gyrus	47.8% Middle Occipital Gyrus, 34.8% Inferior Temporal Gyrus, 8.7% Inferior Occipital Gyrus, 8.7% Middle Temporal Gyrus.	37	-50	-74	0	3.29
	left	Middle Occipital Gyrus		37	-44	-72	6	2.58
808	left	Parahippocampal Gyrus	49.1% Brodmann area 34, 45.5% Amygdala, 3.6% Globus Pallidus, 1.8% Brodmann area 28.	34	-16	-2	-22	2.14
	left	Parahippocampal Gyrus		34	-18	0	-14	1.90
688	right	Superior Frontal Gyrus	73.7% Superior Frontal Gyrus, 26.3% Medial	9	4	60	20	2.45
	right	Superior Frontal Gyrus		9	4	64	16	2.34

				BA	x	y	z	Z
	left	Medial Frontal Gyrus	Frontal Gyrus.	10	0	64	14	2.33
360	left	Inferior Frontal Gyrus	100% Inferior Frontal Gyrus.	47	-28	34	-10	2.23
	left	Claustrum			-28	30	-8	2.13
240	left	Fusiform Gyrus	69.2% Declive, 30.8% Fusiform Gyrus.	37	-40	-66	-12	2.17
	left	Declive			-32	-70	-10	1.87
224	left	Lingual Gyrus	92.9% Lingual Gyrus, 7.1% Inferior Occipital Gyrus.	17	-8	-100	-6	1.89

Hyper-activation (social > core)

				BA	x	y	z	Z
1344	right	Inferior Frontal Gyrus	70.6% Insula, 20.6% Inferior Frontal Gyrus, 8.8% Precentral Gyrus.	44	49	18	8	3.54
	right	Inferior Frontal Gyrus		45	46	22	8	3.35
1128	left	Medial Frontal Gyrus	57% Medial Frontal Gyrus, 31.6% Superior Frontal Gyrus, 11.4% Cingulate Gyrus.	32	-2	14	48	3.54
	right	Cingulate Gyrus		24	5	10	44	3.09
1040	right	Middle Occipital Gyrus	100% Middle Occipital Gyrus.		30	-90	2	3.09
	right	Middle Occipital Gyrus		19	38	-86	10	3.06
1032	right	Declive	81.3% Declive, 18.7% Fusiform Gyrus.		44	-76	-16	3.16
	right	Declive			42	-70	-18	3.04
728	right	Superior Temporal Gyrus	55.6% Middle Temporal Gyrus, 44.4% Superior	22	54.8	-33.6	3.2	3.29
	right	Middle Temporal Gyrus			52	-35	0	3.24

			Temporal Gyrus.					
472	left	Fusiform Gyrus	78.9% Fusiform Gyrus, 21.1% Culmen.	37	-36	-46	-16	2.49

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 6

Network of right inferior frontal gyrus revealed by MACM-A analysis.

Volume (mm³)	Side	Brain region	BA	MNI coordinates (x, y, z)			ALE value (10⁻²)	Z
5352	right	Inferior Frontal Gyrus	47	40	26	-8	3.54	5.85
	right	Insula	13	46	10	-8	2.47	4.46
	right	Inferior Frontal Gyrus	46	38	36	6	2.42	4.39
	right	Inferior Frontal Gyrus	13	38	28	2	2.14	3.99
4808	left	Insula	13	-34	24	-2	3.88	6.25
	left	Inferior Frontal Gyrus	46	-48	38	4	1.98	3.75
1536	right	Anterior Cingulate	32	6	50	-4	2.43	4.41
	right	Medial Frontal Gyrus	11	2	38	-22	2.13	3.97
	left	Medial Frontal Gyrus	11	0	42	-26	2.08	3.90
	right	Anterior Cingulate	32	4	46	-16	1.94	3.69
1192	left	Cingulate Gyrus	32	-8	26	42	3.24	5.48
1104	left	Inferior Frontal Gyrus	9	-46	12	26	2.27	4.18

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

Table 7

Network of left fusiform gyrus revealed by MACM-A analysis.

Volume (mm^3)	Side	Brain region	BA	MNI coordinates (x, y, z)			ALE value (10^{-2})	Z
2776	left	Fusiform Gyrus	37	-44	-50	-14	2.04	5.18
	left	Parahippocampal Gyrus	19	-40	-48	-6	1.58	4.39
2240	right	Amygdala		30	-4	-20	1.88	4.90
	right	Parahippocampal Gyrus	28	20	-16	-24	1.18	3.59
	right	Putamen		34	4	-10	1.16	3.55
1872	left	Amygdala		-26	-2	-22	2.27	5.56
1184	right	Inferior Occipital Gyrus	19	48	-74	-2	2.21	5.46
	right	Fusiform Gyrus	19	40	-72	-14	1.43	4.11

Abbreviations: ALE = Action Likelihood Estimation, BA = Brodmann Area, MNI = Montreal Neurological Institute.

References

Adolphs, R., Damasio, H., Tranel, D., & Damasio, A. R. (1996). Cortical Systems for the Recognition of Emotion in Facial Expressions. *The Journal of Neuroscience, 16*(23), 7678-7687. doi:10.1523/JNEUROSCI.16-23-07678.1996

Adolphs, R., Tranel, D., & Damasio, A. R. (2003). Dissociable neural systems for recognizing emotions. *Brain and Cognition, 52*(1), 61-69. doi:10.1016/S0278-2626(03)00009-5

Ahn, W.-Y., Kishida, K. T., Gu, X., Lohrenz, T., Harvey, A., Alford, J. R., . . . Montague, P. R. (2014). Nonpolitical Images Evoke Neural Predictors of Political Ideology. *Current Biology, 24*(22), 2693-2699. doi:10.1016/j.cub.2014.09.050

Alain, C., Du, Y., Bernstein, L. J., Barten, T., & Banai, K. (2018). Listening under difficult conditions: An activation likelihood estimation meta-analysis. *Human Brain Mapping, 39*(7), 2695-2709. doi:10.1002/hbm.24031

Amting, J. M., Greening, S. G., & Mitchell, D. G. V. (2010). Multiple Mechanisms of Consciousness: The Neural Correlates of Emotional Awareness. *The Journal of Neuroscience, 30*(30), 10039. doi:10.1523/JNEUROSCI.6434-09.2010

Ashworth, F., Pringle, A., Norbury, R., Harmer, C. J., Cowen, P. J., & Cooper, M. J. (2011). Neural response to angry and disgusted facial expressions in bulimia nervosa. *Psychological medicine, 41*(11), 2375-2384. doi:10.1017/S0033291711000626

Baumann, O., & Mattingley, J. B. (2012). Functional topography of primary emotion processing in the human cerebellum. *NeuroImage, 61*(4), 805-811. doi:10.1016/j.neuroimage.2012.03.044

Becker, B., Mihov, Y., Scheele, D., Kendrick, K. M., Feinstein, J. S., Matusch, A., . . . Hurlemann, R. (2012). Fear Processing and Social Networking in the Absence of a Functional Amygdala. *Biological Psychiatry, 72*(1), 70-77. doi:10.1016/j.biopsych.2011.11.024

Behrmann, M., Geng, J. J., & Shomstein, S. (2004). Parietal cortex and attention. *Current Opinion in Neurobiology, 14*(2), 212-217. doi:10.1016/j.conb.2004.03.012

Belyk, M., Brown, S., Lim, J., & Kotz, S. A. (2017). Convergence of semantics and emotional expression within the IFG pars orbitalis. *NeuroImage, 156*, 240-248. doi:10.1016/j.neuroimage.2017.04.020

Benuzzi, F., Lui, F., Duzzi, D., Nichelli, P., & Porro, C. (2008). Does It Look Painful or Disgusting? Ask Your Parietal and Cingulate Cortex. *The Journal of Neuroscience, 28*(4), 923-931. doi:10.1523/JNEUROSCI.4012-07.2008

Berlin, H. A., Stern, E. R., Ng, J., Zhang, S., Rosenthal, D., Turetzky, R., . . . Goodman, W. (2017). Altered olfactory processing and increased insula activity in patients with obsessive-compulsive disorder: An fMRI study. *Psychiatry Research: Neuroimaging, 262*, 15-24. doi:10.1016/j.pscychresns.2017.01.012

Borg, C., de Jong, P. J., Renken, R. J., & Georgiadis, J. R. (2012). Disgust trait modulates frontal-posterior coupling as a function of disgust domain. *Social Cognitive and Affective Neuroscience, 8*(3), 351-358. doi:10.1093/scan/nss006

Britton, J. C., Taylor, S. F., Sudheimer, K. D., & Liberzon, I. (2006). Facial expressions and complex IAPS pictures: Common and differential networks. *NeuroImage, 31*(2), 906-919. doi:10.1016/j.neuroimage.2005.12.050

Burklund, L., Eisenberger, N., & Lieberman, M. (2007). The Face of Rejection: Rejection Sensitivity Moderates Dorsal Anterior Cingulate Activity to Disapproving Facial Expressions. *Social Neuroscience, 2*, 238-253. doi:10.1080/17470910701391711

Calder, A. J., Keane, J., Manes, F., Antoun, N., & Young, A. W. (2000). Impaired recognition and experience of disgust following brain injury. *Nature Neuroscience, 3*(11), 1077-1078. doi:10.1038/80586

Calder, A. J., Lawrence, A. D., & Young, A. W. (2001). Neuropsychology of fear and loathing. *Nature Reviews Neuroscience, 2*(5), 352-363. doi:10.1038/35072584

Cantone, M., Lanza, G., Bella, R., Pennisi, G., Santalucia, P., Bramanti, P., & Pennisi, M. (2019). Fear and disgust: case report of two uncommon emotional disturbances evoked by visual disperceptions after a right temporal-insular stroke. *BMC Neurology, 19*(1), 193. doi:10.1186/s12883-019-1417-0

Caseras, X., Mataix-Cols, D., An, S. K., Lawrence, N. S., Speckens, A., Giampietro, V., . . . Phillips, M. L. (2007). Sex Differences in Neural Responses to Disgusting Visual Stimuli: Implications for Disgust-Related Psychiatric Disorders. *Biological Psychiatry, 62*(5), 464-471. doi:10.1016/j.biopsych.2006.10.030

Caspers, S., Schleicher, A., Bacha-Trams, M., Palomero-Gallagher, N., Amunts, K., & Zilles, K. (2012). Organization of the Human Inferior Parietal Lobule Based on Receptor Architectonics. *Cerebral Cortex, 23*(3), 615-628. doi:10.1093/cercor/bhs048 %J Cerebral Cortex

Chapman, H. A., & Anderson, A. K. (2012). Understanding disgust. *Annals of the New York Academy of Sciences, 1251*(1), 62-76. doi:10.1111/j.1749-6632.2011.06369.x

Chapman, H. A., Kim, D. A., Susskind, J. M., & Anderson, A. K. (2009). In Bad Taste: Evidence for the Oral Origins of Moral Disgust. *Science, 323*(5918), 1222. doi:10.1126/science.1165565

Chen, T., Becker, B., Camilleri, J., Wang, L., Yu, S., Eickhoff, S. B., & Feng, C. (2018). A domain-general brain network underlying emotional and cognitive interference processing: evidence from coordinate-based and functional connectivity meta-analyses. *Brain Structure and Function, 223*(8), 3813-3840. doi:10.1007/s00429-018-1727-9

Cloutman, L. L., Binney, R. J., Drakesmith, M., Parker, G. J. M., & Lambon Ralph, M. A. (2012). The variation of function across the human insula mirrors its patterns of structural connectivity: Evidence from in vivo probabilistic tractography. *NeuroImage, 59*(4), 3514-3521. doi:10.1016/j.neuroimage.2011.11.016

Corradi-Dell'Acqua, C., Tusche, A., Vuilleumier, P., & Singer, T. (2016). Cross-modal representations of first-hand and vicarious pain, disgust and fairness in insular and cingulate cortex. *Nature Communications, 7*(1), 10904. doi:10.1038/ncomms10904

Costafreda, S. G., Brammer, M. J., David, A. S., & Fu, C. H. Y. (2008). Predictors of amygdala activation during the processing of emotional stimuli: A meta-analysis of 385 PET and fMRI studies. *Brain Research Reviews, 58*(1), 57-70. doi:10.1016/j.brainresrev.2007.10.012

Cox, C. L., Gotimer, K., Roy, A. K., Castellanos, F. X., Milham, M. P., & Kelly, C. (2010). Your Resting Brain CAREs about Your Risky Behavior. *PLOS ONE, 5*(8), e12296. doi:10.1371/journal.pone.0012296

Craig, A. D. (2009). How do you feel — now? The anterior insula and human awareness. *Nature Reviews Neuroscience, 10*(1), 59-70. doi:10.1038/nrn2555

Critchley, H. D., Wiens, S., Rotshtein, P., Öhman, A., & Dolan, R. J. (2004). Neural systems supporting interoceptive awareness. *Nature Neuroscience, 7*(2), 189-195. doi:10.1038/nn1176

Curtis, V., de Barra, M., & Aunger, R. (2011). Disgust as an adaptive system for disease avoidance behaviour. *Philosophical Transactions of the Royal Society B: Biological Sciences, 366*(1563), 389-401. doi:10.1098/rstb.2010.0117

Darwin, C. (1965). *The Expression of Emotion in Man and Animals*: University of Chicago Press: Chicago,

IL.

Davey, G. C. L. (2011). Disgust: the disease-avoidance emotion and its dysfunctions. *Philosophical Transactions of the Royal Society B: Biological Sciences, 366*(1583), 3453-3465. doi:10.1098/rstb.2011.0039

Davidson, R. J., & Irwin, W. (1999). The functional neuroanatomy of emotion and affective style. *Trends in Cognitive Sciences, 3*(1), 11-21. doi:10.1016/s1364-6613(98)01265-0

Deeley, Q., Daly, E. M., Azuma, R., Surguladze, S., Giampietro, V., Brammer, M. J., . . . Murphy, D. G. M. (2008). Changes in male brain responses to emotional faces from adolescence to middle age. *NeuroImage, 40*(1), 389-397. doi:10.1016/j.neuroimage.2007.11.023

Deeley, Q., Daly, E. M., Surguladze, S., Page, L., Toal, F., Robertson, D., . . . Murphy, D. G. M. (2007). An Event Related Functional Magnetic Resonance Imaging Study of Facial Emotion Processing in Asperger Syndrome. *Biological Psychiatry, 62*(3), 207-217. doi:10.1016/j.biopsych.2006.09.037

Derntl, B., Habel, U., Windischberger, C., Robinson, S., Kryspin-Exner, I., Gur, R. C., & Moser, E. (2009). General and specific responsiveness of the amygdala during explicit emotion recognition in females and males. *BMC Neuroscience, 10*(1), 91. doi:10.1186/1471-2202-10-91

Dosenbach, N. U. F., Visscher, K. M., Palmer, E. D., Miezin, F. M., Wenger, K. K., Kang, H. C., . . . Petersen, S. E. (2006). A Core System for the Implementation of Task Sets. *Neuron, 50*(5), 799-812. doi:10.1016/j.neuron.2006.04.031

Eickhoff, S. B., Bzdok, D., Laird, A. R., Kurth, F., & Fox, P. T. (2012). Activation likelihood estimation meta-analysis revisited. *NeuroImage, 59*(3), 2349-2361. doi:10.1016/j.neuroimage.2011.09.017

Eickhoff, S. B., Bzdok, D., Laird, A. R., Roski, C., Caspers, S., Zilles, K., & Fox, P. T. (2011). Co-activation patterns distinguish cortical modules, their connectivity and functional differentiation. *NeuroImage, 57*(3), 938-949. doi:10.1016/j.neuroimage.2011.05.021

Eickhoff, S. B., Laird, A. R., Fox, P. M., Lancaster, J. L., & Fox, P. T. (2017). Implementation errors in the GingerALE Software: Description and recommendations. *Human Brain Mapping, 38*(1), 7-11. doi:10.1002/hbm.23342

Eickhoff, S. B., Laird, A. R., Grefkes, C., Wang, L. E., Zilles, K., & Fox, P. T. (2009). Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical estimates of spatial uncertainty. *Human Brain Mapping, 30*(9), 2907-2926. doi:10.1002/hbm.20718

Eickhoff, S. B., Nichols, T. E., Laird, A. R., Hoffstaedter, F., Amunts, K., Fox, P. T., . . . Eickhoff, C. R. (2016). Behavior, sensitivity, and power of activation likelihood estimation characterized by massive empirical simulation. *NeuroImage, 137*, 70-85. doi:10.1016/j.neuroimage.2016.04.072

Ekman, P. (1993). Facial expression and emotion. *American Psychologist, 48*(4), 384-392. doi:10.1037//0003-066x.48.4.384

Fairhall, S. L., & Ishai, A. (2007). Effective Connectivity within the Distributed Cortical Network for Face Perception. *Cerebral Cortex, 17*(10), 2400-2406. doi:10.1093/cercor/bhl148

Feng, C., Becker, B., Huang, W., Wu, X., Eickhoff, S. B., & Chen, T. (2018). Neural substrates of the emotion-word and emotional counting Stroop tasks in healthy and clinical populations: A meta-analysis of functional brain imaging studies. *NeuroImage, 173*, 258-274. doi:10.1016/j.neuroimage.2018.02.023

Fitzgerald, D. A., Angstadt, M., Jelsone, L. M., Nathan, P. J., & Phan, K. L. (2006). Beyond threat: Amygdala reactivity across multiple expressions of facial affect. *NeuroImage, 30*(4), 1441-1448. doi:10.1016/j.neuroimage.2005.11.003

Frith, C. D. (2007). The social brain? *Philosophical Transactions of the Royal Society B: Biological Sciences, 362*(1480), 671-678. doi:10.1098/rstb.2006.2003

Fusar-Poli, P., Placentino, A., Carletti, F., Landi, P., Allen, P., Surguladze, S., . . . Politi, P. (2009). Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *Journal of Psychiatry & Neuroscience, 34*, 418-432. doi:10.1111/j.1365-2850.2009.01434.x

Giovanniello, J., Yu, K., Furlan, A., Nachtrab, G. T., Sharma, R., Chen, X., & Li, B. (2020). A Central Amygdala-Globus Pallidus Circuit Conveys Unconditioned Stimulus-Related Information and Controls Fear Learning. *The Journal of Neuroscience, 40*(47), 9043-9054. doi:10.1523/JNEUROSCI.2090-20.2020

Gorno-Tempini, M. L., Pradelli, S., Serafini, M., Pagnoni, G., Baraldi, P., Porro, C., . . . Nichelli, P. (2001). Explicit and Incidental Facial Expression Processing: An fMRI Study. *NeuroImage, 14*(2), 465-473. doi:10.1006/nimg.2001.0811

Hadjistavropoulos, T., Craig, K. D., Duck, S., Cano, A., Goubert, L., Jackson, P. L., . . . Fitzgerald, T. D. (2011). A biopsychosocial formulation of pain communication. *Psychological Bulletin, 137*(6), 910-939. doi:10.1037/a0023876

Haidt, J., McCauley, C., & Rozin, P. (1994). Individual differences in sensitivity to disgust: A scale sampling seven domains of disgust elicitors. *Personality and Individual Differences, 16*(5), 701-713. doi:10.1016/0191-8869(94)90212-7

Harrison, N. A., Gray, M. A., Gianaros, P. J., & Critchley, H. D. (2010). The Embodiment of Emotional Feelings in the Brain. *The Journal of Neuroscience, 30*(38), 12878. doi:10.1523/JNEUROSCI.1725-10.2010

Heining, M., Young, A., Ioannou, G., Andrew, C., Brammer, M., Gray, J., & Phillips, M. (2004). Disgusting Smells Activate Human Anterior Insula and Ventral Striatum. *Annals of the New York Academy of Sciences, 1000*, 380-384. doi:10.1196/annals.1280.035

Herrington, J. D., Taylor, J. M., Grupe, D. W., Curby, K. M., & Schultz, R. T. (2011). Bidirectional communication between amygdala and fusiform gyrus during facial recognition. *NeuroImage, 56*(4), 2348-2355. doi:10.1016/j.neuroimage.2011.03.072

Holtmann, O., Bruchmann, M., Mönig, C., Schwindt, W., Melzer, N., Miltner, W. H. R., & Straube, T. (2020). Lateralized Deficits of Disgust Processing After Insula-Basal Ganglia Damage. *Frontiers in Psychology, 11*(1429). doi:10.3389/fpsyg.2020.01429

Jabbi, M., Bastiaansen, J., & Keysers, C. (2008). A Common Anterior Insula Representation of Disgust Observation, Experience and Imagination Shows Divergent Functional Connectivity Pathways. *PLOS ONE, 3*(8), e2939. doi:10.1371/journal.pone.0002939

Jiang, X., Li, X., Lv, J., Zhang, T., Zhang, S., Guo, L., & Liu, T. (2015). Sparse representation of HCP grayordinate data reveals novel functional architecture of cerebral cortex. *Human Brain Mapping, 36*(12), 5301-5319. doi:10.1002/hbm.23013

Jiang, X., X., L., J., L., S., Z., S., Z., W., Z., . . . T., L. (2018). Temporal Dynamics Assessment of Spatial Overlap Pattern of Functional Brain Networks Reveals Novel Functional Architecture of Cerebral Cortex. *IEEE Transactions on Biomedical Engineering, 65*(6), 1183-1192. doi:10.1109/TBME.2016.2598728

Jiang, X., Zhang, T., Zhang, S., Kendrick, K. M., & Liu, T. (2021). Fundamental functional differences between gyri and sulci: implications for brain function, cognition, and behavior. *Psychoradiology, 1*(1), 23-41. doi:10.1093/psyrad/kkab005

Kipps, C. M., Duggins, A. J., McCusker, E. A., & Calder, A. J. (2007). Disgust and Happiness Recognition Correlate with Anteroventral Insula and Amygdala Volume Respectively in Preclinical Huntington's Disease. *Journal of Cognitive Neuroscience, 19*(7), 1206-1217. doi:10.1162/jocn.2007.19.7.1206

Kirby, L. A. J., & Robinson, J. L. (2017). Affective mapping: An activation likelihood estimation (ALE) meta-analysis. *Brain and Cognition, 118*, 137-148. doi:10.1016/j.bandc.2015.04.006

Klugah-Brown, B., Di, X., Zweerings, J., Mathiak, K., Becker, B., & Biswal, B. (2020). Common and separable neural alterations in substance use disorders: A coordinate-based meta-analyses of functional neuroimaging studies in humans. *Human Brain Mapping, 41*(16), 4459-4477. doi:10.1002/hbm.25085

Klugah-Brown, B., Zhou, X., Pradhan, B. K., Zweerings, J., Mathiak, K., Biswal, B., & Becker, B. (2021). Common neurofunctional dysregulations characterize obsessive–compulsive, substance use, and gaming disorders—An activation likelihood meta-analysis of functional imaging studies. *Addiction Biology, 26*(4), e12997. doi:10.1111/adb.12997

Kohn, N., Eickhoff, S. B., Scheller, M., Laird, A. R., Fox, P. T., & Habel, U. (2014). Neural network of cognitive emotion regulation — An ALE meta-analysis and MACM analysis. *NeuroImage, 87*, 345-355. doi:10.1016/j.neuroimage.2013.11.001

Kotkowski, E., Price, L. R., Mickle Fox, P., Vanasse, T. J., & Fox, P. T. (2018). The hippocampal network model: A transdiagnostic metaconnectomic approach. *NeuroImage: Clinical, 18*, 115-129. doi:10.1016/j.nicl.2018.01.002

Krolak-Salmon, P., Hénaff, M.-A., Isnard, J., Tallon-Baudry, C., Guénot, M., Vighetto, A., . . . Mauguière, F. (2003). An attention modulated response to disgust in human ventral anterior insula. *Annals of Neurology, 53*(4), 446-453. doi:10.1002/ana.10502

Kupfer, T. R., & Fessler, D. M. T. (2018). Ectoparasite defence in humans: relationships to pathogen avoidance and clinical implications. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences, 373*(1751), 20170207. doi:doi:10.1098/rstb.2017.0207

Laird, A. R., Eickhoff, S. B., Li, K., Robin, D. A., Glahn, D. C., & Fox, P. T. (2009). Investigating the Functional Heterogeneity of the Default Mode Network Using Coordinate-Based Meta-Analytic Modeling. *The Journal of Neuroscience, 29*(46), 14496. doi:10.1523/JNEUROSCI.4004-09.2009

Laird, A. R., Fox, P. M., Price, C. J., Glahn, D. C., Uecker, A. M., Lancaster, J. L., . . . Fox, P. T. (2005). ALE meta-analysis: controlling the false discovery rate and performing statistical contrasts. *Human Brain Mapping, 25*(1), 155-164. doi:10.1002/hbm.20136

Laird, A. R., Robinson, J. L., McMillan, K. M., Tordesillas-Gutiérrez, D., Moran, S. T., Gonzales, S. M., . . . Lancaster, J. L. (2010). Comparison of the disparity between Talairach and MNI coordinates in functional neuroimaging data: Validation of the Lancaster transform. *NeuroImage, 51*(2), 677-683. doi:10.1016/j.neuroimage.2010.02.048

Lancaster, J. L., Tordesillas-Gutiérrez, D., Martinez, M., Salinas, F., Evans, A., Zilles, K., . . . Fox, P. T. (2007). Bias between MNI and Talairach coordinates analyzed using the ICBM-152 brain template. *Human Brain Mapping, 28*(11), 1194-1205. doi:10.1002/hbm.20345

Langner, R., Rottschy, C., Laird, A. R., Fox, P. T., & Eickhoff, S. B. (2014). Meta-analytic connectivity modeling revisited: Controlling for activation base rates. *NeuroImage, 99*, 559-570. doi:10.1016/j.neuroimage.2014.06.007

Lassalle, A., Zürcher, N. R., Porro, C. A., Benuzzi, F., Hippolyte, L., Lemonnier, E., . . . Hadjikhani, N. (2019). Influence of anxiety and alexithymia on brain activations associated with the perception of

others' pain in autism. *Social Neuroscience, 14*(3), 359-377. doi:10.1080/17470919.2018.1468358

LeDoux, J. E., & Pine, D. S. (2016). Using Neuroscience to Help Understand Fear and Anxiety: A Two-System Framework. *American Journal of Psychiatry, 173*(11), 1083-1093. doi:10.1176/appi.ajp.2016.16030353

Li, J., Xu, L., Zheng, X., Fu, M., Zhou, F., Xu, X., . . . Becker, B. (2019). Common and Dissociable Contributions of Alexithymia and Autism to Domain-Specific Interoceptive Dysregulations: A Dimensional Neuroimaging Approach. *Psychotherapy and Psychosomatics, 88*(3), 187-189. doi:10.1159/000495122

Lindquist, D. H. (2020). Emotion in motion: A three-stage model of aversive classical conditioning. *Neuroscience & Biobehavioral Reviews, 115*, 363-377. doi:10.1016/j.neubiorev.2020.04.025

Liu, Z., Rolls, E. T., Liu, Z., Zhang, K., Yang, M., Du, J., . . . Feng, J. (2019). Brain annotation toolbox: exploring the functional and genetic associations of neuroimaging results. *Bioinformatics, 35*(19), 3771-3778. doi:10.1093/bioinformatics/btz128

Malhi, G., Lagopoulos, J., Sachdev, P., Ivanovski, B., Shnier, R., & Ketter, T. (2007). Is a lack of disgust something to fear? An fMRI facial recognition study in euthymic bipolar disorder patients. *Bipolar Disorders, 9*, 345-357. doi:10.1111/j.1399-5618.2007.00485.x

Mattavelli, G., Sormaz, M., Flack, T., Asghar, A. U. R., Fan, S., Frey, J., . . . Andrews, T. J. (2014). Neural responses to facial expressions support the role of the amygdala in processing threat. *Social Cognitive and Affective Neuroscience, 9*(11), 1684-1689. doi:10.1093/scan/nst162

Meier, L., Friedrich, H., Federspiel, A., Jann, K., Morishima, Y., Landis, B. N., . . . Dierks, T. (2015). Rivalry of homeostatic and sensory-evoked emotions: Dehydration attenuates olfactory disgust and its neural correlates. *NeuroImage, 114*, 120-127. doi:10.1016/j.neuroimage.2015.03.048

Mesulam, M. M., & Mufson, E. J. (1982). Insula of the old world monkey. III: Efferent cortical output and comments on function. *Journal of Comparative Neurology, 212*(1), 38-52. doi:10.1002/cne.902120104

Mihov, Y., Kendrick, K. M., Becker, B., Zschernack, J., Reich, H., Maier, W., . . . Hurlemann, R. (2013). Mirroring Fear in the Absence of a Functional Amygdala. *Biological Psychiatry, 73*(7), e9-e11. doi:10.1016/j.biopsych.2012.10.029

Molapour, T., Golkar, A., Navarrete, C. D., Haaker, J., & Olsson, A. (2015). Neural correlates of biased social fear learning and interaction in an intergroup context. *NeuroImage, 121*, 171-183. doi:10.1016/j.neuroimage.2015.07.015

Morawetz, C., Bode, S., Baudewig, J., & Heekeren, H. R. (2017). Effective amygdala-prefrontal connectivity predicts individual differences in successful emotion regulation. *Social Cognitive and Affective Neuroscience, 12*(4), 569-585. doi:10.1093/scan/nsw169

Müller, V. I., Cieslik, E. C., Laird, A. R., Fox, P. T., Radua, J., Mataix-Cols, D., . . . Eickhoff, S. B. (2018). Ten simple rules for neuroimaging meta-analysis. *Neuroscience and Biobehavioral Reviews, 84*, 151-161. doi:10.1016/j.neubiorev.2017.11.012

Murphy, F. C., Nimmo-Smith, I., & Lawrence, A. D. (2003). Functional neuroanatomy of emotions: a meta-analysis. *Cognitive, Affective & Behavioral Neuroscience, 3*(3), 207-233. doi:10.3758/cabn.3.3.207

Oaten, M., Stevenson, R. J., & Case, T. I. (2009). Disgust as a disease-avoidance mechanism. *Psychological Bulletin, 135*(2), 303-321. doi:10.1037/a0014823

Olatunji, B. O., & Sawchuk, C. (2005). Disgust: Characteristic Features, Social Manifestations, and Clinical

Implications. *Journal of Social and Clinical Psychology, 24*(7), 932-962. doi:10.1521/jscp.2005.24.7.932

Olatunji, B. O., Williams, N. L., Tolin, D. F., Abramowitz, J. S., Sawchuk, C. N., Lohr, J. M., & Elwood, L. S. (2007). The Disgust Scale: item analysis, factor structure, and suggestions for refinement. *Psychological assessment, 19*(3), 281-297. doi:10.1037/1040-3590.19.3.281

Oliveri, M., Babiloni, C., Filippi, M. M., Caltagirone, C., Babiloni, F., Cicinelli, P., . . . Rossini, P. M. (2003). Influence of the supplementary motor area on primary motor cortex excitability during movements triggered by neutral or emotionally unpleasant visual cues. *Experimental Brain Research, 149*(2), 214-221. doi:10.1007/s00221-002-1346-8

Page, M. J., Moher, D., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., . . . McKenzie, J. E. (2021). PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ, 372*, n160. doi:10.1136/bmj.n160

Papitto, G., Friederici, A. D., & Zaccarella, E. (2020). The topographical organization of motor processing: An ALE meta-analysis on six action domains and the relevance of Broca's region. *NeuroImage, 206*, 116321. doi:10.1016/j.neuroimage.2019.116321

Phan, K. L., Wager, T., Taylor, S. F., & Liberzon, I. (2002). Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage, 16*(2), 331-348. doi:10.1006/nimg.2002.1087

Phelps, E. A., & LeDoux, J. E. (2005). Contributions of the Amygdala to Emotion Processing: From Animal Models to Human Behavior. *Neuron, 48*(2), 175-187. doi:10.1016/j.neuron.2005.09.025

Phillips, M., Marks, I., Senior, C., Lythgoe, D., O'Dwyer, A., Meehan, O., . . . McGuire, P. (2000). A differential neural response in obsessive-compulsive disorder patients with washing compared with checking symptoms to disgust. *Psychological Medicine, 30*, 1037-1050. doi:10.1017/S0033291799002652

Phillips, M., Williams, L., Heining, M., Herba, C., Russell, T., Andrew, C., . . . Gray, J. (2004). Differential Neural Responses to Overt and Covert Presentations of Facial Expressions of Fear and Disgust. *NeuroImage, 21*, 1484-1496. doi:10.1016/j.neuroimage.2003.12.013

Phillips, M., Williams, L., Senior, C., Bullmore, E. T., Brammer, M. J., Andrew, C., . . . David, A. S. (1999). A differential neural response to threatening and non-threatening negative facial expressions in paranoid and non-paranoid schizophrenics. *Psychiatry Research: Neuroimaging, 92*(1), 11-31. doi:10.1016/S0925-4927(99)00031-1

Phillips, M., Young, A. W., Scott, S. K., Calder, A. J., Andrew, C., Giampietro, V., . . . Gray, J. A. (1998). Neural responses to facial and vocal expressions of fear and disgust. *Proceedings of the Royal Society B: Biological Sciences, 265*(1408), 1809-1817. doi:doi:10.1098/rspb.1998.0506

Phillips, M., Young, A. W., Senior, C., Brammer, M., Andrew, C., Calder, A. J., . . . David, A. S. (1997). A specific neural substrate for perceiving facial expressions of disgust. *Nature, 389*(6650), 495-498. doi:10.1038/39051

Pujol, J., Blanco-Hinojo, L., Coronas, R., Esteba-Castillo, S., Rigla, M., Martínez-Vilavella, G., . . . Caixàs, A. (2018). Mapping the sequence of brain events in response to disgusting food. *Human Brain Mapping, 39*(1), 369-380. doi:10.1002/hbm.23848

Reid, A. T., Hoffstaedter, F., Gong, G., Laird, A. R., Fox, P., Evans, A. C., . . . Eickhoff, S. B. (2017). A seed-based cross-modal comparison of brain connectivity measures. *Brain Structure and Function, 222*(3), 1131-1151. doi:10.1007/s00429-016-1264-3

Robinson, J. L., Laird, A. R., Glahn, D. C., Blangero, J., Sanghera, M. K., Pessoa, L., . . . Fox, P. T. (2012).

The functional connectivity of the human caudate: An application of meta-analytic connectivity modeling with behavioral filtering. *NeuroImage, 60*(1), 117-129. doi:10.1016/j.neuroimage.2011.12.010

Robinson, J. L., Laird, A. R., Glahn, D. C., Lovallo, W. R., & Fox, P. T. (2010). Metaanalytic connectivity modeling: Delineating the functional connectivity of the human amygdala. *Human Brain Mapping, 31*(2), 173-184. doi:10.1002/hbm.20854

Rolls, E. T. (1999). *The Brain and Emotion*. New York: Oxford University Press.

Rolls, E. T. (2000a). The Orbitofrontal Cortex and Reward. *Cerebral Cortex, 10*(3), 284-294. doi:10.1093/cercor/10.3.284

Rolls, E. T. (2000b). Précis of The brain and emotion. *Behavioral and Brain Sciences, 23*(2), 177-191. doi:10.1017/S0140525X00002429

Rolls, E. T. (2004). The functions of the orbitofrontal cortex. *Brain and Cognition, 55*(1), 11-29. doi:10.1016/S0278-2626(03)00277-X

Rozin, P., & Fallon, A. E. (1987). A perspective on disgust. *Psychological Review, 94*(1), 23-41. doi:10.1037/0033-295X.94.1.23

Rozin, P., Haidt, J., & Fincher, K. (2009). From Oral to Moral. *Science, 323*(5918), 1179. doi:10.1126/science.1170492

Rozin, P., Haidt, J., & McCauley, C. (2016). Disgust. In L. F. Barrett, M. Lewis, & J. M. Haviland-Jone (Eds.), *Handbook of Emotions, Fourth Edition* (pp. 815-834). New York: Guilford Publications.

Rozin, P., Lowery, L., & Ebert, R. (1994). Varieties of disgust faces and the structure of disgust. *Journal of Personality and Social Psychology, 66*(5), 870-881. doi:10.1037//0022-3514.66.5.870

Salih, F., Sharott, A., Khatami, R., Trottenberg, T., Schneider, G., Kupsch, A., . . . Grosse, P. (2009). Functional connectivity between motor cortex and globus pallidus in human non-REM sleep. *The Journal of Physiology, 587*(Pt 5), 1071-1086. doi:10.1113/jphysiol.2008.164327

Sambataro, F., Dimalta, S., Di Giorgio, A., Taurisano, P., Blasi, G., Scarabino, T., . . . Bertolino, A. (2006). Preferential responses in amygdala and insula during presentation of facial contempt and disgust. *European Journal of Neuroscience, 24*, 2355-2362. doi:10.1111/j.1460-9568.2006.05120.x

Schäfer, A., Leutgeb, V., Reishofer, G., Ebner, F., & Schienle, A. (2009). Propensity and sensitivity measures of fear and disgust are differentially related to emotion-specific brain activation. *Neuroscience Letters, 465*(3), 262-266. doi:10.1016/j.neulet.2009.09.030

Schäfer, A., Schienle, A., & Vaitl, D. (2005). Stimulus type and design influence hemodynamic responses towards visual disgust and fear elicitors. *International Journal of Psychophysiology, 57*(1), 53-59. doi:10.1016/j.ijpsycho.2005.01.011

Schienle, A., Höfler, C., Keck, T., & Wabnegger, A. (2020). Neural underpinnings of perception and experience of disgust in individuals with a reduced sense of smell: An fMRI study. *Neuropsychologia, 141*, 107411. doi:10.1016/j.neuropsychologia.2020.107411

Schienle, A., Schäfer, A., Hermann, A., & Vaitl, D. (2009). Binge-Eating Disorder: Reward Sensitivity and Brain Activation to Images of Food. *Biological Psychiatry, 65*(8), 654-661. doi:10.1016/j.biopsych.2008.09.028

Schienle, A., Schäfer, A., Hermann, A., Walter, B., Stark, R., & Vaitl, D. (2006). fMRI responses to pictures of mutilation and contamination. *Neuroscience Letters, 393*(2), 174-178. doi:10.1016/j.neulet.2005.09.072

Schienle, A., Schäfer, A., Stark, R., Walter, B., & Vaitl, D. (2005a). Gender differences in the processing of

disgust- and fear-inducing pictures: an fMRI study. *Neuroreport, 16*(3), 277-280. doi:10.1097/00001756-200502280-00015

Schienle, A., Schäfer, A., Stark, R., Walter, B., & Vaitl, D. (2005b). Relationship between Disgust Sensitivity, Trait Anxiety and Brain Activity during Disgust Induction. *Neuropsychobiology, 51*, 86-92. doi:10.1159/000084165

Schienle, A., & Scharmüller, W. (2013). Cerebellar activity and connectivity during the experience of disgust and happiness. *Neuroscience, 246*, 375-381. doi:10.1016/j.neuroscience.2013.04.048

Schienle, A., Stark, R., Walter, B., Blecker, C., Ott, U., Kirsch, P., . . . Vaitl, D. (2002). The insula is not specifically involved in disgust processing: an fMRI study. *Neuroreport, 13*(16), 2023-2026. doi:10.1097/00001756-200211150-00006

Schienle, A., Übel, S., Schöngaßner, F., Ille, R., & Scharmüller, W. (2014). Disgust regulation via placebo: An fMRI study. *Social Cognitive and Affective Neuroscience, 9*(7), 985-990. doi:10.1093/scan/nst072

Schienle, A., Übel, S., & Wabnegger, A. (2016). When opposites lead to the same: a direct comparison of explicit and implicit disgust regulation via fMRI. *Social Cognitive and Affective Neuroscience, 12*(3), 445-451. doi:10.1093/scan/nsw144

Schienle, A., & Wabnegger, A. (2021). The processing of visual food cues during bitter aftertaste perception in females with high vs. low disgust propensity: an fMRI study. *Brain Imaging and Behavior*. doi:10.1007/s11682-021-00455-2

Schroeder, U., Hennenlotter, A., Erhard, P., Haslinger, B., Stahl, R., Lange, K., & Ceballos-Baumann, A. (2005). Functional neuroanatomy of perceiving surprised faces. *Human Brain Mapping, 23*, 181-187. doi:10.1002/hbm.20057

Shapira, N. A., Liu, Y., He, A. G., Bradley, M. M., Lessig, M. C., James, G. A., . . . Goodman, W. K. (2003). Brain activation by disgust-inducing pictures in obsessive-compulsive disorder. *Biological Psychiatry, 54*(7), 751-756. doi:10.1016/S0006-3223(03)00003-9

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., . . . Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during activation and rest. *Proceedings of the National Academy of Sciences, 106*(31), 13040. doi:10.1073/pnas.0905267106

Sprengelmeyer, R., Rausch, M., Eysel, U., & Przuntek, H. (1998). Neural structures associated with recognition of facial expressions of basic emotions. *Proceedings of the Royal Society B-Biological Sciences, 265*, 1927-1931. doi:10.1098/rspb.1998.0522

Sprengelmeyer, R., Young, A. W., Calder, A. J., Karnat, A., Lange, H., Hömberg, V., . . . Rowland, D. (1996). Loss of disgust: Perception of faces and emotions in Huntington's disease. *Brain, 119*(5), 1647-1665. doi:10.1093/brain/119.5.1647

Stark, R., Schienle, A., Girod, C., Walter, B., Kirsch, P., Blecker, C., . . . Vaitl, D. (2005). Erotic and disgust-inducing pictures—Differences in the hemodynamic responses of the brain. *Biological Psychology, 70*(1), 19-29. doi:10.1016/j.biopsycho.2004.11.014

Stark, R., Schienle, A., Walter, B., Kirsch, P., Blecker, C., Ott, U., . . . Vaitl, D. (2004). Hemodynamic Effects of Negative Emotional Pictures – A Test-Retest Analysis. *Neuropsychobiology, 50*(1), 108-118. doi:10.1159/000077948

Stark, R., Schienle, A., Walter, B., Kirsch, P., Sammer, G., Ott, U., . . . Vaitl, D. (2003). Hemodynamic responses to fear and disgust-inducing pictures: an fMRI study. *International Journal of Psychophysiology, 50*(3), 225-234. doi:10.1016/S0167-8760(03)00169-7

Stark, R., Zimmermann, M., Kagerer, S., Schienle, A., Walter, B., Weygandt, M., & Vaitl, D. (2007).

Hemodynamic brain correlates of disgust and fear ratings. *NeuroImage, 37*(2), 663-673. doi:10.1016/j.neuroimage.2007.05.005

Straube, T., Weisbrod, A., Schmidt, S., Raschdorf, C., Preul, C., Mentzel, H.-J., & Miltner, W. H. R. (2010). No impairment of recognition and experience of disgust in a patient with a right-hemispheric lesion of the insula and basal ganglia. *Neuropsychologia, 48*(6), 1735-1741. doi:10.1016/j.neuropsychologia.2010.02.022

Surguladze, S. A., Brammer, M. J., Young, A. W., Andrew, C., Travis, M. J., Williams, S. C. R., & Phillips, M. L. (2003). A preferential increase in the extrastriate response to signals of danger. *NeuroImage, 19*(4), 1317-1328. doi:10.1016/S1053-8119(03)00085-5

Surguladze, S. A., El-Hage, W., Dalgleish, T., Radua, J., Gohier, B., & Phillips, M. L. (2010). Depression is associated with increased sensitivity to signals of disgust: A functional magnetic resonance imaging study. *Journal of Psychiatric Research, 44*(14), 894-902. doi:10.1016/j.jpsychires.2010.02.010

Tao, D., He, Z., Lin, Y., Liu, C., & Tao, Q. (2021). Where does fear originate in the brain? A coordinate-based meta-analysis of explicit and implicit fear processing. *NeuroImage, 227*, 117686. doi:10.1016/j.neuroimage.2020.117686

Tettamanti, M., Rognoni, E., Cafiero, R., Costa, T., Galati, D., & Perani, D. (2012). Distinct pathways of neural coupling for different basic emotions. *NeuroImage, 59*(2), 1804-1817. doi:10.1016/j.neuroimage.2011.08.018

Trautmann, S. A., Fehr, T., & Herrmann, M. (2009). Emotions in motion: Dynamic compared to static facial expressions of disgust and happiness reveal more widespread emotion-specific activations. *Brain Research, 1284*, 100-115. doi:10.1016/j.brainres.2009.05.075

Tu, S., Qiu, J., Martens, U., & Zhang, Q. (2013). Category-selective attention modulates unconscious processes in the middle occipital gyrus. *Consciousness and Cognition, 22*(2), 479-485. doi:10.1016/j.concog.2013.02.007

Turk-Browne, N., Norman-Haignere, S., & McCarthy, G. (2010). Face-Specific Resting Functional Connectivity between the Fusiform Gyrus and Posterior Superior Temporal Sulcus. *Frontiers in Human Neuroscience, 4*(176). doi:10.3389/fnhum.2010.00176

Turkeltaub, P. E., Eickhoff, S. B., Laird, A. R., Fox, M., Wiener, M., & Fox, P. (2012). Minimizing within-experiment and within-group effects in Activation Likelihood Estimation meta-analyses. *Human Brain Mapping, 33*(1), 1-13. doi:10.1002/hbm.21186

Uddin, L. Q. (2015). Salience processing and insular cortical function and dysfunction. *Nature Reviews Neuroscience, 16*(1), 55-61. doi:10.1038/nrn3857

van Overveld, W. J. M., de Jong, P. J., & Peters, M. L. (2009). Digestive and cardiovascular responses to core and animal-reminder disgust. *Biological Psychology, 80*(2), 149-157. doi:10.1016/j.biopsycho.2008.08.002

Vicario, C. M., Rafal, R. D., Martino, D., & Avenanti, A. (2017). Core, social and moral disgust are bounded: A review on behavioral and neural bases of repugnance in clinical disorders. *Neuroscience and Biobehavioral Reviews, 80*, 185-200. doi:10.1016/j.neubiorev.2017.05.008

Viinikainen, M., Jääskeläinen, I. P., Alexandrov, Y., Balk, M. H., Autti, T., & Sams, M. (2010). Nonlinear relationship between emotional valence and brain activity: Evidence of separate negative and positive valence dimensions. *Human Brain Mapping, 31*(7), 1030-1040. doi:10.1002/hbm.20915

von dem Hagen, E. A. H., Beaver, J. D., Ewbank, M. P., Keane, J., Passamonti, L., Lawrence, A. D., & Calder,

A. J. (2009). Leaving a bad taste in your mouth but not in my insula. *Social Cognitive and Affective Neuroscience, 4*(4), 379-386. doi:10.1093/scan/nsp018

Vytal, K., & Hamann, S. (2010). Neuroimaging support for discrete neural correlates of basic emotions: a voxel-based meta-analysis. *Journal of Cognitive Neuroscience, 22*(12), 2864-2885. doi:10.1162/jocn.2009.21366

Wager, T. D., Kang, J., Johnson, T. D., Nichols, T. E., Satpute, A. B., & Barrett, L. F. (2015). A Bayesian Model of Category-Specific Emotional Brain Responses. *PLOS Computational Biology, 11*(4), e1004066. doi:10.1371/journal.pcbi.1004066

Wager, T. D., Lindquist, M., & Kaplan, L. (2007). Meta-analysis of functional neuroimaging data: current and future directions. *Social Cognitive and Affective Neuroscience, 2*(2), 150-158. doi:10.1093/scan/nsm015

Wager, T. D., Phan, K. L., Liberzon, I., & Taylor, S. F. (2003). Valence, gender, and lateralization of functional brain anatomy in emotion: a meta-analysis of findings from neuroimaging. *NeuroImage, 19*(3), 513-531. doi:10.1016/S1053-8119(03)00078-8

Wang, Y., Braver, T. S., Yin, S., Hu, X., Wang, X., & Chen, A. (2019). Reward improves response inhibition by enhancing attentional capture. *Social Cognitive and Affective Neuroscience, 14*(1), 35-45. doi:10.1093/scan/nsy111

Weiner, K. S., & Zilles, K. (2016). The anatomical and functional specialization of the fusiform gyrus. *Neuropsychologia, 83*, 48-62. doi:10.1016/j.neuropsychologia.2015.06.033

Wertheim, J., & Ragni, M. (2018). The Neural Correlates of Relational Reasoning: A Meta-analysis of 47 Functional Magnetic Resonance Studies. *Journal of Cognitive Neuroscience, 30*(11), 1734-1748. doi:10.1162/jocn_a_01311

Whalen, P. J. (2007). The uncertainty of it all. *Trends in Cognitive Sciences, 11*(12), 499-500. doi:10.1016/j.tics.2007.08.016

Wicker, B., Keysers, C., Plailly, J., Royet, J.-P., Gallese, V., & Rizzolatti, G. (2003). Both of Us Disgusted in My Insula: The Common Neural Basis of Seeing and Feeling Disgust. *Neuron, 40*(3), 655-664. doi:10.1016/S0896-6273(03)00679-2

Winston, J. S., O'Doherty, J., & Dolan, R. J. (2003). Common and distinct neural responses during direct and incidental processing of multiple facial emotions. *NeuroImage, 20*(1), 84-97. doi:10.1016/S1053-8119(03)00303-3

Wright, P., He, G., Shapira, N., Goodman, W., & Liu, Y. (2004). Disgust and the Insula: fMRI Responses to Pictures of Mutilation and Contamination. *Neuroreport, 15*, 2347-2351. doi:10.1097/00001756-200410250-00009

Yamasaki, H., LaBar, K. S., & McCarthy, G. (2002). Dissociable prefrontal brain systems for attention and emotion. *Proceedings of the National Academy of Sciences, 99*(17), 11447. doi:10.1073/pnas.182176499

Yao, S., Becker, B., Zhao, W., Zhao, Z., Kou, J., Ma, X., . . . Kendrick, K. M. (2018). Oxytocin Modulates Attention Switching Between Interoceptive Signals and External Social Cues. *Neuropsychopharmacology, 43*(2), 294-301. doi:10.1038/npp.2017.189

Yao, S., Zhao, W., Geng, Y., Chen, Y., Zhao, Z., Ma, X., . . . Kendrick, K. M. (2018). Oxytocin Facilitates Approach Behavior to Positive Social Stimuli via Decreasing Anterior Insula Activity. *International Journal of Neuropsychopharmacology, 21*(10), 918-925. doi:10.1093/ijnp/pyy068

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature Methods, 8*(8), 665-670.

doi:10.1038/nmeth.1635

Yu, F., Barron, D. S., Tantiwongkosi, B., Fox, M., & Fox, P. (2018). Characterisation of meta-analytical functional connectivity in progressive supranuclear palsy. *Clinical Radiology, 73*(4), 415.e411-415.e417. doi:10.1016/j.crad.2017.11.007

Zald, D. H. (2003). The human amygdala and the emotional evaluation of sensory stimuli. *Brain Research Reviews, 41*(1), 88-123. doi:10.1016/S0165-0173(02)00248-5

Zhou, F., Li, J., Zhao, W., Xu, L., Zheng, X., Fu, M., . . . Becker, B. (2020). Empathic pain evoked by sensory and emotional-communicative cues share common and process-specific neural representations. *eLife, 9*, e56929. doi:10.7554/eLife.56929

Zhou, F., Zhao, W., Qi, Z., Geng, Y., Yao, S., Kendrick, K. M., . . . Becker, B. (2020). Beyond fear centers – a distributed fMRI-based neuromarker for the subjective experience of fear. *BioRvix.org*. doi:10.1101/2020.11.23.394973

Ziegler, J., Montant, M., Briesemeister, B., Brink, T., Wicker, B., Ponz, A., . . . Braun, M. (2018). Do Words Stink? Neural Reuse as a Principle for Understanding Emotions in Reading. *Journal of Cognitive Neuroscience, 30*, 1023-1032. doi:10.1162/jocn_a_01268

Zilbovicius, M., Meresse, I., Chabane, N., Brunelle, F., Samson, Y., & Boddaert, N. (2006). Autism, the superior temporal sulcus and social perception. *Trends in Neurosciences, 29*(7), 359-366. doi:10.1016/j.tins.2006.06.004

Zimmer, U., Höfler, M., Koschutnig, K., & Ischebeck, A. (2016). Neuronal interactions in areas of spatial attention reflect avoidance of disgust, but orienting to danger. *NeuroImage, 134*, 94-104. doi:10.1016/j.neuroimage.2016.03.050