1

A generalized cortical activity pattern at internally-generated

mental context boundaries during unguided narrative recall

Hongmi Lee[1*], Janice Chen[1]

[1]Department of Psychological and Brain Sciences,

Johns Hopkins University,

Baltimore, MD 21218, USA

*Correspondence: Hongmi Lee (hongmi.lee@jhu.edu)

Abstract

Current theory and empirical studies suggest that humans segment continuous experiences into

events based on the mismatch between predicted and actual sensory inputs; detection of these

"event boundaries" evokes transient neural responses. However, boundaries can also occur at

transitions between internal mental states, without relevant external input changes. To what

extent do such "internal boundaries" share neural response properties with externally-driven

boundaries? We conducted an fMRI experiment where subjects watched a series of short

movies and then verbally recalled the movies, unprompted, in the order of their choosing.

During recall, transitions between movies thus constituted major boundaries between internal

mental contexts, generated purely by subjects' unguided thoughts. Following the offset of each

recalled movie, we observed stereotyped spatial activation patterns in the default mode

network, especially the posterior medial cortex, consistent across different movie contents and

even across the different tasks of movie watching and recall. Surprisingly, the between-movie

boundary patterns were negatively correlated with patterns at boundaries between events within

a movie. Thus, major transitions between mental contexts elicit neural phenomena shared

across internal and external modes and distinct from within-context event boundary detection,

potentially reflecting a cognitive state related to the flushing and reconfiguration of situation

models.

Introduction

Humans perceive and remember continuous experiences as discrete events (Brunec et al., 2018; Clewett et al., 2019; Shin & DuBrow, 2021; Zacks, 2020). Studies of event segmentation have shown that when participants attend to external information (e.g., watch a video), 1) boundaries between events are detected when mismatches arise between predicted and actual sensory input (Zacks et al., 2007, 2011), and 2) boundary detection evokes transient neural responses in a consistent set of brain areas (Speer et al., 2007; Zacks et al., 2001; Zheng et al., 2021). Among these areas is the default mode network (DMN; Buckner & DiNicola, 2019) proposed to be involved in representing complex mental models of events (Ranganath & Ritchey, 2012; Ritchey & Cooper, 2020). However, a substantial portion of human cognition is internally-driven (Hasselmo, 1995; Honey et al., 2017), and such spontaneous production of thoughts and actions is also punctuated by mental context transitions (Christoff et al., 2016; Mildner & Tamir, 2019; Smallwood & Schooler, 2015; Tseng & Poppenk, 2020). What manner of brain activity marks boundaries between mental contexts when they are internally-generated? Are the brain responses at internal boundaries similar to those at external boundaries?

Here, we used naturalistic movie viewing and free spoken recall with fMRI to characterize neural activity at boundaries between internally-generated mental contexts (Figure 1A). Subjects watched ten short movies (encoding phase), then verbally recounted the movies in any order, in their own words (recall phase). The transitions between recalled movies were determined purely by subjects' internal mentation; no external cues prompted the recall onset or offset of each movie. Moreover, the unguided spoken recall allowed us to identify the exact moments of context transitions and explicitly track shifts in the contents of thoughts (Chen et al., 2017; Sripada & Taxali, 2020), which was not possible in prior studies using silent rest (Karapanagiotidis et al., 2020; Tseng & Poppenk, 2020). We observed limited changes in univariate activity at these internal boundaries between recalled movies. However, we observed transient, highly generalizable and fine-grained activation patterns throughout the DMN, consistent across diverse movie contents and similar to those at external between-movie boundaries during encoding. Moreover, these between-movie boundary patterns were not merely stronger versions of within-movie "event boundaries", but instead manifested as a distinct type of neural transition. We propose that these cortical patterns reflect a cognitive state related to the flushing and reconfiguration of mental context (DuBrow et al., 2017; Manning et al., 2016).

Figure 1. Experimental procedures and univariate responses. (A) In the encoding phase, subjects watched 10 short movies approximately 2 to 8 minutes long. Each movie started with a 6-second title scene. In the free spoken recall phase, subjects verbally recounted each movie plot in as much detail as possible regardless of the order of presentation. After recalling one movie, subjects spontaneously proceeded to the next movie, and the transitions between movies were considered as internally-driven boundaries. Red arrows indicate the boundaries (onsets and offsets) between watched or recalled movies. Black arrows indicate the non-boundary moments (middle) of each watched or recalled movie. (B) Whole-brain maps of mean activation (BOLD signals z-scored across all volumes within a scanning run) following between-movie boundaries during recall (4.5 – 19.5 seconds from the offset of each movie). Blue areas indicate regions with lower-than-average activation. Red areas indicate regions with higher-than-average activation. White dots on the lateral surface of the left hemisphere (the leftmost map) indicate vertices that showed significantly lower activation following between-movie boundaries compared to non-boundary periods (Bonferroni corrected $p < .05$). The non-boundary periods were defined as the middle 15 seconds of each recalled movie, shifted forward by 4.5 seconds. Changes in whole-brain univariate responses across time around the boundaries are shown in Figure 1 supplement video 1 (recall phase) and Figure 1 supplement video 2 (encoding phase).

Results and discussion

We first examined whether internally-driven boundaries evoke changes in blood oxygen level-dependent (BOLD) signals during recall. We observed limited transient changes in activation at the boundaries between recalled movies in multiple cortical regions; a whole-brain analysis with multiple comparisons correction revealed that the mean activation of boundary periods (15 seconds following the offset of each movie) was lower than that of non-boundary periods (middle 15 seconds within each movie) in a small number of vertices within the left motor, auditory, and inferior parietal cortices (Figure 1B), although weaker responses were observed in widespread cortical regions (Figure 1 supplement video 1).

Figure 2. Consistent activation patterns associated with between-movie boundaries. (A) Schematic of the pattern similarity analysis. Boundary patterns were defined as the mean pattern averaged across 15 seconds following the offset of each watched or recalled movie. Non-boundary patterns were defined as the mean pattern averaged across 15 seconds in the middle of each watched or recalled movie. For each subject and cortical parcel (Schaefer et al., 2018; 200 parcels per hemisphere), we computed pairwise between-movie pattern similarity (Pearson correlation), separately for boundary patterns and non-boundary patterns measured during recall (a & b, blue arrows). We also computed between-movie and between-phase (encoding-recall) pattern similarity, again separately for boundary and non-boundary patterns (c & d, red arrows). The time windows for both boundary and non-boundary periods were shifted forward by 4.5 seconds to account for the hemodynamic response delay. (B) Whole-brain *t* statistic map of cortical parcels that showed consistent between-movie boundary patterns during recall. These parcels displayed significantly greater between-movie pattern similarity in the boundary condition compared to the non-boundary condition during recall. The map was masked by parcels that showed significantly positive between-movie pattern similarity in the boundary condition during recall. Both effects were Bonferroni corrected across parcels (*p* < .05). (C) Whole-brain *t* statistic map of cortical parcels that showed consistent between-movie boundary patterns across encoding and recall. These parcels displayed significantly greater between-movie and between-phase pattern similarity in the boundary condition compared to the non-boundary condition. The map was masked by parcels that showed significantly positive between-movie and between-phase pattern similarity in the boundary condition. Both effects were Bonferroni corrected across parcels (*p* < .05).

Next, we tested whether there were neural activation *patterns* specific to internally-driven boundaries and consistent across different movies. We performed a whole-brain pattern similarity analysis on the recall data to identify regions where 1) boundary period activation patterns were positively correlated across different recalled movies, and 2) this correlation was

higher at boundaries than at non-boundaries (Figure 2A, blue arrows). We observed a

consistent boundary pattern, i.e., whenever participants transitioned from talking about one

movie to the next, in several cortical parcels (Schaefer et al., 2018) including the DMN and

auditory/motor areas (Figure 2B), suggesting the involvement of both cognitive and

sensory/motor factors at recall boundaries (e.g., breaks in recall speech generation and memory

retrieval).

To what extent is the internally-driven boundary pattern, measured during recall, similar

to patterns observed at boundaries during encoding? To test this, we again computed between-

movie pattern similarity, but now across the encoding and recall phases (Figure 2A, red arrows).

We found that DMN areas, especially the posterior medial cortex (PMC), showed a consistent

boundary pattern across task phases (encoding and recall) and across movies (Figure 2C).

Thus, we next examined the phenomenon in more detail specifically in PMC. Figures 3A and 3C

visualize the high and consistently positive correlations of PMC boundary patterns across

different movies both within the recall phase (Recall Offset vs. Recall Offset, $t(14) = 11.82$, p

$< .001$, Cohen's $d_z = 3.05$) and even between experimental phases (Recall Offset vs. Encoding

Offset, $t(14) = 14.54$, $p < .001$, Cohen's $d_z = 3.75$). No such correlation was present between

non-boundary patterns ($t(14)$s < 1, ps $> .3$). Individual subjects' activation maps visualize the

similarity between boundary patterns during encoding and recall (Figure 3B, Figure 3

supplement 1). We observed similar results in the lateral parietal DMN sub-region (angular

gyrus; Figure 3 supplement 2). This observation of a generalized boundary pattern in DMN

regions extends prior reports of DMN univariate responses at external event boundaries (Reagh

et al., 2020; Speer et al., 2007) and transitions between tasks (Crittenden et al., 2015; Smith et

al., 2018).

Is the generalized boundary pattern evoked by the offset or onset of a movie? Thus far,

we tested boundary responses following offsets, based on prior findings that post-stimulus

neural responses contribute to memory formation (Ben-Yakov et al., 2013; Ben-Yakov & Dudai,

2011; Medvedeva et al., 2021). However, other studies reported neural responses specific to

the onset of an episode (Bulkin et al., 2020; Fox et al., 2005; Wen et al., 2020). We computed

between-phase (encoding-recall) pattern similarity across individual time points, time-locked to

either the onset or offset of each watched or recalled movie (Figure 3D). We found that

significantly positive between-phase correlations emerged before the encoding and recall

onsets, starting from 4.5 seconds following the offsets of the preceding movie. Thus, it is likely

that boundary patterns were driven primarily by offset responses.

Figure 3. Boundary pattern in the posterior medial cortex (PMC). (A) PMC activation pattern similarity (Pearson correlation) between the 10 movie stimuli (M1 – 10), conditions (Offset = boundary, Middle = non-boundary), and experimental phases (Encoding, Recall), averaged across all subjects. The boundary pattern of a movie was defined as the mean pattern averaged across the 15-second window following the offset of the movie. The non-boundary pattern was defined as the mean pattern averaged across the 15-second window in the middle of a movie. The time windows for both boundary and non-boundary patterns were shifted forward by 4.5 seconds to account for the hemodynamic response delay. PMC regions of interest are shown as white areas on the inflated surface of a template brain. (B) Subject-specific mean activation patterns associated with between-movie boundaries during encoding (left) and recall (right). The boundary patterns were averaged across all movies and then z-scored across vertices within the PMC ROI mask, separately for each experimental phase. PMC (demarcated by black outlines) of four example subjects (S1 – 4) are shown on the medial surface of the right hemisphere of the fsaverage6 template brain. (C) Within-phase (Recall-Recall) and between-phase (Encoding-Recall) pattern similarity across different movies, computed separately for the boundary (Offset) and non-boundary (Middle) patterns in PMC. Bar graphs show the mean across subjects. Circles represent individual subjects. Error bars show SEM across subjects. ***p < .001. (D) Time-point-by-time-point PMC pattern similarity across the encoding phase and recall phase activation patterns around between-movie boundaries, averaged across all subjects. The time series of activation patterns were locked to either the onset (left) or the offset (right) of each movie. During encoding, the onset of a movie and the offset of the preceding movie were separated by a 6-second title scene. During recall, onsets and offsets of recalled movies were separated by, on average, a 9.3-second pause (boundaries concatenated across subjects, SD = 16.8 seconds). Note that in this figure, zero corresponds to the true stimulus/behavior time, with no shifting for hemodynamic response delay. Areas outlined by black lines indicate correlations significantly different from zero after multiple comparisons correction (Bonferroni corrected p < .05). Time-time correlations within each experimental phase can be found in Figure 3 supplement 3.

192
193
Figure 4. Comparisons between template patterns. We measured the similarity (Pearson correlation)
between the average between-movie boundary pattern during recall and average encoding patterns
associated with 1) between-movie boundaries (Movie offset), 2) silence (No sound), and 3) major within-
movie event boundaries (Event offset) in the posterior medial cortex (PMC; left panel) and the auditory
cortex (AUD; right panel). The regions of interest are shown as white areas on the inflated surface of a
template brain. Bar graphs show the mean across subjects. Circles represent individual subjects. Error bars
show SEM across subjects. ***$p < .001$ against zero.
201

We focused our analyses up to this point on transitions between movies because they

provided clear boundaries between mental contexts during recall. Yet, event boundaries in

naturalistic movie stimuli are often defined as transitions between scenes within a movie

(Baldassano et al., 2017; Chen et al., 2017; Zacks et al., 2010). In prior work, it has been shown

that for within-movie event boundaries, neural responses scale positively with human judgments

of the "strength" of scene transitions (Ben-Yakov & Henson, 2018). Thus, we hypothesized that

boundaries between movies (i.e., between mental contexts) would manifest as stronger

versions of within-movie boundaries. We correlated 1) the mean between-movie boundary

pattern during recall and 2) the mean within-movie event boundary pattern during encoding, in

PMC (Figure 4, 'Event offset' condition). Surprisingly, the two were negatively correlated ($t(14) =$

5.10, $p < .001$, Cohen's $d_z = 1.32$), suggesting that the between-movie boundary pattern may

reflect a cognitive state qualitatively different from the state elicited by event boundaries during

movie watching (e.g., attentional engagement; Song et al., 2021).

Is the generalized boundary pattern driven by shared low-level perceptual or motoric

factors rather than cognitive states? Visual features (i.e., black screen) or pauses in speech

cannot explain boundary-specific similarity between encoding and recall phases, because

boundary and non-boundary periods were identical in terms of visual input during recall and

speech generation during movie watching. Moreover, PMC boundary patterns showed positive

between-phase pattern correlations ($t(14)$ = 3.94, p = .003, Cohen's d_z = 1.25) greater than

those of non-boundary patterns ($t(14)$ = 3.22, p = .011, Cohen's d_z = 1.02) even when restricted

to boundaries without pauses between recalled movies. We also ruled out the possibility that

silence during movie title scenes and pauses at recall boundaries drove the generalized

boundary pattern in PMC; the recall boundary pattern was not correlated with the pattern

associated with silent periods during encoding ($t(14)$ = 1.93, p = .074, Cohen's d_z = .498),

whereas the auditory cortex showed a positive correlation between the two ($t(14)$ = 10.31, p

< .001, Cohen's d_z = 2.66) (Figure 4, 'No sound' condition). Likewise, the movies' audio

amplitudes modulated the time course of similarity between the recall boundary pattern and the

encoding data in the auditory cortex, but not in PMC (Figure 4 supplements 1, 2).

Our study demonstrates that internally-generated boundaries between mental contexts

during narrated memory recall evoke generalized neural activation patterns in core posterior-

medial areas of the DMN (Ritchey & Cooper, 2020). These cortical patterns were similar to

those observed at major transitions between externally-driven contexts (different audiovisual

movies), suggesting that they reflect a general cognitive state associated with mental context

transitions. Importantly, these between-context patterns were distinct from within-context event

boundary detection signals. We speculate that the boundary state may be a temporary "relay"

state that occurs when no one mental model wins the competition to receive full attentional

focus (e.g., attention lapse, off-focus, mind-blanking; deBettencourt et al., 2018; Mittner et al.,

2016; Mortaheb et al., 2021; Ward & Wegner, 2013) following the flushing of the prior mental

context. An alternative, but not mutually exclusive, possibility is that the boundary state involves

the recruitment of cognitive control to resolve the competition between mental contexts; the

relative activation of the frontoparietal control network interdigitated with the DMN (Braga &

Buckner, 2017) may contribute to the generalized boundary pattern.

The current study focused on the boundaries between narratives involving multiple

entities and their interactions, and thus the transition may coincide with the reconfiguration of

situation models represented in the DMN (Ranganath & Ritchey, 2012). Future studies may

investigate whether the boundary pattern we observed is associated specifically with

detachment from situation-level information accumulated over long timescales (Hasson et al.,

2015), or also with detachment from relatively simple and brief stimulus or task contexts (e.g.,

Crittenden et al., 2015; Kragel & Polyn, 2015; Schapiro et al., 2013). Our results also lay the

groundwork for future studies to examine the possible utility of these "relay" states in

comprehension, learning, and building durable memories.

Materials and methods

Here, we provide a selective overview of procedures and analysis methods. More detailed

descriptions of participants, stimuli, experimental procedures, fMRI data acquisition and

preprocessing can be found in Lee & Chen (2021).

Participants

Twenty-one subjects (12 females) between the ages of 20 and 33 participated in the study.

Informed consent was obtained in accordance with procedures approved by the Princeton

University Institutional Review Board. Six subjects were excluded from analyses due to

excessive motion.

Stimuli

Ten audiovisual movies (ranged 2.15 – 7.75 minutes) were used in the experiment. The movies

varied in format (animation, live-action) and content. Each movie clip was prepended with a title

scene where the movie title in white letters faded in and out at the center of the black screen.

The movie title was shown approximately for 3 seconds of the 6-second long title scene. At the

beginning of each scanning run, a 39-second long audiovisual introductory cartoon was played

before the movie stimuli. The introductory cartoon was excluded from analyses.

Experimental procedures

The experiment consisted of two phases, encoding and free spoken recall (Figure 1A), both

performed inside the MRI scanner. In the encoding phase, subjects watched a series of ten

short movies. Subjects were instructed to pay attention to the movies, and no behavioral

responses were required. There were two scanning runs, and subjects watched five movies in

each run. Stimulus presentation began 3 seconds after the first volume of each run. In the free

spoken recall phase, subjects were instructed to verbally recount what they remembered from

the movies, regardless of the order of presentation. Subjects were encouraged to describe their

memory in their own words in as much detail as possible. A white dot was presented in the

center of the black screen during the free spoken recall phase, though subjects were not

required to fixate. Subjects' recall speech was audio-recorded using an MR-compatible noise-

cancelling microphone and then manually transcribed. The recall transcripts were also

timestamped to identify the onset and offset of the description of each movie (there were no

intrusions across movies during recall).

fMRI acquisition and preprocessing

Imaging data were collected on a 3T Siemens Prisma scanner at Princeton Neuroscience

Institute. Functional images were acquired using a T2*- weighted multiband accelerated echo-

planar imaging sequence (TR = 1.5 s; TE = 39 ms; flip angle = 50°; acceleration factor = 4; 60

slices; 2 × 2 × 2 mm^3). Whole-brain anatomical images and fieldmap images were also

acquired. Functional images were motion-corrected and unwarped using FSL, and then

coregistered to the anatomical image, resampled to the fsaverage6 cortical surface, and

smoothed (FWHM 4 mm) using FreeSurfer Functional Analysis Stream. The smoothed data

were also high-pass filtered (cutoff = 140 s) and z-scored within each scanning run. The first 5

volumes of encoding scanning runs and the first 3 volumes of free spoken recall scanning runs

were excluded from analyses.

Cortical parcellation and region of interest (ROI) definition

For whole-brain pattern similarity analysis, we used an atlas (Schaefer et al., 2018) which

divided the cortical surface into 400 parcels (200 parcels per hemisphere) based on functional

connectivity patterns (17 networks version). For region-of-interest analyses, we defined the

bilateral posterior-medial cortex (PMC) by combining the parcels corresponding to the

precuneus and posterior cingulate cortex within Default Network A. The bilateral auditory cortex

ROI was defined by combining the parcels corresponding to the primary and secondary auditory

cortices within Somatomotor Network B. The bilateral angular gyrus ROI consisted of the

parcels corresponding to the inferior parietal cortex within Default Network A, B and C.

Univariate activation analysis

We performed whole-brain univariate activation analysis to identify brain areas that show

activation changes at between-movie boundaries compared to non-boundary periods during

recall (Figure 1B). The boundary periods were the first 15 seconds following the offset of each

recalled movie, and the non-boundary periods were the 15 seconds in the middle of each

recalled movie. Both boundary and non-boundary period time windows were shifted forward by

4.5 seconds to account for the hemodynamic response delay. For each vertex in each subject's

brain, we computed the mean boundary activation by first averaging preprocessed BOLD

signals across time points within each boundary period, and then across all recalled movies.

Likewise, we computed the mean non-boundary activation for each subject and vertex by first

averaging preprocessed BOLD signals across time points within each non-boundary period, and
then across all recalled movies. We then computed the difference between the boundary and
non-boundary activation for each subject. Finally, we performed a group-level one-sample *t*-test
against zero (two-tailed). Bonferroni correction was applied to correct for multiple comparisons
across vertices on the resulting whole-brain statistical parametric map.
326
Pattern similarity analysis
We performed whole-brain pattern similarity analysis (Figure 2A) to identify brain areas that
showed content-and task-general neural activation patterns associated with between-movie
boundaries. For each cortical parcel of each subject's brain, we extracted boundary and non-
boundary activation patterns for each movie, separately for the encoding phase and the recall
phase. Boundary patterns were generated by averaging the spatial patterns of activation within
the boundary period (the first 15 seconds following the offset) of each watched or recalled
movie. Non-boundary patterns were generated by averaging spatial patterns within the non-
boundary period (the middle 15 seconds) of each watched or recalled movie. Again, both
boundary and non-boundary time windows were shifted forward by 4.5 seconds to account for
the hemodynamic response delay. We then computed Pearson correlation coefficients between
the patterns within and across different movies, conditions (boundary, non-boundary), and
experimental phases.
Using the resulting correlation matrix (see Figure 3A for an example) for each parcel, we
first identified brain areas that showed boundary patterns which were consistent *across recalled*
movies (Figure 2B). For each subject's recall phase, we computed the mean of all pairwise
between-movie correlations, separately for the boundary patterns and the non-boundary
patterns. We then performed a group-level two-tailed one-sample *t*-test against zero on the
mean boundary pattern correlations to test whether the boundary pattern similarity was overall
positive. We also performed a group-level two-tailed paired-samples *t*-test between the mean
boundary vs. non-boundary pattern correlations to test whether the boundary pattern similarity
was greater than the non-boundary pattern similarity. Each of the resulting whole-brain
statistical parametric maps was corrected for multiple comparisons across parcels using the
Bonferroni method. Finally, we identified parcels that showed significant effects in both tests
after the correction (Figure 2B). Likewise, we identified brain areas that showed boundary
patterns which were consistent *across the encoding and recall phases* as well as across movies
(Figure 2C). This was achieved by repeating the identical analysis procedures using the
boundary and non-boundary pattern correlations computed across the encoding and recall

phases, instead of using the correlations computed within the recall phase.

We also performed the same pattern similarity analysis in the PMC (Figure 3) and

angular gyrus (Figure 3 supplement 2) ROIs, as done for an individual cortical parcel in the

whole-brain analysis.

Comparing the onset- and offset-locked boundary patterns

To test whether the consistent activation patterns associated with between-movie boundaries

were evoked by the onset or offset of a movie, we examined TR-by-TR pattern correlations

across time points around the boundaries. The time points were locked to either the onset or the

offset of 1) each video clip (excluding the title scene) or 2) recall of each movie. For each

subject and ROI, we extracted the time series of activation patterns from 30 seconds before to

60 seconds after the onset/offset of each watched or recalled movie. We averaged the time

series across movies to create a single time series of boundary-related activation patterns per

experimental phase. We then computed Pearson correlation coefficients across different time

points in the time series of mean activation patterns within each experimental phase (i.e.,

encoding-encoding and recall-recall correlation; Figure 3 supplement 3) or between phases (i.e.,

encoding-recall correlation; Figure 3D, Figure 3 supplement 2D). Finally, we performed two-

tailed one-sample t-tests against zero on each cell of the time-time correlation matrices from all

subjects to identify the time points at which positive pattern correlations appeared. Bonferroni

correction was applied to correct for multiple comparisons across all cells in the time-time

correlation matrix.

Comparing the between-movie and within-movie boundary patterns

To test whether the recall activation patterns evoked by between-movie boundaries were similar

to encoding activation patterns evoked by event boundaries within a movie (Figure 4, 'Event

offset' condition), we identified the strongest event boundaries within each movie. We utilized

the fine-grained event boundaries defined in our previous study (Lee & Chen, 2021) which

divided the ten movie stimuli into 202 events excluding title scenes (mean duration = 13.5

seconds, ranged 2 – 42 seconds). We had four independent coders watch the movie stimuli and

then choose which of the fine-grained event boundaries were the most important. The coders

were instructed to select the boundaries such that the ten movies were divided into 60 ± 10

events excluding title scenes. Of these, 25 event boundaries were identified as important by all

four coders, which resulted in 27 "coarse" events in total (ranging between one and five events

per movie; mean duration = 100.9 seconds, ranged 21 – 417 seconds). To mitigate the

possibility of carryover effects from the between-movie boundaries, within-movie event

boundaries that occurred within the first 45 seconds of each movie clip were excluded from the

analysis, leaving 15 within-movie event boundaries in total. For each subject, we generated the

mean within-movie event boundary pattern of an ROI by averaging activation patterns from 4.5

to 19.5 seconds following each of the 15 event boundaries during encoding. Likewise, the mean

between-movie boundary pattern was generated by averaging all activation patterns from 4.5 to

19.5 seconds following the offset of each movie during recall. We then computed a Pearson

correlation coefficient across the mean within-movie event boundary pattern and the mean

between-movie boundary pattern. A two-tailed one-sample *t*-test against zero was performed to

test whether the group-level similarity between the two patterns was positive.

Testing the effect of audio amplitudes

Brief periods of silence were present at transitions between movies during both encoding and

recall. During encoding, the 6-second title period between movies was silent. During recall,

subjects often paused speaking for several seconds between recall of different movies. We

tested whether the between-movie boundary patterns were associated with the absence of

sound in general, as opposed to between-movie transitions specifically.

We first compared the activation pattern associated with any silent periods within movies

during encoding and the activation pattern evoked by between-movie boundaries during recall

(Figure 4, 'No sound' condition). To identify all periods of silence within the movies, we

extracted the audio amplitudes of the movie clips (Figure 4 supplement 1A) by applying a Hilbert

transform to the single-channel audio signals (44.1 kHz). The audio amplitudes were

downsampled to match the temporal resolution of fMRI data (TR = 1.5 s), convolved with a

double-gamma hemodynamic response function, and *z*-scored across time points. The periods

of silence were defined as the within-movie time points (again excluding the first 45 seconds of

each movie) whose audio amplitudes were equal to or lower than the mean amplitude of the

time points corresponding to the silent between-movie title periods. For each subject and ROI,

we averaged the activation patterns across these within-movie silent time periods to produce

the mean activation pattern associated with the absence of sound. The mean pattern was then

correlated with the between-movie boundary pattern produced by averaging 4.5 – 19.5 seconds

following the offset of each movie during recall. A two-tailed one-sample *t*-test was performed to

compare the group-level correlation coefficients against zero.

We additionally tested whether the time course of audio amplitude was correlated with

the time course of pattern similarity (Pearson correlation) between the recall phase between-

movie boundary pattern and each time point of the encoding phase data (Figure 4 supplements

1B, 1C, 2). The time courses were generated for all time points within each movie, excluding the

first 45 seconds of each movie. We first computed each subject's Pearson correlation coefficient

between the two types of time courses. We then performed a group-level one-sample t-test

against zero (two-tailed).

Acknowledgements

We thank Sarah DuBrow, Christopher J. Honey, and Megan T. deBettencourt for scientific

discussions and comments on earlier versions of the manuscript. This work was supported by

the Sloan Research Fellowship (J. C.) and Google Faculty Research Award (J. C.).

Competing interests

The authors declare no competing interests.

Citation diversity statement

Recent work in several fields of science has identified a bias in citation practices such that

papers from women and other minority scholars are under-cited relative to the number of such

papers in the field (Caplar et al., 2017; Dion et al., 2018; Dworkin et al., 2020; Maliniak et al.,

2013; Mitchell et al., 2013). Here we sought to proactively consider choosing references that

reflect the diversity of the field in thought, form of contribution, gender, race, ethnicity, and other

factors. First, we obtained the predicted gender of the first and last author of each reference by

using databases that store the probability of a first name being carried by a woman (Dworkin et

al., 2020; Zhou et al., 2020). By this measure (and excluding self-citations to the first and last

authors of our current paper), our references contain 13.64% woman(first)/woman(last), 15.91%

man/woman, 25.87% woman/man, and 44.59% man/man. This method is limited in that a)

names, pronouns, and social media profiles used to construct the databases may not, in every

case, be indicative of gender identity and b) it cannot account for intersex, non-binary, or

transgender people. Second, we obtained predicted racial/ethnic category of the first and last

author of each reference by databases that store the probability of a first and last name being

carried by an author of color (Ambekar et al., 2009; Sood & Laohaprapanon, 2018). By this

measure (and excluding self-citations), our references contain 3.86% author of color

(first)/author of color(last), 15.58% white author/author of color, 23.87% author of color/white

author, and 56.69% white author/white author. This method is limited in that a) names, Census

entries, and Wikipedia profiles used to make the predictions may not be indicative of

racial/ethnic identity, and b) it cannot account for Indigenous and mixed-race authors, or those

who may face differential biases due to the ambiguous racialization or ethnicization of their

names. We look forward to future work that could help us to better understand how to support

equitable practices in science.

References

Ambekar, A., Ward, C., Mohammed, J., Male, S., & Skiena, S. (2009). Name-ethnicity
classification from open sources. *Proceedings of the 15th ACM SIGKDD International*
Conference on Knowledge Discovery and Data Mining, 49–58.
https://doi.org/10.1145/1557019.1557032
Baldassano, C., Chen, J., Zadbood, A., Pillow, J. W., Hasson, U., & Norman, K. A. (2017).
Discovering event structure in continuous narrative perception and memory. *Neuron*,
95(3), 709-721.e5. https://doi.org/10.1016/j.neuron.2017.06.041
Ben-Yakov, A., & Dudai, Y. (2011). Constructing realistic engrams: Poststimulus activity of
hippocampus and dorsal striatum predicts subsequent episodic memory. *Journal of*
Neuroscience, *31*(24), 9032–9042. https://doi.org/10.1523/JNEUROSCI.0702-11.2011
Ben-Yakov, A., Eshel, N., & Dudai, Y. (2013). Hippocampal immediate poststimulus activity in
the encoding of consecutive naturalistic episodes. *Journal of Experimental Psychology:*
General, *142*(4), 1255–1263. https://doi.org/10.1037/a0033558
Ben-Yakov, A., & Henson, R. N. (2018). The hippocampal film editor: Sensitivity and specificity
to event boundaries in continuous experience. *Journal of Neuroscience*, *38*(47), 10057–
10068. https://doi.org/10.1523/JNEUROSCI.0524-18.2018
Braga, R. M., & Buckner, R. L. (2017). Parallel interdigitated distributed networks within the
individual estimated by intrinsic functional connectivity. *Neuron*, *95*(2), 457-471.e5.
https://doi.org/10.1016/j.neuron.2017.06.038
Brunec, I. K., Moscovitch, M., & Barense, M. D. (2018). Boundaries shape cognitive
representations of spaces and events. *Trends in Cognitive Sciences*, *22*(7), 637–650.
https://doi.org/10.1016/j.tics.2018.03.013
Buckner, R. L., & DiNicola, L. M. (2019). The brain's default network: Updated anatomy,
physiology and evolving insights. *Nature Reviews Neuroscience*, *20*(10), 593–608.
https://doi.org/10.1038/s41583-019-0212-7
Bulkin, D. A., Sinclair, D. G., Law, L. M., & Smith, D. M. (2020). Hippocampal state transitions at
the boundaries between trial epochs. *Hippocampus*, *30*(6), 582–595.
https://doi.org/10.1002/hipo.23180
Caplar, N., Tacchella, S., & Birrer, S. (2017). Quantitative evaluation of gender bias in
astronomical publications from citation counts. *Nature Astronomy*, *1*(6), 1–5.
https://doi.org/10.1038/s41550-017-0141

Chen, J., Leong, Y. C., Honey, C. J., Yong, C. H., Norman, K. A., & Hasson, U. (2017). Shared
memories reveal shared structure in neural activity across individuals. *Nature*
Neuroscience, *20*(1), 115–125. https://doi.org/10.1038/nn.4450
Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-
wandering as spontaneous thought: A dynamic framework. *Nature Reviews*
Neuroscience, *17*(11), 718–731. https://doi.org/10.1038/nrn.2016.113
Clewett, D., DuBrow, S., & Davachi, L. (2019). Transcending time in the brain: How event
memories are constructed from experience. *Hippocampus*, *29*(3), 162–183.
https://doi.org/10.1002/hipo.23074
Crittenden, B. M., Mitchell, D. J., & Duncan, J. (2015). Recruitment of the default mode network
during a demanding act of executive control. *ELife*, *4*, e06481.
https://doi.org/10.7554/eLife.06481
deBettencourt, M. T., Norman, K. A., & Turk-Browne, N. B. (2018). Forgetting from lapses of
sustained attention. *Psychonomic Bulletin & Review*, *25*(2), 605–611.
https://doi.org/10.3758/s13423-017-1309-5
Dion, M. L., Sumner, J. L., & Mitchell, S. M. (2018). Gendered citation patterns across political
science and social science methodology fields. *Political Analysis*, *26*(3), 312–327.
https://doi.org/10.1017/pan.2018.12
DuBrow, S., Rouhani, N., Niv, Y., & Norman, K. A. (2017). Does mental context drift or shift?
Current Opinion in Behavioral Sciences, *17*, 141–146.
https://doi.org/10.1016/j.cobeha.2017.08.003
Dworkin, J. D., Linn, K. A., Teich, E. G., Zurn, P., Shinohara, R. T., & Bassett, D. S. (2020). The
extent and drivers of gender imbalance in neuroscience reference lists. *Nature*
Neuroscience, *23*(8), 918–926. https://doi.org/10.1038/s41593-020-0658-y
Fox, M. D., Snyder, A. Z., Barch, D. M., Gusnard, D. A., & Raichle, M. E. (2005). Transient
BOLD responses at block transitions. *NeuroImage*, *28*(4), 956–966.
https://doi.org/10.1016/j.neuroimage.2005.06.025
Hasselmo, M. E. (1995). Neuromodulation and cortical function: modeling the physiological
basis of behavior. *Behavioural Brain Research*, *67*(1), 1–27.
https://doi.org/10.1016/0166-4328(94)00113-T
Hasson, U., Chen, J., & Honey, C. J. (2015). Hierarchical process memory: memory as an
integral component of information processing. *Trends in Cognitive Sciences*, *19*(6), 304–
313.

Honey, C. J., Newman, E. L., & Schapiro, A. C. (2017). Switching between internal and external

modes: A multiscale learning principle. *Network Neuroscience*, *1*(4), 339–356.

https://doi.org/10.1162/NETN_a_00024

Karapanagiotidis, T., Vidaurre, D., Quinn, A. J., Vatansever, D., Poerio, G. L., Turnbull, A., Ho,

N. S. P., Leech, R., Bernhardt, B. C., Jefferies, E., Margulies, D. S., Nichols, T. E.,

Woolrich, M. W., & Smallwood, J. (2020). The psychological correlates of distinct neural

states occurring during wakeful rest. *Scientific Reports*, *10*(1), 21121.

https://doi.org/10.1038/s41598-020-77336-z

Kragel, J. E., & Polyn, S. M. (2015). Functional interactions between large-scale networks

during memory search. *Cerebral Cortex*, *25*(3), 667–679.

https://doi.org/10.1093/cercor/bht258

Lee, H., & Chen, J. (2021). Narratives as networks: Predicting memory from the structure of

naturalistic events. *BioRxiv*, 2021.04.24.441287.

https://doi.org/10.1101/2021.04.24.441287

Maliniak, D., Powers, R., & Walter, B. F. (2013). The gender citation gap in international

relations. *International Organization*, *67*(4), 889–922.

https://doi.org/10.1017/S0020818313000209

Manning, J. R., Hulbert, J. C., Williams, J., Piloto, L., Sahakyan, L., & Norman, K. A. (2016). A

neural signature of contextually mediated intentional forgetting. *Psychonomic Bulletin &*

Review, *23*(5), 1534–1542. https://doi.org/10.3758/s13423-016-1024-7

Medvedeva, A., Saw, R., Silvestri, C., Sirota, M., Fuggetta, G., & Galli, G. (2021). Offset-related

brain activity in the left ventrolateral prefrontal cortex promotes long-term memory

formation of verbal events. *Brain Stimulation*, *14*(3), 564–570.

https://doi.org/10.1016/j.brs.2021.03.002

Mildner, J. N., & Tamir, D. I. (2019). Spontaneous thought as an unconstrained memory

process. *Trends in Neurosciences*, *42*(11), 763–777.

https://doi.org/10.1016/j.tins.2019.09.001

Mitchell, S. M., Lange, S., & Brus, H. (2013). Gendered citation patterns in international

relations journals. *International Studies Perspectives*, *14*(4), 485–492.

https://doi.org/10.1111/insp.12026

Mittner, M., Hawkins, G. E., Boekel, W., & Forstmann, B. U. (2016). A neural model of mind

wandering. *Trends in Cognitive Sciences*, *20*(8), 570–578.

https://doi.org/10.1016/j.tics.2016.06.004

Mortaheb, S., Klados, M. A., Calster, L. V., Boulakis, P. A., Georgoula, K., Majerus, S., &

Demertzi, A. (2021). Mind blanking is associated with a rigid spatio-temporal profile in

typical wakefulness. *BioRxiv*, 2021.05.10.443428.

https://doi.org/10.1101/2021.05.10.443428

Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour.

Nature Reviews Neuroscience, *13*(10), 713–726. https://doi.org/10.1038/nrn3338

Reagh, Z. M., Delarazan, A. I., Garber, A., & Ranganath, C. (2020). Aging alters neural activity

at event boundaries in the hippocampus and Posterior Medial network. *Nature*

Communications, *11*(1), 3980. https://doi.org/10.1038/s41467-020-17713-4

Ritchey, M., & Cooper, R. A. (2020). Deconstructing the posterior medial episodic network.

Trends in Cognitive Sciences, *24*(6), 451–465. https://doi.org/10.1016/j.tics.2020.03.006

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S.

B., & Yeo, B. T. T. (2018). Local-global parcellation of the human cerebral cortex from

intrinsic functional connectivity MRI. *Cerebral Cortex*, *28*(9), 3095–3114.

https://doi.org/10.1093/cercor/bhx179

Schapiro, A. C., Rogers, T. T., Cordova, N. I., Turk-Browne, N. B., & Botvinick, M. M. (2013).

Neural representations of events arise from temporal community structure. *Nature*

Neuroscience, *16*(4), 486–492. https://doi.org/10.1038/nn.3331

Shin, Y. S., & DuBrow, S. (2021). Structuring memory through inference-based event

segmentation. *Topics in Cognitive Science*, *13*(1), 106–127.

https://doi.org/10.1111/tops.12505

Smallwood, J., & Schooler, J. W. (2015). The science of mind wandering: Empirically navigating

the stream of consciousness. *Annual Review of Psychology*, *66*(1), 487–518.

https://doi.org/10.1146/annurev-psych-010814-015331

Smith, V., Mitchell, D. J., & Duncan, J. (2018). Role of the default mode network in cognitive

transitions. *Cerebral Cortex*, *28*(10), 3685–3696. https://doi.org/10.1093/cercor/bhy167

Song, H., Finn, E. S., & Rosenberg, M. D. (2021). Neural signatures of attentional engagement

during narratives and its consequences for event memory. *Proceedings of the National*

Academy of Sciences, *118*(33). https://doi.org/10.1073/pnas.2021905118

Sood, G., & Laohaprapanon, S. (2018). Predicting race and ethnicity from the sequence of

characters in a name. *ArXiv:1805.02109 [Stat]*. http://arxiv.org/abs/1805.02109

Speer, N. K., Zacks, J. M., & Reynolds, J. R. (2007). Human brain activity time-locked to

narrative event boundaries. *Psychological Science*, *18*(5), 449–455.

https://doi.org/10.1111/j.1467-9280.2007.01920.x

Sripada, C., & Taxali, A. (2020). Structure in the stream of consciousness: Evidence from a
verbalized thought protocol and automated text analytic methods. *Consciousness and*
Cognition, *85*, 103007. https://doi.org/10.1016/j.concog.2020.103007

Tseng, J., & Poppenk, J. (2020). Brain meta-state transitions demarcate thoughts across task
contexts exposing the mental noise of trait neuroticism. *Nature Communications*, *11*(1),
3480. https://doi.org/10.1038/s41467-020-17255-9

Ward, A. F., & Wegner, D. M. (2013). Mind-blanking: when the mind goes away. *Frontiers in*
Psychology, *0*. https://doi.org/10.3389/fpsyg.2013.00650

Wen, T., Duncan, J., & Mitchell, D. J. (2020). Hierarchical representation of multistep tasks in
multiple-demand and default mode networks. *Journal of Neuroscience*, *40*(40), 7724–
7738. https://doi.org/10.1523/JNEUROSCI.0594-20.2020

Zacks, J. M. (2020). Event perception and memory. *Annual Review of Psychology*, *71*(1), 165–
191. https://doi.org/10.1146/annurev-psych-010419-051101

Zacks, J. M., Braver, T. S., Sheridan, M. A., Donaldson, D. I., Snyder, A. Z., Ollinger, J. M.,
Buckner, R. L., & Raichle, M. E. (2001). Human brain activity time-locked to perceptual
event boundaries. *Nature Neuroscience*, *4*(6), 651–655. https://doi.org/10.1038/88486

Zacks, J. M., Kurby, C. A., Eisenberg, M. L., & Haroutunian, N. (2011). Prediction error
associated with the perceptual segmentation of naturalistic events. *Journal of Cognitive*
Neuroscience, *23*(12), 4057–4066. https://doi.org/10.1162/jocn_a_00078

Zacks, J. M., Speer, N. K., Swallow, K. M., Braver, T. S., & Reynolds, J. R. (2007). Event
perception: A mind-brain perspective. *Psychological Bulletin*, *133*(2), 273–293.
https://doi.org/10.1037/0033-2909.133.2.273

Zacks, J. M., Speer, N. K., Swallow, K. M., & Maley, C. J. (2010). The brain's cutting-room floor:
Segmentation of narrative cinema. *Frontiers in Human Neuroscience*, *0*.
https://doi.org/10.3389/fnhum.2010.00168

Zheng, J., Schjetnan, A. G. P., Yebra, M., Mosher, C., Kalia, S., Valiante, T. A., Mamelak, A. N.,
Kreiman, G., & Rutishauser, U. (2021). Cognitive boundary signals in the human medial
temporal lobe shape episodic memory representation. *BioRxiv*, 2021.01.16.426538.
https://doi.org/10.1101/2021.01.16.426538

Zhou, D., Cornblath, E. J., Stiso, J., Teich, E. G., Dworkin, J. D., Blevins, A. S., & Bassett, D. S.
(2020). *Gender diversity statement and code notebook v1.0*. Zenodo.
https://doi.org/10.5281/zenodo.3672110

625