Dysfunction of an energy sensor NFE2L1 triggers uncontrollable AMPK signal and glucose metabolism reprogramming

Qiufang Yang[1,@], Wenshan Zhao[1,@], Yadi Xing[2,@], Peng Li[1], Xiaowen Zhou[1], Haoming Ning[1], Ranran Shi[1], Shanshan Gou[1], Yalan Chen[1], Wenjie Zhai[1], Yahong Wu[1], Guodong Li[1], Zhenzhen Chen[1], Yonggang Ren[3], Yanfeng Gao[1,4], Yiguo Zhang[5], Yuanming Qi[1,6,7,*], Lu Qiu[1,5,*]

[1] School of Life Sciences, Zhengzhou University, Zhengzhou 450001, China; [2] School of Agricultural Sciences, Zhengzhou University, Zhengzhou 450001, China; [3] Department of Biochemistry, North Sichuan Medical College, Nanchong 637000, China; [4] School of Pharmaceutical Sciences (Shenzhen), Sun Yat-sen University, Shenzhen 518107, China; [5] The Laboratory of Cell Biochemistry and Topogenetic Regulation, College of Bioengineering and Faculty of Sciences, Chongqing University, Chongqing 400044, China; [6] Henan Key Laboratory of Bioactive Macromolecules, Zhengzhou University, Zhengzhou 450001, China; [7] International Joint Laboratory for Protein and Peptide Drugs of Henan Province, Zhengzhou University, Zhengzhou 450001, China.

[@] Qiufang Yang, Wenshan Zhao and Yadi Xing have contributed equally to this work.

*Correspondence to Dr. Lu Qiu (qiulu@zzu.edu.cn) or Prof. Yuanming Qi (qym@zzu.edu.cn).

School of Life Sciences, Zhengzhou University, Zhengzhou 450001, China.

Tel.: +86 371 67783235; fax: +86 371 67783235.

Abstract

NFE2L1 (also called Nrf1) acts a core regulator of redox signaling and metabolism homeostasis, and thus its dysfunction results in multiple systemic metabolic diseases. However, the molecular mechanism(s) by which NFE2L1 regulates glycose and lipid metabolism is still elusive. Here, we found that the loss of NFE2L1 in human HepG2 cells led to a lethal phenotype upon glucose deprivation. The uptake of glucose was also affected by NFE2L1 deficiency. Further experiments unveiled that although the glycosylation of NFE2L1 was monitored through the glycolysis pathway, it enabled to sense the energy state and directly interacted with AMPK. These indicate that NFE2L1 can serve as a dual sensor and regulator of glucose homeostasis. In-depth sights into transcriptome, metabolome and seahorse data further unraveled that glucose metabolism was reprogrammed by disruption of NFE2L1, so as to aggravate the Warburg effect in NFE2L1-silenced hepatoma cells, along with the mitochondrial damage observed under the electron microscope. Collectively, these demonstrate that disfunction of NFE2L1 triggers the uncontrollable signaling by AMPK towards glucose metabolism reprogramming in the liver cancer development.

Keywords: NFE2L1/Nrf1; AMPK; LKB1; Reactive Oxygen Species/ROS; Glucose Metabolism; Mitochondria Function

1. Introduction

Redox homeostasis is a necessary prerequisite for maintenance of physiological responses, and the imbalance of redox homeostasis leads to various chronic systemic diseases. Nuclear factor erythroid 2 like 1 (NFE2L1/Nrf1), a core member of the cap'n'collar basic-region leucine zipper (CNC-bZIP) family, plays a critical role in regulating redox homeostasis in eukaryotic cells. Conventional knockout of *NFE2L1* could induce strong oxidative stress injury and result in fetal death. Also, conditional knockout of *NFE2L1* can increase cellular reactive oxygen species (ROS) level [1-4].

Studies have revealed that NFE2L1 plays an important role in metabolic pathway

and influences the development of metabolic diseases. Human GWAS study shows

that the single nucleotide polymorphism re3764400 which located in the 5'-flanking

regions of *NFE2L1* gene is associated with obesity [5]. Overexpression of *NFE2L1*

led to weight loss in transgenic mice, accompanied by insulin resistance symptoms

such as increased pancreatic islets, increased insulin secretion, and increased blood

glucose levels, eventually leading to diabetes mellitus [6]. In mouse pancreatic β cells,

specific knockout of NFE2L1 caused hyperinsulinemia and glucose intolerance, the

early symptoms of type II diabetes [7]. In adipocytes, specific knockout of *NFE2L1*

gene resulted in almost complete disappearance of subcutaneous adipose tissue in

mice, accompanied by insulin resistance, adipocyte hypertrophy and other

obesity-related Inflammation [8]. And knockout of *NFE2L1* gene in the liver quickly

caused non-alcoholic steatohepatitis (NASH) in mice [2,9]. All these results indicated

that *NFE2L1* gene is essential for maintaining the homeostasis of lipid/carbohydrate

metabolism.

Interestingly, research has shown that glycosylation of NFE2L1 protein mediates

the cleavage and nuclear entry of NFE2L1 [10]. O-glycosylation modification in the

Neh6L domain of NFE2L1 can reduce the transcription activity of NFE2L1 protein

[11]. Additionally, in HepG2 cells, deletion of NFE2L1 can cause glucose deprivation,

which finally induces cell death [12]. These studies suggest that NFE2L1 may be a

key factor for regulating glucose metabolism homeostasis by sensing glucose levels.

Consistently, our previous research also showed that loss of NFE2L1 expression can

disrupt AMP-activated protein kinase (AMPK) signaling pathway [13], the core

pathway involved in regulating energy metabolism [14,15]. In addition, metformin

(MET), an AMPK signaling activator and the first-line drug for the treatment of

diabetes, could inhibit NFE2L1 in an AMPK-independent manner, while the NFE2L1

could disruptive the activation of AMPK signal by MET [13], indicating the upstream

regulation effects of NFE2L1 on AMPK signal.

Previous studies have suggested that NFE2L1 is essential for maintaining the

homeostasis of glucose and lipid metabolism. However, the molecular mechanism is

still obscure. Here, we investigated the effects of NFE2L1 deficiency on the glucose

metabolism process in HepG2 cells and the results showed the dual functions of

NFE2L1 as the sensor of glucose level and the regulator of glucose metabolism. At

the same time, we also identified the mechanism of NFE2L1 in regulating glucose

metabolism homeostasis through the verification of the effects of NFE2L1 in

regulating AMPK signal.

2. Materials and Methods

2.1. Cell lines, culture and transfection

The information of the cell lines used in this study was shown in Table S1. Cells

were growing in DMEM supplemented with 10% (v/v) FBS. The experimental cells

were transfected by using Lipofectamine® 3000 Transfection Kit for 8 h, and then

allowed for recovery from transfection in the fresh medium. The information of kits

were shown in Table S1.

2.2. Expression constructs and other plasmids

The expression constructs for NFE2L1 and LKB1 were made by cloning the target

sequences from full-length CDS sequences of *NFE2L1* and *LKB1* into

pLVX-EGFP-puro vector. The primers used for these expression constructs were

shown in Table S1.

2.3. Establishment of lentiviral infected cell lines

Lentiviruses for infection of HepG2^{EGFP} and HepG2^{LKB1} cells were packaged in

HEK293T cells. HEK293T cells (5×10^5) were seeded in 6-well plates and were

transfected with three plasmids (1 μg of *pMD2.G*, 2 μg of *psPAX2*, 3 μg of either

pEGFP or *pLKB1::EGFP* in 1mL of transfection volume) when cell confluence

reached to 80%. The medium was collected and filtered by using 0.45 μm sterile filter

to obtain the virus after 36~48 h. The virus was employed to transduce target cells,

and puromycin (100 μM) was used to screen out the infected cells.

2.4. Cell viability assay

Experimental cells seeded in 96-well plates were processed according to the

experimental design. The cytotoxic effects of indicated compound on experimental

cells were determined by Cell Counting Kit-8 (CCK8) [16]. And the absorbance at

450 nm was measured by a microplate reader (SpectraMax iD5, USA). The

information of kits were shown in Table S1.

2.5. Cellular ROS staining

Experimental cells were allowed for growing to reach an appropriate confluence in

6-well plates and then incubated in a serum-free medium containing 10 μM of

2',7'-Dichlorodihydrofluorescein diacetate (DCFH-DA) [17] at 37□ for 20 min.

Thereafter, the cells were washed three times with serum-free media, before the green

fluorescent images were achieved by microscopy.

2.6. Glucose uptake assay

The uptake of 2-Deoxy-2-[(7-nitro-2,1,3-benzoxadiazol-4-yl)amino]-D-glucose

(2-NBDG), which is a fluorescent glucose analog, is used to visualizing the uptake

capacity of glucose by living cells. Experimental cells were allowed for incubated in a

serum-free medium containing 20 μM of 2-NBDG at 37□ for 10 min. Thereafter, the

cells were washed three times with serum-free media, before the green fluorescent

images were achieved by microscopy.

2.7. RNA isolation and real-time quantitative PCR (qPCR)

Experimental cells were subjected to isolate total RNAs using the RNAsimple Kit.

Then, 1 μg total RNAs were added in a reverse-transcriptase reaction to generate the

first strand of cDNA, by using the RevertAid First Strand Synthesis Kit. The

synthesized cDNA served as the template for qPCR with distinct primers, which

occurred in the GoTaq® qPCR Master Mix. The mRNA expression level of β-actin

was selected as an optimal internal standard control. The primers used for qPCR and

the information of kits were shown in Table S1.

2.8. Western bloting (WB), Co-Immunoprecipitation (Co-IP) and protein

deglycosylation reactions

The total protein extraction buffer and WB were carried out as previously described

[13]. The Co-IP were carried out as previously described [18]. Before visualized by

WB, the deglycosylation reactions of samples with Endo H in vitro were carried out

as described previously [19]. The information of all antibodies used herein were

shown in Table S1.

2.9. Transcriptome sequencing

The cultured cells (HepG2^{shNC} and HepG2^{shNFE2L1}) were lysed by Trizol, and the

transcriptome sequencing by *Beijing Genomics Institute (BGI)* with the Order Number

is F21FTSCCKF2599_HOMdynrN. The samples were measured use the

BGISEQ-500 platform. The transcriptome sequencing results are shown in Table S2.

2.10. Metabolomics testing

The HepG2^{shNC} and HepG2^{shNFE2L1} cells were digested and counted, and the cell

pellets were obtained by low-speed centrifugation. The cell samples were

immediately frozen in liquid nitrogen. Metabolomics testing by *Metabolomics and*

Systems Biology Company, Germany. And the Order Number is POCMTS2016006.

The samples were measured with a Waters ACQUITY Reversed Phase Ultra

Performance Liquid Chromatography (RP-UPLC) coupled to a Thermo-Fisher

Exactive mass spectrometer. For each type of cells, three replicates were extracted and

measured.

2.11. Electron Microscopy

The adherent cells were digested with trypsin and fixed with 2.5% glutaraldehyde.

The ultrastructure of cells was examined by transmission electron microscope

(JEM-HT7700 model, Hitachi, Japan) with 5um, 1um, and 500nm scales,

respectively.

2.12. Seahorse Metabolic Analyzer

The oxygen consumption rate (OCR) of cells was measured by Seahorse XF Cell

Mito Stress Test Kit. The experimental operation was carried out according to the

instructions. The OCR were measured using the Seahorse Metabolic Analyzer to

characterize the cell's oxidative phosphorylation level. The information of Kits were

shown in Table S1.

2.13. Assays of ATP and Lactate Levels in cells

The ATP levels and Lactate levels of cells are determined according to the

instruction of ATP assay kit and Lactic Acid assay kit. The experimental operation

was carried out according to the instructions. The information of Kits were shown in

Table S1.

2.14. Statistical analysis

The data were here presented as a fold change (means ± S.D.), each of which

represents at least three independent experiments that performed in triplicate.

Significant differences were statistically determined by either the Student's t-test or

Multiple Analysis of Variations (MANOVA). The statistical significance was defined

by symbols '*' means $p < 0.05$, whilst the '*n.s.*' letters represent 'no significant'.

3. Results

3.1. NFE2L1 deficiency disrupts cellular energy metabolism signals and induces cell

death by glucose starvation

Previous studies have shown that mice with NFE2L1 overexpression or knockout

exhibit diabetes-like phenotypes [6-8]. Metformin has shown regulatory effects on

NFE2L1 [13]. These studies indicated the involvement of NFE2L1 in maintaining the

homeostasis of glucose metabolism. To further explore the relationship between

NFE2L1 and glucose metabolism, HepG2 cell line with *NFE2L1* knockdown was

constructed (Figure 1A). Through the glucose deprivation experiment, HepG2 cells in

the medium without glucose lacking *NFE2L1* expression were more susceptible to

death (Figure 1B,C). Glucose is the major source of energy for cells, but it is not an

essential nutrient. Here, WZB117 [20], a glucose uptake inhibitor, was used to test the

effects of impaired glucose uptake on cells. The results showed that compared with

the control group, the *NFE2L1* knockdown cells was exhibited enhanced killing

activity by treating with WZB117 (Figure 1D), which indicate that loss of NFE2L1

may disrupt the homeostasis of energy metabolism.

NFE2L1 is a regulator of redox homeostasis, and most of the phenotypes related

to NFE2L1 are related to oxidative stress. Results showed that the ROS level in the control group (HepG2[shNC]) was increased by about 2 times when the glucose was deprived for 6 hours. In *NFE2L1* knockdown cells (HepG2[shNFE2L1]), the level of ROS was increased by about 3 times, and the shrinking of cells has been observed (Figure 1E,F). In *NFE2L1* knockdown cells, the ROS level was increased more than 6 times in comparison with the control group after glucose deprivation for 6 hours, indicating the high levels of oxidative damage of *NFE2L1* knockdown cells. These results were in accordance with previous studies that glucose starvation resulted in strong oxidative stress to induce cell death after *NFE2L1* knockout [3,12].

AMPK, the core signal of energy metabolism, is the guardian of metabolism and mitochondrial homeostasis [21]. Western bloting (WB) results showed knockdown of *NFE2L1* caused a significant increase in the phosphorylation level of AMPK and its downstream protein acetyl-CoA carboxylase alpha (ACC) [22] (Figure 1G), indicating that the cells may be in a starvation state even in the sufficient nutritional conditions. This phenomenon may due to the lack of NFE2L1 in HepG2[shNFE2L1] cells. All these results suggested that NFE2L1 could function through sensing the energy state of the cells.

3.2. Glycolysis promotes glycation of NFE2L1 and lack of NFE2L1 promotes glucose absorption

In the cell culture system, in addition to glucose, other nutrients also contribute to provide the beneficial environment for cell growth, including cytokines and fetal bovine serum (FBS). Studies have shown that NFE2L1 could be activated by FBS/mechanistic target of rapamycin kinase (mTOR)/sterol regulatory element binding transcription factor 1 (SREBF1) pathway [23]. The mTOR and AMPK signaling pathways could be affected by NFE2L1 overexpression or knockdown [13], indicating the correlation of NFE2L1 with energy metabolism. Therefore, the alternation of NFE2L1 in response to the change of glucose and serum concentration was tested in the cell culture medium and the results showed that addition of FBS

could upregulate the total protein content of NFE2L1, and glucose could directly

affect the bands of NFE2L1 (Figure 2A,B). The level of p-AMPK gradually decreased

with the increase of serum concentration in the presence and absence of glucose

(Figure 2A). And the increased glucose concentration slightly altered the p-AMPK

level with and without FBS (Figure 2B). The p-mTOR level could be significantly

elevated with the presence of glucose (Figure 2A) and FBS (Figure 2B) despite the

concentration alternation of glucose and FBS.

Activation of AMPK and mTOR signals can be characterized by p-ACC and

phosphorylated- ribosomal protein S6 kinase B1 (p-S6K1) expression respectively. In

fact, as the results showed, p-AMPK was more sensitive to FBS in HepG2 cells

(Figure 2B), while p-ACC was more sensitive to glucose (Figure 2A). The expression

level of p-S6K1 increased at low concentrations of FBS and decreased when exposed

to higher concentrations of FBS (Figure 2A,B). With the increased concentration of

glucose, the level of p-S6K1 was upregulated (Figure 2A,B). In addition, the bands of

p-AMPK and p-S6K1 moved down with the increase of FBS, which might be due to

the fact that the protein synthesis upregulated in respond to the increased amount of

FBS, as the results showed by coomassie brilliant blue staining of protein samples in

Figure 2C and Figure 2D. It is important to note that the level of p-ACC has a strong

negative relationship with the glycation of NFE2L1 (Figure 2A,B). Combined with

the increase of p-ACC expression after NFE2L1 knockdown (Figure 1G) and the

inhibition of p-ACC by NFE2L1 overexpression, these results together suggested that

NFE2L1 might directly affect the catalytic activity of p-AMPK on substrates.

Studies have shown that glycosylation can regulate the activity of NFE2L1

[10,11,24,25]. Here, the effects of five common monosaccharides on NFE2L1 protein

were tested and the results indicated that fructose (FRU) and mannose (MAN), similar

as glucose (GLU), could induce the production of NFE2L1 with larger molecular

weight (Figure 2E), which could be ascribed to the glycosylation of NFE2L1 protein

as it revealed by the de-glycosylation experiment shown in Figure 2F. On the contrary,

galactose (GAL) and ribose (RIB) showed null effects on the molecular weight of

NFE2L1 protein (Figure 2F).

The catabolism of carbohydrates mainly undergoes anaerobic glycolysis in the

cytoplasm and aerobic oxidation in the mitochondria. In HepG2 cells, treatment with

2-Deoxy-D-glucose (2DG, a glycolysis inhibitor) [26] could effectively inhibit

glycolysis of NFE2L1. Inhibition of oxidative phosphorylation process using

Oligomycin (OM) [27] could decrease the total expression level of NFE2L1 (Figure

2G). Further results showed that fructose-1,6-bisphosphate (FBP), the intermediate

product of the glycolysis process, induced the glycation of NFE2L1, while the

pyruvate (PYR) had no such effects (Figure 2H). These findings implied that

activation of hexokinase (HK) and phosphofructokinase, liver (PFKL), the key

rate-limiting enzymes of glycolysis, or inhibition of pyruvate kinase M1/2 (PKM)

may have the function of activating NFE2L1.

The above results demonstrated that NFE2L1 was a glucose-sensitive protein. As

the results revealed in Figure 1, knockdown of NFE2L1 induced cell death by glucose

starvation and activation of AMPK signaling, indicating that NFE2L1 might be

involved in feedback regulation of energy metabolism. Our previous studies have

shown that NFE2L1 could be inhibited by metformin (MET), and knockdown of

NFE2L1 disrupted the activation of AMPK signal by MET. In addition, studies have

shown that MET promoted the uptake of glucose in HepG2 cells [28]. These studies

suggested the promotion of MET on glucose uptake in HepG2 cells might be related

to its inhibition of NFE2L1.

NFE2L1 deficiency improved the glucose uptake in HepG2 cells. Interestingly,

knockdown of NFE2L1 counteracted the promotion of MET on glucose uptake of

HepG2 cells (Figure 3A,B). Similarly, loss of NFE2L1 expression also neutralized the

ROS elevation induced by MET [13]. These data indicated that the inhibitory effect of

MET on NFE2L1 might be the main reason for the improvement of the glucose

uptake of HepG2 cells. However, it could not be ruled out that the glucose uptake

capacity of cells reached a threshold. Subsequently, the changes of glucose transporter

genes in transcriptome date were checked, and the data showed that the *solute carrier*

family 2 member 1 (*SLC2A1/GLUT1*) and *SLC2A13* were increased, while the *SLC2A3*, *SLC2A6* and *SLC2A8* were depressed after *NFE2L1* knockdown (Figure 3C). And the qPCR results showed that *SLC2A3* was significantly reduced, while *SLC2A4* and *SLC2A6* were obviously increased (Figure 3D). The *SLC2A2*, *SLC2A5*, *SLC2A7* and *SLC2A9* genes were not detected by real-time quantitative PCR (qPCR), which might be due to the low or lack of expression of these genes. By comparing the transcriptome and qPCR results, it was found that only *SLC2A3* was consistently reduced, but it was opposite of the phenotype. Then, the protein levels of glucose transporter were been detected and the results showed that the protein expression level of GLUT3 was significantly increased after NFE2L1 knockdown (Figure 3E), indicating that the increase in glucose uptake caused by NFE2L1 knockdown might be related to GLUT3. In addition, the mRNA level of *SLC2A1*, of which the protein level could hardly been detected, was highest, suggesting that GLUT1 might be severely inhibited at the post-transcriptional level in HepG2 cells.

3.3. NFE2L1 deficiency results in the reprogramming of glucose metabolism and exacerbates the Warburg effect

The above results revealed the function of NFE2L1, as glucose metabolism sensor, to regulate glucose uptake. Next, the effects of NFE2L1 knockdown on the key rate-limiting enzymes implicated in glycolysis, gluconeogenesis and tricarboxylic acid (TCA) cycle were tested (Figure 4A). As the results shown in Figure 4B, increase of *heme oxygenase 1* (*HMOX1*) and *prostaglandin-endoperoxide synthase 2* (*PTGS2/COX2*), and decrease of *PTGS1* were detected accompanied by NFE2L1 knockdown, suggesting the loss of function of NFE2L1 [3]. *HK1*, the key enzyme involved in catalyzing the conversion of glucose to glucose-6-phosphate, was significantly increased. *PFKL* and phosphofructokinase, muscle (*PFKM*), which catalyze the conversion of fructose-6-phosphate to fructose-1,6-bisphosphate, was reduced. The expression of *PKM*, which has been involved in catalyzing the conversion of phosphoenolpyruvate to pyruvate, was increased. No apparent change

of the key enzyme gene *dihydrolipoamide dehydrogenase* (*DLD*) that catalyzes the

entry of pyruvate into the TCA cycle was detected. The expression of the *pyruvate*

carboxylase (*PC*) gene was significantly reduced, indicating that the metabolic flow

from glycolysis to the TCA cycle might be restricted (Figure 4C).

Then the expression of key enzyme genes involved in TCA cycle was detected

and the results showed that *citrate synthase* (*CS*), which catalyzes to produce citrate,

and *oxoglutarate dehydrogenase* (*OGDH*), which catalyzes to produce succinyl-CoA,

exhibited no significant changes in mRNA levels. And *isocitrate dehydrogenase*

(NADP(+)) 1 (*IDH1*), which catalyzes the conversion of isocitrate to a-oxoglutarare,

was reduced. The level of *IDH2* and *IDH3G* was upregulated (Figure 4D). It can be

inferred from these results that the TCA cycle displayed no remarkable changes.

phosphoenolpyruvate carboxykinase 2 (*PCK2*) and *fructose-bisphosphatase 1* (*FBP1*),

the key rate-limiting enzymes in the gluconeogenesis process, were significantly

increased, while no difference of *glucose-6-phosphatase catalytic subunit 3* (*G6PC3*)

was detected (Figure 4E), indicating that the gluconeogenesis process might be

enhanced as NFE2L1 absence.

The western blotting result showed that NFE2L1 knockdown could efficiently

elevate the expression of NFE2L2, HMOX1, and PTGS2 and decrease kelch like

ECH associated protein 1 (KEAP1) (Figure 4F), which were consistent with our

previous results [3]. Then the changes in the protein levels of the key enzymes of

sugar metabolism were detected. The WB results of these key proteins were almost

the same as the gene expression (Figure 4F). The protein expression of IDH1 and

IDH2 both decreased. Combined with the significant decrease of PC and the increase

of PCK2, these results indicated the enhancement of the glycolysis and

gluconeogenesis, while the process from anaerobic glycolysis to aerobic oxidation

was been blockaded. Consistently, the transcriptome data also suggested that

glycolysis and gluconeogenesis were enhanced after knockdown of NFE2L1 in

HepG2 cells (Figure 4G).

In the process of glucose metabolism, loss of NFE2L1 may induce increased

glycolysis and gluconeogenesis and the suppressive oxidative phosphorylation.

These results indicated that NFE2L1 deficiency might trigger the Warburg effect,

consistent with that specific knockout of *NFE2L1* in liver tissue could trigger the

Non-alcoholic fatty liver disease (NAFLD) and hepatocellular carcinoma (HCC) in

the mouse model [2]. Analysis of metabolomic data showed that, in the *NFE2L1*

knockout HepG2 cells, the content of glucose metabolism intermediate products

glucose and pyruvate were increased, and amino acids closely related to glycolysis

intermediate products were significantly increased, such as glycerate, valine,

isoleucine, leucine, cysteate, cysteine, serine, glycine, and threonine (Figure 4H).

Malate, 2-oxoglutarate, and cis-aconitate, the TCA cycle intermediate products, were

significantly reduced (Figure 4H). These results implied that the absence of NFE2L1

produced a phenotype similar to the warburg effect. The enhancement of warburg

effect, one of the metabolic characteristics of tumor cells, often means the malignant

transformation of tumors. It is worth mentioning that the metabolome data showed

that the loss of NFE2L1 expression caused an increase in the levels of various amino

acids in cells, which might be the reason for the activation of mTOR signal caused by

NFE2L1 knockdown [13,29,30].

3.4. NFE2L1 knockdown causes the damage of mitochondrial function

Consistent with our results, previous studies had shown that overexpression and

knockout of NFE2L1 could change glucose metabolism through different ways such

as ROS, insulin secretion, and liver metabolism [6,7,12]. However, the underlying

molecular mechanism by which NFE2L1 affects glucose metabolism remains to be

explored. As a transcription factor, NFE2L1 could influence thousands of downstream

genes with various functions [3,31,32], making it difficult to identify the detailed

mechanism of NFE2L1 in affecting glucose metabolism. Actually, the effect of

NFE2L1 deficiency on glucose metabolic reprogramming is likely to be a systemic

response.

Previous result showed that the TCA cycle in the mitochondria was inhibited

(Figure 4), which made it necessary to explore the effects of NFE2L1 on

mitochondrial function. Under electron microscope, it was found that NFE2L1

knockdown significantly reduced the number of mitochondria. The remaining

mitochondria caused by NFE2L1 knockdown possessed much smaller size and the

ridge-like structure was significantly reduced (Figure 5A,B). In NFE2L1 knockdown

cells, accumulation of lipid droplets could be found, which is consistent with the lipid

accumulation induced by special knockout of NFE2L1 in mouse live [2]. In addition,

more autophagosomes could be found in NFE2L1 knockdown cells (Figure 5A,B),

suggesting that damaged mitochondria could be transferred to autophagic

mitochondria to be degraded. Consistently, studies have shown that AMPK can

promote mitochondrial division and mitochondrial autophagy [33,34], suggesting that

the alternation of mitochondrial caused by NFE2L1 knockdown may be related to

increased AMPK signals.

By testing the oxygen consumption rate (OCR) of the cells, it was found that the

oxygen consumption of the cells was reduced to about 40% of the control group after

NFE2L1 knockdown (Figure 5C). The oxygen consumption of the control cells was

reduced by 41.56%±6.04%, while in the NFE2L1 knockdown cells it was reduced by

22.31%±7.79% (Figure 5C) after adding OM, the oxidative phosphorylation inhibitor,

indicating that the proportion of adenosine triphosphate (ATP) produced by

mitochondrial respiration was significantly reduced after NFE2L1 knockdown. In the

control group, the ATP content was reduced under treatment with 2DG or OM (Figure

5d). In addition, the lactate content significantly increased after NFE2L1 knockdown

(Figure 5E). These results indicated that mitochondrial function was severely

impaired after NFE2L1 knockdown.

In cells lacking NFE2L1 expression, although mitochondrial function was

impaired and the oxidative phosphorylation process was inhibited, the ATP content

increased more than twice (Figure 5D), meaning that compared with the control group,

cells lacking NFE2L1 needed to consume dozens of times more glucose to produce

these ATP through glycolysis. This might be the reason why lack of NFE2L1 led to

406 increased cellular glucose uptake and cell death by glucose deprivation. Loss of
407 NFE2L1 resulted in increased ATP in cells and the changed mitochondria morphology,
408 suggesting the potential changes of ATP synthesis-related genes. Transcriptome data
409 showed that the mRNA level of *ATP5IF1*, *ATP5ME* and *ATP5PD* genes significantly
410 increased, and *ATP5F1E*, *ATP5MC3*, *ATP5MPL* and *ATP5PF* remarkably decreased
411 after NFE2L1 knockdown (Figure 5F). No significant difference of genes related to
412 ATP synthesis was observed in mitochondria (Figure 5G) despite the impacts of
413 NFE2L1 deficiency in the characterization of mitochondria, indicating NFE2L1 might
414 indirectly regulate the total amount of ATP. It is worth noting that NFE2L1
415 knockdown could still activate AMPK signal in the presence of high ATP
416 concentration (Figure 1G), suggesting that the influence of NFE2L1 on AMPK signal
417 may be through interfering with AMPK's perception of ATP and adenosine
418 monophosphate (AMP) levels in cells.

420 *3.5. NFE2L1 serves as an energy-sensitive protein and directly inhibits the*
421 *phosphorylation of AMPK by serine/threonine kinase 11 (STK11/LKB1)*

422  Lack of NFE2L1 could disrupt AMPK signal which is the core energy
423 metabolism regulation pathway [13-15]. Exploring the effect of NFE2L1 on AMPK
424 signal is helpful to understand the influence of NFE2L1 on the metabolic network. To
425 verify whether NFE2L1 is affected by ATP or AMP, exogenous ATP or AMP was used
426 to treat HepG2 cells. The western blotting results suggested that the addition of
427 exogenous ATP could increase the protein level of NFE2L1, while exogenous AMP
428 showed the opposite effects (Figure 6A). This result was consistent with the response
429 of NFE2L1 to glucose and serum (Figure 2), indicating that the response of NFE2L1
430 to the energy state might be due to ATP and AMP content in the cell. Subsequently,
431 the effect of AMP on the phosphorylation of AMPK after NFE2L1 knockdown was
432 tested, and the results indicated that NFE2L1 knockdown can reverse the activation of
433 AMPK signal induced by AMP (Figure 6B,C). Interestingly, previous study showed
434 NFE2L1 knockdown can reverse the activation of AMPK signal induced by MET [13].

435 These results were consistent, which showed the inhibition of NFE2L1 expression,

436 indicating that activation of AMPK signaling by AMP and MET depended on the

437 inhibitory effect of NFE2L1.

438  In the low-energy state, the ATP content in the cell decreases while the AMP

439 content increases, which induces AMPK phosphorylation by LKB1 [35-37]. The

440 activation of AMPK by LKB1 requires the co-localization of the two proteins in the

441 plasma membrane. Here, our results showed that loss of NFE2L1, which is a

442 membrane protein [38], inhibited AMPK signal activation induced by AMP,

443 suggesting that NFE2L1 might be involved in the process of AMPK phosphorylation

444 induced by LKB1.

445  In order to verify the effect of NFE2L1 on LKB1-induced AMPK

446 phosphorylation, LKB1 overexpressing and NFE2L1 knockdown HepG2 cells were

447 constructed. Unexpectedly, in HepG2 cells, overexpression of LKB1 could activate

448 AMPK signal and significantly inhibit the protein level of NFE2L1, but showed no

449 obvious effects on the mRNA level of *NFE2L1* (Figure 6D,E), indicating the

450 coincidence of NFE2L1 reduction and AMPK phosphorylation increase. In cells with

451 NFE2L1 deficiency, overexpression of LKB1 could not further increase AMPK

452 phosphorylation (Figure 6F,G). Glucose starvation (GS) experiment results showed

453 that glucose starvation for 2h significantly increased AMPK phosphorylation, and GS

454 could still further increase AMPK phosphorylation after LKB1 overexpression

455 (Figure 6F,G). The cumulative effects of LKB1 overexpression and GS treatment on

456 AMPK phosphorylation levels suggested that the response of AMPK to low-energy

457 states might not completely depend on LKB1. In the NFE2L1 knockdown cells, with

458 the increase of the basal AMPK phosphorylation, although GS treatment further

459 increased AMPK phosphorylation, the cumulative effect of LKB1 overexpression and

460 GS treatment disappeared (Figure 6F,G), suggesting that the effects of NFE2L1 on

461 AMPK signal might mainly relied on LKB1.

462  Subsequently, protein immunoprecipitation experiments were utilized to explore

463 the action mode of NFE2L1 on LKB1-AMPK. The results showed that AMPK could

interact with NFE2L1 protein rather than LKB1 (Figure 6H), suggesting that NFE2L1

might directly bind to AMPK and inhibit LKB1's phosphorylation. In addition, ACC

could also interact with NFE2L1 (Figure 6H). Combined with the negative correlation

between phosphorylated ACC and NFE2L1 (Figure 2A,B), we could speculate that

NFE2L1 may regulate lipid metabolism by directly binding to ACC and inhibiting its

phosphorylation.

4. Discussion

The rational use of nutrients is a necessary prerequisite for the survival of

multicellular organisms. In this process, the perception and feedback of the energy

state of cells is crucial. NFE2L1 has been considered to play an important role in

regulating system redox homeostasis, carbohydrate and lipid metabolism homeostasis,

embryonic development and cancer [4]. The transcription factor NFE2L1

systematically regulates the transcription of antioxidant-related genes through the

recognition of antioxidant/electrophile-response element (ARE/EPRE) [39]. However,

the molecular mechanism of metabolism-related phenotypes caused by NFE2L1

overexpression or knockout is still elusive. Identification of the molecular mechanism

of NFE2L1 in regulating substance and energy metabolism will deepen the

recognition of the molecular and physiological functions of NFE2L1, which could

benefit for the treatment of obesity, diabetes, non-alcoholic fatty liver disease and

other systemic metabolic diseases.

NFE2L1 is essential for maintaining intracellular redox homeostasis which is a

prerequisite for cells to maintain normal metabolic processes. Loss of NFE2L1 can

not only directly increase ROS levels, but also induce strong oxidative stress to cause

the death of embryos [1]. In the NFE2L1 knockdown cell line, glucose deprivation

could induce cell death, which might be related to the imbalance of redox homeostasis

(Figure 1). These results were consistent with previous studies [12]. NFE2L1 could

negatively regulate glucose uptake (Figure 3). These results demonstrated the dual

functions of NFE2L1 as the sensor and the regulator of glucose homeostasis.

The increased glucose uptake caused by NFE2L1 knockdown might be related to the increased of GLUT3 (Figure 3). Previous reports mostly focused on studying the relationship between NFE2L1 and GLUT1, GLUT2, GLUT4 respectively [6,7,12]. However, the interaction of NFE2L1 with GLUT3, the transporter with the highest affinity for glucose [40], has not been explored. Recently, more and more studies have shown that GLUT3 plays a vital role in regulating glycolysis, cancer stem cell survival, and cancer cell metastasis [41-43]. In addition, GLUT3, which can be activated by AMPK signals [44], is mainly expressed in tissues with high glucose requirements and poor glucose microenvironment, such as the brain [45]. Here, we found that knockdown of NFE2L1 could significantly increase GLUT3 expression, which might be attributed to AMPK activation. NFE2L1 specific knockout led to neurodegenerative diseases in the nervous system [46] and metabolic disorders and hepatocellular carcinogenesis in liver tissue [2]. However, the expression of GLUT3 gene was decreased in NFE2L1 knockdown cells, making it not easy to identify the underlying molecular mechanism. It is worth mentioning that the glycosylation and deglycosylation of NFE2L1 are necessary to activate its transcription factor activity [19]. Meanwhile, the glycation of NFE2L1 strictly depended on the glycolysis process (Figure 2E-H). These results suggested that the NFE2L1 functioned as a key factor in mediating the glucose homeostasis and redox homeostasis in cells.

Studies have shown that NFE2L1 is sensitive to nutrients in the environment [23]. However, the growth of cells lacking NFE2L1 depended almost solely on glucose. By detecting the key rate-limiting enzymes involved in glucose metabolism and the proteomic of glucose metabolites, we found that loss of NFE2L1 significantly inhibited the oxidative phosphorylation process and enhanced the glycolysis and gluconeogenesis process (Figure 4). Obviously, NFE2L1 deficiency in HepG2 cells triggered and enhanced the Warburg effect, which might be the cause of spontaneous NASH that could be finally developed into HCC in the liver of *NFE2L1* knockout mice [2].

Overexpression of NFE2L1 can reverse the mitochondrial damage caused by

N-glycanase 1 (NGLY1) knockout [25], suggesting that NFE2L1 is essential for the maintenance of mitochondrial morphology and function. Here, we also observed that knockdown of NFE2L1 can cause mitochondrial damage (Figure 5). However, the relationship between the Warburg effect and mitochondrial function damage caused by NFE2L1 deficiency is still hard to determine. Once mitochondria, as the most important organelle for the production of ATP, are damaged, cells may need to consume more glucose to produce ATP through glycolysis. As the results showed, glucose uptake capacity increased after NFE2L1 knockdown. And after deprivation of glucose, cells with NFE2L1 knockdown died. Strikingly, The ATP content in HepG2^{shNFE2L1} cells was significantly higher than that in HepG2^{shNC} cells (Figure 5D), which is consistent with the results of Zhu et al. [12]. OM, the inhibitor of oxidative phosphorylation, could hardly inhibit ATP production in HepG2^{shNFE2L1} cells, indicating that these ATP were rarely produced from the oxidative phosphorylation process (Figure 5D). In short, HepG2^{shNFE2L1} cells need to maintain a higher level of ATP content with lower ATP production efficiency. This may be the reason of the dependence of the growth of HepG2^{shNFE2L1} cells on glucose.

AMPK functions as the core regulator of energy metabolism to sense the levels of ATP and AMP. The binding of AMP with AMPK is necessary for the activation of AMPK by phosphorylation [37]. In the low energy state, the increase of AMP caused by the consumption of ATP can activate AMPK signals to promote catabolism and inhibit synthesis reactions which could finally produce more ATP [15]. The increase of phosphorylated AMPK level after NFE2L1 knockdown (Figure 1E) might be responsible for the reprogramming of glucose metabolism and the increase of ATP (Figures 3, 4). Under normal circumstances, high levels of ATP could inhibit AMPK signaling to maintain energy homeostasis. Obviously, NFE2L1 knockdown destroyed the energy homeostasis maintained by this mechanism. High ATP content and highly activated AMPK signals coexist in the HepG2^{shNFE2L1} cells, suggesting the problem of the negative feedback regulation of AMPK signals.

In the HepG2^{shNFE2L1} cells, AMP lost the ability of increasing the level of AMPK

phosphorylation (Figure 6B,C), which once again confirmed that NFE2L1 might be involved in AMPK signal perception of ATP and AMP. In other words, AMP binding is not necessary for the phosphorylation of AMPK in the absence of NFE2L1. Interestingly, various methods that can improve AMPK phosphorylation can also lead to decreased NFE2L1, including FBS starvation [23], glucose starvation [13], MET treatment [13], AMP treatment (Figure 6A), and LKB1 overexpression (Figure 6D). On the contrary, treatments with EGF and insulin [23] that can inhibit AMPK phosphorylation are accompanied by the increase of NFE2L1. In addition, NFE2L1 could interact with AMPK protein (Figure 6H). All these results revealed that NFE2L1 could bind to AMPK in the cytoplasm to inhibit the phosphorylation of AMPK in addition to the function as a transcription factor in the nucleus. In a low-energy state, the binding of NFE2L1 with AMPK reduced due to the lower expression level of NFE2L1 protein, which decreased the inhibition of NFE2L1 to AMPK activity (Figure 6I). Also, AMPK could bind with NFE2L1 and inhibit the degradation of NFE2L1. However, the sequential relationship between the decrease of NFE2L1 protein and the increase of AMPK phosphorylation is still unknown. Whether the effects of AMP/ATP and LKB1 on AMPK depend on NFE2L1 requires to be explored.

As an environment-sensitive protein, NFE2L1 could be changed in response to various stresses even the environmental temperature changes [4,47]. NFE2L1 could function as a transcription factor to directly regulate the transcription and expression of thousands of genes involved in almost all biological processes [32,48]. In addition, with the half-life of about 0.5h, NFE2L1 protein is always in the process of dynamic synthesis and degradation [13]. These characteristics make NFE2L1 own the ability of mediating the interaction and dialogue between the internal homeostasis and the external environment. In this study, we discovered for the first time that NFE2L1 could directly regulate the activity of AMPK signaling in a way independent of the function of transcription factor. The findings could deepen the understanding of the molecular mechanism of NFE2L1 in regulating material and energy metabolism, and

redox homeostasis, which is valuable for targeting NFE2L1 to prevent or treat
systemic diseases.

5. Conclusions

In conclusion, the results of glucose starvation, glucose uptake, selective inhibition
of glucose metabolism and protein deglycosylation experiments suggested that
NFE2L1 could act as the sensor and the regulator of glucose homeostasis.
Transcriptome, metabolome, seahorse and electron microscopy results revealed that
impaired NFE2L1 caused reprogram of glucose metabolism, damaged the
mitochondrial and aggravated the Warburg effect. And the energy signal transmission
and Co-IP experiments found that NFE2L1 could sense the energy state and regulate
AMPK signaling pathway by directly interact with AMPK protein. The novel
AMP/NFE2L1/AMPK signaling pathway that discovered in this study may be the
core mechanism of NFE2L1 in metabolism and metabolic diseases.

Supplementary Materials: The following are available online at xxx. **Table S1.** The
key resources in this study. **Table S2.** The transcriptome sequencing data of
HepG2^{shNC} and HepG2^{shNFE2L1} cell lines.
Author Contributions: LQ and YQ conceived the research and designed the
experiments; QY, WZ, YX, PL, XZ, HN, RS, SG, and YC conducted the experiments,
acquired and analyzed the data with the critical assistance from WJZ, YW, GL, ZC,
YR, YX, YG and YZ; all authors contribute to writing the manuscript; WZ, XZ, YX,
YZ and LQ revised the manuscript. All authors read and approved the final
manuscript.
Funding: This work was supported by the Postdoctoral Research Grant in Henan
Province (201901006, 201902006), the China Postdoctoral Science Foundation
(2020M672286), the Program for Innovative Talents of Science and Technology in
Henan Province (18HASTIT042), Science Foundation for Excellent Young Scholars

Data Availability Statement: The data presented in this study are available in this

manuscript.

Conflicts of Interest: The authors declare no conflict of interests.

References

1. Chen, L.Y.; Kwong, M.; Lu, R.H.; Ginzinger, D.; Lee, C.; Leung, L.; Chan, J.Y. Nrf1 is

critical for redox balance and survival of liver cells during development. *Mol Cell Biol*

2003, *23*, 4673-4686.

2. Xu, Z.R.; Chen, L.Y.; Leung, L.; Yen, T.S.B.; Lee, C.; Chan, J.Y. Liver-specific

inactivation of the nrf1 gene in adult mouse leads to nonalcoholic steatohepatitis and

hepatic neoplasia. *P Natl Acad Sci USA* **2005**, *102*, 4120-4125.

3. Qiu, L.; Wang, M.; Hu, S.; Ru, X.; Ren, Y.; Zhang, Z.; Yu, S.; Zhang, Y. Oncogenic

activation of nrf2, though as a master antioxidant transcription factor, liberated by

specific knockout of the full-length nrf1 alpha that acts as a dominant tumor repressor.

Cancers **2018**, *10*, 520.

4. Zhang, Y.G.; Xiang, Y.C. Molecular and cellular basis for the unique functioning of nrf1,

an indispensable transcription factor for maintaining cell homoeostasis and organ

integrity. *Biochemical Journal* **2016**, *473*, 961-1000.

5. Speliotes, E.K.; Willer, C.J.; Berndt, S.I.; Monda, K.L.; Thorleifsson, G.; Jackson, A.U.;

Lango Allen, H.; Lindgren, C.M.; Luan, J.; Magi, R., *et al.* Association analyses of

249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet*

2010, *42*, 937-948.

6.   Hirotsu, Y.; Higashi, C.; Fukutomi, T.; Katsuoka, F.; Tsujita, T.; Yagishita, Y.;

Matsuyama, Y.; Motohashi, H.; Uruno, A.; Yamamoto, M. Transcription factor

nf-e2-related factor 1 impairs glucose metabolism in mice. *Genes to cells : devoted to*

molecular & cellular mechanisms **2014**, *19*, 650-665.

7.   Zheng, H.Z.; Fu, J.Q.; Xue, P.; Zhao, R.; Dong, J.; Liu, D.X.; Yamamoto, M.; Tong,

Q.C.; Teng, W.P.; Qu, W.D., *et al.* Cnc-bzip protein nrf1-dependent regulation of

glucose-stimulated insulin secretion. *Antioxid Redox Sign* **2015**, *22*, 819-831.

8.   Hou, Y.Y.; Liu, Z.Y.; Zuo, Z.; Gao, T.C.; Fu, J.Q.; Wang, H.H.; Xu, Y.Y.; Liu, D.X.;

Yamamoto, M.; Zhu, B.B., *et al.* Adipocyte-specific deficiency of nfe2l1 disrupts

plasticity of white adipose tissues and metabolic homeostasis in mice. *Biochem Bioph*

Res Co **2018**, *503*, 264-270.

9.   Parola, M.; Novo, E. Nrf1 gene expression in the liver: A single gene linking oxidative

stress to nafld, nash and hepatic tumours. *J Hepatol* **2005**, *43*, 1096-1097.

10.   Zhang, Y.G.; Ren, Y.G.; Li, S.J.; Hayes, J.D. Transcription factor nrf1 is topologically

repartitioned across membranes to enable target gene transactivation through its

acidic glucose-responsive domains. *Plos One* **2014**, *9*, e93458.

11.   Chen, J.Y.; Liu, X.P.; Lue, F.L.; Lu, X.P.; Ru, Y.; Ren, Y.G.; Yao, L.B.; Zhang, Y.G.

Transcription factor nrf1 is negatively regulated by its o-glcnacylation status. *Febs Lett*

2015, *589*, 2347-2358.

12.   Zhu, Y.P.; Zheng, Z.; Xiang, Y.C.; Zhang, Y.G. Glucose starvation-induced rapid

death of nrf1 alpha-deficient, but not nrf2-deficient, hepatoma cells results from its

fatal defects in the redox metabolism reprogramming. *Oxidative Medicine and Cellular*

Longevity **2020**, *2020*, 4959821.

13.   Gou, S.; Qiu, L.; Yang, Q.; Li, P.; Zhou, X.; Sun, Y.; Zhou, X.; Zhao, W.; Zhai, W.; Li,

G., *et al.* Metformin leads to accumulation of reactive oxygen species by inhibiting the

nfe2l1 expression in human hepatocellular carcinoma cells. *Toxicol Appl Pharmacol*

2021, *420*, 115523.

14.   Garcia, D.; Shaw, R.J. Ampk: Mechanisms of cellular energy sensing and restoration

of metabolic balance. *Mol Cell* **2017**, *66*, 789-800.

15.   Lin, S.C.; Hardie, D.G. Ampk: Sensing glucose as well as cellular energy status. *Cell*

Metab **2018**, *27*, 299-313.

16.   Ishiyama, M.; Miyazono, Y.; Sasamoto, K.; Ohkura, Y.; Ueno, K. A highly

water-soluble disulfonated tetrazolium salt as a chromogenic indicator for nadh as well

as cell viability. *Talanta* **1997**, *44*, 1299-1305.

17.   Szychowski, K.A.; Rybczynska-Tkaczyk, K.; Leja, M.L.; Wojtowicz, A.K.; Gminski, J.

Tetrabromobisphenol a (tbbpa)-stimulated reactive oxygen species (ros) production in

cell-free model using the 2',7'-dichlorodihydrofluorescein diacetate (h2dcfda)

assay-limitations of method. *Environ Sci Pollut Res Int* **2016**, *23*, 12246-12252.

18.   Wang, H.F.; Sun, Y.X.; Zhou, X.M.; Chen, C.X.; Jiao, L.; Li, W.Q.; Gou, S.S.; Li, Y.Y.;

Du, J.F.; Chen, G.Y., *et al.* Cd47/sirp alpha blocking peptide identification and

synergistic effect with irradiation for cancer immunotherapy. *J Immunother Cancer*

2020, *8*, e000905.

19.   Zhang, Y.; Lucocq, J.M.; Yamamoto, M.; Hayes, J.D. The nhb1 (n-terminal homology

box 1) sequence in transcription factor nrf1 is required to anchor it to the endoplasmic

reticulum and also to enable its asparagine-glycosylation. *Biochemical Journal* **2007**,

408, 161-172.

20.   Liu, Y.; Cao, Y.Y.; Zhang, W.H.; Bergmeier, S.; Qian, Y.R.; Akbar, H.; Colvin, R.; Ding,

J.; Tong, L.Y.; Wu, S.Y., *et al.* A small-molecule inhibitor of glucose transporter 1

downregulates glycolysis, induces cell-cycle arrest, and inhibits cancer cell growth in

vitro and in vivo. *Mol Cancer Ther* **2012**, *11*, 1672-1682.

21.   Herzig, S.; Shaw, R.J. Ampk: Guardian of metabolism and mitochondrial homeostasis.

Nat Rev Mol Cell Biol **2018**, *19*, 121-135.

22.   Corton, J.M.; Gillespie, J.G.; Hardie, D.G. Role of the amp-activated protein kinase in

the cellular stress response. *Curr Biol* **1994**, *4*, 315-324.

23.   Zhang, Y.N.; Nicholatos, J.; Dreier, J.R.; Ricoult, S.J.H.; Widenmaier, S.B.;

Hotamisligil, G.S.; Kwiatkowski, D.J.; Manning, B.D. Coordinated regulation of protein

synthesis and degradation by mtorc1. *Nature* **2014**, *513*, 440-+.

24.   Lehrbach, N.J.; Breen, P.C.; Ruvkun, G. Protein sequence editing of skn-1a/nrf1 by

peptide:N-glycanase controls proteasome gene expression. *Cell* **2019**, *177*, 737-750

e715.

25.   Yang, K.; Huang, R.; Fujihira, H.; Suzuki, T.; Yan, N. N-glycanase ngly1 regulates

mitochondrial homeostasis and inflammation through nrf1. *J Exp Med* **2018**, *215*,

2600-2616.

26.   Marsh, J.; Mukherjee, P.; Seyfried, T.N. Drug/diet synergy for managing malignant

astrocytoma in mice: 2-deoxy-d-glucose and the restricted ketogenic diet. *Nutr Metab*

(Lond) **2008**, *5*, 33.

27.   Jastroch, M.; Divakaruni, A.S.; Mookerjee, S.; Treberg, J.R.; Brand, M.D.

Mitochondrial proton and electron leaks. *Essays Biochem* **2010**, *47*, 53-67.

28.   Cai, L.Y.; Jin, X.; Zhang, J.N.; Li, L.; Zhao, J.F. Metformin suppresses nrf2-mediated

chemoresistance in hepatocellular carcinoma cells by increasing glycolysis. *Aging-Us*

2020, *12*, 17582-17600.

29.   Jewell, J.L.; Russell, R.C.; Guan, K.L. Amino acid signalling upstream of mtor. *Nat*

Rev Mol Cell Biol **2013**, *14*, 133-139.

30.   Abu-Remaileh, M.; Wyant, G.A.; Kim, C.; Laqtom, N.N.; Abbasi, M.; Chan, S.H.;

Freinkman, E.; Sabatini, D.M. Lysosomal metabolomics reveals v-atpase- and

mtor-dependent regulation of amino acid efflux from lysosomes. *Science* **2017**, *358*,

807-813.

31.   Cui, M.; Wang, Z.; Chen, K.; Shah, A.M.; Tan, W.; Duan, L.; Sanchez-Ortiz, E.; Li, H.;

Xu, L.; Liu, N., *et al.* Dynamic transcriptional responses to injury of regenerative and

non-regenerative cardiomyocytes revealed by single-nucleus rna sequencing. *Dev*

Cell **2020**, *53*, 102-116 e108.

32.   Liu, P.; Kerins, M.J.; Tian, W.; Neupane, D.; Zhang, D.D.; Ooi, A. Differential and

overlapping targets of the transcriptional regulators nrf1, nrf2, and nrf3 in human cells.

J Biol Chem **2019**, *294*, 18131-18149.

33.   Toyama, E.Q.; Herzig, S.; Courchet, J.; Lewis, T.L.; Loson, O.C.; Hellberg, K.; Young,

N.P.; Chen, H.; Polleux, F.; Chan, D.C., *et al.* Amp-activated protein kinase mediates

mitochondrial fission in response to energy stress. *Science* **2016**, *351*, 275-281.

34.   Egan, D.F.; Shackelford, D.B.; Mihaylova, M.M.; Gelino, S.; Kohnz, R.A.; Mair, W.;

Vasquez, D.S.; Joshi, A.; Gwinn, D.M.; Taylor, R., *et al.* Phosphorylation of ulk1 (hatg1)

by amp-activated protein kinase connects energy sensing to mitophagy. *Science* **2011**,

331, 456-461.

35.   Houde, V.P.; Ritorto, M.S.; Gourlay, R.; Varghese, J.; Davies, P.; Shpiro, N.;

Sakamoto, K.; Alessi, D.R. Investigation of lkb1 ser(431) phosphorylation and cys(433)

farnesylation using mouse knockin analysis reveals an unexpected role of prenylation

in regulating ampk activity. *Biochemical Journal* **2014**, *458*, 41-56.

36.   Zhang, C.S.; Jiang, B.; Li, M.Q.; Zhu, M.J.; Peng, Y.Y.; Zhang, Y.L.; Wu, Y.Q.; Li, T.Y.;

Liang, Y.; Lu, Z.L., *et al.* The lysosomal v-atpase-ragulator complex is a common

activator for ampk and mtorc1, acting as a switch between catabolism and anabolism.

Cell Metab **2014**, *20*, 526-540.

37.   Zhang, Y.L.; Guo, H.; Zhang, C.S.; Lin, S.Y.; Yin, Z.; Peng, Y.; Luo, H.; Shi, Y.; Lian,

G.; Zhang, C., *et al.* Amp as a low-energy charge signal autonomously initiates

assembly of axin-ampk-lkb1 complex for ampk activation. *Cell Metab* **2013**, *18*,

546-555.

38.   Zhang, Y.G.; Hayes, J.D. Identification of topological determinants in the n-terminal

domain of transcription factor nrf1 that control its orientation in the endoplasmic

reticulum membrane. *Biochemical Journal* **2010**, *430*, 497-510.

39.   Johnsen, O.; Murphy, P.; Prydz, H.; Kolsto, A.B. Interaction of the cnc-bzip factor

tcf11/lcr-f1/nrf1 with mafg: Binding-site selection and regulation of transcription.

Nucleic Acids Res **1998**, *26*, 512-520.

40.   Gould, G.W.; Holman, G.D. The glucose transporter family: Structure, function and

tissue-specific expression. *Biochem J* **1993**, *295 (Pt 2)*, 329-341.

41.   Flavahan, W.A.; Wu, Q.L.; Hitomi, M.; Rahim, N.; Kim, Y.; Sloan, A.E.; Weil, R.J.;

Nakano, I.; Sarkaria, J.N.; Stringer, B.W.*, et al.* Brain tumor initiating cells adapt to

restricted nutrition through preferential glucose uptake. *Nat Neurosci* **2013**, *16*,

1373-1382.

42.   Wu, D.; Harrison, D.L.; Szasz, T.; Yeh, C.F.; Shentu, T.P.; Meliton, A.; Huang, R.T.;

Zhou, Z.; Mutlu, G.M.; Huang, J.*, et al.* Single-cell metabolic imaging reveals a

slc2a3-dependent glycolytic burst in motile endothelial cells. *Nat Metab* **2021**, *3*,

714-727.

43.   Yang, M.J.; Guo, Y.; Liu, X.Y.; Liu, N.Q. Hmga1 promotes hepatic metastasis of

colorectal cancer by inducing expression of glucose transporter 3 (glut3). *Med Sci*

Monitor **2020**, *26*, e924975.

44.   Dai, W.X.; Xu, Y.; Mo, S.B.; Li, Q.G.; Yu, J.; Wang, R.J.; Ma, Y.L.; Ni, Y.; Xiang, W.Q.;

Han, L.Y.*, et al.* Glut3 induced by ampk/creb1 axis is key for withstanding energy

stress and augments the efficacy of current colorectal cancer therapies. *Signal*

Transduct Tar **2020**, *5*, 177.

45.   Maher, F.; Vannucci, S.; Takeda, J.; Simpson, I.A. Expression of mouse-glut3 and

human-glut3 glucose transporter proteins in brain. *Biochem Biophys Res Commun*

1992, *182*, 703-711.

46.   Lee, C.S.; Lee, C.; Hu, T.; Nguyen, J.M.; Zhang, J.S.; Martin, M.V.; Vawter, M.P.;

Huang, E.J.; Chan, J.Y. Loss of nuclear factor e2-related factor 1 in the brain leads to

dysregulation of proteasome gene expression and neurodegeneration. *P Natl Acad*

Sci USA **2011**, *108*, 8408-8413.

47.   Bartelt, A.; Widenmaier, S.B.; Schlein, C.; Johann, K.; Goncalves, R.L.S.; Eguchi, K.;

Fischer, A.W.; Parlakgul, G.; Snyder, N.A.; Nguyen, T.B., *et al.* Brown adipose tissue

thermogenic adaptation requires nrf1-mediated proteasomal activity. *Nat Med* **2018**,

24, 292-303.

48.   Hamazaki, J.; Murata, S. Er-resident transcription factor nrf1 regulates proteasome

expression and beyond. *Int J Mol Sci* **2020**, *21*, 3683.

784 **Figures and Figure legend**

Figure 1. NFE2L1 knockdown caused HepG2 cells to be more sensitive to glucose starvation. (**A**) The total protein of HepG2^{shNC} and HepG2^{shNFE2L1} cells were collected, then the expression of NFE2L1 and β-actin were detected by WB. The HepG2^{shNC} and HepG2^{shNFE2L1} cells were cultured used the DMEM with or without glucose (2 g/L or 0 g/L) for 18h, the morphology of cells were observed under the microscope (**B**), and the survival rate of cells was detected by CCK8 kit (**C**). (**F**) The HepG2^{shNC} and HepG2^{shNFE2L1} cells were treated with WZB117 (200μM), and the survival rate of cells was detected in 0h, 24h, 48h, 72h by CCK8 kit, respectively. The HepG2^{shNC} and HepG2^{shNFE2L1} cells were cultured used the DMEM without glucose for 6h, then the ROS in cells were staining by DCFH-DA (10μM) for 20min and achieved by microscopy (**E**), and the intensity of fluorescence were counted (**D**), the scale is 50μm. (**G**) The total protein of HepG2^{shNC} and HepG2^{shNFE2L1} cells were collected, then the expression of NFE2L1, pAMPK, pACC and β-actin were detected by WB. n ≥ 3, '*' means $p < 0.05$.

Figure 2. NFE2L1 is a glucose sensitive protein. (**A**) HepG2 cells were cultured used DMEM medium with different volume ratios of FBS (0%, 1.25%, 2.5%, 5%, 10%, 20%), with or without glucose (4g/L or 0g/L) for 16h, and the total protein were collected, then the expression of NFE2L1, p-AMPK, p-ACC, p-mTOR, p-S6K1 and β-actin were detected by WB. (**B**) HepG2 cells were cultured used DMEM medium with different concentrations of glucose (0g/L, 0.5g/L, 1g/L, 2g/L, 4g/L, 8g/L), and with or without FBS (10% or 0%) for 16h, and the total protein were collected, then the expression of NFE2L1, p-AMPK, p-ACC, p-mTOR, p-S6K1 and β-actin were detected by WB. (**C**) The total protein change of the sample in (**A**) staining by coomassie brilliant blue. (**D**) The total protein change of the sample in (**B**) staining by coomassie brilliant blue. (**E**) HepG2 cells were glucose starvation for 4h and then cultured used DMEM with GLU (2g/L), FRU (30mM), GAL (30mM), MAN (30mM) and RIB (30mM) for 4h, the total protein were collected and the expression of NFE2L1 and β-actin were detected by WB. (**F**) The HepG2 cells were cultured used DMEM medium with GLU (0g/L), GLU (2g/L), FRU (30mM) or MAN (30mM) for 4h, the total protein were collected and Endo H were used to deglycosylate the glycated protein in vitro, then the expression of NFE2L1 and β-actin were detected by

WB. (**G**) The HepG2 cells were glucose starvation for 4h and then treated with GLU (2g/L), GLU (2g/L) + 2DG (20mM) or GLU (2g/L) + OM (10μM), and the total protein was collected after 4 hours, then the expression of NFE2L1, p-AMPK, p-ACC, p-mTOR, p-S6K1 and β-actin were detected by WB. (**H**) The HepG2 cells were glucose starvation for 4h and then treated with GLU (2g/L), FBP (30mM), or PYR (30mM) for 4h, the total protein was collected and then the expression of NFE2L1 and β-actin were detected by WB.

Figure 3. NFE2L1 can affect the uptake of glucose for HepG2 cells. (**A**) The HepG2[shNC] and HepG2[shNFE2L1] cells were treated with MET (1mM) for 12h, and then incubated in a serum-free medium containing 20μM of 2-NBDG at 37□ for 10min and the green fluorescent images were achieved by microscopy, the scale is 50μm. (**B**) The statistics show the results in (**A**). (**C**) The TPM value of *SLC2A1*, *SLC2A2*, *SLC2A3*, *SLC2A4*, *SLC2A5*, *SLC2A6*, *SLC2A7*, *SLC2A8*, *SLC2A9*, *SLC2A10*, *SLC2A11*, *SLC2A12*, *SLC2A13*, *SLC2A14* genes that related to glucose uptake of cell, date comes from transcriptome sequencing. (**D**) The expression of *SLC2A1-9* genes in HepG2[shNC] and HepG2[shNFE2L1] cells were detected by qPCR. The *SLC2A2*, *SLC2A5*,

SLC2A7 and *SLC2A9* have not be detected for it not expressed or extremely low, and

β-actin gene was used as the internal control. (**E**) The expression of NFE2L1, GLUT1,

GLUT3, GLUT4 and β-actin in HepG2^{shNC} and HepG2^{shNFE2L1} cells were detected by

WB. n ≥ 3, '*' means $p < 0.05$, '*n.s.*' means 'no significant'.

Figure 4. Knockdown of NFE2L1 leads to reprogramming of glucose metabolism. (**A**)

The schematic diagram of glycolysis, gluconeogenesis and TCA metabolism process

and key enzymes. (**B**) The expression of *NFE2L1*, *HMOX1*, *PTGS1* and *PTGS2* genes

in HepG2^{shNC} and HepG2^{shNFE2L1} cells were detected by qPCR, *β-actin* gene was used

as the internal control. (**C**) The expression of *HK1*, *HK2*, *PFKL*, *PFKM*, *PKM*, *PC*

and *DLD* genes in HepG2^{shNC} and HepG2^{shNFE2L1} cells were detected by qPCR,

β-actin gene was used as the internal control. (**D**) The expression of *CS*, *IDH1*, *IDH2*,

IDH3A, *IDH3B*, *IDH3G* and *OGDH* genes in HepG2^{shNC} and HepG2^{shNFE2L1} cells

were detected by qPCR, *β-actin* gene was used as the internal control. (**E**) The

expression of *NFE2L1*, *G6PC3*, *FBP1* and *PCK2* genes in HepG2^{shNC} and

HepG2^{shNFE2L1} cells were detected by qPCR, *β-actin* gene was used as the internal

control. (**F**) The expression of NFE2L1, KEAP1, NFE2L2, HMOX1, PTGS2, HK1,

PFKL, PKM, PC, IDH1, IDH2, PCK2 and β-actin in HepG2^{shNC} and HepG2^{shNFE2L1}

cells were detected by WB. (**G**) The expression of key enzyme for glucose

metabolism in the transcriptome data. (**H**) The results of the metabolome showed the

effect of NFE2L1 on Central Metabolism. n ≥ 3, '*****' means $p < 0.05$.

Figure 5. The effect of NFE2L1 knockdown in HepG2 cells on the morphology and function of mitochondria. The morphology of mitochondria in HepG2[shNC] cells (**A**) and HepG2[shNFE2L1] cells (**B**) were detected by electron microscope. (**C**) The oxygen consumption rate (OCR) of HepG2[shNC] and HepG2[shNFE2L1] cells were measured by Seahorse XF Cell Mito Stress Test Kit. (**D**) The HepG2[shNC] and HepG2[shNFE2L1] cells were treated with 2DG (20mM) or OM (10μM) for 12h, and the content of ATP in cells were detected by ATP assay kit. (**E**) The content of lactate in HepG2[shNC] and

HepG2^{shNFE2L1} cells were detected by Lactic Acid assay kit. (**F**) The TPM value of

ATPAF1, *ATPAF2*, *ATP5F1A*, *ATP5F1B*, *ATP5F1C*, *ATP5F1D*, *ATP5F1E*, *ATP5IF1*,

ATP5MC1, *ATP5MC2*, *ATP5MC3*, *ATP5MD*, *ATP5ME*, *ATP5MF*, *ATP5MG*,

ATP5MGL, *ATP5MPL*, *ATP5PB*, *ATP5PD*, *ATP5PF*, *ATP5PO* and *CLPB* genes that

related to ATP synthesis in mitochondria in HepG2^{shNC} and HepG2^{shNFE2L1} cells, date

comes from transcriptome sequencing. (**G**) The total TPM value of genes in (**F**). n $\geq$ 3,

'*'means $p < 0.05$, '*n.s.*' means 'no significant'.

Figure 6. NFE2L1 disrupts the phosphorylation of AMPK by LKB1 by directly

interacting with AMPK. (**A**) The HepG2^{shNC} and HepG2^{shNFE2L1} cells were treated

with ATP (500μM) or AMP (500μM) for 1h, 2h, 3h, respectively. The expression of

NFE2L1 and β-actin were detected by WB. (**B**) The HepG2^{shNC} and HepG2^{shNFE2L1}

cells were treated with AMP (500μM) for 2h and the expression of NFE2L1, p-AMPK

and β-actin were detected by WB. (**C**) Count and calculate the relative fold change of

p-AMPK in (**B**). (**D**) The expression of LKB1, p-AMPK, p-ACC, NFE2L1, HO1 and

β-actin in HepG2^{EGFP} and HepG2^{LKB1} cells were detected by WB. HepG2^{EGFP} and

HepG2^{LKB1} cells were constructed by lentiviral infected. (**E**) The expression of *LKB1*

and *NFE2L1* in HepG2^{EGFP} and HepG2^{LKB1} cells were detected by qPCR, with the

β-actin was used as the internal control. (**F**) HepG2^{EGFP}, HepG2^{LKB1}, HepG2^{shNC} and

HepG2$^{shNFE2L1+LKB1}$ cells were treated with glucose starvation (GS) for 4h, and the

expression of NFE2L1, LKB1, p-AMPK, p-ACC and β-actin were detected by WB.

(**G**) Count and calculate the relative fold change of p-AMPK in (**F**). (**H**) The NFE2L1

overexpression plasmid *pLVX-NFE2L1-puro* transfected into HEK293T cells, and the

total protein was collected with non-denatured lysate buffer after 48h. The specific

antibodies of NFE2L1, LKB1, AMPK, and ACC protein were used for Co-IP

experiments, then NFE2L1 and β-actin were detected by WB. 'L' means 'Low

exposure' and 'H' means 'Heavy exposure'. (**I**) The model diagram of NFE2L1

involved in regulating AMPK signal. n ≥ 3, '*' means $p < 0.05$, '*n.s.*' means 'no

significant'.